

RESEARCH ARTICLE

Functional characterization of a new terpene synthase from *Plectranthus amboinicus*


Nur Suhanawati Ashaari¹, Mohd Hairul Ab. Rahim^{1^{aa}}, Suriana Sabri^{2^{ab}}, Kok Song Lai³, Adelene Ai-Lian Song², Raha Abdul Rahim¹, Wan Muhamad Asrul Nizam Wan Abdullah¹, Janna Ong Abdullah^{1*}

1 Department of Cell and Molecular Biology, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Selangor, Malaysia, **2** Department of Microbiology, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Selangor, Malaysia, **3** Health Sciences Division, Abu Dhabi Women's College, Higher Colleges of Technology, Abu Dhabi, United Arab of Emirates

^{aa} Current address: Department of Industrial Biotechnology, Faculty of Industrial Sciences and Technology, Universiti Malaysia Pahang, Kuantan, Pahang, Malaysia

^{ab} Current address: Enzyme and Microbial Technology Research Center, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Selangor, Malaysia

* janna@upm.edu.my



OPEN ACCESS

Citation: Ashaari NS, Ab. Rahim MH, Sabri S, Lai KS, Song AA-L, Abdul Rahim R, et al. (2020) Functional characterization of a new terpene synthase from *Plectranthus amboinicus*. PLoS ONE 15(7): e0235416. <https://doi.org/10.1371/journal.pone.0235416>

Editor: Björn Hamberger, Michigan State University, UNITED STATES

Received: April 7, 2020

Accepted: June 15, 2020

Published: July 2, 2020

Copyright: © 2020 Ashaari et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files. GenBank accession numbers are within the paper, under the 'Isolation of full-length monoterpene synthase gene and sequence characterization' section.

Funding: This work was supported by Ministry of Science, Technology and Innovation, Malaysia (Fundamental Research Grant Scheme 07-02-14-1504FR) and Universiti Putra Malaysia (UPM Putra Grant GP-IPS/2015/9463800). The funders had no

Abstract

Plectranthus amboinicus (Lour.) Spreng is an aromatic medicinal herb known for its therapeutic and nutritional properties attributed by the presence of monoterpene and sesquiterpene compounds. Up until now, research on terpenoid biosynthesis has focused on a few mint species with economic importance such as thyme and oregano, yet the terpene synthases responsible for monoterpene production in *P. amboinicus* have not been described. Here we report the isolation, heterologous expression and functional characterization of a terpene synthase involved in *P. amboinicus* terpenoid biosynthesis. A putative monoterpene synthase gene (*PamTps1*) from *P. amboinicus* was isolated with an open reading frame of 1797 bp encoding a predicted protein of 598 amino acids with molecular weight of 69.6 kDa. *PamTps1* shares 60–70% amino acid sequence similarity with other known terpene synthases of Lamiaceae. The *in vitro* enzymatic activity of *PamTps1* demonstrated the conversion of geranyl pyrophosphate and farnesyl pyrophosphate exclusively into linalool and nerolidol, respectively, and thus *PamTps1* was classified as a linalool/nerolidol synthase. *In vivo* activity of *PamTps1* in a recombinant *Escherichia coli* strain revealed production of linalool and nerolidol which correlated with its *in vitro* activity. This outcome validated the multi-substrate usage of this enzyme in producing linalool and nerolidol both in *in vivo* and *in vitro* systems. The transcript level of *PamTps1* was prominent in the leaf during daytime as compared to the stem. Gas chromatography-mass spectrometry (GC-MS) and quantitative real-time PCR analyses showed that maximal linalool level was released during the daytime and lower at night following a diurnal circadian pattern which correlated with the *PamTps1* expression pattern. The *PamTps1* cloned herein provides a molecular basis for the terpenoid biosynthesis in this local herb that could be exploited for valuable production using metabolic engineering in both microbial and plant systems.

role in study design, data collection and analysis, decision to publish, or preparation of the manuscript for both funds.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Plectranthus amboinicus (Lour.) Spreng, locally known as *Bangun-bangun* in Malaysia, is a medicinal herb that belongs in the Lamiaceae family along with the herbs sage, thyme, basil and oregano. It is found naturally throughout the tropics and warm regions of Africa, Asia and Australia [1]. This plant is characterized by its green, succulent, heart-shaped leaves with scalloped edges and can grow to about 50 cm tall with horizontal stems up to 180 cm long [2]. It possesses a distinctive oregano-like odor and flavor that make it an excellent ingredient for culinary purposes. This herb has been traditionally used for treatment of coughs, sore throats, nasal congestion [2], animal and insect bites and also as a breast milk stimulant for hundreds of years [3]. However, the last decade witnessed a large increase of scientific interest in *P. amboinicus* research, as evidenced by increasing numbers of publications related to the bioactivities of *P. amboinicus*. This plant extract exhibited antibacterial activity against methicillin-resistant *Staphylococcus aureus* in a murine model [4], and was proven effective against reproductive tract infections by *Candida albicans*, *Proteus vulgaris* and *Klebsiella pneumoniae* [5]. It has been reported that *P. amboinicus* possesses anti-inflammatory and antitumor activities [6], larvicidal [7], and antithrombotic and antioxidant activities [8]. These therapeutic and medicinal properties of *P. amboinicus* are mainly attributed to its natural phytochemical compounds present in the essential oils or the plant extract. Essential oils of *P. amboinicus* are known to contain high amounts of bioactive compounds, mainly monoterpenoids such as carvacrol, thymol, γ -terpinene, α -terpineol and p -cymene with various pharmacological properties [1].

Monoterpenes are C_{10} terpenoids, usually produced by plants, with increasing industrial and clinical applications. In higher plants, terpenoids are synthesized *via* two independent pathways located in separate intracellular compartments. The methylerythritol (MEP) pathway is located in the plastid and responsible for production of mono- (C_{10}), di- (C_{20}) and tetraterpenes (C_{40}), while sesqui- (C_{15}), tri- (C_{30}) and polyterpenes (C_n) are synthesized *via* the mevalonate (MVA) pathway in the cytosol. Both pathways produce the universal precursors, isopentenyl pyrophosphate (IPP) and dimethylallyl pyrophosphate (DMAPP), for terpenoids biosynthesis. Further condensation of the two precursors gives rise to linear prenyl pyrophosphate precursors, geranyl pyrophosphate (GPP, C_{10}), farnesyl pyrophosphate (FPP, C_{15}) and geranyl geranyl pyrophosphate (GGPP, C_{20}), and terpene synthases are the key enzymes responsible for catalyzing these substrates into a variety of terpenoids found in plants [9,10]. All terpene synthases have similar properties with respect to their native molecular mass (monomers or homodimers) and requirement for divalent metal ions such as Mg^{2+} and Mn^{2+} for activity. The sequence similarities between terpene synthases are dominated by species relationship regardless of substrate or product specificity, and it was reported that many terpene synthases catalyze formation of multiple products [9,11].

Plants in the Lamiaceae family are known to produce a large variety of terpenoids, and this diversity could be due to expression of multiple terpene synthases and formation of multiple products by individual terpene synthases as previously reported [12–15]. Terpene synthase genes have been previously isolated and characterized from several important Lamiaceae members including *Thymus caespitius* [13,14], *T. vulgaris* [16], *Coriandrum sativum* L. [15], *Origanum vulgare* [12] and *Salvia officinalis* [17]; however the terpene synthases responsible for terpenoids production in *P. amboinicus* have not been functionally characterized. Thus, the main objectives of this present study were to isolate and clone the full-length transcript of *P. amboinicus* monoterpene synthase, and to functionally express and characterize the recombinant terpene synthase in both *in vivo* and *in vitro* systems. The expression pattern of the isolated transcript that was involved in the formation and emission of terpenoids in different plant parts under day/night period was also explained. The information obtained from this

study will benefit future exploitations of the isolated enzyme for terpenes biosynthesis in a simpler heterologous microbial or plant system.

Materials and methods

Plant material

The *P. amboinicus* leaves were collected from plants grown at the Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Selangor, Malaysia. The plant was identified and authenticated taxonomically at the School of Environmental Science and Natural Resources, Universiti Kebangsaan Malaysia, Selangor, Malaysia. The voucher specimen, UKMB40411, was deposited in the herbarium facility. Fresh plant samples were used directly for GC-MS analysis of the volatile compounds. For RNA extraction, the leaves were harvested and immediately frozen in liquid nitrogen and kept at -80°C until ready for the extraction process.

GC-MS analysis of *Plectranthus amboinicus* volatiles

The volatiles released from the *P. amboinicus* leaves and stems harvested at 2.00 AM, 8.00 AM, 2.00 PM and 8.00 PM were collected using headspace solid phase microextraction (HS-SPME) equipped with a 100 μm polydimethylsiloxane (PDMS) fiber (Supelco) and analyzed using an Agilent 7890A gas chromatograph coupled to an Agilent 5975C quadrupole mass detector (Agilent Technologies, Santa Clara, USA). The instrument was equipped with an Agilent HP-5MS capillary column (30 m x 250 μm inner diameter x 0.25 μm film) and helium was used as the carrier gas at a flow rate of 1 ml per min. The SPME fiber was conditioned at the GC injection port at 250°C for 5 min before use. Approximately 1 g of the *P. amboinicus* tissue was placed in a 20 ml headspace vial fitted with a silicone septum screw cap. Following 10 min of sample conditioning at room temperature, the SPME fiber was exposed to the headspace for 30 min at 60°C and immediately desorbed in the gas chromatograph injector at 250°C for 15 min using a splitless mode. The GC oven was maintained at 40°C for 2 min, gradually increased to 175°C at a rate of $5^{\circ}\text{C}/\text{min}$ and then an increment up to 250°C at $90^{\circ}\text{C}/\text{min}$. Linear retention index (LRI) was determined through the injection of a C8 to C20 series of straight chain n-alkanes (Sigma Aldrich, USA) and calculated in accordance to van Den Dool and Kratz [18]. The volatile compounds were identified by mass spectra comparison using a MSD Chemstation Enhanced Data Analysis Software (E.02.02.1431 version, Agilent Technologies) and the National Institute of Standards and Technology library database (NIST 14). In addition, the compounds were tentatively identified by comparing the experimental retention indices with the theoretical ones obtained from the literatures. The relative amount of the individual component was expressed as a percentage of the peak area of respective compound over the total peak areas of all identified volatiles.

Isolation of full-length *P. amboinicus* monoterpene synthase gene (*PamTps1*)

Based on the *P. amboinicus* transcriptome library (SRA Accession No.:SRR7842030) previously created from matured leaves and sequenced using the MiSeq Illumina platform (Mohd Hairul Ab. Rahim, unpublished), a 1361 bp partial terpene synthase transcript (Accession No.: GGXS01005129) that showed the highest sequence similarity to known plant monoterpene synthases was selected and designated as *PamTps1*. Total RNA was extracted using Tri Reagent (MRC, USA) following the manufacturer's protocol. The RNA was quantified and its purity was determined using UV-VIS spectrophotometer (NanoDrop 1000, Thermo Scientific,

USA), and its integrity was assessed on a 1.2% (w/v) agarose gel. Rapid Amplification of cDNA Ends (RACE) was conducted using SMARTer RACE Kit according to the manufacturer's protocol with minor modifications (Clontech, USA). For the first strand cDNA synthesis, 1 µg of total RNA was reverse transcribed using SMARTScribe Reverse Transcriptase (Clontech, USA) according to the manufacturer's instructions. The components of the RACE reactions were 1X Advantage 2 PCR buffer, 0.8 mM dNTP mix, 1 µM gene-specific primer (5' - CCC TATCCCTCACAAATGGGAGTTTCT-3'), 1X Universal Primer A Mix, 2.5 µl of 5'-RACE-Ready cDNA, 1X Advantage 2 Polymerase Mix and PCR-grade water added to a final volume of 50 µl. The gene-specific primer was designed at the conserved region of the partial sequence obtained from the transcriptomic data (SRA Accession No.:SRR7842030 and Accession No.:GGXS01005129) that fulfilled the requirements of 23–28 bp and 50–70% GC contents with a melting temperature ranging from 65 to 72°C. The gene-specific primer was synthesized by Bioneer, Korea. The RACE amplification was conducted using a touch-down program set at 5 cycles of 94°C for 30 sec, 72°C for 3 min, 5 cycles of 94°C for 30 sec, 70°C for 30 sec, 72°C for 3 min, and 25 cycles of 94°C for 30 sec, 68°C for 30 sec, 72°C for 3 min, and a final extension of 72°C for 10 min. The amplified product was cloned into pGEMT-Easy vector (Promega) and sent for sequencing (Bioneer, Korea). Based on the assembled fragments obtained by RACE and the known partial sequence, the full-length cDNA was then amplified in a PCR reaction containing 1X Pfx buffer, 0.3 mM dNTP mixture, 2 mM MgSO₄, 0.5 U of Platinum™ Pfx DNA Polymerase (Thermo Fisher Scientific, USA), 0.8 µM of *PamTps1*-F (5' -CAACGC AGAGTACATGGGATGGAGCAA-3') and *PamTps1*-R (5' -GCATTTGTTTCAGACATATGGA TGGAACAGC-3') primers, and 2.5 µl of 5'-RACE-Ready cDNA. Amplification was done at 94°C for 2 min, followed by 35 cycles of 94°C for 15 sec, 68°C for 1 min, 68°C for 3 min, and a 10 min final extension at 68°C. The successful amplicon was cloned into a pGEMT-Easy vector (Promega) and sent for sequencing (Bioneer, Korea).

Full-length sequence analysis and phylogenetic tree construction

The full-length transcript of *PamTps1* was aligned against the non-redundant protein database via the BLASTx algorithm. The N-terminal signal peptide sequence was predicted using ChloroP 1.1 Server (<http://www.cbs.dtu.dk/services/ChloroP/>) [19] and TargetP 1.1 Server (<http://www.cbs.dtu.dk/services/TargetP/>) [20]. The presence of conserved motifs shared by all known terpene synthases were identified by protein sequence alignments between *PamTps1* and linalool synthase of *Perilla frutescens* var. *hirtella* (ACN42013.2), *P. citriodora* (AAX16075.1) and *Lavandula latifolia* (ABD77417.1), *S. rosmarinus* pinene synthase (ABP01684.1), *P. setoyensis* geraniol synthase (ACN42010.1), γ -terpinene synthase of *T. caespititius* (AID51195.1), *T. serpyllum* (AGT29345.1) and *O. syriacum* (AEO27879.1), *L. x intermedia* 3-carene synthase (ARA91313.1), *T. caespititius* α -terpineol synthase (AGK88250.1) and *O. vulgare* terpene synthase 5 (ADK73617.1) using Clustal Omega (<https://www.ebi.ac.uk/Tools/msa/clustalo/>) [21] and BoxShade version 3.21 server (https://embnet.vital-it.ch/software/BOX_form.html). The evolutionary relationships of *PamTps1* were inferred using the Neighbor-Joining method and the bootstrap consensus tree was inferred with 1000 replicates computed via the PHYLogeny Inference Package version 3.695 (PHYLIP) [22].

Functional expression of *PamTps1*

The ORF of *PamTps1* excluding the N-terminal transit peptide was amplified using *PamTps1* forward (5' -ATTC**AAGCTT**ATGAAGCCCCGCTGTTGAAGCC-3') and reverse (5' -ATTC**CTCGAGTCCAGAGCC**GACATATGGATGGAACAG-3') primers with restriction enzyme (RE) sites shown in italics to create *Hind*III and *Xho*I (New England Biolabs, Canada) overhangs for

use during ligation into a pET32b(+) bacterial expression vector (Merck, Germany), respectively. The three amino acids spacer of GSG (underlined) was also incorporated into the reverse primer. The PCR components and programs were as previously described with an annealing temperature of 64°C. The successful amplicon was cloned into the pET32b(+) expression vector (Merck Millipore, USA) and transformed into *E. coli* Rosetta™ 2 (DE3) competent cells (Merck Millipore, USA), and presence of the insert was confirmed using RE digestions, followed by sequencing.

A single colony of recombinant expression cells harboring pET32b:*PamTps1* and empty vector pET32b, respectively, was inoculated into 10 ml Luria-Bertani (LB) medium containing 50 µg/ml carbenicillin and 34 µg/ml of chloramphenicol and grown at 37°C, 250 rpm overnight. Approximately 5 ml of the overnight culture was transferred into 95 ml of fresh LB medium supplemented with the same antibiotics and grown at 37°C until OD_{600nm} ~ 0.6–0.8 was reached. Protein expression was induced by the addition of isopropyl-β-D-thiogalactopyranoside (IPTG) to a final concentration of 0.5 mM and incubated with shaking at 150 rpm, 28°C for 16 h. After induction, the cells were harvested by centrifugation at 3214 x g, 4°C for 30 min and kept frozen at -20°C until ready for protein extraction. Protein extraction was conducted using the BugBuster Protein Extraction Reagent (Novagen, USA) according to the manufacturer's instructions. The lysate was then centrifuged at 12,633 x g, 4°C for 20 min to obtain the soluble protein fraction while the insoluble pellet was treated with 6 M urea at 4°C overnight followed by centrifugation to recover the inclusion body (IB) fraction. Buffer exchange to assay buffer containing 10 mM Tris-Cl, 10% (v/v) glycerol, 1 mM dithiothreitol (DTT) and 10 mM MgCl₂ at pH 7.5 [14] was conducted using Amicon® Centrifugal Filter (MW cutoff = 30 kDa) (Merck Millipore, USA) following the manufacturer's instructions.

The soluble protein fraction containing the His-tagged protein was purified using a HisTrap™ HP 5 ml column (GE Healthcare, USA) according to the manufacturer's instructions. The eluted fractions were pooled and concentrated to 1 ml using Amicon® Centrifugal Filter and further purified by gel filtration chromatography using Superdex 200 10/300 GL column (GE Healthcare, USA). The fractions containing pure *PamTps1* were eluted isocratically using the assay buffer at a flow rate of 0.5 ml/min.

The purified recombinant *PamTps1* including the soluble and insoluble protein extracts were resolved on 10% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and visualized by Coomassie blue staining. Western blotting for detection of the His-tagged protein was also conducted using anti-polyHistidine-HRP antibody (Sigma-Aldrich, USA) (dilution 1:2000) and detected using the SuperSignal® West Pico Chemiluminescent HRP substrate (Pierce, USA) following the manufacturer's instructions.

Functional characterization of *PamTps1*

Enzyme assays were performed accordingly in a glass vial using 25 µg recombinant protein in 100 µl reaction buffer containing 10 mM Tris-Cl, 10% (v/v) glycerol, 1 mM DTT and 10 mM MgCl₂ [14]. Phosphatase inhibitors of sodium tungstate (Na₂WO₄) and sodium fluoride (NaF) were added at final concentrations of 0.1 mM and 0.05 mM, respectively, to prevent geraniol formation. The enzymatic reaction was initiated by addition of 20 µM GPP or FPP (Sigma Aldrich, USA) and incubated at 30°C with constant shaking for 2 h. Terpene products were collected by using a SPME fiber exposed for 30 min in the headspace above the assay mixture at 60°C in a water bath and analyzed by GC-MS. The oven temperature was programmed at 50°C and gradually increased to 280°C at a rate of 10°C/min for 3 min. The temperature of the ion source and transfer line were set at 220°C and 280°C, respectively, and the electron impact mass spectra were recorded at 70 eV ionization energy. The identities of the terpene

compounds were determined with referral to the NIST 14 library, as well as comparison of mass spectra and retention times with authentic standards (Sigma-Aldrich, USA). The *in vivo* activity of *PamTps1* was conducted using the recombinant *E. coli* strain harboring the expression construct pET32b:*PamTps1* grown under the conditions as described in the protein expression section and the *E. coli* host harboring an empty vector was used as a control. Cultures were sampled at 24 h of post-induction and the volatile terpenoid compounds in the headspace of each culture were analyzed using HS-SPME-GC-MS as described in the assay reactions.

Gene expression analysis of *PamTps1*

For the expression analysis of *PamTps1* in various tissues, leaf and stem samples were collected at 8.00 AM, 2.00 PM, 8.00 PM and 2.00 AM with three biological replicates. The plant samples were immediately frozen and stored at -80°C until the extraction process using the Tri Reagent (MRC, USA). Total RNA of 1 μg was treated for genomic elimination and reverse transcribed using QuantiTect[®] Reverse Transcription kit (Qiagen, USA) according to the manufacturer's protocol. Quantitative RT-PCR was conducted to determine *PamTps1* transcript abundance involved in the production of linalool and nerolidol. A 291 bp fragment in the 3' region of *PamTps1* was amplified using the gene-specific primers (S1 Table). The experiments were performed using QuantiNova SYBR Green PCR Master Mix (Qiagen, USA) as per manufacturer's procedure and the real-time cycler program as follows: 95°C for 2 min, 40 cycles of 95°C for 5 s, 60°C for 10 s, followed by a melting curve analysis of $65\text{--}95^{\circ}\text{C}$ with 0.5°C increments. Sequencing analysis was performed to verify amplification of the expected region of *PamTps1*. Each primers pair was validated using a standard curve of serial cDNA dilutions to calculate the correlation coefficient and amplification efficiency. Three reference genes namely elongation factor G (*EF-G*), tubulin and adenine phosphoribosyl transferase (*APRT*) were used for normalization of the qPCR data. The primers for the qPCR analysis are listed in the S1 Table.

Results and discussion

Volatiles profiling of *Plectranthus amboinicus* leaves

Headspace—solid phase microextraction—gas chromatography—mass spectrometry (HS-SPME-GC-MS) analysis demonstrated that the *P. amboinicus* volatiles were dominated by α -bergamotene (19.5%), carvacrol (19.6%), caryophyllene (19.2%), p-cymene (8.2%), γ -terpinene (10.5%) and humulene (5.6%) (Table 1, Fig 1). Seven terpene alcohols including 1-octen-3-ol, terpinene-4-ol, linalool and nerolidol were detected which formed 2.35% of *P. amboinicus* volatiles. Carvacrol has a characteristic pungent spicy-woody odor which contributes to the strong oregano-like aroma [23], and together with the high relative percentages of citrus odor of α -bergamotene [24] and caryophyllene (spicy-woody) [25] collectively contribute to the unique, strong aromatic odor of *P. amboinicus*. This result is in agreement with the known volatile constituents of *P. amboinicus* previously reviewed by Arumugam et al. [1]. However, the phytochemical composition of plants may vary depending on geographical locations, climatic conditions, methods of extraction and identification. The volatile constituents of *P. amboinicus* leaves collected from Uganda which were also extracted using HS-SPME showed the presence of linalool (50.3%) as the main component with other detected volatiles such as carvacrol (14.34%), geranyl acetate (11.7%), nerol acetate (11.6%) and γ -terpinene (3.2%) [26]. On the other hand, *P. amboinicus* oil from Cambodia was shown to contain dominant constituents of thymol (57.4%), carvacrol (13.5%), γ -terpinene (5.6%) and p-cymene (5.2%) [27]. Interestingly, thymoquinone and thymohydroquinone, phytochemical

Table 1. Compounds identified from *Plectranthus amboinicus* leaf volatiles harvested at 8.00 AM using HS-SPME-GC-MS analysis.

No	RT ^a (min)	Compounds	RI ^b	Relative content ^c (%)
1	8.304	β -Thujene	924	0.432 \pm 0.048
2	10.081	1-Octen-3-ol	983	1.140 \pm 0.054
3	10.279	β -Myrcene	990	1.372 \pm 0.147
4	10.653	α -Phellandrene	1002	0.458 \pm 0.030
5	11.043	α -Terpinene	1015	2.508 \pm 0.212
6	11.434	ρ -Cymene	1027	8.208 \pm 0.445
7	12.069	β -Ocimene	1048	0.149 \pm 0.015
8	12.512	γ -Terpinene	1063	10.475 \pm 0.465
9	13.229	Terpinolene	1086	0.161 \pm 0.003
10	13.683	Linalool	1101	0.163 \pm 0.020
11	15.665	Isoborneol	1169	0.131 \pm 0.012
12	15.951	(-)-Terpinen-4-ol	1178	0.821 \pm 0.026
13	16.370	α -Terpineol	1193	0.021 \pm 0.010
14	18.032	Thymoquinone	1252	2.296 \pm 0.753
15	19.343	Thymol	1294	0.331 \pm 0.013
16	20.171	Carvacrol	1326	19.583 \pm 1.937
17	21.144	Eugenol	1363	0.042 \pm 0.004
18	22.794	Caryophyllene	1430	19.211 \pm 2.355
19	23.213	α -Bergamotene	1447	19.529 \pm 1.255
20	23.347	(+)-Epi- β -Santalene	1453	0.201 \pm 0.014
21	23.627	Humulene	1464	5.636 \pm 0.623
22	24.181	(E)- β -Farnesene	1486	1.014 \pm 0.056
23	24.385	(-)-Zingiberene	1501	0.181 \pm 0.068
24	24.548	α -Muurolene	1508	0.478 \pm 0.077
25	24.758	β -Bisabolene	1517	2.195 \pm 0.079
26	25.108	β -Sesquiphellandrene	1532	1.080 \pm 0.039
27	26.034	(\pm)- <i>trans</i> -Nerolidol	1571	0.033 \pm 0.008
28	26.384	Thymohydroquinone	1586	0.213 \pm 0.056
29	26.576	Caryophyllene oxide	1594	1.419 \pm 0.123
30	27.148	Humulene epoxide II	1615	0.190 \pm 0.022
31	28.797	α -Bisabolol	1688	0.039 \pm 0.010

^a Retention time (RT) in min

^b van den Dool and Kratz retention index calculated for HP-5MS column

^c Relative peak area expressed as percentage of the peak area of corresponding compound over the total peak areas of all identified volatiles.

<https://doi.org/10.1371/journal.pone.0235416.t001>

compounds with potential application as an anti-cancer drug that can be found in *Nigella sativa* were also detected in *P. amboinicus* albeit in small amounts. The presence of thymoquinone in *P. amboinicus* has also been documented by Chen et al. [28] in water-hexane extract with further identification using mass spectrophotometer and NMR analysis.

Isolation of full-length monoterpene synthase gene and sequence characterization

The partial *PamTps1* transcript (Accession No.: GGXS01005129) was identified from the *P. amboinicus* transcriptome (SRA Accession No.:SRR7842030) using the BLAST alignment that showed hits to known terpene synthases and also on the basis of the presence of its conserved sequence characteristics that are shared by all terpene synthases. The full-length *PamTps1*




Fig 1. Total ion chromatogram of *Plectranthus amboinicus* leaf volatiles harvested at 8.00 AM extracted using headspace solid phase microextraction (HS-SPME). The numbers corresponded to the compounds detected as described in [Table 1](#).

<https://doi.org/10.1371/journal.pone.0235416.g001>

transcript (Accession no: MK050501) contained an open reading frame (ORF) of 1797 bp encoding a protein of 598 amino acids with theoretical isoelectric point (pI) and molecular weight of 5.40 and 69.6 kDa, respectively. The transcript was predicted to contain a 52 amino acid N-terminal chloroplast transit peptide, which most likely targets this protein to the plastid, the location of monoterpenes biosynthesis found in other Lamiaceae members. This predicted transit peptide was characterized by high contents of serine and threonine, and low number of acidic residues which in accordance to Williams et al. [29] is expected for the *Tpsb* subfamily.

The sequence identity of *PamTps1* at the amino acid level was compared via BLASTp algorithm against the NCBI non-redundant protein database and showed 60–70% amino acid sequence identity to monoterpene synthases of other Lamiaceae members including *T. caespitius* γ -terpinene synthase (AID51201.1) [14], *S. rosmarinus* pinene synthase (ABP01684.1), *P. frutescens* linalool synthase (AAL38029.1) and *P. setovensis* geraniol synthase (ACN42010.1) [30]. This analysis confirmed that the sequence similarity among Lamiaceae terpene synthases was relatively high but may not necessarily be linked to the catalytic function of the enzyme. Based on the sequence similarity and presence of N-terminal transit peptide, *PamTps1* was designated as a putative monoterpene synthase.

Despite the sequence diversity, terpene synthases share several conserved amino acid residues both at the N- and C-terminal protein domains. These conserved motifs were found in the deduced amino acid sequence of *PamTps1* (Fig 2) notably the tandem arginine motif (RR_xW) located downstream of the transit peptide at the N-terminal region of monoterpene synthases. Deletion of this motif on the *Mentha spicata* limonene synthase was found to affect the ability of the enzyme to utilize geranyl pyrophosphate as a substrate, suggesting that this motif might be involved in the isomerization of geranyl pyrophosphate to a cyclizable intermediate [29]. This arginine pair has also been reported to stabilize the closed active site in the



Fig 2. Alignment of PamTps1 amino acid sequence without transit peptide with other plant monoterpene synthases using Clustal Omega and BoxShade server. Conserved domains of RRx8W, LQLYEASFL, DDxxD, GTLxEL and DTE were labelled. AGK88250.1: *T. caespititius* α -terpineol synthase; AR491313.1: *L. x intermedia* 3-carene synthase; ABD77417.1: *L. latifolia* linalool synthase; AID51195.1: *T. caespititius* γ -terpinene synthase; ADK73617.1: *O. vulgare* terpene synthase 5; AGT29345.1: *T. serpyllum* γ -terpinene synthase; AEO27879.1: *O. syriacum* γ -terpinene synthase; ABP01684.1: *S. rosmarinus* pinene synthase; ACN42010.1: *P. setoyensis* geraniol synthase; ACN42013.2: *P. frutescens* var. *hirtella* linalool synthase and AAX16075.1: *P. citriodora* linalool synthase.

<https://doi.org/10.1371/journal.pone.0235416.g002>

enzyme-ligand complexes reaction [31]. The LQLYEASFL motif which was also found in *PamTps1* but as a LQLYEASFLE sequence is another conserved motif that is believed to be part of the active site and might be involved in substrate binding [15,32]. Another conserved region in terpene synthases is the aspartate-rich DDxxD motif that has been found in almost all isolated plant terpene synthases. This highly conserved motif, which occurs as DDVYD in *P. amboinicus*, is located at the C-terminal domain of all terpene synthases and is known to be involved in the binding of divalent metal ion cofactor such as Mg²⁺ or Mn²⁺ to initiate binding and activation of the diphosphate moiety of the substrate [33–35]. The DDxxD motif located at the entrance of the catalytic site was demonstrated to be critical in positioning the substrate for catalysis as mutation of this motif often led to decreased catalytic activity and abnormal product [36,37]. Besides that, an additional metal cofactor binding motif NSE/DTE which evolved from a second aspartate-rich region to form a consensus sequence of (L,V)(V,L,A)(N,D)D(L,I,V)x(S,T)xxxE occurred as LADDLGTAPFE in *PamTps1*. Both the DDxxD and NSE/DTE motifs bind to a trinuclear magnesium ions cluster and are involved in the fixation of pyrophosphate substrate. *PamTps1* also contains other motifs such as RxR and GTLxEL that are postulated to be part of terpene synthases active site [17,38] and occur as RDR and GTLDEL, located about 35 amino acids upstream and two amino acids downstream of DDxxD, respectively.

Functional expression of *PamTps1*

The N-terminal transit peptide necessary for plastidial targeting of monoterpene synthases has been reported to facilitate formation of inclusion bodies in an *E. coli* expression system. High yield expression of soluble monoterpene synthases can be accomplished by truncation of the transit peptide from the coding region to remove the target sequence [11]. Previous studies on bacterial expression of terpene synthases isolated from other members of Lamiaceae such as *M. spicata* limonene synthase and *T. caespititius* γ -terpinene synthase demonstrated the expression of functional enzymes upon removal of the transit peptide upstream of the double arginine motif [13,29]. Therefore, the signal peptide was removed from *PamTps1* sequence during cloning in order to achieve expression of a soluble and functional putative monoterpene synthase in an *E. coli* system. The complete *PamTps1* ORF was 1797 bp, of which 147 bp corresponding to the putative transit peptide was removed to improve protein solubility during expression. The truncated *PamTps1* was cloned into pET32b(+) expression vector that contained a dual-fusion tag consisting of thioredoxin (TrxA) and histidine (His₆) to give a predicted mass of 79 kDa for the fusion protein. The truncated *PamTps1* was expressed in Rosetta™ 2(DE3) strain which supplied rare tRNAs to cope with the differences of codon usage between *PamTps1* and *E. coli* that may impede translation due to the demands for tRNAs that may be lacking in the bacterial host. Protein expression analysis for the clone harboring the truncated *PamTps1* was compared with the control expression containing the empty pET32b(+) vector. The recombinant protein was then purified using immobilized metal affinity chromatography (IMAC) followed by gel filtration chromatography to yield a pure protein for functional characterization study. The SDS-PAGE and Western blot analyses showed the presence of a corresponding protein band of an estimated size of ~79 kDa (S1 Fig).

Functional characterization of *PamTps1*

Most of the previously described terpene synthases were multi-product enzymes where the identity and relative abundance of the terpene products did not accurately predict the protein function [15,39,40]. Thus, the activity of *PamTps1* was investigated through *in vitro* enzymatic assay and *in vivo* expression in an *E. coli* system. The *PamTps1* enzymatic reaction




Fig 3. GC-MS chromatograms (selected ion, $m/z = 93$) of *PamTps1* products generated both *in vitro* and *in vivo* systems. *PamTps1* enzymatic reaction incubated with (A) GPP or (B) FPP; products generated in *E. coli* harboring (C) empty vector and (D) *PamTps1*; (E) (-)-linalool standard and (F) *cis*- and *trans*-nerolidol standards. *In vitro* *PamTps1* reactions produced exclusively linalool and nerolidol when incubated with GPP and FPP, respectively. In recombinant *E. coli*, *PamTps1* produced both linalool and nerolidol. Only sample peaks higher than the negative control were marked with numbers. Corresponding compounds are: 1 = linalool (retention time = 6.7 min); 2 = *trans*-nerolidol (retention time = 13.0 min) and 3 = *cis*-nerolidol (retention time = 12.6 min).

<https://doi.org/10.1371/journal.pone.0235416.g003>

demonstrated that this protein predominantly catalyzed formation of linalool from GPP confirming that *PamTps1* is a functional monoterpene synthase (Fig 3A). Interestingly, when FPP was provided as a substrate, *PamTps1* was able to synthesize sesquiterpene nerolidol, the C_{15} analogue of linalool (Fig 3B). The identification of linalool and nerolidol produced from *PamTps1* enzymatic reaction was conducted by comparing their retention times and mass spectra with the authentic standard materials and NIST14 library (Fig 3 and S2 Fig). Smaller peaks of linalool and nerolidol were observed in the control reactions without enzyme which could arise as a result of non-enzymatic solvolysis of GPP and FPP in the presence of metal ions (S3 Fig) [41]. However, it was evident from the chromatogram that the significant and higher signal abundances of linalool and nerolidol as compared to the control reactions were the products of *PamTps1* enzymatic reactions, albeit with partial contamination of solvolysed products of the substrates. These enzymatic products of *PamTps1* were also detected in the volatiles composition of *P. amboinicus* although only present at 0.2% of the total volatiles detected (Table 1).

Some terpene synthases may behave differently in *in vivo* as compared with the *in vitro* due to differences in cofactors availability and other biochemical conditions [42,43]. It was previously shown that terpene synthase products can vary depending on the expression host and

the subcellular localization of the protein [42,44,45]. Thus, in order to demonstrate whether multi-substrate activity of *PamTps1* extends to the *in vivo* environment, microbial production of terpenoids in the *E. coli* system was conducted. The terpenoid profile produced in the recombinant *E. coli* host up to 24 h of post-induction was illustrated in Fig 3F. It was observed that the recombinant *E. coli* harboring *PamTps1* was capable of producing both linalool and nerolidol *in vivo* without supplementation of any exogenous GPP or FPP substrates, which correlated with the *in vitro* activity of *PamTps1*. This outcome validated the multi-substrate use of this enzyme in producing linalool and nerolidol in both the *in vivo* and *in vitro* systems. Hence, we designated *PamTps1* as a linalool/nerolidol synthase, and this is the first report describing isolation and functional characterization of such substrate promiscuity of a terpene synthase from *P. amboinicus*. Recent advances and progresses in the characterization of enzymes responsible for terpenoids biosynthesis in plants and bacteria have revealed the existence of multi-substrate terpene synthases capable of synthesizing terpenes of different chain lengths depending on corresponding substrate availability. Pazouki and Niinemets [46] highlighted that there were at least 40 confirmed cases of multi-substrate enzymes among plant terpenoids, suggesting that the substrate promiscuous terpene synthases are prevalent within the plant kingdom.

Based on the functional assay study, it was evident that *PamTps1* is a high-fidelity, multi-substrate enzyme capable of performing both monoterpene and sesquiterpene activities in *in vitro* and *in vivo* systems for the formation of linalool and nerolidol. In *planta*, the formation of linalool and nerolidol requires either two specialized terpene synthases (linalool and nerolidol synthases) or a substrate promiscuous linalool/nerolidol synthase. Linalool synthases are widely identified from both angiosperms and gymnosperms, and are considered a defective monoterpene cyclase that can catalyze the initial ionization and rearrangement of GPP but cannot proceed to the subsequent cyclization steps [47]. It was an interesting observation that most papers reported linalool synthase or nerolidol synthase or linalool/nerolidol synthase showed high-fidelity activity by producing exclusively either linalool or nerolidol instead of multiple compounds as demonstrated by the promiscuous terpene synthases [33,48–52]. The *PamTps1* shared a similarity of less than 36% with other linalool/nerolidol synthase from *V. vinifera* [53,54], *Streptomyces clavuligerus* [55], *A. majus* [48] and *S. lycopersicum*. Sequence comparison of *PamTps1* with plant monoterpene synthases showed that Lamiaceae linalool synthases differ from others by a three-amino acid deletion (i.e. between residues 524–526 in *PamTps1*) at the C-terminal region of the protein (Fig 2) which structurally provided more water access to the active site of linalool synthase and caused premature quenching by water capture [47]. However, no deletion of such amino acids was observed in *PamTps1*, and it was postulated that this premature quenching was due to the less efficient active site closure in the enzyme-ligand complexes which thereby was unable to shield reactive carbocation intermediates from the water molecule [56]. It will be interesting to further explore this *PamTps1* to identify the structural features that confer the multi-substrate function that can be exploited to engineer terpene synthases with high fidelity and specificity for production of terpenoids in a microbial cell factory.

Expressional analysis of *PamTps1*

Volatile terpenoids are often synthesized and emitted from specific plant tissue at a particular time that correlates with the spatio-temporal expression of their terpene synthases, suggesting that the terpenoid biosynthesis is transcriptionally regulated [57]. The leaf and stem tissues of aromatic plants were generally associated with the presence of secretory structures that produced large quantities of volatile terpenoids consisting mostly monoterpenes and

sesquiterpenes [58,59]. Based on our preliminary histomorphology results (unpublished), *P. amboinicus* leaves exhibited high accumulation of essential oils. Since Lamiaceae leaves produce essential oils rich in terpenoids and stems are also frequently used for essential oils extraction, both tissues were selected for *PamTps1* differential expression analysis using quantitative RT-PCR (RT-qPCR). Comparison of volatiles released by *P. amboinicus* leaf and stem tissues during the natural 24 h day/night cycle is displayed in S4 Fig. From this analysis, it was evident that the leaves of this herbal plant emitted a vast array of volatiles throughout the day and the amounts declined towards the night. A similar observation was made in *Lillium* whereby its floral scent emission was significantly influenced by light intensity and temperature [60].

Preferential accumulation of *PamTps1* transcripts was observed in the leaf tissue instead of the stems, and this concurred with the emission of linalool and nerolidol from the leaves (Fig 4). Overall, our findings revealed that the *PamTps1* showed a 42-fold preferential expression in leaves as compared to the stem with a maximal expression in the afternoon. This observation was also noted with the terpene synthases expression in *S. guaranitica* such as linalool synthases 1 and 2, geranyl linalool synthase, selinene synthase and β -caryophyllene synthase that showed high expression levels in the leaf tissue as compared to the stems [61]. Likewise, a *Citrus* terpene synthase exhibited preferential expression in leaves and fruit flavedo which corresponds to the terpenes accumulation and essential oils production by these tissues in the *Citrus* plants [62]. Similarly, germacrene A synthase in *Achillea millefolium* showed high expression levels in the leaves and flowers in contrast to the stem tissue [63]. In contrast, expression of linalool synthase was mostly reported in flower tissues which accounted for the floral scents emission of *Lillium* [64], *A. argute* [65] and *Osmanthus fragrans* [66] that correlated with the volatiles released during flower development. The expression of terpene synthase genes has been reported to be highly up-regulated in specialized cells such as those in glandular trichomes, which were located on the aerial parts of the plants [11,14,67]. A number of Lamiaceae terpene synthases, including linalool synthase, involved in terpenoid biosynthesis in secretory glandular trichomes had been functionally characterized as reviewed by Lange & Turner [68]. In *P. amboinicus*, our preliminary histomorphological results showed that the




Fig 4. Correlation between linalool and nerolidol emissions and *PamTps1* expression in leaves and stems of *P. amboinicus* within a 24 h day/night cycle. Relative expression analysis was performed by qRT-PCR using *EFG*, *TUB* and *APRT* as reference genes. The relative transcription level in tissue with the highest expression quantity was set to 1 (100%). Each bar represents the mean value \pm SE of three biological and three technical replicates. The linalool and nerolidol emission data presented are means \pm SE of duplicate experiments.

<https://doi.org/10.1371/journal.pone.0235416.g004>

glandular trichomes are probably the storage sites for the essential oils produced in the leaves. However, it remains unknown whether *PamTps1* expression is associated with this storage site. The *PamTps1* expression in the leaves followed a diurnal circadian rhythm with increased level exhibited early in the morning at 2.00 AM, and achieved a high point around 2.00 PM before diminishing during the remainder of the day (Fig 4). This expression pattern is similar to *A. chinensis* bifunctional nerolidol synthase (*AcNES1*) expression in the whole flower that showed an increment from 4.00 AM to a maximum point at midday before decreasing afterwards [69]. The *AcNES1* expression accompanied by time point analysis of terpenes suggested that nerolidol was largely accumulated throughout the day with maximal emission at midnight preceded by a steady decrease and remained low until morning. Similar trend was observed in *P. amboinicus* leaf where the highest nerolidol emission occurred early in the morning (2.00 AM) and the amounts reduced thereafter. In contrast, maximal linalool emission from the *P. amboinicus* leaf was observed at 8.00 AM and started to decline in the afternoon till midnight (Fig 4). Based on this pattern, we hypothesized that the accumulation of linalool happened during the night before the morning emission. Similar correlation was observed with Chen et al. [65] study where the *A. argute* linalool synthase (*AaLS1*) displayed constitutive expression with slight reduction in the morning and an increased at midday accompanied by maximal emission rate of linalool at 8.00 AM. This phenomenon could possibly be related to the study previously demonstrated in *Pinus pinea* where large emission of oxygenated monoterpene linalool was controlled by stomata opening, which was influenced by light intensity and temperature [70,71]. This was accompanied by a significant reduction in the emission rate of monoterpenoids during midday which was attributable to diurnal water-stress leaves with closed stomata [70]. It remains uncertain whether the diurnal water-stress leaves with closed stomata could explain the terpenoids emission pattern in *P. amboinicus*.

Phylogenetic analysis of *PamTps1*

Phylogenetic profiling of *PamTps1* protein was conducted to infer its evolutionary relationship with members of the plant terpene synthase subfamilies and other terpene synthases from the Lamiaceae family. Plant terpene synthases share a common evolutionary origin based upon their similar reaction mechanisms and conserved structural and sequence characteristics that include amino acid sequence homology and conserved sequence motifs [9,10,72]. The bifurcation of the terpene synthases involved in primary and secondary metabolism appears to occur before the separation of angiosperms and gymnosperms. Previous phylogenetic analysis of terpene synthases from the gymnosperms and angiosperms delineated the *Tps* gene family into eight subfamilies, designated as *Tpsa* through *Tpsg* and α -farnesene synthase cluster [9,72,73]. Terpene synthases that produce secondary metabolites are classified into subfamilies of *Tpsa* (angiosperm sesquiterpene and diterpene synthases), *Tpsb* (angiosperm monoterpene synthases) and *Tpsd* (gymnosperm monoterpene synthases) are only distantly related to *Tpsc* (copalyl diphosphate synthase) and *Tpse* (kaurene synthase) subfamilies that are rich in specialized (i.e. secondary) metabolism beyond carrying the enzyme involved in gibberellin biosynthesis, and distantly ancient branch *Tpsf* containing linalool synthase [9,11].

Functional characterization of *PamTps1* using *in vitro* and *in vivo* expression systems showed that this multi-substrate enzyme possesses a monoterpene synthase and a sesquiterpene synthase activity producing exclusively linalool and nerolidol, respectively. The phylogenetic tree (Fig 5) reveals that *PamTps1* belongs to the *Tpsb* group as expected for a Lamiaceae monoterpene synthase. The presence of RR_xW motif, which is characteristic of the angiosperm *Tpsb* group [11,73], positioned *PamTps1* in the *Tpsb* cluster. This monophyletic *Tpsb* group contains terpene synthases such as pinene synthase, linalool synthase and limonene




Fig 5. Phylogenetic relationship of *PamTps1* with selected plant terpene synthases from different subfamilies. Target sequences upstream of the RRx₈W motif of the alignment were removed. Selection of terpene synthases subfamilies were based on previous literatures [11,100,101]. The *Tpsc* and *Tpsc* subfamilies were chosen as outgroups. Evolutionary relationship was inferred using Neighbor-Joining method with 1000 replicates for bootstrapping. The numbers indicated were the actual bootstrap values of the branches.

<https://doi.org/10.1371/journal.pone.0235416.g005>

synthase which produce cyclic and acyclic products, which were grouped together on the basis of their sequence similarities and the presence of tandem arginine motif, despite differences in their catalytic functions. It is evident from Fig 5 that the phylogenetically closest sequences to *PamTps1* are *Lavandula* and *Salvia* terpene synthases with more than 67% similarity, signifying a close relationship between these genera that are cluster together in *Tpsb* regardless of the product specificity [67]. This could probably be related to the adaptive evolutionary process of ancestral gene copy that had undergone a divergence in structure and function which contributed to the large diversity of terpene synthases [10]. The other known multi-substrate terpene synthases that belong to the *Tpsb* and have a transit peptide are the *S. lycopersicum* linalool synthase [74], *Hedychium coronarium* terpene synthase [75] and β-ocimene synthase of *Arabidopsis thaliana* [45]. It is noteworthy to mention that most of the reported linalool/nerolidol synthases belong to the *Tpsg* group, a clade closely related to the *Tpsb* which comprises of angiosperm acyclic terpene synthases that produce monoterpenes, sesquiterpenes and diterpenes. However, *Tpsg* group members lack the common structural feature of RRx₈W motif present in the *Tpsb* members [9,54].

The multi-substrate activity of terpene synthases are widely divergent across the *Tps* subfamilies in the multiple plant species. Phylogenetic analysis suggested that all plant terpene synthases originated from the ancient diterpene synthases of the *Tpsc* clade that represents the base of the rooted phylogenetic tree such as illustrated in Fig 5 [9,76,77]. These old diterpene synthases adopt a tri-domain architecture of αβγ proteins that contain a transit peptide [78–80]. Further evolutionary modifications over time have witnessed diversification of product profiles that are not only been associated with changes in the active site center structure, but also loss of γ-domain in the isoprene, monoterpene and sesquiterpene synthases, and transit peptide loss in the sesquiterpene synthases [78–80]. Identification of two plastidic and

cytosolic linalool/nerolidol synthases in *A. majus* and *F. ananassa* further suggests that sesquiterpene synthases might have evolved from the monoterpene synthases ancestor through the loss of plastid signal peptide and acquirement of an active site adaptation to the FPP substrate [46,48,81,82]. The evolution of terpene synthases family shows that these enzymes possess a remarkable flexibility to evolve into new functional diversification and subsequently contribute to the chemical diversity of terpenoids in nature [9].

Conclusions

Initially, *PamTps1* was selected as a terpene synthase candidate among other candidates for functional characterization study based on the highest sequence similarity with available plant terpene synthases in the database. The functional study has undoubtedly classified *PamTps1* as a linalool/nerolidol synthase with the ability to exclusively produce linalool and nerolidol in both *in vivo* and *in vitro* systems, suggesting that this enzyme possesses both monoterpene synthase and sesquiterpene synthase activities. Interestingly, *PamTps1* was clustered with the *Tpsb* subfamily as expected for a predicted monoterpene synthase instead of *Tpsg* or *Tpsf* which was presumed for an acyclic monoterpene synthase or ancient linalool synthase, respectively. Although this study did not clarify the exact role of *PamTps1* in *P. amboinicus*, it did not rule out the possibility that *PamTps1* could function as both a monoterpene synthase and a sesquiterpene synthase *in planta*. The expression analysis showed that this transcript was highly expressed in *P. amboinicus* leaves throughout the day that correlated with its linalool emission following a diurnal circadian pattern. Even though *PamTps1* is only accountable for production of minor volatiles in *P. amboinicus*, it represents the first substrate promiscuous monoterpene synthase that has been cloned and functionally characterized from this herbal plant. The substrate promiscuity activity of *PamTps1* has intrigued us to further study this enzyme for the opportunity to attain additional insight into its catalytic basis of product specificity.

Supporting information

S1 Table. qPCR primers used in this study.

(PDF)

S1 Fig. SDS-PAGE and Western blot profile of purified *PamTps1*. Lane M: Full-Range Rainbow™ Molecular Weight Markers (GE Healthcare, USA). Lanes 1–3: *PamTps1* protein from IMAC fractions and lane 4: *PamTps1* from gel filtration fraction on SDS-PAGE analysis (A) and Western blot analysis (B). The recombinant *PamTps1* band was observed with the expected size of 79 kDa.

(TIF)

S2 Fig. Mass spectra comparison of products generated by recombinant *PamTps1*. Mass spectra of linalool (A) and (B) nerolidol generated by *PamTps1*; mass spectra of authentic (-)-linalool (C) and *trans*-nerolidol (D) standards; mass spectra of linalool (E) and *trans*-nerolidol (F) in the NIST14 library.

(TIF)

S3 Fig. GC-MS chromatograms (selected ion, $m/z = 93$) negative control reactions of *in vitro* assay. Negative control reaction without presence of *PamTps1* (A) and empty vector control reaction (B) after incubation with GPP; negative control reaction without presence of *PamTps1* (C) and empty vector control reaction (D) after incubation with FPP. Corresponding compounds are: 1 = linalool (retention time = 6.7 min); 2 = *trans*-nerolidol (retention

time = 13.0 min) and 3 = *cis*-nerolidol (retention time = 12.6 min).
(TIF)

S4 Fig. Comparison of chromatographic profiles of volatiles emitted from different tissues of *P. amboinicus* within a 24 h day/night cycle. TIC profiles of (A) leaf and (B) stem at 2.00 AM; (C) leaf and (D) stem at 8.00 AM; (E) leaf and (F) stem at 2.00 PM; and (G) leaf and (H) stem at 8.00 PM. Corresponding retention time for linalool and *trans*-nerolidol are 13.7 min and 26.0 min, respectively.
(TIF)

S1 Raw images.
(PDF)

Acknowledgments

We are grateful to Professor Jacqueline Batley from University of Western Australia for her critical comments on our manuscript. We also would like to thank Ms. Teh Kah Yee for helpful discussion on the gene expression study using the qPCR approach.

Author Contributions

Conceptualization: Mohd Hairul Ab. Rahim, Janna Ong Abdullah.

Funding acquisition: Nur Suhanawati Ashaari, Mohd Hairul Ab. Rahim, Suriana Sabri, Kok Song Lai, Adelene Ai-Lian Song, Raha Abdul Rahim, Janna Ong Abdullah.

Investigation: Nur Suhanawati Ashaari, Mohd Hairul Ab. Rahim.

Methodology: Nur Suhanawati Ashaari, Mohd Hairul Ab. Rahim, Wan Muhamad Asrul Nizam Wan Abdullah.

Resources: Nur Suhanawati Ashaari, Mohd Hairul Ab. Rahim, Janna Ong Abdullah.

Supervision: Mohd Hairul Ab. Rahim, Suriana Sabri, Kok Song Lai, Adelene Ai-Lian Song, Janna Ong Abdullah.

Validation: Nur Suhanawati Ashaari, Mohd Hairul Ab. Rahim, Suriana Sabri, Kok Song Lai, Adelene Ai-Lian Song, Janna Ong Abdullah.

Writing – original draft: Nur Suhanawati Ashaari.

Writing – review & editing: Nur Suhanawati Ashaari, Mohd Hairul Ab. Rahim, Suriana Sabri, Kok Song Lai, Adelene Ai-Lian Song, Raha Abdul Rahim, Janna Ong Abdullah.

References

1. Arumugam G, Swamy M, Sinniah U. *Plectranthus amboinicus* (Lour.) Spreng: Botanical, phytochemical, pharmacological and nutritional significance. *Molecules*. 2016 Mar 30; 21(4):369. <https://doi.org/10.3390/molecules21040369> PMID: 27043511
2. Khare RS, Banerjee S, Kundu K. *Coleus aromaticus* benth—A nutritive medicinal plant of potential therapeutic value. *Int J Pharma Bio Sci*. 2011; 2(3):488–500.
3. Damanik R, Wahlqvist M, Wattanapenpaiboon N. Lactagogue effects of *Torbangun*, a Batakese traditional cuisine. *Asia Pac J Clin Nutr*. 2006; 15(2):267–74. PMID: 16672214
4. de Oliveira FFM, Torres AF, Gonçalves TB, Santiago GMP, de Carvalho CBM, Aguiar MB, et al. Efficacy of *Plectranthus amboinicus* (Lour.) Spreng in a murine model of methicillin-resistant *Staphylococcus aureus* skin abscesses. *Evidence-Based Complement Altern Med*. 2013; 2013:1–9. <https://doi.org/10.1155/2013/291592>.

5. Pritima R, Pandian R. Antimicrobial activity of *Coleus aromaticus* (Benth) against microbes of reproductive tract infections among women. *African J Infect Dis*. 2008 Sep 23; 1(1). <https://doi.org/10.4314/ajid.v1i1.42081>.
6. Gurgel APAD, da Silva JG, Grangeiro ARS, Oliveira DC, Lima CMP, da Silva ACP, et al. In vivo study of the anti-inflammatory and antitumor activities of leaves from *Plectranthus amboinicus* (Lour.) Spreng (Lamiaceae). *J Ethnopharmacol*. 2009 Sep; 125(2):361–3. <https://doi.org/10.1016/j.jep.2009.07.006> PMID: 19607901
7. Senthilkumar A, Venkatesalu V. Chemical composition and larvicidal activity of the essential oil of *Plectranthus amboinicus* (Lour.) Spreng against *Anopheles stephensi*: a malarial vector mosquito. *Parasitol Res*. 2010 Oct 29; 107(5):1275–8. <https://doi.org/10.1007/s00436-010-1996-6> PMID: 20668876
8. Bhatt P, Joseph GS, Negi PS, Varadaraj MC. Chemical composition and nutraceutical potential of Indian Borage (*Plectranthus amboinicus*) stem extract. *J Chem*. 2013; 2013:1–7. <https://doi.org/10.1155/2013/320329>.
9. Chen F, Tholl D, Bohlmann J, Pichersky E. The family of terpene synthases in plants: a mid-size family of genes for specialized metabolism that is highly diversified throughout the kingdom. *Plant J*. 2011 Apr; 66(1):212–29. <https://doi.org/10.1111/j.1365-3113X.2011.04520.x> PMID: 21443633
10. Trapp SC, Croteau R. Genomic organization of plant terpene synthases and molecular evolutionary implications. *Genetics*. 2001; 158(2):811–32. PMID: 11404343
11. Bohlmann J, Meyer-Gauen G, Croteau R. Plant terpenoid synthases: Molecular biology and phylogenetic analysis. *Proc Natl Acad Sci*. 1998 Apr 14; 95(8):4126–33. <https://doi.org/10.1073/pnas.95.8.4126> PMID: 9539701
12. Crocoll C, Asbach J, Novak J, Gershenzon J, Degenhardt J. Terpene synthases of oregano (*Origanum vulgare* L.) and their roles in the pathway and regulation of terpene biosynthesis. *Plant Mol Biol*. 2010 Aug 25; 73(6):587–603. <https://doi.org/10.1007/s11103-010-9636-1> PMID: 20419468
13. Lima AS, Schimmel J, Lukas B, Novak J, Barroso JG, Figueiredo AC, et al. Genomic characterization, molecular cloning and expression analysis of two terpene synthases from *Thymus caespititius* (Lamiaceae). *Planta*. 2013 Jul 28; 238(1):191–204. <https://doi.org/10.1007/s00425-013-1884-2> PMID: 23624978
14. Mendes MD, Barroso JG, Oliveira MM, Trindade H. Identification and characterization of a second isogene encoding γ -terpinene synthase in *Thymus caespititius*. *J Plant Physiol*. 2014; 171(12):1017–27. <https://doi.org/10.1016/j.jplph.2014.04.001> PMID: 24974328
15. Galata M, Sarker LS, Mahmoud SS. Transcriptome profiling, and cloning and characterization of the main monoterpene synthases of *Coriandrum sativum* L. *Phytochemistry*. 2014 Jun; 102:64–73. <https://doi.org/10.1016/j.phytochem.2014.02.016> PMID: 24636455
16. Rudolph K, Parthier C, Egerer-Sieber C, Geiger D, Muller YA, Kreis W, et al. Expression, crystallization and structure elucidation of γ -terpinene synthase from *Thymus vulgaris*. *Acta Crystallogr Sect F Struct Biol Commun*. 2016 Jan 1; 72(1):16–23. <https://doi.org/10.1107/S2053230X15023043>.
17. Wise ML, Savage TJ, Katahira E, Croteau R. Monoterpene synthases from common sage (*Salvia officinalis*): cDNA isolation, characterization, and functional expression of (+)-sabinene synthase, 1,8-cineole synthase, and (+)-bornyl diphosphate synthase. *J Biol Chem*. 1998 Jun 12; 273(24):14891–9. <https://doi.org/10.1074/jbc.273.24.14891> PMID: 9614092
18. van Den Dool H, Dec. Kratz P. A generalization of the retention index system including linear temperature programmed gas—liquid partition chromatography. *J Chromatogr A*. 1963 Jan; 11(3):463–71. [https://doi.org/10.1016/S0021-9673\(01\)80947-X](https://doi.org/10.1016/S0021-9673(01)80947-X).
19. Emanuelsson O, Nielsen H, Heijne G Von. ChloroP, a neural network-based method for predicting chloroplast transit peptides and their cleavage sites. *Protein Sci*. 1999; 8(5):978–84. <https://doi.org/10.1110/ps.8.5.978> PMID: 10338008
20. Emanuelsson O, Brunak S, von Heijne G, Nielsen H. Locating proteins in the cell using TargetP, SignalP and related tools. *Nat Protoc*. 2007 Apr 19; 2(4):953–71. <https://doi.org/10.1038/nprot.2007.131> PMID: 17446895
21. Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, et al. The EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic Acids Res*. 2019 Jul 2; 47(W1):W636–41. <https://doi.org/10.1093/nar/gkz268> PMID: 30976793
22. Felsenstein JP. PHYLIP (phylogeny inference package), Version 3.6. Vol. 5, *Cladistics*. 2004.
23. Khan M, Khan ST, Khan NA, Mahmood A, Al-Kedhairy AA, Alkhatlan HZ. The composition of the essential oil and aqueous distillate of *Origanum vulgare* L. growing in Saudi Arabia and evaluation of their antibacterial activity. *Arab J Chem*. 2018 Feb 27; <https://doi.org/10.1016/J.ARABJC.2018.02.008>.

24. Rusdi NA, Goh H-H, Baharum SN. GC-MS/Olfactometric characterisation and aroma extraction dilution analysis of aroma active compounds in *Polygonum minus* essential oil. *Plant Omics*. 2016 Aug 20; 9(4):289–91. <https://doi.org/10.21475/poj.16.09.04.p7901>.
25. El-Zaedi H, Martínez-Tomé J, Calín-Sánchez Á, Burló F, Carbonell-Barrachina Á. Volatile composition of essential oils from different aromatic herbs grown in Mediterranean regions of Spain. *Foods*. 2016 May 25; 5(2):41. <https://doi.org/10.3390/foods5020041>.
26. Asimwe S, Karlsson AB, Azeem M, Mugisha KM, Namutebi A, Gakunga NJ. Chemical composition and toxicological evaluation of the aqueous leaf extracts of *Plectranthus amboinicus* Lour. *Spreng. Int J Pharm Sci Invent*. 2014; 3(2):19–27.
27. Koba K, Garde D, Sanda K, Delgado Raynaud C, Chaumont J-P. Chemical composition and antimicrobial properties of the leaf essential oil of *Coleus aromaticus* Benth. from Cambodia. *Int J Essent Oil Ther*. 2007 Jan 1; 1:16–20.
28. Chen Y-S, Yu H-M, Shie J-J, Cheng T-JR, Wu C-Y, Fang J-M, et al. Chemical constituents of *Plectranthus amboinicus* and the synthetic analogs possessing anti-inflammatory activity. *Bioorg Med Chem*. 2014; 22(5):1766–72. <https://doi.org/10.1016/j.bmc.2014.01.009> PMID: 24491635
29. Williams DC, McGarvey DJ, Katahira EJ, Croteau R. Truncation of limonene synthase preprotein provides a fully active “pseudomature” form of this monoterpene cyclase and reveals the function of the amino-terminal arginine pair. *Biochemistry*. 1998 Sep; 37(35):12213–20. <https://doi.org/10.1021/bi980854k> PMID: 9724535
30. Masumoto N, Korin M, Ito M. Geraniol and linalool synthases from wild species of perilla. *Phytochemistry*. 2010; 71(10):1068–75. <https://doi.org/10.1016/j.phytochem.2010.04.006> PMID: 20447664
31. Hyatt DC, Youn B, Zhao Y, Santhamma B, Coates RM, Croteau RB, et al. Structure of limonene synthase, a simple model for terpenoid cyclase catalysis. *Proc Natl Acad Sci*. 2007 Mar 27; 104(13):5360–5. <https://doi.org/10.1073/pnas.0700915104> PMID: 17372193
32. Mcgeady P, Croteau R. Isolation and characterization of an active-site peptide from a monoterpene cyclase labeled with a mechanism-based inhibitor. *Arch Biochem Biophys*. 1995 Feb; 317(1):149–55. <https://doi.org/10.1006/abbi.1995.1147> PMID: 7872777
33. Karuppiyah V, Ranaghan KE, Leferink NGH, Johannissen LO, Shanmugam M, Ní Cheallaigh A, et al. Structural basis of catalysis in the bacterial monoterpene synthases linalool synthase and 1,8-cineole synthase. *ACS Catal*. 2017 Sep 22; 7(9):6268–82. <https://doi.org/10.1021/acscatal.7b01924> PMID: 28966840
34. Kampranis SC, Ioannidis D, Purvis A, Mahrez W, Ninga E, Katerelos NA, et al. Rational conversion of substrate and product specificity in a *Salvia* monoterpene synthase: structural insights into the evolution of terpene synthase function. *Plant Cell*. 2007; 19(6):1994–2005. <https://doi.org/10.1105/tpc.106.047779> PMID: 17557809
35. Whittington DA, Wise ML, Urbansky M, Coates RM, Croteau RB, Christianson DW. Nonlinear partial differential equations and applications: Bornyl diphosphate synthase: Structure and strategy for carbocation manipulation by a terpenoid cyclase. *Proc Natl Acad Sci*. 2002 Nov 26; 99(24):15375–80. <https://doi.org/10.1073/pnas.232591099> PMID: 12432096
36. Prosser I, Phillips AL, Gittings S, Lewis MJ, Hooper AM, Pickett JA, et al. (+)-(10R)-Germacrene A synthase from goldenrod, *Solidago canadensis*; cDNA isolation, bacterial expression and functional analysis. *Phytochemistry*. 2002 Aug; 60(7):691–702. [https://doi.org/10.1016/s0031-9422\(02\)00165-6](https://doi.org/10.1016/s0031-9422(02)00165-6) PMID: 12127586
37. Cane DE, Xue Q, Fitzsimons BC. Trichodiene synthase. Probing the role of the highly conserved aspartate-rich region by site-directed mutagenesis. *Biochemistry*. 1996 Jan; 35(38):12369–76. <https://doi.org/10.1021/bi961344y> PMID: 8823172
38. Li R, Fan Y. Molecular cloning and expression analysis of a terpene synthase gene, HcTPS2, in *Hedychium coronarium*. *Plant Mol Biol Report*. 2011 Mar 16; 29(1):35–42. <https://doi.org/10.1007/s11105-010-0205-1>.
39. Rusdi N, Goh H-H, Sabri S, Ramzi A, Mohd Noor N, Baharum S. Functional characterisation of new sesquiterpene synthase from the Malaysian herbal plant, *Polygonum minus*. *Molecules*. 2018 Jun 6; 23(6):1370. <https://doi.org/10.3390/molecules23061370>.
40. Despinasse Y, Fiorucci S, Antonczak S, Moja S, Bony A, Nicolè F, et al. Bornyl-diphosphate synthase from *Lavandula angustifolia*: A major monoterpene synthase involved in essential oil quality. *Phytochemistry*. 2017 May; 137:24–33. <https://doi.org/10.1016/j.phytochem.2017.01.015> PMID: 28190677
41. Cori O, Chayet L, De La Fuente M, Fernandez LA, Hashagen U, Perez L, et al. Stereochemical aspects of chain lengthening and cyclization processes in terpenoid biosynthesis. In: Chapeville F, Haenni A., editors. *Chemical Recognition in Biology Molecular Biology, Biochemistry and Biophysics*. Springer, Berlin, Heidelberg; 1980. p. 97–110. https://doi.org/10.1007/978-3-642-81503-4_8.

42. Fischer MJC, Meyer S, Claudel P, Perrin M, Ginglinger JF, Gertz C, et al. Specificity of *Ocimum basilicum* geraniol synthase modified by its expression in different heterologous systems. *J Biotechnol*. 2013 Jan; 163(1):24–9. <https://doi.org/10.1016/j.jbiotec.2012.10.012> PMID: 23108028
43. Su P, Hu T, Liu Y, Tong Y, Guan H, Zhang Y, et al. Functional characterization of NES and GES responsible for the biosynthesis of (E)-nerolidol and (E,E)-geranylinalool in *Tripterygium wilfordii*. *Sci Rep*. 2017 Feb 27; 7(1):40851. <https://doi.org/10.1038/srep40851>.
44. Ginglinger J-F, Boachon B, Hofer R, Paetz C, Kollner TG, Miesch L, et al. Gene coexpression analysis reveals complex metabolism of the monoterpene alcohol linalool in *Arabidopsis* flowers. *Plant Cell*. 2013 Nov 1; 25(11):4640–57. <https://doi.org/10.1105/tpc.113.117382> PMID: 24285789
45. Huang M, Abel C, Sohrabi R, Petri J, Haupt I, Cosimano J, et al. Variation of herbivore-induced volatile terpenes among *Arabidopsis* ecotypes depends on allelic differences and subcellular targeting of two terpene synthases, TPS02 and TPS03. *Plant Physiol*. 2010 Jul 1; 153(3):1293–310. <https://doi.org/10.1104/pp.110.154864> PMID: 20463089
46. Pazouki L, Niinemets Ü. Multi-substrate terpene synthases: their occurrence and physiological significance. *Front Plant Sci*. 2016; 7(July):1019. <https://doi.org/10.3389/fpls.2016.01019>.
47. Crowell AL, Williams DC, Davis EM, Wildung MR, Croteau R. Molecular cloning and characterization of a new linalool synthase. *Arch Biochem Biophys*. 2002 Sep; 405(1):112–21. [https://doi.org/10.1016/S0003-9861\(02\)00348-X](https://doi.org/10.1016/S0003-9861(02)00348-X) PMID: 12176064
48. Nagegowda DA, Gutensohn M, Wilkerson CG, Dudareva N. Two nearly identical terpene synthases catalyze the formation of nerolidol and linalool in snapdragon flowers. *Plant J*. 2008 Jul; 55(2):224–39. <https://doi.org/10.1111/j.1365-3113X.2008.03496.x> PMID: 18363779
49. Arimura G, Garms S, Maffei M, Bossi S, Schulze B, Leitner M, et al. Herbivore-induced terpenoid emission in *Medicago truncatula*: concerted action of jasmonate, ethylene and calcium signaling. *Planta*. 2007 Nov 30; 227(2):453–64. <https://doi.org/10.1007/s00425-007-0631-y> PMID: 17924138
50. Nieuwenhuizen NJ, Green SA, Chen X, Bailleul EJD, Matich AJ, Wang MY, et al. Functional genomics reveals a compact terpene synthase gene family can account for terpene volatile production in apple (*Malus domestica*). *Plant Physiol*. 2013 Feb 1; 161(2):787–804. <https://doi.org/10.1104/pp.112.208249> PMID: 23256150
51. Gao F, Liu B, Li M, Gao X, Fang Q, Liu C, et al. Identification and characterization of terpene synthase genes accounting for volatile terpene emissions in flowers of *Freesia x hybrida*. *J Exp Bot*. 2018 Aug 14; 69(18):4249–65. <https://doi.org/10.1093/jxb/ery224> PMID: 29901784
52. Degenhardt J, Köllner TG, Gershenzon J. Monoterpene and sesquiterpene synthases and the origin of terpene skeletal diversity in plants. *Phytochemistry*. 2009; 70(15–16):1621–37. <https://doi.org/10.1016/j.phytochem.2009.07.030> PMID: 19793600
53. Zhu B-Q, Cai J, Wang Z-Q, Xu X-Q, Duan C-Q, Pan Q-H. Identification of a plastid-localized bifunctional nerolidol/linalool synthase in relation to linalool biosynthesis in young grape berries. *Int J Mol Sci*. 2014 Dec 1; 15(12):21992–2010. <https://doi.org/10.3390/ijms151221992> PMID: 25470020
54. Martin DM, Aubourg S, Schouwey MB, Daviet L, Schalk M, Toub O, et al. Functional annotation, genome organization and phylogeny of the grapevine (*Vitis vinifera*) terpene synthase gene family based on genome assembly, FLcDNA Cloning, and enzyme assays. *BMC Plant Biol*. 2010; 10(1):226. <https://doi.org/10.1186/1471-2229-10-226>.
55. Nakano C, Kim H-K, Ohnishi Y. Identification and characterization of the linalool/nerolidol synthase from *Streptomyces clavuligerus*. *ChemBioChem*. 2011 Nov 4; 12(16):2403–7. <https://doi.org/10.1002/cbic.201100501> PMID: 21910204
56. González V, Grundy DJ, Faraldos JA, Allemann RK. The amino-terminal segment in the β -domain of δ -cadinene synthase is essential for catalysis. *Org Biomol Chem*. 2016; 14(31):7451–4. <https://doi.org/10.1039/c6ob01398h> PMID: 27431578
57. Nagegowda DA. Plant volatile terpenoid metabolism: Biosynthetic genes, transcriptional regulation and subcellular compartmentation. *FEBS Lett*. 2010 Jul 16; 584(14):2965–73. <https://doi.org/10.1016/j.febslet.2010.05.045> PMID: 20553718
58. Schillmiller AL, Schauvinhold I, Larson M, Xu R, Charbonneau AL, Schmidt A, et al. Monoterpenes in the glandular trichomes of tomato are synthesized from a neryl diphosphate precursor rather than geranyl diphosphate. *Proc Natl Acad Sci U S A*. 2009; 106(26):10865–70. <https://doi.org/10.1073/pnas.0904113106> PMID: 19487664
59. Rehman R, Hanif MA, Mushtaq Z, Al-Sadi AM. Biosynthesis of essential oils in aromatic plants: A review. *Food Rev Int*. 2016 Apr 2; 32(2):117–60. <https://doi.org/10.1080/87559129.2015.1057841>.
60. Hu Z, Zhang H, Leng P, Zhao J, Wang W, Wang S. The emission of floral scent from *Lilium 'siberia'* in response to light intensity and temperature. *Acta Physiol Plant*. 2013 May 5; 35(5):1691–700. <https://doi.org/10.1007/s11738-012-1211-8>.

61. Ali M, Hussain RM, Rehman NU, She G, Li P, Wan X, et al. De novo transcriptome sequencing and metabolite profiling analyses reveal the complex metabolic genes involved in the terpenoid biosynthesis in Blue Anise Sage (*Salvia guaranitica* L.). *DNA Res.* 2018 Dec 1; 25(6):597–617. <https://doi.org/10.1093/dnares/dsy028> PMID: 30188980
62. Dornelas MC, Mazzafera P. A genomic approach to characterization of the Citrus terpene synthase gene family. *Genet Mol Biol.* 2007; 30(3 suppl):832–40. <https://doi.org/10.1590/S1415-47572007000500011>.
63. Pazouki L, Memari HR, Kännaste A, Bichele R, Niinemets Ä. Germacrene A synthase in yarrow (*Achillea millefolium*) is an enzyme with mixed substrate specificity: gene cloning, functional characterization and expression analysis. *Front Plant Sci.* 2015 Mar 3; 6(March):1–16. <https://doi.org/10.3389/fpls.2015.00111>.
64. Abbas F, Ke Y, Yu R, Fan Y. Functional characterization and expression analysis of two terpene synthases involved in floral scent formation in *Lilium* 'Siberia.' *Planta.* 2019 Jan 14; 249(1):71–93. <https://doi.org/10.1007/s00425-018-3006-7>.
65. Chen X, Yauk Y, Nieuwenhuizen NJ, Matich AJ, Wang MY, Perez RL, et al. Characterisation of an (S)-linalool synthase from kiwifruit (*Actinidia arguta*) that catalyses the first committed step in the production of floral lilac compounds. *Funct Plant Biol.* 2010; 37:232–43.
66. Zheng R, Liu C, Wang Y, Luo J, Zeng X, Ding H, et al. Expression of MEP pathway genes and non-volatile sequestration are associated with circadian rhythm of dominant terpenoids emission in *Osmanthus fragrans* Lour. flowers. *Front Plant Sci.* 2017 Oct 30; 8(October):1–10. <https://doi.org/10.3389/fpls.2017.01869>.
67. Tholl D. Terpene synthases and the regulation, diversity and biological roles of terpene metabolism. *Curr Opin Plant Biol.* 2006 Jun; 9(3):297–304. <https://doi.org/10.1016/j.pbi.2006.03.014> PMID: 16600670
68. Markus Lange B, Turner GW. Terpenoid biosynthesis in trichomes-current status and future opportunities. *Plant Biotechnol J.* 2013 Jan; 11(1):2–22. <https://doi.org/10.1111/j.1467-7652.2012.00737.x> PMID: 22979959
69. Green SA, Chen X, Nieuwenhuizen NJ, Matich AJ, Wang MY, Bunn BJ, et al. Identification, functional characterization, and regulation of the enzyme responsible for floral (E)-nerolidol biosynthesis in kiwifruit (*Actinidia chinensis*). *J Exp Bot.* 2012 Mar; 63(5):1951–67. <https://doi.org/10.1093/jxb/err393> PMID: 22162874
70. Niinemets U. Stomatal constraints may affect emission of oxygenated monoterpenoids from the foliage of *Pinus pinea*. *Plant Physiol.* 2002 Nov 1; 130(3):1371–85. <https://doi.org/10.1104/pp.009670> PMID: 12428002
71. Staudt M, Bertin N, Hansen U, Seufert G, Cicciolij P, Foster P, et al. Seasonal and diurnal patterns of monoterpene emissions from *Pinus pinea* (L.) under field conditions. *Atmos Environ.* 1997 Jan; 31 (SUPPL. 1):145–56. [https://doi.org/10.1016/S1352-2310\(97\)00081-2](https://doi.org/10.1016/S1352-2310(97)00081-2).
72. Bohlmann J, Crock J, Jetter R, Croteau R. Terpenoid-based defenses in conifers: cDNA cloning, characterization, and functional expression of wound-inducible (E)- α -bisabolene synthase from grand fir (*Abies grandis*). *Proc Natl Acad Sci.* 1998 Jun 9; 95(12):6756–61. <https://doi.org/10.1073/pnas.95.12.6756> PMID: 9618485
73. Dudareva N, Martin DM, Kish CM, Kolosova N, Gorenstein N, Fäldt J, et al. (E)- β -ocimene and myrcene synthase genes of floral scent biosynthesis in snapdragon: function and expression of three terpene synthase genes of a new terpene synthase subfamily. *Plant Cell.* 2003 May 1; 15(5):1227–41. <https://doi.org/10.1105/tpc.011015> PMID: 12724546
74. Falara V, Akhtar TA, Nguyen TTH, Spyropoulou EA, Bleeker PM, Schauvinhold I, et al. The tomato terpene synthase gene family. *Plant Physiol.* 2011; 157(2):770–89. <https://doi.org/10.1104/pp.111.179648> PMID: 21813655
75. Yue Y, Yu R, Fan Y. Characterization of two monoterpene synthases involved in floral scent formation in *Hedychium coronarium*. *Planta.* 2014; 240(4):745–62. <https://doi.org/10.1007/s00425-014-2127-x> PMID: 25056927
76. Keeling CI, Dullat HK, Yuen M, Ralph SG, Jancsik S, Bohlmann J. Identification and functional characterization of monofunctional *ent*-copalyl diphosphate and *ent*-kaurene synthases in White Spruce reveal different patterns for diterpene synthase evolution for primary and secondary metabolism in gymnosperm. *Plant Physiol.* 2010 Mar; 152(3):1197–208. <https://doi.org/10.1104/pp.109.151456> PMID: 20044448
77. Hayashi K, Kawaide H, Notomi M, Sakigi Y, Matsuo A, Nozaki H. Identification and functional analysis of bifunctional *ent*-kaurene synthase from the moss *Physcomitrella patens*. *FEBS Lett.* 2006 Nov 13; 580(26):6175–81. <https://doi.org/10.1016/j.febslet.2006.10.018> PMID: 17064690

78. Hillwig ML, Xu M, Toyomasu T, Tiernan MS, Wei G, Cui G, et al. Domain loss has independently occurred multiple times in plant terpene synthase evolution. *Plant J*. 2011 Dec; 68(6):1051–60. <https://doi.org/10.1111/j.1365-313X.2011.04756.x> PMID: 21999670
79. Köksal M, Jin Y, Coates RM, Croteau R, Christianson DW. Taxadiene synthase structure and evolution of modular architecture in terpene biosynthesis. *Nature*. 2011 Jan 6; 469(7328):116–20. <https://doi.org/10.1038/nature09628> PMID: 21160477
80. Memari HR, Pazouki L. Biology, controls and models of tree volatile organic compound emissions [Internet]. Vol. 5. 2013. 47–93 p. <https://doi.org/10.1007/978-94-007-6606-8>.
81. Jones CG, Moniodis J, Zulak KG, Scaffidi A, Plummer JA, Ghisalberti EL, et al. Sandalwood fragrance biosynthesis involves sesquiterpene synthases of both the terpene synthase (TPS)-a and TPS-b sub-families, including santalene synthases. *J Biol Chem*. 2011 May 20; 286(20):17445–54. <https://doi.org/10.1074/jbc.M111.231787> PMID: 21454632
82. Aharoni A, Giri AP, Verstappen FWA, Berteaux CM, Sevenier R, Sun Z, et al. Gain and loss of fruit flavor compounds produced by wild and cultivated strawberry species. *Plant Cell*. 2004 Nov 1; 16(11):3110–31. <https://doi.org/10.1105/tpc.104.023895> PMID: 15522848