

Functional genetic polymorphisms and female reproductive disorders: Part I: polycystic ovary syndrome and ovarian response

M. Simoni^{1,5}, C.B. Tempfer², B. Destenaves³ and B.C.J.M. Fauser⁴

¹Department of Medicine, Endocrinology, Metabolism and Geriatrics, University of Modena and Reggio Emilia, I-41100 Modena, Italy;

²Department of Obstetrics and Gynecology, Medical University, 1090 Vienna, Austria; ³Stratified Medicine Group, Merck Serono International S.A., 1202 Geneva, Switzerland; ⁴Department of Reproductive Medicine and Gynecology, University Medical Center, 3508 GA Utrecht, The Netherlands

⁵Correspondence address. Tel: +39-059-4224563; Fax: +39-059-4222528; E-mail: manuela.simoni@unimore.it

BACKGROUND: The identification of polymorphisms associated with a disease can help to elucidate its pathogenesis, and this knowledge can be used to improve prognosis for women with a particular disorder, such as polycystic ovary syndrome (PCOS). Since an altered response to ovarian stimulation is also a characteristic of the disease, further knowledge about its aetiology could help in defining the parameters that determine the response of an individual to ovarian stimulation. **METHODS:** PubMed and EMBASE databases were systematically searched for gene association studies published until the end of August 2007, using search criteria relevant to PCOS and ovarian response to stimulation. Data from additional papers identified through hand searches were also included; 139 publications were reviewed. **RESULTS:** Several genes involved in ovarian function and metabolism are associated with increased susceptibility to PCOS, but none is strong enough to correlate alone with susceptibility to the disease, or response to therapy. A single-nucleotide polymorphism in exon 10 of the FSH receptor (FSHR) gene, *FSHR* p.N680S, was consistently identified as having a significant association with ovarian response to FSH. **CONCLUSIONS:** No consistent association between gene polymorphism and PCOS could be identified. The *FSHR* gene may play a significant role in the success of ovarian stimulation, and can be used as a marker to predict differences in FSHR function and ovarian response to FSH. Genotyping the *FSHR* p.N680S polymorphism may provide a means of identifying a population of poor responders before *in vitro* fertilization procedures are initiated.

Keywords: female reproduction; genetic polymorphisms; polycystic ovary syndrome

Introduction

Genetic studies contribute towards our understanding of disease pathogenesis and hold the promise of improving our ability to individualize treatment for patients. In this review, we first explain the rationale and methodology of studies that identify gene–disease associations. We then review the extensive body of research aimed at identifying contributing genetic factors in polycystic ovary syndrome (PCOS) and assess genetic studies that have investigated patients' ovarian response to gonadotrophin treatment using assisted reproductive techniques. This provides the first systematically selected summary of published studies about all genes

that have so far been investigated in these areas. We then discuss the limitations of such studies and provide recommendations for using this information in the future, with the aim of guiding the design and interpretation of future studies.

Genetic association studies

Genetic association studies are used to investigate genes or genetic markers that might be associated with a particular disease phenotype or trait. These studies rely on the identification and characterization of natural variants or polymorphisms in the DNA sequence among individuals. If an association is present, a particular variant

© The Author 2008. Published by Oxford University Press on behalf of the European Society of Human Reproduction and Embryology. All rights reserved.

For Permissions, please email: journals.permissions@oxfordjournals.org

The online version of this article has been published under an open access model. Users are entitled to use, reproduce, disseminate, or display the open access version of this article for non-commercial purposes provided that: the original authorship is properly and fully attributed; the Journal and Oxford University Press are attributed as the original place of publication with the correct citation details given; if an article is subsequently reproduced or disseminated not in its entirety but only in part or as a derivative work this must be clearly indicated. For commercial re-use, please contact journals.permissions@oxfordjournals.org 459




Figure 1: Graphic representation of types of genetic variants, showing insertion/deletion (ins/del) polymorphisms, both coding and non-coding SNPs, and repeat polymorphisms such as tandem repeats or VNTR. Variants are shown occurring within a gene (in this example the *INS* gene), but can also occur outside of genes. Other types of genetic variations that affect larger regions, such as copy number variations, are not shown. SNP, single-nucleotide polymorphism; VNTR, variable number of tandem repeats.

will be seen more often than expected by chance in an individual carrying the trait. Variants fall into several categories, depending on where they occur relative to a gene. First, variants can occur within or outside a gene. Those that occur within a gene may occur within an exon, an intron or regulatory regions. Regulatory regions control gene transcription and include the promoter region upstream of the gene and the 3'-untranslated region downstream of the protein-coding region (Fig. 1). Variants within a gene can be functional or silent. Functional variation within a gene can be the direct cause of a phenotype abnormality or may increase susceptibility to a disease. Functional variants in coding regions can change the protein sequence; in non-coding regions they may have effects on RNA transcription and processing (Kim *et al.*, 2005; Wang *et al.*, 2005). Silent variants occur in protein-coding DNA but do not change the sequence of a gene product. In the majority of cases, variation that occurs outside a gene is used as

a tag to identify nearby functional variation within a gene (Fig. 2). Such analyses are based on a chromosomal property called linkage disequilibrium. Linkage disequilibrium refers to the observation that in the general population, two DNA variants that are located close to each other tend to be observed together more frequently than two variants that are located further apart. Variants may be single-base changes, known as single-nucleotide polymorphisms (SNPs), or they may occur as insertions or deletions of one base or more.

Until recently, the high cost of testing for genetic variation has meant that most analyses have concentrated on the study of a limited number of functional genetic variants, primarily SNPs, in specific genes. Candidate genes for genotyping are selected according to their function: they encode proteins that are thought to have a role in the disease or response to treatment. Variants within the candidate genes are most often selected because they occur within exons and would result in a change in amino acid sequence in the protein. Alternatively, they are located in non-coding regions, but change a transcription factor-binding site or influence splicing efficiency, affecting the expression of a protein. The main advantage of the candidate SNP approach is that such studies are affordable, as only a limited number of variants are studied and a relatively small sample size can be used. In the context of genome-wide analyses, the main advantages of using SNPs are their abundance in the genome, and the possibility of conducting genotyping in a high-throughput manner. To date, a wealth of results has been obtained from studies addressing the problem of potential associations between genetic polymorphisms and PCOS or ovarian response to gonadotrophins.




Figure 2: Principles of genetic association, and possible explanations for an observed association.

Materials and Methods

We systematically searched the PubMed and EMBASE databases for gene association studies published until the end of August 2007, using the terms ‘PCOS’, ‘polycystic and (ovary or ovaries)’, ‘ovarian and response’, ‘OHSS’ or ‘ovarian and hyperstimulation’, combined with ‘polymorphism or polymorphisms’ or ‘mutation or mutations’. The search was not limited by language of publication. Two authors (B.C.J.M.F. and M.S.) then selected relevant studies using the

following criteria: more than one patient, inclusion of a control group and with at least the abstract written in English. Also included were additional papers identified through hand searches carried out by the same authors.

All results of selected studies are comprehensively summarized in functional group-specific tables by gene of interest, with a brief description in the text. As the frequency of genetic differences varies between ethnic or geographic populations, each study is based on a specific patient population to minimize heterogeneity. Therefore, the tables also include critical details that give information about the context (ethnic background) and likely strength (based on sample size) of the study, to provide a source of reference.

Results

Polycystic ovary syndrome

PCOS affects about 1 in 10 women of reproductive age, and is the most common endocrine condition in this group. The syndrome is associated with multiple endocrinological and metabolic abnormalities, with hyperandrogenaemia and anovulation as the central hallmarks (The Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group, 2004a, b). About 80% of women with PCOS have abnormal menstrual cycles (Conway *et al.*, 1989), many are anovulatory (Laven *et al.*, 2002) and, in those who manage to conceive successfully after treatment, there is an increased risk of complications during pregnancy as well as neonatal complications (Boomsma *et al.*, 2006).

Twin and familial studies suggest that there is a genetic component to PCOS, with a polygenic pattern of inheritance (Jahanfar and Eden, 1996; Diamanti-Kandarakis and Piperi, 2005; Fratantonio *et al.*, 2005). There is also a strong environmental component (e.g. diet and exercise) (Abbott *et al.*, 2002). A series of familial studies conducted in the USA have demonstrated a strong linkage between susceptibility to PCOS and the dinucleotide marker D19S884 in the chromosomal region 19p13.2 (Urbanek *et al.*, 2000a, 2005, 2003; Stewart *et al.*, 2006). The identity of the gene or genetic element responsible for this effect is not yet known. As complex traits such as PCOS result from the interaction of several genetic variations with environmental factors, individual genetic variations have a modest effect that is more difficult to detect than for monogenic traits.

The search for PCOS susceptibility genes has focused mainly on genes involved in sex hormone regulation, insulin sensitivity, cardiovascular risk or steroid biosynthesis and detoxification (Ehrmann, 2005; Escobar-Morreale *et al.*, 2005; Fratantonio *et al.*, 2005). The results from these studies are summarized in Tables I–IV and are described below.

Sex hormones and hormone regulators

Because PCOS is characterized by endocrinological abnormalities, polymorphisms in genes encoding sex hormones or regulators of their activity have been investigated (Table I).

Hormone receptors. Two variants in the gene encoding the $\beta 2$ adrenergic receptor (*ADRB2*) were studied in Japanese women: the p.Q27E variant was linked to PCOS susceptibility, but the p.R16G variant was not (Kurabayashi *et al.*, 2006). Within the $\beta 3$ adrenergic receptor (*ADRB3*) gene, the p.W64R polymorphism does not appear to confer an increased risk of PCOS (Perez-Bravo

et al., 2005; Kurabayashi *et al.*, 2006), although it may affect triglyceride regulation in women with this condition (Perez-Bravo *et al.*, 2005). Conflicting findings have been reported from studies investigating the association between the length of the CAG repeat microsatellite in the androgen receptor (*AR*) gene and PCOS susceptibility (Hickey *et al.*, 2002; Jaaskelainen *et al.*, 2005); however, short alleles of this microsatellite appear to be associated with increased androgenic activity (Mifsud *et al.*, 2000), and this may lead to PCOS. One study has shown that an AccI restriction fragment length polymorphism (RFLP) in the gene that encodes the beta subunit of the gonadotrophin follicle-stimulating hormone (*FSHB*) may be associated with PCOS susceptibility (Tong *et al.*, 2000). A larger number of studies have investigated polymorphisms in the gene that encodes the receptor for FSH rather than the gene for the hormone itself. The *FSHR* gene contains two important SNPs in exon 10, which are in linkage disequilibrium and change two amino acids at positions 307 and 680. However, although the p.N680S genotype influences ovarian response (see section on ovarian response to gonadotrophins), this common polymorphism does not seem to play a major role in PCOS. In Japanese women with PCOS, the S allele was more prevalent than in normal subjects (Sudo *et al.*, 2002), but no differential allelic distribution was found in Caucasian (Conway *et al.*, 1999) or Chinese women (Tong *et al.*, 2001).

Sex hormone-binding globulin. Women with PCOS have subnormal levels of SHBG, and a microsatellite repeat in the promoter of the *SHBG* gene has been linked to PCOS susceptibility in Greek (Xita *et al.*, 2003) but not in Slovenian (Ferk *et al.*, 2007) women, although in both populations, serum SHBG levels were strongly influenced by the genotype. The p.D327N polymorphism in this gene was not found to influence PCOS susceptibility in studies in Slovenia or the Czech Republic (Bendlova *et al.*, 2007; Ferk *et al.*, 2007).

Regulators of hormone activity. Studies of the following genes that also encode hormones or proteins regulating hormone activity have failed to show an association with PCOS susceptibility: Dopamine D3 receptor (*DRD3*) (Kahsar-Miller *et al.*, 1999), follistatin (*FST*) (Urbanek *et al.*, 2000b; Calvo *et al.*, 2001; Jones *et al.*, 2007), guanine nucleotide-binding protein G (s) subunit alpha (*GNAS*) (Hahn *et al.*, 2006), leptin receptor (*LEPR*) (Oksanen *et al.*, 2000) and luteinizing hormone (*LHB*) (Rajkhowa *et al.*, 1995; Elter *et al.*, 1999). However, there was an association between phenotype and two of the SNPs investigated in the *FST* gene (Jones *et al.*, 2007). Furthermore, although not associated with susceptibility, the genotype distribution of the signal transduction protein gene *GNAS* was found to be associated with body mass index (BMI) and insulin resistance in women with PCOS (Hahn *et al.*, 2006; Jones *et al.*, 2007).

Steroid metabolism and biosynthesis

Several studies have investigated whether polymorphisms in enzymes involved in the biosynthesis and metabolism of sex steroids confer PCOS susceptibility (Table II).

Sex steroid synthesis. Type 5 17β -hydroxysteroid dehydrogenase is an important enzyme for testosterone biosynthesis, and a polymorphism in the gene encoding this enzyme (*AKR1C3*) has been

Table I. Polymorphisms of genes encoding sex hormones and hormone regulators.

Gene (locus, protein name and its function)	Variant		Association with susceptibility		Phenotype
	Name	dbSNP ID	Positive (number of cases, number of controls)	Negative (number of cases, number of controls)	
<i>ADRB2</i> (5q31-q32, β 2 adrenergic receptor: calcium signalling, may play a role in insulin resistance)	p.R16G	rs1042713		Japanese women (59, 97) (Kurabayashi <i>et al.</i> , 2006)	
	p.Q27E	rs1042714	Japanese women (59, 97) (Kurabayashi <i>et al.</i> , 2006)		
<i>ADRB3</i> (8p12-p11.2, β 3 adrenergic receptor: hormone receptor)	p.W64R	rs4994		Chilean women (106, 82) (Perez-Bravo <i>et al.</i> , 2005)	Chilean women—triglyceride levels (106, 82) (Perez-Bravo <i>et al.</i> , 2005)
<i>AR</i> (Xq11.2-q12, androgen receptor: hormone receptor)	CAG repeat	rs5902610, rs25885096	Australian women (122, 83) (Hickey <i>et al.</i> , 2002)	Japanese women (59, 97) (Kurabayashi <i>et al.</i> , 2006)	Asian women—T levels (91, 12) (Mifsud <i>et al.</i> , 2000)
<i>DRD3</i> (3q13.3, dopamine D3 receptor: regulates sex hormone secretion)	MscI RFLP (p.S9G)	rs6280		Finnish women (106, 112) (Jaaskelainen <i>et al.</i> , 2005)	
<i>FSHB</i> (11p13, follicle-stimulating hormone β : sex hormone)	AccI RFLP (1736T/C)	rs6169	Chinese women (135, 105) (Tong <i>et al.</i> , 2000)	Caucasian women (152, 96) (Kahsar-Miller <i>et al.</i> , 1999)	
<i>FSHR</i> (2p21-p16, follicle-stimulating hormone receptor: hormone receptor)	p.A307T	rs6165	Japanese women (18, 258) (Sudo <i>et al.</i> , 2002)	Chinese women (124, 236) (Tong <i>et al.</i> , 2001)	Caucasian women (93, 51) (Conway <i>et al.</i> , 1999)
	p.N680S	rs6166	Japanese women (18, 258) (Sudo <i>et al.</i> , 2002)	Chinese women (124, 236) (Tong <i>et al.</i> , 2001)	
<i>FST</i> (5q11.2, follistatin: inhibits follicle-stimulating hormone release)	951G/A	rs1746136		Caucasian women (93, 51) (Conway <i>et al.</i> , 1999)	Spanish women (34, 15) (Calvo <i>et al.</i> , 2001)
	343xT/A	rs722910		Families, mixed descent (249 ^a) (Urbanek <i>et al.</i> , 2000b)	
	D5S474, D5S822, D5S628			Families, mixed descent (249 ^a) (Urbanek <i>et al.</i> , 2000b)	

		rs3797297		Australian women (173, 107) (Jones <i>et al.</i> , 2007)	Australian women—FAI and SHBG levels (173, 107) (Jones <i>et al.</i> , 2007)
		rs11745088		Australian women (173, 107) (Jones <i>et al.</i> , 2007)	Australian women—DHES-S level (173, 107) (Jones <i>et al.</i> , 2007)
		rs1423560, rs3203788, rs1062809, rs1127760, rs1127761, rs7121		Australian women (173, 107) (Jones <i>et al.</i> , 2007)	
<i>GNAS</i> (20q13.32, guanine nucleotide-binding protein G(s) subunit alpha isoform: GPCR signal transduction)	393T/C			German caucasian women (278, 209) (Hahn <i>et al.</i> , 2006)	German caucasian women—BMI/weight, insulin resistance (278, 209) (Hahn <i>et al.</i> , 2006)
<i>LEPR</i> (1p31, leptin receptor: regulates hormone activity)	p.K109R	rs1137100		Finnish women (38, 122) (Oksanen <i>et al.</i> , 2000)	
	p.R223Q	rs1137101		Finnish women (38, 122) (Oksanen <i>et al.</i> , 2000)	
	p.N656K	rs8179183		Finnish women (38, 122) (Oksanen <i>et al.</i> , 2000)	
	ins/del CTTTA in 3'-UTR			Finnish women (38, 122) (Oksanen <i>et al.</i> , 2000)	
<i>LHB</i> (19q13.32, luteinizing hormone: sex hormone)	p.W8R (p.W28R)	rs1800447		Turkish women (30, 30) (Elter <i>et al.</i> , 1999), UK women (153, 212) (Rajkhowa <i>et al.</i> , 1995)	
	p.I15T (p.I35T)	rs3434826		Turkish women (30, 30) (Elter <i>et al.</i> , 1999), UK women (153, 212) (Rajkhowa <i>et al.</i> , 1995)	
<i>SHBG</i> (17p13-p12, sex hormone-binding globulin: mediates hormonal activity)	TAAAA repeat	rs35785886	Greek women (185, 324) (Xita <i>et al.</i> , 2003)	Slovenian women (123, 110) (Ferk <i>et al.</i> , 2007)	Greek women—FAI and SHBG levels (185, 324) (Xita <i>et al.</i> , 2003)
	p.D327 N	rs6259		Czech Republic women (248, 109) (Bendlova <i>et al.</i> , 2007) Slovenian women (123, 110) (Ferk <i>et al.</i> , 2007)	Slovenian women—SHBG levels (123, 110) (Ferk <i>et al.</i> , 2007)

The list (ordered alphabetically) shows genes that have been investigated for their role in polycystic ovary syndrome. Studies showing positive or no associations of these genes with disease susceptibility and positive associations with phenotype (clinical characteristics of the condition, e.g. endocrinological abnormalities) are presented. Genes are listed alphabetically. ^aNumber of PCOS families. BMI, body mass index; PCOS, polycystic ovary syndrome; RFLP, restriction fragment length polymorphism.

Table II. Polymorphisms of genes encoding enzymes involved in steroid metabolism and biosynthesis.

Gene (locus, protein name and its function)	Variant		Association with susceptibility		Phenotype	Treatment response
	Name	dbSNP ID	Positive (number of cases, number of controls)	Negative (number of cases, number of controls)		
<i>AKR1C3</i> (17q11-q21, 5 17β -hydroxysteroid dehydrogenase: testosterone biosynthesis, oestrogen metabolism)	- 71G/A	rs3763676	North American women (121, 128) (Qin <i>et al.</i> , 2006)			
<i>CYP11A1</i> (15q22-q24, cytochrome P450 11A1 enzyme: steroid biosynthesis, oestrogen metabolism, Phase I detoxification)	MspI RFLP (6235T/C) (3801T/C)	rs4646903	South Indian women (180, 72) (Babu <i>et al.</i> , 2004)			
<i>CYP11A1</i> (15q23-q24, cytochrome P450 11A1 enzyme: steroid biosynthesis)	TTTTA VNTR (D15S520)		Greek women (80, 90) (Diamanti-Kandarakis <i>et al.</i> , 2000)		Chinese women— BMI (201, 147) (Hahn <i>et al.</i> , 2006; Wang <i>et al.</i> , 2006a), UK women—T levels (97, 51aPCO, 59) (Gharani <i>et al.</i> , 1997)	
			UK (371, 350) and Finnish women (1589 ^a) (Gaasenbeek <i>et al.</i> , 2004)			
			Chinese women (201, 147) (Hahn <i>et al.</i> , 2006; Wang <i>et al.</i> , 2006a)			
			UK women (97, 51aPCO, 59) (Gharani <i>et al.</i> , 1997)			
	AAAT repeat (intron 1)			UK (371, 350) and Finnish women (1589 ^a) (Gaasenbeek <i>et al.</i> , 2004)		
<i>CYP11B2</i> (8q21-q22, aldosterone synthetase: steroid biosynthesis, regulates ovarian renin-angiotensin system)	- 344C/T	rs1799998	Chinese women (92, 60) (Zhao <i>et al.</i> , 2003)			
<i>CYP17A1</i> (10q24.3, 17α -hydroxylase: oestrogen biosynthesis)	MspA1 RFLP (- 34T/C)	rs743572		Polish women (55, 56) (Marszalek <i>et al.</i> , 2001)		
				Greek women (50, 50) (Diamanti-Kandarakis <i>et al.</i> , 1999)		
				UK women (69, 124) (Techatrasak <i>et al.</i> , 1997)		
				UK women (97, 59) (Gharani <i>et al.</i> , 1997)		
<i>CYP19A1</i> (15q21.1, aromatase: steroid biosynthesis)	TTTTA VNTR (D15S103)			Spanish women (186, 71) (Petry <i>et al.</i> , 2005)		
	SNP44	rs12907866				
	SNP50	rs2414096				Spanish women—'PCOS score' (186, 71) (Petry <i>et al.</i> , 2005)
	SNP60	rs17601241				
	SNP64	rs4646				
<i>EPHX1</i> (1q42.1, microsomal epoxide hydrolase: detoxification, component of the anti-oestrogen binding site complex)	p.Y113H	rs1051740	Finnish women (rs1051740, rs2234922 haplotype) (112, 115) (Korhonen <i>et al.</i> , 2003b)			
	p.H139R	rs2234922	Finnish women (rs1051740, rs2234922 haplotype) (112, 115) (Korhonen <i>et al.</i> , 2003b)			

<i>GATA6</i> (18q11.2, GATA binding protein 6: transcription factor)		rs7235350, rs9957475, rs9944560, rs3764962, rs1941083		Australian women (173, 107) (Jones <i>et al.</i> , 2006)	
<i>GSTM1</i> (1p13.3, glutathione S-transferase M1: phase 2 detoxification)	Null deletion			South Indian women (180, 72) (Babu <i>et al.</i> , 2004)	
<i>GSTT1</i> (22q11.23, glutathione S-transferase T1: phase 2 detoxification)	Null deletion			South Indian women (180, 72) (Babu <i>et al.</i> , 2004)	
<i>H6PD</i> (1p36, hexose-6 phosphate dehydrogenase: steroid biosynthesis)	p.R453Q	rs6688832	Spanish women (116, 76) (San Millan <i>et al.</i> , 2005)	UK women (191, 261) (Draper <i>et al.</i> , 2006)	Spanish women—basal F and 17OHP levels (116, 76) (San Millan <i>et al.</i> , 2005)
<i>HSD11B1</i> (1q32-q41, 11 β -hydroxysteroid dehydrogenase: steroid biosynthesis)	83557ins/delA			Spanish women (116, 76) (San Millan <i>et al.</i> , 2005)	
	83597T/G	rs12086634		UK women (202,263) (Draper <i>et al.</i> , 2006) UK women (202, 263) (Draper <i>et al.</i> , 2006)	
<i>HSD17B6</i> (12q13, 17 β -hydroxysteroid dehydrogenase 6: steroid biosynthesis)		rs898611	Australian women (173, 107) (Jones <i>et al.</i> , 2006)		Australian women—fasting insulin and glucose levels, BMI (173, 107) (Jones <i>et al.</i> , 2006)
		rs2277339, rs10459246		Australian women (173, 107) (Jones <i>et al.</i> , 2006)	
		rs7967600		Australian women (173, 107) (Jones <i>et al.</i> , 2006)	Australian women—fasting insulin and glucose levels, BMI (173, 107) (Jones <i>et al.</i> , 2006)
		rs1870673		Australian women (173, 107) (Jones <i>et al.</i> , 2006)	Australian women—BMI, T levels (173, 107) (Jones <i>et al.</i> , 2006)
<i>SRD5A1</i> (5p15, 5 α -reductase isoform: steroid biosynthesis)		rs1227117		Australian women (173, 107) (Jones <i>et al.</i> , 2006)	Australian women—BMI (173, 107) (Jones <i>et al.</i> , 2006)
		Haplotype: rs3797179, rs39848 Haplotype: rs472402, rs2677933, rs248805, rs3822430, rs10060745	North American women (287, 187) (Goodarzi <i>et al.</i> , 2006)		North American women—hirsutism, mFG score (287, 187) (Goodarzi <i>et al.</i> , 2006) North American women—hirsutism, mFG score (287, 187) (Goodarzi <i>et al.</i> , 2006)
<i>SRD5A2</i> (2p23, 5 α -reductase isoform: steroid biosynthesis)		Haplotype: rs11889731, rs7571644, rs12470143, rs12467911, rs2300697	North American women (287, 187) (Goodarzi <i>et al.</i> , 2006)		

Continued

Table II. *Continued*

Gene (locus, protein name and its function)		Variant		Association with susceptibility		Phenotype	Treatment response
Name	dbSNP ID	Positive (number of cases, number of controls)	Negative (number of cases, number of controls)				
	rs11675297, rs2754530		North American women (287, 187) (Goodarzi <i>et al.</i> , 2006)	North American women (287, 187) (Goodarzi <i>et al.</i> , 2006)			
p.V89L	rs523349	North American women (287, 187) (Goodarzi <i>et al.</i> , 2006)					

The list (ordered alphabetically) shows genes that have been investigated for their role in polycystic ovary syndrome. Studies showing positive or no associations of these genes with disease susceptibility and positive associations with phenotype (clinical characteristics of the condition, e.g. endocrine abnormalities) and/or treatment response are presented. ^aNot a case-control study.

linked to PCOS susceptibility in North American women (Qin *et al.*, 2006). A polymorphism in the *CYP11A1* gene, which encodes the enzyme cytochrome P450 11A1, has shown an association with PCOS susceptibility in Indian women (Babu *et al.*, 2004). Cytochrome P450 11A (*CYP11A1*) is a rate-limiting enzyme involved in the synthesis of androgens. Studies have indicated that a pentanucleotide repeat in the gene is associated with PCOS susceptibility (Gharani *et al.*, 1997; Diamanti-Kandarakis *et al.*, 2000; Gaasenbeek *et al.*, 2004; Wang *et al.*, 2006a; Jones *et al.*, 2007). However, the large study of Gaasenbeek *et al.* found only a weak association between the pentanucleotide repeat and PCOS, and no association with another promoter microsatellite (Gaasenbeek *et al.*, 2004). In Chinese women, a polymorphism in the promoter region of the aldosterone synthase gene (*CYP11B2*), which affects the balance of the ovarian renin-angiotensin system, has been linked to PCOS susceptibility (Zhao *et al.*, 2003). A polymorphism in the gene for 17 α -hydroxylase (*CYP17A1*), which is active in estrogen biosynthesis, was not associated with PCOS susceptibility (Techatraisak *et al.*, 1997; Diamanti-Kandarakis *et al.*, 1999; Marszalek *et al.*, 2001). The *CYP19A1* gene encodes a key component of aromatase, which catalyses the production of estrogens from androgens. Although polymorphisms in this gene were not found to directly affect susceptibility to PCOS in women from the UK (Gharani *et al.*, 1997) or Spain (Petry *et al.*, 2005), the SNP50 polymorphism in this gene did influence PCOS severity in the Spanish study (Petry *et al.*, 2005).

Sex steroid metabolism. The low activity haplotype (H113-R139) of the gene encoding the detoxification enzyme microsomal epoxide hydrolase (*EPHX1*) was significantly associated with PCOS susceptibility in Finnish women (Korhonen *et al.*, 2003b). A variant of the hexose-6-phosphate dehydrogenase gene (*H6PD*), which has been implicated in a rare cortisone reductase deficiency that is characterized by a PCOS-like phenotype, has been linked to PCOS susceptibility in Spanish (San Millan *et al.*, 2005), but not in UK women (Draper *et al.*, 2006). Polymorphisms in the gene encoding 17 β -hydroxysteroid dehydrogenase 6 (*HSD17B6*) have been found to be associated with either PCOS or with the clinical phenotype (Jones *et al.*, 2006). A North American genetic association study of the two isoforms of the steroid biosynthesis enzyme 5 α -reductase found that haplotypes in both *SRD5A1* and *SRD5A2* were associated with PCOS susceptibility, but that only variants in the *SRD5A1* gene were associated with severity of hirsutism (Goodarzi *et al.*, 2006).

Polymorphisms in genes encoding 11 β -hydroxysteroid dehydrogenase (*HSD11B1*) (San Millan *et al.*, 2005; Draper *et al.*, 2006), the glutathione S-transferases M1 or T1 (*GSTM1*, *GSTT1*) (Babu *et al.*, 2004) and the transcription factor GATA-binding protein 6 (*GATA6*) (Jones *et al.*, 2006) have also been investigated but failed to show an association with PCOS.

Genes involved in type 2 diabetes and cardiovascular disease

The increased risk of type 2 diabetes and cardiovascular disease in women with PCOS has led to numerous association studies in genes related to these diseases (Table III).

Adiponectin. Patients with obesity, type 2 diabetes, insulin resistance or PCOS have abnormally low levels of adiponectin, and therefore several studies have investigated the effect of two

Table III. Polymorphisms of genes encoding proteins involved in type 2 diabetes and cardiovascular disease.

Gene (locus, protein name and its function)	Variant		Association with susceptibility		Phenotype	Treatment response
	Name	dbSNP ID	Positive (number of cases, number of controls)	Negative (number of cases, number of controls)		
Genes encoding proteins involved in insulin pathways <i>ADIPOQ</i> (3q27, adiponectin: mediates insulin resistance)	45G/T	rs2241766	German women (57, 567) (Haap <i>et al.</i> , 2005)	Greek women (132, 100; 100, 140) (Panidis <i>et al.</i> , 2004; Xita <i>et al.</i> , 2005) Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004)	Greek women—insulin level (100, 140) (Panidis <i>et al.</i> , 2004; Xita <i>et al.</i> , 2005)	
	276G/T	rs1501299	Finnish women (143, 245) (Heinonen <i>et al.</i> , 2005)	Greek women (132, 100; 100, 140) (Panidis <i>et al.</i> , 2004; Xita <i>et al.</i> , 2005) Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004) Finnish women (143, 245) (Heinonen <i>et al.</i> , 2005) Finnish women (58, 91) (Heinonen <i>et al.</i> , 2001)	Greek women—insulin exposure, adiponectine levels, BMI (100, 140) (Panidis <i>et al.</i> , 2004; Xita <i>et al.</i> , 2005)	
<i>APOE</i> (19q13.2, apolipoprotein E: mediates insulin resistance)	3937C/T and 4075C/T (e2, e3 and e4 alleles)	rs429358+rs7412				
<i>CAPN10</i> (2q37.3, calpain-10: mediates insulin resistance)	4841T/C (UCSNP44)	rs2975760	Spanish women (148, 93) (Gonzalez <i>et al.</i> , 2003)	German women (57, 567) (Haap <i>et al.</i> , 2005)	Spanish women—hirsutism score (81, 37) (Escobar-Morreale <i>et al.</i> , 2002)	
	4852G/A (UCSNP43)	rs3792267		Spanish women (81, 37) (Escobar-Morreale <i>et al.</i> , 2002) German women (57, 567) (Haap <i>et al.</i> , 2005)	Spanish women—hirsutism score (81, 37) (Escobar-Morreale <i>et al.</i> , 2002)	
	7920ins/de132 bp (UCSNP19)			Spanish women (148, 93; 81, 37) (Gonzalez <i>et al.</i> , 2003) North American women (181, 422) (Ehrmann <i>et al.</i> , 2002a) Spanish women (148, 93) (Gonzalez <i>et al.</i> , 2003) North American women (181, 422) (Ehrmann <i>et al.</i> , 2002a)		
	UCSNP45	rs41266971	Spanish women (81, 37) (Escobar-Morreale <i>et al.</i> , 2002)	German women (57, 567) (Haap <i>et al.</i> , 2005)		

Continued

Table III. Continued

Gene (locus, protein name and its function)	Variant		Association with susceptibility		Phenotype	Treatment response
	Name	dbSNP ID	Positive (number of cases, number of controls)	Negative (number of cases, number of controls)		
	UCSNP63	rs5030952		Spanish women (148, 93) (Gonzalez <i>et al.</i> , 2003)	Spanish women (148, 93)—cholesterol level (Gonzalez <i>et al.</i> , 2003)	
<i>ENPP1</i> (6q22-q23, plasma cell differentiation antigen glycoprotein (PC-1): mediates insulin resistance)	p.K121Q	rs1044498	Finnish women (143, 115) (Heinonen <i>et al.</i> , 2004)	North American women (181, 422) (Ehrmann <i>et al.</i> , 2002a) Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004)		
<i>FOXC2</i> (16q22-16q24, forkhead box C2: transcription factor, mediates insulin resistance)	-512C/T	rs34221221		Japanese women (123, 180) (Baba <i>et al.</i> , 2007) German women (57, 567) (Haap <i>et al.</i> , 2005)		
<i>GYS1</i> (19q13.3, glycogen synthetase 1 (muscle): mediates insulin resistance)	XbaI RFLP (C/T)	rs8103451		UK women (90, 62) (Rajkhowa <i>et al.</i> , 1996)		
<i>IGF1</i> (12q22-q23, insulin-like growth factor-1: insulin signalling)	CA repeat (promoter)			Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004)		
<i>IGF2</i> (11p15.5, insulin-like growth factor-2: insulin signalling)	ApaI RFLP (17200G/A)	rs680	Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004)			
<i>IGF1R</i> (15q26.3, insulin-like growth factor-1 receptor: insulin signalling)	AGG repeat at position 967			Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004)		
<i>IGF2R</i> (6q26, insulin-like growth factor-2 receptor: insulin signalling)	ins/delACAA in 3' UTR			Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004)		
<i>INS</i> (11p15.5, insulin: hormone)	VNTR		UK women, linkage studies (54, 78 and 2 family-based linkage studies; 74, 150) (Franks <i>et al.</i> , 1999; Michelmore <i>et al.</i> , 2001)	UK (255 families+185, 1062) and Finnish women (530 [72 confirmed by US], 1069) (Powell <i>et al.</i> , 2005) Spanish women (40, 38) (Calvo <i>et al.</i> , 2002) Czech women (38, 22) (Vankova <i>et al.</i> , 2002)		
<i>INSR</i> (19p13.3-p13.2, insulin receptor: hormone regulation)	p.H1058H T/C (3364T/C)	rs1799817	Chinese women (120, 40) (Chen <i>et al.</i> , 2004) Non-obese North American women (99, 136) (Siegel <i>et al.</i> , 2002) Korean women (134, 100) (Lee <i>et al.</i> , 2007)	Korean women (174, 93) (Lee <i>et al.</i> , 2006)	Chinese women—weight/BMI (120, 40) (Chen <i>et al.</i> , 2004)	
	p.C1008C T/C (3128T/C)	No dbSNP ID	Chinese women (109, 107) (Jin <i>et al.</i> , 2006)		Chinese women—insulin sensitivity (109, 107) (Jin <i>et al.</i> , 2006)	
	109482A/G, 109665C/T,	No dbSNP ID, rs6510959,		Korean women (134, 100) (Lee <i>et al.</i> , 2007)		

	125498A/G, 127527G/A, 143485G/C, 161822G/A, 168828T/A 176477C/T	rs2303672, rs2059806, rs2252673, rs2860175, No dbSNP ID No dbSNP ID				
<i>IRS1</i> (2q36, insulin receptor substrate-1: insulin signalling)	p.G972R	rs1801278	Korean women (134, 100) (Lee <i>et al.</i> , 2007) Chilean women (146, 97) (Sir-Petermann <i>et al.</i> , 2004)	German women (57, 567) (Haap <i>et al.</i> , 2005)	Chilean women—insulin resistance (146, 97) (Sir-Petermann <i>et al.</i> , 2004)	Metformin response in Turkish women (60 ^a) (Ertunc <i>et al.</i> , 2005)
			Japanese women (123, 180) (Baba <i>et al.</i> , 2007)	North American women (227 ^a) (Ehrmann <i>et al.</i> , 2002b)	North American women, in carriers of CYP21 heterozygous mutation (114, 95) (Witchel <i>et al.</i> , 2005)	
			Turkish women (60, 60) (Dilek <i>et al.</i> , 2005)	Spanish caucasian women (103, 48) (Villuendas <i>et al.</i> , 2005) Caucasian women (53, 102) (El Mkaem <i>et al.</i> , 2001)	Spanish caucasian women—BMI, insulin resistance (103, 48) (Villuendas <i>et al.</i> , 2005) Caucasian women—insulin resistance (53, 102) (El Mkaem <i>et al.</i> , 2001)	
<i>IRS2</i> (13q34, insulin receptor substrate-2: insulin signalling)	p.G1057D	rs1805097		German women (57, 567) (Haap <i>et al.</i> , 2005)	Caucasian American women—blood glucose level (227 ^a) (Ehrmann <i>et al.</i> , 2002b)	
				Spanish caucasian women (103, 48) (Villuendas <i>et al.</i> , 2005) Caucasian women (53, 102) (El Mkaem <i>et al.</i> , 2001)	Caucasian women—blood glucose level (53, 102) (El Mkaem <i>et al.</i> , 2001)	
<i>PONI</i> (7q21.3, paraoxonase 1: antioxidant enzyme, linked to insulin resistance and cardiovascular disease)	–108C/T	rs705379	Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004)			
	p.L55M	rs854560		Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004)	Caucasian women—BMI and insulin resistance (72, 42) (San Millan <i>et al.</i> , 2004)	
	p.Q192R	rs662		Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004)		
<i>PPARG</i> (3p25, peroxisome proliferator-activated receptor- γ : transcription factor, mediates insulin resistance, regulates <i>CCL5</i> expression)	p.P12A	rs1801282	Finnish women (135, 115) (Korhonen <i>et al.</i> , 2003a)	German women (57, 567) (Haap <i>et al.</i> , 2005)	Turkish women—glucose metabolism (60, 60) (Tok <i>et al.</i> , 2005)	
				Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004)	German women—insulin sensitivity; hirsutism score (102, 104) (Hahn <i>et al.</i> , 2005)	

Continued

Table III. Continued

Gene (locus, protein name and its function)	Variant		Association with susceptibility		Phenotype	Treatment response
	Name	dbSNP ID	Positive (number of cases, number of controls)	Negative (number of cases, number of controls)		
				Chinese women (201, 147) (Wang <i>et al.</i> , 2006b)	North American women—insulin sensitivity (218 ^a) (Hara <i>et al.</i> , 2002)	
	p.H447H, 1431C/T	rs3856806		North American women (285, 187) (Antoine <i>et al.</i> , 2007) Italian women (100, 100) (Orio <i>et al.</i> , 2003a)	Italian women—BMI (100, 100) (Orio <i>et al.</i> , 2003a)	
<i>PPARGC1A</i> (4p15.2, peroxisome proliferator-activated receptor gamma, coactivator 1 alpha: transcriptional coactivator for steroid receptors and nuclear receptors)	p.S482G	rs8192678		Chinese women (201, 147) (Wang <i>et al.</i> , 2006b)		
<i>PPP1R3A</i> (7q31.1, protein phosphatase-1: mediates insulin resistance)	5 bp ins/del in the 3'UTR (ARE-1/ARE-2)	rs5886683+rs5886684			North American women—insulin sensitivity (186 ^a) (Alcoser <i>et al.</i> , 2004)	
<i>PTPN1</i> (20q13.1-q13.2, protein tyrosine phosphatase 1B: insulin signalling)	981C/T	rs11575938		Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004)		
	1484ins/delG			Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004)		
<i>RETN</i> (19p13.2, resistin: mediates insulin resistance)	-179C/G (-180C/G)	rs1862513		Greek women (320, 180) (Xita <i>et al.</i> , 2004)	Greek women—BMI (320, 180) (Xita <i>et al.</i> , 2004)	
<i>SORBS1</i> (10q23.3-q24.1, sorbin: may play a role in insulin resistance)	p.T228A	rs2281939		Caucasian women (72, 42) (San Millan <i>et al.</i> , 2004)	Caucasian women—BMI (72, 42) (San Millan <i>et al.</i> , 2004)	
Mediators of vascular function or genes linked to cardiovascular risk						
<i>AGT</i> (1q42-q43, angiotensin I: mediates vascular homeostasis)	p.M235T	rs699	Italian women (95, 64) (Zulian <i>et al.</i> , 2005)		Italian women—insulin sensitivity (95, 64) (Zulian <i>et al.</i> , 2005)	
<i>MTHFR</i> (1p36.3, methylenetetrahydrofolate reductase: olate metabolism, linked to cardiovascular disease)	677C/T (p.A222V)	rs1801133		Italian women (70, 70) (Orio <i>et al.</i> , 2003b)		
<i>PLAT</i> (8p12, tissue plasminogen activator: fibrinolysis system, linked to cardiovascular disease)	Alu repeat ins/del 311 bp	rs4646972		Turkish women (91, 100) (Karadeniz <i>et al.</i> , 2007)	Turkish women—cholesterol and LDL levels (91, 100) (Karadeniz <i>et al.</i> , 2007)	

<i>SERPINE1</i> (7q21.3-q22, plasminogen-activator inhibitor-1 (PAI-1): fibrinolysis system, linked to cardiovascular disease)	–675ins/deG (4G/5G)	Several IDs: rs1799768, rs34857375, rs1799762, rs1799889	Greek women (98, 64) (Diamanti-Kandarakis <i>et al.</i> , 2004)	Caucasian women (72, 42; 106, 102) (San Millan <i>et al.</i> , 2004; Walch <i>et al.</i> , 2005a)	North American women—rate of miscarriages (149, 234) (Glueck <i>et al.</i> , 1999) Greek women—SERPINE1 (PAI-1) levels (98, 64) (Diamanti-Kandarakis <i>et al.</i> , 2004)
			Turkish women (91, 100) (Karadeniz <i>et al.</i> , 2007)		

The list shows genes that have been investigated for their role in polycystic ovary syndrome. Genes are grouped according to their function (and then alphabetically). Studies showing positive or no associations of these genes with disease susceptibility and positive associations with phenotype (clinical characteristics of the condition, e.g. endocrine abnormalities) are presented. *Replicated in a family study (52 probands). NB: when only the number of cases is given, the study was not case control. BMI, body mass index; *CCL5*, formally known as *RANTES*, Regulation upon Activation Normal T cell Expressed and Secreted; LDL, low-density lipoprotein; PAI-1, plasminogen activator inhibitor-1; RFLP, restriction fragment-length polymorphism; VNTR, variable number of tandem repeats.

SNPs in the adiponectin gene (*ADIPOQ*) on PCOS susceptibility. Whereas there was an association with the 45G/T polymorphism in German and Finnish women (Haap *et al.*, 2005; Heinonen *et al.*, 2005), there was no association for this polymorphism or the 276G/T polymorphism in Greek or Caucasian women (Panidis *et al.*, 2004; San Millan *et al.*, 2004; Heinonen *et al.*, 2005; Xita *et al.*, 2005) (NB: 276G/T was not investigated by Haap *et al.*). Both polymorphisms may, however, affect insulin levels (Xita *et al.*, 2005).

Calpain-10. The gene for the cysteine protease calpain-10 (*CAPN10*) is associated with type 2 diabetes, and potential associations between five different polymorphisms and PCOS have been investigated. In the *CAPN10* gene, two different Spanish studies found an association between either the SNP UCSNP44 (Gonzalez *et al.*, 2003) or UCSNP45 (Escobar-Morreale *et al.*, 2002) and PCOS susceptibility. These associations were not observed in a German study (Haap *et al.*, 2005). No associations were found between the UCSNP43, UCSNP19, UCSNP63 polymorphisms and PCOS (Ehrmann *et al.*, 2002a; Gonzalez *et al.*, 2003), although there were some links with the symptoms of PCOS (Escobar-Morreale *et al.*, 2002; Gonzalez *et al.*, 2003).

Plasma cell differentiation antigen. Genes encoding components of the insulin signalling system have been investigated in relation to PCOS across a number of studies. Plasma cell differentiation antigen glycoprotein (*PC-1*, also known as *ENPPI*, ectonucleotide pyrophosphatase/phosphodiesterase 1) is a mediator of insulin receptor signalling and has been implicated in insulin resistance. However, studies investigating an association between PCOS susceptibility and the p.K121Q polymorphism have reported both positive (Heinonen *et al.*, 2004) and negative (San Millan *et al.*, 2004; Baba *et al.*, 2007) results. Although early case-control studies in the UK indicated a link between PCOS susceptibility and the insulin gene variable number of tandem repeats microsatellite length (Franks *et al.*, 1999; Michelmore *et al.*, 2001), a large subsequent study in more than 3000 women from the UK and Finland found no such association (Powell *et al.*, 2005). In addition, studies conducted in the Czech Republic (Vankova *et al.*, 2002) and Spain (Calvo *et al.*, 2002) have found no association between this variant and PCOS.

Insulin receptor gene. Three polymorphisms in the insulin receptor gene (*INSR*) have shown a positive association with PCOS: 3364T/C (p.H1058H) in Chinese (Chen *et al.*, 2004) and American women (Siegel *et al.*, 2002), but not Korean women (Lee *et al.*, 2006); 3128T/C (p.C1008C) in Chinese women (Jin *et al.*, 2006); and 176477C/T in Korean women (Lee *et al.*, 2007). The 2007 study by Lee *et al.* investigated a total of nine polymorphisms but only found an association with 176477C/T. The 3364T/C and 3128T/C polymorphisms were also associated with the phenotype of PCOS (Chen *et al.*, 2004; Jin *et al.*, 2006).

Insulin receptor substrates. The insulin receptor substrates IRS-1 and IRS-2 are proteins that participate in the insulin signal transduction pathway. Extensive investigations on a possible association between the p.G972R polymorphism in the *IRS-1* gene and PCOS have been carried out, with both positive (Sir-Petermann *et al.*, 2004; Dilek *et al.*, 2005; Baba *et al.*, 2007) and negative (El Mkaem *et al.*, 2001; Ehrmann *et al.*, 2002b; Haap *et al.*, 2005; Villuendas *et al.*, 2005) results. The polymorphism was

Table IV. Polymorphisms of genes encoding proteins involved in tissue remodelling or inflammatory processes.

Gene (locus, protein name and its function)	Variant		Association with susceptibility		Phenotype
	Name	dbSNP ID	Positive (number of cases, number of controls)	Negative (number of cases, number of controls)	
Genes involved in tissue remodeling					
<i>MMP1</i> (11q22.3, matrix metalloproteinase-1: endometrial lesion formation)	1607ins/delG	rs11292517	Caucasian women (62, 94) (Walch <i>et al.</i> , 2005b)		
Inflammatory mediators					
<i>HLA-A</i> (6p21.3, major histocompatibility complex class I, A: antigen processing and presentation)	A11		Japanese women (56, 237) (Kaibe <i>et al.</i> , 2006)		
<i>HLA-B</i> (6p21.3, major histocompatibility complex class I, B: antigen processing and presentation)	B39		Protective effect in Japanese women (56, 237) (Kaibe <i>et al.</i> , 2006)		
<i>HLA-C</i> (6p21.3, major histocompatibility complex class I, C: antigen processing and presentation)	All alleles			Japanese women (56, 237) (Kaibe <i>et al.</i> , 2006)	
<i>HLA-DRB1</i> (6p21.3, major histocompatibility complex class I, DR beta 1: antigen processing and presentation)	*0403		Japanese women (68, 292) (Kaibe <i>et al.</i> , 2006)		
<i>HFE</i> (6p21.3, haemochromatosis protein (MHC class I family): antigen processing and presentation, iron binding)	p.H63D	rs1799945		Spanish women (78, 43) (Botella-Carretero <i>et al.</i> , 2006)	
	p.C282Y	rs1800562		Spanish women (78, 43) (Botella-Carretero <i>et al.</i> , 2006)	
<i>IL1A</i> (2q14, interleukin-1 alpha: cytokine)	– 889C/T	rs1800587	Austrian caucasian women (105, 102) (Kolbus <i>et al.</i> , 2007)		Austrian caucasian women—FSH serum level, LH/FSH ratio (105, 102) (Kolbus <i>et al.</i> , 2007)
<i>IL1B</i> (2q14, interleukin-1 beta: cytokine)	– 511C/T, exon 5	rs16944		Austrian caucasian women (105, 102) (Kolbus <i>et al.</i> , 2007)	
	3953G/A (exon5 <i>TaqI</i> RFLP, E1/E2)	rs1143634		Austrian caucasian women (105, 102) (Kolbus <i>et al.</i> , 2007)	
<i>IL6</i> (7p21, interleukin-6: cytokine)	– 597G/A	rs1800797	Spanish caucasian women (85, 25) (Villuendas <i>et al.</i> , 2002)		
	– 572G/C	rs1800796		Spanish caucasian women (85, 25) (Villuendas <i>et al.</i> , 2002)	
	– 174G/C	rs1800795	Spanish caucasian women (85, 25) (Villuendas <i>et al.</i> , 2002)	Austrian caucasian women (62, 94) (Walch <i>et al.</i> , 2004)	Austrian caucasian women—BMI, T levels, OGTT results (62, 94) (Walch <i>et al.</i> , 2004) German women—androstendione levels (50, 0) (Mohlig <i>et al.</i> , 2004)
<i>IL6R</i> (1q22, interleukin-6 repector alpha: cytokine receptor)	CA repeat			Spanish women (88, 45) (Escobar-Morreale <i>et al.</i> , 2003)	
<i>IL6ST</i> (5q11.2, gp130, interleukin-6 repector beta: cytokine receptor)	p.G148R	rs3729960	Spanish women (88, 45) (Escobar-Morreale <i>et al.</i> , 2003)		
<i>TNF</i> (6p21.3, tumour necrosis factor alpha: cytokine)	– 850C/T (– 857C/T)	rs1799724		Cauasian Finnish women (87, 115) (Korhonen <i>et al.</i> , 2002)	Australian women—OGTT results (122, 28) (Milner <i>et al.</i> , 1999)

	-308G/A	rs1800629	Chinese women (118, 54) (Mao <i>et al.</i> , 2000) Australian women (122, 28) (Milner <i>et al.</i> , 1999)	Spanish women (87, 36) and Italian women (64, 29)—hyperandrogenism (Peral <i>et al.</i> , 2002)
<i>TNFRSF1B</i> (1p36.3-p36.2, tumour necrosis factor receptor 2: cytokine receptor)	p.M196R	rs1061622	Spanish women (87, 36) and Italian women (64, 29) (Peral <i>et al.</i> , 2002)	
	CA repeat		Spanish women (87, 36) and Italian women (64, 29) (Peral <i>et al.</i> , 2002)	
	1663G/A	rs1061624	Spanish women (87, 36) and Italian women (64, 29) (Peral <i>et al.</i> , 2002)	
	1668T/G	rs5030792	Spanish women (87, 36) and Italian women (64, 29) (Peral <i>et al.</i> , 2002)	
	1690T/C	rs3397	Spanish women (87, 36) and Italian women (64, 29) (Peral <i>et al.</i> , 2002)	

The list shows genes that have been investigated for their role in polycystic ovary syndrome. Genes are grouped according to their function (and then alphabetically), and studies showing positive or no associations of these genes with disease susceptibility and positive associations with phenotype (clinical characteristics of the condition, e.g. endocrine abnormalities) are presented.

associated with symptoms of PCOS across four different studies (El Mkadem *et al.*, 2001; Sir-Petermann *et al.*, 2004; Villuendas *et al.*, 2005; Witchel *et al.*, 2005). Metformin, an insulin-sensitizing drug that is often used in women with PCOS to induce ovulation, is believed to act via phosphorylation of IRS proteins. This could explain why a study in Turkish women with PCOS indicated that the IRS-1 genotype may influence a patient's response to metformin therapy (Ertunc *et al.*, 2005). In contrast, investigations into the p.G1057D polymorphism in the *IRS-2* gene have failed to show an association with susceptibility to PCOS (El Mkadem *et al.*, 2001; Haap *et al.*, 2005; Villuendas *et al.*, 2005), but have linked this polymorphism with blood glucose levels in women with PCOS (El Mkadem *et al.*, 2001; Ehrmann *et al.*, 2002b).

Insulin-like growth factors. An ApaI RFLP in the insulin-like growth factor-2 (*IGF-2*) gene has been linked to PCOS susceptibility in Caucasian women (San Millan *et al.*, 2004), whereas polymorphisms in genes encoding IGF-1, IGF-1 receptor and IGF-2 receptor were not associated (San Millan *et al.*, 2004). Of three variants investigated in the paraoxonase gene (*PON1*) (San Millan *et al.*, 2004), the -108C/T variant was linked to PCOS susceptibility in Caucasian women. The p.L55M polymorphisms was not associated with PCOS, but was linked with a higher BMI and greater insulin resistance in women who were homozygotic for the 55M polymorphism compared with carriers of the common 55L allele. This polymorphism is believed to contribute to impaired insulin function by increasing levels of oxidative stress in women with PCOS. The remaining polymorphism investigated (p.Q192R) was associated with neither susceptibility nor phenotype.

Peroxisome proliferator-activated receptor-γ. Peroxisome proliferator-activated receptor-γ (*PPARG*) regulates the expression of several anti-atherosclerotic proteins. One study of the polymorphism p.P12A showed an association with PCOS susceptibility in Finnish women (Korhonen *et al.*, 2003a), but five other studies found no such association (Orio *et al.*, 2003a; San Millan *et al.*, 2004; Haap *et al.*, 2005; Wang *et al.*, 2006b; Antoine *et al.*, 2007). Three studies found effects on glucose metabolism or insulin sensitivity in carriers of this polymorphism (Hara *et al.*, 2002; Hahn *et al.*, 2005; Tok *et al.*, 2005) and indications that genotype also determines levels of hirsutism (Hahn *et al.*, 2005). The 1431C/T (p.H447H) polymorphism in this gene has been linked to obesity in women with PCOS (Orio *et al.*, 2003a) but not to susceptibility to PCOS (Antoine *et al.*, 2007). So far, there has only been a single study of the *PPARG* coactivator 1 alpha (*PPARGC1A*) gene and no association was shown between the polymorphism studied (p.S482G) and PCOS (Wang *et al.*, 2006b).

Glucose metabolism and insulin resistance. An insertion/deletion polymorphism in the AT-rich element within the 3'-untranslated region of the muscle-specific glycogen-targeting subunit of the protein phosphatase-1 gene (*PPP1R3A*), which has been associated with insulin resistance and diabetes, has been shown to influence insulin resistance and androgen levels in women with PCOS (Alcoser *et al.*, 2004). Also related to the phenotype of PCOS is the -179C/G polymorphism in the promoter region of the resistin gene (*RETN*) (Xita *et al.*, 2004) and the p.T228A polymorphism in the sorbin and SH3 domain containing 1 (*SORBS1*) gene (San Millan *et al.*, 2004). Both were linked to higher BMI in carriers,

but neither was linked with susceptibility (San Millan *et al.*, 2004; Xita *et al.*, 2004).

A number of variants of other genes involved in insulin resistance, type 2 diabetes and/or obesity have been investigated but failed to show an association with PCOS: apolipoprotein E (*APOE*) (Heinonen *et al.*, 2001), forkhead box C2 (*FOXC2*) (Haap *et al.*, 2005), glycogen synthetase 1 (*GYS1*) (Rajkhowa *et al.*, 1996) and protein tyrosine phosphatase 1B (*PTPN1*) (San Millan *et al.*, 2004).

Genes linked to cardiovascular risk. Polymorphisms in a limited number of genes linked to cardiovascular risk have been studied to assess a possible role in PCOS (Table III). In one study, the p.M235T polymorphism in the gene encoding angiotensin I (*AGT*), a potent vasoconstrictor, was linked to PCOS susceptibility and a genotype-dependent difference was seen in insulin sensitivity in women with PCOS (Zulian *et al.*, 2005). Methylene-tetrahydrofolate reductase (*MTHFR*) is an enzyme involved in the homocysteine pathway, and the p.A222V polymorphism in the *MTHFR* gene increases circulating homocysteine levels. As elevated levels of plasma homocysteine are associated with an increased risk of cardiovascular disease, a study investigated whether this polymorphism increases PCOS susceptibility in Italian women, but failed to show an effect (Orio *et al.*, 2003b). As part of the fibrinolytic system, the gene encoding tissue plasminogen activator (*PLAT*) has been investigated in relation to PCOS. Although no association between the Alu repeat insertion/deletion and susceptibility was found, PCOS patients that were homozygous for the insertion (i/i genotype) had lower levels of low-density and total cholesterol than other genotypes (Karadeniz *et al.*, 2007). Plasminogen activator inhibitor-1 (*SERPINE1*) is also involved in blood coagulation and has been implicated in cardiovascular disease. The 4G/5G polymorphism in the *SERPINE1* gene has been linked with PCOS susceptibility in one study (Diamanti-Kandarakis *et al.*, 2004), but no association was found in three other studies (San Millan *et al.*, 2004; Walch *et al.*, 2005a; Karadeniz *et al.*, 2007). Women with PCOS who are carriers of the polymorphism were, however, found to have an increased risk of miscarriage (Glueck *et al.*, 1999) and higher plasminogen activator inhibitor-1 levels (Diamanti-Kandarakis *et al.*, 2004).

Genes involved in tissue remodelling and inflammatory processes

Tissue remodelling. Matrix metalloproteinases (MMPs) are thought to play a critical role in the remodelling of the extracellular matrix during follicular development (Goldman and Shalev, 2004). The 1607G insertion/deletion polymorphism in the *MMP1* gene has been associated with an increased risk of PCOS in Caucasian women (Walch *et al.*, 2005b) (Table IV). The association of genes of the histocompatibility family with PCOS has been studied in Japanese women. Whereas women with PCOS were more likely to be carriers of the *HLA-A11* or *HLA-DRB1*0403* alleles, they were less likely to be carriers of the *HLA-B39* allele when compared with healthy individuals (indicating a protective effect). No association was found for the alleles tested within *HLA-C* (Kaibe *et al.*, 2006). No association was found between two polymorphisms in the gene encoding haemochromatosis protein, in the MHC Class I family (*HFE*), and PCOS (Botella-Carretero *et al.*, 2006).

Inflammatory cytokines. Circulating levels of several inflammatory cytokines, including tumour necrosis factor- α (TNF- α) and interleukin-6 (IL-6), correlate with insulin resistance and obesity (Milner *et al.*, 1999; Mohlig *et al.*, 2004). It has also been proposed that another inflammatory cytokine, IL-1, influences the processes of ovulation, fertilization and implantation. A study in Austrian women showed that the -889C/T polymorphism in the *IL1A* gene is associated with PCOS, which was not the case for the -511C/T and 3953G/A polymorphisms in the *IL1B* gene (Kolbus *et al.*, 2007). Within the interleukin-6 (*IL6*) gene, the 597G/A polymorphism has been shown to be associated with PCOS susceptibility (Villuendas *et al.*, 2002); the 572G/C polymorphism not associated (Villuendas *et al.*, 2002) and the -174G/C was both associated with susceptibility (Villuendas *et al.*, 2002) and not associated with (Walch *et al.*, 2004), but related to, the PCOS phenotype (Mohlig *et al.*, 2004; Walch *et al.*, 2004). An investigation into polymorphisms in the genes of the IL-6 receptor complex found no association between the presence of a microsatellite CA repeat in the *IL6R* gene, but an association between PCOS and the p.G148R polymorphism in the IL-6 signal transducer (*IL6ST*) gene (Escobar-Morreale *et al.*, 2003).

Polymorphisms in the promoter region of the TNF- α gene (*TNF*) do not appear to affect PCOS susceptibility (Milner *et al.*, 1999; Mao *et al.*, 2000; Korhonen *et al.*, 2002), but the -308G/A polymorphism is linked with glucose tolerance in women with PCOS (Milner *et al.*, 1999). The actions of TNF- α are mainly mediated by type 1 or type 2 TNF receptors. Of the five polymorphisms in the TNF receptor 2 gene (*TNFRSF1B*) investigated in relation to PCOS in Spanish and Italian women, only p.M196R was found to be associated with PCOS susceptibility and hyperandrogenism, but the other four polymorphisms investigated were not (Peral *et al.*, 2002).

Ovarian response to gonadotrophins

Ovarian stimulation is usually performed by administering exogenous FSH following pituitary down-regulation. Ovarian response to FSH, however, varies widely among women undergoing ovarian stimulation (Georgiou *et al.*, 1997) with wide-reaching clinical consequences. If ovarian response to stimulation is insufficient, the cycle may need to be cancelled. Similarly, if ovarian response is excessive, cycle cancellation may be necessary to avoid the risk of ovarian hyperstimulation syndrome (OHSS), a potentially life-threatening complication. Application of pharmacogenetics to the problem of predicting ovarian response utilizes information about the genetic make-up of an individual patient. This information may then be used in conjunction with existing clinical factors to predict treatment response (Fauser *et al.*, 2008).

FSH receptor

FSH induces the proliferation of granulosa cells and the synthesis of the androgen-converting enzyme aromatase, and also plays a crucial role in secondary follicle recruitment and selection of the dominant follicle (Fauser and Van Heusden, 1997). The action of FSH is mediated by FSHR, a member of the G-protein-coupled receptor family, which is expressed solely by granulosa cells. The *FSHR* gene harbours more than 900 SNPs, arranged in two major linkage disequilibrium blocks (Fig. 3). Two non-synonymous SNPs in strong linkage disequilibrium (p.A307T and p.N680S)




Figure 3: LD plot of the genomic region on chromosome 2 harbouring the *FSHR* gene, obtained from the HapMap website (www.hapmap.org). More than 900 SNPs are currently listed in the National Centre for Biotechnology Information SNP database. These SNPs are grouped into two major LD blocks (red triangles). *FSHR*, follicle-stimulating hormone receptor; LD, linkage disequilibrium.

have been identified in exon 10 of the *FSHR* gene (Simoni *et al.*, 1999). As the interaction between FSH and its receptor plays a key role in ovarian stimulation, a number of groups have investigated the effect of polymorphisms in the *FSHR* gene on ovarian response (Table V).

Studies in women with normal ovarian function demonstrate convincingly that SNPs in exon 10 modulate *FSHR* function and the ovarian response to FSH. This effect was first observed in a partly retrospective, non-randomized study of German women undergoing controlled ovarian hyperstimulation for assisted reproduction. The amount of FSH needed for controlled ovarian hyperstimulation to achieve similar peak estradiol levels was significantly lower in women with the N/N genotype at position 680 of the *FSHR* gene compared with women carrying the S/S or N/S genotypes, indicating a lower ovarian sensitivity to FSH *in vivo* for the S680 allele (Perez Mayorga *et al.*, 2000). Similar results were later obtained by other investigators who studied populations from different ethnic backgrounds (Sudo *et al.*, 2002; de Castro *et al.*, 2003, 2004; Behre *et al.*, 2005; Falconer *et al.*, 2005; Jun *et al.*, 2006; Loutradis *et al.*, 2006). In accordance with a genetic control of *FSHR* p.N680S genotype status on *in vitro* fertilization (IVF) outcome, de Castro *et al.* (2003) reported that women with the S/S genotype have significantly higher rates of cycle cancellation and poor response compared with carriers of the N/S or N/N genotypes. These consistent findings in various populations indicate that the effects of the *FSHR* p.N680S polymorphism are independent of ethnic background and may also be present in other, previously uninvestigated, populations. Recently, it was shown that in women undergoing IVF treatment, the clinical pregnancy rate in women with the N/N genotype is

significantly higher compared with those with the S/S genotype (Jun *et al.*, 2006). However, another study using a similar study design showed opposite results, with higher pregnancy rates in women with the S/S genotype (Klinkert *et al.*, 2006) (Table VI). These contrasting data should be interpreted with caution, and larger, well-designed and properly powered studies should be conducted before drawing conclusions about the effects of the *FSHR* genotype on pregnancy rates.

In a study involving menstrual cycle monitoring in women with normal, mono-ovulatory cycles, Greb *et al.* (2005) were able to show that during the luteo-follicular transition, serum levels of estradiol, progesterone and inhibin A were significantly lower and FSH started to rise earlier in women with the S/S genotype compared with women who carried the N/N genotype for the *FSHR* p.N680S polymorphism (Fig. 4). In addition, FSH levels were steadily and significantly higher during the follicular phase in the S/S genotype group, whereas no differences were observed between groups with regard to estradiol, inhibin B and growth velocity of the dominant follicle, showing that higher levels of endogenous FSH are necessary to achieve ovulation in carriers of the S/S genotype. Menstrual cycles were significantly longer in women with the S/S genotype, with a difference of about 2 days between these women and those with the N/N genotype. Thus, this study demonstrated that the S/S genotype results in a higher ovarian threshold for FSH, decreased negative feedback to the pituitary and a longer menstrual cycle (Greb *et al.*, 2005). Other studies have observed that the genotype for the codon 680 polymorphism was associated with differences in cycle length (Perez Mayorga *et al.*, 2000), basal FSH levels (Sudo *et al.*, 2002; Laven *et al.*, 2003; Falconer *et al.*, 2005; Jun *et al.*, 2006; Loutradis *et al.*,

Table V. Polymorphisms in genes encoding sex hormones and hormone regulators, enzymes involved in metabolism and biosynthesis, or paracrine factors.

Gene	Locus	Protein name	Protein function	Variant		Association with ovarian response		Phenotype (cases, controls)
				Name	dbSNP ID	Positive (cases, controls)	Negative (cases, controls)	
Sex hormones and hormone regulators								
<i>AMH</i>	19p13.3	Anti-Müllerian hormone	Hormone	p.I49S	rs10407022			Caucasian women—E ₂ levels (53, 45) (Kevenaar <i>et al.</i> , 2007)
<i>AMHR2</i>	12q13	Anti-Müllerian hormone type II receptor	Hormone receptor	−482A/G	rs2002555			Caucasian women—E ₂ levels levels (53, 45) (Kevenaar <i>et al.</i> , 2007)
<i>ESR1</i>	6q25.1	Oestrogen receptor α	Hormone receptor	PvuII RFLP (−397T/C) (g.938T/C)	rs2234693		Spanish women (170) (Mao <i>et al.</i> , 2000; de Castro <i>et al.</i> , 2004)	Caucasian women—follicule/oocyte ratio, pregnancy rate (100, 100) (Georgiou <i>et al.</i> , 1997)
							Chinese women (200, 200) (Sundarajan <i>et al.</i> , 1999)	Chinese women—serum oestradiol levels, follicle/oocyte ratio, pregnancy rate (200, 200) (Sundarajan <i>et al.</i> , 1999)
<i>ESR2</i>	14q23.2	Oestrogen receptor β	Hormone receptor	AluI RFLP (1730A/G) (39A/G)	rs4986938		Spanish women (170) (Mao <i>et al.</i> , 2000; de Castro <i>et al.</i> , 2004)	
<i>FSHR</i>	2p21-p16	Follicle-stimulating hormone receptor	Hormone receptor	p.N680S (in complete LD with p.A307T, rs6165)	rs6166	German women (93) (Perez Mayorga <i>et al.</i> , 2000; Behre <i>et al.</i> , 2005)	Dutch women (105) (Klinkert <i>et al.</i> , 2006)	German women—peak oestradiol levels (93) (Behre <i>et al.</i> , 2005)
						Japanese women (58) (Sudo <i>et al.</i> , 2002)		German women—circulating FSH, levels, number of follicles, luteal phase and menstrual cycle length (23) (Perez Mayorga <i>et al.</i> , 2000; Greb <i>et al.</i> , 2005)
						Spanish women (102) (de Castro <i>et al.</i> , 2003; de Castro <i>et al.</i> , 2004)		Japanese women (58)—basal FSH (Sudo <i>et al.</i> , 2002)
						Korean women (263) (Jun <i>et al.</i> , 2006)		Dutch women (148) (Laven <i>et al.</i> , 2003)
						Greek women (125) (Loutradis <i>et al.</i> , 2006)		Swedish women (68) (Falconer <i>et al.</i> , 2005)
								Korean women—basal FSH, pregnancy rate (263) (Jun <i>et al.</i> , 2006)
								Greek women (125)—FSH levels, follicle and oocyte number (Loutradis <i>et al.</i> , 2006)
								Dutch women—pregnancy rate (105) (Klinkert <i>et al.</i> , 2006)
Enzymes involved in metabolism and biosynthesis								
<i>CYP19A1</i>	15q21.1	Aromatase	Steroid biosynthesis	1672C/T	rs10046			Spanish women (170) (de Castro <i>et al.</i> , 2004)
Paracrine factors								

<i>BMP15</i>	Xp11.2	Bone morphogenic protein 15	Oocyte and follicular development	-673C/T	No dbSNP ID	Spanish women (307) (Moron <i>et al.</i> , 2006) rs3810682	Spanish women (307) (Moron <i>et al.</i> , 2006)
				905G/A	rs3897937	Spanish women (307) (Moron <i>et al.</i> , 2006)	
				308A/G (p.N103S)	rs41308602	Spanish women (307) (Moron <i>et al.</i> , 2006)	
<i>MTHFR</i>	1p36.3	Methylenetetrahydrofolate reductase	Folate metabolism, linked to cardiovascular disease	677C/T (p.A222V)	rs1801133	German women (105) (Thaler <i>et al.</i> , 2006)	German women—E ₂ levels, oocyte number (105) (Thaler <i>et al.</i> , 2006)
				1298A/C (p.E429A)	rs1801131	North American women (223) (Rosen <i>et al.</i> , 2007)	

The list shows genes that have been investigated for their role in ovarian response. Genes are grouped according to their function (and then alphabetically), and studies showing positive or no associations of these genes with ovarian response and/or phenotype (e.g. basal FSH, follicular oestradiol levels) are presented.

2006), pregnancy rates (Jun *et al.*, 2006; Klinkert *et al.*, 2006) and follicular/oocyte number (Loutradis *et al.*, 2006) (Table V).

Behre *et al.* (2005) conducted a prospective interventional study using a randomized controlled trial design and showed that the codon 680 polymorphism of the *FSHR* gene caused a differential estradiol response to FSH. In this study, the same FSH dose for ovarian hyperstimulation resulted in significantly lower serum levels of estradiol in women with the S/S genotype than in women with the N/N genotype. This lower response could be overcome by increasing the FSH dose (Behre *et al.*, 2005) (Fig. 5). Despite differences in estradiol levels, no significant differences were detected in the number of follicles or retrieved oocytes, fertilization rate, cumulative embryo score or pregnancy rate. This finding suggests that, using standard protocols, FSH might be overdosed in individual women, which may put them at risk of OHSS. Indeed, a retrospective association study demonstrated that the *FSHR* S680 allele was represented to a higher degree in women developing OHSS, and that the N680 allele was associated significantly with the severity of OHSS (Daelemans *et al.*, 2004). Since OHSS is relatively rare, none of the studies performed so far in individual centres have sufficient power to demonstrate or refute convincingly any association with the *FSHR* gene; this is an issue that should be analysed retrospectively in a multicentre study.

Taken together, these data indicate that the *FSHR* gene may play a significant role in the success of ovarian stimulation. Women with the *FSHR* S680 allele comprise 60–75% of women undergoing IVF (allelic distribution is similar to that of the general population), and are characterized by higher basal FSH serum concentrations, the need for a higher amount of exogenous FSH and a higher risk of hypo- or hyper-response. Thus, genotyping the *FSHR* p.N680S polymorphism may provide a means of identifying a population of poor responders before IVF procedures are initiated. Since the basis of the poor responder status of these women is a reduced response of the *FSHR* to FSH stimulation, a stimulation protocol designed to overcome the partial resistance to FSH response should be sufficient to improve significantly the success of IVF in these women. The clinical effectiveness of such an approach should be confirmed in randomized controlled trials.

It is important to note that studies in women with ovarian dysfunction did not find any association between *FSHR* polymorphisms and ovarian response to FSH; e.g. in premature ovarian failure (Conway *et al.*, 1999; Sundblad *et al.*, 2004). No significant difference in the FSH dose needed for monofollicular development was detected between women with different *FSHR* genotypes in a non-randomized trial and normogonadotrophic anovulatory women treated with low-dose FSH for ovulation induction (Laven *et al.*, 2003). Therefore, it seems that the differential estradiol response caused by different *FSHR* alleles is evident so far only in women with normal ovarian function. Another study demonstrated that the S680 allele is significantly more frequent in women with a normal menstrual cycle and elevated FSH levels than in women with normal FSH levels, corroborating the idea of a higher FSH threshold (de Koning *et al.*, 2006). Finally, an intriguing association between the A307–S680 haplotype and ovarian cancer susceptibility was recently reported (Yang *et al.*, 2006).

Table VI. Comparison of the study designs and outcomes of two conflicting studies investigating the relation between the FSHR polymorphism at amino acid position 680 (Asn/Ser) and pregnancy rate.

	Jun <i>et al.</i> (2006)	Klinkert <i>et al.</i> (2006)
Number of patients	263	105
Status regarding ICSI patients	Not excluded	Excluded
Frequency of unexplained infertility (%)	35.6	25.0
Genotype associated with significantly higher FSH levels	Ser/Ser	No difference
Genotype associated with significantly higher clinical pregnancy rate	Asn/Asn	Ser/Ser

ICSI, intracytoplasmic sperm injection.

Other genes

In addition to studies investigating the influence of *FSHR* genotype on ovarian responsiveness, other genetic factors have been analysed and reported to be involved in modulating ovarian sensitivity to FSH. Important candidates are polymorphisms in the *ER1* (Georgiou *et al.*, 1997; Sundarajan *et al.*, 1999; de Castro *et al.*, 2004), and *ER2* genes (de Castro *et al.*, 2004) as well as in the anti-Müllerian hormone (AMH) and AMH type II receptor genes (Kevenaar *et al.*, 2007) and in *MTHFR* (Thaler *et al.*, 2006; Rosen *et al.*, 2007). As ovarian responsiveness to FSH may be a polygenic trait, future studies should investigate the combined role of all these factors in large numbers of well-characterized patients undergoing ovarian hyperstimulation, using stringent genetic epidemiological criteria.

Finally, several studies have explored the effects of polymorphisms in genes for metabolic enzymes on ovarian response (Table V). De Castro *et al.* (2004) studied the 1672C/T polymorphism in the *CYP19A1* gene, which encodes an enzyme involved in estrogen synthesis, and found that it does not influence ovarian response to exogenous FSH. Bone morphogenetic protein 15 (BMP-15), a member of the transforming growth factor- β superfamily, is a paracrine factor expressed exclusively in the ovaries and is involved in oocyte and follicular development. A recent study investigated the association between OHSS and polymorphisms in the *BMP15* gene (Moron *et al.*, 2006). Although individual polymorphisms did not show a strong association with the risk of OHSS, the TGGG haplotype was a risk factor and the CCAA haplotype was found to confer protection against OHSS (Moron *et al.*, 2006).


Discussion

This systematic review emphasizes the inconsistency of results obtained by studies that have investigated associations between polymorphisms and PCOS or ovarian response. This may be due to methodological reasons, such as population stratification (genetic heterogeneity, population history), inappropriate disease/trait definition, subject selection issues or chance findings. One of the major problems with candidate gene studies concerns sample size, which is frequently insufficient. The sample sizes of many such studies have been calculated based on knowledge acquired while studying monogenic traits, which have a very strong effect. Unlike monogenetic traits, complex traits result from the interaction of several genetic variations and environmental factors, thus individual genetic variations have a much more modest effect (Lander and Schork, 1994). This may not have been taken into consideration, resulting in studies that are too small to reveal a genetic

association in the context of a complex trait. Furthermore, it has been clearly demonstrated that studies with small populations, those that investigate a genetic variation with a small effect, or those that have a flexible design that is prone to bias are less likely to be replicated (Ioannidis, 2005). Similar conclusions were drawn by Gorroochurn *et al.* (2007), who showed that for commonly observed *P*-value thresholds ($P = 0.02$ – 0.01 , when $\alpha = 0.05$), replication probabilities are surprisingly low (around 60–70% chance of replication).

Inconsistent results may also be due to characteristics inherent to polymorphisms, such as incomplete penetrance, genetic heterogeneity and gene–gene or gene–environment interactions. Other shortcomings of the use of candidate genes/markers to study association include the fact that only a very tiny part of the genome is being studied, and this is done independently of any interactions that may be involved. In addition, candidate gene selection relies on prior knowledge, making it impossible to reveal an association with genes that have unknown function or that have not been known to be implicated in the disease/trait being studied.

To overcome some of these limitations, the use of genome-wide scans is an experimental tool that is becoming an increasingly realistic option. Over the past 5–10 years, refinement of technology involving polymerase chain reaction, development of microarray technology and the remarkable progress in the characterization of the human genome sequence have enabled the study of thousands of DNA variations in a single experiment. Commercial genotyping tools currently allow the study of almost a million SNPs per sample in a single assay, representing roughly 10% of the estimated total number of SNPs in the human genome. Although not all known SNPs are represented on one genotyping microarray, linkage disequilibrium allows nearly 90% of the human genome to be studied with current technology (Schork *et al.*, 2000; Locke *et al.*, 2006; Roeder *et al.*, 2006). Therefore, by performing an assay for a particular SNP, it is also possible to indirectly test for the presence of other variants that are in linkage disequilibrium. Although less advanced, the same technology is also being used to study quantitative (or copy number) gene variation on a genome-wide basis; this type of variation has recently been found to impact a much larger part of the genome than originally thought (Iafrate *et al.*, 2004). Before this technique offers clinically useful data, however, key methodological points must be addressed. These include increasing the sample size compared with single gene/marker studies and resolving statistical problems inherent to multiple testing. Although it is obvious that a number of the studies reviewed had a small sample size, it is quite difficult to give a generic sample size that will fit all studies, as this depends on the frequency and number of the markers being tested and the importance of the genetic effect of the associated marker; genome-



H, luteinizing hormone. *P < 0.05; **P < 0.005.

Figure 4: Menstrual cycle-dependent serum levels of LH (A), FSH (B), estradiol (C), progesterone (D), inhibin A (E), and inhibin B (F) referenced to the day of the LH surge (0) in women with the Asn680/Asn680 (n=12) and the Ser680/Ser680 (n=9) genotype from the luteo-follicular transition phase (LH -25) until ovulation (LH +2). Means are displayed as bars, and error bars show SEM; *P<0.05; **P<0.005.

Serum hormone concentrations during the menstrual cycle in women carrying either N⁶⁸⁰/N⁶⁸⁰ or S⁶⁸⁰/S⁶⁸⁰ allele variants of the N680S polymorphism in the FSH receptor gene [Reproduced with permission from Greb *et al.* (2005), © 2005 The Endocrine Society].




Figure 5: Serum levels of oestradiol before ovulation induction were significantly lower in women with the Ser/Ser allele variant (group I, n=24) compared to the Asn/Asn allele variant (group III, n=44) of the FSH receptor (lower panel: *significant difference between group I and III). This difference in ovarian response could be overcome by increasing the daily FSH dose from 150U/day to 225U/day (upper panel:*significant higher total FSH dose) in women with the Ser/Ser allele variant (group II, n=25); lower panel: no significant difference between group II and III.

Total FSH dose required (upper) and serum estradiol concentration (lower) in normo-ovulatory women undergoing controlled ovarian hyperstimulation, grouped according to N680S genotype for the FSH receptor gene [Reproduced with permission from Behre *et al.* (2005), ©2005 Lippincott Williams & Wilkins].

wide studies are often carried out on more than 1000 individuals. If the association studies have been properly designed, replication of the results in an independent population is one of the best methods to confirm an association.

Regardless of whether a candidate gene or a genome-wide approach is used, we suggest that in order to improve the quality of results, genetic association studies should be carried out in two phases: an exploratory phase, followed by a validation phase. Whenever possible, exploratory studies should be carried

out using a genome-wide approach to maximize the chances of identifying an association. Ideally, these exploratory studies should be replicated, and only markers that are positive in at least two exploratory studies should be considered for validation. As explained previously, the identification of an association between a certain allele or genotype and a given disease/trait does not necessarily mean that there is an aetiological link. Due to the occurrence of linkage disequilibrium, in a number of cases, the associated genetic variation will only be a marker, and not the direct cause, of the trait; for this reason, care must be taken when drawing conclusions from the results. The recommended second phase consists of validating the small set of markers that were associated with the disease/trait in the exploratory studies. Depending on the objective, validation of the markers can take several forms. In some cases, it may be enough to replicate the association with this marker in a different study population (e.g. if a genetic marker is predictive of response to treatment or of disease state). On the other hand, if the aim is to study the impact of genetic variants on molecular mechanisms, then it will be necessary to first either sequence or screen the DNA region around the associated variant with a higher density of markers in order to pinpoint the particular variation that is the causative variation (resulting in, e.g. changes in amino acid sequence or changes to the promoter region affecting gene expression). Once identified, this causative variation will only be truly validated by results of *in vitro* and *in vivo* assays, such as site-directed mutagenesis or reporter gene assays.

On the basis of the data we have summarized, there is currently only one polymorphism for which a sufficient number of studies have consistently identified a significant association. Studies in women from various ethnic backgrounds with normal ovarian function demonstrate convincingly that SNPs in exon 10 of the FSH receptor gene can be used as markers to predict differences in FSHR function and ovarian response to FSH (Perez Mayorga *et al.*, 2000; Sudo *et al.*, 2002; de Castro *et al.*, 2003, 2004; Falconer *et al.*, 2005; Jun *et al.*, 2006; Loutradis *et al.*, 2006), although ovarian response is a polygenic trait and the interaction with other gene polymorphisms remains to be investigated. On the contrary, no consistent association between gene polymorphism and PCOS could be identified.

Conclusions

By pursuing the most promising polymorphisms identified in this review, we hope that the key factors involved in the pathogenesis of PCOS will be identified in the not-too-distant future, and that this knowledge can be used to improve prognosis for women with this disorder. These women are also part of a wider group who would potentially benefit from further analyses to confirm the link between variants in the *FSHR* gene and ovarian response to gonadotrophins. Strong evidence for an association could ultimately lead to stimulation protocols that are carefully personalized to each woman's individual needs, according to the particular combination of polymorphisms inherited.

Acknowledgements

The authors would like to thank Drs Polly Field, Imogen Horsey and Kay Elder for their assistance in drafting the manuscript.

Funding

The preparation of this manuscript was sponsored by an unrestricted educational grant from Merck Serono International S.A., Geneva, Switzerland (an affiliate of Merck KGaA, Darmstadt, Germany). Funding to pay the Open Access publication charges for this article was provided by Merck Serono International S.A., an affiliate of Merck KGaA, Darmstadt, Germany.

References

- Abbott DH, Dumesic DA, Franks S. Developmental origin of polycystic ovary syndrome—a hypothesis. *J Endocrinol* 2002;**174**:1–5.
- Alcoser SY, Hara M, Bell GI, Ehrmann DA. Association of the (AU)AT-rich element polymorphism in PPP1R3 with hormonal and metabolic features of polycystic ovary syndrome. *J Clin Endocrinol Metab* 2004;**89**:2973–2976.
- Antoine HJ, Pall M, Trader BC, Chen YD, Azziz R, Goodarzi MO. Genetic variants in peroxisome proliferator-activated receptor gamma influence insulin resistance and testosterone levels in normal women, but not those with polycystic ovary syndrome. *Fertil Steril* 2007;**87**:862–869.
- Baba T, Endo T, Sata F, Honma H, Kitajima Y, Hayashi T, Manase K, Kanaya M, Yamada H, Minakami H *et al.* Polycystic ovary syndrome is associated with genetic polymorphism in the insulin signaling gene IRS-1 but not ENPP1 in a Japanese population. *Life Sci* 2007;**81**:850–854.
- Babu KA, Rao KL, Kanakavalli MK, Suryanarayana VV, Deenadayal M, Singh L. CYP11A1, GSTM1 and GSTT1 genetic polymorphism is associated with susceptibility to polycystic ovaries in South Indian women. *Reprod Biomed Online* 2004;**9**:194–200.
- Behre HM, Greb RR, Mempel A, Sonntag B, Kiesel L, Kaltwasser P, Seliger E, Ropke F, Gromoll J, Nieschlag E *et al.* Significance of a common single nucleotide polymorphism in exon 10 of the follicle-stimulating hormone (FSH) receptor gene for the ovarian response to FSH: a pharmacogenetic approach to controlled ovarian hyperstimulation. *Pharmacogenet Genomics* 2005;**15**:451–456.
- Bendlova B, Zavadilova J, Vankova M, Vejrazkova D, Lukasova P, Vcelak J, Hill M, Cibula D, Vondra K, Starka L *et al.* Role of D327N sex hormone-binding globulin gene polymorphism in the pathogenesis of polycystic ovary syndrome. *J Steroid Biochem Mol Biol* 2007;**104**:68–74.
- Boomsma CM, Eijkemans MJ, Hughes EG, Visser GH, Fauser BC, Macklon NS. A meta-analysis of pregnancy outcomes in women with polycystic ovary syndrome. *Hum Reprod Update* 2006;**12**:673–683.
- Botella-Carretero JJ, Luque-Ramirez M, varez-Blasco F, San Millan JL, Escobar-Morreale HF. Mutations in the hereditary hemochromatosis gene are not associated with the increased body iron stores observed in overweight and obese women with polycystic ovary syndrome. *Diabetes Care* 2006;**29**:2556.
- Calvo RM, Villuendas G, Sancho J, San Millan JL, Escobar-Morreale HF. Role of the follistatin gene in women with polycystic ovary syndrome. *Fertil Steril* 2001;**75**:1020–1023.
- Calvo RM, Telleria D, Sancho J, San Millan JL, Escobar-Morreale HF. Insulin gene variable number of tandem repeats regulatory polymorphism is not associated with hyperandrogenism in Spanish women. *Fertil Steril* 2002;**77**:666–668.
- Chen ZJ, Shi YH, Zhao YR, Li Y, Tang R, Zhao LX, Chang ZH. Correlation between single nucleotide polymorphism of insulin receptor gene with polycystic ovary syndrome. *Zhonghua Fu Chan Ke Za Zhi* 2004;**39**:582–585.
- Conway GS, Honour JW, Jacobs HS. Heterogeneity of the polycystic ovary syndrome: clinical, endocrine and ultrasound features in 556 patients. *Clin Endocrinol (Oxf)* 1989;**30**:459–470.
- Conway GS, Conway E, Walker C, Hoppner W, Gromoll J, Simoni M. Mutation screening and isoform prevalence of the follicle stimulating hormone receptor gene in women with premature ovarian failure, resistant ovary syndrome and polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 1999;**51**:97–99.
- Daelemans C, Smits G, de M V, Costagliola S, Englert Y, Vassart G, Delbaere A. Prediction of severity of symptoms in iatrogenic ovarian hyperstimulation syndrome by follicle-stimulating hormone receptor Ser680Asn polymorphism. *J Clin Endocrinol Metab* 2004;**89**:6310–6315.
- de Castro F, Ruiz R, Montoro L, Perez-Hernandez D, Sanchez-Casas PE, Real LM, Ruiz A. Role of follicle-stimulating hormone receptor Ser680Asn polymorphism in the efficacy of follicle-stimulating hormone. *Fertil Steril* 2003;**80**:571–576.
- de Castro F, Moron FJ, Montoro L, Galan JJ, Hernandez DP, Padilla ES, Ramirez-Lorca R, Real LM, Ruiz A. Human controlled ovarian hyperstimulation outcome is a polygenic trait. *Pharmacogenetics* 2004;**14**:285–293.
- de Koning CH, Benjamins T, Harms P, Homburg R, van Montfrans JM, Gromoll J, Simoni M, Lambalk CB. The distribution of FSH receptor isoforms is related to basal FSH levels in subfertile women with normal menstrual cycles. *Hum Reprod* 2006;**21**:443–446.
- Diamanti-Kandarakis E, Bartzis MI, Zapanti ED, Spina GG, Filandra FA, Tsianateli TC, Bergiele AT, Kouli CR. Polymorphism T→C (–34 bp) of gene CYP17 promoter in Greek patients with polycystic ovary syndrome. *Fertil Steril* 1999;**71**:431–435.
- Diamanti-Kandarakis E, Bartzis MI, Bergiele AT, Tsianateli TC, Kouli CR. Microsatellite polymorphism (ttta)(n) at –528 base pairs of gene CYP11alpha influences hyperandrogenemia in patients with polycystic ovary syndrome. *Fertil Steril* 2000;**73**:735–741.
- Diamanti-Kandarakis E, Palioniko G, Alexandraki K, Bergiele A, Koutsouba T, Bartzis M. The prevalence of 4G5G polymorphism of plasminogen activator inhibitor-1 (PAI-1) gene in polycystic ovarian syndrome and its association with plasma PAI-1 levels. *Eur J Endocrinol* 2004;**150**:793–798.
- Diamanti-Kandarakis E, Piperi C. Genetics of polycystic ovary syndrome: searching for the way out of the labyrinth. *Hum Reprod Update* 2005;**11**:631–643.
- Dilek S, Ertunc D, Tok EC, Erdal EM, Aktas A. Association of Gly972Arg variant of insulin receptor substrate-1 with metabolic features in women with polycystic ovary syndrome. *Fertil Steril* 2005;**84**:407–412.
- Draper N, Powell BL, Franks S, Conway GS, Stewart PM, McCarthy MI. Variants implicated in cortisone reductase deficiency do not contribute to susceptibility to common forms of polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 2006;**65**:64–70.
- Ehrmann DA. Polycystic ovary syndrome. *N Engl J Med* 2005;**352**:1223–1236.
- Ehrmann DA, Schwarz PE, Hara M, Tang X, Horikawa Y, Imperial J, Bell GI, Cox NJ. Relationship of calpain-10 genotype to phenotypic features of polycystic ovary syndrome. *J Clin Endocrinol Metab* 2002a;**87**:1669–1673.
- Ehrmann DA, Tang X, Yoshiuchi I, Cox NJ, Bell GI. Relationship of insulin receptor substrate-1 and –2 genotypes to phenotypic features of polycystic ovary syndrome. *J Clin Endocrinol Metab* 2002b;**87**:4297–4300.
- El Mkaed SA, Lautier C, Macari F, Molinari N, Lefebvre P, Renard E, Gris JC, Cros G, Daures JP, Bringer J *et al.* Role of allelic variants Gly972Arg of IRS-1 and Gly1057Asp of IRS-2 in moderate-to-severe insulin resistance of women with polycystic ovary syndrome. *Diabetes* 2001;**50**:2164–2168.
- Elter K, Erel CT, Cine N, Ozbek U, Hacıhanefioglu B, Ertungealp E. Role of the mutations Trp8=> Arg and Ile15=> Thr of the human luteinizing hormone beta-subunit in women with polycystic ovary syndrome. *Fertil Steril* 1999;**71**:425–430.
- Ertunc D, Tok EC, Aktas A, Erdal EM, Dilek S. The importance of IRS-1 Gly972Arg polymorphism in evaluating the response to metformin treatment in polycystic ovary syndrome. *Hum Reprod* 2005;**20**:1207–1212.
- Escobar-Morreale HF, Peral B, Villuendas G, Calvo RM, Sancho J, San Millan JL. Common single nucleotide polymorphisms in intron 3 of the calpain-10 gene influence hirsutism. *Fertil Steril* 2002;**77**:581–587.
- Escobar-Morreale HF, Calvo RM, Villuendas G, Sancho J, San Millan JL. Association of polymorphisms in the interleukin 6 receptor complex with obesity and hyperandrogenism. *Obes Res* 2003;**11**:987–996.
- Escobar-Morreale HF, Luque-Ramirez M, San Millan JL. The molecular-genetic basis of functional hyperandrogenism and the polycystic ovary syndrome. *Endocr Rev* 2005;**26**:251–282.
- Falconer H, Andersson E, Aanesen A, Fried G. Follicle-stimulating hormone receptor polymorphisms in a population of infertile women. *Acta Obstet Gynecol Scand* 2005;**84**:806–811.
- Fauser BC, Van Heusden AM. Manipulation of human ovarian function: physiological concepts and clinical consequences. *Endocr Rev* 1997;**18**:71–106.
- Fauser BC, Diedrich K, Devroey P. Predictors of ovarian response: progress towards individualized treatment in ovulation induction and ovarian stimulation. *Hum Reprod Update* 2008;**14**:1–14.
- Ferk P, Teran N, Gersak K. The (TAAAA)n microsatellite polymorphism in the SHBG gene influences serum SHBG levels in women with polycystic ovary syndrome. *Hum Reprod* 2007;**22**:1031–1036.

- Franks S, Gharani N, McCarthy M. Genetic abnormalities in polycystic ovary syndrome. *Ann Endocrinol (Paris)* 1999;**60**:131–133.
- Fratantonio E, Vicari E, Pafumi C, Calogero AE. Genetics of polycystic ovarian syndrome. *Reprod Biomed Online* 2005;**10**:713–720.
- Gaasenbeek M, Powell BL, Sovio U, Haddad L, Gharani N, Bennett A, Groves CJ, Rush K, Goh MJ, Conway GS *et al.* Large-scale analysis of the relationship between CYP11A promoter variation, polycystic ovarian syndrome, and serum testosterone. *J Clin Endocrinol Metab* 2004;**89**:2408–2413.
- Georgiou I, Konstantelli M, Syrou M, Messinis IE, Lolis DE. Oestrogen receptor gene polymorphisms and ovarian stimulation for in-vitro fertilization. *Hum Reprod* 1997;**12**:1430–1433.
- Gharani N, Waterworth DM, Batty S, White D, Gilling-Smith C, Conway GS, McCarthy M, Franks S, Williamson R. Association of the steroid synthesis gene CYP11a with polycystic ovary syndrome and hyperandrogenism. *Hum Mol Genet* 1997;**6**:397–402.
- Glueck CJ, Wang P, Fontaine RN, Sieve-Smith L, Tracy T, Moore SK. Plasminogen activator inhibitor activity: an independent risk factor for the high miscarriage rate during pregnancy in women with polycystic ovary syndrome. *Metabolism* 1999;**48**:1589–1595.
- Goldman S, Shalev E. MMPs and TIMPs in ovarian physiology and pathophysiology. *Front Biosci* 2004;**9**:2474–2483.
- Gonzalez A, Abril E, Roca A, Aragon MJ, Figueroa MJ, Velarde P, Ruiz R, Faye O, Galan JJ, Herreros JA *et al.* Specific CAPN10 gene haplotypes influence the clinical profile of polycystic ovary patients. *J Clin Endocrinol Metab* 2003;**88**:5529–5536.
- Goodarzi MO, Shah NA, Antoine HJ, Pall M, Guo X, Azziz R. Variants in the 5 α -reductase type 1 and type 2 genes are associated with polycystic ovary syndrome and the severity of hirsutism in affected women. *J Clin Endocrinol Metab* 2006;**91**:4085–4091.
- Gorroochurn P, Hodge SE, Heiman GA, Durner M, Greenberg DA. Non-replication of association studies: 'pseudo-failures' to replicate? *Genet Med* 2007;**9**:325–331.
- Greb RR, Grieshaber K, Gromoll J, Sonntag B, Nieschlag E, Kiesel L, Simoni M. A common single nucleotide polymorphism in exon 10 of the human follicle stimulating hormone receptor is a major determinant of length and hormonal dynamics of the menstrual cycle. *J Clin Endocrinol Metab* 2005;**90**:4866–4872.
- Haap M, Machicao F, Stefan N, Thamer C, Tschrutter O, Schnuck F, Wallwiener D, Stumvoll M, Haring HU, Fritsche A. Genetic determinants of insulin action in polycystic ovary syndrome. *Exp Clin Endocrinol Diabetes* 2005;**113**:275–281.
- Hahn S, Fingerhut A, Khomtsiv U, Khomtsiv L, Tan S, Quadbeck B, Herrmann BL, Knebel B, Muller-Wieland D, Mann K *et al.* The peroxisome proliferator activated receptor gamma Pro12Ala polymorphism is associated with a lower hirsutism score and increased insulin sensitivity in women with polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 2005;**62**:573–579.
- Hahn S, Frey UH, Siffert W, Tan S, Mann K, Janssen OE. The CC genotype of the GNAS T393C polymorphism is associated with obesity and insulin resistance in women with polycystic ovary syndrome. *Eur J Endocrinol* 2006;**155**:763–770.
- Hara M, Alcoser SY, Qadir A, Beiswenger KK, Cox NJ, Ehrmann DA. Insulin resistance is attenuated in women with polycystic ovary syndrome with the Pro(12)Ala polymorphism in the PPARgamma gene. *J Clin Endocrinol Metab* 2002;**87**:772–775.
- Heinonen S, Korhonen S, Hippelainen M, Hiltunen M, Mannermaa A, Saarikoski S. Apolipoprotein E alleles in women with polycystic ovary syndrome. *Fertil Steril* 2001;**75**:878–880.
- Heinonen S, Korhonen S, Helisalmi S, Koivunen R, Tapanainen JS, Laakso M. The 121Q allele of the plasma cell membrane glycoprotein 1 gene predisposes to polycystic ovary syndrome. *Fertil Steril* 2004;**82**:743–745.
- Heinonen S, Korhonen S, Helisalmi S, Koivunen R, Tapanainen J, Hippelainen M, Laakso M. Associations between two single nucleotide polymorphisms in the adiponectin gene and polycystic ovary syndrome. *Gynecol Endocrinol* 2005;**21**:165–169.
- Hickey T, Chandy A, Norman RJ. The androgen receptor CAG repeat polymorphism and X-chromosome inactivation in Australian Caucasian women with infertility related to polycystic ovary syndrome. *J Clin Endocrinol Metab* 2002;**87**:161–165.
- Iafate AJ, Feuk L, Rivera MN, Listewnik ML, Donahoe PK, Qi Y, Scherer SW, Lee C. Detection of large-scale variation in the human genome. *Nat Genet* 2004;**36**:949–951.
- Ioannidis JP. Why most published research findings are false. *PLoS Med* 2005;**2**:e124.
- Jaaskelainen J, Korhonen S, Voutilainen R, Hippelainen M, Heinonen S. Androgen receptor gene CAG length polymorphism in women with polycystic ovary syndrome. *Fertil Steril* 2005;**83**:1724–1728.
- Jahanfar S, Eden JA. Genetic and non-genetic theories on the etiology of polycystic ovary syndrome. *Gynecol Endocrinol* 1996;**10**:357–364.
- Jin L, Zhu XM, Luo Q, Qian Y, Jin F, Huang HF. A novel SNP at exon 17 of INSR is associated with decreased insulin sensitivity in Chinese women with PCOS. *Mol Hum Reprod* 2006;**12**:151–155.
- Jones MR, Italiano L, Wilson SG, Mullin BH, Mead R, Dudbridge F, Watts GF, Stuckey BG. Polymorphism in HSD17B6 is associated with key features of polycystic ovary syndrome. *Fertil Steril* 2006;**86**:1438–1446.
- Jones MR, Wilson SG, Mullin BH, Mead R, Watts GF, Stuckey BG. Polymorphism of the follistatin gene in polycystic ovary syndrome. *Mol Hum Reprod* 2007;**13**:237–241.
- Jun JK, Yoon JS, Ku SY, Choi YM, Hwang KR, Park SY, Lee GH, Lee WD, Kim SH, Kim JG *et al.* Follicle-stimulating hormone receptor gene polymorphism and ovarian responses to controlled ovarian hyperstimulation for IVF-ET. *J Hum Genet* 2006;**51**:665–670.
- Kahsar-Miller M, Boots LR, Azziz R. Dopamine D3 receptor polymorphism is not associated with the polycystic ovary syndrome. *Fertil Steril* 1999;**71**:436–438.
- Kaibe M, Takakuwa K, Murakawa H, Ishii K, Tamura M, Tanaka K. Studies on the human leukocyte antigens in patients with polycystic ovary syndrome in a Japanese population—possible susceptibility of HLA-A11 and -DRB1*0403 to patient population with polycystic ovary syndrome. *Am J Reprod Immunol* 2006;**55**:301–306.
- Karadeniz M, Erdogan M, Berdeli A, Saygili F, Yilmaz C. 4G/5G Polymorphism of PAI-1 gene and Alu-repeat I/D polymorphism of TPA gene in Turkish patients with polycystic ovary syndrome. *J Assist Reprod Genet* 2007;**24**:412–418.
- Kevenaar ME, Themmen AP, Laven JS, Sonntag B, Fong SL, Uitterlinden AG, de Jong FH, Pols HA, Simoni M, Visser JA. Anti-Mullerian hormone and anti-Mullerian hormone type II receptor polymorphisms are associated with follicular phase estradiol levels in normo-ovulatory women. *Hum Reprod* 2007;**22**:1547–1554.
- Kim JY, Moon SM, Ryu HJ, Kim JJ, Kim HT, Park C, Kimm K, Oh B, Lee JK. Identification of regulatory polymorphisms in the TNF-TNF receptor superfamily. *Immunogenetics* 2005;**57**:297–303.
- Klinkert ER, te Velde ER, Weima S, van Zandvoort PM, Hanssen RG, Nilsson PR, de Jong FH, Looman CW, Broekmans FJ. FSH receptor genotype is associated with pregnancy but not with ovarian response in IVF. *Reprod Biomed Online* 2006;**13**:687–695.
- Kolbus A, Walch K, Nagele F, Wenzl R, Unfried G, Huber JC. Interleukin-1 alpha but not interleukin-1 beta gene polymorphism is associated with polycystic ovary syndrome. *J Reprod Immunol* 2007;**73**:188–193.
- Korhonen S, Romppanen EL, Hiltunen M, Mannermaa A, Punnonen K, Hippelainen M, Heinonen S. Lack of association between C-850T polymorphism of the gene encoding tumor necrosis factor-alpha and polycystic ovary syndrome. *Gynecol Endocrinol* 2002;**16**:271–274.
- Korhonen S, Heinonen S, Hiltunen M, Helisalmi S, Hippelainen M, Koivunen R, Tapanainen JS, Laakso M. Polymorphism in the peroxisome proliferator-activated receptor-gamma gene in women with polycystic ovary syndrome. *Hum Reprod* 2003a;**18**:540–543.
- Korhonen S, Romppanen EL, Hiltunen M, Helisalmi S, Punnonen K, Hippelainen M, Heinonen S. Two exonic single nucleotide polymorphisms in the microsomal epoxide hydrolase gene are associated with polycystic ovary syndrome. *Fertil Steril* 2003b;**79**:1353–1357.
- Kurabayashi T, Yahata T, Quan J, Tanaka K. Association of polymorphisms in the beta2 and beta3 adrenoceptor gene with polycystic ovary syndrome. *J Reprod Med* 2006;**51**:389–393.
- Lander ES, Schork NJ. Genetic dissection of complex traits. *Science* 1994;**265**:2037–2048.
- Laven JS, Imani B, Eijkemans MJ, Fauser BC. New approach to polycystic ovary syndrome and other forms of anovulatory infertility. *Obstet Gynecol Surv* 2002;**57**:755–767.
- Laven JS, Mulders AG, Suryandari DA, Gromoll J, Nieschlag E, Fauser BC, Simoni M. Follicle-stimulating hormone receptor polymorphisms in women with normogonadotropic anovulatory infertility. *Fertil Steril* 2003;**80**:986–992.
- Lee EJ, Oh B, Lee JY, Kimm K, Lee SH, Baek KH. A novel single nucleotide polymorphism of INSR gene for polycystic ovary syndrome. *Fertil Steril* 2007 (in press).
- Lee EJ, Yoo KJ, Kim SJ, Lee SH, Cha KY, Baek KH. Single nucleotide polymorphism in exon 17 of the insulin receptor gene is not associated

- with polycystic ovary syndrome in a Korean population. *Fertil Steril* 2006;**86**:380–384.
- Locke DP, Sharp AJ, McCarroll SA, McGrath SD, Newman TL, Cheng Z, Schwartz S, Albertson DG, Pinkel D, Altshuler DM *et al.* Linkage disequilibrium and heritability of copy-number polymorphisms within duplicated regions of the human genome. *Am J Hum Genet* 2006;**79**:275–290.
- Loutradis D, Patsoula E, Minas V, Koussidis GA, Antsaklis A, Michalas S, Makrigiannakis A. FSH receptor gene polymorphisms have a role for different ovarian response to stimulation in patients entering IVF/ICSI-ET programs. *J Assist Reprod Genet* 2006;**23**:177–184.
- Mao W, Yu L, Chen Y. Study on the relationship between a polymorphism of tumor necrosis factor- α gene and the pathogenesis of polycystic ovary syndrome. *Zhonghua Fu Chan Ke Za Zhi* 2000;**35**:536–539.
- Marszalek B, Lacinski M, Babych N, Capla E, Biernacka-Lukanty J, Warenik-Szymankiewicz A, Trzeciak WH. Investigations on the genetic polymorphism in the region of CYP17 gene encoding 5'-UTR in patients with polycystic ovarian syndrome. *Gynecol Endocrinol* 2001;**15**:123–128.
- Michelmore K, Ong K, Mason S, Bennett S, Perry L, Vessey M, Balen A, Dunger D. Clinical features in women with polycystic ovaries: relationships to insulin sensitivity, insulin gene VNTR and birth weight. *Clin Endocrinol (Oxf)* 2001;**55**:439–446.
- Mifsud A, Ramirez S, Yong EL. Androgen receptor gene CAG trinucleotide repeats in anovulatory infertility and polycystic ovaries. *J Clin Endocrinol Metab* 2000;**85**:3484–3488.
- Milner CR, Craig JE, Hussey ND, Norman RJ. No association between the -308 polymorphism in the tumour necrosis factor alpha (TNF α) promoter region and polycystic ovaries. *Mol Hum Reprod* 1999;**5**:5–9.
- Mohlig M, Spranger J, Osterhoff M, Ristow M, Pfeiffer AF, Schill T, Schlosser HW, Brabant G, Schöf C. The polycystic ovary syndrome per se is not associated with increased chronic inflammation. *Eur J Endocrinol* 2004;**150**:525–532.
- Moron FJ, de CF, Royo JL, Montoro L, Mira E, Saez ME, Real LM, Gonzalez A, Manes S, Ruiz A. Bone morphogenetic protein 15 (BMP15) alleles predict over-response to recombinant follicle stimulation hormone and iatrogenic ovarian hyperstimulation syndrome (OHSS). *Pharmacogenet Genomics* 2006;**16**:485–495.
- Oksanen L, Tiitinen A, Kaprio J, Koistinen HA, Karonen S, Kontula K. No evidence for mutations of the leptin or leptin receptor genes in women with polycystic ovary syndrome. *Mol Hum Reprod* 2000;**6**:873–876.
- Orio F, Jr, Matarese G, Di BS, Palomba S, Labella D, Sanna V, Savastano S, Zullo F, Colao A, Lombardi G. Exon 6 and 2 peroxisome proliferator-activated receptor- γ polymorphisms in polycystic ovary syndrome. *J Clin Endocrinol Metab* 2003a;**88**:5887–5892.
- Orio F, Jr, Palomba S, Di BS, Colao A, Tauchmanova L, Savastano S, Labella D, Russo T, Zullo F, Lombardi G. Homocysteine levels and C677T polymorphism of methylenetetrahydrofolate reductase in women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 2003b;**88**:673–679.
- Panidis D, Kourtis A, Kukuviitis A, Farmakiotis D, Xita N, Georgiou I, Tsatsoulis A. Association of the T45G polymorphism in exon 2 of the adiponectin gene with polycystic ovary syndrome: role of Delta4-androstenedione. *Hum Reprod* 2004;**19**:1728–1733.
- Peral B, San Millan JL, Castello R, Moghetti P, Escobar-Morreale HF. Comment: the methionine 196 arginine polymorphism in exon 6 of the TNF receptor 2 gene (TNFRSF1B) is associated with the polycystic ovary syndrome and hyperandrogenism. *J Clin Endocrinol Metab* 2002;**87**:3977–3983.
- Perez Mayorga M, Gromoll J, Behre HM, Gassner C, Nieschlag E, Simoni M. Ovarian response to follicle-stimulating hormone (FSH) stimulation depends on the FSH receptor genotype. *J Clin Endocrinol Metab* 2000;**85**:3365–3369.
- Perez-Bravo F, Echiburru B, Maliqueo M, Santos JL, Sir-Petermann T. Tryptophan 64 \rightarrow arginine polymorphism of beta-3-adrenergic receptor in Chilean women with polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 2005;**62**:126–131.
- Petry CJ, Ong KK, Michelmore KF, Artigas S, Wingate DL, Balen AH, de ZF, Ibanez L, Dunger DB. Association of aromatase (CYP 19) gene variation with features of hyperandrogenism in two populations of young women. *Hum Reprod* 2005;**20**:1837–1843.
- Powell BL, Haddad L, Bennett A, Gharani N, Sovio U, Groves CJ, Rush K, Goh MJ, Conway GS, Ruokonen A *et al.* Analysis of multiple data sets reveals no association between the insulin gene variable number tandem repeat element and polycystic ovary syndrome or related traits. *J Clin Endocrinol Metab* 2005;**90**:2988–2993.
- Qin K, Ehrmann DA, Cox N, Refetoff S, Rosenfield RL. Identification of a functional polymorphism of the human type 5 β -hydroxysteroid dehydrogenase gene associated with polycystic ovary syndrome. *J Clin Endocrinol Metab* 2006;**91**:270–276.
- Rajkhowa M, Talbot JA, Jones PW, Pettersson K, Haavisto AM, Huhtaniemi I, Clayton RN. Prevalence of an immunological LH beta-subunit variant in a UK population of healthy women and women with polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 1995;**43**:297–303.
- Rajkhowa M, Talbot JA, Jones PW, Clayton RN. Polymorphism of glycogen synthetase gene in polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 1996;**44**:85–90.
- Roeder K, Bacanu SA, Wasserman L, Devlin B. Using linkage genome scans to improve power of association in genome scans. *Am J Hum Genet* 2006;**78**:243–252.
- Rosen MP, Shen S, McCulloch CE, Rinaudo PF, Cedars MI, Dobson AT. Methylenetetrahydrofolate reductase (MTHFR) is associated with ovarian follicular activity. *Fertil Steril* 2007;**88**:632–638.
- San Millan JL, Corton M, Villuendas G, Sancho J, Peral B, Escobar-Morreale HF. Association of the polycystic ovary syndrome with genomic variants related to insulin resistance, type 2 diabetes mellitus, and obesity. *J Clin Endocrinol Metab* 2004;**89**:2640–2646.
- San Millan JL, Botella-Carretero JI, varez-Blasco F, Luque-Ramirez M, Sancho J, Moghetti P, Escobar-Morreale HF. A study of the hexose-6-phosphate dehydrogenase gene R453Q and 11 β -hydroxysteroid dehydrogenase type 1 gene 83557insA polymorphisms in the polycystic ovary syndrome. *J Clin Endocrinol Metab* 2005;**90**:4157–4162.
- Schork NJ, Nath SK, Fallin D, Chakravarti A. Linkage disequilibrium analysis of biallelic DNA markers, human quantitative trait loci, and threshold-defined case and control subjects. *Am J Hum Genet* 2000;**67**:1208–1218.
- Siegel S, Futterweit W, Davies TF, Concepcion ES, Greenberg DA, Villanueva R, Tomer Y. A C/T single nucleotide polymorphism at the tyrosine kinase domain of the insulin receptor gene is associated with polycystic ovary syndrome. *Fertil Steril* 2002;**78**:1240–1243.
- Simoni M, Gromoll J, Höppner W, Kamischke A, Krafft T, Stähle D, Nieschlag E. Mutational analysis of the follicle-stimulating hormone (FSH) receptor in normal and infertile men: identification and characterization of two discrete FSH receptor isoforms. *J Clin Endocrinol Metab* 1999;**84**:751–755.
- Sir-Petermann T, Angel B, Maliqueo M, Santos JL, Riesco MV, Toloza H, Perez-Bravo F. Insulin secretion in women who have polycystic ovary syndrome and carry the Gly972Arg variant of insulin receptor substrate-1 in response to a high-glycemic or low-glycemic carbohydrate load. *Nutrition* 2004;**20**:905–910.
- Stewart DR, Dombroski BA, Urbanek M, Ankener W, Ewens KG, Wood JR, Legro RS, Strauss JF, III, Dunaif A, Spielman RS. Fine mapping of genetic susceptibility to polycystic ovary syndrome on chromosome 19p13.2 and tests for regulatory activity. *J Clin Endocrinol Metab* 2006;**91**:4112–4117.
- Sudo S, Kudo M, Wada S, Sato O, Hsueh AJ, Fujimoto S. Genetic and functional analyses of polymorphisms in the human FSH receptor gene. *Mol Hum Reprod* 2002;**8**:893–899.
- Sundararajan C, Liao W, Roy AC, Ng SC. Association of oestrogen receptor gene polymorphisms with outcome of ovarian stimulation in patients undergoing IVF. *Mol Hum Reprod* 1999;**5**:797–802.
- Sundblad V, Chiauzzi VA, Escobar ME, Dain L, Charreau EH. Screening of FSH receptor gene in Argentine women with premature ovarian failure (POF). *Mol Cell Endocrinol* 2004;**222**:53–59.
- Techaraisak K, Conway GS, Rumsby G. Frequency of a polymorphism in the regulatory region of the 17 α -hydroxylase-17,20-lyase (CYP17) gene in hyperandrogenic states. *Clin Endocrinol (Oxf)* 1997;**46**:131–134.
- Thaler CJB, Budiman H, Ruebsamen H, Nagel D, Lohse P. Effects of the common 677C>T mutation of the 5,10-methylenetetrahydrofolate reductase (MTHFR) gene on ovarian responsiveness to recombinant follicle-stimulating hormone. *Am J Reprod Immunol* 2006;**55**:251–258.
- The Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome. *Fertil Steril* 2004a;**81**:19–25.
- The Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome (PCOS). *Hum Reprod* 2004b;**19**:41–47.
- Tok EC, Aktas A, Ertunc D, Erdal EM, Dilek S. Evaluation of glucose metabolism and reproductive hormones in polycystic ovary syndrome

- on the basis of peroxisome proliferator-activated receptor (PPAR)-gamma2 Pro12Ala genotype. *Hum Reprod* 2005;**20**:1590–1595.
- Tong Y, Liao WX, Roy AC, Ng SC. Association of AccI polymorphism in the follicle-stimulating hormone beta gene with polycystic ovary syndrome. *Fertil Steril* 2000;**74**:1233–1236.
- Tong Y, Liao WX, Roy AC, Ng SC. Absence of mutations in the coding regions of follicle-stimulating hormone receptor gene in Singapore Chinese women with premature ovarian failure and polycystic ovary syndrome. *Horm Metab Res* 2001;**33**:221–226.
- Urbanek M, Legro RS, Driscoll D, Strauss JF, III, Dunaif A, Spielman RS. Searching for the polycystic ovary syndrome genes. *J Pediatr Endocrinol Metab* 2000a;**13**(Suppl 5):1311–1313.
- Urbanek M, Wu X, Vickery KR, Kao LC, Christenson LK, Schneyer A, Legro RS, Driscoll DA, Strauss JF, III, Dunaif A *et al.* Allelic variants of the follistatin gene in polycystic ovary syndrome. *J Clin Endocrinol Metab* 2000b;**85**:4455–4461.
- Urbanek M, Woodroffe A, Ewens KG, amanti-Kandarakis E, Legro RS, Strauss JF, III, Dunaif A, Spielman RS. Candidate gene region for polycystic ovary syndrome on chromosome 19p13.2. *J Clin Endocrinol Metab* 2005;**90**:6623–6629.
- Urbanek M, Du Y, Silander K, Collins FS, Stepan CM, Strauss JF, III, Dunaif A, Spielman RS, Legro RS. Variation in resistin gene promoter not associated with polycystic ovary syndrome. *Diabetes* 2003;**52**:214–217.
- Vankova M, Vrbikova J, Hill M, Cinek O, Bendlova B. Association of insulin gene VNTR polymorphism with polycystic ovary syndrome. *Ann N Y Acad Sci* 2002;**967**:558–565.
- Villuendas G, San Millan JL, Sancho J, Escobar-Morreale HF. The -597 G->A and -174 G->C polymorphisms in the promoter of the IL-6 gene are associated with hyperandrogenism. *J Clin Endocrinol Metab* 2002;**87**:1134–1141.
- Villuendas G, Botella-Carretero JI, Roldan B, Sancho J, Escobar-Morreale HF, San Millan JL. Polymorphisms in the insulin receptor substrate-1 (IRS-1) gene and the insulin receptor substrate-2 (IRS-2) gene influence glucose homeostasis and body mass index in women with polycystic ovary syndrome and non-hyperandrogenic controls. *Hum Reprod* 2005;**20**:3184–3191.
- Walch K, Grimm C, Zeillinger R, Huber JC, Nagele F, Hefler LA. A common interleukin-6 gene promoter polymorphism influences the clinical characteristics of women with polycystic ovary syndrome. *Fertil Steril* 2004;**81**:1638–1641.
- Walch K, Grimm C, Huber JC, Nagele F, Kolbus A, Hefler LA. A polymorphism of the plasminogen activator inhibitor-1 gene promoter and the polycystic ovary syndrome. *Eur J Obstet Gynecol Reprod Biol* 2005a;**123**:77–81.
- Walch K, Nagele F, Zeillinger R, Vytiska-Binstorfer E, Huber JC, Hefler LA. A polymorphism in the matrix metalloproteinase-1 gene promoter is associated with the presence of polycystic ovary syndrome in Caucasian women. *Fertil Steril* 2005b;**83**:1565–1567.
- Wang D, Johnson AD, Papp AC, Kroetz DL, Sadee W. Multidrug resistance polypeptide 1 (MDR1, ABCB1) variant 3435C>T affects mRNA stability. *Pharmacogenet Genomics* 2005;**15**:693–704.
- Wang Y, Wu X, Cao Y, Yi L, Chen J. A microsatellite polymorphism (tttta)_n in the promoter of the CYP11a gene in Chinese women with polycystic ovary syndrome. *Fertil Steril* 2006a;**86**:223–226.
- Wang Y, Wu X, Cao Y, Yi L, Fan H, Chen J. Polymorphisms of the peroxisome proliferator-activated receptor-gamma and its coactivator-1alpha genes in Chinese women with polycystic ovary syndrome. *Fertil Steril* 2006b;**85**:1536–1540.
- Witchel SF, Kahsar-Miller M, Aston CE, White C, Azziz R. Prevalence of CYP21 mutations and IRS1 variant among women with polycystic ovary syndrome and adrenal androgen excess. *Fertil Steril* 2005;**83**:371–375.
- Xita N, Tsatsoulis A, Chatzikiyriakidou A, Georgiou I. Association of the (TAAAA)_n repeat polymorphism in the sex hormone-binding globulin (SHBG) gene with polycystic ovary syndrome and relation to SHBG serum levels. *J Clin Endocrinol Metab* 2003;**88**:5976–5980.
- Xita N, Georgiou I, Tsatsoulis A, Kourtis A, Kukuvtis A, Panidis D. A polymorphism in the resistin gene promoter is associated with body mass index in women with polycystic ovary syndrome. *Fertil Steril* 2004;**82**:1466–1467.
- Xita N, Georgiou I, Tsatzikiyriakidou A, Vounatsou M, Papassotiriou GP, Papassotiriou I, Tsatsoulis A. Effect of adiponectin gene polymorphisms on circulating adiponectin and insulin resistance indexes in women with polycystic ovary syndrome. *Clin Chem* 2005;**51**:416–423.
- Yang CQ, Chan KY, Ngan HY, Khoo US, Chiu PM, Chan QK, Xue WC, Cheung AN. Single nucleotide polymorphisms of follicle-stimulating hormone receptor are associated with ovarian cancer susceptibility. *Carcinogenesis* 2006;**27**:1502–1506.
- Zhao SP, Tang XM, Shao DH, Dai HY, Dai SZ. Association study between a polymorphism of aldosterone synthetase gene and the pathogenesis of polycystic ovary syndrome. *Zhonghua Fu Chan Ke Za Zhi* 2003;**38**:94–97.
- Zulian E, Sartorato P, Schiavi F, Moghetti P, Castello R, Mantero F, Opocher G, Scaroni C. The M235T polymorphism of the angiotensinogen gene in women with polycystic ovary syndrome. *Fertil Steril* 2005;**84**:1520–1521.

Submitted on May 7, 2007; resubmitted on May 2, 2008; accepted on May 30, 2008