

Functional genetic variation of human miRNAs and phenotypic consequences

BOREL, Christelle, ANTONARAKIS, Stylianos

Abstract

A large number of human protein-coding genes are finely regulated by one or more microRNAs. Members of this small noncoding RNA family have emerged as important post-transcriptional regulators of gene expression and are involved in a number of disease phenotypes. Variability in the human genome is extensive and includes the common and rare single nucleotide polymorphisms (SNPs) and copy number variations (CNVs). The functional significance of the genome's variability is under intense investigation. In this article we review the emerging literature on how human genomic variation influences the outcome of microRNA targeting and the associated phenotypic effects. Illustrative examples are discussed that demonstrate the biological importance of functional polymorphisms affecting miRNA-mediated gene regulation.

Reference

BOREL, Christelle, ANTONARAKIS, Stylianos. Functional genetic variation of human miRNAs and phenotypic consequences. *Mammalian genome*, 2008, vol. 19, no. 7-8, p. 503-9

DOI : 10.1007/s00335-008-9137-6

PMID : 18787897

Available at:

<http://archive-ouverte.unige.ch/unige:1319>

Disclaimer: layout of this document may differ from the published version.



UNIVERSITÉ
DE GENÈVE

Functional genetic variation of human miRNAs and phenotypic consequences

Christelle Borel · Stylianos E. Antonarakis

Received: 28 April 2008 / Accepted: 28 July 2008 / Published online: 12 September 2008
© Springer Science+Business Media, LLC 2008

Abstract A large number of human protein-coding genes are finely regulated by one or more microRNAs. Members of this small noncoding RNA family have emerged as important post-transcriptional regulators of gene expression and are involved in a number of disease phenotypes. Variability in the human genome is extensive and includes the common and rare single nucleotide polymorphisms (SNPs) and copy number variations (CNVs). The functional significance of the genome's variability is under intense investigation. In this article we review the emerging literature on how human genomic variation influences the outcome of microRNA targeting and the associated phenotypic effects. Illustrative examples are discussed that demonstrate the biological importance of functional polymorphisms affecting miRNA-mediated gene regulation.

Introduction

The human genome contains many forms of genetic variation. One class of variants comprises the millions of single nucleotide polymorphisms (SNPs) first identified more than 30 years ago in the β -globin gene cluster (Kan and Dozy 1978), and recently cataloged in different populations (Hinds et al. 2005; The International HapMap Consortium 2003). A second common class of variation are copy number variants (CNVs), also identified about 30 years ago in the α -globin gene cluster (Goossens et al. 1980; Kan

et al. 1975). A considerable effort is now devoted toward understanding the extent and magnitude of CNVs in human populations (Wheeler et al. 2008). Here we review the emerging evidence of the effect of human genetic variation on microRNA-mediated gene regulation and discuss how human interindividual variability can influence the outcome of microRNA targeting.

miRNAs and their functions

MicroRNAs (miRNAs, miR), which are short, noncoding RNA molecules (~21 nt), have revealed a new dimension in the complexity of translation control. Our current understanding of the functions of miRNAs is incomplete (Bushati and Cohen 2007). There is considerable evidence that miRNAs function by binding to complementary sequences that usually lie in the 3' untranslated regions (3'UTR) of target mRNAs to induce cleavage, repression of productive translation, and regulation of mRNA stability in the cytoplasm (Ambros 2004; Bartel 2004; Bushati and Cohen 2007; Carrington and Ambros 2003; Filipowicz et al. 2008; Jackson and Standart 2007; Nilsen 2007). Recently, studies revealed additional functions of miRNAs such as upregulation of target mRNA translation upon cell-cycle arrest (Vasudevan et al. 2007), import into the nucleus (Hwang et al. 2007), or even the secretion of the miRNAs from the cell (Valadi et al. 2007). A novel aspect of miRNA-mediated repression with a noncoding transcript in *Arabidopsis thaliana* that can regulate the activity of an miRNA by mimicking its target site has been demonstrated (Franco-Zorrilla et al. 2007). Some authors suggest that some RNAs have a nonproductive interaction with a complementary miRNA to inhibit and regulate the activity of the miRNA. These examples indicate that miRNA

C. Borel · S. E. Antonarakis (✉)
Department of Genetic Medicine and Development, University
of Geneva Medical School and University Hospitals of Geneva,
Geneva 1211, Switzerland
e-mail: stylianos.antonarakis@medecine.unige.ch

molecules have very diverse functions that remain to be explored.

To date, the miRNA registry (MiRBase) contains 6397 mature miRNAs, among which 678 are human miRNAs (April 2008, release 11.0). miRNAs can be encoded in independent transcription units, in polycistronic clusters, or within the introns of protein-coding genes (Bartel 2004). Thus far, most registered miRNAs are widely expressed and highly conserved. Deep sequencing of small RNA libraries and cell-type-specific analyses are currently valuable approaches to uncovering miRNAs with less abundant expression; thus, the catalog of miRNAs is expected to grow substantially in the future (Bar et al. 2008; Glazov et al. 2008; Sunkar et al. 2008).

miRNA expression profiles indicated that some miRNAs are under complex control during development and in a variety of tissues (Farh et al. 2005; Lim et al. 2005; Stark et al. 2005). These studies suggest that miRNAs define highly specific cell identities and have a critical role in cell differentiation. In *C. elegans*, the miRNA lousy-6 (*lisy-6*) is specifically activated by the transcription factor *die-1* in ASE left cells, whereas *lisy-6* is downregulated by miR-273 and then activated by the transcription factor *cog-1* in ASE right cells (Chang et al. 2004; Johnston and Hobert 2003; Johnston et al. 2005). Genetic data demonstrated that *lisy-6*, *cog-1*, miR-273, and *die-1* act in a regulatory double-negative feedback loop (Johnston et al. 2005). This inverse expression of *lisy-6* controls laterality of the nematode chemosensory system and is an excellent illustration of the role of miRNAs in controlling terminally differentiated cellular states.

Although the proteins mediating miRNA biogenesis and function have been clarified, the precise mechanism by which miRNAs regulate the expression of target mRNAs remains unclear. Mature miRNAs preferentially form nonperfect duplexes with the target mRNA (usually in the mRNA 3'UTR) and recruit a repressing complex termed the RNA-induced silencing complex (RISC). This regulation can be highly pleiotropic, with one miRNA able to target several hundred different transcripts. In fact, bioinformatic prediction indicates that 30% of animal genes may be miRNA targets.

Principles of miRNA biogenesis

miRNAs are transcribed by RNA polymerase II (Bartel 2004) as long primary transcripts (pri-miRNAs, 100 nt–10 kb) and are then processed within the nucleus into hairpin-shaped precursor miRNAs (pre-miRNAs, ~70 nt) by the RNase III enzyme Droscha (Lee et al. 2003) and the double-stranded RNA-binding domain cofactor DGCR8/Pasha (DiGeorge syndrome critical region gene 8) (Denli et al. 2004; Gregory et al. 2004; Han et al. 2004; Landthaler

et al. 2004; Lee et al. 2003). An alternative miRNA processing pathway has recently been described that uses splicing processes to bypass Droscha cleavage and to generate miRNA precursors from short intronic sequences (Berezikov et al. 2007; Okamura et al. 2007; Ruby et al. 2007). These small intronic sequences of noncoding RNA are called mirtrons.

In the cytoplasm, Dicer (RNase III) processes the pre-miRNAs to generate ~22-nt miRNA duplexes (Lund et al. 2004). One strand remains a mature miRNA and is then assembled into the effector complex RISC (Khvorova et al. 2003; Schwarz et al. 2003).

Principles of miRNA–mRNA interactions

To specify repression, miRNAs seem to require only short stretches of complementarity to a mRNA, following a broad set of principles that have been identified experimentally and by bioinformatic studies (Brennecke et al. 2005; Doench and Sharp 2004; Grimson et al. 2007; Lewis et al. 2005; Lytle et al. 2007; Nielsen et al. 2007). However, the mechanistic details of miRNA–mRNA interactions are poorly understood and there are always exceptions that make the predictions of miRNA targets difficult. Many studies have confirmed that the interaction is initiated by a continuous and perfect base pairing of seven contiguous nucleotides at positions 2–8 from the 5' end of the miRNA, representing the “seed region” (Lewis et al. 2003, 2005). Mismatches, a guanine–uracil pairing (GU), or bulges in the seed sequences disturb the binding of miRNA with this target site and consequently affect function (Brennecke et al. 2005). However, it has been observed that an A on position 1 of the miRNA (5') and either A or U at position 9 can boost miRNA targeting, although these do not need to base pair with target nucleotides (Grimson et al. 2007). Other characteristics specify targeting: mismatches and bulges are present in the central part of the miRNA–target site duplex, a core region of 3' pairing to residues 13–16 within the miRNA 3' end. However, the degree of repression is also related to UTR contexts and factors that can improve site efficiency (Grimson et al. 2007): multiple, closely spaced target sites act cooperatively, high local AU content in the vicinity of the miRNA target site improves site efficiency, and effective sites are located preferentially near both the poly(A) tail and the stop codon. As part of the different models proposed for miRNA–mRNA interactions, miRNA target sites can be categorized into two distinct classes: 5'-dominant and 3'-compensatory. 5'-Dominant target sites have perfect base pairing with at least 7–8 nt at the 5' end of the miRNA; the base pairing to the rest of the miRNA is considered irrelevant (Chen and Rajewsky 2006; Lewis et al. 2003). 3'-Compensatory target sites have extended base pairing to the 3' end of the miRNA in order to

compensate for an imperfect or shorter stretch of base pairing with the 5' end of the miRNA (Gupta et al. 2006; Kiriakidou et al. 2004; Schratt et al. 2006; Stark et al. 2005).

Genetic variation

Genetic variations range from large chromosome anomalies to single-nucleotide changes. Recently, multiple studies have identified a large number of submicroscopic CNVs of DNA segments ranging from kilobases (kb) to megabases (Mb) in size (Beckmann et al. 2007). CNVs consist of deletions, insertions, duplications, and complex multisite variants in comparison with a reference genome. All those DNA sequence variations could potentially affect the maturation of miRNAs, the silencing machinery, the structure or the expression level of mature miRNA, and the base pairing at the target site, and have functional role for miRNA-mediated gene regulation. We focus on human miRNAs and review all the evidence for human regulatory polymorphisms perturbing miRNA function and thus in turn causing phenotypic variations and disorder.

Polymorphisms and heterogeneity of miRNA sequences

Polymorphism in pri-, pre-, and mature miRNA sequences

A representative mammalian pri-miRNA comprises a stem (~33 bp) with a terminal loop and flanking sequences (~100 bp). Experiments show that a part of the terminal loop (Zeng and Cullen 2005; Zeng et al. 2005) and sequences flanking the stem loop (Han et al. 2006; Lee et al. 2003; Zeng and Cullen 2003) are critical for processing by Drosha and Dicer. It has been hypothesized that polymorphisms in pre-miRNA may influence the miRNA maturation and thereby modulate miRNA expression.

A total of 173 human pre-miRNAs in 96 Japanese individuals have been sequenced (Iwai and Naraba 2005). The study identified ten SNPs in ten pre-miRNA hairpins among which only one SNP in the mature sequence of miR-30c-2 is likely to affect stem integrity (this prediction has not yet been confirmed experimentally).

In another study, a bioinformatic search selected 323 known SNPs located within 227 human pre-miRNA sequences (Duan et al. 2007). Twelve of these SNPs were within pre-miRNA sequences and one, a G/U polymorphism, located at the eighth nucleotide within the mature sequence of miR-125a, has been functionally characterized. This SNP (rs12975333) blocks in vitro the processing

of pri-miRNA into pre-miRNA and alters the translation suppression by miR-125a on a well-known *Lin-28* target mRNA. The biological consequences of these miRNA SNPs remain to be determined.

In a similar study, the occurrence of common SNPs in 474 human pre-miRNAs found by screening the dbSNP database was studied (Saunders et al. 2007). Sixty-five SNPs (including indel polymorphisms) in 49 pre-miRNAs were found, thus exhibiting a SNP density of ~1.3 SNPs per kilobase pair. Only three miRNAs (hsa-mir-125a, hsa-mir-627, and hsa-mir-662) were found to have SNPs within the seed region (rs12975333, rs2620381, and rs9745376, respectively). The conclusion of the study was that miRNA genes have low polymorphism density and that most identified polymorphisms are not within the seed region, indicative of strong selective constraint on human pre-miRNAs. None of these polymorphisms has been tested experimentally to determine whether these SNPs might affect the structure or processing of miRNAs.

A germline DNA variant (C/T) in the primary sequence of the miRNA cluster encoding for miR-16-1 and miR-15a (7 bp in the 3' direction after the precursor) has been identified (Calin et al. 2005). This DNA variant was found in 11 of 75 patients with chronic lymphocytic leukemia, but not seen in 160 subjects without cancer. This nucleotide variation results in reduced mature expression of miR-16-1 and miR-15 in vitro and in vivo and is associated with deletion of the normal allele containing this miR cluster.

A recent example illustrates how a common polymorphism in a miR precursor can consequently affect the mature miRNA expression level. Jazdzewski et al. (2008) reported a G/C (rs 2910164) SNP in pre-miR-146a that results in vitro in reduced amounts of mature miR-146a and contributes to the genetic predisposition to papillary thyroid carcinoma.

A recent genome-wide analysis in healthy individuals revealed 43 miRNAs mapping in CNV regions in one study (Redon et al. 2006), while another study identified 14 CNVs encompassing 21 known miRNAs (Wong et al. 2007). It is not known if the variation in copy number of the miRNAs alters their function and contributes to the phenotypic diversity in humans.

3'- and 5'-end heterogeneity of miRNAs—*isomiRs*

Detailed analysis of mature miRNA sequences revealed an extensive degree of variation at the terminal nucleotides compared to the current miRBase reference sequences (Cummins et al. 2006; Lagos-Quintana et al. 2002; Landgraf et al. 2007; Morin et al. 2008; Ruby et al. 2006; Wu et al. 2007). This population of variants in known miRNAs is collectively termed *isomiRs*. All cited studies observed in most cases nonrandom and conserved 3'-end

heterogeneity with 3'-end deletions and extensions of one to three nucleotides. The most prevalent type of modifications generated single-nucleotide 3' extensions that differ from the genomic sequence. In contrast, 5' ends were more homogeneous, with only a few cases of variation (Landgraf et al. 2007; Morin et al. 2008). These results predict that a particular miRNA hairpin may generate numerous mature miRNAs.

Although the biological consequence of this observation remains to be determined, such variants might act by influencing the miRNA half-life, subcellular localization, and miRNA target specificity (particularly 5'-end variation). In addition, the most abundant isomiRs' sequences need to be determined for each miRNA, and it is essential to verify whether any of these variants associate within RISC and function as gene silencers.

Polymorphisms affecting miRNA–mRNA interactions

Polymorphism in miRNA target sites

The thermodynamics of RNA–RNA interaction plays an essential role in the binding of a miRNA with its target mRNA; it is proposed that SNPs at miRNA binding sites may alter the expression of target genes. Polymorphism density is significantly lower in conserved target site regions that are complementary to the seed region of the miRNA (Chen and Rajewsky 2006). SNPs in this region are expected to be functional and they are candidates for causal variants of human disease. Several studies scanned for the occurrence of SNPs in human 3' UTRs with the purpose of identifying SNPs that may modulate expression of computationally predicted miRNA target sites (Bao et al. 2007; Chen and Rajewsky 2006; Georges et al. 2007; Saunders et al. 2007). Among the 120,000 known 3' UTR SNPs, 19,913 modified putative miRNA targets either by destroying predicted conserved target sites (785) or predicted nonconserved target sites (9470) or by creating 10,283 novel predicted target sites (<http://www.patrocles.org>; Georges et al. 2007).

The first evidence that mutations in a miRNA target site might affect the phenotype came from a study of the gene *SLITRK1*, considered a strong candidate for Tourette's syndrome (Abelson et al. 2005). Mutation screening in 174 unrelated individuals with Tourette's syndrome revealed a G-to-A transition in the 3' UTR of *SLITRK1* that was not present in 1800 normal individuals. This variant is predicted to stabilize the interaction of 3' UTR of *SLITRK1* with miR-189 by changing a G:U wobble pair to an A:U Watson-Crick pair at position 9 of the miRNA. It was demonstrated in vitro that the 3' UTR of *SLITRK1* indeed

facilitates miR-189-mediated downregulation of a luciferase reporter in Neuro2a cells, and the G-to-A variant resulted in a repression of luciferase expression compared to the wild type. There is a developmentally regulated and overlapping expression pattern of *SLITRK1* mRNA and miR-189 in brain regions previously implicated in Tourette's syndrome (Abelson et al. 2005). Moreover, *SLITRK1* promotes dendritic growth while the mutation likely results in a loss of function. These results suggest an association of rare *SLITRK1* sequence variants with Tourette's syndrome (Abelson et al. 2005).

Another example of a functional SNP in the target miRNA sequence is related to hypertension. A SNP rs5186 (1166A/C) in the 3' UTR for the angiotensin receptor-1 (*AGTR1*) gene has been characterized (Sethupathy et al. 2007). The 1166C allele of *AGTR1* has been shown to increase the risk of essential hypertension in several association studies in various populations (EHT [MIM 145500]). Moreover, it has been shown independently that increased levels of *AGTR1* contribute to cardiovascular disease (Song and White 2002; Van Geel et al. 2000) and antagonists of *AGTR1* are now widely used in the treatment of hypertension (Burnier and Brunner 2000). The likely mechanism through which SNP rs5186 regulates *AGTR1* protein levels has recently been elucidated (Sethupathy et al. 2007). This SNP has been localized in a target site for human miR-155 within the 3' UTR of *AGTR1* (Martin et al. 2006). Using reporter silencing assays, it was shown that miR-155 downregulates the expression of only the A and not the C allele of rs5186. Thus, the C allele may be functionally associated with hypertension by abrogating regulation by miR-155, thereby increasing *AGTR1* levels (Sethupathy et al. 2007). Moreover, miR-155 maps to human chromosome 21 and it was observed by quantitative real-time PCR that miR-155 is overexpressed in trisomy 21 individuals. Fibroblasts from monozygotic twins discordant for trisomy 21 showed that the *AGTR1* protein is lower in trisomy 21 fibroblasts compared to normal. Remarkably, it has been reported that individuals with trisomy 21 have lower diastolic and systolic blood pressure levels than age- and gender-matched controls (Draheim et al. 2002; Morrison et al. 1996). Overexpression of miR-155 in trisomy 21 is likely to excessively suppress the *AGTR1* common allele; this may contribute to the lower blood pressure in trisomy 21 individuals.

There are four additional examples of loss of miRNA target site sequences due to SNPs. The presence of SNP 829C/T (rs34764978) in the 3' UTR of *DHFR* led to a loss of miR-24 function and resulted in dihydrofolate reductase overexpression and methotrexate resistance (Mishra et al. 2007). Furthermore, SNP 3142C/G (rs1063320) in the *HLA-G* 3' UTR affects the targeting of miR-148a, miR-148b, and miR-152 and is associated with risk of asthma in

children (Tan et al. 2007). Wang et al. (2008) demonstrated a direct link between miR-433 and a common SNP (rs12720208, T/C) located in the 3' UTR of *fibroblast growth factor 20* (*FGF20*), already associated with an increased risk for Parkinson's disease. The risk allele for this SNP abolished the miR-433 functional activity and increased the translation activity of *FGF20* and α -synuclein expression.

Another recent example of loss of miRNA target site sequences due to SNPs is that type 2 diabetes implicated miR-657 and the IGF-II receptor gene (*IGF2R*). *IGF2R* is located in a region that has been shown to be related to insulin resistance and obesity-related metabolic phenotypes (Duggirala et al. 2001). A case control association study concluded that the ACAA insertion/deletion (144/140 bp) polymorphism at the 3' UTR of the *IGF2R* might participate in the pathogenesis of type 2 diabetes (Villuendas et al. 2006). Recently, new evidence showed that hsa-miR-657 can post-transcriptionally regulate the *IGF2R* expression levels in Hep G2 cells by targeting its 3' UTR. Moreover, it has been demonstrated that the ACAA insertion/deletion polymorphism alters the interaction between *IGF2R* and miR-657 (Lv et al. 2008).

On the other hand, a DNA variant can create a gain of a miRNA target site. QTL mapping and genetic analysis reveal a G-to-A substitution in the 3' UTR of the *GDF8* gene implicated in the polygenic hypermuscularity of Texel sheep (Clop et al. 2006). This point mutation creates two novel target sites for miR-1 and miR-206 (sharing a common seed sequence), which are highly expressed in skeletal muscle. In the presence of the mutation, there was a threefold reduction of circulating *myostatin* and 1.5-fold reduction in the *GDF8* transcript level. The novel miRNA target sites were also experimentally validated.

Polymorphisms affecting miRNA expression

To date there are no SNPs described that affect miRNA expression levels. It is anticipated that such variation will be discovered in the near future as the miRNAs are better characterized, and more genomic variants, common or rare, will be functionally analyzed and be associated with phenotypes.

Conclusions

The study of SNPs has revealed that any two randomly selected human genomes differ by 0.1% (Frazer et al. 2007). A fraction of this genetic variation would affect functional elements directly and will have a causative role in the phenotype. It is therefore expected that genetic

variability would also affect the function of miRNAs. This review points out the emergence of a new class of regulatory variability that might substantially contribute to the heritability of complex traits. Some illustrative examples have been discussed here. However, the majority of this functional variability (particularly rare private variants) may remain unknown. The miRNA function provides excellent opportunities for the discovery of potential epistatic interactions between polymorphisms in miRNA genes and their targets as important risk allele combinations. Because miRNAs affect the fine-tuning of gene expression, we predict that functional variants related to miRNA biology will likely be involved in the complex multifactorial phenotypes.

Acknowledgments The authors' lab is supported by the Swiss National Science Foundation, the NCCR Frontiers in Genetics, The European Union, the NIH, and the Lejeune and ChildCare Foundations. We thank our laboratory colleagues for discussions, critical reading, and numerous suggestions.

Web Resources

dbSNP: <http://www.ncbi.nlm.nih.gov>
 SeattleSNPS database: <http://pga.gs.washington.edu>
 International HapMap project: <http://www.hapmap.org>
 Perlegen Sciences: <http://genome.perlegen.com>
 Patrocles: <http://www.patrocles.org>
 PolymiRTS: <http://compbio.utm.edu/miRSNP/>
 Tarbase: <http://www.diana.pcbi.upenn.edu/tarbase.html>

References

- Abelson JF, Kwan KY, O'Roak BJ, Baek DY, Stillman AA et al (2005) Sequence variants in *SLITRK1* are associated with Tourette's syndrome. *Science* 310:317–320
- Ambros V (2004) The functions of animal microRNAs. *Nature* 431:350–355
- Bao L, Zhou M, Wu L, Lu L, Goldowitz D et al (2007) PolymiRTS Database: linking polymorphisms in microRNA target sites with complex traits. *Nucleic Acids Res* 35:D51–D54
- Bar M, Wyman SK, Fritz BR, Qi J, Garg KS et al (2008) MicroRNA discovery and profiling in human embryonic stem cells by deep sequencing of small RNA libraries. *Stem Cells*
- Bartel DP (2004) MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116:281–297
- Beckmann JS, Estivill X, Antonarakis SE (2007) Copy number variants and genetic traits: closer to the resolution of phenotypic to genotypic variability. *Nat Rev Genet* 8:639–646
- Berezikov E, Chung WJ, Willis J, Cuppen E, Lai EC (2007) Mammalian mirtron genes. *Mol Cell* 28:328–336
- Brennecke J, Stark A, Russell RB, Cohen SM (2005) Principles of microRNA-target recognition. *PLoS Biol* 3:e85
- Burnier M, Brunner HR (2000) Angiotensin II receptor antagonists. *Lancet* 355:637–645
- Bushati N, Cohen SM (2007) microRNA functions. *Annu Rev Cell Dev Biol* 23:175–205
- Calin GA, Ferracin M, Cimmino A, Di Leva G, Shimizu M et al (2005) A microRNA signature associated with prognosis and

- progression in chronic lymphocytic leukemia. *N Engl J Med* 353:1793–1801
- Carrington JC, Ambros V (2003) Role of microRNAs in plant and animal development. *Science* 301:336–338
- Chang S, Johnston RJ Jr, Frokjaer-Jensen C, Lockery S, Hobert O (2004) MicroRNAs act sequentially and asymmetrically to control chemosensory laterality in the nematode. *Nature* 430:785–789
- Chen K, Rajewsky N (2006) Natural selection on human microRNA binding sites inferred from SNP data. *Nat Genet* 38:1452–1456
- Clop A, Marcq F, Takeda H, Pirottin D, Tordoir X et al (2006) A mutation creating a potential illegitimate microRNA target site in the myostatin gene affects muscularity in sheep. *Nat Genet* 38:813–818
- Cummins JM, He Y, Leary RJ, Pagliarini R, Diaz LA Jr et al (2006) The colorectal microRNAome. *Proc Natl Acad Sci USA* 103:3687–3692
- Denli AM, Tops BB, Plasterk RH, Ketting RF, Hannon GJ (2004) Processing of primary microRNAs by the microprocessor complex. *Nature* 432:231–235
- Doench JG, Sharp PA (2004) Specificity of microRNA target selection in translational repression. *Genes Dev* 18:504–511
- Draheim CC, McCubbin JA, Williams DP (2002) Differences in cardiovascular disease risk between nondiabetic adults with mental retardation with and without Down syndrome. *Am J Ment Retard* 107:201–211
- Duan R, Pak C, Jin P (2007) Single nucleotide polymorphism associated with mature miR-125a alters the processing of pri-miRNA. *Hum Mol Genet* 16:1124–1131
- Duggirala R, Blangero J, Almasy L, Arya R, Dyer TD et al (2001) A major locus for fasting insulin concentrations and insulin resistance on chromosome 6q with strong pleiotropic effects on obesity-related phenotypes in nondiabetic Mexican Americans. *Am J Hum Genet* 68:1149–1164
- Farh KK, Grimson A, Jan C, Lewis BP, Johnston WK et al (2005) The widespread impact of mammalian microRNAs on mRNA repression and evolution. *Science* 310:1817–1821
- Filipowicz W, Bhattacharyya SN, Sonenberg N (2008) Mechanisms of post-transcriptional regulation by microRNAs: are the answers in sight? *Nat Rev Genet* 9:102–114
- Franco-Zorrilla JM, Valli A, Todesco M, Mateos I, Puga MI et al (2007) Target mimicry provides a new mechanism for regulation of microRNA activity. *Nat Genet* 39:1033–1037
- Frazer KA, Ballinger DG, Cox DR, Hinds DA, Stuve LL et al (2007) A second generation human haplotype map of over 3.1 million SNPs. *Nature* 449:851–861
- Georges M, Coppieters W, Charlier C (2007) Polymorphic miRNA-mediated gene regulation: contribution to phenotypic variation and disease. *Curr Opin Genet Dev* 17:166–176
- Glazov EA, Cottee PA, Barris WC, Moore RJ, Dalrymple BP et al (2008) A microRNA catalog of the developing chicken embryo identified by a deep sequencing approach. *Genome Res* 18:957–964
- Goossens M, Dozy AM, Embury SH, Zachariades Z, Hadjiminis MG et al (1980) Triplicated alpha-globin loci in humans. *Proc Natl Acad Sci USA* 77:518–521
- Gregory RI, Yan KP, Amuthan G, Chendrimada T, Doratotaj B et al (2004) The Microprocessor complex mediates the genesis of microRNAs. *Nature* 432:235–240
- Grimson A, Farh KK, Johnston WK, Garrett-Engle P, Lim LP et al (2007) MicroRNA targeting specificity in mammals: determinants beyond seed pairing. *Mol Cell* 27:91–105
- Gupta A, Gartner JJ, Sethupathy P, Hatzigeorgiou AG, Fraser NW (2006) Anti-apoptotic function of a microRNA encoded by the HSV-1 latency-associated transcript. *Nature* 442:82–85
- Han J, Lee Y, Yeom KH, Kim YK, Jin H et al (2004) The Drosha-DGCR8 complex in primary microRNA processing. *Genes Dev* 18:3016–3027
- Han J, Lee Y, Yeom KH, Nam JW, Heo I et al (2006) Molecular basis for the recognition of primary microRNAs by the Drosha-DGCR8 complex. *Cell* 125:887–901
- Hinds DA, Stuve LL, Nilsen GB, Halperin E, Eskin E et al (2005) Whole-genome patterns of common DNA variation in three human populations. *Science* 307:1072–1079
- Hwang HW, Wentzel EA, Mendell JT (2007) A hexanucleotide element directs microRNA nuclear import. *Science* 315:97–100
- Iwai N, Naraba H (2005) Polymorphisms in human pre-miRNAs. *Biochem Biophys Res Commun* 331:1439–1444
- Jackson RJ, Standart N (2007) How do microRNAs regulate gene expression? *Sci STKE* 2007:re1
- Jazdzewski K, Murray EL, Franssila K, Jarzab B, Schoenberg DR et al (2008) Common SNP in pre-miR-146a decreases mature miR expression and predisposes to papillary thyroid carcinoma. *Proc Natl Acad Sci USA* 105:7269–7274
- Johnston RJ, Hobert O (2003) A microRNA controlling left/right neuronal asymmetry in *Caenorhabditis elegans*. *Nature* 426:845–849
- Johnston RJ Jr, Chang S, Etchberger JF, Ortiz CO, Hobert O (2005) MicroRNAs acting in a double-negative feedback loop to control a neuronal cell fate decision. *Proc Natl Acad Sci USA* 102:12449–12454
- Kan YW, Dozy AM (1978) Polymorphism of DNA sequence adjacent to human beta-globin structural gene: relationship to sickle mutation. *Proc Natl Acad Sci USA* 75:5631–5635
- Kan YW, Holland JP, Dozy AM, Charache S, Kazazian HH (1975) Deletion of the beta-globin structure gene in hereditary persistence of foetal haemoglobin. *Nature* 258:162–163
- Khvorova A, Reynolds A, Jayasena SD (2003) Functional siRNAs and miRNAs exhibit strand bias. *Cell* 115:209–216
- Kiriakidou M, Nelson PT, Kouranov A, Fitziev P, Bouyioukos C et al (2004) A combined computational-experimental approach predicts human microRNA targets. *Genes Dev* 18:1165–1178
- Lagos-Quintana M, Rauhut R, Yalcin A, Meyer J, Lendeckel W et al (2002) Identification of tissue-specific microRNAs from mouse. *Curr Biol* 12:735–739
- Landgraf P, Rusu M, Sheridan R, Sewer A, Iovino N et al (2007) A mammalian microRNA expression atlas based on small RNA library sequencing. *Cell* 129:1401–1414
- Landthaler M, Yalcin A, Tuschl T (2004) The human DiGeorge syndrome critical region gene 8 and its *D. melanogaster* homolog are required for miRNA biogenesis. *Curr Biol* 14:2162–2167
- Lee Y, Ahn C, Han J, Choi H, Kim J et al (2003) The nuclear RNase III Drosha initiates microRNA processing. *Nature* 425:415–419
- Lewis BP, Shih IH, Jones-Rhoades MW, Bartel DP, Burge CB (2003) Prediction of mammalian microRNA targets. *Cell* 115:787–798
- Lewis BP, Burge CB, Bartel DP (2005) Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* 120:15–20
- Lim LP, Lau NC, Garrett-Engle P, Grimson A, Schelter JM et al (2005) Microarray analysis shows that some microRNAs down-regulate large numbers of target mRNAs. *Nature* 433:769–773
- Lund E, Guttinger S, Calado A, Dahlberg JE, Kutay U (2004) Nuclear export of microRNA precursors. *Science* 303:95–98
- Lv K, Guo Y, Zhang Y, Wang K, Jia Y et al (2008) Allele-specific targeting of hsa-miR-657 to human IGF2R creates a potential mechanism underlying the association of ACAA-insertion/deletion polymorphism with type 2 diabetes. *Biochem Biophys Res Commun* 374:101–105

- Lytle JR, Yario TA, Steitz JA (2007) Target mRNAs are repressed as efficiently by microRNA-binding sites in the 5' UTR as in the 3' UTR. *Proc Natl Acad Sci USA* 104:9667–9672
- Martin MM, Lee EJ, Buckenberger JA, Schmittgen TD, Elton TS (2006) MicroRNA-155 regulates human angiotensin II type 1 receptor expression in fibroblasts. *J Biol Chem* 281:18277–18284
- Mishra PJ, Humeniuk R, Mishra PJ, Longo-Sorbello GS, Banerjee D et al (2007) A miR-24 microRNA binding-site polymorphism in dihydrofolate reductase gene leads to methotrexate resistance. *Proc Natl Acad Sci USA* 104:13513–13518
- Morin RD, O'Connor MD, Griffith M, Kuchenbauer F, Delaney A et al (2008) Application of massively parallel sequencing to microRNA profiling and discovery in human embryonic stem cells. *Genome Res* 18:610–621
- Morrison RA, McGrath A, Davidson G, Brown JJ, Murray GD et al (1996) Low blood pressure in Down's syndrome, A link with Alzheimer's disease? *Hypertension* 28:569–575
- Nielsen CB, Shomron N, Sandberg R, Hornstein E, Kitzman J et al (2007) Determinants of targeting by endogenous and exogenous microRNAs and siRNAs. *RNA* 13:1894–1910
- Nilsen TW (2007) Mechanisms of microRNA-mediated gene regulation in animal cells. *Trends Genet* 23:243–249
- Okamura K, Hagen JW, Duan H, Tyler DM, Lai EC (2007) The mirtron pathway generates microRNA-class regulatory RNAs in *Drosophila*. *Cell* 130:89–100
- Redon R, Ishikawa S, Fitch KR, Feuk L, Perry GH et al (2006) Global variation in copy number in the human genome. *Nature* 444:444–454
- Ruby JG, Jan C, Player C, Axtell MJ, Lee W et al (2006) Large-scale sequencing reveals 21U-RNAs and additional microRNAs and endogenous siRNAs in *C. elegans*. *Cell* 127:1193–1207
- Ruby JG, Jan CH, Bartel DP (2007) Intronic microRNA precursors that bypass Droscha processing. *Nature* 448:83–86
- Saunders MA, Liang H, Li WH (2007) Human polymorphism at microRNAs and microRNA target sites. *Proc Natl Acad Sci U S A* 104:3300–3305
- Schratt GM, Tuebing F, Nigh EA, Kane CG, Sabatini ME et al (2006) A brain-specific microRNA regulates dendritic spine development. *Nature* 439:283–289
- Schwarz DS, Hutvagner G, Du T, Xu Z, Aronin N et al (2003) Asymmetry in the assembly of the RNAi enzyme complex. *Cell* 115:199–208
- Sethupathy P, Borel C, Gagnebin M, Grant GR, Deutsch S et al (2007) Human microRNA-155 on chromosome 21 differentially interacts with its polymorphic target in the *AGTRI* 3' untranslated region: a mechanism for functional single-nucleotide polymorphisms related to phenotypes. *Am J Hum Genet* 81:405–413
- Song JC, White CM (2002) Clinical pharmacokinetics and selective pharmacodynamics of new angiotensin converting enzyme inhibitors: an update. *Clin Pharmacokinet* 41:207–224
- Stark A, Brennecke J, Bushati N, Russell RB, Cohen SM (2005) Animal microRNAs confer robustness to gene expression and have a significant impact on 3'UTR evolution. *Cell* 123:1133–1146
- Sunkar R, Zhou X, Zheng Y, Zhang W, Zhu JK (2008) Identification of novel and candidate miRNAs in rice by high throughput sequencing. *BMC Plant Biol* 8:25
- Tan Z, Randall G, Fan J, Camoretti-Mercado B, Brockman-Schneider R et al (2007) Allele-specific targeting of microRNAs to HLA-G and risk of asthma. *Am J Hum Genet* 81:829–834
- The International HapMap Consortium (2003) The International HapMap Project. *Nature* 426:789–796
- Valadi H, Ekstrom K, Bossios A, Sjostrand M, Lee JJ et al (2007) Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol* 9:654–659
- Van Geel PP, Pinto YM, Voors AA, Buikema H, Oosterga M et al (2000) Angiotensin II type 1 receptor A1166C gene polymorphism is associated with an increased response to angiotensin II in human arteries. *Hypertension* 35:717–721
- Vasudevan S, Tong Y, Steitz JA (2007) Switching from repression to activation: microRNAs can up-regulate translation. *Science* 318:1931–1934
- Villuendas G, Botella-Carretero JI, Lopez-Bermejo A, Gubern C, Ricart W et al (2006) The ACAA-insertion/deletion polymorphism at the 3' UTR of the IGF-II receptor gene is associated with type 2 diabetes and surrogate markers of insulin resistance. *Eur J Endocrinol* 155:331–336
- Wang G, van der Walt JM, Mayhew G, Li YJ, Zuchner S et al (2008) Variation in the miRNA-433 binding site of *FGF20* confers risk for Parkinson disease by overexpression of alpha-synuclein. *Am J Hum Genet* 82:283–289
- Wheeler DA, Srinivasan M, Egholm M, Shen Y, Chen L et al (2008) The complete genome of an individual by massively parallel DNA sequencing. *Nature* 452:872–876
- Wong KK, de Leeuw RJ, Dosanjh NS, Kimm LR, Cheng Z et al (2007) A comprehensive analysis of common copy-number variations in the human genome. *Am J Hum Genet* 80:91–104
- Wu H, Neilson JR, Kumar P, Manocha M, Shankar P et al (2007) miRNA profiling of naive, effector and memory CD8 T cells. *PLoS ONE* 2:e1020
- Zeng Y, Cullen BR (2003) Sequence requirements for microRNA processing and function in human cells. *RNA* 9:112–123
- Zeng Y, Cullen BR (2005) Efficient processing of primary microRNA hairpins by Droscha requires flanking nonstructured RNA sequences. *J Biol Chem* 280:27595–27603
- Zeng Y, Yi R, Cullen BR (2005) Recognition and cleavage of primary microRNA precursors by the nuclear processing enzyme Droscha. *EMBO J* 24:138–148