

# Functional quasi-likelihood regression models with smooth random effects

Jeng-Min Chiou

*National Health Research Institutes, Taipei, Taiwan*

and Hans-Georg Müller and Jane-Ling Wang

*University of California, Davis, USA*

[Received March 2001. Final revision December 2002]

**Summary.** We propose a class of semiparametric functional regression models to describe the influence of vector-valued covariates on a sample of response curves. Each observed curve is viewed as the realization of a random process, composed of an overall mean function and random components. The finite dimensional covariates influence the random components of the eigenfunction expansion through single-index models that include unknown smooth link and variance functions. The parametric components of the single-index models are estimated via quasi-score estimating equations with link and variance functions being estimated nonparametrically. We obtain several basic asymptotic results. The functional regression models proposed are illustrated with the analysis of a data set consisting of egg laying curves for 1000 female Mediterranean fruit-flies (medflies).

**Keywords:** Estimating equations; Functional data analysis; Functional regression; Principal components; Semiparametric quasi-likelihood regression; Single-index model; Smoothing

## 1. Introduction

Methods for the analysis of data which consist of curves or similar infinite dimensional objects are in demand, as the collection of data of this type is becoming increasingly common. Curve data are being collected for instance in endocrinological (Brumback and Rice, 1998) and in growth studies (Gasser *et al.*, 1984; Gasser and Kneip, 1995; Kneip and Gasser, 1992). Overviews of different methods and approaches that have been used for functions viewed as data can be found in Ramsay and Silverman (1997, 2002).

The point of view that we adopt here is that each observed curve is a realization of a stochastic process reflecting the random nature of the individual curves contained in a sample of curves. For curve data this approach was pioneered by Castro *et al.* (1986) and was further developed in the seminal work of Rice and Silverman (1991), who emphasized the importance of smoothing. For many applications, it is of particular interest to model the relationship between covariates or associated variables as predictors and a response curve obtained for each subject. The incorporation of covariate effects in the form of time shifts was considered previously in Capra and Müller (1997), where a covariate  $Z$  is assumed to influence the timescale via a multiplicative scheme,  $t' = \psi(Z)t$ , for an unknown but smooth function  $\psi$  that can be estimated from the sample of curves. Other recent approaches for incorporating covariates into functional data

*Address for correspondence:* Hans-Georg Müller, Department of Statistics, University of California at Davis, One Shields Avenue, Davis, CA 95616-8705, USA.  
E-mail: mueller@wald.ucdavis.edu

models include the development of functional analysis-of-variance approaches (Brumback and Rice, 1998; Staniswalis and Lee, 1998) and adaptations of varying-coefficient regression models (Hoover *et al.*, 1998). A related random-effects model where the observed curves are assumed to be regression splines with random coefficients has been proposed in recent work of Shi *et al.* (1996) and Rice and Wu (2001).

Our starting-point is a sample of stochastic processes  $X_i(t)$ ,  $i = 1, \dots, n$ ,  $t \in T$ , where  $T$  is an interval. In practice, processes  $X_i$  are usually recorded on a grid of time points  $t_{ij}$ ,  $1 \leq j \leq m_i$ , where we assume that these are equidistant or if not equidistant then on a regular grid generated by a design density. For each process  $X_i$ , we observe a vector of covariates or associated variables  $\mathbf{Z}_i$ ,  $\mathbf{Z}_i \in \mathbb{R}^p$ ,  $p \geq 1$ . The observations  $\{\mathbf{Z}_i, X_i(\cdot)\}$  are assumed to be independent and identically distributed. The processes  $X_i(\cdot)$  assume the role of response curves, which are influenced by predictors  $\mathbf{Z}_i$ . It is assumed that the overall mean function  $\mu(\cdot)$  and the eigenfunctions or principal component functions  $\rho_k(\cdot)$ ,  $k = 1, 2, \dots$ , defined via the autocovariance structure of processes  $X_i$  (see Appendix A for details) do not depend on the covariates  $\mathbf{Z}_i$ , whereas the principal components (also called principal component scores) and the conditional means of the processes  $X_i$  do. The case of a high dimensional covariate vector  $\mathbf{Z}_i$  is handled by reducing the covariate vector to a single index. This single index is then embedded in a semiparametric quasi-likelihood regression (SPQR) model. We use the quasi-likelihood with unknown link and variance function estimation or QLUE approach (see Chiou and Müller (1998)), which provides an estimating equation with nonparametrically estimated components, asymptotic distribution theory and an algorithm for fitting the parametric and nonparametric components. We demonstrate the effectiveness of the proposed semiparametric approach through fertility data where the sample of curves consists of the daily egg production observed individually for each of 1000 female Mediterranean fruit-flies (medflies); for an illustration, see Fig. 1.


The paper is organized as follows: The proposed *functional smooth random-effects model* for modelling covariate effects on curve data is presented in Section 2. The estimation of the components of the model, smoothing procedures and asymptotic results are discussed in Section 3. Applying this approach to the egg laying data is the theme of Section 4, which also includes a discussion of various practical issues such as the choice of the number of eigenfunctions and smoothing parameters. Additional technical material such as process representations in terms of eigenfunctions, proofs and details on the estimating equations for SPQR can be found in Appendices A–C.

## 2. Functional smooth random-effects model

Principal components analysis of data vectors is a commonly used multivariate technique. An analysis of a sample of curves may be carried out in a similar way. Castro *et al.* (1986) defined the concept of the best  $K$ -dimensional linear model for stochastic processes and showed that this model corresponds to the first  $K$  terms of the Karhunen–Loève expansion, which is based on the eigenfunctions of the covariance kernel of the process. Rice and Silverman (1991) discussed the application of smoothing techniques for implementing this concept in curve data analysis and proposed a leave one curve out cross-validation technique to select smoothing parameters.

The predictor variable  $\mathbf{Z} \in \mathbb{R}^p$  is observed along with the process  $X(t)$ ,  $t \in T$ , where  $T$  is an interval. We assume that  $\mathbf{Z}$  influences the random components  $A_k$  in the following class of functional smooth random-effects models (model 1).

Given a smooth function  $\mu(t)$ ,  $t \in T$ , random variables  $A_k$ ,  $k = 1, 2, \dots$ , and eigenfunctions  $\rho_k : T \rightarrow \mathbb{R}$ , let



**Fig. 1.** 20 randomly selected smoothed egg laying curves for each of the four quartiles of the total number of eggs: (a) first quartile; (b) second quartile; (c) third quartile; (d) fourth quartile

$$X(t) = \mu(t) + \sum_{k=1}^{\infty} A_k \rho_k(t),$$

and assume for the observed random curves, conditional on the covariates,

$$\mu_Z(t) = E\{X(t)|\mathbf{Z}\} = \mu(t) + \sum_{k=1}^{\infty} E(A_k|\mathbf{Z}) \rho_k(t).$$

The following assumptions are made.

- (a) There is an integer  $K < \infty$  such that the eigenvalues  $\lambda_k = E(A_k^2)$  (see equation (17) in Appendix A) satisfy  $\lambda_k < \infty$  for  $k = 1, \dots, K$ , and  $\lambda_k = 0$ , for  $k > K$ , implying that only the first  $K$  terms of the expansion above matter.
- (b) The eigenfunctions  $\rho_k(\cdot)$  are orthonormal in  $L^2([0, T])$  and are twice continuously differentiable.
- (c) For all  $k$  and  $l$ , using the Kronecker symbol  $\delta_{kl} = 1$  if  $k = l$  and  $\delta_{kl} = 0$  if  $k \neq l$ ,

$$E(A_k) = 0 \quad \text{and} \quad \text{cov}(A_k, A_l) = \lambda_k \delta_{kl}.$$

From assumption (c), we find for the conditional covariance (and setting  $s = t$ , also the conditional variance)

$$\text{cov}\{X(s), X(t)|\mathbf{Z}\} = \sum_{k,l} \text{cov}(A_k, A_l|\mathbf{Z}) \rho_k(s) \rho_l(t). \tag{1}$$

- (d) Each of the conditional link functions for the random effects is a smooth function of a single index formed from the data; i.e. there is a parameter vector  $\beta_k \in \mathbb{R}^p$ ,  $\|\beta_k\| = 1$ , and a twice continuously differentiable function  $\alpha_k : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$\alpha_k(\beta'_k \mathbf{z}) = E(A_k|\mathbf{Z} = \mathbf{z}),$$

for  $k = 1, 2, \dots$

These single-index models serve as a device for dimension reduction.

The proposed functional smooth random-effects model is a natural extension of the classical Karhunen–Loève representation for stochastic processes (see equation (18) in Appendix A). It incorporates the influence of the covariate effects  $\mathbf{Z}$  through the conditional distribution of the principal components scores, which are the random effects in the model, via the smooth regression relationship (d).

The link functions  $\alpha_k$  in assumption (d) are of intrinsic interest as they, jointly with the eigenfunctions  $\rho_k$ , determine the nature of the dependence of the process  $X$  on the predictors  $\mathbf{Z}$ . For a given eigenfunction, an interpretation in terms of the underlying subject-matter problem may be found by checking the processes which have large principal components for this eigenfunction; these ‘principal component scores’ are in turn determined by the shape of the eigenfunction. Based on an interpretation of the  $k$ th eigenfunction  $\rho_k$ , the link function  $\alpha_k$  then reveals the changes in relative importance of this eigenfunction in the representation of assumption (d) and model 1 as the covariate  $\mathbf{Z}$  varies.

We conclude this section with three remarks. First, the conditional covariance structure of the model as given in equation (1) is quite adaptable, while keeping the dimension of the non-parametric components low to avoid the ‘curse of dimensionality’ that is associated with high dimensional nonparametric smoothing.

Second, if the domain is  $T = (t_0)$ , a singleton, then  $X$  is just a real random variable and there is only one eigenfunction:  $\rho_1(t_0) = 1$ . In this case, model 1 collapses to a semiparametric regression model with an unknown link function  $g(\cdot)$  and a parameter vector  $\theta \in \mathbb{R}^p$  such that  $E\{X(t_0)|\mathbf{Z} = \mathbf{z}\} = g(\theta' \mathbf{z})$ , which is determined by the joint distribution of  $(\mathbf{Z}, X(t_0))$ . This connects our approach to quasi-likelihood regression models with unknown link and variance functions (see Appendix C). In this sense, model 1 provides a natural extension of SPQR to the case where the responses are functions.

Third, consider a modified model with an explicit error process, given by  $\tilde{X}(t) = X(t) + \varepsilon(t)$ , where the error process  $\varepsilon(t)$  is independent of  $X$  and  $\mathbf{Z}$  and satisfies  $E\{\varepsilon(t)\} = 0$  and  $\text{cov}\{\varepsilon(s), \varepsilon(t)\} = \sigma(s, t)$ , where  $\sigma(\cdot, \cdot)$  is square integrable. This implies that the processes  $\tilde{X}$  have eigenfunctions  $\tilde{\rho}_k$ ,  $k = 1, 2, \dots$ , and that there are uncorrelated zero-mean random variables  $\tilde{A}_k$  such that

$$\sum \tilde{A}_k \tilde{\rho}_k(t) = \sum A_k \rho_k(t) + \varepsilon(t).$$

Therefore, model 1 still applies, based on random effects  $\tilde{A}_k$  and eigenfunctions  $\tilde{\rho}_k$ , with corresponding altered interpretations.

### 3. Estimation of the model components

#### 3.1. Preliminaries

For estimation and prediction, we have assumed that the number  $K$  of eigenfunctions with non-zero eigenvalues in model 1 is finite. If in reality there are more than the assumed number of  $K$  components, the target is the projection on the space spanned by the first  $K$  eigenfunctions. We also assume from now on that the domain  $T$  of the processes is  $T = [0, \tau]$ , and that discretized measurements  $X_i(t_{ij})$  of processes  $X_i(\cdot)$  are available,  $i = 1, \dots, n$  and  $j = 1, \dots, m_i$ . We develop our arguments for equispaced designs, for which  $t_{ij} = t_j, j = 1, \dots, m$ , and the  $t_j$  are generated by a smooth design density (see, for example, Müller (1984)), but extensions to situations that are not equispaced and for which the number of measurements varies from subject to subject can be covered by adding a presmoothing step, if the number of measurements  $m_i$  for the  $i$ th subject satisfies  $0 < c_1 \leq m_i/m \leq c_2 < \infty, i = 1, \dots, n$ , for constants  $c_1, c_2 > 0$ .

Since smoothing plays a central role in the approach proposed, we briefly discuss some relevant issues. Given any scatterplot data  $(W_i, Y_i)_{i=1, \dots, s}$ , with underlying regression function  $m(w) = E(Y|W = w)$ , we define the smoothed estimate  $\hat{m}(\cdot)$  by

$$\hat{m}(w) = S\{w, b, (W_i, Y_i)_{i=1, \dots, s}\}.$$

Here,  $s$  is the number of data in the scatterplot and  $b$  is the smoothing parameter of the smoother  $S$ , which is evaluated at the argument  $w$ .

For our analysis, it does not matter which particular smoothing method is applied, as long as some basic consistency properties are satisfied and the smoother is linear in the data. Kernel estimators, smoothing splines or local polynomial fitting by locally weighted least squares are among the possible choices. We choose here the locally weighted least squares smoother, denoted by  $S_L$ , fitting local lines to the data. A formal definition is

$$S_L\{w, b, (W_i, Y_i)_{i=1, \dots, s}\} = \arg \min_{a_0} \left\{ \min_{a_1} \left( \sum_{i=1}^s K\left(\frac{w - W_i}{b}\right) [Y_i - \{a_0 + a_1(w - W_i)\}]^2 \right) \right\}.$$

This means that the value of the smoother at the argument  $w$  is obtained as the estimated intercept of a regression line at  $w$ , where the line is fitted to those data falling into the window  $[w - b, w + b]$  by weighted least squares. Here,  $K(\cdot) \geq 0$  is a non-negative kernel function, common choices being  $K(x) = (1 - x^2)\mathbf{1}_{(|x| \leq 1)}$  or  $K(x) = \exp(-x^2/2)$ .

#### 3.2. Estimating overall means and eigenfunctions

A crude estimate of the overall mean function  $\mu$  can be obtained by smoothing pointwise averages  $\tilde{\mu}(\cdot)$ . For equispaced designs, these pointwise averages are

$$\tilde{\mu}(t_j) = \frac{1}{n} \sum_{i=1}^n X_i(t_j), \tag{2}$$

given  $n$  sample curves  $X_1, X_2, \dots, X_n$ . To obtain a smooth function in  $t$ , we may add a smoothing step with a small bandwidth  $b_\mu$ ,

$$\hat{\mu}(t) = S[t, b_\mu, \{t_j, \tilde{\mu}(t_j)\}_{j=1, \dots, m}]. \tag{3}$$

For the case of designs that are not equispaced, a presmoothing step can be carried out first with the aim of sampling all curves at the same time points.

Let  $\Sigma$  be the sample  $m \times m$  variance–covariance matrix of the stochastic process  $X$ , with elements

$$v_{rs} = \frac{1}{n} \sum_{i=1}^n \{X_i(t_r) - \hat{\mu}(t_r)\} \{X_i(t_s) - \hat{\mu}(t_s)\}, \quad 1 \leq r, s \leq m. \tag{4}$$

Denote by  $\tilde{\rho}_k = \{\tilde{\rho}_k(t_1), \tilde{\rho}_k(t_2), \dots, \tilde{\rho}_k(t_m)\}$  the eigenvector corresponding to the  $k$ th largest eigenvalue of the matrix  $\Sigma$ . Following the method used in Capra and Müller (1997), which is a variant of a proposal of Rice and Silverman (1991), a smooth estimate  $\hat{\rho}_k(\cdot)$  of the eigenfunction  $\rho_k(\cdot)$  is obtained simply by smoothing the vector  $\tilde{\rho}_k$ ,

$$\hat{\rho}_k(t) = S[t, b_{\rho_k}, \{t_j, \tilde{\rho}_k(t_j)\}_{j=1, \dots, m}], \quad k = 1, \dots, K. \tag{5}$$

### 3.3. Estimating random effects

Estimation of the link functions  $\alpha_k$  and parameter vectors  $\beta_k$  is based on the estimated overall mean function  $\hat{\mu}$  and eigenfunctions  $\hat{\rho}_k$ . A discrete approximation of  $A_k = \langle \rho_k, X - \mu \rangle$  (see expression (19) in Appendix A) then motivates the updated random-effects estimates

$$\hat{A}_{ik} = \frac{\Delta}{m_i} \sum_{j=1}^{m_i} \{X_i(t_{ij}) - \hat{\mu}(t_{ij})\} \hat{\rho}_k(t_{ij}), \quad k = 1, \dots, K, \tag{6}$$

where  $\Delta = t_{ij} - t_{i(j-1)}$ . To update estimates  $\hat{\alpha}_k(\hat{\beta}'_k \mathbf{Z})$ , for  $\mathbf{Z} \in \mathbb{R}^p$  and  $p > 1$ , we adopt

$$\{\hat{\beta}_k, \hat{\alpha}_k(\cdot)\} = \text{QLUE}\{(\mathbf{Z}_i, \hat{A}_{ik})_{i=1, \dots, n}\}, \tag{7}$$

using SPQR in the form of the QLUE approach of Chiou and Müller (1998) which is summarized in Appendix C. For the special case where the covariate  $\mathbf{Z}$  is one dimensional, we may obtain  $\hat{\alpha}_k(\cdot)$  by a simple nonparametric regression step. In this case, there is no parameter vector, and we can regress the  $\hat{A}_{ik}$  nonparametrically on the  $Z_i$ ,  $i = 1, \dots, n$ , to obtain the smooth function estimates

$$\hat{\alpha}_k(v) = S\{v, b_{\alpha_k}, (Z_i, \hat{A}_{ik})_{i=1, \dots, n}\}. \tag{8}$$

### 3.4. Prediction and estimation algorithm

For prediction, given a vector of covariates  $\mathbf{Z}$  as described in Section 2, we substitute the unknown functions  $\mu$ ,  $\{\alpha_k\}_{k=1, \dots, K}$  and  $\{\rho_k\}_{k=1, \dots, K}$  in the model with the estimates  $\hat{\mu}$ ,  $\hat{\alpha}_k$  and  $\hat{\rho}_k$  respectively. The prediction for  $\mu_{\mathbf{Z}}(t) = E\{X(t)|\mathbf{Z}\}$  is then

$$\hat{\mu}_{\mathbf{Z}}(t) = \hat{\mu}(t) + \sum_{k=1}^K \hat{\alpha}_k(\hat{\beta}'_k \mathbf{Z}) \hat{\rho}_k(t). \tag{9}$$

The estimation procedure of the model components is summarized as follows.

- (a) Obtain the estimates  $\hat{\mu}(t)$  of the overall mean function  $\mu(t)$  by equations (2) and (3).
- (b) Obtain the estimates  $\hat{\rho}_k(t)$  of the eigenfunctions  $\rho_k(t)$  by equation (5).
- (c) Obtain the link function fits  $\hat{\alpha}_k(\cdot)$  and the parameter vector estimates  $\hat{\beta}_k$  via equation (7), the QLUE approach.

The smoothing and QLUE steps require the choice of a smoothing parameter. The choice that is used in the QLUE algorithm is based on special features of quasi-likelihood; see Chiou and Müller (1998). Smoothing parameter selection for the function estimates  $\hat{\rho}_k$  and  $\hat{\mu}_0$  requires

other methods. We shall discuss practical choices coupled to the leave one curve out technique in Sections 4.2 and 4.4.

3.5. Some basic asymptotic results

Following arguments of Pezzulli and Silverman (1993), it is possible to obtain consistency properties for eigenfunction estimates  $(\hat{\rho}_k)_{k=1,\dots,K}$ , under mild regularity conditions. For a function  $\varphi \in L^2(d\nu)$  on domain  $T$ , set

$$\|\varphi\| = \left\{ \int_T \varphi^2(t) d\nu(t) \right\}^{1/2}.$$

We consider here a simplified version of model 1 where  $\mathbf{Z} \in \mathbb{R}^1$  and it follows that  $\beta_k = 1$  for  $k = 1, \dots, K$ .

For the smoother  $S$  that is used in the smoothing steps, we require the following two minimal properties, which are satisfied by virtually all commonly used smoothing methods.

- (a) Given independent and identically distributed random pairs  $(W_i, Y_i)_{i=1,\dots,s}$ , from a distribution with regression function  $m(w) = E(Y|W = w)$ , if the regression function  $m(\cdot)$  is twice continuously differentiable, and the probability distribution function  $f_W$  of the  $W_i$ s is continuous at a point  $w$  and satisfies  $f_W(w) > 0$ , then

$$S\{w, b, (W_i, Y_i)_{i=1,\dots,s}\} - m(w) = O_p(\tau_n),$$

for a sequence  $\tau_n \rightarrow 0$ , which depends on the particular smoother and smoothing parameter choice.

- (b) The smoother is linear in the data  $Y_i$ ,

$$S\{w, b, (W_i, Y_i)_{i=1,\dots,s}\} = \sum_{i=1}^s G_i(w) Y_i$$

for (possibly random) weight functions  $G_i$ , independent of  $Y_i$ , and satisfies

$$\left\{ \sum_{i=1}^s G_i^2(w) \right\} \sum_{i=1}^s \mathbf{1}_{\{G_i(w) \neq 0\}} = O_p(1).$$

Examples of smoothers satisfying these properties are local linear fits or kernel smoothers, among others.

*Theorem 1.* If the smoothers  $S$  that are employed in estimating the nonparametric model components satisfy properties (a) and (b), and the covariate  $Z$  has a density which is continuous and positive at any given point  $z$ , then

$$\|\hat{\mu} - EX\| = O_p(n^{-1/2}). \tag{10}$$

If in addition  $\|\hat{\rho}_k - \rho_k\| = O_p(\sigma_n)$ ,  $k = 1, \dots, K$ , for a sequence  $\sigma_n \rightarrow 0$ , then

$$|\hat{\alpha}(z) - \alpha(z)| = O_p(\sigma_n + \tau_n + n^{-1/2} + m^{-1}). \tag{11}$$

The proof of this result is in Appendix B. Consistency requires, not surprisingly, that both the number  $n$  of observed sample curves as well as the number of measurements  $m$  taken per curve satisfy  $n \rightarrow \infty$  and  $m \rightarrow \infty$ .

We note that a uniform result is possible for equation (11):

$$\sup_z |\hat{\alpha}(z) - \alpha(z)| = O_p\{\nu_n(\sigma_n + \tau_n + n^{-1/2} + m^{-1})\},$$

if instead of property (b) we require

$$\sup_z \left\{ \sum_{i=1}^n G_i^2(z) \right\} \sum_{i=1}^n \mathbf{1}_{\{G_i(z) \neq 0\}} = O_p(\nu_n).$$

A typical value which is achieved by most smoothing methods is  $\nu_n = \{\log(n)/nb\}^{-1/2}$ , where  $b$  is the bandwidth equivalent. Uniform consistency therefore requires in addition that  $\nu_n(\sigma_n + \tau_n + n^{-1/2} + m^{-1}) \rightarrow 0$ . Under certain assumptions, Pezzulli and Silverman (1993) have shown that  $\|\hat{\rho}_k - \rho_k\| = o_p(1)$ , which leads to the consistency result  $|\hat{\alpha}(z) - \alpha(z)| = o_p(1)$ .

These results can be extended to the case of multivariate predictors, using the asymptotic consistency properties of QLUE as described in Chiou and Müller (1998). For the following result on the QLUE estimators for the unknown link functions  $\alpha_k$  and parameter vectors  $\beta_k$  we assume for simplicity that the true  $\mu$  and  $\rho_k$  are known, so that the  $\{A_{ik}\}$  play the role as the usual ‘response’ variables in the SPQR step.

*Theorem 2.* Suppose that the overall mean function  $\mu$  and the eigenfunctions  $\rho_k$  in model 1 are given. Then, under regularity conditions, the following event holds for each  $k, k = 1, \dots, K$ , with probability  $1 - \delta$ , for a given arbitrarily small  $\delta > 0$ : there exist QLUE estimates  $\hat{\beta}_k$  of parameter vectors  $\beta_k$ , satisfying  $\|\hat{\beta}_k\| = 1$ , and  $\hat{\alpha}_k$  of link functions  $\alpha_k$ , such that, as  $n \rightarrow \infty$ ,

$$\sup_z |\hat{\alpha}(z) - \alpha(z)| = o_p(1), \tag{12}$$

$$n^{1/2}\{f(\hat{\beta}_k) - f(\beta_k)\} \xrightarrow{D} N_p\{\mathbf{0}, ((Df)(\beta_k))\Sigma^{-1}((Df)(\beta_k))^T\}, \tag{13}$$

where

$$\Sigma = \lim_{n \rightarrow \infty} \left( \frac{1}{n} \sum_{i=1}^n D_i^T V_i^{-1} D_i \right)$$

with  $D_i = (\partial\alpha(\beta_k^T \mathbf{x}_i)/\partial\beta_k^T)$ ,  $f(\beta_k) = (f_1(\beta_k), \dots, f_{p-1}(\beta_k))^T$  with  $f_j(\beta_k) = \beta_{kj}/\|\beta_k\|$  and

$$(Df)(\beta_k) = \left( \frac{\partial f_i(\beta_k)}{\partial \beta_{kj}} \right)_{1 \leq i \leq p-1, 1 \leq j \leq p},$$

defined as a  $(p - 1) \times p$  matrix.

We note that the degrees of freedom for the regression parameter estimates as well as the rank of the corresponding asymptotic covariance estimates are reduced by 1 because of the identifiability constraint in assumption (d) in Section 2. Details of the proof of the theorem including the adjustment of the asymptotic covariance estimates are similar to those in Chiou and Müller (1998, 2002) and are therefore omitted.

## 4. Application to egg laying curves for medflies

### 4.1. Background

Recently, increased interest in research on aging and quantitative biodemography has focused on the relationship between reproductive and aging patterns. Evolution is driven by reproductive success, and the connection between the evolution of aging and reproduction is intriguing not least because it has proved to be elusive. Unravelling this connection may aid our understanding of the aging process. Patterns of reproduction are typically inferred from experiments which involve large experimental cohorts. The study of the reproductive behaviour of large cohorts of



medflies (*Ceratitis capitata*) has led to several interesting findings. For instance, an interaction between nutritional and gender effects results in a reversal of the female–male life expectancy differential under protein deprivation, as reported in Müller *et al.* (1997), and two different modes of aging and reproduction in conjunction with diet were identified in Carey, Liedo, Müller, Wang and Vaupel (1998). It has been conjectured that reproductive activity has a negative effect on longevity, and the term ‘cost of reproduction’ was coined by Partridge and Harvey (1985). Flies are ideal for the study of reproduction and longevity, as large cohorts can be reared with recordings of both daily egg laying and survival. We focus here on the relationship between patterns in the reproduction curves, which are viewed as dependent data, with the total number of eggs produced by a fly and its lifetime as predictors. The total number of eggs produced is a measure of reproductive success, and it is of interest to find out how reproductive patterns are associated with overall reproductive success and lifetime.

The experiment which provides the data for our analysis was carried out in 1992–1995 at the medfly mass rearing and sterilization facility (Moscamed) at Metapa, Chiapas, Mexico, and consisted of  $n = 1000$  female medflies for which daily egg reproduction was recorded. More details about the biological features of the experiment can be found in Carey, Liedo, Müller, Wang and Chiou (1998).

Let  $X_i(\cdot)$  denote the egg laying curve of the  $i$ th fly and  $\mathbf{Z}_i$  the bivector of lifetime and total number of eggs. Randomly subsampled egg laying curves are visualized in Fig. 1, displaying 20 curves for each quartile of the total number of eggs. From the initial sample of 1000 medflies, those who did not lay any eggs were discarded as those curves have constant value 0, leaving egg laying data for  $n = 936$  medflies in the analysis. We restricted the analysis to the first 50 days of egg laying, as we found large variability of egg laying at higher ages which would entirely dominate the eigenfunctions, and as an earlier analysis in Carey, Liedo, Müller, Wang and Chiou (1998) had shown that the total number of eggs as a function of lifetime showed a marked change point at 51 days. Thus, the egg laying curves  $X_i(t)$ ,  $i = 1, \dots, 936$ , are considered as realizations of a stochastic process on  $T = [0, \tau]$  with  $\tau = 50$  days.


As many of the flies died before 50 days, it is of interest to consider whether informative drop-out occurs in the sense that the mean function of egg laying is affected by the pattern of drop-outs. An approximate parametric model of the egg laying process was developed and used in Müller *et al.* (2001) to define a putative egg laying random process after time of death, and in particular a remaining egg laying potential random function. Although interesting associations were established between lifetime and remaining egg laying potential at the time of death, the association between lifetime and the imputed remaining egg laying process after death was found to be at best very weak, providing some justification for the assumption that the drop-out is non-informative in this example.

#### 4.2. Eigenfunctions

The first four estimated eigenfunctions  $\hat{\rho}_k(5)$  are shown in Fig. 2 with the proportion of variance explained for each principal component indicated in the caption. The corresponding eigenvalues range from 0 to above 6000, which indicates large variation, in accordance with the variability seen in Fig. 1. The first eigenfunction is roughly similar to the overall mean shape of the egg laying curves, with a steep rise in the first 10 days and a subsequent relatively flat downward pointing slope.

If we suppose that in fact  $\rho_1(t) = \mu(t)$ , and that only one principal component matters, we obtain the multiplicative effects model

$$E\{X(t)|\mathbf{Z} = \mathbf{z}\} = \mu(t)\{1 + E(A_1|\mathbf{Z} = \mathbf{z})\}$$



**Fig. 2.** First four eigenfunctions  $\{\hat{\rho}_k\}_{k=1,\dots,4}$  (5) of the egg laying data: the first eigenfunction explains 34.38%, the second an additional 16.41%, the third an additional 8.82% and the fourth an additional 4.93% of the total variance of the data (the bandwidths are selected by cross-validation individually by curve)

and

$$\text{var}\{X(t)|\mathbf{Z} = \mathbf{z}\} = \mu^2(t) \text{var}\{A_1|\mathbf{Z} = \mathbf{z}\}.$$

Conditionally on  $\mathbf{Z} = \mathbf{z}$ , we therefore observe quasi-gamma-type behaviour. Unconditionally,

$$\begin{aligned} E\{X(t)\} &= \mu(t), \\ \text{var}\{X(t)\} &= \mu^2(t)\lambda_1, \end{aligned}$$

and the ‘first mode of variation’ is ‘in the direction of’ the mean function  $\mu(\cdot)$ . Thus, unconditionally, we obtain a constant coefficient of variation model in  $t$ ,  $\sqrt{\lambda_1}$  being the coefficient of variation.

We find that the second eigenfunction differentiates this shape into a very early peak at around 5 days and a subsequent much faster decline, leaving room for further peaks later. The third

eigenfunction depicts a broad peak at later ages, and the higher order eigenfunctions resolve this into a series of increasingly complex oscillations.

The mean function  $\hat{\mu}(3)$  does not depend on the covariate and is shown in Fig. 3. The smoothing parameter is determined by minimizing the squared prediction error obtained by the leave one curve out technique,

$$PE = \sum_{i=1}^n \sum_{j=1}^{m_i} \{ \hat{X}_i^{(-i)}(t_{ij}) - X_i(t_{ij}) \}^2 / nm_i, \tag{14}$$

where  $\hat{X}_i^{(-i)}$  is the leave one curve out function estimate obtained by omitting the  $i$ th sample curve in the estimation. Since for the classical Karhunen–Loève model without covariates  $\hat{X}_i^{(-i)}(t) = \hat{\mu}^{(-i)}(t) = \hat{\mu}^{(-i)}(t, b)$ , we can compute the optimal prediction smoothing parameter  $\hat{b}_\mu = \arg \min_b \{ PE(b) \}$ ; this produced  $\hat{b}_\mu = 2.5$  days.

4.3. Total number of eggs as a single covariate

We first consider the case of a single covariate. The covariate  $Z$  is the total number of eggs, which is the predictor for the random components  $A_k$ . In addition to the estimation of the eigenfunctions  $\rho_k$  and the mean function  $\mu$  we need to obtain estimates  $\hat{\alpha}_k$  (8) for the random-effect link functions  $\alpha_k$ . Implementing these steps leads to the estimated functions  $\hat{\alpha}_k$ , shown in Fig. 4 for  $k = 1-4$ . We find that  $\hat{\alpha}_1$  is monotone increasing throughout and almost linear, whereas  $\hat{\alpha}_2$  is slightly increasing at the beginning and then monotone decreasing throughout with increasingly negative slope. The functions  $\hat{\alpha}_k$  of higher order ( $k = 3, 4$  and higher) appear flat, indicating that the covariate affects only the scores for the first two eigenfunctions. Indeed our final model will not use all four random components.

Given function estimates  $\hat{\mu}(\cdot)$  and  $\{ \hat{\alpha}_k(\cdot) \}_{k=1, \dots, K}$ , we obtain the fitted surface

$$\hat{\mu}_Z(t) = \hat{\mu}(t) + \sum_{k=1}^K \hat{\alpha}_k(Z) \hat{\rho}_k(t), \tag{15}$$





Fig. 3. Estimated overall mean egg laying function of time  $\hat{\mu}(t)$  (3), with a leave one curve out cross-validated bandwidth of 2.5 days




**Fig. 4.** Smoothed random-effects link functions  $\{\hat{\alpha}_k(Z)\}_{k=1,\dots,4}$  (8), in the dependence on the total number of eggs, based on a sample of 936 medflies (the bandwidths selected by cross-validation are 500 for all four curves)

according to equation (9). The fitted surface  $\hat{\mu}_Z(t)$  (15), shown in Fig. 5(a), exhibits an interesting sloping ridge pattern, and the cross-sections, in Fig. 5(b), reveal that, although the peaks are slightly drifting to the right as the total number of eggs increases, the graphs are, for the most part, being shifted upwards as the total number of eggs increases. We note that the number of eigenfunctions is chosen as  $K = 2$  for Fig. 5. This choice is based on visual inspection of the functions  $\hat{\alpha}_k$  in Fig. 4, as well as the predictive quality of the fitted model.

The leave one curve out prediction error (14) is a useful quantification of the predictive quality of a model. The leave one curve out predictors are

$$\hat{\mu}_{Z_i}^{(-i)}(t) = \hat{\mu}^{(-i)}(t) + \sum_{k=1}^K \hat{\alpha}_k^{(-i)}(Z_i) \hat{\rho}_k^{(-i)}(t). \tag{16}$$

Here,  $\hat{\mu}^{(-i)}$ ,  $\hat{\rho}_k^{(-i)}$  and  $\hat{\alpha}_k^{(-i)}$  are all obtained in the same way as before, but omitting the observed process  $X_i$  itself. This functional form of the cross-validation sum of squares was introduced by Rice and Silverman (1991). Prediction errors in the dependence on the number of fitted eigenfunctions are shown in Table 1, supporting the choice  $K = 2$ . The leave one curve out prediction error for a model without the covariate effect, i.e. the model specified by equation (18) in



**Fig. 5.** (a) Fitted surface  $\hat{\mu}_Z(t)$  (15) with the total number of eggs as the covariate and (b) cross-sections through the fitted surface  $\hat{\mu}_Z(t)$  for the total number of eggs fixed at 400 (—), 800 (·····), 1200 (---) and 1600 (— · —)

Appendix A, was found to be 462.54. The large reduction in prediction error when including covariates demonstrates the effectiveness of including the covariate effect in model 1.

#### 4.4. The case of multiple covariates

For the case of multiple covariates, we consider both the total number of eggs laid and lifetime as the predictors for the random components  $A_k$ . The estimation procedure described in Section 3 was implemented for standardized covariates. The dimension for the random components that was used in the analysis is again chosen as  $K = 2$ . The first four estimated random-effects link functions  $\hat{\alpha}_k$  are in Fig. 6.

The estimated coefficients  $\hat{\beta}_1 = (\hat{\beta}_{11}, \hat{\beta}_{12})$  and  $\hat{\beta}_2 = (\hat{\beta}_{21}, \hat{\beta}_{22})$  for the linear predictors in

**Table 1.** Prediction errors PE (14) of fitted model (16) with  $K$  eigenfunctions

$K$	1	2	3	4	5	6	7	8
PE	333.71	313.22	315.22	316.41	315.64	315.62	315.62	315.49

**Table 2.** QLUE for the random components  $\alpha_k(\beta'_k \mathbf{V})$  corresponding to Fig. 6

Function	Coefficient	QLUE (standard error)	Bandwidth	Goodness of fit
$\alpha_1$	$\beta_{11}$ (TotalEgg)	0.8842 (0.0031)	83.78 (variance)	$D = 940.36$
	$\beta_{12}$ (Lifetime)	-0.4671 (0.0059)	1.16 (link)	$P = 916.20$
$\alpha_2$	$\beta_{21}$ (TotalEgg)	0.6730 (0.0520)	28.22 (variance)	$D = 1250.75$
	$\beta_{22}$ (Lifetime)	0.7397 (0.0471)	1.91 (link)	$P = 936.60$

functions  $\alpha_1$  and  $\alpha_2$  are presented in Table 2. These estimates are obtained from the SPQR approach implemented with QLUE. The associated standard errors are useful for asymptotic inference. For the linear predictor  $\beta_1$ , the total number of eggs is relatively more important than lifetime, and both predictors are significant and of opposite sign, forming a contrast in the covariates. For linear predictor  $\beta_2$ , the influence of the two predictors is about the same, forming an average of the covariates, and both predictors are significant.

The effects of individual predictors on the response are a consequence of the coefficients listed in Table 2, and the eigenfunctions in Fig. 2. For the particular application at hand we propose here a criterion for the necessary smoothing parameter selection for the estimation of the link functions that is easy to implement and has good practical properties. It is based on pseudolikelihood (see Davidian and Carroll (1988)),


$$PL\{y; \mu(\beta), \sigma^2(\mu)\} = -\frac{1}{2} \sum_{i=1}^n \log\{2\pi \sigma^2(\mu_i)\} - \frac{1}{2} \sum_{i=1}^n \frac{(y_i - \mu_i)^2}{\sigma^2(\mu_i)},$$

where  $y$ ,  $\mu$  and  $\sigma^2$  respectively represent the vectors of response, mean and variance. Here  $\mu$  and  $\sigma^2$  are replaced by their estimates  $\hat{\mu}$  and  $\hat{\sigma}^2$  in the pseudolikelihood and the optimal bandwidths are chosen by maximizing the pseudolikelihood  $PL(y; \mu, \sigma^2)$ .

The resulting bandwidths for our data analysis are shown in the fourth column of Table 2. In addition, in Table 2 we present the goodness-of-fit statistics for SPQR. Here, we provide the ‘nonparametric’ quasi-deviance  $D$  and Pearson statistic  $P$ , which are asymptotically  $\chi^2$  distributed with the degrees of freedom approximately equal to the number of observations minus the number of parameters to be estimated in the linear predictor. The values in Table 2 do not indicate a lack of fit, as  $D \approx P$  (compare Chiou and Müller (1998), for more details).

### 5. Discussion and concluding remarks

We have developed an approach to extend principal component analysis for curve data to the situation when covariates are present. This approach extends previous work of Rice and Silverman (1991) and yields a flexible functional regression model, the functional smooth random-effects model. This model allows for substantial flexibility in mean regression and heteroscedasticity



**Fig. 6.** Estimated link functions for the random effects  $\{\hat{\alpha}_k(\beta'_k \mathbf{Z})\}_{k=1,\dots,4}$  (7), depicted in dependence on the linear predictors, obtained by the QLUE implementation of SPQR

structure. Flexibility is retained for high dimensions through the single-index feature of the model. Through an example of egg laying curves for 1000 female medflies, this method is shown to be effective in reducing the prediction errors. The approach proposed is simple both conceptually and computationally. The nonparametric components of the SPQR model are structured in such a way that only one-dimensional smoothing steps are needed.

Our main results and techniques have been derived for equispaced designs, where the available measurements for the curve data are made on a regular grid. Non-equispaced data are

clearly important in practice. If a design is ‘mildly’ non-equidistant in the sense that there are no persistent gaps, the methods described in Section 3.1 can still be applied; for persistent gaps in measurements and sparse designs, extensions of the method proposed need to be developed.

An extension to the case that is relevant in practice where the predictor is a time-varying function rather than a time invariant vector is possible through various approaches. One such approach is to discretize a covariate function  $Z(t)$  into a (possibly high dimensional) vector  $Z(s_1), \dots, Z(s_M)$  by specifying an equidistant grid of  $M$  points  $s_1, \dots, s_M$  and then fitting the functional smooth random-effects model as described above, using these vectors as covariates. An alternative approach is to project the random covariate functions onto their first  $K'$  eigenfunctions for some suitable integer  $K'$  and to use the resulting principal component score vectors as  $K'$ -dimensional covariate vectors in the functional smooth random-effects model.

In the approach proposed, for a one-dimensional covariate, acting only on the random effect, all model components can be estimated easily and consistently in one step, and we provide an asymptotic consistency result. The extension to multivariate predictors is achieved by combining these ideas with a single-index quasi-likelihood approach for multivariate and high dimensional regression, using the QLUE technique. This allows us to include both unknown link and variance functions. Our implementation also provides for automatic choices of the smoothing parameters, so that only the covariates need to be identified and no specifications need to be made regarding the nature of the distribution of the random effects or the shape of the link functions.

**Acknowledgements**

We thank Professor J. Carey for making the medfly fecundity data available to us. We are also grateful for the most helpful remarks provided by two referees, an Associate Editor and the Joint Editor. This research was supported in part by National Health Research Institutes grant BS-090-PP-07, National Science Council grant 89-2118-M-194-001, National Science Foundation grants DMS-99-71602 and DMS-02-04869 and National Institutes of Health grant 99-SC-NIH-1028.

**Appendix A: Stochastic processes in  $L^2$**

Suppose that the observed curves  $X_1, X_2, \dots, X_n$  are independent and identically distributed realizations of a stochastic process  $X$  on a domain  $T \subset \mathbb{R}$ . The process is assumed to have mean and covariance functions

$$E\{X(t)\} = \mu(t),$$

$$\text{cov}\{X(s), X(t)\} = \gamma(s, t).$$

Given functions  $f$  and  $g$ , let  $\langle f, g \rangle$  denote the  $L^2(d\nu)$  inner product,

$$\langle f, g \rangle = \int_T f(t) g(t) d\nu(t),$$

where  $d\nu$  is a measure on  $T$  which normally is chosen as the Lebesgue measure.

It is assumed that there is an expansion of  $\gamma$  into orthonormal eigenfunctions  $\rho_k(\cdot)$  on  $L^2(d\nu)$  such that

$$\gamma(s, t) = \sum_{k=1}^{\infty} \lambda_k \rho_k(s) \rho_k(t), \tag{17}$$



with ordered eigenvalues  $\lambda_1 \geq \lambda_2 \geq \dots \geq 0$ . The eigenvalues are non-negative because the covariance kernel  $\gamma(s, t)$  is symmetric and non-negative definite. The orthonormal eigenfunctions satisfy  $\langle \rho_i, \rho_i \rangle = 1$  and  $\langle \rho_i, \rho_j \rangle = 0$  for  $i \neq j$ , and  $\langle \gamma(s, \cdot), \rho_k \rangle = \lambda_k \rho_k(s)$ ,  $s \in T$ ,  $k = 1, 2, \dots$ . The eigenvalues may then be expressed as

$$\lambda_k = \int_T \int_T \rho_k(s) \gamma(s, t) \rho_k(t) \, d\nu(s) \, d\nu(t),$$

$k = 1, 2, \dots$ . Under assumption (17), the stochastic process model then provides the following Karhunen–Loève representation for a randomly selected curve:

$$X(t) = \mu(t) + \sum_{k=1}^{\infty} A_k \rho_k(t), \tag{18}$$

where  $\rho_k$  is the  $k$ th eigenfunction. By Mercer’s theorem (see Riesz and Nagy (1990)), the right-hand side of equation (18) converges uniformly in  $t \in T$ . The random variables  $A_k$  correspond to the principal components and are given by

$$A_k = \langle \rho_k, X - \mu \rangle. \tag{19}$$

The principal components  $A_k$  are uncorrelated random variables and satisfy

$$\begin{aligned} E(A_k) &= 0, \\ \text{var}(A_k) &= \lambda_k, \\ \sum_{k=1}^{\infty} \lambda_k &< \infty, \end{aligned}$$

i.e. the  $k$ th eigenvalue corresponds to the variance of the  $k$ th principal component. From equation (18), we find that, under expression (17),  $X(t)$  is composed of a mean function  $\mu$  plus additive noise, the stochastic part.

The principal components  $A_k$  and basis functions  $\rho_k$  can be interpreted as defining the variation of the stochastic process about its mean function. According to expression (19), the random variable  $A_1$  is the length of the projection of  $X - \mu$  onto  $\rho_1$  for each sample curve  $X$ , and  $A_1 \rho_1$  explains the maximum amount of variation in  $X$  among all functions which involve a single real-valued random variable. Similarly, the function  $A_2 \rho_2$  explains the maximum additional amount of process variation which is unexplained by  $A_1 \rho_1$ , and so forth for  $k = 3, 4, \dots$ .

**Appendix B: Proof of theorem 1**

For equation (10), we observe that

$$E(\|\tilde{\mu} - EX\|^2) = \frac{1}{n^2} \sum_{i,j,k,l} E(A_{ik} A_{jl}) \langle \rho_k, \rho_l \rangle = \frac{1}{n^2} \sum_{i,k} E(A_{ik}^2) \langle \rho_k, \rho_k \rangle = O(n^{-1}).$$

This implies that  $E(\|\tilde{\mu} - \mu\|) = O(n^{-1/2})$  by Jensen’s inequality, and therefore  $\|\tilde{\mu} - \mu\| = O_p(n^{-1/2})$ . The assertion follows from property (b) in Section 3.5 and

$$\|\hat{\mu} - \mu\| \leq \|\tilde{\mu} - \mu\| \sup_x \left\{ \sum_{i=1}^n G_i^2(x) \sum_{i=1}^n \mathbf{1}_{\{G_i(x) \neq 0\}} \right\}. \tag{20}$$

For equation (11), let  $A_{ik}^{(1)} = \langle X_i - \hat{\mu}, \hat{\rho}_k \rangle$ ,  $A_{ik}^{(2)} = \langle X_i - \mu, \rho_k \rangle$ ,  $\hat{\alpha}_k^{(1)}(z) = S\{z, b_{\alpha_k}, (Z_i, A_{ik}^{(1)})_{i=1, \dots, n}\}$  and  $\hat{\alpha}_k^{(2)}(z) = S\{z, b_{\alpha_k}, (Z_i, A_{ik}^{(2)})_{i=1, \dots, n}\}$ . Noting that  $E(A_{ik}^{(2)}|Z) = \alpha_k(Z)$ , the consistency assumption (a) in Section 3.5 on the smoother implies that  $|\hat{\alpha}_k^{(2)}(z) - \alpha_k(z)| = O_p(\tau_n)$ . Furthermore, observing the linearity assumption (b),

$$\begin{aligned} |\hat{\alpha}_k^{(2)}(z) - \hat{\alpha}_k^{(1)}(z)| &= \left| \sum_{i=1}^n G_i(z) \{A_{ik}^{(2)} - A_{ik}^{(1)}\} \right| \\ &\leq \left\{ \sum_{i=1}^n G_i^2(z) \right\}^{1/2} \left\{ \sum_{i=1}^n (A_{ik}^{(2)} - A_{ik}^{(1)})^2 \mathbf{1}_{\{G_i(z) \neq 0\}} \right\}^{1/2}. \end{aligned} \tag{21}$$

Note that

$$\begin{aligned} \sup_i |A_{ik}^{(2)} - A_{ik}^{(1)}| &\leq \sup_i |\langle X_i, \rho_k - \hat{\rho}_k \rangle| + |\langle \hat{\mu}, \hat{\rho}_k - \rho_k \rangle| + |\langle \rho_k, \hat{\mu} - \mu \rangle| \\ &= O_p(\|\hat{\mu} - \mu\| + \|\hat{\rho}_k - \rho_k\|), \end{aligned}$$

since  $K$  is finite, so  $\sup_i \|X_i\| < \infty$ . Then expression (21) implies that  $|\hat{\alpha}_k^{(2)}(z) - \hat{\alpha}_k^{(1)}(z)| = O_p(\|\hat{\mu} - \mu\| + \|\hat{\rho}_k - \rho_k\|)$ , using assumption (b) in Section 3.5 in the same way as in inequality (20). A second application of the same argument, using  $\sup_i |A_{ik}^{(1)} - \hat{A}_{ik}| = O_p(m^{-1})$ , which follows from the smoothness properties of  $X_i$ ,  $\hat{\rho}_k$  and  $\hat{\mu}$ , and equation (6) then implies the result.

### Appendix C: Semiparametric quasi-likelihood regression

SPQR is an estimation procedure for estimating the unknown model components: these consist of two smooth functions, the link function  $g(\cdot)$  and the variance function  $\sigma^2(\cdot)$ , and a vector of regression parameter  $\beta_0$ , with  $\|\beta_0\| = 1$ . Given the observations  $y_i$  and the predictors  $\mathbf{x}_i$ , the model assumptions are

$$\begin{aligned} E(y_i) &= g(\mathbf{x}_i^T \beta_0), & i = 1, \dots, n, \\ \text{var}(y_i) &= \text{var}(\varepsilon_i) = \sigma^2\{g(\mathbf{x}_i^T \beta_0)\} = \sigma^2\{E(y_i)\}. \end{aligned}$$

A three-stage iterative QLUE procedure was proposed by alternating the parametric and nonparametric estimation steps. The procedure can be summarized as follows; see Chiou and Müller (1998, 1999) for additional details.

Let  $S^{(\nu)}\{z, h; (z_i, y_i)_{i=1, \dots, n}\}$ ,  $\nu \geq 0$ , be generic notation for a nonparametric estimator for the  $\nu$ th derivative of a regression function,  $d^\nu E(Y|Z = z)/dz^\nu$ , based on scatterplot data  $(z_i, y_i)_{i=1, \dots, n}$ . Here the  $z_i$ s are design points,  $y_i$ s are raw data to be smoothed,  $h$  denotes a bandwidth and  $z$  is a target level at which the function is to be evaluated. Let  $g_\nu$  be the  $\nu$ th derivative of the link function  $g$ . The various updating steps are as follows.

*Step 1*—nonparametric estimation step for the link function: for given  $\hat{\beta}$ ,  $\|\hat{\beta}\| = 1$ , estimates of the link function and its first derivative are updated by

$$\hat{g}_\nu(t; \hat{\beta}) = S^{(\nu)}\{t, b_\nu; (\mathbf{x}_i^T \hat{\beta}, y_i)_{i=1, \dots, n}\}, \quad \nu = 0, 1.$$

*Step 2*—nonparametric estimation step for the variance function: for given  $\hat{\beta}$ ,  $\|\hat{\beta}\| = 1$  and  $\hat{g}_0(\cdot; \hat{\beta})$ , an updated nonparametric variance estimate  $\hat{\sigma}^2(\cdot)$  is obtained by

$$\hat{\sigma}^2(u) = S^{(0)}\{u, b; (\hat{\mu}_i, \hat{\varepsilon}_i^2)_{i=1, \dots, n}\}$$

where  $\hat{\mu}_i = \hat{g}_0(\mathbf{x}_i^T \hat{\beta}; \hat{\beta})$  and  $\hat{\varepsilon}_i^2 = (y_i - \hat{\mu}_i)^2$  are squared residuals which serve as the ‘raw’ variance estimates and are based on a current model fit.

*Step 3*—parametric estimation step: for given  $\hat{g}_0(\cdot; \hat{\beta})$ ,  $\hat{g}_1(\cdot; \hat{\beta})$  and  $\hat{\sigma}^2(\cdot)$ ,  $\hat{\beta}$  is updated by solving the quasi-likelihood estimating equation with respect to  $\beta$ , inserting the current estimates for the link and variance function. The estimated estimating equation for the score is

$$U^*(\beta) = \sum_{i=1}^n \frac{y_i - \hat{g}_0(\eta_i; \beta)}{\hat{\sigma}^2\{\hat{g}_0(\eta_i; \beta)\}} \hat{g}_1(\eta_i; \beta) \mathbf{x}_i,$$

where  $\mathbf{x}_i = (x_{i1}, x_{i2}, \dots, x_{ip})^T$ ,  $\eta_i = \mathbf{x}_i^T \beta$  and  $\|\beta\| = 1$ .

The iteration is run until convergence. Sliced inverse regression (Li, 1991) estimates provide satisfactory starting values for  $\beta$ . To simplify the notation, we denote the estimated vector of regression parameters and the estimated link function obtained at convergence of the iteration by

$$\{\hat{\beta}, \hat{g}(\cdot)\} = \text{QLUE}\{(\mathbf{x}_i, y_i)_{i=1, \dots, n}\}.$$

### References

Brumback, B. and Rice, J. (1998) Smoothing spline models for the analysis of nested and crossed samples of curves (with discussion). *J. Am. Statist. Ass.*, **93**, 961–994.

- Capra, W. B. and Müller, H. G. (1997) An accelerated time model for response curves. *J. Am. Statist. Ass.*, **92**, 72–83.
- Carey, J., Liedo, P., Müller, H. G., Wang, J. L. and Chiou, J. M. (1998) Relationship of age patterns of fecundity to mortality, longevity and lifetime reproduction in a large cohort of Mediterranean fruit fly females. *J. Geront. A*, **53**, B245–B251.
- Carey, J., Liedo, P., Müller, H. G., Wang, J. L. and Vaupel, J. (1998) Dual modes of aging in Mediterranean fruit fly females. *Science*, **281**, 996–998.
- Castro, P. E., Lawton, W. H. and Sylvestre, E. A. (1986) Principal modes of variation for processes with continuous sample curves. *Technometrics*, **28**, 329–337.
- Chiou, J. M. and Müller, H. G. (1998) Quasi-likelihood regression with unknown link and variance functions. *J. Am. Statist. Ass.*, **93**, 1376–1387.
- Chiou, J. M. and Müller, H. G. (1999) Nonparametric quasi-likelihood. *Ann. Statist.*, **27**, 36–64.
- Chiou, J. M. and Müller, H. G. (2002) Quasi-likelihood regression with multiple indices and smooth link and variance functions. To be published.
- Davidian, M. and Carroll, R. J. (1988) A note on extended quasi-likelihood. *J. R. Statist. Soc. B*, **50**, 74–82.
- Gasser, T. and Kneip, A. (1995) Searching for structure in curve samples. *J. Am. Statist. Ass.*, **90**, 1179–1188.
- Gasser, T., Müller, H. G., Köhler, W., Molinari, L. and Prader, A. (1984) Nonparametric regression analysis of growth curves. *Ann. Statist.*, **12**, 210–229.
- Hoover, D., Rice, J., Wu, C. and Yang, L.-P. (1998) Nonparametric smoothing estimates of time-varying coefficient models with longitudinal data. *Biometrika*, **85**, 809–822.
- Kneip, A. and Gasser, T. (1992) Statistical tools to analyze data representing a sample of curves. *Ann. Statist.*, **20**, 1266–1305.
- Li, K. C. (1991) Sliced inverse regression for dimension reduction. *J. Am. Statist. Ass.*, **86**, 316–327.
- Müller, H. G. (1984) Optimal designs for nonparametric kernel regression. *Statist. Probab. Lett.*, **2**, 285–290.
- Müller, H. G., Carey, J. R., Wu, D., Liedo, P. and Vaupel, J. W. (2001) Reproductive potential predicts longevity of female Mediterranean fruit flies. *Proc. R. Soc. Lond. B*, **268**, 445–450.
- Müller, H. G., Wang, J. L., Capra, W. B., Liedo, P. and Carey, J. R. (1997) Early mortality surge in protein-deprived females causes reversal of sex differential of life expectancy in Mediterranean fruit flies. *Proc. Acad. Sci. USA*, **94**, 2762–2765.
- Partridge, L. and Harvey, P. H. (1985) Costs of reproduction. *Nature*, **316**, 20–21.
- Pezzulli, S. and Silverman, B. W. (1993) Some properties of smoothed principal components analysis for functional data. *Comput. Statist.*, **8**, 1–16.
- Ramsay, J. O. and Silverman, B. W. (1997) *Functional Data Analysis*. New York: Springer.
- Ramsay, J. O. and Silverman, B. W. (2002) *Applied Functional Data Analysis*. New York: Springer.
- Rice, J. A. and Silverman, B. W. (1991) Estimating the mean and covariance structure nonparametrically when the data are curves. *J. R. Statist. Soc. B*, **53**, 233–243.
- Rice, J. A. and Wu, C. (2001) Nonparametric mixed effects models for unequally sampled noisy curves. *Biometrics*, **57**, 253–259.
- Riesz, F. and Nagy, B. S. (1990) *Functional Analysis*. New York: Dover Publications.
- Shi, M., Weiss, R. E. and Taylor, J. M. G. (1996) An analysis of paediatric CD4 counts for acquired immune deficiency syndrome using flexible random curves. *Appl. Statist.*, **45**, 151–163.
- Staniswalis, J. G. and Lee, J. J. (1998) Nonparametric regression analysis of longitudinal data. *J. Am. Statist. Ass.*, **93**, 1403–1418.