

Functional Status Outperforms Comorbidities in Predicting Acute Care Readmissions in Medically Complex Patients

Shirley L. Shih, MD^{1,2}, Paul Gerrard, MD³, Richard Goldstein, PhD¹, Jacqueline Mix, MPH⁴, Colleen M. Ryan, MD^{5,6}, Paulette Niewczyk, PhD^{4,7}, Lewis Kazis, ScD⁸, Jaye Hefner, MD⁹, D. Clay Ackerly, MD, MSc^{10,11}, Ross Zafonte, DO^{1,2}, and Jeffrey C. Schneider, MD^{1,2}

¹Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital, Harvard Medical School, Boston, MA, USA; ²Department of Physical Medicine and Rehabilitation, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA; ³New England Rehabilitation Hospital of Portland, Portland, ME, USA; ⁴Uniform Data System for Medical Rehabilitation, Amherst, NY, USA; ⁵Sumner Redstone Burn Center, Surgical Services, Massachusetts General Hospital, Boston, MA, USA; ⁶Shriners Hospital for Children@-Boston, Boston, MA, USA; ⁷Daemen College, Health Care Studies Department, Amherst, NY, USA; ⁸Department of Health Policy and Management, Boston University School of Public Health, Boston, MA, USA; ⁹Department of Internal Medicine, Spaulding Rehabilitation Hospital, Harvard Medical School, Boston, MA, USA; ¹⁰Department of Internal Medicine, Massachusetts General Hospital, Boston, MA, USA; ¹¹Department of Medicine and Department of Health Care Policy, Harvard Medical School, Boston, MA, USA.

OBJECTIVE: To examine functional status versus medical comorbidities as predictors of acute care readmissions in medically complex patients.

DESIGN: Retrospective database study.

SETTING: U.S. inpatient rehabilitation facilities.

PARTICIPANTS: Subjects included 120,957 patients in the Uniform Data System for Medical Rehabilitation admitted to inpatient rehabilitation facilities under the medically complex impairment group code between 2002 and 2011.

INTERVENTIONS: A Basic Model based on gender and functional status was developed using logistic regression to predict the odds of 3-, 7-, and 30-day readmission from inpatient rehabilitation facilities to acute care hospitals. Functional status was measured by the FIM[®] motor score. The Basic Model was compared to six other predictive models—three Basic Plus Models that added a comorbidity measure to the Basic Model and three Gender-Comorbidity Models that included only gender and a comorbidity measure. The three comorbidity measures used were the Elixhauser index, Deyo-Charlson index, and Medicare comorbidity tier system. The c-statistic was the primary measure of model performance.

MAIN OUTCOME MEASURES: We investigated 3-, 7-, and 30-day readmission to acute care hospitals from inpatient rehabilitation facilities.

RESULTS: Basic Model c-statistics predicting 3-, 7-, and 30-day readmissions were 0.69, 0.64, and 0.65, respectively. The best-performing Basic Plus Model (Basic+Elixhauser) c-statistics were only 0.02 better than the Basic Model, and the best-performing Gender-Comorbidity Model (Gender+Elixhauser) c-statistics were more than 0.07 worse than the Basic Model.

CONCLUSIONS: Readmission models based on functional status consistently outperform models based on medical comorbidities. There is opportunity to improve current national readmission risk models to more accurately predict readmissions by incorporating functional data.

KEY WORDS: functional status; care transitions; statistical modeling; risk assessment; outcomes.

J Gen Intern Med 30(11):1688–95

DOI: 10.1007/s11606-015-3350-2

© Society of General Internal Medicine 2015

INTRODUCTION

Hospitalizations account for the largest share of healthcare costs in the U.S., comprising nearly one-third of all healthcare expenditures.^{1,2} In 2011, readmissions within 30 days of hospital discharge represented more than \$41 billion in hospital costs, with Medicare patients accounting for more than one-half of readmissions.³ Financial penalties for excess 30-day hospital readmissions were instituted by the Centers for Medicare and Medicaid Services (CMS) in 2012⁴; more than 2,200 hospitals were fined a total of \$280 million in reduced Medicare payments in fiscal year 2013.⁵ Despite a growing body of literature describing risk factors associated with hospital readmissions,⁶ it is unclear whether investigative efforts have reduced hospital readmissions.^{5,7,8}

Most readmission risk prediction models have targeted specific medical diagnoses and have utilized comorbidities and demographic data as the central risk factors for hospital readmission. Large U.S. administrative datasets have demonstrated poor discriminative ability (c-statistics: 0.55–0.65).^{6,7} However, few studies have considered functional status or illness severity as potential readmission risk factors.⁶ There is increasing evidence that functional status improves readmission model performance.⁹

Most of the literature on function and hospital readmissions comes from the inpatient rehabilitation setting. Inpatient

Electronic supplementary material The online version of this article (doi:10.1007/s11606-015-3350-2) contains supplementary material, which is available to authorized users.

Received October 21, 2014

Revised April 2, 2015

Accepted April 6, 2015

Published online May 9, 2015

rehabilitation is an ideal population in which to examine the impact of functional status on readmission risk, because (1) inpatient rehabilitation patients often have complex care transitions after acute care discharge, and represent a significant proportion of total readmissions¹⁰; (2) inpatient rehabilitation facilities (IRFs) routinely document functional status using a valid instrument—the FIM^{®11}; and (3) a majority of U.S. IRFs participate in one of the only national datasets that contain standardized functional data—the Uniform Data System for Medical Rehabilitation (UDSMR). Furthermore, there is increasing recognition that a standardized inpatient assessment tool is necessary to improve the discharge planning process and transitions to post-acute care.¹² Models predicting readmission in the stroke,^{13–18} burn,¹⁹ cancer,²⁰ and general inpatient rehabilitation populations²¹ have identified functional status as an important predictor of hospital readmission. Limitations of prior work include small and single-center study designs, narrowly defined patient populations, and defining readmissions beyond the 30-day period.⁶

Overall, there is a lack of literature on the utility of function as a readmission predictor in a large population of medical patients. Moreover, function is a modifiable risk factor with potential to impact readmission outcomes if function-based interventions are instituted early. Therefore, the objective of this study was to compare functional status with medical comorbidities as predictors of acute care readmissions in the medically complex rehabilitation population. We hypothesized that acute care readmission prediction models based on functional status would outperform models based on comorbidities, and that the addition of comorbidity variables to function-based models would not significantly enhance predictive performance.

METHODS

Study Design, Setting, and Population

This was a retrospective database study using data from the UDSMR. The UDSMR is a database of demographic, medical, functional, and facility characteristics of IRF patients, and includes data collected from the Inpatient Rehabilitation Facility Patient Assessment Instrument (IRF-PAI) from approximately 70 % of IRFs in the U.S.

The study population includes adult patients admitted to an IRF for at least one night between 2002 and 2011 under the medically complex impairment code (17.1-17.9). Impairment codes are assigned to each patient upon IRF admission and reflect the primary reason for admission to the IRF. The medically complex impairment code comprises a heterogeneous group of conditions in patients with complex and often multiple diagnoses, including certain infections, neoplasms, nutritional or fluid derangements, circulatory disorders, ventilator- and non-ventilator-dependent respiratory disorders, skin disorders, and medical or surgical complications.²² Patients were excluded from analysis if they were in terminal

care or died while in rehabilitation (they would not have been eligible for readmission to acute care), or if they were not admitted to an IRF directly from acute care. Patients from IRFs where more than 5 % of patients had zero-onset days (defined as zero days between acute care hospital admission and transfer to an IRF) were also excluded from the study, as previous studies have identified these as atypical IRF patients.²³ The unit of analysis for this study was an individual IRF admission.

Primary Outcome and Study Variables

The primary outcome measure was acute care readmission (i.e., transfer from an IRF to an acute care facility with subsequent admission) at 3, 7, and 30 days. While previous studies have largely investigated 30-day readmissions based on the CMS guidelines,²⁴ this study also examined earlier time frames to target preventable readmissions. Early readmissions at 3 days, and possibly at 7 days, tend to reflect premature acute care discharges, inadequate handoff communication, or medication discrepancies.²⁵

Predictor variables included patient age, gender, medical comorbidities, and functional status. The UDSMR captures up to ten medical comorbidities as coded by the International Classification of Diseases, Ninth Edition, Clinical Modification (ICD-9-CM). Comorbidities were subsequently assessed using three different comorbidity classification systems: the Elixhauser comorbidity index,^{26,27} Deyo-Charlson comorbidity index,^{28,29} and the Medicare comorbidity tier system^{30,31} (Table 1).

Functional status was assessed using the FIM[®] instrument (FIM), a validated instrument that measures function using 18 items subcategorized into motor (13 items) and cognitive (5

Table 1 Comorbidity Classification Systems: Summary of the Comorbidity Indices Used in the Analysis

Index	Description
Deyo-Charlson	The Charlson comorbidity index was one of the first methods devised for grouping medical comorbidities. ²⁹ It was developed for use with comorbidity data abstracted from patient charts and subsequently adapted for use with ICD-9-CM codes. ²⁸ The Deyo-Charlson index places ICD-9 codes into one of 17 comorbidity categories and assigns weights to each category. It was developed to identify comorbid illnesses that increase the risk of acute hospital mortality. This index is best suited to detect the most severely ill patients.
Elixhauser	The Elixhauser index is an alternative scheme developed for use with ICD-9 data to address some of the perceived limitations of the Charlson index. ²⁶ Following a number of modifications, it now incorporates 29 disease categories to assess the impact of each disease category on outcomes.
CMS Comorbidity Tiers	The CMS Comorbidity Tiers rely on a four-tiered system for grading medical complexity as part of the prospective payment system developed by RAND for the Medicare prospective payment system of IRFs. ^{30,31} The most complex tier receives the highest payment for a given diagnosis and age. Comorbidity data is obtained for up to 10 ICD-9-CM codes.

items) domains³² (Table 2). The FIM was developed for tracking rehabilitation outcomes and was subsequently incorporated into the IRF-PAI for use in Medicare's prospective payment system for IRFs. Because FIM scores are obtained upon admission to the IRF, they serve as a surrogate assessment of patient functional status at the time of discharge from acute care facilities. FIM scores upon IRF admission have been used in previous research as a proxy for acute care discharge functional status.²¹

Statistical Analysis

Several logistic regression models were created at each readmission time point (3, 7, and 30 days)—a basic model based on functional status (Basic Model), an alternative model based on comorbidities (Gender-Comorbidity Model), and a composite model that included both (Basic Plus Model).

To develop the Basic Model, an exploratory model was first created with the variables age, gender, admission FIM motor score, and admission FIM cognitive score as predictors of 3-, 7-, and 30-day readmission. Linearity checks were made for age and admission FIM motor and cognitive scores. The FIM cognitive score was not statistically significant and therefore was excluded from our final models. A decision was also made to drop age from our models because it did not change model c-statistics, and when used with predictiveness curves, it did not change predicted probabilities and was not better calibrated than models without the age variable. Gender and the quadratic form of the admission FIM motor score were used in the final model.

Table 2 FIM Instrument: a Standardized Measure of Functional Status at the Time of Admission to an Inpatient Rehabilitation Facility

Motor Items	Cognitive Items
Eating	Comprehension
Grooming	Expression
Bathing	Social Interaction
Dressing, upper body	Problem Solving
Dressing, lower body	Memory
Toileting	
Bladder management	
Bowel management	
Transfers- Bed/Chair/Wheelchair	
Transfers- Toilet	
Transfers- Bath/Shower	
Walk/Wheelchair	
Stairs	
Scoring	
Total assistance with helper	1
Maximal assistance with helper	2
Moderate assistance with helper	3
Minimal assistance with helper	4
Supervision or setup with helper	5
Modified independence with no helper	6
Complete independence with no helper	7
Motor Subscale Score (5 items)	13 to 91
Cognitive Subscale Score (5 items)	5 to 35
Total Score (18 items)	18 to 126

There are 18 FIM items—13 motor and 5 cognitive. Each item is scored on an ordinal scale from 1 (dependent) to 7 (independent). Scores are summed for the Motor (13 items) and Cognitive (5 items) sub-scores as well as the total (18 items) FIM

The final regression models included a Basic Model with the variables gender and admission FIM motor scores, three Gender-Comorbidity Models with the variables gender and each of the three comorbidity classification systems, and three Basic Plus Models with all variables (gender, admission FIM motor score, and the three comorbidity classification systems). Each of these models was assessed at each readmission time point. In summary, for each readmission time point, there was one Basic Model, three Gender-Comorbidity Models, and three Basic Plus Models, yielding 21 different models (Table 3). Bootstrapping was used for internal validation of the models.³³ Given the large dataset and small number of study variables, there was little concern of model overfitting, which was also confirmed using bootstrapping and shrinkage coefficient estimates.

To test the study hypothesis, Basic Models were compared to Gender-Comorbidity Models to determine whether function-based models outperformed comorbidity-based readmission models. Also, Basic Models were compared to Basic Plus Models to determine whether any clinically meaningful improvement in predictive value was achieved by adding comorbidity data to a function-based model. Model comparisons were performed at each readmission time point primarily by calculating the c-statistic, or area under the receiver operating characteristic (ROC) curve. The c-statistic is a common measure to compare the goodness of fit of logistic regression models. Due to the large sample size, p values were not used for between-model comparisons, as even negligible differences are likely to be statistically significant.³⁴ Instead, as adjunctive measures of model performance, the Akaike information criterion (AIC), Bayesian information criterion (BIC), pseudo R², and shrinkage coefficients were also calculated. Smaller AIC and BIC values and larger pseudo R² and shrinkage coefficients indicate better model performance.³⁵ Adjustments for multiple comparisons were not made because p value based tests were not used. Interactions between functional status and comorbidity measures at each readmission time point were examined.

Table 3 Overview of Predictive Variables Included in Each Logistic Regression Model

Regression Model	Variables
Basic Model	Gender, admission FIM motor score
Basic Plus Models	
Basic+Elixhauser	Gender, FIM motor score, 29 Elixhauser index comorbidities
Basic+Deyo	Gender, FIM motor score, 2 Deyo-Charlson index comorbidity sum scores*
Basic+CMS Tiers	Gender, FIM motor score, CMS comorbidity tier classification
Gender-Comorbidity Models	
Gender+Elixhauser	Gender, 29 Elixhauser index comorbidities
Gender+Deyo	Gender, 2 Deyo-Charlson index comorbidity sum scores*
Gender+CMS Tiers	Gender, CMS comorbidity tier classification

*Deyo-Charlson sum scores are calculated as follows: The first sum score is based on summing the total number of comorbidities that a subject has that are on the Deyo-Charlson index. The second sum score is the total number of points from the Charlson index that the patient has

In this analysis, standard errors were also adjusted to control for clustering effects.³⁶ However, in the UDSMR dataset, there were at least two dozen IRFs with only one medically complex patient, thus making it difficult to reliably account for random effects.³⁷ To account for repeat admissions to an IRF, a sensitivity analysis was performed examining model c-statistics for the entire study population and comparing them to c-statistics for patients that only presented for an initial rehabilitation admission.

All data analyses were performed using Stata software version 12.1.

RESULTS

The UDSMR database included a total of 134,730 adult patients admitted under the medically complex impairment code to an IRF for at least one night between 2002 and 2011. We excluded 13,773 patients from analysis for the following reasons: terminal care (n=231); died in rehabilitation (n=575); were not admitted to an IRF directly from acute care (=6147); patients from IRFs where more than 5 % of patients had zero onset days (n=6820). Our final sample size was 120,957 patients from 1041 IRFs.

Patient demographics and IRF characteristics are shown in Table 4. In general, the study population was older (mean age 71 years), predominantly white (84 %) and female (54 %), unmarried (54 %), and unemployed (91 %). Medicare patients accounted for 78 % of the study sample. On average, patients

Table 4 Patient Demographics and IRF Characteristics

Number of subjects	120,957
Number of facilities	1041
Age, mean years (SD)	71.4 (13.68)
Male, number (percent)	55,719 (46.07)
Race, number (percent)	
Caucasian	100,342 (84.17)
African American	12,399 (10.40)
Latino/Hispanic	4165 (3.49)
Asian	1389 (1.17)
American Indian/Alaskan	347 (0.29)
Hawaiian/Pacific Islander	318 (0.27)
Multiracial	248 (0.21)
Married, number (percent)	54,933 (45.75)
Living alone, number (percent)	37,434 (32.35)
Employed premorbid, number (percent)	11,164 (9.35)
Primary payer source, number (percent)	
Medicare	94,744 (78.33)
Medicaid	4745 (3.92)
Workers Compensation	354 (0.29)
Unreimbursed	1277 (1.06)
Commercial	18,596 (15.37)
Other	1240 (1.03)
Number of comorbidities, mean (SD)	8.05 (2.42)
Onset days, mean (SD)	17.40 (23.79)
Length of IRF stay, mean days (SD)	13.21 (8.66)
Operating beds, mean (SD)	54.14 (40.92)
FIM Motor admission rating, mean (SD)	40.62 (12.40)
FIM Motor discharge rating, mean (SD)	61.00 (17.02)
Discharge disposition, number (percent)	
Community	88,223 (72.94)
Acute Facility	19,857 (16.42)
Skilled Nursing/Subacute	9317 (7.70)
Other	3560 (2.94)

had eight ICD-9 comorbidities, acute care stays of 17.4 days prior to transfer to an IRF, IRF lengths of stay of 13.2 days, admission FIM motor score of 40.6 (maximum possible score of 91.0), and discharge FIM motor score of 61.0. Only 6440 (5.32 %) patients had more than one admission to the IRF. A total of 19,857 (16.42 %) patients were readmitted to the acute care hospital, of whom 4252 (3.52 %) were readmitted within 3 days, 9560 (7.90 %) within 7 days, and 19,372 (16.02 %) within 30 days of IRF admission. Examination of readmission rates over time demonstrated essentially no change in the rates over the study period.

Regression Model Results

Table 5 shows the odds ratios (ORs) for transfer to an acute care hospital from an IRF at each of the discharge time points based on gender and admission FIM motor score (Basic Model). Female gender and higher FIM motor scores were associated with significantly lower odds of readmission at 3 days (OR_{gender} 0.84 [95 % CI 0.79, 0.90]; OR_{FIM} 0.94 [95 % CI 0.94, 0.95]), 7 days (OR_{gender} 0.83 [95 % CI 0.79, 0.88]; OR_{FIM} 0.96 [95 % CI 0.96, 0.96]), and 30 days (OR_{gender} 0.84 [95 % CI 0.81, 0.87]; OR_{FIM} 0.96 [95 % CI 0.96, 0.96]).

C-statistics for each model and time point are shown in Table 6. Basic Model c-statistics at 3, 7, and 30 days were 0.69, 0.64, and 0.65, respectively. The Basic Plus Models had only marginally better c-statistics. Of the three Basic Plus Models, the best-performing Basic+Elixhauser Model c-statistics were only 0.01, 0.02, and 0.02 greater than the Basic Model c-statistics at the 3-, 7-, and 30-day time points, respectively. In comparison, the Basic Model c-statistics outperformed the Gender-Comorbidity Models at each time point. Of the three Gender-Comorbidity Models, the best-performing Gender+Elixhauser Model c-statistics were 0.12, 0.07, and 0.08 less than the Basic Model c-statistics at 3, 7, and 30 days, respectively. ROC curves for the Basic Model and the best-performing Basic Plus and Gender-Comorbidity Models are provided as supplementary materials (Figs. 1, 2, and 3, available online). As expected, higher c-statistics corresponded with higher pseudo R² and shrinkage coefficients and with lower AIC and BIC estimates (*Supplementary Materials*, Table 7, available online). In general, interactions were not statistically significant and did not affect the results as measured by change in c-statistic. A sensitivity analysis examining model c-statistics for the 5.32 % of patients who had multiple IRF admissions also revealed

Table 5 Logistic Regression Coefficients for the Basic Model at Each Time Point

	3 days	7 days	30 days
Female Gender	0.84 (0.79, 0.90)	0.83 (0.79, 0.88)	0.84 (0.81, 0.87)
Admission FIM Motor	0.94 (0.94, 0.95)	0.96 (0.96, 0.96)	0.96 (0.96, 0.96)
Constant	0.036 (0.034, 0.039)	0.095 (0.090, 0.10)	0.23 (0.22, 0.24)

Data presented as coefficient (95 % confidence interval)

Table 6 Comparison of the Basic Model with Basic Plus and Gender-Comorbidity Readmission Models at All Time Points Using C-Statistics (see Table 1 for model descriptions)

	Basic Model	Basic Plus Models			Gender-Comorbidity Models		
	Gender+FIM	Basic+Elixhauser	Basic+Deyo-Charlson	Basic+CMS Tiers	Gender+Elixhauser	Gender+Deyo-Charlson	Gender+CMS Tiers
3 days	0.69	0.70	0.69	0.69	0.57	0.54	0.54
7 days	0.64	0.65	0.64	0.64	0.57	0.54	0.53
30 days	0.65	0.66	0.65	0.65	0.57	0.54	0.56

no significant difference compared to the overall model c-statistics of the entire study population.

DISCUSSION

To our knowledge, this is the first large national database study examining the role of functional status as the primary predictor of acute care readmission risk in the subacute medically complex population. Our results demonstrate that acute care readmissions at 3, 7, and 30 days are reasonably well predicted based on gender and functional status alone. Importantly, the addition of medical comorbidities did not contribute any further predictive value. However, predictive models that considered only comorbidities without accounting for functional status had inferior performance compared to function-based models. Our results were similar across all three discharge time points, suggesting that functional status is a fairly consistent predictor of readmission risk.

This study is also novel in that it examines not only 30-day hospital readmission, but also early readmissions at 3 and 7 days. Prior national models have traditionally focused on hospital readmissions at 30 days or beyond but have not addressed early readmissions.⁶ However, early readmissions tend to represent more preventable readmissions, as they are more likely secondary to premature hospital discharge, inadequate handoff communication, or medication discrepancies, compared to late readmissions at 30 days.²⁵ Factors surrounding readmissions to acute care from a subacute facility within 7 days deserve further investigation in future readmissions research.

The results of this study suggest that medical comorbidities are insufficient to predict readmission risk for the medically complex IRF population. Recent work by Donzé et al. (2013) suggests that factors such as specific lab values (hemoglobin and sodium level), length of hospital stay, admissions within the last 12 months, and type of admission are helpful predictors of readmission risk in the acute care medical population, presumably because this information addresses the severity of illness or impact of disease on the patient.³⁸ Importantly, Donzé et al. did not include functional status as a variable in their readmissions model. Functional status might also act as a non-disease-specific surrogate marker of illness severity. For instance, prior studies suggest that functional status as measured by the FIM might be considered an indirect measure of a patient's burden of care upon a caregiver.^{39–43} Alternatively,

poor functional status may be an additional risk factor for readmissions by predisposing patients to complications related to impaired mobility such as infections,⁴⁴ venous thromboembolic events,⁴⁵ and falls.⁴⁶ This is consistent with the finding that FIM motor scores were significant predictors of increased readmission risk, but FIM cognitive scores did not significantly improve our readmission models. Although the data presented demonstrates an association between functional status and readmission, further study is needed to determine whether there is a causal relationship between improvements in functional status and reductions in hospital readmissions.

Recent research has identified early rehabilitation of patients during the acute care period as a feasible practice, even in the intensive care setting.^{47,48} Early mobilization of acute respiratory failure patients in the intensive care unit (ICU) has been associated with improvements in functional status and reductions in ICU and overall hospital length of stay.⁴⁹ These studies suggest that, unlike medical comorbidities and patient demographics, functional status is a modifiable risk factor that can be substantially improved with early, concentrated physical and occupational therapy and consultation with a physical medicine and rehabilitation specialist for a comprehensive rehabilitation plan.⁴⁹

Similarly, there are potential clinical applications for utilizing functional status as a means of predicting high-readmission-risk patients. In the acute care setting, development and validation of a simplified functional status measure may prove to be a valuable means of identifying patients at high risk of readmission, such that appropriate preventative steps can be taken prior to discharge. The 6-item AlphaFIM[®] instrument, an abbreviated version of the FIM, is currently being utilized as a functional status measure in the acute care setting, with promising results.^{50,51} Similarly, the 6-item AcuteFIM[™] is an abbreviated version that is being piloted for acute discharge planning.⁵² The Improving Medicare Post-Acute Care Transformation Act (IMPACT) of 2014 is advocating for a standardized inpatient assessment tool to improve quality comparisons across different post-acute care settings, facilitate post-acute discharge planning, and utilize this information for future reforms in hospital reimbursement.¹² Based on our study findings, there is growing evidence supporting the utility of functional status measures as a component of the standardized inpatient assessment.

Despite the increasing body of evidence showing that functional status occupies an important role in health outcomes^{53–59} and that it is a better predictor of mortality than disease-

specific measures and comorbidities,^{60–62} prior research has often overlooked the role of functional status on hospital readmissions. Of the 26 distinct readmission models examined in a 2011 systematic review, only eight models considered functional status as a predictor variable, and only two studies ultimately included overall health and functional status, activities of daily living dependence, and mobility in their final models.⁶ In a recent single-center study of medical patients, Hoyer et al. (2014) found that functional status at the time of acute care discharge was a significant predictor of 30-day acute care readmission, where lowest-functioning patients had the highest rates of readmission (OR 29 %, 95 % CI 25–32 %).²¹ Coupled with our findings that the presence of medical comorbidities does not further improve the prediction of readmission risk, these studies suggest that emphasis on functional status will enhance national readmission models, and interventions aimed at improving function may impact readmission rates.

Placed in the national context of reduced reimbursement for hospital readmissions, this study identifies new areas that can be targeted to reduce readmission rates, ultimately translating into healthcare cost savings. Prior CMS models for hospital readmission risk considered only medical comorbidities and basic demographic factors,^{63–65} with a relatively poor ability to predict hospital readmissions (c-statistics 0.60–0.63). The addition of functional measures can improve current CMS readmission risk models to more accurately identify patients at high risk of readmission and more fairly reimburse hospitals based on performance.

Study Limitations

The limitations of this study should be considered when interpreting the clinical significance of our results. This was a retrospective cohort analysis of previously collected data that was not designed for this study. For instance, the primary reasons for patient readmission to acute care were unavailable in the UDSMR dataset, and therefore were not considered in our analysis. Important prediction variables such as the lab tests included by Donzé et al. (2013)³⁸ and similar surrogates of disease severity or functional status may not be included in our models. Similarly, difficult-to-measure variables such as social support, treatment adherence, quality of post-discharge care, presence of depressive symptoms, and patient cognitive factors outside the scope of the Cognitive FIM are not well captured in the dataset, but may act as confounding factors in our analysis. However, the fact that only key variables were included in our models, and that the choice of variables was hypothesis-driven rather than exploratory, can also be considered a relative strength of this study. Second, the comorbidity data included a maximum of only ten ICD-9-CM codes, and therefore may not have accounted for all patient medical comorbidities. To address this limitation, we included in our models three different comorbidity scoring systems (Elixhauser, Deyo-Charlson, and CMS comorbidity tiers) with established validity in order to minimize bias. Third, unlike well-defined diagnoses such as

stroke or burn, the medically complex category comprises a heterogeneous population with widely varying medical diagnoses. Patient functional status in this context is likely to have wider variability, which may tend to dilute any associations. Nonetheless, the heterogeneity of the medical population represents the variety of medical patients treated in the healthcare system, and therefore the results are applicable to a broad patient base. Finally, the data for this study was obtained from an IRF population and thus may not be directly applicable to medical patients in the acute care setting. However, the UDSMR dataset is one of the largest national datasets with available functional data, and functional data was collected at the time of IRF admission, thereby serving as a reasonable surrogate of functional status at the time of acute care discharge.²¹ Nonetheless, future work is needed to examine and test these findings in the acute care setting.

CONCLUSIONS

Functional status is a frequently overlooked risk factor for readmissions and is a more valuable predictor of readmission risk than medical comorbidities in the medically complex inpatient rehabilitation population. Our results add to the growing body of evidence that functional status is an important predictor of readmissions. There is opportunity to improve current national readmission risk models to more accurately predict readmissions and more fairly reimburse hospitals based on performance. Future efforts are needed to explore early clinical assessment and treatment of functional impairments to reduce hospital readmissions.

Acknowledgments: This paper was presented as an abstract at the American Academy of Physical Medicine and Rehabilitation 2014 Annual Assembly on November 11, 2014.

Conflicts of Interest: There are no conflicts of interest to disclose.

Author Contributions: The contributions of the authors are as follows: design and conceptualization of the study; Paul Gerrard, Jeffrey C. Schneider, Ross Zafonte; analysis and interpretation of the data; Paul Gerrard, Richard Goldstein, Lewis Kazis, Jeffrey C. Schneider, Shirley L. Shih; drafting and revising the manuscript for intellectual content; Shirley L. Shih, Paul Gerrard, Jeffrey C. Schneider, Richard Goldstein, Jacqueline Mix, Colleen M. Ryan, Paulette Niewczyk, Jaye Hefner, D. Clay Ackerty.

Corresponding Author: Jeffrey C. Schneider, MD; Department of Physical Medicine and Rehabilitation Spaulding Rehabilitation Hospital, Harvard Medical School, Boston, MA, USA (e-mail: jcschneider@partners.org).

REFERENCES

1. **Pfuntner A, Wier LM, Steiner C.** Costs for Hospital Stays in the United States, 2010. HCUP Statistical Brief #146. Rockville: Agency for Healthcare Research and Quality; 2013.
2. **Pfuntner A, Wier LM, Steiner C.** Costs for Hospital Stays in the United States, 2011. HCUP Statistical Brief #168. Rockville: Agency for Healthcare Research and Quality; 2013.

3. **Hines AL, Barrett ML, Jiang HJ, Steiner CA.** Conditions with the Largest Number of Adult Hospital Readmissions by Payer, 2011. HCUP Statistical Brief #172. Rockville: Healthcare Research and Quality; 2014.
4. "Readmission Reductions Program." Centers for Medicare and Medicaid Services. <<http://cms.gov/Medicare/Medicare-Fee-for-Service-Payment/AcuteInpatientPPS/Readmissions-Reduction-Program.html/>>. Aug 04, 2014. Last accessed April 1, 2015.
5. **James J.** Health Policy Brief: Medicare Hospital Readmission Reduction Program. Health Aff. 2013;32:1-5.
6. **Kansagara D, Englander H, Salanitro A, et al.** Risk Prediction Models for Hospital Readmission: A Systematic Review. JAMA. 2011;306(15):1688-98.
7. **Marks E.** Complexity Science and the Readmission Dilemma: Comment on "Potentially Avoidable 30-Day Hospital Readmissions in Medical Patients" and "Association of Self-Reported Hospital Discharge Handoffs With 30-Day Readmissions". JAMA Intern Med. 2013;173(8):629-31.
8. **Gerhardt G, Yemane A, Hickman P, et al.** Data Shows Reduction in Medicare Hospital Readmission Rates During 2012. Medicare Medicaid Res Rev. 2013;3(2):E1-12.
9. **Coleman EA, Min SJ, Chomiak A, Kramer AM.** Posthospital Care Transitions: Patterns, Complications, and Risk Identification. Health Serv Res. 2004;39(5):1449-65.
10. **Burke RE, Coleman EA.** Interventions to Decrease Hospital Readmissions: Keys for Cost-Effectiveness. JAMA Intern Med. 2013;173(8):695-8.
11. **Hamilton BB, Laughlin JA, Fiedler RC, Granger CV.** Interrater Reliability of the 7-Level Functional Independence Measure (FIM). Scand J Rehabil Med. 1994;26(3):115-9.
12. "Bipartisan, Bicameral Effort Underway to Advance Medicare Post-Acute Reform." U.S. House of Representatives Committee on Ways and Means. 2014 Mar 18. Available at: <http://waysandmeans.house.gov/news/documentsingle.aspx?DocumentID=373213>. Last accessed April 1, 2015.
13. **Roberts PS, DiVita MA, Riggs RV, et al.** Risk factors for discharge to an acute care hospital from inpatient rehabilitation among stroke patients. PM&R. 2014;6(1):50-5.
14. **Chung DM, Niewczyk P, DiVita M, et al.** Predictors of discharge to acute care after inpatient rehabilitation in severely affected stroke patients. Am J Phys Med Rehabil. 2012;91(5):387-92.
15. **Ottenbacher KJ, Graham JE, Ottenbacher AJ, et al.** Hospital readmission in persons with stroke following postacute inpatient rehabilitation. J Gerontol A Biol Sci Med Sci. 2012;67(8):875-81.
16. **Roth EJ, Lovell L, Harvey RL, et al.** Incidence of and risk factors for medical complications during stroke rehabilitation. Stroke. 2001;32(2):523-9.
17. **Stineman MG, Ross R, Maislin G, et al.** Risks of acute hospital transfer and mortality during stroke rehabilitation. Arch Phys Med Rehabil. 2003;84(5):712-8.
18. **Gerrard P, Black-Schaffer R, DiVita M, et al.** Functional status as a predictor of acute care readmissions in the stroke population [Abstract]. Am J Phys Med Rehabil. 2014;93(3)(Supplement 2):a7.
19. **Schneider JC, Gerrard P, Goldstein R, et al.** Predictors of transfer from rehabilitation to acute care in burn injuries. J Trauma Acute Care Surg. 2012;73(6):1596-601.
20. **Asher A, Roberts PS, Bresee C, et al.** Transferring inpatient rehabilitation facility cancer patients back to acute care (TRIPBAC). PM&R. 2014;6(9):808-13.
21. **Hoyer EH, Needham DM, Atanelov L, et al.** Association of impaired functional status at hospital discharge and subsequent rehospitalization. J Hosp Med. 2014;9(5):277-82.
22. UB Foundation Activities. The Inpatient Rehabilitation Facility-Patient Assessment Instrument (IRF-PAI) Training Manual. 2004. Available at: <http://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/InpatientRehabFacPPS/downloads/irfpaimanual040104.pdf>. Last accessed April 1, 2015.
23. **Schneider JC, Tan WH, Goldstein R, et al.** Time to Rehabilitation in the Burn Population: Incidence of Zero Onset Days in the UDSMR National Dataset. J Burn Care Res. 2013;34(6):666-73.
24. Centers for Medicare and Medicaid Services. Medicare Hospital Quality Chartbook: Performance Report on Outcome Measures. Sept 2013. Yale New Haven Health Services Corporation, Centers for Outcomes Research and Evaluation.
25. **Ouslander JG, Diaz S, Hain D, Tappen R.** Frequency and Diagnoses Associated with 7- and 30-Day Readmission of Skilled Nursing Facility Patients to a Nonteaching Community Hospital. J Am Med Dir Assoc. 2011;12:195-203.
26. **Elixhauser A, Steiner C, Harris DR, Coffey RM.** Comorbidity measures for use with administrative data. Med Care. 1998;36(1):8-27.
27. **Comorbidity Software** [computer program]. Version 3.5. Available at: <http://www.hcup-us.ahrq.gov/toolssoftware/comorbidity/comorbidity.jsp>. 2010. Last accessed April 1, 2015.
28. **Deyo RA, Cherkin DC, Ciol MA.** Adapting a clinical comorbidity index for use with ICD-9-CM administrative databases. J Clin Epidemiol. 1992;45(6):613-9.
29. **Charlson M, Szatrowski TP, Peterson J, Gold J.** Validation of a combined comorbidity index. J Clin Epidemiol. 1994;47(11):1245-51.
30. Centers for Medicare and Medicaid Services. Medicare Program; Prospective Payment System for Inpatient Rehabilitation Facilities. Fed Regist. 2001;66:41316-430.
31. **Carter GM, Buntin MB, Hayden O, et al.** In: Carter GM, Buntin BM, Hayden O, et al., eds. Analyses for the Initial Implementation of the Inpatient Rehabilitation Facility Prospective Payment System. Santa Monica: RAND Corporation; 2002:99-154.
32. **Stineman MG, Shea JA, Jette A, et al.** The Functional Independence Measure: Tests of Scaling, Assumptions, Structure, and Reliability Across 20 Diverse Impairment Categories. Arch Phys Med Rehabil. 1996;77(11):1101-8.
33. **Grunkeimer GL, Wu YX.** Bootstrap resampling methods: Something for nothing? Ann Thorac Surg. 2004;77:1142-4.
34. **Lin M, Lucas, Jr. HC, Shmueli G.** Too Big to Fail: Large Samples and the *p*-Value Problem. Inform Syst Res. 2013;24(4):906-17.
35. **Harrell FE Jr.** Regression Modeling Strategies: With Application to Linear Models, Logistic Regression, and Survival Analysis. New York: Springer-Verlag New York; 2001.
36. **Williams R.** A Note on Robust Variance Estimation for Cluster-Correlated Data. Biometrics. 2000;56:645-6.
37. **McNeish DM.** Modeling sparsely clustered data: design-based, model-based, and single-level methods. Psychol Methods. 2014;19(4):552-63.
38. **Donzé J, Aujesky D, Williams D, Schnipper JL.** Potentially avoidable 30-day hospital readmissions in medical patients: derivation and validation of a prediction model. JAMA Intern Med. 2013;173(8):632-8.
39. **Corrigan JD, Smith-Knapp K, Granger CV.** Validity of the Functional Independence Measure for Persons with Traumatic Brain Injury. Arch Phys Med Rehabil. 1997;78(8):828-34.
40. **Disler PB, Roy CW, Smith BP.** Predicting hours of care needed. Arch Phys Med Rehabil. 1993;74(2):139-43.
41. **Granger CV, Cotter AC, Hamilton BB, Fiedler RC.** Functional Assessment Scales: A Study of Persons after Stroke. Arch Phys Med Rehabil. 1993;74(2):133-8.
42. **Granger CV, Cotter AC, Hamilton BB, et al.** Functional Assessment Scales: A Study of Persons with Multiple Sclerosis. Arch Phys Med Rehabil. 1990;71(11):870-5.
43. **Hamilton BB, Deutsch A, Russell C, et al.** Relation of Disability Costs to Function: Spinal Cord Injury. Arch Phys Med Rehabil. 1999;80(4):385-91.
44. **Anderson DJ, Chen LF, Schmader KE, et al.** Poor Functional Status as a Risk Factor for Surgical Site Infection Due to Methicillin-Resistant *Staphylococcus aureus*. Infect Control Hosp Epidemiol. 2008;29:832-9.
45. **Sellier E, Labarere J, Sevestre A-M, et al.** Risk Factors for Deep Vein Thrombosis in Older Patients: A Multicenter Study with Systematic Compression Ultrasonography in Postacute Care Facilities in France. J Am Geriatr Soc. 2008;56:224-30.
46. **Robbins AS, Rubenstein LZ, Josephson KR, et al.** Predictors of Falls Among Elderly People: Results of Two Population-Based Studies. Arch Intern Med. 1989;149(7):1628-33.
47. **Adler J, Malone D.** Early Mobilization in the Intensive Care Unit: A Systematic Review. Cardiopulm Phys Ther J. 2012;23(1):5-13.
48. **Needham DM.** Mobilizing Patients in the Intensive Care Unit: Improving Neuromuscular Weakness and Physical Function. JAMA. 2008;300(14):1685-90.
49. **Needham DM, Korupolu R, Zanni JM, et al.** Early Physical Medicine and Rehabilitation for Patients with Acute Respiratory Failure: A Quality Improvement Project. Arch Phys Med Rehabil. 2010;91:536-42.
50. **Stillman G, Granger C, Niewczyk P.** Projecting function of stroke patients in rehabilitation using the AlphaFIM instrument in acute care. PM&R. 2009;1(3):234-9.
51. **Lo A, Tahair N, Sharp S, Bayley MT.** Clinical utility of the AlphaFIM[®] instrument in stroke rehabilitation. Int J Stroke. 2012;7(2):118-24.
52. Uniform Data System for Medical Rehabilitation. The FIM[®] Instrument: Its Background, Structure, and Usefulness. Buffalo: UDSMR; 2012.
53. **Brezinski D, Stone PH, Muller JE, et al.** Prognostic significance of the Kamoisky Performance Status score in patients with acute myocardial infarction: comparison with the left ventricular ejection fraction and the exercise treadmill test performance. The MILIS Study Group. Am Heart J. 1991;121(5):1374-81.

54. **Cesari M, Kritchevsky SB, Penninx BW, et al.** Prognostic value of usual gait speed in well-functioning older people—results from the Health, Aging and Body Composition Study. *J Am Geriatr Soc.* 2005;53(10):1675–80.
55. **Cesari M, Onder G, Zamboni V, et al.** Physical function and self-rated health status as predictors of mortality: results from longitudinal analysis in the iSIRENTE study. *BMC Geriatr.* 2008;8:34.
56. **Cesari M, Kritchevsky SB, Newman AB, et al.** Added value of physical performance measures in predicting adverse health-related events: results from the Health, Aging And Body Composition Study. *J Am Geriatr Soc.* 2009;57(2):251–9.
57. **Penninx BW, Ferrucci L, Leveille SG, et al.** Lower extremity performance in nondisabled older persons as a predictor of subsequent hospitalization. *J Gerontol A Biol Sci Med Sci.* 2000;55(11):M691–7.
58. **Stommel M, Given BA, Given CW.** Depression and functional status as predictors of death among cancer patients. *Cancer.* 2002;94(10):2719–27.
59. **Inouye SK, Peduzzi PN, Robison JT, et al.** Importance of functional measures in predicting mortality among older hospitalized patients. *JAMA.* 1996;279(15):1187–93.
60. **Landi F, Liperoti R, Russo A, et al.** Disability, more than multimorbidity, was predictive of mortality among older persons aged 80 years and older. *J Clin Epidemiol.* 2010;63:752–9.
61. **Marengoni A, von Strauss E, Rizzuto D, et al.** The impact of chronic multimorbidity and disability on functional decline and survival in elderly persons. A community-based, longitudinal study. *J Intern Med.* 2009;265(2):288–95.
62. **Lee SJ, Go AS, Lindquist K, et al.** Chronic conditions and mortality among the oldest old. *Am J Public Health.* 2008;98(7):1209–14.
63. **Krumholz HM, Lin Z, Drye EE, et al.** An administrative claims measure suitable for profiling hospital performance based on 30-day all-cause readmission rates among patients with acute myocardial infarction. *Circ Cardiovasc Qual Outcomes.* 2011;4(2):243–52.
64. **Keenan PS, Normand SL, Lin Z, et al.** An administrative claims measure suitable for profiling hospital performance on the basis of 30-day all-cause readmission rates among patients with heart failure. *Circ Cardiovasc Qual Outcomes.* 2008;1:29–37.
65. **Lindenauer PK, Normand SL, Drye EE.** Development, validation, and results of a measure of 30-day readmission following hospitalization for pneumonia. *J Hosp Med.* 2011;6(3):142–50.