

Further Indo-West Pacific palaemonoid shrimps (Crustacea: Decapoda: Palaemonoidea), principally from the New Caledonian region

XINZHENG LI¹ & ALEXANDER J. BRUCE²

¹Institute of Oceanology, Chinese Academy of Sciences, Qingdao, China, and ²Queensland Museum, South Brisbane, QLD, Australia

(Accepted 14 April 2006)

Abstract

Based on the material deposited in the Muséum national d'Histoire naturelle, Paris, collected from the Indo-West Pacific, principally from the New Caledonian region, the present paper reports 117 palaemonoid shrimp species, which belong, respectively, to Anchistioididae (one genus, one species), Gnathophyllidae (one genus, one species), Palaemonidae Palaemoninae (seven genera, nine species), and Palaemonidae Pontoniinae (30 genera, 106 species), including eight new species. The new species are all Pontoniinae: *Mesopontonia brevicarpalis* sp. nov., *Palaemonella komaii* sp. nov., *Periclimenes crosnieri* sp. nov., *Periclimenes forgesi* sp. nov., *Periclimenes loyautensis* sp. nov., *Periclimenes paralcocki* sp. nov., *Periclimenes paraleator* sp. nov., and *Periclimenes pseudalcocki* sp. nov. The last six new species are members of the deep-water “*Periclimenes alcocki* species complex”, which has more than two (usually four) pairs of dorsolateral telson spines anterior to the posterior telson margin, the cornea is usually reduced, the dactyl of the major second chela is generally flanged and the chela is sometimes covered with small tubercles. The complex is usually found at more than 200 m depth in the West Pacific. The species can be distinguished from each other by the armature of ambulatory propod and dactyl, diameter of cornea, rostrum shape and the number of pairs of dorsolateral telson spines. *Mesopontonia brevicarpalis* sp. nov., from the southeast coast of Africa, is the seventh species of the genus. *Palaemonella komaii* sp. nov. is very similar to *Palaemonella dolichodactylus* Bruce, 1991 and *Palaemonella hachijo* Okuno, 1999. These three species share the features of very long and slender ambulatory pereopods with the dactyl more than eight times longer than its basal depth and with several long setae on the dorsal dactylar margin.

Résumé

Basé sur le matériel déposé au Muséum national d'Histoire naturelle, à Paris, récolté dans l'Indo-ouest Pacifique, principalement dans la région néo-calédonienne, la présente étude porte sur 117 espèces de crevettes Palaemonoidea appartenant respectivement aux familles Anchistioididae (un genre, une espèce), Gnathophyllidae (un genre, une espèce), et Palaemonidae (37 genres et 115 espèces). Chez les Palaemonidae sept genres et neuf espèces appartiennent à la sous-famille Palaemoninae et 30 genres et 106 espèces, donc huit nouvelles, à la sous-famille Pontoniinae. Les espèces sont: *Mesopontonia brevicarpalis* sp. nov., *Palaemonella komaii* sp. nov., *Periclimenes crosnieri* sp. nov., *Periclimenes forgesi* sp. nov., *Periclimenes loyautensis* sp. nov., *Periclimenes paralcocki* sp. nov.,

Correspondence: Xinzheng Li, Institute of Oceanology, Chinese Academy of Sciences, 7 Nanhai Road, Qingdao 266071, China.
Email: lixzh@ms.qdio.ac.cn

Published 12 July 2006

ISSN 0022-2933 print/ISSN 1464-5262 online © 2006 Taylor & Francis
DOI: 10.1080/00222930600763627

Periclimenes paraleator sp. nov, *Periclimenes pseudalcocki* sp. nov. Les six dernières appartiennent à ce que l'on peut appeler "le complexe *Periclimenes alcocki*" donc les espèces ont un telson portant, en avant de son bord postérieur, plus de deux (habituellement quatre) paires d'épines dorsolatérales, une cornée habituellement réduite, la plus grande des pinces des deuxième périoopodes parfois couverte de petits tubercules et avec des dactyles généralement munis d'un rebord. Les espèces de ce complexe sont habituellement trouvées à plus de 200 m de profondeur dans l'Ouest Pacifique. Elles peuvent être séparées les unes des autres par l'armature du propode et du dactyle des pattes ambulatoires, le diamètre de la cornée, le rostre et le nombre des paires d'épines dorsolatérales du telson. *Mesopontonia brevicarpalis* sp. nov. de la côte sud-est de l'Afrique est la septième espèce du genre, *Palaemonella komaii* sp. nov. est très semblable à *Palaemonella dolichodactylus* Bruce, 1991 et *Palaemonella hachijo* Okuno, 1999; ces trois espèces présentent des périoopodes ambulatoires très longs et grêles, terminés par un dactyle plus de huit fois plus long que large à sa base et orné de plusieurs longues soies sur son bord dorsal. Si l'on considère les espèces mentionnées par Bruce (1981a, 1985a, 1990a, 1991a, 1991b, 1996) d'après les récoltes faites dans Indo-Ouest Pacifique et déposées au Muséum national d'Histoire naturelle, c'est 169 espèces de crevettes Palaemonoidea appartenant à cinq familles ou sous-familles et 48 genres qui sont actuellement déposés à Paris.

Keywords: *Indo-West Pacific, New Caledonia, Palaemonoidea, shrimps, new species*

Introduction

The shrimp material of the superfamily Palaemonoidea deposited principally in the collections of the Muséum national d'Histoire naturelle, Paris, collected mainly from the Indo-Pacific areas, has been reported by Bruce (1981a, 1985a, 1990a, 1991a, 1991b, 1996). The present study, which is a continuation of the series reports of the superfamily, started in 1995 by A. J. Bruce when he visited the museum. X. Li restarted this study when he visited the museum in the fall of 2004. The present paper reports the combined results of this research based on the Indo-Pacific material from the east coast of Africa to southwestern Pacific, but principally from the New Caledonian region, collected by a series of campaigns by the Muséum national d'Histoire naturelle, and the Office de la Recherche Scientifique et Technique Outre-Mer (now Institut de Recherche pour le Développement).

The names, research vessels, dates and localities of expeditions (campagnes) which produced samples in this study are listed as following:

MD 32: "Marion Dufresne", 11 August to 10 September 1982, La Réunion.

MUSORSTOM 3: "Coriolis", 31 May to 7 June 1985, Philippines.

BIOCAL: "Jean-Charcot", 9 August to 10 September 1985, New Caledonia (Loyalty Basin and Norfolk Ridge).

KARUBAR: "Baruna Jaya 1", 21 October to 5 November 1991, Indonesia (Kai Islands and Tanimbar).

MUSORSTOM 7: "Alis", 5 May to 4 June 1992, Wallis Islands and Futuna.

BERYX 11: "Alis", 13–23 October 1992, New Caledonia (Norfolk and Loyalty Ridges).

SMIB 8: "Alis", 26 January to 3 February 1993, New Caledonia (Norfolk Ridge).

BATHUS 1: "Alis", 9–20 March 1993, New Caledonia (east coast).

BATHUS 2: "Alis", 10–18 May 1993, New Caledonia (southwest coast).

BATHUS 3: "Alis", 22 November to 2 December 1993, New Caledonia (south of Loyalty Islands ridge, Norfolk ridge, southwest New Caledonia).

HALIPRO 1: "Alis", 18–25 March and 29 March to 1 April 1994, New Caledonia (east and south coast).

BATHUS 4: "Alis", 1–12 August 1994, New Caledonia (north and west coast).

MUSORSTOM 8: "Alis", 19 September to 14 October 1994, Vanuatu.

- HALIPRO 2: “Tangaroa”, 4–28 November 1996, Norfolk Ridge and south Loyalty Ridge.
 AMUSIUM 1: “Alis”, 5–14 June 1998, Lagoons of Landsdowne and Bellona.
 MUSORSTOM 10: “Alis”, 5–19 August 1998, Fiji.
 SUVA 2: “Alis”, 13–22 October 1998, Fiji (Viti Levu, north and west lagoons).
 BORDAU 1: “Alis”, 22 February to 14 March 1999, Fiji.
 SURPRISES: “Alis”, 12 April to 15 May 1999, New Caledonia (Surprise Atoll, northwest coast of New Caledonia).
 LITHIST: “Alis”, 10–12 August 1999, New Caledonia.
 SUVA 4: “Alis”, 21–26 September 1999, Fiji (Suva Harbour, Lauthana Bay, Beqa Lagoon, Pacific Harbour).
 BORDAU 2: “Alis”, 31 May to 22 June 2000, Tonga.
 NORFOLK 1: “Alis”, 19–26 June 2001, Norfolk Ridge.
 SALOMON 1: “Alis”, 23 September to 7 October 2001, Solomon Islands.
 NORFOLK 2: “Alis”, 20 October to 6 November 2003, Norfolk Ridge.
 SALOMON 2: “Alis”, 20 October to 8 November 2004, Solomon Islands.

In the following account, carapace length (cl) refers to the postorbital carapace length; the rostral formula is presented as “a+b/c” (a=the number of dorsal rostral teeth posterior to the orbital margin, b=the number of dorsal rostral teeth anterior to the orbital margin, c=the number of the ventral rostral teeth) or “b/c” if there is no dorsal rostral tooth on the carapace posterior to the orbital margin; stn=station. The specimens are all deposited in the collections of the Muséum national d’Histoire naturelle, Paris.

Species list

(Three families, 39 genera, 117 species.)

Family ANCHISTIOIDIDAE Borradaile, 1915

Anchistioides willeyi (Borradaile, 1899) 616

Family GNATHOPHYLLIDAE Dana, 1852

Gnathophyllum americanum Guérin-Méneville, 1855 617

Family PALAEMONIDAE Rafinesque, 1815

Subfamily PALAEMONINAE Rafinesque, 1815

Brachycarpus biunguiculatus (Lucas, 1846) 618
Leander tenuicornis (Say, 1818) 618
Macrobrachium australe (Guérin-Méneville, 1838) 619
Macrobrachium equidens (Dana, 1852) 619
Nematopalaemon tenuipes (Henderson, 1893) 620
Palaemon debilis Dana, 1852 620
Palaemonetes atrinubes Bray, 1976 621
Urocaridella antonbruunii (Bruce, 1967) 621
Urocaridella urocaridella (Holthuis, 1950) 623

Subfamily PONTONIINAE Kingsley, 1878

<i>Allopontonia iaini</i> Bruce, 1972	623
<i>Altopontonia disparostris</i> Bruce, 1990	624
<i>Amphipontonia kanak</i> Bruce, 1991	625
<i>Anchistus custos</i> (Forskål, 1775)	625
<i>Anchistus miersi</i> (De Man, 1888)	626
<i>Anchistus pectinis</i> Kemp, 1922	626
<i>Climeniperæus truncoideus</i> (Chace and Bruce, 1993)	627
<i>Conchodytes biunguiculatus</i> (Paulson, 1875)	627
<i>Conchodytes maculatus</i> Bruce, 1989	628
<i>Conchodytes meleagrinae</i> Peters, 1852	628
<i>Coralliocaris</i> sp.	629
<i>Coralliocaris macrophthalma</i> (H. Milne-Edwards, 1837)	629
<i>Coralliocaris superba</i> (Dana, 1852)	630
<i>Coralliocaris viridis</i> Bruce, 1974	630
<i>Dasycaris ceratops</i> Holthuis, 1952	631
<i>Dasycaris symbiotes</i> Kemp, 1922	631
<i>Dasycaris zanzibarica</i> Bruce, 1973	632
<i>Exochlomenella denticulata</i> (Nobili, 1906)	632
<i>Exochlomenella</i> cf. <i>denticulata</i> (Nobili, 1906)	633
<i>Exochlomenella maldivensis</i> Duris and Bruce, 1995	633
<i>Hamodactylus boschmai</i> Holthuis, 1952	634
<i>Hamodactylus noumeae</i> Bruce, 1970	634
<i>Hamopontonia corallicola</i> Bruce, 1970	635
<i>Harpiliopsis beaupresii</i> (Audouin, 1825)	635
<i>Harpiliopsis depressa</i> (Stimpson, 1860)	636
<i>Harpiliopsis spinigera</i> (Ortmann, 1890)	636
<i>Harpilius lutescens</i> Dana, 1852	637
<i>Ischnopontonia lophos</i> (Barnard, 1962)	638
<i>Jocaste japonica</i> (Ortmann, 1890)	638
<i>Jocaste lucina</i> (Nobili, 1901)	639
<i>Kemponia agag</i> (Kemp, 1922)	639
<i>Kemponia amymone</i> (De Man, 1902)	641
<i>Kemponia anacanthus</i> (Bruce, 1988)	641
<i>Kemponia andamanensis</i> (Kemp, 1922)	642
<i>Kemponia darwiniensis</i> (Bruce, 1987)	643
<i>Kemponia elegans</i> (Paulson, 1875)	643
<i>Kemponia ensifrons</i> (Dana, 1852)	644
<i>Kemponia grandis</i> (Stimpson, 1860)	644
<i>Kemponia kororensis</i> (Bruce, 1977)	645
<i>Kemponia lacertae</i> (Bruce, 1992)	647
<i>Kemponia nilandensis</i> (Borradaile, 1915)	647
<i>Kemponia seychellensis</i> (Borradaile, 1915)	648
<i>Kemponia</i> cf. <i>suwadiensis</i> (Borradaile, 1915)	649
<i>Kemponia tenuipes</i> (Borradaile, 1898)	650
<i>Manipontonia psamathe</i> (De Man, 1902)	651
<i>Mesopontonia brevicarpalis</i> sp. nov.	652

<i>Mesopontonia gracilicarpus</i> Bruce, 1990	655
<i>Metapontonia fungiacola</i> Bruce, 1967	655
<i>Palaemonella crosnieri</i> Bruce, 1978	656
<i>Palaemonella dolichodactylus</i> Bruce, 1991	656
<i>Palaemonella hachijo</i> Okuno, 1999	657
<i>Palaemonella komaii</i> sp. nov.	657
<i>Palaemonella pottsii</i> (Borradaile, 1915)	663
<i>Palaemonella pusilla</i> Bruce, 1975	666
<i>Palaemonella rotumana</i> (Borradaile, 1898)	666
<i>Palaemonella spinulata</i> Yokoya, 1936	668
<i>Paraclimenes franklinae</i> (Bruce, 1990) nom. nov.	668
<i>Paranchistus nobilii</i> Holthuis, 1952	669
<i>Paranchistus ornatus</i> Holthuis, 1952	669
<i>Periclimenella petitthouarsi</i> (Audouin, 1825)	670
<i>Periclimenella spinifera</i> (De Man, 1902)	670
<i>Periclimenes affinis</i> (Zehntner, 1894)	671
<i>Periclimenes alcocki</i> Kemp, 1922	672
<i>Periclimenes aleator</i> Bruce, 1991	673
<i>Periclimenes amboinensis</i> (De Man, 1888)	674
<i>Periclimenes attenuatus</i> Bruce, 1971	675
<i>Periclimenes brevicarpalis</i> (Schenkel, 1902)	676
<i>Periclimenes brevirostris</i> Bruce, 1991	677
<i>Periclimenes calcaratus</i> Chace and Bruce, 1993	677
<i>Periclimenes ceratophthalmus</i> Borradaile, 1915	679
<i>Periclimenes commensalis</i> Borradaile, 1915	679
<i>Periclimenes cristimanus</i> Bruce, 1965	680
<i>Periclimenes crosnieri</i> sp. nov.	681
<i>Periclimenes forcipulatus</i> Bruce, 1991	686
<i>Periclimenes forgesi</i> sp. nov.	686
<i>Periclimenes foveolatus</i> Bruce, 1981	691
<i>Periclimenes hertzvigi</i> Balss, 1913	692
<i>Periclimenes imperator</i> Bruce, 1967	694
<i>Periclimenes incertus</i> Borradaile, 1915	695
<i>Periclimenes involens</i> Bruce, 1996	696
<i>Periclimenes kempi</i> Bruce, 1969	697
<i>Periclimenes laccadivensis</i> (Alcock and Anderson, 1884)	697
<i>Periclimenes lanipes</i> Kemp, 1922	698
<i>Periclimenes latipollex</i> Kemp, 1922	699
<i>Periclimenes lepidus</i> Bruce, 1978	699
<i>Periclimenes loyautensis</i> sp. nov.	700
<i>Periclimenes macropthalmus</i> Fujino and Miyake, 1970	703
<i>Periclimenes magnificus</i> Bruce, 1979	705
<i>Periclimenes novaecaledoniae</i> Bruce, 1967	706
<i>Periclimenes paralcocki</i> sp. nov.	707
<i>Periclimenes paraleator</i> sp. nov.	711
<i>Periclimenes pilipes</i> Bruce and Zmarzley, 1983	715
<i>Periclimenes platyrhynchus</i> Bruce, 1991	716
<i>Periclimenes pseudalcocki</i> sp. nov.	716

<i>Perichlimeses soror</i> Nobili, 1904	721
<i>Perichlimeses tosaensis</i> Kubo 1951	721
<i>Perichlimeses uniunguiculatus</i> Bruce, 1990	722
<i>Perichlimeses vaubani</i> Bruce, 1990.	722
<i>Perichlimeses venustus</i> Bruce, 1990	723
<i>Perichlimeses</i> sp.	723
<i>Philarius imperialis</i> (Kubo, 1940)	725
<i>Platycaris latirostris</i> Holthuis, 1952	726
<i>Pliopontonia furtiva</i> Bruce, 1973	726
<i>Pontoniopsis comanthi</i> Borradaile, 1915	727
<i>Thaumastocaris streptopus</i> Kemp, 1922	727
<i>Zenopontonia noverca</i> (Kemp, 1922)	728

Systematic account

Family ANCHISTIOIDIDAE Borradaile, 1915

Anchistioides willeyi (Borradaile, 1899)

Palaemonopsis willeyi Borradaile 1899, p 410, Plates 36, 37, Figure 7.

Amphipalaemon willeyi: Borradaile 1917, p 407, Plate 59, Figure 13.

Anchistioides willeyi: Gordon 1935, p 344, 345, Figures 23a, 24a; Holthuis 1952a, p 18, 214–219, Figures 106, 107 (*partim*); Bruce 1990a, p 211; Bruce 1991a, p 400, Figure 70; Bruce 1991b, p 269, Figures 3g, 29, 30; Bruce 1996, p 261.

Material examined

Comoro Islands: (i) Mayotte, lagoon, dredge, 55 m, coll. A. Crosnier, September 1959, 1 ovig. ♀ (MNHN-Na 141804). **Madagascar:** (ii) Pracel Bank, west coast, 55 m, sand, trawl, coll. A. Crosnier, June 1959, 1♀ (MNHN-Na 14805); (iii) Nosy Be, shallow water, coll. A. Crosnier, 1966, 1♀ (MNHN-Na 15965); (iv) 13°40.3'S, 47°48.0'E, 32 m, sand, trawl, coll. A. Crosnier, 5 December 1972, 1 juvenile (MNHN-Na 14803). **Philippines:** (v) MUSORSTOM 2, stn CP8, 13°55'N, 120°20'E, 85–90 m, 21 November 1980, 2♀♀, 2 juveniles (MNHN-Na 14802). **New Caledonia:** (vi) lagoon, Five Miles Channel, scuba, 15–20 m, in sponge, coll. C. Vadon, 20 September 1978, 1 ovig. ♀ (MNHN-Na 15849); (vii) BATHUS 1, stn CP680, 20°48'S, 165°18'E, 86–92 m, 15 March 1993, 1♀ (MNHN-Na 14801); (viii) Nouville, washings from eunicid tubes, 18–20 m, coll. P. Bouchet, 1 April 1993, 1♂ (MNHN-Na 15812); (ix) Îlot Maitre, 22°19.35'S, 166°25.85'E, 20 m, scuba, in sponge, coll. I. Takeuchi, 10 November 1995, 1♂ (MNHN-Na 15808); (x) Îlot Maitre, stn 82, 22°19.61'S, 166°24.07'E, 10 m, coll. I. Takeuchi, 14 November 1995, 1♂ (MNHN-Na 15844); (xi) Surprise Atoll, North New Caledonia, stn CP 1378, 18°26.3'S, 163°08.2'E, 39 m, 9 May 1999, coll. Richer de Forges, 1♂ (MNHN-Na 15814); (xii) Sarcelle Channel, 30–45 m, in boulders, 1 ovig. ♀ (MNHN-Na 14800). **Fiji:** (xiii) MUSORSTOM 10, south of Viti Levu, stn CP 1364, 18°11.9'S, 178°34.5'E, 80–86 m, 15 August 1998, 2♀♀ (MNHN-Na 15547); (xiv) SUVA 2, Viti Levu, South Lagoon, stn BS 18, 18°11.4'S, 178°28.2'E, 83 m, 15 October 1998, 1 ovig. ♀ (MNHN-Na 15546); (xv) SUVA 4, Viti Levu, Bequ Lagoon, stn DW 07, 28–32 m, 18°22.1'S, 178°02.5'E, 24 September 1999, 1♂ (MNHN-Na 15545).

Remarks

Specimens (iv)–(vii) are of the long-fingered second pereopod type, (ix) with short fingers and in (ii) the fingers are subequal to the palm. The specimens (viii) may have been derived from sand-burrowing sponges rather than the eunicid tubes as this species is normally associated with sponges.

Distribution

Type locality: Ralun, New Britain. Also known from Kenya, Zanzibar, Tanzania, Madagascar, Seychelles, Maldives, Singapore, South China Sea, Philippines, Indonesia (Borneo Bank), Australia (Northern Territory, Queensland), and New Caledonia.

Family GNATHOPHYLLIDAE Dana, 1852***Gnathophyllum americanum*** Guérin-Méneville, 1855

Gnathophyllum americanum Guérin-Méneville 1855, p viii, Plate 2, Figure 14; Chace and Bruce 1993, p 136; Li 1997, p 226, Figure 2; Davie 2002, p 250.

Material examined

Philippines: MUSORSTOM II, shore station, (i) stn 12, opposite Mactan Marine Station, high tide, late afternoon, 0.2–2 m, 9 December 1980, 1 ovig. ♀ (MNHN-Na 15913); (ii) island opposite Mactan Marine Station, reef flat, 2♀♀ (ovig.) (MNHN-Na 15914). **New Caledonia:** (iii) lagoon, reef, Poindimié, north of Tié, associated with anemone, 1 ovig. ♀ (MNHN-Na 14832). **Loyalty Islands:** (iv) ATELIER LIFOU, Lifou, Santal Bay: Easo, near wharf, stn 1406, 20°46.85'S, 167°07.75'E, intertidal, hard substrate, 10/13–14/16/18 November 2000, 1 ovig. ♀ (MNHN-Na 15544).

Parasites

(ii) Bopyrid in branchiae chamber of non-ovigerous female.

Remarks

Not previously recorded from Loyalty Islands. The association with a sea anemone (iii) may have been accidental.

Distribution

Type locality: Cuba. Also known from Red Sea to South Africa, Mauritius, and eastward through Indo-Pacific region to South China Sea (Xisha Islands), Australia (Queensland, New South Wales, Tasman Sea), Loyalty Islands, French Polynesia (Tuamotu Islands, Tahiti, Rikitea); western Atlantic from Bermuda and southern Florida throughout Gulf of Mexico and Caribbean Sea; eastern Atlantic in Canary Islands.

Family PALAEMONIDAE Rafinesque, 1815
Subfamily PALAEMONINAE Rafinesque, 1815
Brachycarpus biunguiculatus (Lucas, 1846)

Palaemon biunguiculatus Lucas 1846, p 45, Plate 4, Figure 4.

Brachycarpus biunguiculatus: Kemp 1925, p 312–314; Holthuis 1952b, p 3–10, Plate 1, Figures a–q; Fransen 1987, p 509, Figure 5; Bruce 1996, p 200, Figures 1a–c, 28a, 30.

Material examined

La Réunion: (i) MD 32, stn CP97, 19°41.4'S, 54°08.7'E, 55 m, pebbles, large slabs of dead reef corals, 28 August 1982, 1♂, 1♀ (MNHN-Na 14942). **New Caledonia:** (ii) MONTROUZIER, Touho (Tié), 5–7 m, coll. B. Richer de Forges, September 1993, 1 ovig. ♀ (MNHN-Na 14808).

Distribution

Type localities: Oran and Bone, Algeria. Also known from the Red Sea, Zanzibar, Sri Lanka, Ryukyu Islands, Caroline Islands, Chesterfield Islands, New Caledonia, Loyalty Islands, Wake Island, and Hawaii. Also known extensively in the Eastern Pacific, Eastern and Western Atlantic, and western Mediterranean region.

Leander tenuicornis (Say, 1818)

Palaemon tenuicornis Say 1818, p 249.

Leander tenuicornis: Kingsley 1878, p 66; Kemp 1925, p 302–304, Figure 11; Holthuis 1952b, p 155–167, Plates 41–42; Bruce 1991b, p 223, Figures 1b, 2; Chace and Bruce 1993, p 6; Bruce 1996, p 202; Bruce 2002a, p 80; Davie 2002, p 292; Li et al. 2004, p 516, Figure 3.

Material examined

Madagascar: Tuléar, coll. R. Hipeau-Jacquotte, 1960, (i) 1♂ (MNHN-Na 15926); (ii) 1 ovig. ♀ (MNHN-Na 15927); (iii) 2 ovig. ♀♀ (MNHN-Na 15925). **New Caledonia:** (iv) MONTROUZIER, Koumac, Infernet Reef, coll. B. Richer de Forges, scuba, 13 m, 5 October 1993, 1 ovig. ♀ (MNHN-Na 14820); (v) Ricaudy Reef, intertidal, coll. B. Richer de Forges, 12 July 1995, 1 ovig. ♀ (MNHN-Na 15875); (vi) lagoon, Ricaudy reef flat, low tide, coll. B. Richer de Forges, 22 June 1997, 1♂ (MNHN-Na 15821); (vii) lagoon (Seiche Croissant), seagrass, small trawl, coll. B. Richer de Forges, 7 May 1998, 3♀♀ (1 ovig.) (MNHN-Na 15815). **Loyalty Islands:** (viii) ATELIER LIFOU, Lifou, Santal Bay: Enu, near wharf, stn 1417, 20°46.9'S, 167°08.3'E, 1–4 m, slabs with sediment cover and coral blocks, 8–24 November 2000, 1♀ (MNHN-Na 15431).

Remarks

Not previously recorded from Loyalty Islands. The association with anemones may have been accidental. The Madagascar material (i–iii) was collected from shelter of an actinarian *Actinodendron* sp.

Distribution

Type localities: Newfoundland Bank. Also known from Red Sea and South Africa to South China Sea, Japan, Philippines, Indonesia, Australia, Caroline Islands, New Zealand, and the Atlantic Ocean from Newfoundland to Brazil, and Mediterranean. Associated with floating weed in the open sea and with attached plants in shallow water.

Macrobrachium australe (Guérin-Méneville, 1838)

Palaemon australis Guérin-Méneville 1838, p 37.

Macrobrachium australis: Chace and Bruce 1993, p 23, Figure 2; Davie 2002, p 293.

Material examined

New Caledonia: west coast (Le Cap), mangroves, coll. B. Richer de Forges, 5 December 1992, 1♂ (MNHN-Na 14821).

Remarks

Not previously recorded from New Caledonia.

Distribution

Type locality: Tahiti. Also known from Madagascar, Seychelles through the Indian Ocean to China (Taiwan), Philippines, Indonesia, and the Pacific islands as far as the Marshall Islands in the North Pacific and the Marquesas Islands in the South Pacific.

Macrobrachium equidens (Dana, 1852)

Palaemon equidens Dana 1852, p 26.

Macrobrachium equidens: Holthuis 1950, p 162, Figure 36; Chace and Bruce 1993, p 25, Figure 4; Davie 2002, p 295.

Material examined

New Caledonia: west coast (Le Cap), mangrove (in burrows), coll. B. Richer de Forges, 4 December 1992, 2♂♂ (MNHN-Na 14822).

Remarks

Not previously recorded from New Caledonia. Usually found in high salinity brackish and salt water, rarely in pure fresh water.

Distribution

Type locality: Singapore. Also known from South Africa, southern India to China (Fujian), Ryukyu Islands, Philippines, Indonesia, Solomon Islands, and Nigeria (possibly introduced; Chace and Bruce 1993).

Nematopalaemon tenuipes (Henderson, 1893)

Leander tenuipes Henderson 1893, p 440, Plate 40, Figures 14, 15.

Palaemon (Nematopalaemon) tenuipes: Holthuis 1950, p 44, Figure 7.

Nematopalaemon tenuipes: Holthuis 1980, p 108; Chace and Bruce 1993, p 39.

Material examined

Madagascar: (i) west coast, in region of Maintirano, coll. A. Crosnier, 1959, 1 ovig. ♀ (MNHN-Na 14828); (ii) same, 1♀ (MNHN-Na 14829). **Indonesia:** (iii) Moluccas, Ambon, Baguala Bay, 19 October 1982, 3♀♀ (2 ovig.) (MNHN-Na 14827).

Remarks

Indonesian specimens have the rostral formula $2+5+?/6^+$ in the female and $2+3+1/6$, $2+4+1/5$ in the ovigerous females. Both of the Maintirano specimens, with the rostrum broken, with seven rostral crest teeth. The ovigerous female (i) with the distal part distal to antennular peduncle lost, the female (ii) with the distal part except the apex linked with the basal part. Not previously recorded from Madagascar and Indonesia.

Distribution

Type localities: Bombay and Madras, India and Gulf of Martaban, Burma. Also known from South Africa, Madagascar, India, Thailand, China, Philippines, and Indonesia.

Palaemon debilis Dana, 1852

Palaemon debilis Dana 1852, p 26; Bruce 1991b, p 227, Figures 1d, 3f; Chace and Bruce 1993, p 40; Bruce 1996, p 202; Davie 2002, p 299; Li et al. 2004, p 521, Figure 9.

Palaemon (Palaemon) debilis: Holthuis 1950, p 66–70, Figure 13.

Material examined

Île Europa: (i) lagoon, coll. P. Fourmanoir, 4♂♂, 13♀♀ (5 ovig.) (MNHN-Na 8460). **Philippines:** (ii) MUSORSTOM II, shore station, Silot Bay, mangrove, 15 December 1980, 1♂, 1♀ (MNHN-Na 15920). **New Caledonia:** (iii) west coast (Le Cap), mangrove, coll. B. Richer de Forges, 2 December 1992, 3 ovig. ♀♀ (MNHN-Na 14831); (iv) west coast (Le Cap), mangrove (in burrows), coll. B. Richer de Forges, 3 December 1992, 6 juveniles (MNHN-Na 14833); (v) south coast, Goro Reef flat, at low tide, coll. B. Richer de Forges, 11 January 1993, 1♂ (with bopyrid), 1 ovig. ♀ (MNHN-Na 14830).

Parasites

(i) *Diplophryxus jordani* Richardson (Isopoda: Bopyridae).

Remarks

Not previously recorded from Île Europa.

Distribution

Type locality: Hilo, Hawaii. Common species throughout most of the Indo-West Pacific region from the Gulf of Suez to French Polynesia (Tuamotu Islands).

Palaemonetes atrinubes Bray, 1976

(Figure 1)

Palaemonetes atrinubes Bray 1976, p 76–82, Figures 23–42; Bruce 1988b, p 115–117, Figure 1; Davie 2002, p 301.

Material examined

New Caledonia: “New Caledonia”, 1903, 11 specimens (dried) (MNHN-Na 14843).

Remarks

There were no collection data with the specimens, other than “New Caledonia”. The specimen label was in the handwriting of A. Milne Edwards (J. Forest, personal communication), all the material is dried, but they can still be checked.

Previously recorded from New Caledonia by Bruce (1995).

Distribution

Type locality: Lower Swan River. Also known from Australia (Western Australia, Queensland) and New Caledonia.

Urocaridella antonbruunii (Bruce, 1967)

Periclimenes antonbruunii Bruce 1967a, p 45–53, Figures 19–22.

Leandrites cyrtorhynchus Fujino and Miyake 1969, p 143–149, Figures 1–3.

Urocaridella antonbruunii: Chace and Bruce 1993, p 7; Bruce 1996, p 203, Figures 1f, 29a, 31; Davie 2002, p 301; Li et al. 2004, p 525, Figure 13.

Material examined

Madagascar: (i) Nosy Be, shallow waters, coll. A. Crosnier, 1966, 1♂, 1♀ (MNHN-Na 15953). **La Réunion:** (ii) MD 32, stn CA88, 19°44.1'S, 54°08.9'E, 55–60 m, 28 August 1982, 1♂ (MNHN-Na 14933). **New Caledonia:** (iii) MONTROUZIER, Touho Bay, coll. B. Richer de Forges, September 1993, 1 ovig. ♀ (MNHN-Na 14932).

Remarks

Not previously recorded from Madagascar and La Réunion. The collection depth of the material from La Réunion is much deeper than the records from Northern Territory (1–3 m, Bruce and Coombes 1995; 12 m, Bruce 1983a), New Caledonia (4–12 m, Bruce 1996), Japan (5 m, Minemizu et al. 2000, p 42), Comoro Islands (20 m, Bruce 1967a),




Figure 1. *Palaemonetes atrinubes* Bray, 1976, ovigerous female (MNHN-Na 14843). (a) Carapace and anterior appendages; (b) anterior carapace; (c) distal part of proximal peduncular segment; (d) antennal scale; (e) distal telson; (f) mandible; (g) first pereiopod; (h) same, chela; (i) second pereiopod; (j) same, chela; (k) third pereiopod; (l) same, propod and dactyl; (m) same, distal propod and dactyl.

Bali, Indonesia (26 m, Debelius 1999). However, Li et al. (2004) recorded that the collection depth of the species from northern South China Sea is 60 m, as well as the present record from La Réunion.

Distribution

Type locality: Dzaoudzi, Mayotte (Pamanzi Island), Comoro Islands. Also known from Kenya, La Réunion, South China Sea, Japan, Australia, New Caledonia, and Hawaii.

Urocaridella urocaridella (Holthuis, 1950)

Urocaridella gracilis Borradaile 1915, p 210.

Leander urocaridella Holthuis 1950, p 28.

Urocaridella urocaridella: Chace and Bruce 1993, p 42–44, Figure 16; Bruce 1996, p 203; Davie 2002, p 302; Li et al. 2004, p 527, Figure 14.

Material examined

Philippines: (i) MUSORSTOM 3, stn CP142, 11°47'N, 123°02'E, 26–27 m, 6 June 1985, 2♂♂, 1 ovig. ♀ (MNHN-Na 14936); (ii) same, 2♂♂, 1 ovig. ♀ (MNHN-Na 14935). **New Caledonia:** (iii) BATHUS 1, stn CP680, 20°48'S, 165°18'E, 86–92 m, 15 March 1993, 1♂, 1 ovig. ♀ (MNHN-Na 14934). **Solomon Islands:** (iv) SALOMON 1, stn CP 1823, 9°50.4'S, 160°53.2'E, 82–83 m, 4 October 2001, 1♂ (MNHN-Na 15434).

Distribution

Type locality: Maldives. Also known from northeastern India, Andaman Islands, Burma, Malaya, Singapore, northern South China Sea, Philippines, Indonesia, and New Caledonia.

Subfamily PONTONIINAE Kingsley, 1878

Allopontonia iaini Bruce, 1972

Allopontonia iaini Bruce 1972, p 7–11, Figures 1–5; Li 2000, p 1, Figure 1; Davie 2002, p 304.

Material examined

New Caledonia: (i) CORAIL 1, Lansdowne Bank, August 1988, 1 ovig. ♀ (MNHN Na-14798); (ii) MONTROUZIER, Koumac, reef outer slope, 10 October 1993, 1 ovig. ♀ (MNHN-Na 14799).

Host

(ii) *Pseudoboletia indiana* (Michelin, 1862) [Toxopneustidae, Echinoidea].

Remarks

The Lansdowne Bank specimen closely resembles those previously described. It is of normal size and in good condition and lacks only the tip of the rostrum. The rostrum has a dentition of at least eight dorsal and two ventral teeth and is of normal shape. Both second pereopods are preserved and are markedly unequal, the minor second pereopod slightly less than half the length of the very robust major pereopod, which is about 1.75 times the carapace length of 4.75 mm.

The Koumac specimen is rather distorted and is noticeably smaller, with carapace length of about 2.76 mm. The body appears to have been rather more strongly compressed than usual. One detached second pereopod is preserved: all pereopods are missing from the right side of the body.

The association with *Pseudoboletia* represents a new host record. The type material was collected in association with *Salmaciella erythracis* (H. L. Clark, 1912). Other specimens have been associated with the genera *Salmacis* and *Asthenosoma*.

Not previously recorded from New Caledonia.

Distribution

Type locality: Heron Island. Also reported from Kenya, Zanzibar, Australia (Queensland), New Caledonia, and Mexico (Baja California).

Altopontonia disparostris Bruce, 1990

Altopontonia disparostris Bruce 1990a, p 192–202, Figures 25–33, 39k; Bruce 1991a, p 390; Bruce 1996, p 204; Bruce 2005, p 2, Figure 8A; Li 2000, p 2, Figure 2.

Material examined

Vanuatu: MUSORSTOM 8, (i) stn CP1049, 16°39.43'S, 168°02.97'W, 469–525 m, 1 October 1994, 1 ovig. ♀ (MNHN-Na 15931); (ii) stn CP1088, 15°09.23'S, 167°15.13'W, 425–455 m, 6 October 1994, 2 ovig. ♀♀ (MNHN-Na 15933). **New Caledonia:** NORFOLK 2, (iii) Île des Pins, stn DW 2156, 22°54'S, 167°15'W, 468–500 m, 5 November 2003, 1 ovig. ♀ (MNHN-Na 15932). **Fiji:** BORDAU 1, (iv) stn DW 1492, 18°43'S, 178°23'W, 430–450 m, 11 March 1999, 2♀♀ (1 ovig.) (MNHN-Na 15436); (v) stn CP 1505, 18°12'S, 178°37'W, 420–450 m, 13 March 1999, 10 ovig. ♀♀ (MNHN-Na 15435).

Remarks

Not previously recorded from Vanuatu and Fiji. Bruce (2005) illustrated the coloration.

Distribution

Type locality: off New Caledonia. Also known from Tasman Sea, Vanuatu and Fiji; 322–525 m.

Amphipontonia kanak Bruce, 1991

Amphipontonia kanak Bruce 1991a, p 382–390, Figures 58–63.

Material examined

New Caledonia: BIOCAL, stn DW37, 23°00'S, 167°16'E, 350 m, 30 August 1985, 1♀ (MNHN-Na 15633).

Remarks

The present record slightly extends this species bathymetric range, from 300 to 350 m.

Distribution

Type locality: New Caledonia and Loyalty Islands.

Anchistus custos (Forskål, 1775)

Cancer custos Forskål 1775, p xxi, 94.

Harpilius inermis Miers 1884, p 291, Plate 32, Figure B.

Anchistus inermis: Borradaile 1898, p 387.

Anchistus custos: Holthuis 1952a, p 105–109, Figures 43, 44; Chace and Bruce 1993, p 72; Bruce 1996, p 205; Li 2000, p 7, Figure 8; Davie 2002, p 305; Li and Liu 2002, p 371, Figure 1; Li et al. 2004, p 528.

Material examined

Madagascar: Tuléar, coll. R. Hipeau-Jacquotte, (i) 19 March 1968, 84 specimens (spms) (36 ovig. ♀♀) (MNHN-Na 15958); (ii) 50 spms (MNHN-Na 15970). **New Caledonia:** (iii) MONTROUZIER, Koumac, scuba, coll. B. Richer de Forges, 6 September 1993, 2♂♂, 3 ovig. ♀♀ (MNHN-Na 14806); (iv) lagoon, Île Ouen-Prony Bay, stn 69, 22°23'S, 166°32'E, 13 m, coll. B. Richer de Forges, no date record, 1♂ (MNHN-Na 11433[e]); (v) Nouméa, associated with “grand *Pinna*”, coll. J. Risbec (Th. Monod Collection number 1779), 3♂♂, 2 ovig. ♀♀ (MNHN-Na 15941).

Host

(i) (ii) *Atrina* sp. [Pinnidae, Mollusca].

Remarks

Previously recorded from New Caledonia at Saint Vincent Bay (Bruce 1996).

Distribution

Type locality: Al Luhayyah, Yemen. Known from Red Sea and eastern Africa to Philippines, southward to Australia (South Australia), and eastward to the Caroline Islands and Fiji.

Anchistus miersi (De Man, 1888)

Harpilius Miersi De Man 1888, p 274, Plate 17, Figures 6–10.

Anchistus miersi: Borradaile 1898, p 387; Bruce 1978a, p 279; Chace and Bruce 1993, p 72; Li 2000, p 11, Figure 11; Davie 2002, p 305; Li 2004a, p 67.

Material examined

Madagascar: northwest coast, Nosy Be (Andilah), intertidal, in *Tridacna* spp., coll. A Crosnier, 1959, 2♂♂, 2♀♀ (1 ovig.), 1 juvenile (MNHN-Na 14807).

Remarks

In all specimens, the lateral posterior telson spines were situated subterminally on the dorsum of the telson, and the female specimen with two pairs of median posterior telson spines. Previously recorded from Madagascar (Nosy Be) by Bruce (1978a).

Distribution

Type locality: Mergui Archipelago. Known from Red Sea and eastern Africa to the Philippines, Indonesia, Australia (Queensland), and eastward to French Polynesia (Tuamotu Islands and Gambier Islands).

Anchistus pectinis Kemp, 1922

Anchistus pectinis Kemp 1925, p 327; Bruce 1991a, p 378, Figures 56–57, 71b–f; Bruce 1991b, p 261, Figure 24; Bruce 1996, p 205, Figures 2–3; Li 2000, p 12, Figure 12.

Material examined

New Caledonia: BATHUS 4, 19°49.00'S, 163°48.40'E, 37 m, 9 August 1994, 1♂, 1♀, 4 ovig. ♀♀ (MNHN-Na 15894).

Remarks

Previously recorded from New Caledonia by Bruce (1991b).

Distribution

Type locality: Nicobar Islands. Also known from Zanzibar, Nicobar Islands, Japan, northeast Australia, and New Caledonia, 10–110 m. Associated with pectinid mollusks *Amusium*, *Pecten*, and *Semipallium*.

Climeniperaeus truncoideus (Chace and Bruce, 1993)

Periclimenaeus truncatus: Holthuis 1952a, p 117, Figures 48–50; Bruce 1981a, p 211, Figures 16, 17d, 18c, f (non *Coralliocaris truncata* Rathbun 1906).

Periclimenaeus truncoideus Chace and Bruce 1993, p 93–94.

Climeniperaeus truncoideus: Bruce 1996, p 210, Figures 4, 5; Li 2000, p 22, Figure 23.

Material examined

Providence Island: (1) ?stn D11, 91.5 m, 4 October ?2003, 2 spms (dried) (MNHN-Na 15945). **Philippines:** (ii) MUSORSTOM 3, stn DR117, 12°31'N, 120°39'E, 92–95 m, 3 June 1985, 1♀ (MNHN-Na 11333[f]).

Remarks

Previously recorded from Philippines by Bruce (1981a, 1996). Not previously recorded from Providence Island.

Distribution

Type locality: Indonesia. Also known from Zanzibar and Philippines.

Conchodytes biunguiculatus (Paulson, 1875)

Pontonia biunguiculata Paulson 1875, p 111, Plate 15, Figure 1.

Conchodytes biunguiculatus: Kemp 1922, p 279, 280–282, Figure 102; Holthuis 1952a, p 17, 199–200; Jacquotte 1963, p 61; Franssen 1994, p 89, Figures 3–7, 12–15, 23, 27, 30, 35; Li 2000, p 24, Figure 24; Li 2001, p 76; Li and Liu 2002, p 372, Figure 2a–f; Bruce 2003, p 212.

Conchodytes kempii Bruce 1989a, p 183.

Material examined

Madagascar: Tuléar, coll. R. Hipeau-Jacquotte, (i) 45 spms (MNHN-Na 15968); (ii) 19 March 1968, 2♂♂, 3 ovig. ♀♀, 1 juvenile (MNHN-Na 15915); (iii) 12 April 1968, 1♂, 1 ovig. ♀ (MNHN-Na 15923).

Hosts

(i) (ii) *Atrina* sp.; (iii) *Atrina vexillum* (Born, 1778) [Pinnidae, Mollusca].

Remarks

First reported from Madagascar at Toliara (Tuléar) by Jacquotte (1963) and subsequently studied in detail.

Distribution

Type locality: Red Sea. Also known from Western Indian Ocean to South China Sea, Philippines, Indonesia, and Marshall Islands.

Conchodytes maculatus Bruce, 1989

Conchodytes maculatus Bruce 1989b, p 182, Figures 1–6; Li 2000, p 25, Figure 25.

Material examined

New Caledonia: MONTROUZIER, Touho Sector, Sand Islet, Touho Pass, stn 1272, 20°49.5'S, 165°19.6'E, 10 m, hard bottom with mud, September 1993, 1 ovig. ♀ (MNHN-Na 15885).

Remarks

Not previously recorded from New Caledonia. Known from intertidal zone to 40 m depth, in association with pearl oysters.

Distribution

Type locality: Australian Northwest Shelf. Also known from Philippines and New Caledonia.

Conchodytes meleagrinae Peters, 1852

Conchodytes meleagrinae Peters 1852, p 594; Bruce 1978a, p 279; Bruce 1991b, p 262, Figure 25a–d; Chace and Bruce 1993, p 74; Li 2000, p 25, Figure 26; Davie 2002, p 307; Li 2004a, p 67.

Material examined

Île Europa: (i) 1♂ (MNHN-Na 14810). **Malaya (or Vietnam):** (ii) associated with *Meleagrina* sp., 1♂ (MNHN-Na 15939). **New Caledonia:** (iii) reef inside Abore, 4 m, with *Pinctada margaritifera*, coll. P. Laboute, 8 November 1987, 1♂, 1♀ (MNHN-Na 15949); (iv) MONTROUZIER, Grand Récif, Mangalia, in pearl oysters, coll. B. Richer de Forges, 15 September 1993, 1♂, 2 ovig. ♀♀ (MNHN-Na 14809). **Loyalty Islands:** (v) ATELIER LIFOU, Lifou, Santal Bay: west or southwest of Easo Point, stn 1429, 20°47.5'S, 167°07.1'E, 8–18 m, sedimentary channels, 3/5/23–24 November 2000, 1 ovig. ♀ (MNHN-Na 15437).

Remarks

Previously recorded from New Caledonia on Touaorou Reef, Yaté (Bruce 1991b). Previously recorded from Madagascar: Île Europa by Bruce (1978a). Not previously recorded from Loyalty Islands.

Distribution

Type locality: Mozambique (Ibo, Cabo Delgado), southeastern coast of Africa. Also known from Red Sea eastward to Hawaii.

***Coralliocaris* sp.** Mitsuhashi and Takeda, 2006*Material examined*

Loyalty Islands: ATELIER LIFOU, Lifou. Chateaubriand Bay: (i) Gaatcha Bay, stn 1463, 20°55.05'S, 167°03.35'E, 30 m, dredge, sand and coral debris, 10 November 2000, 1♀ (MNHN-Na 15442); (ii) Wé Beach, stn 1474, 20°54.8'S, 167°16.1'E, 30 m, fine sand and heads of *Porites*, 11 November 2000, 1 ovig. ♀ (MNHN-Na 15439). Santal Bay: (iii) northeast Chateaubriand Bay, Cila, stn 1456, 20°49.3'S, 167°10.4'E, slope, 25–30 m, 26 November 2000, 1 ovig. ♀ (MNHN-Na 15444).

Remarks

Only the ovigerous female of (ii) has the second pereopods present. Rostrum reaches distal margin of first antennular peduncle segment, armed with two or three dorsal and one ventral tooth, the ventral tooth minute and subapical. Antennal spine on carapace relatively stout and long, reaches almost to the distal margin of basicerite. The depth of 30 m is a particularly deep record for this genus.

Dr. M. Mitsuhashi is reviewing the *Coralliocaris nudirostris* (Heller, 1861) and its closely related species. After checking the present specimens, she considers that it should be a species close to *C. nudirostris* (personal communication).

Distribution

Type locality: Japan (Ryukyu Islands), Papua New Guinea, Loyalty Islands, Australia (Queensland). The other localities outside the type localities should be re-examined (see Remarks).

Coralliocaris macrophthalma (H. Milne-Edwards, 1837)

Pontonia macrophthalma H. Milne Edwards 1837, p 359.

Coralliocaris macrophthalma Borradaile 1917, p 383; Holthuis 1952a, p 17; Chace and Bruce 1993, p 47, 77; Bruce 1998, p 30.

Coralliocaris graminea: Bruce 1977a, p 204, Figure 1; Li 2000, p 33.

Material examined

New Caledonia: lagoon, Amédée Reef flat, from seagrass, coll. C. Vadon, 17 September 1978, 1 ovig. ♀ (MNHN-Na 15862).

Remarks

Not previously recorded from New Caledonia.

Distribution

Type locality: seas of Asia. Also known from Israel, Red Sea, Seychelles, Saya de Malha Australia (Great Barrier Reef (?)), and New Caledonia.

Coralliocaris superba (Dana, 1852)

Oedipus superbus Dana 1852, p 25.

Oedipus dentirostris: Paulson 1875, p 112, Plate 14, Figure 7.

Coralliocaris superba: Stimpson 1860, p 38; Hipeau-Jacquotte 1973, p 104; Bruce 1991b, p 264, Figure 26; Chace and Bruce 1993, p 77; Li 2000, p 35, Figure 35; Davie 2002, p 309; Li and Liu 2002, p 375, Figure 4a–d.

Material examined

Madagascar: (i) west coast, Toliara (Tuléar), intertidal, coll. A. Crosnier, October 1958, 1 ovig. ♀ (MNHN-Na 14812). **New Caledonia:** (ii) MONTROUZIER, Touho (Tié), coll. B. Richer de Forges, scuba, 5–7 m, 1 ovig. ♀ (MNHN-Na 14811).

Remarks

The two specimens, which both lack second pereopods, have rostral dentitions of 5/2 and 6/5, respectively. The ambulatory dactyls have a particularly acute tip to the hoof-shaped ventral process. Previously recorded from New Caledonia in Nouméa and Île Ouen, Baie du Prony (Monod 1976; Bruce 1991b). Previously recorded from Madagascar (Tuléar) by Hipeau-Jacquotte (1973).

Distribution

Type locality: Tongatabu Island, Tonga. Also known from Red Sea, Mozambique and Madagascar to Indonesia, Australia (Western Australia, Queensland, New South Wales, Tasman Sea), eastward to French Polynesia (Society Islands).

Coralliocaris viridis Bruce, 1974

Coralliocaris viridis Bruce 1974a, p 222–224, Figure 1; Chace and Bruce 1993, p 78; Li 2000, p 38, Figure 38; Davie 2002, p 309.

Material examined

Madagascar: (i) northwest coast, Nosy Be (Ambatoloaka), coll. A. Crosnier, September 1958, 1 ovig. ♀ (MNHN-Na 14813); (ii) Îles Glorieuses, intertidal, coll. A. Crosnier, September 1958, 1♂, 1 ovig. ♀ (MNHN-Na 14814).

Remarks

The Îles Glorieuses specimens lack second pereopods but have very slender rostra with a dentition of 4/1 and 5/1, respectively. The Nosy Be specimen has both second pereopods and a rostral dentition of 5/2, all teeth distally situated. Not previously recorded from Madagascar.

Distribution

Type locality: Mombasa Island, Kenya. Also known from Mozambique, Seychelles, Maldive Islands, Sri Lanka, Vietnam, Ryukyu Islands, Indonesia, Papua-New Guinea, and Australia (Northern Territory and Queensland).

Dasycaris ceratops Holthuis, 1952

Dasycaris ceratops Holthuis 1952a, p 176, Figures 87, 88; Bruce 1977b, p 173, Figure 5; Chace and Bruce 1993, p 80; Li 2000, p 43, Figure 44.

Material examined

Madagascar: Nosy Be, shallow water, coll. A. Crosnier, 1966, 1♂, 1 ovig. ♀ (MNHN-Na 15964).

Remarks

Not previously recorded from Madagascar.

Distribution

Type localities: Boeo Bank, Indonesia. Also known from Zanzibar, Madagascar, China, Indonesia, and Australia (Queensland).

Dasycaris symbiotes Kemp, 1922

Dasycaris symbiotes Kemp 1922, p 240, Figures 76, 77, Plate 9; Monod 1976, p 147, Figures 54–60; Li 1997, p 230, Figure 6; Li 2000, p 44, Figure 45.

Material examined

New Caledonia: Dumbéa Bay, 22°13.38'S, 166°22.17'E, scuba, 20 m, coll. Bargibant, 30 November 1995, 1♂ (MNHN-Na 15893).

Remarks

Previously recorded from New Caledonia by Monod (1976).

Distribution

Type localities: India and Mergui Archipelago, 27–31 m, 64 m. Also known from South China Sea (Xisha Islands) and New Caledonia. Associated with sea pen, *Pteroeides* sp.

Dasycaris zanzibarica Bruce, 1973

Dasycaris zanzibarica Bruce 1973b, p 247–257, Figures 1–6; Bruce 1991b, p 265, Figure 27; Bruce 1996, p 214; Li 2000, p 45, Figure 46; Davie 2002, p 310.

Material examined

Madagascar: west coast, Pracel Bank, 25 m, beam trawl, on brown mud, coll. A. Crosnier, June 1959, 1 ovig. ♀ (MNHN-Na 14815).

Remarks

Not previously recorded from Madagascar. Also reported from 39 m at Atoll de Surprise, New Caledonia, and otherwise recorded only from shallow water.

Distribution

Type locality: Changu Island, Zanzibar. Also known from Japan, Taiwan, Philippines, Papua New Guinea, Australia (Western Australia, Great Barrier Reef), and New Caledonia.

Exoclimenella denticulata (Nobili, 1906)

Periclimenes Petitthouarsi var. *denticulata* Nobili 1906a, p 257; Nobili 1907, p 358–359.

Periclimenes (Falciger) denticulatus: Borradaile 1917, p 368, 372.

Periclimenes denticulatus: Bruce 1992, p 59, Figures 12–14.

Exoclimenella denticulata: Duris and Bruce 1995, p 637, Figures 9–11; Li 2000, p 49, Figure 52.

Material examined

Loyalty Islands: ATELIER LIFOU, Lifou, Xepenehe Point, in coral mats, scuba, 12–13 m, 11 November 2000, 1 ♂ (MNHN-Na 15446).

Remarks

The specimen has the dorsal surface of fixed finger of major second pereiopod with a very deep longitudinal groove. Not previously recorded from Loyalty Islands.

Distribution

Type locality: Gatavake, Mangareva Atoll, Gambier Islands. Also known from South China Sea, Australia (Great Barrier Reef), Marshall Islands, and French Polynesia (Tuamotu Islands).

Exoclimenella cf. denticulata (Nobili, 1906)*Material examined*

Loyalty Islands: ATELIER LIFOU, Lifou, Santal Bay: in front of Peng, stn 1464, 20°54.5'S, 167°05.9'E, 30–50 m, dredges, blocks of calcareous algae, 14 November 2000, 1 ovig. ♀ (MNHN-Na 15447).

Remarks

Carpus of second pereopods is distinctly longer than that of the original description.

Exoclimenella maldivensis Duris and Bruce, 1995

Exoclimenella maldivensis Duris and Bruce 1995, p 622, Figures 1–5; De Grave 2000, p 123.

Material examined

Loyalty Islands: ATELIER LIFOU, Lifou, Santal Bay, (i) west or southwest of Easo Point, stn 1429, 20°47.5'S, 167°07.1'E, 8–18 m, sedimentary channels, 3/5/23–24 November 2000, 1♂, 1♀, 1 juvenile (MNHN-Na 15449); (ii) east of bay, in front of Huca Hutighé, stn 1460, 20°52.4'S, 167°08.0'E, 40–60 m, dredge, 6 November 2000, 1♂ (MNHN-Na 15454); (iii) Gaatcha Bay, stn 1463, 20°55.05'S, 167°03.35'E, 20–30 m, dredge, sand and coral debris, 10 November 2000, 1♂ (MNHN-Na 15455); (iv) east of Santal Bay: Mepinyo, stn 1446, 20°50.8'S, 167°09.7'E, 36–40 m, bottom of slope, 16 November 2000, 1 ovig. ♀ (MNHN-Na 15452); (v) north of Cap Aimé Martin (=Acadro), stn 1450, 20°45.8'S, 167°01.65'E, 27–31 m, washings, 21 November 2000, 4♂?♂, 4♀♀ (2 ovig.) (MNHN-Na 15457); (vi) between Cap Mandé and Cap Lefèvre (=Nem), stn 1452, 20°54.6'S, 167°02.1'E, 2–25 m, scuba, 22 November 2000, 1 juvenile (MNHN-Na 15456); (vii) between Cap Wekutr and Cap Wajez, stn 1455, 20°56.8'S, 167°02.7'E, slope, 15–20 m, 25 November 2000, 1♂, 1 ovig. ♀ (MNHN-Na 15450); (viii) northeast Santal Bay, in region of Cila, stn 1456, 20°49.3'S, 167°10.4'E, slope, 25–30 m, 26 November 2000, 2♂♂, 1 ovig. ♀ (MNHN-Na 15451); (ix) in front of Ngoni, stn 1457, 20°46.8'S, 167°02.75'E, 5–10 m, dark overhang and boulders, 27 November 2000, 1♂, 1 ovig. ♀ (MNHN-Na 15453).

Remarks

Not previously recorded from Loyalty Islands. The specimen (i) with 2+7/5 rostral dentition, the subapical dorsal tooth tiny; the female and juvenile of (vii), male of (iii) lack both second pereopods, the male also lacks the carapace and the appendages anterior to the second pereopods, but both the pereopods are present; specimens of (iii) and (vi) lack both second pereopods and (vii) the rostrum is broken.

Distribution

Type locality: Maldives, Timor Sea. Also known from Thailand, Japan, and Loyalty Islands.

Hamodactylus boschmai Holthuis, 1952

Hamodactylus boschmai Holthuis 1952a, p 209, Figures 102–104; Bruce 1970a, p 538, Figure 1; Bruce 1982, p 272, Figures 25, 26; Bruce and Coombes 1995, p 108; Li 2000, p 58, Figure 61.

Material examined

New Caledonia: lagoon, with crinoids, 30 August 1993, 1 spm (MNHN-Na 15840).

Remarks

The only specimen was damaged. Rostrum, telson, all pleopods and most of ambulatory pereopods lost, but both second, left first and left fifth pereopods are present. The species was first reported from New Caledonia near Nouméa, by Bruce (1970a). The association with crinoids is probably accidental.

Distribution

Type locality: Ternate, Aru Islands, Indonesia, 2–13 m. Also known from Zanzibar, Kenya, Madagascar, Indonesia, and New Caledonia.

Hamodactylus noumeae Bruce, 1970

Hamodactylus noumeae Bruce 1970a, p 539–541, Figure 2; Chace and Bruce 1993, p 80; Li 2000, p 59, Figure 62; Davie 2002, p 312.

Material examined

New Caledonia: MONTROUZIER, Koumac Pass, 35 m, gorgonian washings, 28 October 1993, 2♂♂, 1♀ (with branchial bopyrid) (MNHN-Na 14816).

Remarks

The female specimen has five dorsal rostral teeth and both males have four. This species was first described from Nouméa, New Caledonia (Bruce 1970a), but there have been no subsequent reports from these islands.

Distribution

Type locality: Nouméa, New Caledonia. Also known from Zanzibar, Tanzania, Kenya, Seychelles, Japan, and Australia (Western Australia, Northern Territory, Queensland).

***Hamopontonia corallicola* Bruce, 1970**

Hamopontonia corallicola Bruce 1970b, p 41, Figures 1–4; Bruce 1983b, p 896, Figure 10G; Li 2000, p 59, Figure 63.

Material examined

Philippines: MUSORSTOM II, shore station, stn 11, opposite Mactan Marine Station, scuba, 5–18 m, with fungiid coral, 9 December 1980, 1♂ (MNHN-Na 15960).

Remarks

Not previously reported from the Philippines.

Distribution

Type locality: Hong Kong. Also known from Japan, Philippines, Indonesia, and Australia (Great Barrier Reef).

***Harpiliopsis beaupresii* (Audouin, 1825)**

Palaemon Beaupresii Audouin 1825, p 91.

Harpilius Beaupresii: Heller 1861, p 27.

Harpiliopsis beaupresi: Borradaile 1917, p 324, 379, Plate 55, Figure 21; Chace and Bruce 1993, p 82; Li 2000, p 61, Figure 65; Davie 2002, p 312; Li and Liu 2003, p 154, Figure 1.

Material examined

Madagascar: (i) Ambataloaka, Nosy Be, coll. A. Crosnier, September 1958, 1 ovig. ♀ (MNHM-Na 14817). **La Réunion:** (ii) no date, coll. S. Ribes, 1 juvenile (MNHN-Na 14818). **Loyalty Islands:** ATELIER LIFOU, (iii) Lifou, Santal Bay: in front of Huca Hutighé Islet, stn 1434, 20°52.5'S, 167°08.1'E, 5–20 m, hard bottom, 6 November 2000, 2♂♂ (MNHN-Na 15445).

Host

Pocillopora damicornis (L., 1758) [Pocilloporidae, Scleractinia].

Remarks

Both specimens had a rostral dentition of 5/2. Previously reported from both La Réunion (Bruce 1983c) and Madagascar (Hipeau-Jacquotte 1973), but not Loyalty Islands.

Distribution

Type locality: Egyptian Red Sea. Also known from Gulf of Aqaba, Sudan, Ethiopia (Eritrea), Yemen, Kenya, Zanzibar, Mozambique, Madagascar, Seychelles, La Réunion, Mauritius, Maldives, Chagos Islands, Sri Lanka, Andaman Islands, Thailand, Singapore, Vietnam, South China Sea, Japan, Philippines, Indonesia, Australia (Western Australia, Northern Territory, Queensland), Coral Sea, Loyalty Islands, Fiji, Marshall Islands, USA (French Frigate Shoal, Johnson Atoll, Hawaii), and Easter Island.

Harpiliopsis depressa (Stimpson, 1860)

Harpilius depressus Stimpson 1860, p 38; Kemp 1922, p 231, Figures 69, 70.

Harpiliopsis depressus: Borradaile 1917, p 380; Holthuis 1951, p 70, Plates 21, 22, Figures a–f; Holthuis 1952a, p 182, Figure 90; Bruce 1970d, p 306; Bruce 1991b, p 263; Franssen 1994, p 107, Figures 59, 61; Li 1997, p 233; Li 1998, p 222, Figures 8–11; De Grave 2000, p 124.

Periclimes pusillus Rathbun 1906, p 921, Figure 71, Plate 24, Figure 7.

Material examined

New Caledonia: Boulari Pass, stn 505, scuba, on *Pocillopora* sp., coll. J.-L. Menou, 12 April 1994, 1♂ (MNHN-Na 15854).

Remarks

Previously recorded from New Caledonia by Bruce (1991b).

Distribution

Type locality: Hawaii. Common and widespread throughout most of the Indo-Pacific region: Egypt, Israel, Saudi Arabia, Sudan, Yemen, Kenya, Zanzibar, Mozambique, Comoro Islands, Madagascar, Seychelles, La Réunion, Maldives, Chagos Islands, Sri Lanka, Andaman Islands, Nicolar Islands, Indonesia, Papua New Guinea, South China Sea, Japan, Philippines, Australia (Western Australia, Queensland), Mariana Islands, New Caledonia, Loyalty Islands, Marshall Islands, Ellice Islands (Rotuma Island), Fiji, Samoan Islands, Kiribati, Line Islands (Palmyra Island), Hawaii, Johnson Atoll; also Galapagos Islands, Mexico, Costa Rica, Panama, Colombia, and Ecuador. Associated with *Pocillopora*, *Seriatopora*, *Stylophora*, rarely *Acropora* and *Porites*.

Harpiliopsis spinigera (Ortmann, 1890)

(Figure 2)

Anchistia spinigera Ortmann 1890, p 511, Plate 36, Figure 23.

Harpilius depressus var. *gracilis* Kemp 1922, p 234, Figure 71.

Harpiliopsis depressus var. *spinigerus*: Holthuis 1952a, p 184–185.

Harpiliopsis spinigera: Chace and Bruce 1993, p 82; De Grave 2000, p 124; Li 2000, p 64, Figure 67; Davie 2002, p 313; Marin et al. 2004, p 201, Figure 2.

Material examined

New Caledonia: Croissant Reef, dive station 533, scuba, 20 m, coll. Bargibant and Menou, 18 April 1994, 1♂ (MNHN-Na 15838).

Remarks

Not previously recorded with certainty from New Caledonia. Associated with *Pocillopora* and *Stylophora*.




Figure 2. *Harpiliopsis spinigera* (Ortmann, 1880), male (MNHN-Na 15838). (a) Second pereopod; (b) same, fingers.

Distribution

Type locality: Samoa. Also known from Kenya, Zanzibar, Comoro Islands, Seychelles, La Réunion, Maldives, Andaman Islands, Vietnam, Philippines, Indonesia, Australia (Queensland), New Caledonia, Fiji, Marshall Islands, and, in the East Pacific region, Panama and Colombia.

Harpilius lutescens Dana, 1852

Harpilius lutescens Dana 1852, p 25; Bruce 2004, p 6.

Periclimenes (Ancylocaris) amamiensis Kubo 1940b, p 44–46, Figures 11, 12.

Periclimenes (Harpilius) lutescens: Holthuis 1952a, p 88–91, Figure 35.

Periclimenes lutescens: Chace and Bruce 1993, p 117; Li 2000, p 209, Figure 271; Davie 2002, p 329.

Material examined

Loyalty Islands: MUSORSTOM 6, stn DW431, 20°22.25'N, 166°10.00'E, 21 m, 18 February 1989, 1♂ (MNHN-Na 14916).

Remarks

This common and widely distributed coral associate, one of the first pontonine shrimps to be described, has not been previously reported from Loyalty Islands. The specimen has a rostral dentition of 1(?) + 6/2, and lacks both second pereopods. The second maxilliped has the characteristic diagnostic distal segments and the third maxilliped has a well-developed arthrobranch. The ambulatory dactyls are also typical of this species. The present depth record may represent the maximum depth from which this species has so far been recorded.

Distribution

Type locality: Tongatapu Island, Tonga. Also known from the Red Sea, Egypt, Israel, Saudi Arabia, Sudan, Ethiopia (Eritrea), Kenya, Zanzibar, Tanzania, Comoro Islands,

Madagascar, Seychelles, Maldives, Singapore, Vietnam, Japan, Indonesia, Australia (Northern Territory, Queensland), Coral Sea, Solomon Islands, Loyalty Islands, Tonga, and Samoan Islands. Also, French Polynesia (Society Islands) and Marquesas Islands.

Ischnopontonia lophos (Barnard, 1962)

Philarius lophos Barnard 1962, p 242, Figure 2.

Ischnopontonia lophos: Bruce 1966a, p 584, Figures 1–5; Bruce 1976a, p 120; Bruce and Coombes 1995, p 109; De Grave 2000, p 125, Figure 2; Li 2000, p 67, Figure 68; Li and Liu 2003, p 156, Figure 2.

Material examined

Madagascar: Tuléar, Grand Récif, coll. R. Hipeau-Jacquotte, 1968, 1♂, 3♀♀ (2 ovig.) (MNHN-Na 15832).

Host

Galaxea sp. [Oculinidae, Scleractinia].

Remarks

First recorded from Madagascar by Bruce (1976a).

Distribution

Type locality: Inhaca Island, Mozambique. Also known from Kenya, Zanzibar, Tanzania, Comoro Islands, Madagascar, Seychelles, Malaya, Singapore, South China Sea, Ryukyu Islands, Australia (Northern Territory, Queensland), Caroline Islands, and Fiji.

Jocaste japonica (Ortmann, 1890)

Coralliocaris superba var. *japonica* Ortmann 1890, p 509.

Jocaste japonica: Holthuis 1952a, p 193–195, Figure 94 (*partim*); Patton 1966, p 279–280, Figure 3b; Bruce 1969a, p 300, Figure 1; Bruce 1979a, p 239; Li 1996, p 225, Figure 4; Li 1997, p 233; Li 1998, p 222; Li 2000, p 69–70, Figure 70; Li 2001, p 79; Li and Liu 2003, p 157, Figure 3; Bruce 2003, p 219; Li 2004a, p 68.

Material examined

Loyalty Islands: ATELIER LIFOU, Lifou, Santal Bay: near Huca Hutighé Islet, stn 1434, 20°52.5'S, 167°08.1'E, 5–20 m, hard bottom, 6 November 2000, 1♂ (MNHN-Na 15458).

Remarks

Not previously recorded from Loyalty Islands. Associated with *Acropora*.

Distribution

Type locality: Kagoshima, Japan. Also known from Kenya, Zanzibar, Tanzania, Mozambique, Comoro Islands, Madagascar, Seychelles, Mauritius, La Réunion,

Maldives, Chagos Islands, Vietnam, South China Sea, Japan, Philippines, Indonesia, Papua New Guinea, Australia (Western Australia, Queensland), Mariana Islands, Caroline Islands, New Caledonia, Loyalty Islands, Marshall Islands, Cook Islands, and Fiji.

Jocaste lucina (Nobili, 1901)

Coralliocaris lamellirostris Stimpson 1860, p 38; De Man 1902, p 842, Plate 26, Figure 55.

Coralliocaris lucina Nobili 1901, p 5; Borradaile 1917, p 383; Kemp 1922, p 276, Figure 102; Barnard 1950, p 799, Figure 151i–m.

Jocaste lucina: Holthuis 1952a, p 193–195, Figure 94 (*partim*); Patton 1966, p 278, Figure 3a; Bruce 1969a, p 301, Figure 2; Bruce 1974b, p 199; Bruce 1979a, p 239; Bruce 1981b, p 89; Franssen 1994, p 110, Figure 64; Li 1997, p 233; Li 1998, p 222, Figures 12, 13; Li 2000, p 70, Figure 71; De Grave 2000, p 126.

Material examined

Loyalty Islands: ATELIER LIFOU, Lifou, Santal Bay, (i) between Huca Hutighé Islet and the shore, stn 1421, 20°52.4'S, 167°08.5'E, 4 m, coarse sand on flatstone, 26–27 November 2000, 1 ovig. ♀ (MNHN-Na 15460); (ii) in front of Ngoni, stn 1457, 20°46.8'S, 167°02.75'E, 5–10 m, dark overhang and boulders, 27 November 2000, 1 ovig. ♀ (MNHN-Na 15459).

Remarks

Not previously recorded from Loyalty Islands. Associated with *Acropora*, rarely *Pocillopora*, *Stylophora*.

Distribution

Type locality: Eritrea. Also known from Egypt, Israel, Sudan, Yemen, Oman, Kenya, Zanzibar, Tanzania, Mozambique, Comoro Islands, Madagascar, Seychelles, La Réunion, Maldives, Sri Lanka, Andaman Islands, Nicobar Islands, Singapore, Vietnam, South China Sea, Indonesia, Papua New Guinea, Australia (Western Australia, Queensland), Coral Sea, Marianas Islands, New Caledonia, Loyalty Islands, Marshall Islands, Fiji, Cook Islands and Johnson Atoll.

Kemponia agag (Kemp, 1922)

(Figure 3)

Periclimenes (Ancylocaris) agag Kemp 1922, p 197, Figures 47–50, Plate VII, Figure 9.

Periclimenes agag: Li 2000, p 150, Figure 181.

Kemponia agag: Bruce 2004, p 10.

Material examined

New Caledonia: (i) Senez Reef, 7 m, coll. P. Bouchet, 7 September 1992, 1 ♂ (MNHN-Na 14877); (ii) Ouano. scuba, 0.1 m, April 1995, 2 ovig. ♀♀ (MNHN-Na 15810).

Remarks

Previously reported from New Caledonia (Nouméa) by Ledoyer (1984).



Figure 3. *Kemponia agag* (Kemp, 1922), male (MNHN-Na 14877). (a) Carapace and anterior appendages; (b) antennal scale; (c) second pereiopod, ischium, merus, and carpus; (d) same, chela; (e) same, fingers; (f) same, article of carpus and chela; (g) proximal fingers; (h) third pereiopod, propod and dactyl; (i) same, distal propod and dactyl.

Distribution

Type locality: Port Blair, Andaman Islands. Also known from Red Sea, Australia (Queensland), New Caledonia and Marshall Islands.

Kemponia anymone (De Man, 1902)

Periclimenes anymone De Man 1902, p 829–833, Plate 25, Figure 53; Bruce 1991b, p 235; Chace and Bruce 1993, p 102; Li 2000, p 155, Figure 190; Davie 2002, p 323; Li and Liu 2004, p 90, Figure 1g–k.

Kemponia anymone: Bruce 2004, p 11.

Material examined

New Caledonia: (i) lagoon, Îlot Maitre, reef flat, scuba, 3 m, coll. C. Vadon, 29 August 1978, 1 ovig. ♀ (MNHN-Na 15834).

Remarks

The specimen has a rostral dentition of 1+7/3 and has both second pereiopods. The species was previously recorded from New Caledonia by Bruce (1967b, 1991b) and Ledoyer (1984).

Distribution

Type locality: Ternate, Indonesia. Also known from the Nicobar Islands, Singapore, Vietnam, Philippines, Indonesia, Papua New Guinea, Australia (Western Australia, Northern Territory, Queensland), Solomon Islands (New Georgia Islands) New Caledonia, and Samoan Islands.

Kemponia anacanthus (Bruce, 1988)

Periclimenes anacanthus Bruce 1988a, p 105, Figures 1–5; Bruce and Coombes 1995, p 125, Figure 12a; Li 2000, p 156, Figure 191.

Kemponia anacanthus: Bruce 2004, p 12.

Material examined

New Caledonia: (i) stn LF3 “83”, 20°47.27'S, 167°07.34'E, 12 m, on red algae, 27 November 1995, 1♂, 2♀♀ (MNHN-Na 15905); (ii) lagoon, 20°42.15'S, 167°09.90'E, scuba, 1 m, on red algae, 28 November 1995, 1♂ (juvenile?, both second pereiopods lost) (MNHN-Na 15909). **Loyalty Islands:** ATELIER LIFOU, Lifou, Santal Bay, (iii) near Kiki, stn 1411, 20°47.6'S, 167°10.35'E, 4–8 m, sand between coral heads, 13 November 2000, 1♂, 2 juveniles (MNHN-Na 15462); (iv) in front of Peng, stn 1423, 20°54.0'S, 167°07.3'E, 12 m, sandy channels between coral heads, 14 November 2000, 1♀ (MNHN-Na 15461); (v) between Cap Wekutr and Cap Wajez, stn 1455, 20°56.8'S, 167°02.7'E, slope, 15–20 m, 25 November 2000, 1 ovig. ♀ (MNHN-Na 15463); (vi) between Huca

Hutighé Islet and shore, stn 1421, 20°52.4'S, 167°08.5'E, 4 m, coarse sand on flat-stone, 27 November 2000, 1♂ (MNHN-Na 15464).

Remarks

Only the right second pereopod of specimen (iv) and one of the juveniles of (iii) present, all other specimens lack their second pereopods. The unarmed merus and carpus and the long carpus (much longer than palm) of second pereopod of the specimens indicate they are identical to the present species. Not previously recorded from Loyalty Islands.

Distribution

Type locality: Queensland. Also known from Cobourg Peninsula (Australia), New Caledonia, and Loyalty Islands.

Kemponia andamanensis (Kemp, 1922)

Periclimenes (Ancylocaris) andamanensis Kemp 1922, p 204, Figures 54–57.

Periclimenes (Harpilius) andamanensis: Holthuis 1952a, p 79.

Periclimenes andamanensis: Bruce 1977c, p 269; Bruce 2003, p 229; Li 1996, p 229, Figure 7; Li 1997, p 238; Li 2000, p 156, Figure 192; Li 2001, p 81; Li 2004a, p 68; Li and Liu 2004, p 91, Figure 2.

Kemponia andamanensis: Bruce 2004, p 12.

Kemponia cf. *andamanensis*: Li et al. 2004, p 529, Figure 16.

Material examined

Madagascar: (i) Nosy Be, shallow water, coll. A. Crosnier, 1966, 2♂♂, 3 ovig. ♀♀ (MNHN-Na 15969); (ii) same, 1♂, 1♀ (MNHN-Na 15967). **New Caledonia:** (iii) Station Saint Vincent (SASV2), Canal d'Amenée, coll. IFREMER, 5 August 1996, 4♂♂, 9 ovig. ♀♀ (MNHN-Na 15824); (iv) western coast, Ouano, trawl, 1 m, seagrass, 1–5 m, 22 December 1998, 9♂♂, 16♀♀ (4 ovig.) (MNHN-Na 15828). **Fiji:** (v) Viti Levu, West Lagoon, SUVA 2, stn CP66, 37 m, 17°45.1'S, 177°13.7'E, 21 October 1998, 1♂, 1 ovig. ♀ (MNHN-Na 15465).

Remarks

Not previously recorded from Fiji and New Caledonia. The present record extends its bathymetric range to 37 m depth.

Distribution

Type locality: Andaman Islands. Also known from Madagascar, Indonesia, China, South China Sea, Japan, Australia (Queensland), New Caledonia, and Fiji.

Kemponia darwiniensis (Bruce, 1987)

Periclimenes darwiniensis Bruce 1987, p 29, Figures 1–5; Li 2000, p 174, Figure 217.

Kemponia darwiniensis: Bruce 2004, p 13.

Material examined

New Caledonia: St Marie Bay, 14 m, scuba, coll. P. Bouchet, 4 January 1993, 1 ovig. ♀ (MNHN-Na 15911).

Remarks

This is the first record outside its type locality Darwin Harbour, Australia.

Distribution

Type locality: Darwin Harbour, Darwin, Northern Territory, Australia. Also known now from New Caledonia.

Kemponia elegans (Paulson, 1875)

Anchistia elegans Paulson 1875, p 113, Plate 17, Figure 1.

Periclimenes (Falciger) elegans: Borradaile 1917, p 371.

Periclimenes (Ancylocaris) elegans: Kemp 1922, p 215, Figures 60–62.

Periclimenes (Harpilius) elegans: Holthuis 1952a, p 81, Figure 31.

Periclimenes elegans: Bruce 1971a, p 7; Bruce 1983b, p 884; Fransen 1994, p 122; Bruce and Coombes 1995, p 129; Li 1997, p 238; Li 1998, p 222, Figures 14–17; Li 2000, p 178, Figure 225; De Grave 2000, p 135.

Kemponia elegans: Bruce 2004, p 14.

Material examined

New Caledonia: (i) Kuendu Bay, 0.5 m, coll. B. Richer de Forges, 5 May 1996, 1♂, 1 ovig. ♀ (MNHN-Na 15826); (ii) lagoon, Poingam, low tide, coll. B. Richer de Forges, 10 May 1997, 1♂ (MNHN-Na 15829). **Fiji:** (iii) MUSORSTOM 10, south of Vitu Levu, flat reef, low tide, 3 August 1998, 1♂ (MNHN-Na 15467).

Remarks

Not previously recorded from Fiji and New Caledonia.

Distribution

Type locality: Red Sea. Also known from Egypt, Saudi Arabia, Yemen, Kenya, Zanzibar, Tanzania, Madagascar, Seychelles, Kuwait, Laccadive Islands (Minikoi), India, Sri Lanka, Andaman Islands, Nicobar Islands, Singapore, South China Sea, Ryukyu Islands, Japan, Philippines, Indonesia, Papua New Guinea, Timor Sea (Hibernia Reef), Australia (Western Australia, Northern Territory, Queensland), Caroline Islands, Solomon Islands, New Caledonia, Marshall Islands, Fiji, French Polynesia (Society Islands).

Kemponia ensifrons (Dana, 1852)

(Figures 4, 5)

Anchistia ensifrons Dana 1852, p 25.

Periclimeses (Falciger) ensifrons: Borradaile 1917, p 367, 370.

Periclimeses ensifrons: Chace and Bruce 1993, p 111; Li 2000, p 180.

Kemponia ensifrons: Bruce 2004, p 15.

Material examined

New Caledonia: MONTROUZIER, Touho, sandy islet, scuba, coll. B. Richer de Forges, scuba, 7 September 1993, 1 ovig. ♀ (MNHN-Na 14891).

Remarks

Not previously reported from New Caledonia. The species is possibly synonymous with *Kemponia grandis* (Stimpson) (Bruce 2004, p 15).

Distribution

Type locality: Balabac Strait, North Borneo. Also reported from the Egyptian Red Sea, Zanzibar, Comoro Islands, Seychelles, Burma, China, Papua New Guinea, New Caledonia, Marshall Islands, and French Polynesia (Tuamotu Islands).

Kemponia grandis (Stimpson, 1860)

Anchistia grandis Stimpson 1860, p 39.

Periclimeses grandis: Borradaile 1898, p 382; Chace and Bruce 1993, p 112; Li 2000, p 186, Figure 235; Davie 2002, p 327; Li and Liu 2004, p 93, Figure 4; Li 2004a, p 69.

Kemponia grandis: Bruce 2004, p 16; Li et al. 2004, p 530.

Material examined

Madagascar: (i) Nosy Be, shallow waters, coll. A. Crosnier, 1966, 2♂♂, 2 ovig. ♀♀ (MNHN-Na 15956). **New Caledonia:** (ii) MONTROUZIER, Touho, Grand Récif, Mangalia, intertidal zone, coll. B. Richer de Forges, 15 October 1993, 1 ovig. ♀ (MNHN-Na 14893); (iii) Ouano reef flat, low tide, coll. B. Richer de Forges, 26 April 1998, 1 ovig. ♀ (MNHN-Na 15813). **Loyalty Islands:** ATELIER LIFOU, Lifou, (iv) Santal Bay: between Huca Hutighé Islet and the shore, stn 1421, 20°52.4'S, 167°08.5'E, 4 m, coarse sand on flatstone, 26–27 November 2000, 1♂ (MNHN-Na 15471).

Remarks

Previously reported from Madagascar by Bruce (1978a). Not previously recorded from New Caledonia and Loyalty Islands, although it is one of the commonest and most widely distributed Indo-West Pacific pontoniine species.

Distribution

Type locality: Amami O Shima, Ryukyu Islands. Also known from Egypt, Israel, Djibouti, Aden, Kenya, Zanzibar, Tanzania, Mozambique, Comoro Islands, Madagascar, Seychelles,

Sri Lanka, Burma, Malaya, Singapore, Japan, Indonesia, Australia (Northern Territory, Queensland), New Caledonia, Loyalty Islands, Marshall Islands, Tuvalu, and Fiji.

Kemponia kororensis (Bruce, 1977)

Periclimenes kororensis Bruce 1977d, p 33, Figures 1–4; Bruce and Svoboda 1984, p 94, Figures 5, 6; Li 2000, p 201, Figure 258.

Kemponia kororensis: Bruce 2004, p 16.

Material examined

Philippines: MUSORSTOM II, shore station, stn 11, opposite Mactan Marine Station, scuba, 5–18 m, with fungiid coral, 9 December 1980, 1♂ (MNHN-Na 15961).

Remarks

The single specimen with rostral formula of 2+5/4, second to fourth dorsal teeth are grouped over orbital region, but not forming a crest as in the type. Both second pereopods




Figure 4. *Kemponia ensifrons* (Dana, 1852), ovigerous female (MNHN-Na 14891). (a) Carapace and anterior appendages; (b) anterior carapace; (c) antennule; (d) antenna.




Figure 5. *Kemponia ensifrons* (Dana, 1852), ovigerous female (MNHN-Na 14891). (a) Second pereopod; (b) same, chela and carpus; (c) third pereopod, propod and dactyl; (d) same, distal propod and dactyl.

lost. Ambulatory pereopods with flexor margin of propod unarmed except for the distoventral spine. Previously recorded from Philippines by Bruce and Svoboda (1984) from Cebu.

Distribution

Type locality: Koror, Palau Islands, Caroline Islands. Also known from Malaysia, Japan, Philippines, Indonesia, Papua New Guinea, Australia (Queensland), and Marshall Islands.

Kemponia lacertae (Bruce, 1992)

Periclimenes lacertae Bruce 1992, p 46, Figures 1–6; Li 2000, p 203, Figure 260.

Kemponia lacertae: Bruce 2004, p 17.

Material examined

Loyalty Islands: ATELIER LIFOU, Lifou, Chateaubriand Bay: Wé Beach, stn 1474, 20°54.8'S, 167°16.1'E, 0–3 m, fine sand and heads of *Porites*, 10 and 21 November 2000, 1♂ (MNHN-Na 15473).

Remarks

The single specimen with the rostrum inclined slightly upwards, instead of horizontal. Not previously recorded from Loyalty Islands.

Distribution

Type locality: Chinaman's Ridge, Mrs Watson's Bay, Lizard Island, Queensland, Australia, and Liang Island; Hansa Bay, Papua New Guinea. Now also known from Loyalty Islands.

Kemponia nilandensis (Borradaile, 1915)

Periclimenes (Falciger) nilandensis Borradaile 1915, p 211; Borradaile 1917, p 324, 372, Plate 54, Figure 13.

Periclimenes nilandensis: Bruce 1978a, p 222, Figures 8, 9; Chace and Bruce 1993, p 118; Bruce 1996, p 237; Li 2000, p 214, Figure 278; Davie 2002, p 330.

Kemponia nilandensis: Bruce 2004, p 18; Li et al. 2004, p 531.

Material examined

Madagascar: (i) Nosy Be, shallow waters, coll. A. Crosnier, 1966, 2♂♂, 3 ovig. ♀♀ (MNHN-Na 15955). **La Réunion:** (ii) MD32, stn CP43, 21°20.7'S, 55°26.9'E, 73–77 m, shelly basaltic sand, 27 August 1982, 3♂♂, 27 ♀♀ (22 ovig.) (MNHN-Na 14918). **New Caledonia:** (iii) MONTROUZIER, Touho Channel, 35 m, from antipatharians, coll. B. Richer de Forges, 31 August 1993, 4♂♂, 10♀♀ (9 ovig.) (MNHN-Na 14917).

Remarks

Previously recorded from Madagascar (Nosy Bé) by Bruce (1978a). Not previously recorded from La Réunion. Reported from Anaa Reef, New Caledonia by Bruce (1996).

Distribution

Type locality: South Nilandu Atoll, Maldives. Also known from Kenya, Zanzibar, Madagascar, Maldives, South China Sea, Indonesia, Australia (Northern Territory, Queensland), and New Caledonia; to depths of 133 m.

Kemponia seychellensis (Borradaile, 1915)

Periclímenes (Falciger) seychellensis Borradaile 1915, p 212.

Periclímenes (Ancylocaris) seychellensis: Kemp 1922, p 176, Figures 34, 35, Plate 6, Figure 7.

Periclímenes (Harpilius) seychellensis: Holthuis 1952a, p 66, Figure 25; Ledoyer 1984, p 33, Figure 14.

Periclímenes seychellensis: Bruce 1971a, p 8; Bruce 1991b, p 238, Figure 8; Li 2000, p 235, Figure 313.

Kemponia seychellensis: Bruce 2004, p 19.

Material examined

New Caledonia: (i) lagoon, Îlot Maitre, 22°19.70'S, 166°23.30'E, 10 m, on seagrass, scuba, coll. I. Takeuchi, 6 November 1995, 3 ovig. ♀♀ (MNHN-Na 15904); (ii) Rocher à la Voile, 22°18.23'S, 166°26.15'E, 0.5 m, scuba, 18 November 1995, 1 ♂ (MNHN-Na 15906); (iii) lagoon, Ricaudy reef flat, low tide, coll. B. Richer de Forges, 22 June 1997, 19 ♂♂, 14 ovig. ♀♀ (MNHN-Na 15811); (iv) Ouano reef flat, low tide, coll. B. Richer de Forges, 26 April 1998, 4 ovig. ♀♀ (MNHN-Na 15830).

Remarks

The tubercle on the dorsal surface of eyestalk makes the identification of the species simple. The species has been previously recorded from New Caledonia by Ledoyer (1984) and Bruce (1991b).

Distribution

Type locality: Praslin, Seychelles. Also reported from Kenya, Zanzibar, Tanzania, Mozambique, Madagascar, Seychelles, Pakistan, India, Andaman Islands, Singapore, China, Japan, Indonesia, Papua New Guinea, Australia (Western Australia, Northern Territory, Great Barrier Reef), and New Caledonia.

Kemponia cf. suvativensis (Borradaile, 1915)

(Figure 6)

Periclimenes (Falciger) suvativensis Borradaile 1915, p 212; Borradaile 1917, p 375, Plate 55, Figure 16.

Periclimenes suvativensis: Li 2000, p 239, Figure 317; Davie 2002, p 332.

Kemponia suvativensis: Bruce 2004, p 19.

Material examined

New Caledonia: west coast of Le Cap, in mangroves, coll. B. Richer de Forges, 2 December 1992, 1♂, 2 ovig. ♀♀ (MNHN-Na 14926).

Remarks

The specimens were collected with *Palaemon* sp. All the specimens lack second pereiopods, so the identification must be considered as provisional. Not previously recorded from New




Figure 6. *Kemponia cf. suvativensis* (Borradaile, 1915), ovigerous female (MNHN-Na 14926). (a) Carapace; (b) antennal scale; (c) third pereiopod, carpus, propod, and dactyl; (d) same, distal propod and dactyl.

Caledonia. The specimens have rostral dentition of $1+7/3$, the rostrum greatly exceeds the antennular peduncle, particularly in the male, is slender, about 1.35 times the carapace length, as opposed to 1.15 times in the *K. suwadiwensis* type material, the scaphocerite lamella is subquadrate distally and well exceeded by the distolateral tooth. The third pereopod dactyl is about 9.0 times longer than the basal width and about 0.36 of the propod length, as opposed to 0.2 times in the type material; the proportion of length to breadth of the propod are similar but the present material has fewer ventral spines, four as opposed to six.

Distribution

Type locality: Suwadiva Atoll, Maldives. Also reported from Australia (Northern Territory) and New Caledonia.

Kemponia tenuipes (Borradaile, 1898)

Periclimenes tenuipes Borradaile 1898, p 384; Bruce 1978b, p 261, Figures 6A, 7; Bruce 1991b, p 240; Li 2000, p 240, Figure 319.

Periclimenes (Ancylocaris) tenuipes: Kemp 1922, p 220, Plate 8, Figure 11.

Periclimenes (Harpilius) tenuipes: Holthuis 1952a, p 84.

Kemponia tenuipes: Bruce 2004, p 19.

Material examined

New Caledonia: (i) Marie Bay, 14 m, scuba, coll. P. Bouchet, 4 January 1993, 1♂, 2♀♀ (1 ovig.) (MNHN-Na 15918). **Loyalty Islands:** ATELIER LIFOU, Lifou, Santal Bay, (ii) Cap Lefèvre, stn 1469, 20°54.2'S, 167°00.4'E, 70–130 m, dredge, 22–23 November 2000, 1 ovig. ♀ (MNHN-Na 15591); (iii) Peng, stn 1464, 20°54.5'S, 167°05.9'E, 30–50 m, 14 November 2000, 1♂, 1♀ (MNHN-Na 15589); (iv) north of Cape Aimé Martin (=Acadro), stn 1450, 20°45.8'S, 167°01.65'E, 27–31 m, washings, 17 and 21 November 2000, 1♀ (MNHN-Na 15624). **Fiji:** (v) SUVA 2. Viti Levu, West Lagoon, Stn CP 66, 17°45.1'S, 177°13.7'E, 37 m, coll. Richer-ORSTOM, 21 October 1998, 1 ovig. ♀ (MNHN-Na 15614).

Remarks

Not previously recorded from Loyalty Islands and Fiji. Previously recorded from New Caledonia by Bruce (1991b).

Distribution

Type locality: New Britain, Papua New Guinea. Also reported from Jordan, Kenya, Zanzibar, Madagascar, Seychelles, La Réunion, Maldives, Sri Lanka, Andaman Islands,

China, Japan, Philippines, Indonesia, Papua New Guinea, Timor Sea (Hibernia Reef), Australia (Northern Territory, Queensland), Caroline Islands, New Caledonia, Marshall Islands, and Fiji.

Manipontonia psamathe (De Man, 1902)

Urocaris psamathe De Man 1902, p 816–822, Plate 25, Figure 51.

Periclimenes (Ancylocaris) psamathe: Kemp 1922, p 173.

Periclimenes (Harpilius) psamathe: Holthuis 1952a, p 61, Figure 23.

Periclimenes psamathe: Bruce 1991b, p 238, Figures 1a, 3a; Chace and Bruce 1993, p 120; Bruce 1996, p 238; Li 2000, p 229, Figure 304; Davie 2002, p 331; Li et al. 2004, p 545, Figure 31.

Manipontonia psamathe: Bruce et al. 2005, p 6, Figures 1–3.

Material examined

Madagascar: (i) SE coast, Ste Luce, trawl, shelly sand, 50 m, coll. A. Crosnier, October 1958, 1♂, 2 ovig. ♀♀ (MNHN-Na 12904); (ii) Nosy Be, shallow waters, coll. A. Crosnier, 1966, 1♂, 2 ovig. ♀♀, 3 spms (MNHN-Na 15952). **New Caledonia:** (iii) Iré Bay, scuba, night dive, 10 m, coll. P. Laboute, 7 June 1990, 6♂♂, 7♀♀ (6 ovig.) (MNHN-Na 12905). **MONTROUZIER,** (iv) Touho Channel, 48 m, scuba, coll. B. Richer de Forges, 31 August 1993, 1 ovig. ♀ (MNHN-Na 14922); (v) same, 35 m, from antipatharians, 31 August 1993, 2 ovig. ♀♀, 1 juvenile (MNHN-Na 14921); (vi) same, 1 ovig. ♀ (MNHN-Na 14923); (vi) same, 35 m, 1♂ (MNHN-Na 15902).

Hosts

(i) Hydroids and sponges; (iv) (v) antipatharians.

Remarks

First recorded from New Caledonia at Nouméa by Bruce (1970a). Previously recorded from Madagascar by Bruce (1977c, 1978a). The association of (i) with sponges is probably accidental.

Distribution

Type locality: Ternate, Indonesia. Also known from Kenya, Zanzibar, Tanzania, Madagascar, Seychelles, Maldives, Chagos Islands, Thailand, Malaya, Singapore, China, Japan, Philippines, Indonesia, Papua New Guinea, Australia (Western Australia, Northern Territory, Queensland), Caroline Islands, New Caledonia, and Marshall Islands.

Mesopontonia brevicarpus sp. nov.
(Figures 7, 8)

Material examined

La Réunion: MD 32, stn DS173, 20°51.5'S, 55°36.87'E, 270 m, 8 September 1982, 1 ovig. ♀ (cl 2.56 mm), holotype; 1 ♂ (cl 2.53 mm), 1 ovig. ♀ (cl 2.77 mm), paratypes; 3 juveniles (MNHN-Na 14824).

Description

A small-sized pontoniine shrimp of slender, subcylindrical body shape, closely resembling other species of the genus, such as *M. gracilicarpus* Bruce, 1990 and *M. verrucimanus* Bruce, 1996.

Rostrum well developed, horizontal, compressed, extending anteriorly near to the end (ovig. ♀) or to the proximal third (♂) of distal segment of antennular peduncle, about 0.9 of carapace length, dorsal carina well developed, posteriorly elevated, dorsal margin convex, with eight evenly distributed acute teeth, first tooth with (ovig. ♀) or without (♂) basal suture, situated behind posterior orbital margin, distal tooth minute, lateral carinae




Figure 7. *Mesopontonia brevicarpus*, new species. (a, d-g) Holotype ovigerous female; (b) paratype male; (c) juvenile (MNHN-Na 14824). (a) Carapace and anterior appendages; (b, c) carapace; (d) antennule; (e) first pereopod; (f) same, chela; (g) same, apexes of fingers.




Figure 8. *Mesopontonia brevicarpus*, new species, holotype ovigerous female (MNHN-Na 14824). (a) Major second pereiopod; (b) same, chela; (c) same, fingers; (d) minor second pereiopod; (e) same, chela; (f) third pereiopod; (g) same, propod and dactyl; (h) same, distal propod and dactyl.

obsolete, ventral carina feebly developed, margin slightly convex. Carapace smooth, glabrous, with epigastric spine at anterior 0.3 of carapace length, basal suture distinct, supraorbital and antennal spines absent, inferior orbital angle produced, hepatic spine well developed, slender, acute, reaching or extending to anterior margin of carapace, at about 0.12 of carapace length, well below level of inferior orbital angle, anterolateral angle of branchiostegite bluntly rounded. Abdomen and caudal fan without special features (the ovig. ♀ holotype lacks the caudal fan), as in *M. gracilicarpus*.

Antennae as in *M. gracilicarpus*.

Eye with well-developed large, globular, pigmented cornea, diameter about 0.23 of carapace length.

Mouthparts mainly undissected. Third maxilliped without exopod, ischiomerus and basis feebly separated, with a rudimentary arthrobranch.

First pereiopod with finger about 0.7 of palm length, tips hooked, cutting edge entire except convex subapically with fine pectinated serrates; carpus about 1.1 times chela length, 0.9 of merus length, coxa with small ventromedial process.

Second pereiopods well developed, markedly unequal, dissimilar. Major pereiopod with chela about 1.0–1.1 of carapace length, palm subcylindrical, 3.9 times longer than deep, covered with uniformly distributed small blunt tubercles, many with short simple setae distally, fingers about 0.5 of palm length, dactyl slender, about 4.0 times longer than proximal depth, tapering distally, with acute hooked tip, cutting edge with distal half sharp, with one low acute tooth at proximal 0.3 and a notch at proximal 0.38 of the cutting length, without dorsolateral flange; fixed finger with two acute teeth at proximal 0.25 and 0.38 on the cutting edge; carpus short, stout, unarmed, about 0.34 of palm length, sparsely tuberculate, about 2.0 times longer than distal width, merus about 0.85 of palm length, 7.0 times longer than wide, uniform, smooth, unarmed; ischium about 0.9 times merus length, 0.73 of palm length, 7.0 times longer than distal width, smooth, unarmed; basis and coxa without special features. Minor pereiopod with chela about 0.7 of carapace length, 0.95 of

major chela palm length, palm subcylindrical, smooth, about 4.0 times longer than distal width, slightly tapering proximally, fingers subequal to palm, slender, with small hooked tips, cutting edges distally sharp, entire; carpus slender, about 0.75 of chela length, 8.0 times longer than distal width, smooth, unarmed; merus about 1.1 times carpus length, 11.0 times longer than distal width, subuniform, unarmed; basis and coxa without special features.

Ambulatory pereopods slender. Third pereopod with propod about 0.7 of carapace length, 17 times longer than wide, uniform, with a pair of long and slender distoventral spines, ventral border with similar spinules, size decreasing proximally, with long setae distodorsally; dactyl about 0.17 of propod length, slender, biunguiculate, with clearly demarcated unguis, about 3.3 times longer than proximal width, 0.70 of corpus length, corpus 2.3 times longer than proximal depth, compressed.

Measurements (mm)

Carapace length, 2.53–2.77; carapace and rostrum, 4.75–4.81; second pereopod, major chela, 2.75; minor chela, 1.74.

Systematic position

The present species is closely related to *M. gracilicarpus* Bruce, 1990; it can be distinguished from the latter by the tuberculate major second pereopod chela, relatively shorter carpus of minor second pereopod (0.75 of chela length), and the hepatic spine on carapace reaching the anterior margin of carapace. In *M. gracilicarpus*, the major second pereopod chela smooth, the carpus of minor second pereopod 1.5 times chela length, and the hepatic spine not reaching to the anterior margin of carapace. *Mesopontonia brevicarpus* shares the tuberculate major chela of second pereopod with *M. verrucimanus* Bruce, 1996. It can be immediately distinguished from that species by the biunguiculate dactyls of ambulatory pereopods.

Etymology

From *brevis* (Latin), short; *carpus* (Latin), wrist, with reference to the short carpus of major second pereopod.

Remarks

With the addition of the new species, there are now seven species in the genus *Mesopontonia*, including the unnamed taxon, *Mesopontonia* sp. Bruce, 1996. The key to *Mesopontonia* species provided by Bruce (1996) can be modified as following:

- 3. Second pereopods markedly unequal 3'
- Second pereopods at most moderately unequal; carpus of minor second pereopod shorter than chela 4
- 3'. Palm of major second pereopod tuberculate; carpus of minor second pereopod shorter than chela; R. 1+8/2 *M. brevicarpus* sp. nov.
- Palm of major second pereopod smooth; carpus of minor second pereopod about 1.5 times length of chela; R. 1+8/2 *M. gracilicarpus* Bruce, 1990

Mesopontonia gracilicarpus Bruce, 1990

Mesopontonia gracilicarpus Bruce 1990a, p 202–211, Figures 34–37, 39l, m; Bruce 1991a, p 391, Figure 64; Li 2000, p 76, Figure 76.

Material examined

New Caledonia: (i) SMIB 8, Azteque Bank, stn DW181, 23°17.7'S, 168°04.8'E, 311–330 m, 31 January 1993, 1 ovig. ♀ (MNHN-Na 14825); (ii) BATHUS 4, stn CP922, 18°48.04'S, 163°18.58'E, 600 m, 6 August 1994, 1♀ (MNHN-Na 15966); (iii) HALIPRO 2, Norfolk Ridge, stn BT70, 24°46'S, 168°09'E, 226–238 m, 20 November 1996, 1 ovig. ♀ (MNHN-Na 15974); (iv) NORFOLK 1, Antigonina Bank, stn CP1714, 23°22'S, 168°03'E, 257–269 m, 26 June 2001, 1♂ (MNHN-Na 15971).

Remarks

The single specimen of (i) is in poor condition, with the major second pereopod only, damaged, lacking the fingers, sixth abdominal segment and caudal fan and left fifth and right third and fourth pereopods missing. The specimen (ii) with rostral formula 1+11/3, palm of major second chela covered with distinct fine tubercles. Specimen (iii) damaged, both second pereopods lost, rostrum broken. The specimen (iv) with rostral formula 1+9/2, right (?minor) second pereopod lost.

Distribution

Type locality: New Caledonia, 22°56'S, 167°14'E.

Metapontonia fungiacola Bruce, 1967

Metapontonia fungiacola Bruce 1967a, p 24–32, Figures 10–12; Li 2000, p 78, Figure 79.

Material examined

New Caledonia: MONTROUZIER, Touho, scuba, sand islet, coll. B. Richer de Forges, 7 September 1993, 1♀ (MNHN-Na 14826).

Remarks

Not previously reported from New Caledonia and, outside the Indian Ocean, known only from the Ryukyu Islands. The single specimen, which has both second pereopods, agrees well with previous descriptions. The host was not recorded, but the species is usually associated with fungiid corals.

Distribution

Type locality: Pamanzi Reef, Mayotte, Comoro Islands. Also known from Kenya, Tanzania, Seychelles, La Réunion and Ryukyu Islands.

***Palaemonella crosnieri* Bruce, 1978**

Palaemonella crosnieri Bruce 1978a, p 210, Figures 2–4; Li 2000, p 101, Figure 108.

Material examined

Loyalty Islands: ATELIER LIFOU, Lifou, Santal Bay, (i) east of Santal Bay: Mepinyo, stn 1446, 20°50.8'S, 167°09.7'E, 36–40 m, bottom of slope, 16 November 2000, 1♂ (MNHN-Na 15475); (ii) west of Easo Point, stn 1451, 20°47.3'S, 167°06.8'E, 10–21 m, coral heads, 19 November 2000, 1♂ (MNHN-Na 15476); (iii) between Cap Mandé and Cap Lefèvre (=Nem), stn 1452, 20°54.6'S, 167°02.1'E, 2–25 m, scuba, 20 and 22 November 2000, 2♂♂ (MNHN-Na 15477); (iv) between Cap Wekutr and Cap Wajez, stn 1410, 20°56.7'S, 167°03.1'E, 2–4 m, reef flat edge, 25 November 2000, 1♂ (MNHN-Na 15474).

Remarks

The specimen (i) lacks the right pereopod 1 and 2, with a regenerative left second pereopod, and the rostrum broken and the part anterior to orbital margin lost. Previously only known from the type locality, Îles Glorieuses, 11°28.1'S, 47°21.1'E (27°21.1'E in the original reference must be a mistake). The present records extend the bathymetric range to 40 m depth.

Distribution

Type locality: Îles Glorieuses (Inchim Ouan, 11°34'S, 47°19'E), from 20 m. Also known from Kenya, Western Australia, and Loyalty Islands.

***Palaemonella dolichodactylus* Bruce, 1991**

Palaemonella dolichodactylus Bruce 1991b, p 232, Figures 6f–l, 7; Bruce 1991a, p 301; Li 2000, p 101, Figure 110.

Material examined

Fiji: SUVA 2, Viti Levu, (i) Northeast Lagoon, stn BS 20, 18°10.2'S, 178°30.9'E, 14 m, 15 October 1998, 1 ovig. ♀ (MNHN-Na 15481); (ii) South Lagoon, stn BS 35, 18°08.1'S, 178°25.3'E, 16 m, coll. B. Richer de Forges, 18 October 1998, 1 ovig. ♀ (MNHN-Na 15480); (iii) West Lagoon, stn CP 66, 17°45.1'S, 177°13.7'E, 37 m, coll. B. Richer de Forges, 21 October 1998, 1 ovig. ♀ (MNHN-Na 15479).

Remarks

The specimens with the dactyl of ambulatory pereopods about 12 times longer than proximal deep, propod obscurely segmented, but all of them with the posteroventral angle of fifth abdominal somite armed with acute tooth. Not previously recorded from Fiji. See also remarks on *Palaemonella komaii* sp. nov.

Distribution

Type locality: Lagoon Est, New Caledonia, 65–70 m. Also known from the Norfolk Ridge, 250 m (Bruce 1991b) and Fiji.

Palaemonella hachijo Okuno, 1999

Palaemonella hachijo Okuno 1999, p 739, Figures 1–3; Okuno 2000, p 1297, Figure 1.

Material examined

New Caledonia: west coast, mangrove, coll. B. Richer de Forges, 2 December 1992, 1 spm (MNHN-Na 15628).

Remarks

See also remarks on *Palaemonella komaii* sp. nov. The specimen is not in good condition, dried, the right second and fifth pereopods, left fourth and fifth pereopods lost, left second and third pereopods detached. But, the long and large antennal spine which is larger than hepatic spine and distinctly exceeds the antennal basicerite, relatively shorter ambulatory dactyl which is about eight times proximal depth and with only one row of dorsal setae, the propod is not segmented and the distoventral spine is relatively longer which is about one-third of dactyl, show that the specimen is closer to *P. hachijo* than to *P. dolichodactylus*. The rostral formula is 2+6/3, similar to the specimens of Okuno (2000), the posteroventral angle of fifth abdominal somite armed with acute tooth. Previously known only from its type locality Hachijo-jima Island, and Okinawa, Japan.

Distribution

Type locality: Hachijo-jima Island, Japan, 15–45 m. Now also known from New Caledonia, mangrove.

Palaemonella komaii sp. nov.

(Figures 9–11)

Material examined

Tonga: BORDAU 2, Vava'u group, stn DW 1583, 327–360 m, 18°37'S, 174°03'W, 13 June 2000, 1♂ holotype (MNHN-Na 15482). **Fiji:** stn CP 1445, 17°10'S, 178°42'W, 350–365 m, coll. Bouchet, Waren and Richer de Forges, 3 March 1999, 1♂ paratype (MNHN-Na 15600).

Description

Holotype. A small pontoniine shrimp of subcylindrical body form. Most of pereopods lost, only detached right first and both fifth pereopods present.

Carapace smooth, glabrous. Rostrum well developed, shallow, compressed, straight and horizontal, extending beyond anterodorsal margin of second segment of antennular peduncle, about 0.8 of carapace length; dorsal margin with seven (excluding epigastric spine) acute teeth distributed along whole length, interdental spaces smaller anteriorly, first tooth situated slightly posterior to posterior orbital margin; lateral carinae indistinct; ventral margin with two small acute teeth on distal third, smaller than dorsal teeth; interdental spaces and proximal ventral carina feebly setose. Supraorbital spine absent; epigastric spine similar to dorsal rostral teeth, situated at anterior 0.3 of the carapace length; orbit feebly developed, inferior orbital angle feebly produced, bluntly round in lateral view, postorbital carina distinct; antennal spine large slender marginal, distinct below the inferior orbital angle, far exceeding inferior orbital angle, reaching to distal margin of antennal basicerite; hepatic spine distinctly smaller than antennal spine, slender, situated far below level of




Figure 9. *Palaemonella komaii*, new species, holotype male (MNHN-Na 15482). (a) Carapace and anterior appendages; (b) anterior carapace; (c) telson; (d) eye, dorsal view; (e) antennule; (f) same, distal shorter free ramus of upper flagellum; (g) antenna.




Figure 10. *Palaemonella komaii*, new species, holotype male (MNHN-Na 15482). (a) Mandible; (b) maxillula; (c) maxilla; (d) first maxilliped; (e) second maxilliped; (f) third maxilliped.

antennal spine and near to anterolateral angle of carapace, below first dorsal rostral tooth in lateral view; anterolateral angle of carapace not produced, bluntly rounded.

Abdominal segments smooth, glabrous; sixth segment about 1.85 times length of fifth, subcylindrical, about 1.80 times longer than deep, subuniform, posterolateral angle acute, posteroventral angle with small acute tooth; pleura of first three segments broadly rounded, fourth and fifth posteriorly produced, fourth with posteroventral angle rounded, fifth with posteroventral angle with acute tooth. Telson about 1.19 times sixth abdominal segment length, about 3.4 times longer than anterior width, lateral margins curved ventrally, convergent, posterior margin about 0.4 of anterior margin width, rounded; dorsal surface with two pairs of developed acute dorsolateral spines at 0.33 and 0.67 of telson length, the spines 0.085 of telson length; three pairs of posterior spines, lateral spines short, similar to dorsal spines, about 0.05 of telson length, intermediate spines long, robust, about 0.22 of telson length, submedian spines slender, about 0.5 of intermediate spine length.

Eye well developed, cornea and stalk distinct compressed, width of cornea 0.28 of postorbital carapace length, accessory pigment spot distinct, fused with cornea; stalk thinner than cornea, length 0.9 of corneal width, 1.16 of its width.

Antennular peduncle slender, exceeding rostrum; proximal segment about 2.24 times longer than width, with slender acute stylocerite laterally, reaching to about 0.42 of segment length, anterolateral margin feebly produced, with well-developed slender acute lateral tooth;




Figure 11. *Palaemonella komaii*, new species. (a–c, g–i) Holotype male (MNHN-Na 15482); (d–f) paratype male (MNHN-Na 15600). (a) First pereopod; (b) same, chela; (c) same, coxa, basis and ischium; (d) second pereopod, right; (e) same, chela; (f) third pereopod, left; (g) fifth pereopod, left; (h) same, distal propodal subsegment and dactyl; (i) uropod.

statocyst small; medial margin setose, with ventromedial tooth at about 0.5 of length; intermediate segment obliquely articulated with distal segment dorsal length about 0.3 of proximal segment length, 2.1 of width; distal segment subequal to intermediate segment, slender, about 2.6 times longer than distal width; upper flagellum biramous, with 18 proximal segments of rami fused, shorter free ramus with three segments, with about seven groups of aesthetascs; longer ramus, slender, filiform; lower flagellum slender, filiform.

Antennal basicerite robust, with acute distolateral tooth; carpocerite about 0.29 of scaphocerite length, about 3.2 times longer than width, subcylindrical; flagellum lost; scaphocerite well developed, distinctly exceeding antennular peduncle, slender, about 4.0 times longer than wide, greatest width at about 0.4 of length, distal margin bluntly rounded, lateral margin feebly concave, with strong acute distolateral tooth, overreaching distal lamella.

Epistome with median carina. Fourth thoracic sternite with slender finger-like median process, fifth with well-developed posterior transverse ridge separated by deep median notch, with pair of large acute spines on ridge adjacent to notch; sixth and seventh sternites narrow and unarmed; eighth armed with short stout anteriorly directed median process.

Mouthparts typical of the genus. Mandible with corpus moderately robust, with developed palp with three distal setae and one lateral seta; molar process normal, with four strong blunt teeth, lower inner tooth bilobed; incisor process robust, distally with three (right) or four (left) stout acute teeth, central tooth smaller than outer teeth. Maxillula with feebly bilobed palp, upper lobe tapering tip acute, recurved; upper lacinia robust, distal margin feebly rounded with two rows of acute spines; lower lacinia robust, distally rounded,

with simple setae. Maxilla with short simple non-setose palp, tip acute; basal endite deeply bilobed, upper lobe stouter than lower, with simple setae distally; coxal endite obsolete; scaphognathite well developed, about 3.3 times longer than central width. First maxilliped with simple seta on palp subapically; caridean lobe well developed, with simple setae laterally, exopod with well-developed flagellum, with three long distal plumose setae; basal endite with medial margin with dense simple short setae, coxal endite with medial margin with numerous spines; epipod large, triangular, feebly bilobed. Second maxilliped with normal endopod, dactylar segment about 3.96 times longer than broad, with numerous serrulate spines medially; propodal segment broad, subtriangular, with distal margin with numerous long slender setae; carpus and ischiomerus without special features; basis with distomedial angle acute, with several short setae; coxa feebly produced medially, with four to five simple setae; exopod with slender flagellum with three simple setae distally; epipod oval, with rectangular bilobed podobrach. Third maxilliped with robust endopod, extending distally to end of the carpocerite, ischiomerus and basis fused incompletely, combined segment length about 8.3 times central width, setose medially; intermediate segment about 0.77 of combined proximal segment length, 6.8 times longer than distal width, with eight groups of slender setae medially; terminal segment about 0.44 of combined proximal segment length, distally tapering, about 5.1 times longer than proximal width, with eight dense groups of setae medially and laterally and distal setae; exopod with robust flagellum reaching to 0.83 of combined segment, with four simple plumose setae distally; coxa feebly produced medially, with lobed lateral plate; arthrobranch distinct.

First pereopods slender, exceeding tip of rostrum by length of chela and distal 0.8 carpus; chela slender, with palm feebly compressed, about 2.7 times longer than maximal depth, at about proximal 0.27 of length with five transverse rows of short cleaning setae proximoventrally; fingers 1.1 times palm length, slender, tapering, cutting edges sharp, entire, tips hooked, with several groups of setae laterally; carpus slender, distally feebly swollen, about 1.13 of chela length, 7.8 times longer than distal width, with row of ventral serrulate cleaning setae subdistally; merus subuniform, about 1.2 times chela length, 11.8 times longer than central width, ventral surface plain, with distinct distoventral groove; ischium 0.63 times length of chela, 6.1 times longer than distal, subuniform, with several long setae proximoventrally; basis with ventral margin sinuous, with two short blunt processes, proximal process setose; coxa with well-developed ventromedial process, setose and lobed.

Ambulatory pereopods long, slender. Fifth pereopod exceeds carpocerite by dactyl, propod and carpus, exceeds scaphocerite by dactyl and propod; dactyl long and slender, simple, feebly curved, about 0.17 of propod length, 10.6 times proximal depth, external margin with six long simple setae from 0.4 to 0.7 of dactyl length, outer side with lateral sensory seta at 0.74 of dactyl length; propod uniform, about 1.22 of carapace length, 41.4 times longer than distal width, subdivided into six segments, with pair of short slender simple distoventral spines, three single spines distributed respectively at distoventral margin of third to fifth subsegments, two long setae on distodorsal margin of propod; carpus slender, uniform, about 0.6 of propod length, 21.15 times longer than distal width, unarmed; merus 0.8 of propod length, 27.9 times longer than wide, uniform, unarmed; ischium 0.4 of propod length, 11.9 times longer than distal width, uniform; basis and coxa without special features.

Uropod distinctly exceeding telson; protopodite with posterolateral angle acute; exopod 3.0 times longer than central width, lateral border near straight, with small acute distal tooth and long and acute mobile spine medially, diaeresis distinct; endopod about 0.88 of exopod length, 3.3 times longer than wide.

Paratype. Right second and left fifth pereopods detached.

Similar to holotype. Rostrum dorsal margin with five acute teeth (except epigastric spine). First pereopods exceeding tip of rostrum by length of chela and distal two-thirds of carpus. Second pereopods similar and equal, very slender and elongate, exceeding tip of rostrum by length of chela, carpus and distal two-thirds of merus; chela 1.7 times carapace length, palm subcylindrical, subuniform, smooth, distinctly curved at fingers articulate, 4.3 times longer than central depth, fingers smooth, 0.8 of palm length, with hooked tips, dactyl 7.2 times longer than proximal depth, cutting edge entire with two small teeth at proximal 0.25 and 0.35 length, distal tooth larger than proximal tooth, distal 0.6 cutting edge entire, fixed finger similar to dactyl, cutting edge with five to six small teeth on proximal 0.34 length, distal two teeth larger than proximal teeth, distal three-quarters cutting edge entire; carpus unarmed, distally swollen, 0.9 of palm length, 6.2 times longer than distal width; merus uniform, unarmed, 1.18 of palm length, subuniform, 10.7 times longer than width; ischium unarmed, slightly dorsoventrally compressed, slightly proximally tapered, about 0.71 of palm length, 9.1 times longer than distal width; basis and coxa normal. Third pereopod exceeds scaphocerite tip by propod and dactyl; dactyl long and slender, simple, feebly curved, 0.21 of propod length, 8.0 times proximal depth, external margin with eight long simple setae from 0.4 to 0.75 of dactyl length; propod uniform, 0.97 of carapace length, 34.4 times longer than width, subdivided into four (left) or five (right) segments, with pair of short slender simple distoventral spines, three single spines distributed respectively at distoventral margin of third to fifth subsegments, two long setae on distodorsal margin of propod and two long setae on distodorsal margin of subsegments; carpus slender, uniform, 0.6 of propod length, 16.6 times longer than distal width, unarmed; merus, slender, uniform, subequal to propod, 23.4 times longer than wide, unarmed; ischium 0.44 of propod length, 10.1 times longer than distal width, uniform; basis and coxa without special features. Fourth and fifth pereopods similar to third, fourth exceeds scaphocerite tip by dactyl and distal three-fifths of propod, propod divided into five (left) or four (right) subsegments; fifth pereopod longer than fourth, exceeds scaphocerite tip by propod and dactyl, propod divided into five subsegments.

Measurements (mm)

Holotype. Carapace length, 3.39; carapace and rostrum, 6.06; total body length (proximally), 16.8; first pereopod, chela, 1.95; third pereopod, propod, 4.14.

Paratype. Carapace length, 3.62; carapace and rostrum, 6.22; first pereopod, chela, 1.92; second pereopod, chela, 6.04, third pereopod, propod, 3.51.

Systematic position

Palaemonella komaii sp. nov. is very close to *P. dolichodactylus* Bruce, 1991 and *P. hachijo* Okuno, 1999, sharing the features of very long and slender ambulatory pereopods with the dactyl more than 8.0 times longer than basal depth and with several long setae on dorsal dactyl margin, propod more than 26 times longer than width. The new species can be easily distinguished from *P. hachijo* by the subsegmented and more slender and longer ambulatory propod, and the position of hepatic spine on the carapace which is lower and more anterior, close to the anteroventral angle of the carapace. It differs from *P. dolichodactylus* in the position of the hepatic spine on the carapace which is lower and more anterior, close to the anteroventral angle of the carapace, the longer antennal spine, which

is longer than the hepatic spine and extends to the anterior margin of the basicerite, the ambulatory propod is more slender and longer (41.4 to 26 times longer than distal width).

Etymology

The specific name is given in honour of the Japanese carcinologist, Dr Tomoyuki Komai.

Remarks

The extremely long and slender ambulatory pereopods with the dactyl more than eight times longer than its depth and with several long setae on dorsal dactyl margin, the propod more than 26 times longer than wide separate the new species and its allied species, *P. dolichodactylus* Bruce, 1991 and *P. hachijo* Okuno, 1999, from the other species of the genus *Palaemonella*. They may form a “*P. dolichodactylus* species complex”. The new species looks like an intermediate form between *P. dolichodactylus* and *P. hachijo*, in the features of antennal spine and segmented ambulatory propod. But the ambulatory pereopod varies with increasing habitat depth: shortest and most robust in *P. hachijo* (shallow waters, 0–45 m); moderately long and slender in *P. dolichodactylus* (shallow to deep waters, 16–250 m); longest and most slender in *P. komaii* (deep water, 327–360 m). The posteroventral angle of the fifth abdominal somite is not a stable character to distinguish *P. dolichodactylus* and *P. hachijo*, as used by Okuno (1999) and Bruce (2002b), because all specimens of *P. dolichodactylus* we have checked (including the two specimens of Bruce 1991a) have acute tooth on the posteroventral angle of fifth abdominal pleuron.

Palaemonella pottsi (Borradaile, 1915)

Periclimenes (Falciger) pottsi Borradaile 1915, p 212.

Palaemonella pottsi: Kemp 1922, p 126–127; Chace and Bruce 1993, p 89; Li 2000, p 103, Figure 113; Davie 2002, p 316; Li and Liu 2003, p 159, Figure 5a, b; Li 2004a, p 68.

Material examined

Philippines: (i) MUSORSTOM II, shore station, stn 11, opposite Mactan Marine Station, scuba, 5–18 m, 9 December 1980, 1♂ (MNHN-Na 15954); (ii) Dumaran Channel, NE Palawan Islands, 2–3 m, coll. P. Bouchet, May 1985, 6♂♂, 8♀♀ (6 ovig.) (MNHN-Na 14834); (iii) same, 1♂, 5♀♀ (1 ovig.) (MNHN-Na 14838); (iv) Maribago, 10°17'N, 124°00'E, Mactan Island, Cebu Prov., 3–5 m, coll. P. Bouchet, 9 June 1985, 2♂♂, 3♀♀ (2 ovig.) (MNHN-Na 14835); (v) same, 2–3 m, 1♂ (MNHN-Na 14836); (v) same, 1♂, 2 juveniles (MNHN-Na 14837). **New Caledonia:** (vi) Touho Bank, 28 August 1993, 1 ovig. ♀ (MNHN-Na 15863); (vii) lagoon, 30 August 1993, 1♂, 5♀♀ (2 ovig.) (MNHN-Na 15889). **Loyalty Islands:** ATELIER LIFOU, Lifou, Santal Bay (viii) Gaatcha Bay, stn 1436, 20°55.5'S, 167°04.2'E, 10–20 m, coral head, 10 November 2000, 1♂ (MNHN-Na 15596); (ix) between Cap Mandé and Cap Lefèvre, stn 1452, 20°54.6'S, 167°02.1'E, 2–25 m, scuba, 20 and 22 November 2000, 1 ovig. ♀ (MNHN-Na 15483).

Hosts

(i) (ii) (v) *Comanthina schlegeli* (P. H. Carpenter, 1881); (iv) *Comatella ? stelligera* (P. H. Carpenter, 1888); (iii) washings from *Comatella nigra* (P. H. Carpenter, 1888), *Comanthus bennetti* (J. Müller, 1841), *Comaster multibrachiatus* (P. H. Carpenter, 1888). Identified by A. M. Clark. (vi) (vii) crinoids.




Figure 12. *Palaemonella pusilla* Bruce, 1975, male (MNHN-Na 14839). (a) Carapace; (b) rostrum; (c) telson; (d) same, distal; (e) antennule; (f) antenna.

Remarks

Previously reported from Moalboal Island, Philippines (Bruce 1989a; Markham 1989), and from Nouméa, New Caledonia (Bruce 1970c). Not previously recorded from Loyalty Islands. A well-known associate of the crinoid *Comanthina schlegeli*.

Distribution

Type locality: Mabuag and Murray Islands, Torres Strait. Also known from Zanzibar, Singapore, Japan, Philippines, Indonesia, Australia (Western Australia, Northern Territory, Queensland), New Caledonia, Loyalty Islands, and Marshall Islands.




Figure 13. *Palaemonella pusilla* Bruce, 1975, male (MNHN-Na 14839). (a) Mandible; (b) same, incisor process; (c) first pereiopod; (d) second pereiopod; (e) same, fingers; (f) same, carpus; (g) endopod of second pleopod.

Palaemonella pusilla Bruce, 1975

(Figures 12, 13)

Palaemonella pusilla Bruce 1975c, p 169–177, Figures 1–5; Li 2000, p 104, Figure 114.*Material examined*

New Caledonia: MONTROUZIER, Koumac, (i) Koumac Pass, outer reef, scuba, 20 m, coll. B. Richer de Forges, 6 October 1993, 1♂ (MNHN-Na 14839); (ii) Grand Récif, outer slope, scuba, 12 m, coll. B. Richer de Forges, 7 October 1993, 1♂ (MNHN-Na 14840). **Loyalty Islands:** ATELIER LIFOU, Lifou, Santal Bay, (iii) Pointe Lefèvre, stn 1435, 20°55.2'S, 167°00.7'E, 5–30 m, steep drop-offs and overhangs, 8 November 2000, 2♂♂ (MNHN-Na 15486); (iv) between Huca Hutighé Islet and shore, stn 1421, 20°52.4'S, 167°08.5'E, 4 m, coarse sand on flatstone, 26–27 November 2000, 1♀ (MNHN-Na 15485); (v) west of Easo Point, stn 1451, 20°47.3'S, 167°06.8'E, 10–21 m, coral heads, 19 November 2000, 1♂, 3♀♀ (2 ovig.) (MNHN-Na 15484).

Remarks

Previously known only from the type material. The specimens are not in good condition but are adequate for identification. The Koumac specimen lacks all second to fifth pereopods; the Grand Récif specimen has the right second pereopod still preserved. The rostrum is short and acute, not exceeding the end of the second segment of the antennular peduncle, with a dentition of 1+5/1, without supraorbital ridges or tubercles; antennal spine large, laterally projecting; hepatic spine normal. In the Koumac specimen, the mandible with a small single-segmented palp. Cornea large, about 0.24 of carapace length. Merus of second pereopod without distoventral spine, carpus distally swollen, with strong dorsal tooth.

Distribution

Type locality: Kisiti Island, Kenya. Now also known from New Caledonia and Loyalty Islands.

Palaemonella rotumana (Borradaile, 1898)*Periclimenes rotumanus* Borradaile 1898, p 383.*Palaemonella vestigialis* Kemp 1922, p 123–126, Figures 1, 2, Plate 3, Figure 2.

Palaemonella rotumana: Bruce 1970c, p 276–279, Plate 1e–f; Bruce 1991b, p 229, Figures 5, 6a–e; Chace and Bruce 1993, p 89; Bruce 1996, p 220; Li 2000, p 105, Figure 115; Davie 2002, p 317; Li and Liu 2003, p 160, Figure 5c–q; Li et al. 2004, p 534; Marin et al. 2004, p 203, Figure 4.

Material examined

New Caledonia: (i) Récif Mbere, 22°19.9'S, 166°13.2'E, outer slope, 10 m, coll. P. Bouchet, 5 May 1993, 1♀ (MNHN-Na 15804). MONTROUZIER, (ii) Touho, scuba, 10 m, coll. B. Richer de Forges, 8 September 1993, 1♂, 2 ovig. ♀♀ (MNHN-Na 14841); (iii) 22°19.35'S, 166°25.85'E, 20 m, scuba, on sponge?, coll. I. Takeuchi, 10 November 1995, 1 ovig. ♀ (MNHN-Na 15910); (iv) 22°56.30'S, 167°20.47'E, 18 m, scuba, on *Halimeda*, 29 November 1995, 1 juvenile (MNHN-Na 15851); (v) Thio reef flat, Saint Gabriel, 0.8 m, coll. B. Richer de Forges, 19 May 1996, 1♀ (MNHN-Na 15856); (vi) Thio

reef flat, low tide, 1 m, coll. B. Richer de Forges, 1 July 1996, 1♂ (MNHN-Na 15857); (vii) lagoon, Poingam, low tide, coll. B. Richer de Forges, 10 May 1997, 1♂, 1 ovig. ♀ (MNHN-Na 15807). SURPRISES, North New Caledonia, (viii) stn CP 1376, 18°27.2'S, 163°09.2'E, 39 m, 9 May 1999, coll. Richer de Forges, 1 ovig. ♀ (MNHN-Na 15817); (ix) stn CP 1388, 18°23.8'S, 163°06.9'E, 40 m, 11 May 1999, coll. Richer de Forges, 1♂ (MNHN-Na 15818). **Loyalty Islands:** ATELIER LIFOU, Lifou, Santal Bay, (x) Ngoni Beach stn 1459, 20°47.0'S, 167°03.0'E, 55–80 m, trawl, 5 and 13 November 2000, 3♂♂, 3♀♀ (2 ovig.) (MNHN-Na 15488); (xi) same, 5 November 2000, 1♂ (MNHN-Na 15489); (xii) near Huca Hutighé Islet, stn 1434, 20°52.5'S, 167°08.1'E, 5–20 m, hard bottom, 6 November 2000, 2♂♂ (MNHN-Na 15494); (xiii) Récif Shelter, stn 1461, 20°54.0'S, 167°02.1'E, 100–120 m, trawl, 7, 19, and 23 November 2000, 2 ovig. ♀♀ (MNHN-Na 15491); (xiv) ridge SE of Pointe Aimé Martin (=Acadro), stn 1462, 20°47.1'S, 167°02.1'E, 70–120 m, trawl, 9 and 21 November 2000, 1♀ (MNHN-Na 15487); (xv) east of Santal Bay: Mepinyo, stn 1446, 20°50.8'S, 167°09.7'E, 36–40 m, bottom of slope, 16 November 2000, 1♂, 1♀, 3 juveniles (MNHN-Na 15498); (xvi) north of Cap Aimé Martin (=Acadro), stn 1449, 20°45.8'S, 167°01.65'E, 17 m, brushing, 17 November 2000, 2♀♀ (1 ovig.) (MNHN-Na 15502); (xvii) west of Easo Point, stn 1451, 20°47.3'S, 167°06.8'E, 10–21 m, coral heads, 19 November 2000, 2♂♂ (MNHN-Na 15495); (xviii) east of Easo Point, stn 1418, 20°46.9'S, 167°07.9'E, 1–5 m, sand and sea grasses around yacht wharf, 21 November 2000, 1 ovig. ♀ (MNHN-Na 15500); (xix) between Cap Mande and Cap Lefèvre, stn 1452, 20°54.6'S, 167°02.1'E, 2–25 m, scuba, 22 November 2000, 1♂ (MNHN-Na 15501); (xx) south of Cap Lefèvre (=Nem), stn 1454, 20°56.65'S, 167°02.0'E, 15–18 m, dark drop-off, 23 November 2000, 1♂ (MNHN-Na 15496); (xxi) west to southwest Easo Point, stn 1429, 20°47.5'S, 167°07.1'E, 8–18 m, coral heads, sedimentary channels, 24 November 2000, 2 ovig. ♀♀ (MNHN-Na 15592); (xxii) between Cap Wekutr and Cap Wajez, stn 1455, 20°56.8'S, 167°02.7'E, slope, 15–20 m, 25 November 2000, 2 ovig. ♀♀ (MNHN-Na 15492); (xxiii) between Cap Wekutr and Cap Wajez, stn 1410, 20°56.7'S, 167°03.1'E, 2–4 m, reef edge, 25 November 2000, 1♂ (MNHN-Na 15497); (xxiv) NE of Bay, near Cila, stn 1456, 20°49.3'S, 167°10.4'E, slope, 25–30 m, 26 November 2000, 6♂♂, 1♀ (MNHN-Na 15490); (xxv) between Huca Hutighé Islet and shore, stn 1421, 20°52.4'S, 167°08.5'E, 4 m, coarse sand on flatstone, 26–27 November 2000, 1 ovig. ♀ (MNHN-Na 15499); (xxvi) near Ngoni, stn 1457, 20°46.8'S, 167°02.75'E, 5–10 m, dark overhang and boulder, 27 November 2000, 1♂, 1 ovig. ♀ (MNHN-Na 15493). Chateaubriand Bay, (xxvii) Wé Beach, stn 1474, 20°54.8'S, 167°16.1'E, 0–3 m, fine sand and *Porites* heads, 11 November 2000, 3 ovig. ♀♀ (MNHN-Na 15503).

Remarks

Mandible of male with small two-segmented palp. Specimens with most of second to fifth pereopods missing. Previously reported from New Caledonia, from Nouméa (Bruce 1970c, 1991b). Not previously recorded from Loyalty Islands.

Distribution

Type locality: Rotuma Island, Fiji. Widely distributed from the northern Red Sea to Mozambique, including Australia (Western Australia, Northern Territory, Queensland) and east to Fiji and Hawaii, north to Japan, south to New Caledonia, also found in the Eastern Mediterranean Sea.

Palaemonella spinulata Yokoya, 1936

Palaemonella spinulata Yokoya 1936, p 135, Figure 4; Li 2000, p 106, Figure 116; Davie 2002, p 317.

Material examined

New Caledonia: (i) MONTROUZIER, Touho, sandy island, scuba, coll. B. Richer de Forges, 7 September 1993, 1 ovig. ♀ (MNHN-Na 14842). **Loyalty Islands:** ATELIER LIFOU, Lifou, Santal Bay, (ii) west-southwest of Easo Point, stn 1429, 20°47.5'S, 167°07.1'E, 8–18 m, coral heads, sedimentary channels, 5 November 2000, 1 ovig. ♀? (MNHN-Na 15510); (iii) in front of Ngoni beach, stn 1459, 20°47.0'S, 167°03.0'E, 55–80 m, trawl, 5 and 13 November 2000, 1 ♂ (MNHN-Na 15506); (iv) north of Cap Aimé Martin (=Acadro), stn 1449, 20°45.8'S, 167°01.65'E, 17 m, washing, 17 November 2000, 3 ♂♂, 2 ovig. ♀♀ (MNHN-Na 15512); (v) north of Cap Aimé Martin (=Acadro), stn 1449, 20°45.8'S, 167°01.65'E, 17 m, 17 November 2000, 1 ovig. ♀ (MNHN-Na 15594); (vi) Chépénéhé Point, stn 1420, 20°47.7'S, 167°09.35'E, 4–5 m, flatstone with sediment cover, 18–19 November 2000, 1 ovig. ♀ (MNHN-Na 15507); (vii) between Cap Mandé and Cap Lefèvre (=Nem), stn 1452, 20°54.6'S, 167°02.1'E, 2–25 m, scuba, 20 and 22 November 2000, 3 ovig. ♀♀ (MNHN-Na 15511); (viii) south of Cap Lefèvre (=Nem), stn 1454, 20°56.65'S, 167°02.0'E, 15–18 m, 23 November 2000, 1 ovig. ♀ (MNHN-Na 15508); (ix) west-southwest of Easo Point, stn 1429, 20°47.5'S, 167°07.1'E, 8–18 m, coral heads, sedimentary channels, 24 November 2000, 3 ♂♂, 2 ♀♀, 1 juvenile (MNHN-Na 15509); (x) near Ngoni, stn 1457, 20°46.8'S, 167°02.75'E, 5–10 m, 27 November 2000, 1 ovig. ♀ (MNHN-Na 15505); (xi) Kuendu Bay, 0.5 m, coll. B. Richer de Forges, 5 May 1996, 3 ♂♂, 1 ovig. ♀ (MNHN-Na 15827).

Remarks

Not previously recorded from New Caledonia and Loyalty Islands.

Distribution

Type locality: Misaki, Japan. Otherwise known from Kenya, Tanzania, La Réunion, Japan and Australia (Northern Territory, Queensland), New Caledonia, and Loyalty Islands.

Paraclimenes franklinae (Bruce, 1990) nom. nov.

Periclimenes franklini Bruce 1990b, p 55, Figures 1–5; Bruce 1991a, p 314, Figure 9; Li 2000, p 182, Figure 230.

Paraclimenes franklini: Bruce 1994, p 99, Figure 45; Li 2004b, p 820, Figures 1, 2.

Paraclimenes cf. *franklini*: Li et al. 2004, p 534, Figure 20.

Material examined

Tonga: BORDAU 2, (i) “seamount”, N Ha’apai group, stn CP 1575, 19°42'S, 174°21'W, 232–295 m, 11 June 2000, 1 ovig. ♀ (MNHN-Na 15516); (ii) same, stn CP 1626, 23°20'S, 176°14'W, 220–249 m, 19 June 2000, 2 ovig. ♀♀ (MNHN-Na 15517). **Vanuatu:** (iii) MUSORSTOM 8, stn CP1135, 15°40.50'S, 167°02.43'W, 282–375 m, 11 October 1994, 1 ovig. ♀ (MNHN-Na 15871).

Remarks

Not previously recorded from Tonga. The spelling of the specific name is now corrected according to the ICZN rules.

Distribution

Type locality: Coral Sea. Also reported from South China Sea, New Caledonia, and Tonga.

Paranchistus nobilii Holthuis, 1952

Anchistus Miersi Nobili 1906b, p 48 (non *Harpilius Miersi* De Man 1888).

Paranchistus nobilii Holthuis 1952a, p 100, Figures 41, 42; Bruce 1983b, p 890, Figures 6E, 8I, J; Li 2000, p 111, Figure 120.

Material examined

New Caledonia: lagoon, Five Mile Chenal, scuba, 15–20 m, coll. C. Vadon, 20 September 1978, 1♂, 2 ovig. ♀♀ (MNHN-Na 15886).

Remarks

Not previously recorded from New Caledonia.

Distribution

Type locality: Persian Gulf. Also known from Indonesia, New Caledonia, and Kiribati Islands (Gilbert Islands).

Paranchistus ornatus Holthuis, 1952

Paranchistus ornatus Holthuis 1952a, p 97–100, Figures 39, 40; Hipeau-Jacquotte 1974, p 403; Bruce 1975b, p 162; Fransen 1994, p 117, Plate 2D; Li 2000, p 111, Figure 121.

Material examined

Madagascar: Toliara (Tuléar), Grand Récif, coll. A. Crosnier, October 1958, 1♂ (MNHN-Na 14844).

Host

Pinna sp. [Pinnidae, Mollusca].

Remarks

Well known from Madagascar, where it has been studied in detail by Hipeau-Jacquotte (1974).

Distribution

Type locality: Mozambique. Also known from Kenya, Zanzibar, Comoro Islands, Madagascar, and Seychelles.

Periclimenella petitthouarsi (Audouin, 1825)

Palaemon Petitthouarsi Audouin 1825, p 91.

Periclimenes Petitthouarsi: Borradaile 1898, p 381.

Periclimenella petitthouarsi: Duris and Bruce 1995, p 645–655, Figures 13–18; Li 2000, p 143, Figure 177.

Material examined

Madagascar: Ambalatoaka, Nosy Be, coll. A. Crosnier, September 1958, 1♂ (MNHN-Na 14846).

Remarks

Previously recorded from Madagascar, at Nosy Be by Bruce (1977c, 1978a). The specimen has a rostral dentition of 1+6/4, and possesses both second pereopods.

Distribution

Type locality: Egypt. Also known from Red Sea, Egypt, Israel, Saudi Arabia, Sudan, Eritrea, Yemen, Djibouti, Kenya, Zanzibar, Tanzania, Comoro Islands, Madagascar, and Persian Gulf.

Periclimenella spinifera (De Man, 1902)

Periclimenes Petitthouarsi var. *spinifera* De Man 1902, p 824–826.

Periclimenes (Ancylocaris) spiniferus: Kemp 1922, p 195–196.

Periclimenes (Harpilius) spiniferus: Holthuis 1952a, p 76–78, Figure 30.

Periclimenes spiniferus: Chace and Bruce 1993, p 122.

Periclimenella spinifera: Duris and Bruce 1995, p 656–661, Figures 19, 20; Bruce 1996, p 227; Li 2000, p 144, Figure 178; Davie 2002, p 321; Li and Liu 2003, p 163, Figure 8; Li et al. 2004, p 541.

Material examined

New Caledonia: (i) MONTROUZIER, Toumo Bay, coll. B. Richer de Forges, September 1993, 2♂♂ (MNHN-Na 14847); (ii) lagoon, Ricaudy Reef, low tide, 0.5 m, coll. B. Richer de Forges, 30 June 1996, 1♀ (MNHN-Na 15852); (iii) lagoon, Ricaudy reef flat, low tide, coll. B. Richer de Forges, 22 June 1997, 1♀ (with hemiarthrinid bopyrid), 1 ovig. ♀ (MNHN-Na 15822). **Loyalty Islands:** ATELIER LIFOU, Lifou, Santal Bay, (iv) near Huca Hutighé Islet, stn 1434, 20°52.5'S, 167°08.1'E, 5–20 m, hard bottom, 6 November 2000, 4♀♀ (3 ovig.) (MNHN-Na 15518); (v) Easo, near wharf, stn 1406, 20°46.85'S, 167°07.75'E, intertidal, hard bottom, 10, 13, 14, 16, and 18 November 2000, 2♂♂, 3 ovig. ♀♀ (MNHN-Na 15519); (vi) Gaatcha Bay, stn 1436, 20°55.5'S, 167°04.2'E, 10–20 m, coral head on slope, 10 November 2000, 1♂, 1 ovig. ♀ (MNHN-Na 15520); (vii) east of Easo Point, stn 1418, 20°46.9'S, 167°07.9'E, 1–5 m, sand and sea grasses around yacht wharf, 21 November 2000, 1♀ (MNHN-Na 15521).

Remarks

Previously reported from New Caledonia, from Senez Reef (Bruce 1996). Not previously recorded from Loyalty Islands.

Distribution

Type locality: Ternate, Ambon, Indonesia. Also known from Kenya, Tanzania, Madagascar, Seychelles, La Réunion, Maldives, Chagos Islands, Gulf of Mannar, Andaman Islands, Nicobar Islands, Burma, Malaya, Singapore, Vietnam, South China Sea, Japan, Philippines, Indonesia, Papua New Guinea, Australia (Western Australia, Northern Territory, Queensland), Mariannas Islands, New Caledonia, Loyalty Islands, Marshall Islands, Fiji, Samoan Islands, and French Polynesia (Society Islands).

Periclimenes affinis (Zehntner, 1894)

Palaemonella affinis Zehntner 1894, p 208.

Periclimenes (Harpilius) affinis: Holthuis 1958, p 6–8, Figure 2.

Periclimenes affinis: Bruce 1980b, p 2–8, Figures 1–3; Chace and Bruce 1993, p 99; Li 2000, p 149, Figure 180; Davie 2002, p 323; Li and Liu 2004, p 89, Figure 1a–f.

Material examined

Philippines: (i) Maribago, Mactan Island, Cebu, 10°17'N, 124°00'E, 2–3 m, crinoid washings, coll. P. Bouchet, 9 June 1985, 1♂, 1♀?, 3 juveniles (MNHN-Na 14849); (ii) same, 1♀, 2 juveniles (MNHN-Na 14850); (iii) Dumaran Channel, NE Palawan Islands, 2–3 m, crinoid washings, coll. P. Bouchet, May 1985, 3♂♂, 15♀♀ (9 ovig.), 3 juveniles (MNHN-Na 14848). **New Caledonia:** (iv) Touho Bank, on crinoids, 28 August 1993, 4♂♂, 4♀♀ (ovig.) (MNHN-Na 15858); (v) lagoon, on crinoids, 30 August 1993, 29 spms (7 ovig. ♀♀) (MNHN-Na 15865); (vi) Touho, Kohe, 8 September 1993, 2♂♂ (MNHN-Na 15901).

Hosts

(i) (iii) *Comanthina schlegeli* (P. H. Carpenter, 1882); (ii) *Comatella ?stelligera* (P. H. Carpenter, 1880), det. A. M. Clark. (vi) *Comatella* sp. [Crinoidea, Echinoderma].

Remarks

Previously reported from the Philippines, with 3 ovig. ♀♀ collected in 1908, from Siasi, Sulu Islands (Chace and Bruce 1993).

Distribution

Type locality: Ambon, Indonesia. Also known from South China Sea, Japan, Philippines, Indonesia, Australia (Northern Territory, Queensland), and New Caledonia.

Periclimenes alcocki Kemp, 1922

Palaemon (*Brachcarpus*) *laccadivensis* Alcock 1901, p 138–139 (*partim*).

Periclimenes (*Periclimenes*) *alcocki* Kemp 1922, p 154–156, Figures 21–24.

Periclimenes alcocki: Bruce 1978a, p 227–228, Figure 10; Bruce 1991a, p 302–308, Figures 2–5; Chace and Bruce 1993, p 102; Bruce 1996, p 228, Figure 11b–d; Li 2000, p 151, Figure 184; Davie 2002, p 323.

Material examined

Philippines: MUSORSTOM 3, (i) stn CP105, 13°52'N, 120°30'E, 398–417 m, 1 June 1985, 1 ovig. ♀ (MNHN-Na 14853); (ii) stn CP125, 11°57'N, 121°28'E, 388–404 m, 4 June 1985, 1 ovig. ♀ (MNHN-Na 14851); (iii) same, 1 ovig. ♀ (MNHN-Na 14852). **Indonesia:** KARUBAR, Tanimbar Islands, (iv) stn CC21, 5°14'S, 133°00'E, 688–694 m, 25 October 1991, 1 ♂ (MNHN-Na 14860); (v) stn CP59, 8°20'S, 132°11'E, 405–399 m, 31 October 1991, 2 ovig. ♀♀ (MNHN-Na 14856); (vi) stn CP69, 8°42'S, 131°53'E, 356–368 m, 2 November 1991, 1 ♀ (MNHN-Na 14857); (vii) same, 2 ♀♀ (MNHN-Na 14858); (viii) stn CP75, 8°46'S, 131°36'E, 451–452 m, 3 November 1991, 1 ovig. ♀ (MNHN-Na 14862); (ix) same, 1 ♀ (MNHN-Na 14859); (x) stn CP76, 8°50'S, 131°33'E, 401–400 m, 3 November 1991, 2 ovig. ♀♀ (MNHN-Na 14854); (xi) stn CP83, 9°23'S, 131°00'E, 285–292 m, 4 November 1991, 1 ovig. ♀ (MNHN-Na 14861). **New Caledonia:** BIOCAL, (xii) stn CP78, 22°16.25'S, 167°15.53'E, 445–450 m, 5 September 1985, 1 ♀ (MNHN-Na 14867). MUSORSTOM 4, (xiii) stn 202, 18°58.0'S, 169°59.3'E, 560 m, 20 September 1985, 1 ♂ (MNHN-Na 14864); (xiv) stn 236, 22°11.3'S, 167°15.0'E, 495–550 m, 2 October 1985, 1 ♂ (MNHN-Na 14866). MUSORSTOM 5, (xv) stn 384, 19°42.4'S, 158°50.8'E, 772–756 m, 21 October 1986, 1 ♀ (MNHN-Na 14863). BATHUS 4, (xvi) stn CP911, 18°57.80'S, 163°08.47'E, 566–558 m, 5 August 1994, 2 ♀♀ (MNHN-Na 15846); (xvii) stn CP921, 18°46.72'S, 163°17.01'E, 613–610 m, 6 August 1994, 1 ♂ (MNHN-Na 15631). NORFOLK 2, (xviii) Île des Pins, stn DW 2147, 22°50'S, 167°16'W, 496 m, 4 November 2003, 1 ♂, 6 ovig. ♀♀ (MNHN-Na 15523). **Tonga:** (xix) BORDAU 2, “seamount”, stn CP 1625, 23°28'S, 176°22'W, 824 m, 19 June 2000, 1 ♀ (MNHN-Na 15522). **Fiji:** BORDAU 1, (xx) stn CP 1446, 17°11'S, 178°42'W, 350–367 m, 3 March 1999, 1 ♂, 1 ovig. ♀ (MNHN-Na 15524); (xxi) stn DW 1497, 18°44'S, 178°25'W, 335–350 m, 12 March 1999, 1 ovig. ♀ (MNHN-Na 15527). **Solomon Islands:** SALOMON 1, (xxii) stn CP 1804, 9°32.0'S, 160°37.4'E, 609–328 m, 2 October 2001, 1 ♂, 1 ovig. ♀ (MNHN-Na 15529); (xxiii) stn CP 1860, 9°22'S, 160°31'E, 620 m, 7 October 2001, 1 ovig. ♀ (MNHN-Na 15528).

Remarks

Previously recorded from Philippines, from 14°00'N, 120°16'E (Bruce 1981a), Indonesia, from the Tanimbar Islands (Bruce 1996), and Loyalty Islands and New Caledonia (Bruce 1991a).

Distribution

Type locality: Laccadive Sea, 9°34'57"N, 70°36'30"E, 930 m (erroneously given as 90°N in Bruce 1996). Also previously reported from Philippines, Indonesia, Australia (New South Wales), and New Caledonia.

***Periclimenes aleator* Bruce, 1991**

(Figure 14)

Periclimenes aleator Bruce 1991a, p 315–322, Figures 10–14; Li 2000, p 152, Figure 185.*Material examined*

Indonesia: KARUBAR, (i) Kai Islands, stn CP16, 5°17'S, 132°50'E, 315–349 m, 24 October 1991, 1 ovig. ♀ (MNHN-Na 14869); (ii) Tanimbar Islands, stn CP38, 7°40'S, 132°27'E, 620–666 m, 28 October 1991, 1♀ (MNHN-Na 14868). **Solomon Islands:** (iii) SALOMON 1, stn CP 1786, 9°21.3'S, 160°24.6'E, 387 m, 30 September 2001, 1♂ (MNHN-Na 15532). SALOMON 2, (iv) Vella Gulf, stn CP2260, 8°03.5'S, 156°54.5'E, 399–427 m, November 2004, 2 ovig. ♀♀ (MNHN-Na 15946). **Vanuatu:** (v) MUSORSTOM 8, stn CP1136, 15°40.62'S, 167°01.60'W, 398–400 m, 11 October 1994, 1 ovig. ♀ (MNHN-Na 15873). **New Caledonia:** SMIB 8, (vi) stn DW146, 24°55.20'S, 168°21.73'E, Banc Éponge (Mont B), 514–522 m, 27 January 1993, 1 ovig. ♀ (MNHN-Na 14946). BATHUS 4, (vii) stn DW899, 20°16.68'S, 163°50.26'W, 500–600 m, 3 August 1994, 1 ovig. ♀ (MNHN-Na 15870); (viii) stn DW912, 18°55.61'S, 163°07.68'W, 702–690 m, 5 August 1994, 1♀ (MNHN-Na 15867); (ix) stn DW923, 18°51.51'S, 163°24.17'W, 502–470 m, 6 August 1994, 1 ovig. ♀ (MNHN-Na 15868). LITHIST, (x) stn CP09, 24°52.8'S, 168°21.8'E, 518–540 m, Banc Éponge, 11 August 1999, 1 ovig. ♀ (MNHN-Na 15891). NORFOLK 2, (xi) Norfolk Ridge, Kaimon Maru Bank, stn DW 2098, 24°42'S, 168°06'W, 550–668 m, 29 October 2003, 1 ovig. ♀ (MNHN-Na 15934); (xii) Norfolk Ridge, Antigonina Bank, stn CP 2122, 23°22'S, 168°00'W, 560–577 m, 1 November 2003, 1 ovig. ♀ (MNHN-Na 15935). **Loyalty**




Figure 14. *Periclimenes aleator* Bruce, 1991, ovigerous female (MNHN-Na 14946). (a) Carapace and anterior appendages; (b) telson; (c) right (major) second pereiopod; (d) same, fingers; (e) third pereiopod, distal propod and dactyl.

Islands: (xiii) MUSORSTOM 6, stn CP467, 21°05.13'S, 167°32.11'E, 575 m, 21 February 1989, 1♂ (MNHN-Na 14870). **Fiji:** BORDAU 1, (xiv) stn DW 1499, 18°40'S, 178°27'W, 389–400 m, 12 March 1999, 1 ovig. ♀ (MNHN-Na 15525); (xv) stn CP 1504, 18°13'S, 178°34'W, 427–440 m, 13 March 1999, 1 ovig. ♀ (MNHN-Na 15526).

Parasites

The Loyalty Islands male (xiii) had a ventral abdominal hemiarthrinid parasite, *Paraphrixus nigrocinctus* (Chopra) (Isopoda: Bopyridae).

Remarks

Previously known only from the type material from the Loyalty Islands from 570–610 m. The record from 315–349 m depth from the Kai Islands (i) represents not only a new locality, but also a much shallower bathymetric range than that of type locality.

The ovigerous female from stn DW146 (vi) with the cornea large, about 0.18 of posterior orbital carapace length, pereopod with the distoventral and subdistoventral spines of propod finely pectinate distodorsally, the second pereopods similar and subequal.

Distribution

Type locality: Loyalty Islands. Now also known from Indonesia, Solomon Islands, Vanuatu, New Caledonia, and Fiji.

Periclimenes amboinensis (De Man, 1888)

Anchistia amboinensis De Man 1888, p 546–548, Plate 22a, Figure 2.

Periclimenes amboinensis: Borradaile 1898, p 385; Bruce 1991b, p 235; Chace and Bruce 1993, p 102; Bruce 1996, p 231, Figures 11h, i, 28f; Li 2000, p 153, Figure 187; Davie 2002, p 323.

Material examined

Phillipines: (i) Maribago, Mactan Island, Cebu, 10°17'N, 124°50'E, 3–5 m, from mixed crinoid washings, coll. P. Bouchet, 9 June 1985, 1♂ (MNHN-Na 14873); (ii) same, 3♂♂, 8♀♀ (4 ovig.) (MNHN-Na 14874); (iii) same, 10°17'N, 124°00'E, 2–3 m, 3♂♂, 1 ovig. ♀ (MNHN-Na 14875); (iv) same, 1♂, 1♀ (MNHN-Na 14876). **New Caledonia:** (v) Touho Bank, on crinoids, 28 August 1993, 2♂♂, 1♀ (MNHN-Na 15898). **MONTROUZIER,** Touho, (vi) 5–7 m, coll. B. Richer de Forges, September 1993, 1 ovig. ♀ (MNHN-Na 14872); (vii) 10 m, scuba, from crinoids, 8 September 1993, 1♂ (MNHN-Na 15809); (viii) slope before Kobe, associated with comatulids, 8 September 1993, 19♂♂, 41♀♀ (15 ovig.) (MNHN-Na 15805). **Loyalty Islands:** **ATELIER LIFOU,** Lifou, Santal Bay, (ix) Chépenhe Bay, stn 1440, 20°47.2'S, 167°08.6'E, 15–35 m, coral heads and pass sedimentary near wharf, 11, 13, and 16 November 2000, 4♂♂, 4♀♀ (3 ovig.) (MNHN-Na 15872); (x) between Cap Mandé and Cap Lefèvre (=Nem), stn 1452, 20°54.6'S, 167°02.1'E, 2–25 m, scuba, 20 and 22 November 2000, 1 ovig. ♀ (MNHN-Na 15533).

Hosts

(i) (ii) (iii) *Comanthina schlegeli* (P. H. Carpenter, 1882), *Comatella nigra schlegeli* (P. H. Carpenter, 1882), *Comaster multibrachiata schlegeli* (P. H. Carpenter, 1888), *Comanthus*

bennetti (J. Müller, 1841); (iv) *Comatella ?stelligera schlegeli* (P. H. Carpenter, 1888), det. A. M. Clarke; (viii) comatulids; (ix) *Tropiometra afra* (Hartlaub, 1890) [Crinoidea, Echinoderma].

Parasites

One male of (v) and one specimen of (x) have branchial bopyrid isopod parasites.

Remarks

Not previously reported from the Philippines. Recorded from Récif Amere, New Caledonia by Bruce (1991b). All the specimens have the telson with two pairs of dorsolateral spines located marginally, from barely discernible to very distinct, although all are very small. The wide range of morphological variation also present in rostral formula, hepatic spine and ambulatory propod. The rostral formula in the samples of (viii) varies from 0/0 (3♂♂, 4♀♀), 2/0 (1♀), 3/0 (1 ovig. ♀), to 4/1 (3 ovig. ♀♀), 5/1 (2♂♂, 2♀♀, 2 ovig. ♀♀), 5/2 (1♂) and 6–7/1–2, most samples have the formula 7/1. The hepatic spines vary from failing to reach the anterior carapace margin to reaching the margin (most samples), to exceeding the margin by distal one-quarter of the spine (most of the samples without or with fewer rostral teeth). Because of the series variation in the rostral formula and the hepatic spines, it is difficult to identify the samples of (x) as *P. amboinensis* (De Man, 1888) or *P. ceratophthalmus* Borradaile, 1915. As most samples have a ventral rostral tooth, we treat the samples as *P. amboinensis* instead of *P. ceratophthalmus* temporarily until we check the type material of the two forms and more material. The wide range of variation and the diversity of host species suggest that a complex of closely related species exists possibly in association with the wide variety of host crinoids.

Distribution

Type locality: Ambon, Indonesia. Also known from Maldives, Indonesia, Australia (Western Australia, Queensland), New Caledonia, Loyalty Islands, and Marshall Islands.

Periclimenes attenuatus Bruce, 1971

Periclimenes attenuatus Bruce 1971b, p 533–543, Figures 1–5; Chace and Bruce 1993, p 103; Li 2000, p 159, Figure 196; Davie 2002, p 324; Li 2004a, p 68, Figures 1, 2.

Material examined

Philippines: Maribago, Mactan Island, Cebu, 10°17'N, 124°00'E, 2–3 m, from mixed crinoid washings, coll. P. Bouchet, 9 June 1985, 1 ovig. ♀ (MNHN-Na 14878).

Host

Comanthina schlegeli (P. H. Carpenter, 1881), det. A. M. Clark [Crinoides, Echinoderma].

Remarks

Not previously reported from the Philippines. The single specimen has a rostral dentition of 4/1, with a very small ventral tooth. The first pereopods have the carpus about three times longer than the chela, rather swollen distally; the fingers broad, densely setose laterally. Only the extremely slender minor second pereopod is preserved.

Distribution

Type locality: Duke of York Islands. Also known from Philippines, Indonesia, and Australia (Queensland).

Perclimenes brevicarpalis (Schenkel, 1902)

Ancylocaris brevicarpalis Schenkel 1902, p 563, Plate 13, Figure 21.

Perclimenes (Ancylocaris) brevicarpalis: Kemp 1922, p 185–191, Figures 40–42, Plate 67.

Perclimenes (Harpilius) brevicarpalis: Holthuis 1952a, p 69–73, Figure 27.

Perclimenes brevicarpalis: Bruce 1991b, p 236; Chace and Bruce 1993, p 104; Li 2000, p 161, Figure 199; Davie 2002, p 324; Li and Liu 2004, p 92, Figure 3.

Material examined

Madagascar: (i) northwest coast, Ambalatoaka, Nosy Be, coll. A. Crosnier, December 1958, 1♂, 1♀ (MNHN-Na 14883); (ii) Nosy Be, intertidal, coll. A. Crosnier, September 1959, 1♂, 1♀ (MNHN-Na 14882); (iii) same, 2♂♂?, 3♀♀ (2 ovig.) (MNHN-Na 14881); (iv) Tuléar, Grand Récif, with anemone *Stoichactis* sp., coll. R. Hipeau-Jacquotte, 1964, 2♂♂, 1 ovig. ♀ (MNHN-Na 15921); (v) Tuléar, Grand Récif, with anemone *Stoichactis* sp., coll. R. Hipeau-Jacquotte, 1968, 8♂♂, 1♀ (MNHN-Na 15831); (vi) same, 7♂♂, 6 ovig. ♀♀ (MNHN-Na 15916); (vii) same, 1♂, 4 ovig. ♀♀ (MNHN-Na 15917); (viii) same, 4♂♂, 5 ovig. ♀♀ (MNHN-Na 15919); (ix) Tuléar, Grand Récif, with anemone, coll. R. Hipeau-Jacquotte, no date record, 1♀ (MNHN-Na 15924). **New Caledonia:** MONTROUZIER, (x) Ouanap, Koumac, 1 m, on “choux-fleur (cabbage)” anemone, coll. S. Gofas, 19 October 1993, 1♀ (MNHN-Na 14879); (xi) Ouanap Bay, Koumac, intertidal, 22 October 1993, 1♂, 1♀ (MNHN-Na 114880). **Loyalty Islands:** ATELIER LIFOU, Lifou, (xii) Santal Bay: Gaatcha Bay, stn 1463, 20°55.05'S, 167°03.35'E, 20–30 m, dredge, sand and corals, 10 November 2000, 1♀ (MNHN-Na 15534).

Host

(iv)–(viii) *Stoichactis* sp. (anemone) [Actinaria, Coelenterata].

Remarks

The specimens have a rostral dentition of either 5/1 or 6/1. The species has been previously recorded from Madagascar, from Toliara (Tuléar), by Hipeau-Jacquotte (1973) and Bruce (1978a), and from the Northwest Lagoon, New Caledonia, by Bruce (1991b).

Distribution

Type locality: “Makassar” Indonesia. Also known from Gulf of Aqaba, Red Sea, Djibouti, Kenya, Zanzibar, Tanzania, Mozambique, Madagascar, Seychelles, Mauritius, Persian Gulf, Maldives, south India, Andaman Islands, Malaya, Singapore, Vietnam, South China Sea, Ryukyu Islands, Japan, Philippines, Indonesia, Papua New Guinea, Australia

(Western Australia, Northern Territory, Queensland), Caroline Islands, Solomon Islands, New Caledonia, Loyalty Islands, and Marshall Islands.

Periclimeses brevirostris Bruce, 1991

Periclimeses brevirostris Bruce 1991a, p 321–330, Figures 15–20; Li 2000, p 163, Figure 200.

Material examined

Solomon Islands: (i) southwest Santa Isabel, stn CP2194, 8°24.8'S, 159°26.7'E, 440–521 m, November 2004, 1 ovig. ♀ (MNHN-Na 15951). **Vanuatu:** (ii) MUSORSTOM 8, stn CP1028, 17°54.01'S, 168°40.42'W, 624–668 m, coll. B. Richer de Forges, 28 September 1994, 1♀ (MNHN-Na 15882). **New Caledonia:** (iii) BATHUS 1, stn CP709, 21°41.78'S, 166°37.88'E, 650–800 m, 19 March 1993, 1 ovig. ♀ (MNHN-Na 14884).

Remarks

Previously known only from the type material.

Distribution

Type locality: Île des Pins, New Caledonia, 500 m. Known only from New Caledonia.

Periclimeses calcaratus Chace and Bruce, 1993

(Figure 15)

Periclimeses calcaratus Chace and Bruce 1993, p 104–105, Figure 21; Li 2000, p 165, Figure 203.

Material examined

Philippines: MUSORSTOM 3, stn CP103, 14°00'N, 120°18'E, 193–200 m, 1 June 1985, 1♂ (MNHN-Na 14886).

Remarks

Previously known only from the type specimen from the Philippines.

Distribution

Type locality: Albay Gulf, Philippines, 267 m. Not known outside Philippines.




Figure 15. *Perichlimes calcaratus* Chace and Bruce, 1993, male (MNHN-Na 14886). (a) Anterior carapace; (b) antennule; (c) same, distolateral angle of proximal peduncular segment; (d) antenna; (e) third pereiopod, distal propod and dactyl; (f) same, distal corpus and unguis; (g) distoventral propodal spine.

Periclimenes ceratophthalmus Borradaile, 1915

Periclimenes (Corniger) ceratophthalmus Borradaile 1915, p 211.

Periclimenes (Ancylocaris) ceratophthalmus: Kemp 1922, p 172.

Periclimenes (Periclimenes) ceratophthalmus: Kemp 1925, p 324, Figure 18.

Periclimenes ceratophthalmus: Chace and Bruce 1993, p 106; Li 2000, p 167, Figure 206; Davie 2002, p 325.

Material examined

New Caledonia: (i) MONTROUZIER, 1993, Touho-Kohe, 15 m, scuba, on comatulid crinoid, coll. B. Richer de Forge, 7 September 1993, 1 ovig. ♀ (MNHN-Na 14887).

Loyalty Islands: (ii) ATELIER LIFOU, Lifou, Santal Bay: north of Cap Aimé Martin (=Acadro), stn 1447, 20°45.8'S, 167°01.65'E, 17–31 m, scuba, 17 and 22 November 2000, 1 ovig. ♀ (MNHN-Na 15535).

Remarks

Not previously recorded from New Caledonia and Loyalty Islands. The single sample of (i) has a rostral dentition of 4/1 and has a single detached second pereopod only. The cornea has an elongated terminal papilla and the dorsal telson spines are minute.

Distribution

Type locality: Hulule, Malé Atoll, Maldives. Also reported from Kenya, Zanzibar, Seychelles, Maldives, Indonesia, Australia (Queensland), Caroline Islands, Solomon Islands, New Caledonia, and Loyalty Islands.

Periclimenes commensalis Borradaile, 1915

Periclimenes (Cristiger) commensalis Borradaile 1915, p 211; Borradaile 1917, p 364.

Periclimenes commensalis: Chace and Bruce 1993, p 107; Bruce 1996, p 232, Figure 12a–g; Li 2000, p 169, Figure 209; Davie 2002, p 325.

Material examined

Philippines: (i) Dumarán Channel, NE Palawan Islands, 2–3 m, washings from crinoids, coll. P. Bouchet, May 1985, 1♂, 4♀♀ (2 ovig.) (MNHN-Na 14888); (ii) Maribago, Mactán Island, Cebu, 10°17'N, 124°00'E, 3–5 m, washings from crinoids, coll. P. Bouchet, 9 June 1985, 1♂, 1♀?, 2 juveniles (MNHN-Na 14889). **New Caledonia:** (iii) Touho Bank, on crinoids, 28 August 1993, 6♂♂, 9♀♀ (4 ovig.) (MNHN-Na 15896); (iv) lagoon, on crinoids, 30 August 1993, 8♀♀ (2 ovig.) (MNHN-Na 15847); (v) Touho, Kohe, 8 September 1993, 3♂♂, 2♀♀ (MNHN-Na 15888).

Hosts

(i) (ii) *Comanthina schlegeli* (P. H. Carpenter, 1881); (ii) also *Comatella nigra* (P. H. Carpenter, 1888), *Comanter multibrachiata* (P. H. Carpenter, 1888), *Comanthus bennetti* (J. Müller, 1841); det. A. M. Clark. (v) *Comatella* sp. [Crinodea, Echinoderma].

Parasites

A branchial bopyrid is present on one of the specimens from Maribago. A bopyrid is also present beneath the abdomen of one male (iii). A couple of bopyrids are present beneath the abdomen of one female (v).

Remarks

Not previously recorded from the Philippines. Previously reported from Nouméa, New Caledonia, by Monod (1976). The Dumaran Channel specimens were collected in association with *Palaemonella pottsii* and *Periclimenes affinis*. The Touho specimens were found with *Periclimenes amboinensis*.

The majority of specimens still have both second pereopods attached. The chelae are subequal and similar, short and stout, about 1.2 times the carapace length, with a well-marked “ball and socket” articulation with the similarly stout carpus. The distal cutting edges of the fingers are minutely serrate. The ambulatory dactyl of the third pereopod is biunguiculate with a clearly demarkated unguis, the proximal dorsal aspect of which bears a strong central non-articulated spiniform tooth, flanked medially and laterally by smaller similar teeth, not always readily visible. The propod is sparsely setose and the spines are distally dentate as reported in Bruce (1980a, 1982, 1996). Borradaile (1915, 1917) did not provide any illustrations with his description of the type material of this species. The second specimen discovered unfortunately lacked the second pereopods (Holthuis 1952a). Miyake and Fujino (1968), reporting on specimens from the Palau Islands, were the first to describe and illustrate the second pereopods but they did not comment on the carpo-propodal articulation. Monod (1976) described and illustrated a specimen from New Caledonia but did not provide information on the second pereopods. Bruce (1980a) similarly omitted reference to these appendages in reporting on a juvenile associated with a holothurian host in the Solomon Islands. These descriptions also illustrate the ambulatory dactyls, but show or describe only a single simple dorsal spine and not a trifold spine as in the present specimens.

Distribution

Type locality: Murray Island, Torres Strait. Also known from Kenya, Zanzibar, Mozambique, China (Hong Kong), Japan, Philippines, Indonesia, Australia (Northern Territory, Queensland, Elizabeth and Middleton Reefs), Tasman Sea, Caroline Islands, Solomon Islands, New Caledonia, Marshall Islands, and Fiji.

Periclimenes cristimanus Bruce, 1965

Periclimenes cristimanus Bruce 1965, p 487–493, Figures 1, 2; Chace and Bruce 1993, p 108; Li 2000, p 173, Figure 215; Davie 2002, p 326.

Material examined

Philippines: Maribago, Mactan Island, Cebu, 10°17'N, 124°00'E, 0.5–2 m, washings from crinoids, coll. P. Bouchet, 10 June 1985, 1 ovig. ♀, 2 juveniles (MNHN-Na 14890).

Host

Echinothrix calamaris (Pallas, 1774) [Echinoidea, Echinoderma].

Remarks

Not previously recorded from the Philippines.

Distribution

Type locality: Pulau Sudong, Singapore. Also known from Malaya, China (Hong Kong), Philippines, Australia (Queensland), and the Marshall Islands.

Periclimenes crosnieri sp. nov.

(Figures 16–18)

Material examined

Indonesia: KARUBAR, Tanimbar Islands, stn CP45, 7°54'S, 132°47'E, 302–305 m, 29 October 1991, ♀ holotype (cl 5.81 mm); ♀ paratype (cl 5.68 mm) (MNHN-Na 14943).

Description

Holotype. A medium body size pontoniine shrimp of subcylindrical body form.

Carapace smooth, glabrous. Rostrum well developed, moderately deep, compressed, extending beyond antennular peduncle, about 0.67 of carapace length, not reaching distal margin of scaphocerite, anteroventrad proximally, upcurved distally; dorsal carina deep, with seven subequal, evenly distributed along whole length, teeth acute, first tooth situated just above posterior orbital margin; lateral carinae feebly developed, slightly upcurved distally; ventral carina deep distal half, with two low acute teeth on distal third, smaller than dorsal teeth; interdental spaces and proximal ventral carina feebly setose. Supraorbital spine absent, epigastric spine situated at anterior 0.25 of the carapace length; orbit feebly developed, inferior orbital angle strongly produced, bluntly round in dorsal view; antennal spine small slender marginal, distinctly below inferior orbital angle, not exceeding inferior orbital angle; hepatic spine larger than antennal spine, slender, below level of antennal spine, below and between epigastric spine and first dorsal rostral tooth in lateral view; anterolateral angle of carapace not produced, bluntly rounded.




Figure 16. *Periclimenes crosnieri*, new species, holotype female (MNHN-Na 14943), body, lateral view.




Figure 17. *Periclimenes crosnieri*, new species. (a, c–k) Holotype female; (b) paratype female (MNHN-Na 14943). (a) Anterior carapace; (b) carapace; (c) right eye, lateral view; (d) telson; (e) right antennule; (f) same, peduncle, medial view; (g) right antenna; (h) mandible; (i) maxilla; (j) second maxilliped; (k) third maxilliped.

Abdominal segments smooth, glabrous; sixth segment about 1.57 times length of fifth, subcylindrical, about 1.53 times longer than deep, subuniform, posterolateral angle acute, posteroventral angle rounded; pleura of first three segments small, broadly rounded, fourth and fifth posteriorly produced, bluntly rounded. Telson about 1.64 times sixth segment length, about 3.6 times longer than anterior width, lateral margins with anterior third curved ventrally, posterior two-thirds straight, convergent, posterior margin about 0.36 of anterior margin width, rounded, without posterior median point; dorsal surface with sparse long setae, with four pairs of small but well-developed dorsolateral spines at 0.41, 0.59, 0.76, and 0.88 of telson length; three pairs of posterior spines, lateral spines short, similar to dorsal spines, about 0.02 of telson length, intermediate spines long, robust, about 0.08 of telson length, submedian spines robust, about 0.43 of intermediate spine length.

Eye small, with feebly pigmented globular cornea, diameter of cornea 0.14 of posterior orbital carapace length, without accessory pigment spot; stalk subequal to corneal diameter, subequal to its width.

Antennular peduncle overreaching to seventh dorsal rostral tooth; proximal segment about 2.0 times longer than central width, with slender acute stylocerite laterally, reaching to about 0.7 of segment length, anterolateral margin feebly produced, setose, with long slender acute lateral tooth reaching to the end of dorsal margin of intermediate segment; statocyst small; medial margin setose, with ventromedial tooth at about 0.5 of length; intermediate segment obliquely articulated with distal segment, dorsal length about 0.25 of proximal segment length, 0.7 of width, lateral margin expanded, setose; distal segment




Figure 18. *Periclimenes crosnieri*, new species. (a–f, i) Holotype female; (g, h) paratype female (MNHN-Na 14943). (a) First pereiopod; (b) same, chela; (c) major second pereiopod; (d) same, fingers; (e) minor second pereiopod; (f) third pereiopod; (g) same, propod and dactyl; (h) same, distal propod and dactyl; (i) uropod.

about 2.1 times intermediate segment length, 0.46 of proximal segment length, slender, about 1.9 times longer than distal width; upper flagellum biramous, with eight proximal segments of rami fused, shorter free ramus with five segments, length about 0.46 of fused portion, total length about 0.38 of carapace length, with about 17 groups of aesthetascs; longer ramus, slender, filiform, about 0.45 times carapace length; lower flagellum slender, filiform, about 0.73 times carapace length.

Antennal basicerite robust, with acute lateral tooth; carapocerite about 0.45 of length of lateral margin of scaphocerite, about 3.8 times longer than width, subcylindrical, reaching about 0.42 of total length of scaphocerite; flagellum well developed, slender, about 2.6 times postorbital carapace length; scaphocerite well developed, distinctly exceeding tip of rostrum, broad, about 2.1 times longer than wide, greatest width at about 0.4 of length, distal margin bluntly rounded, lateral margin feebly convex, with strong acute distolateral tooth, reaching near to distal lamella.

Epistome unarmed. Fourth thoracic sternite without slender median process, with low transverse ridge with shallow median notch; posterior sternites narrow.

Mouthparts typical of the genus. Mandible with corpus moderately robust, without palp; molar process normal, with four strong blunt teeth, lower inner tooth bilobed; incisor process robust, obliquely truncate distally with three stout acute teeth, central tooth smaller than outer teeth. Maxilla with short simple non-setose palp; basal endite deeply bilobed, upper lobe stouter than lower, with simple setae distally; coxal endite obsolete, medial

margin convex; scaphognathite well developed, about 3.2 times longer than central width, posterior lobe about 0.34 of scaphocerite length, 1.5 times longer than wide, anterior lobe with median margin concave, about 1.7 times longer than wide. Second maxilliped with normal endopod, dactylar segment about 4.0 times longer than broad, with numerous serrulate spines medially; propodal segment broad, with distal margin with numerous long finely serrulate spiniform setae; carpus, ischiomerus and basis without special features; coxa angularly produced medially, with four to five simple setae; exopod with slender flagellum with eight to nine plumose setae distally; epipod simple, without podobranch. Third maxilliped with slender endopod, extending distally to the middle of the carpocerite, ischiomerus and basis fused incompletely, basal portion medially expanded, convex, combined segment length about 5.0 times central width, compressed, setose medially, with submarginal row of 15–17 very short small spines on proximal medial ischial portion; intermediate segment about 0.45 of combined proximal segment length, 3.2 times longer than central width, with several groups of long finely serrulate spiniform setae medially; terminal segment about 0.3 of combined proximal segment length, distally tapering, about 3.2 times longer than proximal width, with sparse spiniform setae medially, with long distal spines; exopod with slender flagellum with 12–14 plumose setae distally; coxa feebly produced medially, with oval lateral plate; arthrobranch distinct.

First pereopods moderately slender, exceeding tip of rostrum by length of chela; chela with palm subcylindrical, slightly compressed, about 2.0 times longer than maximal depth, at about proximal 0.4 of length with several transverse rows of short cleaning setae proximovertrally; finger subequal to palm length, slender, tapering, cutting edges sharp, entire, tips hooked, base slightly expanded, surrounded by palisade of short curved setae medially and laterally, carpus about 1.3 of chela length, 5.0 times longer than distal width, tapering proximally, with seven to eight serrulate cleaning setae distovertrally; merus about 1.4 times chela length, 5.4 times longer than central width; ichium 1.1 of palm length, 2.75 times longer than distal width, not strongly carinate distovertrally, obliquely articulated with basis; basis about 0.34 of carpus length; coxa without ventromedial process.

Second pereopods well developed, generally similar, unequal. Major (right) second pereopod exceeding carpocerite by chela and carpus; chela about 1.33 times carapace length, palm subcylindrical, subuniform, with small tubercles, some of the tubercles with short and fine setae on tip, about 3.4 times longer than maximal depth, fingers about 0.45 of palm length, with stout hooked tips and both with well-developed lateral flanges, lateral surfaces with small tubercles similar to those of palm; dactyl about 3.8 times longer than proximal depth, with developed lateral flange, 0.6 distal cutting edge entire, large triangular acute tooth at proximal quarter, notch present distal to and proximal to tooth, respectively; fixed finger stouter than dactyl, with well-developed lateral flange, cutting edge with large pit at base, large tooth with rounded tip present in pit, distal three-quarters cutting edge entire; carpus about 0.22 of palm length, 1.1 times longer than distal width, cup-like, feebly excavate distally, proximally tapered, armed with small tubercles similar to palm; merus about 0.5 of palm length, subuniform, about 3.65 times longer than distal depth, armed with sparse small tubercles, without distovertrally tooth; ischium about 0.5 of palm length, subequal to merus length, 3.66 times longer than distal width, slightly dorsoventrally compressed, slightly proximally tapered, armed with sparse small tubercles; basis and coxa normal. Minor (left) pereopod exceeding carpocerite by chela and distal two-thirds of carpus; chela about 0.81 times postorbital carapace length, 0.62 times length of major chela; palm about 3.5 times longer than maximal depth, with very sparse small tubercles; fingers about 0.5 of palm length, with strongly hooked tips, cutting edges entire, sharp, with

feebly dentition proximally; dactyl about 4.8 times longer than proximal depth, with feebly developed lateral flange; carpus cup-like, about 0.33 of palm length, 1.2 times the distal width, with scattered small tubercles; merus about 0.84 of palm length, subequal to ischium, with tubercles ventrally; basis and coxa normal.

Ambulatory pereopods robust. Third pereopod exceeds carpocerite by dactyl, propod, and distal quarter of carpus; dactyl compressed, curved, about 0.22 of propod length, unguis distinct, about 0.75 of dorsal length of corpus, corpus about 2.5 times longer than proximal depth, ventral margin unarmed, without distal accessory tooth, with two distolateral sensory setae; propod about 0.51 of carapace length, 6.8 times longer than maximum wide, slightly tapered distally, with pair of short slender simple distoventral spines, three single spines distributed on ventral margin, and transverse rows of long setae distolaterally; carpus about 0.52 of propod length, 3.65 times longer than distal width, unarmed; merus 1.08 of propod length, 6.5 times longer than wide, uniform, unarmed; ischium subequal to carpus length, 0.5 of merus length, 3.45 times longer than distal width, slightly tapered proximally; basis and coxa without special features. Fourth and fifth pereopods similar to third, fourth propod subequal, fifth propod 1.1 times third propod length, less strongly spinose distoventrally; fifth exceeds carpocerite by dactyl and distal three-fifths propod, with propod with transverse rows of cleaning setae distolaterally.

Uropod distinctly exceeding telson; protopodite with posterolateral angle short, rounded; exopod 2.4 times longer than central width, lateral border distinctly convex, with small acute distal tooth, with large mobile spine medially, diaeresis distinct; endopod about 0.9 of exopod length, 3.0 times longer than wide.

Paratype. Similar to holotype. Rostrum longer than that of the holotype, 0.77 of carapace length, but not reaching the distal end of scaphocerite, lower than that of holotype, dorsal margin slightly concave, with eight dorsal and four ventral teeth, interdental spaces and with distinct short setas. Telson regenerated and soft except the extreme base, the four dorsolateral spines distinct along right margin, obscure along left margin.

Measurements (mm)

Holotype. Female, carapace length, 5.81; carapace and rostrum, 10.19; second pereopod, major chela, 7.74; minor chela, 4.80.

Paratype. Female, carapace length, 5.68; carapace and rostrum, 10.03; second pereopod, major chela, 8.47; minor chela, 5.57.

Systematic position

In its general morphology, particularly the telson with four pairs of dorsolateral spines, major second pereopods with lateral flange on the fingers and palms covered with fine tubercles, and shape of the rostrum, *Periclimenes crosnieri* sp. n. shows a close resemblance to *P. aleator* Bruce, 1991 and other species of the “*P. alcocki* species group”. It can immediately be distinguished from these species by the presence of a simple non-biunguiculate dactyl on the ambulatory pereopods. It is similar to *P. uniunguiculatus* Bruce, 1990, *P. granuloides* Hayashi, 1986, *P. foresti* Bruce, 1981, or other deep-water species with a simple dactyl on the ambulatory pereopods. It can easily be distinguished from those species by the telson with four pairs of dorsolateral spines, and also the well-developed lateral flange on the major second pereopod.

Etymology

The specific name is given in honour of the French carcinologist, Alain Crosnier.

Remarks

Telson with four pairs of dorsolateral spines may show that *Periclimenes crosnieri* n. sp. belongs to the “*Periclimenes alcocki* species group”, but the simple dactyl on the ambulatory pereopods indicates that it is separated from the other members of the group. Captured at a depth of 302–305 m, occurs in shallower depth than *P. alcocki* and *P. aleator*, but at similar depth with *P. albatrossae* Chace and Bruce, 1993, which has seven pairs of dorsolateral spines on telson.

***Periclimenes forcipulatus* Bruce, 1991**

Periclimenes forcipulatus Bruce 1991a, p 330, Figures 21–25; Li 2000, p 181, Figure 227.

Material examined

Solomon Islands: SALOMON 1, stn DW 1854, 9°46.4'S, 160°52.9'E, 229–260 m, 7 October 2001, 1♀ (MNHN-Na 15536).

Remarks

Not previously recorded from Solomon Islands. The present record extends the bathymetric range to the shallower depth of 229 m.

Distribution

Type locality: Loyalty Islands, New Caledonia, 460 m. Now also known from Solomon Islands.

***Periclimenes forgesi* sp. nov.**
(Figures 19, 20)

Material examined

New Caledonia: (i) BATHUS 3, stn CP846, 23°02.90'S, 166°57.97'E, 500–514 m, 1 December 1993, 1 ovig. ♀ holotype (MNHN-Na 14928); (ii) NORFOLK 1, Norfolk Ridge, stn CP1670, 23°39'S, 167°59'E, 382–386 m, Banc Jumeau-ouest, 21 June 2001, 1 ovig. ♀ paratype (MNHN-Na 15869).

Description

A medium to large body size pontoniine shrimp of subcylindrical body form. Holotype with right scaphocerite, first, second, fourth and fifth pereopods, most of telson except proximal part anterior to anterior pair of dorsolateral spines, uropod with distal part posterior to diaeresis lost; right third pereopod and left fourth merus to dactyl detached; both hepatic spines and first three dorsal rostral teeth broken basally. Paratype with the tip of rostrum broken, right fourth, left fourth and fifth pereopods lost; dactyl of left (major) second pereopod lacking.




Figure 19. *Periclimenes forgesi*, new species, holotype ovigerous female (MNHN-Na 14928). (a) Carapace and anterior appendages; (b) anterior carapace; (c) eye; (d) antennule; (e) antenna; (f) mandible; (g) maxillula, upper lacinia and palp; (h, i) same, lower lacinia; (j) maxilla; (k) first maxilliped; (l) second maxilliped.

Holotype. Carapace smooth, glabrous. Rostrum short, slender, compressed, reaching to distodorsal end of second segment of antennular peduncle, about 0.42 of carapace length, feebly expanded laterally proximally, with feeble crest on proximal part and tapering distally; dorsal margin with seven small subequal acute teeth, more or less evenly distributed along whole length, second tooth situated just above posterior orbital margin; lateral carinae feebly developed; ventral margin straight, with one small acute tooth at distal 0.18 of rostral length, anterior to the distal dorsal tooth. Supraorbital spine absent, epigastric spine situated at anterior 0.21 of carapace length, distinctly smaller than dorsal rostral teeth; orbit feebly developed, inferior orbital angle produced, bluntly round; antennal spine small, slender, marginal, distinct below inferior orbital angle, not exceeding inferior orbital angle; hepatic spine feebly larger than antennal spine, small, slender, slightly below level of antennal spine, at level of first dorsal rostral tooth in lateral view; anterolateral angle of carapace not produced, bluntly rounded.

Abdominal segments smooth, glabrous; pleura of first three segments broadly rounded, fourth and fifth posteriorly produced, bluntly rounded; third segment without dorsal prominence; sixth segment about 1.7 times length of fifth, subcylindrical, about 1.75 times longer than deep, subuniform, posterolateral and posteroventral angles bluntly angulate.

Eye normally developed, cornea globular, well pigmented, diameter 0.16 of postorbital carapace length, without accessory pigment spot; stalk feebly compressed, feebly tapered proximally, 0.63 of corneal diameter and 0.74 of central width.




Figure 20. *Periclimenes forgesi*, new species, holotype ovigerous female (MNHN-Na 14928). (a) Third maxilliped; (b) first pereiopod; (c) same, chela; (d) second pereiopod; (e) same, fingers; (f) third pereiopod; (g, h) same, distal propod and dactyl; (i) uropod.

Antennular peduncle overreaching rostrum by third segment; proximal segment 1.85 times longer than central width, with slender acute stylocerite laterally, reaching to about 0.7 of segment length, anterolateral margin produced, setose, with long slender acute lateral tooth overreaching middle dorsal length of intermediate segment; statocyst small; medial margin straight, with plumose setose, ventromedial tooth minute, at about 0.5 of length. Intermediate segment obliquely articulated with distal segment, dorsal margin oblique proximolaterally, middle dorsal length about 0.26 of proximal segment length, 1.1 times width (not including lateral lobe), lateral margin expanded, with long robust plumose setae, medial margin similarly setose to lateral margin. Distal segment with mid-dorsal length 0.42 of proximal segment length, 1.4 times longer than distal width; upper flagellum biramous, with 10 proximal segments of rami fused, shorter free ramus with six segments, length about 0.64 of fused portion, total length about 0.53 of carapace length, with 22 groups of aesthetascs; longer ramus (free portion) slender, filiform, about 0.83 times carapace length; lower flagellum slender, filiform, subequal to carapace length.

Antennal basicerite robust, with acute distolateral tooth; carpocerite robust, 2.7 times longer than width, subcylindrical, reaching about 0.46 of total length of scaphocerite; scaphocerite well developed, exceeding antennular peduncle by distal third of lamella, broad, 2.5 times longer than wide, greatest width at about 0.4 of length, distal margin produced, bluntly rounded, distinctly overreaching distolateral tooth, lateral margin straight, with strong acute distolateral tooth.

Epistome unarmed. Fourth thoracic sternite without slender median process, with low transverse ridge with a median notch posteriorly; fifth thoracic sternite with similar ridge, posterior sternites unarmed.

Mouthparts typical of the genus. Mandible without palp; molar process normal, with four strong blunt teeth, lower inner tooth bilobed; incisor process robust, obliquely truncate distally with three (left) or four (right) stout teeth, central tooth smaller than outer teeth. Maxillula with palp bilobed, lower lobe shorter, acute tip with very short seta, upper lobe longer, with rounded tip; upper lacinia oval, with 10 spines distally and scattered setae; lower lacinia slender, tapered with numerous spiniform setae distally and simple setae on outer surface. Maxilla with short simple non-setose palp; basal endite deeply bilobed, upper lobe stouter than lower, with simple setae distally; coxal endite obsolete, medial margin convex; scaphognathite well developed, about 2.9 times longer than central width. First maxilliped with elongate slender palp, with short setulose preterminal seta; basal endite large, broad, setose medially; coxal endite distinct, convex; exopod with normal flagellum with four plumose distal setae and simple setae on distal half laterally; caridean lobe well developed, broad; epipod large, triangular, bilobed. Second maxilliped with normal endopod, dactylar segment about 3.5 times longer than broad, with numerous serrulate spines medially; propodal segment broad, with distal margin with long spiniform setae; carpus, ischiomerus and basis without special features; coxa angularly produced medially, with four to five simple setae; exopod with slender flagellum with eight to nine plumose setae distally; epipod simple, without podobrach. Third maxilliped with slender endopod, exceeding carpocerite by distal quarter of terminal segment; ischiomerus and basis incompletely fused, border indicated only by medial notch, basal portion feebly expanded medially, combined segment length about 6.0 times central width, compressed, setose medially, with submarginal row of very short small spines on proximal medial ischial portion; intermediate segment about 0.58 of combined proximal segment length, 6.0 times longer than central width, with six groups of long finely serrulate spiniform setae medially; terminal segment about 0.44 of combined proximal segment length, distally tapering, about 4.8 times longer than proximal width, with seven groups of spiniform setae medially, and long spiniform distal setae; exopod with slender flagellum with about 10 plumose setae distally; coxa feebly produced medially, with oval lateral plate; arthrobranch distinct.

First pereopods moderately slender, exceeding carpocerite by chela and distal five-sixths carpus; chela with palm subcylindrical, slightly compressed, about 2.25 times longer than depth, with six transverse rows of short cleaning setae proximally; finger subequal to palm length, slender, tapering, cutting edges sharp, entire, tips hooked, surrounded by four groups of palisade setae medially and laterally on distal half, setae shorter distally; carpus about 0.93 of chela length, 5.45 times longer than distal width, tapering proximally, with one row two groups serrulate cleaning setae distally, outer group with eight long setae, inner group with six short setae, about half as long as the long setae; merus about 1.2 times chela length, 7.0 times longer than central width; ichium 0.6 of chela length, 3.3 times longer than distal width, not strongly carinate distally, obliquely articulated with basis; basis and coxa without special features.

Left (minor) second pereopod well developed, exceeding carpocerite by chela, carpus and distal third of merus, rostrum by chela and carpus; chela about 0.95 times carapace length, palm subcylindrical, subuniform, with sparse small tubercles on medial (fixed finger side) surface, about 4.8 times longer than maximal depth, fingers about 0.6 of palm length, with stout hooked tips and feebly developed lateral flanges, fixed finger stouter than dactyl, with few scattered small tubercles on lateral surfaces similar to those of palm, cutting edge

with distal 0.72 entire, small tooth on proximal 0.23 and small blunt rounded tooth in proximal pit; dactyl about 3.6 times longer than proximal depth, cutting edge with distal 0.64 entire, two small teeth and two notches on proximal 0.36, proximal tooth larger and triangular; carpus cup-like, about 0.34 of palm length, 1.76 times longer than distal width, feebly excavate distally, proximally tapered; merus about 0.83 of palm length, subuniform, about 5.5 times longer than distal depth, armed with sparse small tubercles on flexor margin, without distoventral tooth; ischium about 0.8 of palm length, subequal to merus length, 4.7 times longer than distal width, feebly compressed, slightly proximally tapered; basis and coxa normal.

Ambulatory pereopods robust. Third pereopod exceeds carpocerite by dactyl, propod, and distal three-quarters of carpus; dactyl compressed, about 0.19 of propod length, unguis distinct, about 0.4 of dorsal length of corpus, corpus about 2.5 times longer than proximal depth, with one distolateral sensory seta, ventral margin feebly concave, with acute distal accessory tooth, distal accessory tooth reaching at about proximal 0.4 of unguis; propod about 0.64 of carapace length, 10.5 times longer than maximum wide, slightly tapered distally, with one pair of slender simple distoventral spines, one pair of similar subdistoventral spines, three single spines distributed on ventral margin, one small transverse row of long setae distolaterally and scattered setae on distodorsal and ventral margin; carpus about 0.47 of propod length, 4.2 times longer than distal width, unarmed; merus 0.94 of propod length, 8.85 times longer than wide, uniform, unarmed; ischium 0.42 of propod length, 3.62 times longer than distal width, slightly proximally tapered; basis and coxa without special features. Fourth and fifth pereopods similar to third, fourth propod subequal, fifth propod 1.1 times third propod length, fifth exceeds carpocerite by dactyl and distal 9/10 propod.

Uropod with protopodite with posterolateral angle short, rounded; exopod 2.5 times longer than central width, lateral border feebly convex, with small acute distal tooth, mobile spine medially lost, diaeresis distinct; endopod subequal to exopod length, 3.3 times longer than wide.

Paratype. Similar to holotype. Rostrum with six dorsal teeth. Telson slender, 1.69 times sixth abdominal somite, 4.1 times proximal width, with four pairs of dorsolateral spines at 0.41, 0.58, 0.72 and 0.88 of telson length, right side with one additional spine at 0.95 of telson length, the spines about 0.056 of telson length; lateral margins straight, tapered distally, posterior margin 0.4 of proximal width, with three pairs of spines, lateral spines similar to dorsolateral spines, intermediate spines long and robust, about 0.12 of telson length, submedian spines slender, non-setae, about 0.36 of intermediate spine length. Uropod exceeding telson tip.

Measurements (mm)

Holotype. Carapace length, 7.3; carapace and rostrum, 10.3; total animal length (proximally), 29.5; left (minor) second pereopod chela, 7.4; ovum length, 0.73.

Paratype. Carapace length, 6.8; total animal length (proximally), 28.6; left (major) second pereopod chela, 10.4; right (minor) second pereopod chela, 6.1.

Systematic position

In its general morphology, especially the raised proximal dorsal rostral carina, *Periclimenes forgesi* sp. n. shows a close resemblance to *P. vaubani* Bruce, 1990a. *Periclimenes forgesi* may be distinguished from that species easily by the four pairs of dorsolateral telson spines, and also the following features: (1) rostrum with deeper lamina; (2) carapace without epigastric tubercle; (3) hepatic spine on carapace subequal to antennal spine; (4) ambulatory propods lacking transverse rows of setae ventrally except the small row distoventrally; (5) scaphocerite with distal lamella distinctly exceeding distolateral tooth; and (6) first pereiopod with carpus shorter than chela. *Periclimenes forgesi* differs from the other deep-water *Periclimenes* species with four pairs of dorsolateral telson spines by the short and slender rostrum with a distinct proximal elevated dorsal margin over orbital region, ventral margin straight, with one tooth, cornea well developed, about 0.16 of the postorbital margin carapace length, hepatic not noticeably larger than antennal spine, ambulatory propods lacking transverse rows of setae on ventral margin.

Etymology

The specific name is given in honour of the specimen collector, Dr Bertrand Richer de Forges.

Remarks

The small hepatic spines on both sides of the holotype are broken basally, appearing as if articulated on the carapace, making the species appear similar to *Allopontonia iaini* Bruce, 1972. But, the armature of ambulatory pereiopods is not as in *Allopontonia iaini*, in which the ventral border of ambulatory propods are strongly spinulate, the ventral border of dactyl is denticulate, not armed with only one accessory tooth. In addition, *Allopontonia* is an echinoid-associated genus in shallow water.

Periclimenes foveolatus Bruce, 1981

Periclimenes foveolatus Bruce 1981a, p 196–201, Figures 6–9, 17a, b, 18b, e; Bruce 1991a, p 312, Figures 8, 72; Chace and Bruce 1993, p 111; Li 2000, p 182, Figure 229.

Material examined

Philippines: MUSORSTOM 3, stn CP91, 14°00'N, 120°18'E, 190–203 m, 31 May 1985, 1♂ (MNHN-Na 14892).

Remarks

Previously known only from the type material. The single specimen has a rostral dentition of 9/4 and lacks any trace of an epigastric spine or tubercle. The cornea is moderately large. The minor second pereiopod only is preserved, the palm of the chela is feebly, minutely tuberculate ventrally. The third pereiopod has the dactyl with a small, slender accessory tooth and the telson has two pairs of medium-sized dorsal spines.

Distribution

Type locality: SW of Manila Bay, Philippines, 191–188 m. Also recorded from Loyalty Islands.

Periclimenes hertwigi Balss, 1913

(Figure 21)

Periclimenes hertwigi Balss 1913, p 235; Bruce 1990a, p 151, Figures 1–2, 39c; Chace and Bruce 1993, p 113; Li 2000, p 189, Figure 239; Davie 2002, p 327.

Periclimenes Hertwigi: Balss 1914, p 49–51, Figures 28–30.

Periclimenes (Ancylocaris) gracilirostris Kubo 1940b, p 41–44, Figures 8–10.

Periclimenes (Periclimenes) hertwigi: Holthuis 1952a, p 43–47, Figures 11, 12.

Material examined

Philippines: MUSORSTOM 3, (i) stn CP90, 14°00'N, 120°19'E, 195 m, 31 May 1985, 1 ovig. ♀ (MNHN-Na 14898). **Indonesia:** KARUBAR, (ii) Tanimbar Islands, stn CP45, 7°54'S, 132°47'E, 302–305 m, 29 October 1991, 1 ♂ (MNHN-Na 14895); (iii) Kai Islands, stn CP27, 5°33'S, 132°51'E, 304–314 m, 26 October 1991, 1 ♂ (MNHN-Na 15879). **Solomon Islands:** SALOMON 1, (iv) stn CP1792, 9°15.4'S, 160°08.9'E, 477–505 m, 30 September 2001, 1 ♀ (MNHN-Na 15538); (v) stn CP1831, 10°12.1'S, 161°19.2'E, 135–325 m, 5 October 2001, 1 ovig. ♀ (MNHN-Na 15537). SALOMON 2, (vi) Rendova, stn CP2284, 8°38.4'S, 157°21.5'E, 195–197 m, November 2004, 1 ovig. ♀ (MNHN-Na 15947); (vii) northwest Isabel, stn CP2210, 7°34.2'S, 157°41.8'E, 240–305 m, November 2004, 1 ovig. ♀ (MNHN-Na 15948); (viii) northwest Isabel, stn CP2199, 7°43.1'S, 158°29.6'E, 296–304 m, November 2004, 2 ovig. ♀♀ (MNHN-Na 15957); (ix) Vella Gulf, stn CP2260, 8°03.5'S, 156°54.5'E, 399–427 m, November 2004, 1 ovig. ♀ (MNHN-Na 15943). **New Caledonia:** BIOCAL, (x) stn CP45, 22°47'S, 167°15'E, 430–465 m, 30 August 1985, 1 ♂, 1 ovig. ♀ (MNHN-Na 14901). MUSORSTOM 4, (xi) stn 156, 18°54.0'S, 163°18.8'E, 525 m, 15 September 1985, 1 ovig. ♀ (MNHN-Na 14904); (xii) stn 169, 18°54.3'S, 163°11.2'E, 390 m, 17 September 1985, 1 juvenile ♂ (MNHN-Na 14903); (xiii) stn 194, 18°52.8'S, 163°21.7'E, 545 m, 19 September 1985, 1 ♂, 1 ovig. ♀ (MNHN-Na 14899); (xiv) stn 221, 22°58.6'S, 167°36.8'E, 535–560 m, 29 September 1985, 2 ♂♂ (MNHN-Na 14902); (xv) stn 236, 2°11.3'S, 163°15.0'E, 495–550 m, 2 October 1985, 1 ♀ (MNHN-Na 14900). HALIPRO 1, (xvi) stn CP866, 21°26.91'S, 166°17.23'E, 550–600 m, 22 March 1993, 1 ♂ (MNHN-Na 14897). BATHUS 2, (xvii) stn CP770, 22°09.56'S, 166°04.09'E, 400–402 m, 18 May 1993, 1 ovig. ♀ (MNHN-Na 14894). BATHUS 3, (xviii) stn CP813, 23°45.23'S, 168°16.57'E, 410–415 m, 28 November 1993, 1 ovig. ♀ (MNHN-Na 14896). NORFOLK 1, (xix) Norfolk Ridge, stn CP1708, 23°43'S, 168°16'E, 381–384 m, Banc Jumeau-est, 25 June 2001, 1 ♂ (MNHN-Na 15866).

Parasites

Specimen (xv) had a branchial parasite, *Bopyrinina paucimaculata* Markham [Isopoda: Bopyridae].




Figure 21. *Periclimenes hertwigi* Balss, 1913. (a, h) Ovigerous female (MNHN-Na 14899); (b, e, g, i) male (MNHN-Na 14899); (c, d, f) ovigerous female (MNHN-Na 14896); j, k, male (MNHN-Na 14901). (a) Anterior carapace; (b) rostrum; (c) distal antennal scale; (d-f) first pereiopod, chela, lateral view; (g, h) second pereiopod, chela; (i) same, fingers; (j) third pereiopod, distal propod and dactyl; (k) same, distoventral angle of corpus.

Remarks

Not previously reported from the Philippines or Indonesia. The Philippines specimen (i) has the rostrum with three ventral rostral teeth.

Distribution

Type locality: Sagami Bay, Japan, 120 m. Now known from East China Sea, Japan, Philippines, Indonesia, Australia (Queensland), and New Caledonia.

Periclimentes imperator Bruce, 1967

Periclimentes imperator Bruce 1967a, p 53–62, Figures 23–25; Bruce 1991b, p 237; Bruce 1996, p 234; Li 2000, p 192, Figure 243; Davie 2002, p 327; Li and Liu 2004, p 95, Figure 6.

Material examined

Madagascar: (i) Pracel Bank, west coast, 55 m, sand, trawl, coll. A. Crosnier, June 1959, 1♂ (MNHN-Na 14909); (ii) Nosy Be, stn 10, 26 m, coll. P. Laboute, 9 September 1992, phot. CD03, 1♂, 1 ovig. ♀ (MNHN-Na 14939). **New Caledonia:** (iii) scuba stn 535, coll. J.-L. Menou, no date record, 1♂, 1♀ (MNHN-Na 14907); (iv) MONTROUZIER, Koumac: Ilot Tangdion, dredge, coll. B. Richer de Forges, 8 October 1993, 1 ovig. ♀ (MNHN-Na 14908).

Hosts

(ii) *Melibe* sp.; (iii) *Thelonota ananas* (Jaeger) [Holothuroidea, Echinoderma].v

Remarks

First reported from Nouméa (New Caledonia) by Bruce (1967b) and subsequently also in New Caledonia from Nouméa, Île Ouen, Baie du Prony, Île des Pins, and Lagoon Nord Ouest by Bruce (1991b). Previously reported from Madagascar, from Baie d'Ambaro (Bruce 1977c) and also from Mitsio Island, by Bruce (1978a). The specimen from Pracel Bank at the depth of 55 m represents a new bathymetric record for this species, previously known only to 30 m (Bruce 1991b). Unfortunately this identification cannot be considered absolutely certain as the specimen lacks a rostrum.

Distribution

Type locality: Chumbe Island, Zanzibar. Also known from Egypt, Israel, Jordan, Kenya, Zanzibar, Mozambique, Comoro Islands, Madagascar, Seychelles, Japan, Indonesia, Australia (Western Australia, Queensland), Caroline Islands, New Caledonia, and Hawaii.

Periclimenes incertus Borradaile, 1915

Periclimenes (Cristiger) incertus Borradaile 1915, p 210; Borradaile 1917, p 364, Plate 53, Figure 7.

Periclimenes (Periclimenes) impar Kemp 1922, p 147–149, Figures 16, 17, Plate 3, Figure 1.

Periclimenes (Periclimenes) incertus: Holthuis 1959, p 193–194.

Periclimenes incertus: Chace and Bruce 1993, p 114; Bruce 1996, p 234; Li 2000, p 193, Figure 244; Davie 2002, p 327.

Material examined

Madagascar: (i) northwest coast, S. W. Îles Mitsio, 30 m, sand, trawl, coll. A. Crosnier, February 1960, 3 ovig. ♀♀ (MNHN-Na 14910); (ii) northwest coast, Tany Kely, Nosy Be' 20 m, on sponge, coll. P. Laboute, 19 July 1970, 41 spms (8 ovig. ♀♀) (MNHN-Na 14911). **Solomon Islands:** (iii) SALOMON 1, stn CP 1810, 9°47.7'S, 160°50.5'E, 53 m, 3 October 2001, 1 ovig. ♀ (MNHN-Na 15586). **New Caledonia:** (iv) lagoon, Îlot Maitre, scuba, 25 m, with sponge, coll. C. Vadon, 5 September 1978, 3♀♀ (2 ovig.) (MNHN-Na 15836); (v) same, 2♂♂, 9 ovig. ♀♀ (MNHN-Na 15899); (vi) lagoon, channel of Îlot Maitre, scuba, 25 m, with sponge, coll. C. Vadon, 5 September 1978, 1♂, 8 ovig. ♀♀ (MNHN-Na 15861); (vii) lagoon, Five Mile Channel, scuba, 15–20 m, associated with alcyonarian *Nephtia* sp., coll. C. Vadon, 13 September 1978, 1♂, 4 ovig. ♀♀ (MNHN-Na 15816); (viii) lagoon, channel of Îlot Maitre, stn 21, scuba, 25 m, with sponge, coll. C. Vadon, 19 September 1978, 3♂♂, 6 ovig. ♀♀ (MNHN-Na 15855); (ix) lagoon, channel of Îlot Maitre, scuba, 25 m, with sponge, coll. C. Vadon, 19 September 1978, 1 ovig. ♀ (MNHN-Na 15835); (x) same, 4♂♂, 8♀♀ (4 ovig.) (MNHN-Na 15845); (xi) lagoon, Seiche croissant Reef, scuba, 9 m, with coral, coll. C. Vadon, 29 September 1978, 1♂, 1 ovig. ♀ (MNHN-Na 15900); (xii) lagoon, coll. C. Vadon, 4 October 1978, 6 ovig. ♀♀ (MNHN-Na 15844); (xiii) lagoon, southwest, 22°19.41'S, 166°20.89'E, 20 m, scuba, with sponge, coll. I. Takeuchi, 9 November 1995, 1♀, 1 spm (abdomen lost) (MNHN-Na 15907); (xiv) Îlot Maitre, 22°19.35'S, 166°25.85'E, 20 m, scuba, with sponge, coll. I. Takeuchi, 10 November 1995, 17♂♂, 13♀♀ (2 ovig.) (MNHN-Na 15820); (xv) Îlot Maitre, stn 82, 22°19.61'S, 166°24.07'E, 10 m, coll. I. Takeuchi, 14 November 1995, 33 spms (13 ovig. ♀♀) (MNHN-Na 15877); (xvi) lagoon (Seiche Croissant), seagrass, small trawl, coll. Richer de Forges, 6 May 1998, 4♂♂, 1♀ (with a bopyrid), 12 ovig. ♀♀ (MNHN-Na 15806).

Hosts

(ii) *Rhizochalina* sp.; (viii) ?*Siphonochalina* sp.; (xii) *Clathrya astroderma* [Porifera]. (vii) *Nephtia* sp. [Alcyonaria].

Remarks

Previously reported from Madagascar, from Baie d'Ambaro (Bruce 1977c) and from Tany Kely (Bruce 1978a). The Solomon Islands record represents the greatest depth from which this species has so far been reported. The Tany Kely specimens are largely juveniles: the second pereopods, in those specimens in which they are still both attached, are distinctly unequal. The Ampombilava specimen lacks both second pereopods but can be safely identified by the characteristic ambulatory dactyl. The association with *Rhizochalina*

represents a new host record. *Periclimenes incertus* has been frequently noted as an associate of sponges. Previously reported hosts include species of the genera *Acarinus*, *Clathria*, *Desmopsamma*, *Dysidea*, *Oceanapta*, *Petrosia*, *Siphonochalina*, and *Spirastrella* (Bruce 1976b); *Arenochalina* and *Leucetta* (Bruce 1983d); and *Callyspongia* (Bruce 1988c).

Distribution

Type locality: South Nilandu Atoll, Maldives. Also known from Yemen, Kenya, Zanzibar, Tanzania, Madagascar, Sri Lanka, Andaman Islands, Singapore, Philippines, Indonesia, Australia (Western Australia, Northern Territory, Queensland), and New Caledonia.

Periclimenes involens Bruce, 1996

(Figure 22)

Periclimenes involens Bruce 1996, p 234, Figures 13, 28h; Li 2000, p 198, Figure 251.

Material examined

La Réunion: MD 32, stn CP55, 21°05.3'S, 55°12.5'E, 97–110 m, 22 August 1982, 1 ovig. ♀ (MNHN-Na 14912).

Remarks

Previously known only from the type specimens from the Philippines. The present specimen agrees closely with the original description and is in good condition with intact rostrum and both second pereopods. The specimen has a carapace length of 2.3 mm, the rostrum reaching the end of the antennular peduncle, with a dentition of 1+7/2, the first tooth situated slightly posterior to the level of the orbital margin. The specimen lacks the elevated posterior dorsal rostral carina found in *P. vaubani* and the ventral teeth are distinctly larger. The ventral border of the rostrum is straight, and not convex as in *P. richeri*. The second pereopods are similar, unequal, both dactyls with a dorsal flange, the major carpus is about 0.3 of the palm length and the minor about 0.5. This new record indicates a large increase in the range of this small species. The only previous record was from a similar depth, 92–97 m.



Figure 22. *Periclimenes involens* Bruce, 1996, ovigerous female (MNHN-Na 14912), carapace.

Distribution

Type locality: 12°31'N, 120°39'E, off Mindoro, Philippines. Now also known from La Réunion.

Periclimenes kemp Bruce, 1969

Periclimenes kemp Bruce 1969b, p 260; Bruce 1981b, p 80, Figure 2; Fransen 1994, p 125; Bruce and Coombes 1995, p 131; De Grave 2000, p 136; Li 2000, p 200, Figure 256.

Material examined

New Caledonia: lagoon, Five Mile Channel, scuba, 15–20 m, associated with alcyonarian *Nephtya* sp., coll. C. Vadon, 13 September 1978, 2♂♂, 1♀ (MNHN-Na 15823).

Host

Nephtya sp. [Alcyonacea].

Distribution

Type locality: Red Sea coast of Egypt, 1 m. Known also from Kenya, Zanzibar, Seychelles, Andaman Islands, Singapore, Papua New Guinea, Australia (Cobourg Peninsula, Great Barrier Reef), New Caledonia, and Fiji.

Periclimenes laccadivensis (Alcock and Anderson, 1884)

Palaemonella laccadivensis Alcock and Anderson 1894, p 157.

Periclimenes laccadivensis: Kemp 1922, p 152, Figure 19; Bruce 1979a, p 225; Bruce 1991a, p 301, Figure 1; Bruce 1992, p 69, Figure 20; Li 2000, p 202, Figure 259.

Material examined

New Caledonia: (i) LITHIST, stn DW05, 23°38.2'S, 167°42.9'E, 433–500 m, Stylaster Bank, 10 August 1999, 1 ovig. ♀ (MNHN-Na 15881); (ii) NORFOLK 1, Norfolk Ridge, Introuvable Bank, stn DW1697, 24°39'S, 168°38'E, 569–616 m, coll. Lozouet, Boisselier, Richer-IRD, 24 June 2001, 1 ovig. ♀ (MNHN-Na 15930).

Remarks

Previously recorded from New Caledonia by Bruce (1991a).

Distribution

Type locality: Laccadive Sea, 770–1353 m. Known also from western Indian Ocean, South China Sea, New Caledonia, and off Australia (Tasmania).

***Periclimenes lanipes* Kemp, 1922**

(Figure 23)

Periclimenes (Periclimenes) lanipes Kemp 1922, p 156–158, Plate 4, Figure 4.

Periclimenes lanipes: Bruce 1991b, p 237; Chace and Bruce 1993, p 116; Li 2000, p 203, Figure 261; Davie 2002, p 329.

Material examined

Philippines: (i) Maribago, Mactan Island, Cebu, washings, coll. P. Bouchet, 9 June 1985, 1♂, 5 ♀♀ (3 ovig.), 17 juveniles (MNHN-Na 14913). **Fiji:** (ii) southeast Viti Levu, stn CP 1358, 17°48.5'S, 178°46.7'E, 80–120 m, coll. P. Bouchet, B. Richer de Forges, 13 August 1998, 1♂ (MNHN-Na 15587).

Hosts

(i) *Euryale aspera* Lamarck, 1816 and *Astroboa nuda* (Lyman, 1874), det. A. Gilles [Ophiuroidea, Echinoderma].




Figure 23. *Periclimenes lanipes* Kemp, 1922. a, c, ovigerous female; (b) male (MNHN-Na 14913). (a, b) Carapace; (c) eye.

Remarks

Previously reported from Jolo Island and Siasi, Sulu Islands from Philippines (Chace and Bruce 1993). Not previously reported from Fiji.

Distribution

Type locality: 12°48'N, 98°16'10"E, Mergui Islands, Burma, 44 m. Also known from Somalia, Kenya, Zanzibar, Mozambique, Madagascar, Singapore, South China Sea, Philippines, Australia (Queensland), New Caledonia, and Fiji.

Periclimenes latipollex Kemp, 1922

Periclimenes (*Periclimenes*) *latipollex* Kemp 1922, p 150, Figure 18, Plate 4, Figure 3; Holthuis 1952a, p 47–48, Figures 13, 14.

Periclimenes latipollex: Bruce 1971a, p 8; Bruce 1981a, p 195, Figure 3; Bruce 1991a, p 309, Figure 6; Li 2000, p 204, Figure 262.

Material examined

New Caledonia: NORFOLK 2, Norfolk Ridge, Jumeau Est Bank, stn DW 2108, 23°47'S, 168°17'W, 403–440 m, 31 October 2003, 1 ovig. ♀ (MNHN-Na 15937).

Remarks

The dactyl of ambulatory pereopods with accessory spine is minute, as shown by Bruce (1991a, p 309, Figure 6f), but the rostrum is very slender, rod-like as shown by Kemp (1922) and Holthuis (1952a), with rostral formula 2+6/2. Previously recorded from New Caledonia by Bruce (1991a). The present record extends its bathymetric range from 73–304 m to 440 m.

Distribution

Type locality: Mergui Archipelago. Also known from Somalia, Kenya, Philippines, Indonesia, and New Caledonia.

Periclimenes lepidus Bruce, 1978

Periclimenes lepidus Bruce 1978a, p 244–252, Figures 20–24; Li 2000, p 205, Figure 264.

Material examined

Comoro Islands: (i) BENTHEDI, north Pamanzi Island, stn 32R, 12°45.1'S, 45°17.9'E, 15–20 m, slope with corals going to a sandy area with Heterocongridae, 1 ovig. ♀ (MNHN-Na 15942). **Madagascar:** (ii) northwest coast, near Nosy Be, Entrée Bank of SE slope, 40 m, coll. P. Laboute, 10 September 1970, 1♂, 6♀♀ (3 ovig.) (MNHN-Na 14915). **New Caledonia:** (iii) MONTROUZIER, Touho Channel, 35 m, from antipatharians, coll. B. Richer de Forges, 31 August 1993, 9 spms (7 ovig. ♀♀) (MNHN-Na 14914).

Remarks

Not previously recorded from Comoro Islands and New Caledonia. Previously recorded from Madagascar, from Nosy Be (Bruce 1978a), so the Madagascar specimens are topotypic.

Distribution

Type locality: Nosy Be, Madagascar, 40 m. Also known from Kenya, Seychelles, and Maldives.

Periclimeses loyautensis sp. nov.

(Figure 24)

Material examined

Loyalty Islands: MUSORSTOM 6, stn CP464, 21°02.30'S, 167°31.60'E, 430 m, coll. B. Richer de Forges, 21 February 1989, 1 ovig. ♀ holotype (MNHN-Na 14947).

Description

A medium size, robust pontonine shrimp of subcylindrical body form; lacking right second (major ?), fifth and left third pereiopods, left second (minor ?) pereiopod detached.

Carapace smooth, glabrous. Rostrum well developed, moderately deep, compressed, extending well beyond antennular peduncle, about 0.66 of carapace length, not reaching distal scaphocerite, anteroventrad except the anterad tip; dorsal carina deep, with seven




Figure 24. *Periclimeses loyautensis*, new species, holotype ovigerous female (MNHN-Na 14947). (a) Carapace; (b) telson; (c) same, distal part; (d) eye, distal view; (e) minor second pereiopod; (f) same, fingers; (g) third pereiopod, propod and dactyl; (h) same, distal propod and dactyl.

subequal acute teeth along all length, proximal three more closely grouped than distal teeth, first tooth situated just posterior to posterior orbital margin; lateral carinae feebly developed, slightly upcurved distally and extending posteriorly on carapace posterodorsally; ventral carina feebly developed, with three low acute teeth on distal half margin, smaller than dorsal teeth; interdental spaces and proximal ventral carina feebly setose. Supraorbital spine absent; epigastric spine situated at anterior 0.3 of the carapace length, with basal suture; orbit feebly developed, inferior orbital angle strongly produced, round; antennal spine small slender marginal, distinct below the inferior orbital angle, not exceeding inferior orbital angle; hepatic spine larger than antennal one, slender, very slightly lower than, almost at same level of antennal spine, below between the epigastric spine and the first dorsal rostral tooth and closer to first dorsal rostral spine in lateral view; anterolateral angle of carapace not produced, bluntly rounded.

Abdominal segments smooth, glabrous; third abdominal segment not produced, non-carinate; pleura of first three segments broadly rounded, fourth and fifth posteriorly produced rounded, sixth segment about 1.8 times length of fifth, subcylindrical, about 1.73 times longer than deep, subuniform, posterolateral angle acute, posteroventral angle rounded. Telson about 1.53 times sixth segment length, about 3.7 times longer than anterior width, lateral margins with posterior two-thirds straight, convergent, posterior margin about 0.32 of anterior margin width, rounded, without posterior median point; dorsal surface with four pairs of small but well-developed dorsolateral spines at 0.4, 0.6, 0.75, and 0.9 of telson length, spines about 0.044 of telson length; posterior margin with three pairs of spines, lateral spines short, similar to dorsolateral spines, intermediate spines long, robust, about 0.12 of telson length, submedian spines slender, about 0.41 of intermediate spine length.

Eye small, with globular cornea, corneal diameter 0.13 of posterior orbital carapace length, without accessory pigment spot; stalk subequal to corneal diameter, subequal to corneal width, feebly compressed.

Antennular peduncle overreaching distal ventral rostral tooth; proximal segment about 2.0 times longer than central width, with slender acute stylocerite laterally, reaching to about 0.7 of segment length, anterolateral margin feebly produced, setose, with long slender acute lateral tooth reaching near to the end of dorsal margin of intermediate segment; statocyst small; medial margin setose, with ventromedial tooth at about 0.5 of length; intermediate segment with dorsal length about 0.3 of proximal segment length, 1.1 of width, lateral margin expanded, setose; distal segment about 0.43 times proximal segment length, slender, about 2.0 times longer than distal width; upper flagellum biramous, with 10 proximal segments of rami fused, shorter free ramus with five segments, length about 0.5 of fused portion, total length about 0.54 of carapace length, with about 23 groups of aesthetascs; longer ramus, slender, filiform, distal segments lost; lower flagellum slender, filiform, about 0.87 times carapace length.

Antennal basicerite robust, with acute distolateral tooth; carpocerite about 0.49 of length of lateral margin of scaphocerite, about 3.2 times longer than width, subcylindrical, reaching about 0.41 of total length of scaphocerite; flagellum well developed, slender, about 2.2 times postorbital carapace length; scaphocerite well developed, distinctly exceeding tip of rostrum, broad, about 2.5 times longer than wide, greatest width at about 0.4 of length, distal margin bluntly rounded, distinctly exceeding distolateral tooth, lateral margin nearly straight, with strong acute distolateral tooth.

Epistome unarmed. Fourth thoracic sternite without slender median process; posterior sternites unarmed.

Mouthparts mainly undissected. Mandible without palp; incisor process distally with three (right) or four (left) stout acute teeth, central teeth smaller than outer teeth. Third maxilliped with slender endopod, extending distally to three-quarters length of carpocerite, ischiomerus and basis incompletely fused, basal portion medially expanded, convex, combined segment compressed, setose medially; intermediate segment about 0.43 of combined proximal segment length, with several groups of long finely serrulate spiniform setae medially; terminal segment about 0.33 of combined proximal segment length, distally tapering, with seven transverse groups of spiniform setae ventromedially, with long distal spiniform setae; exopod with slender flagellum extending to the distal end of ischiomerus, with eight to nine plumose setae distally; arthrobranch distinct.

First pereopods moderately slender, exceeding tip of rostrum by length of chela; chela with palm subcylindrical, slightly compressed, about 2.33 times longer than maximal depth, at about proximal 0.3 of length with several transverse rows of short cleaning setae proximoventrally; fingers 0.83 of palm length, slender, tapering, cutting edges sharp, entire, tips hooked, base slightly expanded, surrounded by palisade of long setae medially and laterally at about 0.3, 0.7 and subterminally of fingers; carpus about 1.22 of chela length, 5.4 times longer than distal width, subuniform, with seven to eight serrulate cleaning subdistal setae ventrally; merus about 1.4 times chela length, 7.0 times longer than distal width; ichium 0.78 of chela length, 4.5 times longer than distal width, not strongly carinate distoventrally, obliquely articulated with basis.

Left (minor) second pereopod well developed, exceeding carpocerite by carpus and chela; chela about 0.93 times postorbital carapace length; palm about 4.0 times longer than maximal depth, with very small tubercles; fingers about 0.7 of palm length, with strongly hooked tips, cutting edges entire, sharp, proximal 0.3 of fixed finger cutting edge with two teeth, proximal one larger and round, distal one smaller and acute, proximal 0.4 of dactyl cutting edge with two acute teeth, proximal one larger than distal; dactyl about 5.3 times longer than proximal depth, with feebly developed lateral flange; carpus cup-like, about 0.43 of palm length, 1.87 times distal width, with scattered small tubercles; merus about 0.9 of palm length, subequal to ischium, with tubercles ventrally; basis and coxa normal.

Ambulatory pereopods slender. Third pereopod exceeds carpocerite by dactyl, propod, and distal two-thirds of carpus; dactyl compressed, curved, about 0.18 of propod length, unguis distinct, about 0.61 of dorsal length of corpus, 3.7 times longer than proximal width; corpus about 2.5 times longer than proximal depth, dorsal margin feebly convex, non-setate, ventral margin straight, with acute distal accessory tooth, about 0.29 of unguis length, with two distolateral sensory setae; propod about 0.58 of carapace length, 13 times longer than maximum width, subuniform, with slender simple distoventral spine and five single spines distributed along ventral margin, and scattered long simple setae distally; carpus about 0.53 of propod length, 6.1 times longer than distal width, with distodorsal lobe, unarmed; merus subequal to propod length, 9.2 times longer than central width, uniform, unarmed; ischium subequal to carpus length, 0.52 of propod length, 4.9 times longer than distal width, slightly tapered proximally; basis and coxa without special features. Fourth and fifth pereopods similar to third, fourth propod subequal, fifth propod 1.1 times third propod length; fifth exceeds carpocerite by dactyl and almost all propod.

Pleopods without special features.

Uropod distinctly exceeding distal end of intermediate posterior spine of telson; protopodite with posterolateral angle short, rounded; exopod 2.7 times longer than central width, lateral border distinctly convex, with small acute distal tooth, with large mobile spine

medially, diaeresis distinct; endopod about 0.97 of exopod length, 3.6 times longer than wide.

Ova small, numerous, more than 400.

Measurements (mm)

Carapace length, 5.81; carapace and rostrum, 9.86; total body length (approximately), 26.6; second pereiopod chela, 5.46; length of ovum, 0.45.

Systematic position

Periclimenes loyautensis is very close to *P. aleator* Bruce, 1991 and other species of “*P. alcocki* species group” which have more than two (usually four) pairs of dorsolateral telson spines, second pereiopods with lateral flange and covered with fine tubercles. It can be distinguished from *P. aleator* by the hepatic spine very slightly below the level of the antennal spine, at almost the same level, also the relatively more slender ambulatory pereiopods and longer uropods which overreach the distal end of intermediate posterior spine of telson. In *P. aleator*, the hepatic spine is distinctly lower than antennal spine, the ambulatory pereiopods are relatively robust, uropods reach to about middle of intermediate posterior spine of telson. *Periclimenes loyautensis* differs from *P. alcocki* Kemp, 1992, by the larger cornea, the diameter is 0.13 of carapace length, and larger dorsolateral telson spines, about 0.044 of telson length, and the third pereiopod without transverse rows of setae distally, propod with well-developed spinulation, dactyl with longer and more slender accessory tooth. In latter species, the corneal diameter is about 0.1 or less of carapace length, the dorsolateral telson spines minute, and the posterior telson spines are also much shorter. *Periclimenes loyautensis* is also similar to *P. crosnieri* n. sp., the biunguiculate dactyl of ambulatory pereiopods easily distinguish it from that species, in which the ambulatory dactyl is simple.

Etymology

The specific name is given from its type locality, the Loyalty Islands.

***Periclimenes macrophthalmus* Fujino and Miyake, 1970**

(Figure 25)

Periclimenes (Harpilius) macrophthalmus Fujino and Miyake 1970, p 250, Figures 3–5.

Periclimenes macrophthalmus: Li 2000, p 211, Figure 272.

Material examined

Fiji: BORDAU 1, stn CP 1445, 350–365 m, 17°10'S, 178°42'W, 3 March 1999, 1 ovig. ♀ (MNHN-Na 15620).

Remarks

The single female accords well with the original description, although it is different in the following aspects: the proximal part of rostrum looks deeper than that of the holotype; the first tooth (epigastric spine) of the dorsal rostral series teeth is similar to others in size, not




Figure 25. *Perichlimes macrophthalmus* Fujino and Miyake, 1970, ovigerous female (MNHN-Na 14620). (a) Carapace and anterior appendages; (b) right second pereiopod; (c) fingers of left pereiopod. Scale bars: 1.41 mm (a); 2.24 mm (b); 0.5 mm (c).

distinctly stronger than others; the rostrum is armed with at least two ventral teeth, the first minute; the upper flagellum of the antennule is fused for six basal segments. The present specimen is larger than the holotype, with the postorbital carapace length 4.90 mm, telson 3.45 mm. Both second pereiopods and most of the ambulatory pereiopods are lost in the holotype. The present specimen has both the second pereiopods and the ambulatory pereiopods on right side. Second pereiopods are similar and equal, overreaching the end of the scaphocerite by carpus and chela; the chela is subequal to the postorbital carapace length; the palm elongate, about 6.7 times longer than the maximal depth, with very sparse small tubercles; the fingers about 0.46 of the palm length, with strongly hooked tips, proximal 0.3 of the cutting edges of both fingers armed with four to five minute teeth, distal 0.7 of cutting edges entire, sharp; the dactyl about 5.5 times longer than the proximal depth; the carpus distinctly swollen distally, about 0.77 of the palm length, 6.07 times the distal width, smooth; the merus unarmed distoventrally, subequal to palm length; ischium 0.95 of palm length; basis and coxa normal. The ambulatory pereiopods are slender. The third pereiopod overreaching the end of the scaphocerite by the dactyl and distal three-fifths of the propod; the propod 0.65 of the postorbital carapace length, armed with three

pairs of long spines and long setae distoventrally; the dactyl 0.17 of the propod length, slender and hooked, the flexor margin concave, unarmed, the unguis distinct; the carpus, merus and ischium 0.62, 1.05 and 0.45 times the palm length, respectively. The fourth and fifth pereopods similar to the third pereopod. The diameter of egg is 0.45 and length 0.57 mm.

Previously only known from its type locality. The present record extends the range far eastward to Fiji, and the bathymetric range to 350–365 m.

Distribution

Type locality: East China Sea west of Goto Retto, Kyushu, Japan, 145 m. Now also known from Fiji, 350–365 m.

Periclimenes magnificus Bruce, 1979

Periclimenes magnificus Bruce 1979b, p 195, Figures 1–5; Bruce 1996, p 237; Fransen 1989, p 143, Figures 4b, c, 5e, 6i–m, 7i–p; Li 2000, p 212, Figure 274; Li et al. 2004, p 545, Figure 30; Marin et al. 2004, p 209, Figure 11.

Material examined

New Caledonia: (i) Vie Bay, scuba, 10 m, coll. P. Laboute, 8 June 1990, 1 ovig. ♀ (MNHN-Na 15841); (ii) Îlot Sainte Marie, cf. antenne radio, scuba, 11 m, coll. P. Laboute, 8 June 1990, 1 ovig. ♀; (iii) Îlot Maitre, stn 64, 22°19.35'S, 166°25.85'W, 21 m, coll. Alan Myers, 10 November 1995, 1♂, 1 ovig. ♀ (MNHN-Na 15864); (iv) Îlot Crouy, 3 m, coll. Berthault, 15 June 1996, 1 ovig. ♀ (MNHN-Na 15878).

Hosts

(ii) *Goniopora* sp.; (iii) sea anemone [Scleractinia; Actinaria, Coelenterata].

Remarks

Previously recorded from New Caledonia by Bruce (1996).

Distribution

Type locality: Heron Island, Queensland, 26–29 m. Also known from Vietnam, South China Sea, Japan, Philippines, Indonesia, Australia (Northern Territory, Queensland), and New Caledonia.

Periclimenes novaecaledoniae Bruce, 1967

(Figure 26)

Periclimenes novaecaledoniae Bruce 1967b, p 1157–1165, Figures 6–9; Bruce 1991b, p 237; Bruce 1996, p 237; Li 2000, p 215, Figure 279; Davie 2002, p 330.

Material examined

New Caledonia: (i) Touho Bank, with crinoids, 28 August 1993, 2♂♂, 1♀ (MNHN-Na 15895); (ii) MONTROUZIER, Koumac, Chenal Infernet, associated with crinoids, coll. B. Richer de Forges, photo. B. Rudman, 5 October 1993, 1 ovig. ♀ (MNHN-Na 14919).




Figure 26. *Periclimenes novaecaledoniae* Bruce, 1967, ovigerous female (MNHN-Na 14919), third pereiopod. (a) Distal propod and dactyl; (b) distal corpus and unguis, dactyl; (c) distoventral propodal spine; (d) distal second ventral spine.

Remarks

Originally described from New Caledonia. The specimen has a rostral dentition of 6/1 and has both second pereopods, which are subequal, with the fingers strongly concave, subequal to the palm length, tridentate proximally, distally finely serrulate. Third pereopod with dactyl biunguiculate with a dorsal spinule; propod spinulate, sparsely setose. The proximal segment of the antennular peduncle is strongly bidentate distolaterally.

Distribution

Type locality: Îlot Maitre, New Caledonia. Otherwise known only from Australia (Northern Territory), and possibly from Madagascar.

Periclimenes paralcocki sp. nov.

(Figure 27)

Material examined

Tuvalu: MUSORSTOM 7, Banc Bayonnaise, stn CP631, 11°54'S, 179°32'W, 600 m, coll. 29 May 1992, 1♀ holotype (MNHN-Na 14865).




Figure 27. *Periclimenes paralcocki*, new species, holotype ovigerous female (MNHN-Na 14865). (a) Carapace and anterior appendages; (b) anterior carapace; (c) eye, dorsal view; (d) antennular peduncle; (e) antenna; (f) left third pereopod; (g) same, distal propod and dactyl; (h) telson; (i) same, distal part.

Description

A large pontoniine shrimp of subcylindrical body form, very similar to *Periclimenes alcocki* Kemp, 1922; lacking right (major?) second pereopod.

Carapace smooth, glabrous, with feeble dorsal hump on posterior two-fifths. Rostrum well developed, compressed, moderately deep, anteroventrad mostly and slightly upcurved apically, with feeble dorsal curvature, distinctly overreaching antennular peduncle, extending near to tip of scaphocerite, about 0.7 of carapace length; dorsal carina distinct, with seven large, acute teeth evenly distributed along dorsal margin, distal tooth much smaller than proximal teeth, first tooth situated on carapace just behind level of posterior orbital margin, with feeble basal suture; lateral carinae well developed, horizontal except up-curved distal part; ventral margin with three teeth on distal half between fifth and seventh dorsal teeth; interdental spaces and proximal ventral carina feebly setose. Supraorbital spine absent; epigastric spine situated at anterior 0.25 of carapace length, with basal suture; orbit feebly developed, inferior orbital angle produced, oval rounded; antennal spine small slender marginal, distinctly below inferior orbital angle, not exceeding inferior orbital angle; hepatic spine larger than antennal spine, slender, slightly lower than antennal spine, below the level between epigastric spine and first dorsal rostral tooth, remote from anterior carapace margin; anterolateral angle of carapace not produced, bluntly rounded.

Abdominal segments smooth, glabrous; third abdominal segment not produced, non-carinate; pleura of first three segments broadly rounded, fourth and fifth posteriorly produced rounded, sixth segment about 1.75 times length of fifth, subcylindrical, about 1.54 times longer than central depth, posterolateral angle acute, posteroventral angle rounded. Telson about 1.9 times sixth segment length, slender, about 4.3 times longer than anterior width, lateral margins sublinear, convergent, dorsal surface with three pairs of dorsolateral spines at 0.41, 0.73, and 0.87 of telson length, spines about 0.034 of telson length; posterior margin about 0.34 of anterior margin width, rounded, without acute posterior median point; posterior margin with normal three pairs of spines, lateral spines short, similar to dorsolateral spines, intermediate spines long, robust, about 0.067 of telson length, submedian spines about 0.45 of intermediate spine length, with sparse plumose setose.

Eye moderately reduced, with globular cornea, corneal diameter 0.125 of postorbital carapace length, without accessory pigment spot; stalk feebly compressed, subuniform, diameter subequal to cornea, length subequal to corneal diameter, subequal to dorsal width.

Antennular peduncle overreaching second ventral rostral tooth; proximal segment about 1.75 times longer than central width, with slender acute stylocerite laterally, reaching to about 0.7 of segment length, anterolateral margin feebly produced, setose, with long slender acute lateral tooth overreaching distodorsal end of intermediate segment; statocyst small; medial margin near straight, setose, with ventromedial tooth at about 0.5 of length; intermediate segment with dorsal length about 0.26 of proximal segment length, subequal to width, lateral margin expanded, setose; distal segment about 0.43 times proximal segment length, slender, about 1.77 times longer than distal width; upper flagellum biramous, with 11 proximal segments of rami fused, shorter free ramus with five segments, length about 0.4 of fused portion, total length about 0.48 of carapace length, with about 26 groups of aesthetascs; longer ramus slender, filiform, about 0.93 of carapace length; lower flagellum slender, filiform, about 0.93 times carapace length.

Antennal basicerite robust, with acute distolateral tooth; carpocerite subcylindrical, about 0.39 of scaphocerite length, 3.25 times longer than width; flagellum well developed,

slender, about 3.5 times postorbital carapace length; scaphocerite well developed, distinctly exceeding distal end of antennular peduncle, slightly overreaching rostral tip, broad, about 2.37 times longer than wide, greatest width at about 0.3 of length, distal margin bluntly rounded, slightly exceeding the distolateral tooth, lateral margin feebly convex, with strong acute distolateral tooth.

Epistome unarmed. Fourth thoracic sternite without slender median process, with broad triangular plates, separated by deep median fissure, fifth sternite similar, subequal to those of fourth; posterior sternites unarmed.

Mouthparts mainly undissected. Mandible without palp; incisor process distally with three (right) or four (left) stout acute teeth, central teeth smaller than outer teeth. Third maxilliped with slender endopod, extending distally to about 0.6 of carapocerite length, ischiomerus and basis fused incompletely, combined segment compressed, subuniform in ischiomerus portion and feebly produced medially in basal portion, setose medially, 5.4 times longer than central width; penultimate segment slightly compressed, about 0.53 of combined proximal segment length, with six groups of long finely spiniform setae medially; terminal segment about 0.37 of combined proximal segment length, 3.6 times longer than proximal width, distally tapering, with seven transverse groups of spiniform setae ventromedially, with long distal spiniform setae; exopod with slender flagellum just overreaching distal end of combined proximal segment, with nine plumose setae distally; coxa with small subacute setose medial process, rounded lateral plate, arthrobranch distinct.

First pereopods moderately slender, extending slightly beyond distal margin of scaphocerite, exceeding carapocerite by chela and distal half carpus; chela with palm subcylindrical, slightly compressed, about 2.25 times longer than maximal depth, at about proximal 0.3 of length with six transverse rows of short cleaning setae proximoventrally; fingers 0.86 of palm length, robust, tapering, subspatulate, surrounded by five groups of long setae medially and laterally along the fingers length, distal group shorter, cutting edges sharp, entire; carpus about 1.27 of chela length, 5.4 times longer than distal width, tapering proximally, with transverse row of serulate setae distoventrally; merus about 1.42 times chela length, 6.6 times longer than distal width, subuniform; ichium 0.64 of chela length, 2.7 times longer than distal width, feebly carinate distoventrally, obliquely articulated with basis; basis and coxa normal, no special feature.

Left (? minor) second pereopod well developed, exceeding carapocerite by chela, carpus and distal sixth merus, scaphocerite by fingers and distal two-thirds of palm; chela subequal to postorbital carapace length; palm about 4.2 times longer than maximal depth, with very small tubercles; fingers about 0.6 of palm length, with strongly hooked tips, cutting edges entire, sharp, proximal 0.3 of fixed finger cutting edge with two small teeth, proximal one smaller, rounded, distal one larger, acute, proximal 0.4 of dactyl cutting edge with two small acute teeth, proximal tooth larger than distal; dactyl about 4.3 times longer than proximal depth, with feebly developed lateral flange; carpus cup-like, about 0.37 of palm length, 1.67 times distal width, with scattered small tubercles; merus about 0.8 of palm length, covered with scattered small tubercles; ischium 0.66 of palm length; basis and coxa normal.

Ambulatory pereopods robust. Third pereopod exceeds carapocerite by dactyl and propod, scaphocerite by dactyl; dactyl compressed, curved, about 0.15 of propod length, unguis distinct, about 0.54 of dorsal length of corpus, 4.2 times longer than proximal width; corpus about 2.38 times longer than proximal depth, dorsal margin feebly convex, non-setose, ventral margin largely straight, with acute distal accessory tooth, about 0.31 of

unguis length, with short distolateral sensory setae; propod about 0.56 of carapace length, 9.3 times longer than maximum wide, subuniform, with pair of distoventral spines and one pair of sub-distoventral spines, and three spines distributed along distal 0.6 of ventral margin, with row of long simple setae distolaterally, and group of long simple setae distodorsally, small groups of long simple setae along ventral and dorsal margins; carpus about 0.51 of propod length, 4.3 times longer than distal width, with long produced distodorsal lobe, unarmed; merus subequal to propod, 7.9 times longer than central width, uniform, unarmed; ischium 0.49 of propod length, 3.4 times longer than distal width, feebly tapered proximally; basis and coxa without special features. Fourth and fifth pereopods similar to third, fourth propod subequal, fifth 1.09 times third propod length; fifth exceeds carpocerite by dactyl and distal one-third propod.

Pleopods without special features.

Uropod distinctly exceeding distal end of extended intermediate posterior spine of telson; protopodite with posterolateral angle short, rounded; exopod 2.59 times longer than central width, lateral border distinctly convex, with small acute distal tooth, and small mobile spine medially, diaeresis distinct; endopod about 0.93 of exopod length, 3.3 times longer than central width.

Measurements (mm)

Carapace length, 8.38; carapace and rostrum, 14.4; total body length (approximately), 40.0; left (? minor) second pereopod chela, 8.23.

Systematic position

Perichlimes *paralcocki* n. sp. is very close to *P. alcocki* Kemp, 1922. The telson with three pairs of dorsolateral spines distinguishes the new species from *P. alcocki* and other related species immediately. Most of the other *P. alcocki* allied species, which are distributed in deep water and with more than two pairs of dorsolateral telson spines, have four pairs of dorsolateral telson spines except *P. albatrossae* Chace and Bruce, 1993, which has seven pairs of those spines. *Perichlimes paralcocki* can be distinguished from *P. alcocki* also by the relatively larger cornea, telson spines and accessory tooth of ambulatory pereopod dactyl, and the rostrum distinctly reaching beyond the antennular peduncle, the flexor margin of ambulatory pereopod propod is armed with spines and setae along the whole length. In *P. alcocki*, the rostrum does not exceed the antennular peduncle, the flexor margin of ambulatory pereopod propod is unarmed except for the distal part. The number of dorsal telson spines of *P. alcocki* may show considerable variation (Bruce 1996, p 230), so the corneal diameter, length of rostrum, length of dorsolateral telson spines, and armature of ambulatory propod should be used to distinguish the allied species in the *Perichlimes alcocki* species group as well as the number of dorsolateral telson spines.

Etymology

Para (Greek), near, and the specific name *alcocki*, referring to the close relationship between the two species.

Periclimenes paraleator sp. nov.
(Figure 28)

Material examined

New Caledonia: MUSORSTOM 4, stn 178, 18°56.3'S, 163°12.9'E, 520 m, 18 September 1985, 1 ovig. ♀ holotype (MNHN-Na 14945).

Description

A medium-sized pontoniine shrimp of subcylindrical body form, very similar to *Periclimenes aleator* Bruce, 1991; both second pereopods lost.

Carapace smooth, glabrous. Rostrum well developed, compressed, shallow, horizontal, with distinct dorsal curvature, slightly extending antennal scale, about 0.86 of carapace length; dorsal carina distinct, with seven long, acute subequal teeth along proximal 0.6 rostral length and anterior dorsal carapace, and minute subapical tooth, first tooth situated on carapace, second just above posterior orbital margin; lateral carinae feebly developed; ventral margin with four teeth on distal 0.6 ventral rostral length, first two long, similar to dorsal teeth, third tooth much smaller than first two, distal tooth subapical, minute, smaller than subapical dorsal tooth; interdental spaces and proximal ventral carina feebly setose. Supraorbital spine absent; epigastric spine situated at anterior 0.3 of the carapace length, with basal suture; orbit feebly developed, inferior orbital angle acutely produced, oval; antennal spine small, slender, marginal, distinctly below inferior orbital angle, not exceeding inferior orbital angle; hepatic spine larger than antennal spine, slender, slightly lower than antennal spine, below first dorsal rostral tooth, remote from anterior carapace margin; anterolateral angle of carapace not produced, bluntly rounded.

Abdominal segments smooth, glabrous; third abdominal segment not produced, non-carinate; pleura of first three segments broadly rounded, fourth and fifth posteriorly produced rounded, sixth segment about 1.6 times length of fifth, subcylindrical, about 1.8 times longer than deep, subuniform, posterolateral angle acute, posteroventral angle rounded. Telson about 1.62 times sixth segment length, slender, about 4.4 times longer than anterior width, lateral margins sublinear, with anterior third curved ventrally, subparallel, posterior two-thirds straight, convergent, dorsal surface with four pairs of well-developed dorsolateral spines at 0.39, 0.54, 0.67, and 0.84 of telson length, spines about 0.044 of telson length; posterior margin about 0.39 of anterior margin width, rounded, without acute posterior median point; posterior margin with normal three pairs of spines, lateral spines short, similar to dorsolateral spines, intermediate spines long, robust, about 0.09 of telson length, submedian spines about 0.47 of intermediate spine length.

Eye moderate large, with globular cornea, corneal diameter 0.16 of posterior orbital carapace length, without accessory pigment spot; stalk compressed, distinctly narrower than cornea in lateral view, 0.8 of corneal diameter, 1.22 times lateral width and 0.93 of dorsal maximum width.

Antennular peduncle overreaching second ventral rostral tooth; proximal segment about 2.0 times longer than central width, with slender acute stylocerite laterally, reaching to about 0.7 of segment length, anterolateral margin feebly produced, setose, with long slender acute lateral tooth reaching to about 0.6 of intermediate segment length; statocyst small; medial margin near straight, setose, with ventromedial tooth at about 0.5 of length; intermediate segment with dorsal length about 0.26 of proximal segment length, subequal to width, lateral margin expanded, setose; distal segment about 0.41 times proximal




Figure 28. *Perichimenes paraleator*, new species, holotype ovigerous female (MNHN-Na 14945). (a) Carapace and anterior appendages; (b) eye, dorsal view; (c) same, lateral view; (d) antennular peduncle; (e) antenna; (f) third pereopod, propod and dactyl; (g) same, distal propod and dactyl.

segment length, slender, about 1.75 times longer than distal width; upper flagellum biramous, with 11 proximal segments of rami fused, shorter free ramus with five segments, length about 0.42 of fused portion, total length about 0.54 of carapace length, with about 25 groups of aesthetascs; longer ramus slender, filiform, about 0.7 of carapace length; lower flagellum slender, filiform, about 0.95 times carapace length.

Antennal basicerite robust, with acute distolateral tooth; carpocerite subcylindrical, about 0.35 of scaphocerite length, 3.0 times longer than width; flagellum well developed, slender, more than 2.4 times postorbital carapace length; scaphocerite well developed, distinctly exceeding distal end of antennular peduncle, reaching beyond third ventral rostral tooth, broad, about 2.85 times longer than wide, greatest width at about 0.35 of length, distal margin bluntly rounded, distinctly exceeding distolateral tooth, lateral margin nearly straight, with strong acute distolateral tooth.

Epistome unarmed. Fourth thoracic sternite without slender median process, with broad triangular plates, separated by deep median fissure, fifth sternite similar, lateral plates smaller; posterior sternites unarmed.

Mouthparts mainly undissected. Mandible without palp; incisor process distally with three (right) or four (left) stout acute teeth, central teeth smaller than outer teeth. Third maxilliped with slender endopod, extending distally to middle of carpocerite, ischimerus and basis fused completely, combined segment compressed, subuniform, setose medially, 6.1 times longer than central width; penultimate segment slightly compressed, about 0.57 of combined proximal segment length, with five groups of long finely spiniform setae medially; terminal segment about 0.45 of combined proximal segment length, 4.7 times longer than proximal width, distally tapering, with six transverse groups of spiniform setae ventromedially, with long distal spiniform setae; exopod with slender flagellum extending to distal end of combined proximal segment, with seven plumose setae distally; coxa with small subacute setose medial process, rounded lateral plate, arthrobranch distinct.

First pereopods moderately slender, extending to tip of rostrum, exceeding carpocerite by chela and distal two-thirds carpus; chela with palm subcylindrical, slightly compressed, about 2.0 times longer than maximal depth, at about proximal 0.3 of length with six transverse rows of short cleaning setae proximoventrally; fingers 0.93 of palm length, robust, tapering, surrounded by three groups of long setae medially and laterally, distally with shorter setae, cutting edges sharp, entire; carpus about 1.28 of chela length, 5.0 times longer than distal width, tapering proximally, with transverse row of serrulate setae distoventrally; merus about 1.5 times chela length, 7.4 times longer than distal width, subuniform; ichium 0.68 of chela length, 3.6 times longer than distal width, feebly carinate distoventrally, obliquely articulated with basis; basis 0.47 of chela length; coxa normal, with small, non-setose distoventral process.

Ambulatory pereopods slender. Third pereopod exceeds carpocerite by dactyl and propod, scaphocerite by dactyl; dactyl compressed, curved, about 0.19 of propod length, unguis distinct, about 0.51 of dorsal length of corpus, 3.64 times longer than proximal width; corpus about 2.2 times longer than proximal depth, dorsal margin feebly convex, non-setose, ventral margin largely straight, with acute distal accessory tooth, about 0.35 of unguis length, with short distolateral sensory setae; propod about 0.48 of carapace length, 10.5 times longer than maximum wide, subuniform, with pair of distoventral spines and four spines distributed along distal three-quarters ventral margin, with row of long simple setae distolaterally, and group of long simple setae distodorsally, small groups of long simple setae along ventral and dorsal margins; carpus about 0.56 of propod length, 5.1 times longer than distal width, with long produced distodorsal lobe, unarmed; merus 1.16

of propod length, 10.0 times longer than central width, uniform, unarmed; ischium 0.64 of propod length, 4.9 times longer than distal width, feebly proximally tapered; basis and coxa without special features. Fourth and fifth pereopods similar to third, fourth propod 1.08, fifth propod 1.14 times third propod length; fifth exceeds carpocerite by dactyl and half propod.

Pleopods without special features.

Uropod distinctly exceeding distal end of extending intermediate posterior spine of telson; protopodite with posterolateral angle short, rounded; exopod 2.76 times longer than central width, lateral border distinctly convex, with small acute distal tooth, with large mobile spine medially, diaeresis distinct; endopod about 0.97 of exopod length, 3.5 times longer than wide.

Ova small, numerous, more than 300.

Measurements (mm)

Carapace length, 6.03; carapace and rostrum, 11.06; total body length (approximately), 30.1; length of ovum, 0.65.

Systematic position

Periclimenes paraleator n. sp. is very close to *P. aleator* Bruce, 1991. It can be distinguished from the latter by the longer rostrum which distinctly exceeds the scaphocerite. In *P. aleator*, the rostrum also reaches slightly beyond the antennular peduncle. Also, the new species has a more robust body and relatively shorter ambulatory pereopods than *P. aleator*. *Periclimenes paraleator* is also similar to *P. alcocki* Kemp, 1922. It can be distinguished from the latter by the longer rostrum, larger cornea and more developed dorsolateral telson spines and the propods of ambulatory pereopods with spinules along most ventral margin.

Etymology

Para (Greek), near, and the specific name *aleator*, referring to the close relationship between the two species.

Remarks

In the deep-water *Periclimenes* species with more than two pairs of dorsolateral telson spines, only *P. albatrossae* Chace and Bruce, 1993 has the rostrum exceeding the scaphocerite. *Periclimenes paraleator* can be distinguished immediately from that species by having only four pairs of dorsolateral telson spines. In *P. albatrossae*, seven pairs of those spines are present. In addition, the propod of ambulatory pereopods in *P. albatrossae* is unarmed on the flexor margin.

Periclimenes pilipes Bruce and Zmarzley, 1983

(Figure 29)

Periclimenes pilipes Bruce and Zmarzly 1983, p 644–654, Figures 1–6; Chace and Bruce 1993, p 119; Li 2000, p 226, Figure 297.

Material examined

Philippines: Maribago, Mactan Island, Cebu, 10°17'N, 124°00'E, 3–5 m, washings from mixed crinoids, coll. P. Bouchet, 9 June 1985, 1♂, 1 juvenile ♀ (MNHN-Na 14920).

Hosts

Comanthina schlegeli (P. H. Carpenter, 1881), *Comatella nigra* (P. H. Carpenter, 1888), *Comanter multibrachiata* (P. H. Carpenter, 1888), *Comanthus bennetti* (J. Müller, 1841), det. A. M. Clark [Crinoidea, Echinoderma].




Figure 29. *Periclimenes pilipes* Bruce and Zmarzley, 1983 (MNHN-Na 14920). (a) Third pereiopod, distal propod and dactyl; (b) same, distal corpus and unguis.

Remarks

Previously reported from Moalboal Island in the Philippines by Bruce (1989a). The male specimen, cl 2.5 mm, has a rostral dentition of 6/2 and the female, 5/1. The first pereopod has the fingers feebly subspatulate; second pereopods stout, carpus short, merus unarmed; third pereopod with dactyl biunguiculate, with conspicuous trifid dorsal spine on proximal unguis, propod with long straight stiff setae. Proximal segment of antennular peduncle strongly bidentate distolaterally.

Distribution

Type locality: Eniwetak Atoll, Marshall Islands. So far, known only from Marshall Islands and Philippines.

***Periclimenes platyrhynchus* Bruce, 1991**

Periclimenes platyrhynchus Bruce 1991a, p 358, Figures 41–44; Li 2000, p 227, Figure 300.

Material examined

New Caledonia: BATHUS 4, stn DW942, 19°04.26'S, 163°27.36'E, 270–264 m, 8 August 1994, 2♂♂, 3 ovig. ♀♀ (MNHN-Na 15922).

Remarks

Previously known only from the type locality New Caledonia.

Distribution

Type locality: only, New Caledonia.

***Periclimenes pseudalcocki* sp. nov.**

(Figure 30)

Material examined

Indonesia: KARUBAR, Kai Islands, stn CP16, 5°17'S, 132°50'E, 315–349 m, 24 October 1991, 1♀ holotype (MNHN-Na 14855).

Description

A large pontoniine shrimp of subcylindrical body form, very similar to *Periclimenes alcocki* Kemp, 1922.

Carapace smooth, glabrous. Rostrum well developed, compressed, deep, especially anteroventrally and distinctly upcurved distally, distinctly overreaching antennular peduncle, extending near to tip of scaphocerite, about 0.68 of carapace length; dorsal margin with seven low acute teeth, first tooth situated on carapace just behind level of posterior orbital margin; lateral carinae well developed, except upcurved distal part; ventral margin with two small acute teeth on distal third and subapical small tubercle, first tooth at level of last dorsal tooth. Supraorbital spine absent; epigastric spine similar to dorsal rostral




Figure 30. *Periclimenes pseudalcocki*, new species, holotype female (MNHN-Na 14855). (a) Carapace and anterior appendages; (b) telson; (c) same, distal part; (d) eye, dorsal view; (e) antennular peduncle; (f) antenna; (g) third pereiopod; (h) same, distal propod and dactyl, lateral view; (i) same, medial view.

tooth, situated at anterior 0.23 of carapace length; orbit feebly developed, inferior orbital angle produced, oval rounded; antennal spine small slender marginal, distinctly below the inferior orbital angle, not exceeding inferior orbital angle; hepatic spine larger than antennal spine, stout, slightly lower than antennal spine, below level between epigastric spine and first dorsal rostral tooth, remote from anterior carapace margin; anterolateral angle of carapace not produced, bluntly rounded.

Abdominal segments smooth, glabrous; third abdominal segment not produced, non-carinate; pleura of first three segments broadly rounded, fourth and fifth posteriorly produced rounded, sixth segment about 2.0 times length of fifth, subcylindrical, about 1.52 times longer than central depth, posterolateral angle acute, posteroventral angle rounded. Telson about 1.64 times sixth segment length, slender, about 3.35 times longer than anterior width, lateral margins sublinear, convergent, dorsal surface with four pairs of dorsolateral spines at 0.45, 0.64, 0.78 (average), and 0.90 (average) of telson length, dorsal spines about 0.045 of telson length; posterior margin about 0.40 of anterior margin width, rounded, with minute sharp posterior median point; posterior margin with normal three pairs of spines, lateral spines short and robust, similar to dorsolateral spines, intermediate spines long, robust, about 0.075 of telson length, submedian spines about 0.41 of intermediate spine length, non-setulose.

Eye reduced, with globular cornea, corneal diameter 0.11 of posterior orbital carapace length, without accessory pigment spot; stalk feebly compressed, subuniform, dorsal width subequal to corneal diameter, length 0.95 of corneal diameter.

Antennular peduncle overreaching second ventral rostral tooth; proximal segment about 1.43 times longer than central width, with slender acute stylocerite laterally, reaching to about 0.7 of segment length, anterolateral margin feebly produced, setose, with long slender acute lateral tooth overreaching distodorsal end of intermediate segment; statocyst small; medial margin near straight, setose, with ventromedial tooth at about 0.54 of length; intermediate segment with dorsal length about 0.3 of proximal segment length, subequal to width, lateral margin expanded, setose, medial margin strongly setose; distal segment about 0.54 times proximal segment length, slender, about 1.7 times longer than distal width; upper flagellum well developed, biramous, with proximal 12 segments fused, shorter free ramus with four segments, length about 0.28 of fused portion, total length about 0.49 of carapace length, with about 28 groups of aesthetascs; longer ramus slender, filiform, about 0.55 of carapace length; lower flagellum slender, filiform, about 0.83 times carapace length.

Antennal basicerite robust, with strong acute distolateral tooth; carpocerite subcylindrical, about 0.39 of scaphocerite length, 2.67 times longer than width; flagellum well developed, slender, about 2.0 times postorbital carapace length; scaphocerite well developed, distinctly exceeding rostral tip, broad, about 2.16 times longer than wide, greatest width at about 0.3 of length, distal margin bluntly rounded, slightly exceeding the distolateral tooth, lateral margin feebly convex, with strong acute distolateral tooth.

Epistome unarmed. Fourth thoracic sternite without slender median process, with broad triangular plates, separated by deep median notch, fifth sternite similar, subequal to those of fourth; posterior sternites unarmed.

Mouthparts mainly undissected. Mandible without palp; incisor process distally with three (right) or four (left) stout acute teeth, central teeth smaller than outer teeth. Third maxilliped reaching to distal end of carpocerite, similar to that of *P. alcocki*, coxa with arthrobranch distinct.

First pereopods moderately slender, exceeding distal margin of antennal scale by chela, exceeding carpocerite by chela and five-sixths carpus; similar to that of *P. alcocki*.

Second pereopods well developed, subsimilar except the fingers, chelae covered with small tubercles. Major (right) pereopod exceeding distal end of antennal scale by chela, carpocerite by chela, carpus and distal third merus; minor (left) pereopod also exceeding distal end of antennal scale by chela, carpocerite by chela, carpus and distal third merus. Other features very similar to those of *P. alcocki*.

Ambulatory pereopods slender. Third pereopod exceeds carpocerite by dactyl, propod and distal two-fifths carpus, scaphocerite by dactyl and half propod; dactyl compressed, slender, curved, about 0.18 of propod length, unguis distinct, about 0.62 of dorsal length of corpus, 5.5 times longer than proximal width; corpus about 2.57 times longer than proximal depth, dorsal margin feebly convex, non-setose, ventral margin largely feebly concave, except feebly convex proximal portion, distal accessory tooth acute, minute, about 0.11 of unguis length, with single short distolateral sensory seta; propod compressed, subuniform, about 0.52 of carapace length, 8.3 times longer than wide, with three transverse rows of long simple setae distoventrally and one row of similar setae distodorsally, without distoventral spines, major part of ventral border unarmed, non-setose; carpus feebly compressed, tapering feebly proximally, about 0.59 of propod length, 4.9 times longer than distal width, with long produced distodorsal lobe, unarmed; merus

compressed and feebly twisted, 1.16 of propod length, 7.9 times longer than central width, uniform, unarmed; ischium compressed, 0.64 of propod length, 4.0 times longer than distal width, feebly proximally tapered; basis and coxa without special features. Fourth and fifth pereopods similar to third, fourth propod 1.1 of third propod length, fifth propod subequal to fourth; fourth and fifth dactyl with the accessory tooth very minute, like tubercle; fifth exceeds carpocerite by dactyl and distal two-thirds propod, scaphocerite by dactyl.

Pleopods without special features.

Uropod distinctly exceeding distal end of telson, exopod reaching the distal tip of extended intermediate posterior spines; protopodite with distolateral lobe broadly rounded; exopod 2.17 times longer than central width, lateral border feebly convex, with small distal tooth, and larger mobile spine medially, diaeresis distinct, at 0.75 of length; endopod subequal to exopod length, 2.7 times longer than central width.

Measurements (mm)

Carapace length, 8.64; carapace and rostrum, 14.2; total body length (approximately), 40.0; right (major) second pereopod chela, 10.32; left (minor) second pereopod chela, 7.88.

Systematic position

Periclimenes pseudalcocki n. sp. is very close to *P. alcocki* Kemp, 1922. It can be distinguished from the latter by the smaller accessory tooth of ambulatory dactyl which is at most 0.11 of unguis length (in third pereopod, it is like a tubercle in the fourth and fifth pereopod), and the propod has no distoventral or ventral spines. In *P. alcocki*, the accessory tooth of ambulatory dactyl is about 0.2 of unguis length, and propod has one distoventral spine and two ventral spines. In addition, the rostrum of *P. pseudalcocki* is deeper, and dorsal telson spines larger (about 0.045 of telson length, it is 0.03 in *P. alcocki*). *Periclimenes pseudalcocki* can be distinguished from other allied deep-water species with four pairs of dorsolateral telson spines by the markedly reduced cornea, the corneal diameter 0.11 of carapace length. *Periclimenes pseudalcocki* is also very similar to *P. crosnieri* n. sp. It can be distinguished from that species by the biunguiculate third pereopod dactyl, non-spinules propod, reduced cornea, and other detailed features.

Etymology

Pseudos (Greek), fallacy, and the specific name *alcocki*, referring to the close relationship between the two species.

Key to the species of *Periclimenes alcocki* species group

1. Telson with seven pairs of dorsolateral spines; rostrum overreaching scaphocerite; third pereopod with dactyl truncate subdistally, propod without spinules on flexor margin *P. albatrossae* Chace and Bruce, 1993

- Telson with three to five pairs of dorsolateral spines 2.
- 2. Telson with three pairs of dorsolateral spines; rostrum not overreaching scaphocerite; corneal diameter about 0.125 of carapace length, eye stalk not distinctly narrower than cornea; dorsal rostral margin concave; third pereopod with propod with spinules on flexor margin, and transverse setal rows distally and spines; R. 1+7/3 *P. paralcocki* sp. n.
- Telson with four to five pairs of dorsolateral spines 3
- 3. Rostrum overreaching scaphocerite; corneal diameter about 0.16 of carapace length, eye stalk distinctly narrower than cornea lateral view; dorsal rostral margin concave; third pereopod with propod with spinules on flexor margin, and transverse setal rows distally and spines; R. 1+7+1/4 *P. paraleator* sp. n.
- Rostrum not overreaching scaphocerite. 4
- 4. Cornea markedly reduced, diameter about 0.1 of carapace length 5
- Cornea not markedly reduced, diameter more than 0.13 of carapace length; dorsal telson spines not minute 7
- 5. Ambulatory pereopods with propod without distoventral or ventral spines, dactyl with accessory tooth minute, at most 0.11 of unguis length; rostrum relatively deep, distally with a strong upward curve; dorsal telson spines relatively large, 0.045 of telson length; R. 1+7/2 *P. pseudalcocki* sp. n.
- Ambulatory pereopods with propod with distoventral and ventral spines 6
- 6. Corneal diameter about 0.075 of carapace length; hepatic spine at about same level of posterior rostral series (epigastric) tooth; third pereopod dactyl 0.17 of propod length, accessory tooth about 0.2 of unguis length; dorsal telson spines minute, about 0.03 telson length; R. 2+6–8/2–4 (1+9/3) *P. alcocki* Kemp, 1922
- Corneal diameter about 0.1 of carapace length; hepatic spine anterior to level of posterior rostral series (epigastric) tooth; third pereopod dactyl about 0.28 of propod length, accessory tooth about 0.07 of unguis length; dorsal telson spines minute, about 0.04 telson length; R. 1+7/1 *P. tangeroa* Bruce, 2005
- 7. Ambulatory dactyl with accessory tooth almost as long as unguis, laterally twisted; R. 1+7–9/2–3 *P. poupini* Bruce, 1990
- Ambulatory dactyl with accessory tooth not unusually long or twisted. 8
- 8. Rostrum distinctly exceeding antennular peduncle 9
- Rostrum not reaching end of antennal peduncle; pigmented cornea large, diameter about 0.16–0.18 of carapace length 10
- 9. Hepatic spine distinctly lower than antennal spine; R. 1–2+5–9/3–4
. *P. aleator* Bruce, 1991
- Hepatic spine very slightly lower than, almost at same level of antennal spine; R. 2+6/3 *P. loyautensis* sp. n.
- 10. Rostrum deeper, proximally not elevated over orbital region, upturned in female, ventral margin convex, with two teeth; ambulatory dactyl with accessory tooth about 0.3 of unguis length; R. 1+5–6/2 *P. brevirostris* Bruce, 1991
- Rostrum shallower, proximally elevated over orbital region, tapered distally, ventral margin straight, with one tooth; ambulatory dactyl with accessory tooth reaching about proximal 0.4 of unguis length; R. 1+6–7/1 *P. forgesi* sp. n.

Periclimenes soror Nobili, 1904

Periclimenes soror Nobili 1904, p 232.

Periclimenes (Periclimenes) soror: Holthuis 1952a, p 51–53, Figure 17.

Periclimenes soror Bruce 1976c, p 299–306, Figures 1–6; Wicksten and Hendrickx 1985, p 571; Chace and Bruce 1993, p 122; Li 2000, p 237, Figure 316; Davie 2002, p 332; Li and Liu 2004, p 95, Figure 7.

Material examined

New Caledonia: (i) MONTROUZIER, Koumac, 25 m, 26 October 1993, 1♂, 1♀ (MNHN-Na 14924). **Loyalty Islands:** (ii) ATELIER LIFOU, Lifou, Santal Bay: west-southwest of Easo Point, stn 1429, 20°47.5'S, 167°07.1'E, 8–18 m, coral heads, sedimentary channels, 3, 5, 23, and 24 November 2000, 2♂♂, 2 ovig. ♀♀ (MNHN-Na 15540).

Host

Echinaster luzonicus (Grey, 1849) [Asteroidea, Echinoderma].

Remarks

The specimens had the accessory tooth on the ambulatory dactyls minute, virtually obsolete. The distolateral angle of the proximal segment of the antennular peduncle was strongly bidentate, but tridentate on the left side in the female.

Distribution

Type locality: Djibouti. Also known from the Red Sea, Kenya, Zanzibar, Tanzania, Madagascar, Comoro Islands, Seychelles, Chagos Islands, Malaysia (Sabah), China (Hong Kong), Japan, Philippines, Indonesia, Australia (Western Australia, Northern Territory, Queensland, New South Wales), Marianas Islands, Caroline Islands, Solomon Islands, New Caledonia, Loyalty Islands, Marshall Islands, Fiji, French Polynesia (Society Islands, Tuamotu Islands), and Hawaii. Also known from the East Pacific region, the Gulf of California, Mexico, Panama, and Columbia.

Periclimenes tosaensis Kubo, 1951

Periclimenes (Ancylocaris) tosaensis Kubo 1951, p 268, Figures 7, 8.

Periclimenes (Harpilius) tosaensis: Bruce 1966b, p 15, Figures 1, 2, 3a, 4a, b.

Periclimenes tosaensis: Bruce 1976a, p 106, Figure 12; Bruce 1981a, p 196, Figure 5; Li 2000, p 243, Figure 323; Li 2004b, p 821; Li et al. 2004, p 548, Figure 33.

Material examined

Fiji: MUSORSTOM 10, south Viti Levu, stn CP 1363, 18°12.4'S, 178°33.0'E, 144–150 m, 15 August 1998, 1 ovig. ♀ (MNHN-Na 15541).

Remarks

Not previously recorded from Fiji.

Distribution

Type locality: Tosa Bay, Shikoku, Japan. Also known from Seychelles, South China Sea, East China Sea, Philippines, and Fiji.

***Periclimeses uniunguiculatus* Bruce, 1990**

Periclimeses uniunguiculatus Bruce 1990a, p 167, Figures 12–15, 39e; Bruce 1996, p 239–241, Figures 14d–f, 28g; Li 2000, p 244, Figure 324.

Periclimeses sp. A Bruce 1991a, p 371.

Material examined

New Caledonia: (i) BERYX 11, south New Caledonia, stn DW27, 23°37'S, 167°41'E, 460–470 m, 18 October 1992, 1 ovig. ♀ (MNHN-Na 14885); (ii) BATHUS 3, stn CP832, 23°03.07'S, 166°53.70'E, 650–669 m, 30 November 1993, 1♀ (MNHN-Na 14941); (iii) HALIPRO 2, Norfolk Ridge, stn BT60, 24°52'S, 168°44'E, 1133–1280 m, 18 November 1996, 1♀ (MNHN-Na 15880).

Remarks

The specimens fit the original description and illustrations of the species very well, except for the larger body size with the postorbital carapace length 6.1 mm. The smaller body size of the holotype male may be caused by the bopyrid parasites present.

One of the authors (X. Li) checked the ovigerous female (MNHN-Na 12870) reported by Bruce (1996, under the name of *P. uniunguiculatus*, from Comoro Islands, at a depth of 500 m) and the adult female (MNHN-Na 12042) reported by Bruce (1991a, under the name of *P. sp. A*, from New Caledonia, at a depth of 825–860 m), during his visit to the museum (MNHN, Paris, 2004), and found that they are conspecific. Unfortunately, the holotype male was absent then. Anyway, both specimens mostly fit Bruce's (1990a) description. We are now sure the specimen of Bruce's (1991a) *P. sp. A* is identical to the present species. The present records extend the bathymetric range of the species from 460 to 1280 m.

Distribution

Type locality: New Caledonia. Also known from Comoro Islands (Bruce, 1996).

***Periclimeses vaubani* Bruce, 1990**

Periclimeses vaubani Bruce 1990a, p 174, Figures 16–19, 38a–d; Bruce 1991a, p 315; Bruce 2005, p 4, Figure 1; Li 2000, p 245, Figure 325.

Material examined

New Caledonia: (1) NORFOLK 1, Norfolk Ridge, Introuvable Bank, stn DW1697, 24°39'S, 168°38'E, 569–616 m, 24 June 2001, 1♂, 1 ovig. ♀ (MNHN-Na 15929); (ii) Norfolk Ridge, Antigonina Bank, stn CP 2122, 23°22'S, 168°00'W, 560–577 m, 1 November 2003, 1♀ (MNHN-Na 15940).

Distribution

Type locality: New Caledonia. Previously recorded from Norfolk Ridge by Bruce (2005) from West Norfolk Ridge in the Tasman Sea.

Periclimenes venustus Bruce, 1990

Periclimenes venustus Bruce 1990c, p 230–240, Figures 1–6, 7a, 8a; Chace and Bruce 1993, p 124; Li 2000, p 246, Figure 327; Davie 2002, p 333.

Material examined

New Caledonia: PLOUVEAL, stn DW1222, Loyalty Islands, Uvea Lagoon, 20°28'S, 166°30'E, 15 m, 12 September 1992, 1♀ (MNHN-Na 14871).

Remarks

Not previously reported from New Caledonia. The single specimen has both second pereiopods but lacks a rostrum, with an epigastric spine on the carapace.

Distribution

Type locality: Coral Bay, Port Essington, Northern Territory, Australia. Also known from Japan, Philippines, Indonesia, Papua New Guinea, Australia (Western Australia), and New Caledonia.

***Periclimenes* sp.**

(Figure 31)

Material examined

La Réunion: MD 32, stn DS173, 20°51.5'S, 55°36.8'E, 270 m, 8 September 1982, 1♀ (MNHN-Na 14927).

Description

The single specimen, cl 2.7 mm, unfortunately lacks both second pereiopods and the posterior part of the telson; right first pereiopod, third and fifth pereiopods detached. It is a small pontoniine shrimp with subcylindrical body form.

Carapace smooth, glabrous. The rostrum is long and slender, horizontal mostly and feebly upcurved distally, well exceeding the antennular peduncle and subequal to the postorbital carapace length, with a dentition of 1+6/5, epigastric spine and dorsal rostral teeth are large, long, slender and acute, as are ventral teeth, except for minute distal tooth, epigastric spine with basal suture, situated at anterior 0.4 of carapace length, first dorsal tooth on carapace, second dorsal tooth at level of posterior orbital margin. Supraorbital spine absent; antennal spine long, slender, marginal, close to inferior orbital angle, distinctly exceeding inferior orbital angle; hepatic spine large, as long as antennal spine, more robust, at same level as antennal spine, below the level between epigastric spine and first dorsal rostral tooth.

Third abdominal segment with feebly dorsal prominence; sixth segment about 2.1 times length of fifth, subcylindrical, about 1.7 times longer than central depth, posterolateral angle acute, posteroventral angle rounded.




Figure 31. *Periclimenes* sp. 1, female (MNHN-Na 14927). (a) Body, lateral view; (b) rostrum; (c) first pereiopod; (d) same, propod and dactyl.

Eye well developed, with globular cornea, corneal diameter 0.18 of posterior orbital carapace length, feebly pigmented, without accessory pigment spot; stalk subcylindrical, slightly wider than cornea in dorsal view.

Antennular peduncle overreaching third ventral rostral tooth; upper flagellum biramous, with proximal four segments fused, shorter free ramus with four segments, with nine groups of aesthetascs. Antennal basicerite robust, with strong acute lateral tooth; scaphocerite exceeding antennular peduncle, reaching basal fourth ventral rostral tooth; lateral margin straight, distolateral tooth almost reaching distal end of lamella.

Epistome with well-developed rounded bosses.

Fourth thoracic sternite without slender median process, with broad shallow transverse plates, separated by deep median fissure, fifth sternite similar, transverse plates subequal to those of fourth; posterior sternites unarmed.

Mandible without palp; incisor process distally with three (right) or four (left) stout acute teeth, central teeth smaller than outer teeth. Third maxilliped reaching to basal carapocerite, coxa with arthrobranch distinct.

First pereiopods moderately slender, reaching distal end of scaphocerite, exceeding carapocerite by chela and half carpus. Ambulatory pereiopods robust, third pereiopod exceeds carapocerite by dactyl and distal two-thirds propod; dactyl compressed, long, slender, simple, unguis distinct; propods with long distoventral spine and three small

ventral spinules. Uropod slender; protopodite with distolateral lobe broadly rounded; exopod with lateral border feebly convex, with small distal tooth, and larger mobile spine medially, diaeresis distinct.

Remarks

Although the peculiar rostrum, armature on carapace, epistome with well-developed rounded bosses, and the feebly dorsal prominence on third abdominal segment show that the present taxon is distinguishable from almost all of the members in the genus, it is difficult to present it as a new species as it lacks both second pereopods and the posterior part of the telson. These deficiencies make its exact systematic position unclear. More material is necessary to decide its precise systematic position. The armature of rostrum and carapace, particularly the long and acute rostral teeth and the hepatic spine which is at same level as antennal spine, looks somewhat like that of *P. latipollex* Kemp, 1922. The present form can be easily distinguished from that species by the simple dactyls of ambulatory pereopods and deeper rostrum (not rod-like, although shallow).

Philarius imperialis (Kubo, 1940)

Harpilius imperialis Kubo 1940a, p 1, Figures 1–3.

Philarius imperialis: Holthuis 1952a, p 15; Miyake and Fujino 1968, p 420, Figure 6; Bruce 1976b, p 482; Franssen 1994, p 132; Bruce and Coombes 1995, p 137; Li 1996, p 230, Figure 9; Li 1998, p 224; Li 2000, p 252, Figure 335; De Grave 2000, p 142.

Material examined

New Caledonia: (i) lagoon, Îlot Maitre, scuba, 5 m, with coral, coll. C. Vadon, 13 September 1978, 1 ovig. ♀ (abdomen damaged, tail-fan lost) (MNHN-Na 15848). **Loyalty Islands:** ATELIER LIFOU, Lifou, Santal Bay, (ii) opposite Ngoni beach, stn 1459, 20°47.0'S, 167°03.0'E, 55–80 m, trawl, 5 and 13 November 2000, 1 ovig. ♀ (MNHN-Na 15542); (iii) in the region of Easo Point, stn 1465, 20°47.7'S, 167°07.0'E, 35–45 m, trawl, 16 November 2000, 1 ovig. ♀ (MNHN-Na 15543).

Host

(i) *Acropora* sp. [Scleractinia].

Remarks

Not previously recorded from New Caledonia and Loyalty Islands.

Distribution

Type locality: Nankin-hama, Hahajima, Ogasawa Islands. Also known from Arabia, Israel, Kenya, Zanzibar, Tanzania, Seychelles, La Réunion, Singapore, Vietnam, China, Japan, Papua New Guinea, Australia (Western Australia, Northern Territory, Great Barrier Reef), Coral Sea, Caroline Islands, New Caledonia, Loyalty Islands, and Marshall Islands.

Platycaris latirostris Holthuis, 1952

Platycaris latirostris Holthuis 1952a, p 173, Figures 85, 86; Barnard 1962, p 243; Bruce 1966c, p 1, Figures 1–5; Bruce 1976d, p 47, Figure 4; Bruce 1985b, p 5, Figures 4, 5; Li 2000, p 254, Figure 337.

Material examined

Madagascar: Tuléar, Grand Récif, coll. R. Hipeau-Jacquotte, 1968, 10♂♂, 8♀♀ (6 ovig.) (MNHN-Na 15833).

Host

Galaxea sp. [Scleractinia].

Remarks

First recorded from Madagascar by Barnard (1962).

Distribution

Type locality: Indonesia. Also known from Zanzibar, Tanzania, Madagascar, Indonesia, Australia (Queensland), and Fiji.

Pliopontonia furtiva Bruce, 1973

Pliopontonia furtiva Bruce 1973a, p 99–108, Figures 1–5, Plate 1; Bruce 1991b, p 266; Chace and Bruce 1993, p 128; Li 2000, p 258, Figure 341; Davie 2002, p 336.

Material examined

New Caledonia: Canal Woodin, 15 m, scuba, coll. Hamel, 19 November 1992, 1♂ (MNHN-Na 14929).

Host

Ricordia florida (Duchassaing and Michelotti, 1860) [Corallimorpharia, Anthozoa].

Remarks

Previously reported from New Caledonia in Atoll de Huon (Bruce, 1991b). The specimen has a rostral dentition of 7/0 and has both second pereiopods: the ambulatory pereiopods are held in the dorsiflexed position as previously noted. The association with *Ricordia* represents a new host record.

Distribution

Type locality: Ras Iwatine, Kenya. Also known from the Seychelles, Japan, Philippines, Indonesia, Australia (Queensland, New South Wales), Solomon Islands, New Caledonia, and Marshall Islands.

***Pontoniopsis comanthi* Borradaile, 1915**

Pontoniopsis comanthi Borradaile 1915, p 213; Borradaile 1917, p 377, Plate 57, Figure 27; Holthuis 1952a, p 153–156, Figures 70, 71; Chace and Bruce 1993, p 130; Li 2000, p 276, Figure 369; Davie 2002, p 337.

Material examined

Philippines: (i) Maribago, Mactan Island, Cebu Province, 10°17'N, 124°00'E, 2–3 m, washings from crinoids, coll. P. Bouchet, 9 June 1985, 2♂♂, 1 ovig. ♀? (MNHN-Na 14930); (ii) same, 3–5 m, washings from mixed crinoids, 6♂♂, 5♀♀ (3 ovig.) (MNHN-Na 14931). **New Caledonia:** (iii) Touho Bank, on crinoid, 1♀ (MNHN-Na 15843); (iv) lagoon, on crinoids, 30 August 1993, 3♂♂, 1♀ (MNHN-Na 15883).

Hosts

(i) (ii) *Comanthina schlegeli* (P. H. Carpenter, 1881); (ii) also *Comatella nigra* (P. H. Carpenter, 1888), *Comanter multibrachiata* (P. H. Carpenter, 1888) *Comanthus bennetti* (J. Müller, 1841); det. A. M. Clark [Crinoides, Echinoderma].

Parasites

One specimen from (ii) has a hemiarthrinid parasite, *Paraphrixus nigrocinctus* (Chopra) [Bopyridae, Isopoda].

Remarks

Previously recorded from the Philippines, from Moalboal Island (Bruce 1989a). The specimens present no special features.

Distribution

Type locality: Mabuaig Island, Torres Strait. Also known from Israel, Red Sea, Kiribati, Kenya, Zanzibar, Seychelles, Japan, Philippines, Indonesia, Australia (Western Australia, Queensland), Coral Sea, Tasman Sea, Marianas Islands, Marshall Islands, and Fiji.

***Thaumastocaris streptopus* Kemp, 1922**

Thaumastocaris streptopus Kemp 1922, p 244, Figures 78–80; Bruce 1991b, p 258, Figure 21; Bruce 1996, p 250–251, Figure 21; Li 2000, p 286, Figure 380.

Material examined

Chesterfield: (i) AMUSIUM 1, stn CC16, 20°20'S, 160°48'E, 68–70 m, 6 May 1998, 1♀ (MNHN-Na 15819). **New Caledonia:** Îlot Maitre, (ii) lagoon, scuba, 25 m, with sponge, coll. C. Vadon, 5 September 1978, 1♂ (MNHN-Na 15837); (iii) Chenal of Îlot Maitre, scuba, 22 m, with sponge, coll. C. Vadon, 3 October 1978, 1♀ (MNHN-Na 15859); (iv) 22°19.35'S, 166°25.85'E, 20 m, scuba, with sponge, coll. I. Takeuchi, 10 November 1995, 1♂ (MNHN-Na 15825).

Remarks

The specimen of (i) has a carapace length 6.65 mm, rostral formula is 9/3, the first pereopods are equal, the carpa six subsegmented; (ii) has a carapace length 3.59 mm, rostral formula is 11/4, the carpa of first pereopods are equal but dissimilar, right with four subsegments, left with two subsegments, and right merus with two subsegments; (iii) has a

carapace length 6.90 mm, rostral formula is 9/3, the first pereopods are equal, with the carpa five-subsegmented and merus two-subsegmented; (iv) has a carapace length 7.43 mm, rostral formula is 10/3, the first pereopods are equal, with the carpa five-subsegmented.

Distribution

Type locality: New Caledonia. Also known from Jordan, Sudan, Somalia, Kenya, Zanzibar, Tanzania, Madagascar, Philippines, Indonesia, Australia (Queensland), Caroline Islands, Chesterfield Islands, New Caledonia, and Marshall Islands.

Zenopontonia noverca (Kemp, 1922)

Periclimenes (*Periclimenes*) *noverca* Kemp 1922, p 162–165, Figures 28–30.

Zenopontonia noverca: Bruce 1975a, p 277–280, Figure 1; Li 2000, p 307, Figure 408; Davie 2002, p 341.

Material examined

Madagascar: (i) Nosy Be' Tany Kely, 25 m, photo, CD 25, coll. P. Laboute, 21 September 1994, 1♂, 4 ovig. ♀♀ (MNHN-Na 14937); (ii) same, photo, CD 26, 2 ovig. ♀♀ (MNHN-Na 14938). **New Caledonia:** (iii) lagoon, entre of Seiche croissant Reef and Îlot Maitre, scuba, 12 m, with coral, coll. C. Vadon, 10 October 1978, 1 ovig. ♀ (MNHN-Na 15890); (iv) Nouméa (Anse Vata), 2 m, coll. Berthault, 1 May 2000, 1 ovig. ♀ (MNHN-Na 15892).

Hosts

(i) (ii) *Pentaceraster* sp. [Asteroïdes, Echinoderma]; (iii) *Pocillopora* sp. [Scleractinia].

Remarks

Previously reported from Madagascar, from Nosy Be, by Bruce (1975a, 1978a). (i) (ii) Five females have a rostral dentition of 8/0, one has 10/0, and the single male has 9/0; (iii) rostral formula 7/1. This species is usually associated with asteroids, so the association with *Pocillopora* is probably accidental.

Distribution

Type locality: New Caledonia. Also known from Zanzibar, Madagascar, Australia (Queensland), and New Caledonia.

Summary

By the studies of Bruce (1981a, 1983b, 1985a, 1990a, 1991a, 1991b, 1996) and the present paper, 48 genera 169 species of the superfamily Palaemonoidea have been reported based on the material deposited in the collections of the Muséum national d'Histoire naturelle, Paris, collected mainly from the Indo-Pacific areas. The appendix shows the geographical distribution of the 117 presently reported species.

Acknowledgements

We are most grateful to Alain Crosnier for providing the opportunity to study the material and for the numerous improvements to the draft manuscript, and writing the French abstract. Special thanks are due to him from both authors for his kind assistance during their visits to Paris. Thanks are due to B. Richer de Forges, who was the main collector of the present research samples from the New Caledonia area. J. C. Markham kindly identified the bopyrid parasites, Régis Cleva helped to collect some references, both are also greatly appreciated. This study was partly supported by the National Natural Science Foundation of China (no. 40276044 (XL)), and facilitated by support from the Australian Biological Resources Study (A.J.B.).

References

- Alcock A. 1901. A descriptive catalogue of the Indian deep-sea Crustacea Decapoda Macrura and Anomala, in the Indian Museum. Being a revised account of the deep-sea species collected by the Royal Indian Marine Survey Ship *Investigator*. Calcutta: Trustees of the Indian Museum, 286 p, 3 plates.
- Alcock A, Anderson AR. 1894. Natural history notes from H. M. Indian Marine Survey Steamer *Investigator*, Commander C. F. Oldham, R. N., commanding, Series II, 14: an account of a recent collection of deep sea Crustacea from the Bay of Bengal and Laccadive Sea. *Journal of the Asiatic Society of Bengal* 63:141–185. Plate 9.
- Audouin V. 1825. Explication sommaire des planches de Crustacés de l'Égypte et de la Syrie, publiées par Jules-César Savigny, membre de l'Institut: offrant un exposé des caractères naturels des genres avec la distinction des espèces. Description de l'Égypte ou recueil des observations et des recherches qui ont été faites en Égypte pendant l'expédition de l'armée française. *Histoire Naturelle* 1:77–98.
- Balss H. 1913. Diagnosen neuer ostasiatischer Macruren. *Zoologischer Anzeiger* 42:234–239.
- Balss H. 1914. Ostasiatische Decapoden II. Die Natantia und Reptantia. *Abhandlungen der Bayerischen Akademie der Wissenschaften Supplement* 2:1–101. Figures 1–50, Plate 1.
- Barnard KH. 1950. Descriptive catalogue of South African Decapoda Crustacea. *Annals of the South African Museum* 38:1–837. Figures 1–154.
- Barnard KH. 1962. New records of marine Crustacea from the East African Region. *Crustaceana* 3:239–245. Figures 1–3.
- Borradaile LA. 1898. A revision of the Pontoniidae. *Annals and Magazine of Natural History, Series 7* 2:376–391.
- Borradaile LA. 1899. On the Stomatopoda and Macrura brought by Dr. Willey from the South Seas. In: Willey A, editor. *Zoological results based on materials from New Britain, New Guinea, Loyalty Islands and elsewhere, collected during the years 1895, 1896 and 1897, part 4*. Cambridge: Cambridge University Press. p 395–428. Plates 36–39.
- Borradaile LA. 1915. Notes on Carides. *Annals and Magazine of Natural History, Series 8* 15:205–213.
- Borradaile LA. 1917. The Percy Sladen Trust Expedition to the Indian Ocean in 1905, under the leadership of Mr. J. Stanley Gardiner, M. A., Vol. 6, VIII: On the Pontoniinae. *The Transactions of the Linnean Society of London*, (2)17:323–396, pls. 52–57.
- Bray DM. 1976. A review of two Western Australian shrimps of the genus *Palaemonetes*, *P. australis* Dakin, 1915 and *P. atrinubes* sp. nov. (Decapoda, Palaemonidae). *Record of the Western Australian Museum* 4:65–84. Figures 1–42.
- Bruce AJ. 1965. Notes on some Indo-Pacific Pontoniinae, X. *Periclimenes cristimanus* sp. nov., a new pontoniinid shrimp from Singapore. *Annals and Magazine of Natural History, Series 13* 8:487–493. Figures 1, 2.
- Bruce AJ. 1966a. Notes on some Indo-Pacific Pontoniinae. XI. A re-examination of *Philarius lophos* Barnard, with the designation of a new genus *Ischnopontonia*. *Bulletin of Marine Science* 16:584–598. Figures 1–5.
- Bruce AJ. 1966b. *Periclimenes tosaensis* Kubo. Notes on some Indo-Pacific Pontoniinae, I. *Crustaceana* 10:15–22. Figures 1–4.
- Bruce AJ. 1966c. Notes on some Indo-Pacific Pontoniinae, II. *Platycaris latirostris* Holthuis. *Crustaceana* 11:1–9. Figures 1–5.
- Bruce AJ. 1967a. Notes on some Indo-Pacific Pontoniinae III–IX. Descriptions of some new genera and species from the Western Indian Ocean and the South China Sea. *Zoologischen Verhandlungen, Leiden* 87:1–73. Figures 1–29.

- Bruce AJ. 1967b. A report on some pontoniid shrimps from New Caledonia. (Crustacea Decapoda Natantia). Bulletin du Muséum National d'Histoire Naturelle, Paris, Série 2 39:1148–1171. Figures 1–10.
- Bruce AJ. 1969a. Observations upon the host specificity and distribution of *Jocaste japonica* (Ortmann) and *Jocaste lucina* (Nobili) (Decapoda Natantia, Pontoniinae). Crustaceana 17:298–302. Figures 1, 2.
- Bruce AJ. 1969b. Preliminary descriptions of sixteen new species of the genus *Periclimenes* Costa, 1844 (Crustacea, Decapoda Natantia, Pontoniinae). Zoologische Mededelingen, Leiden 43:253–278.
- Bruce AJ. 1970a. Report on some commensal pontoniid shrimps (Crustacea: Palaemonidae) associated with an Indo-Pacific gorgonian host (Coelenterata: Gorgonacea) Journal of Zoology, London 160:537–544. Figures 1–3.
- Bruce AJ. 1970b. Notes on some Indo-Pacific Pontoniinae, XV. *Hamopontonia corallicola* gen. nov., sp. nov., a new pontoniid shrimp from Hong Kong. Crustaceana 18:37–48. Figures 1–4.
- Bruce AJ. 1970c. Observations on the Indo-West Pacific species of the genus *Palaemonella* Dana, 1852 (Decapoda, Pontoniinae). Crustaceana 19:273–287. Figures 1–7, Plate 1.
- Bruce AJ. 1970d. On the identity of *Periclimenes pusillus* Rathbun, 1906 (Decapoda, Pontoniinae). Crustaceana 19:306–310. Figure 1.
- Bruce AJ. 1971a. *Palaemonella* Dana and *Periclimenes* Costa. Pontoniid shrimps from the ninth cruise of R/V Anton Bruun, IIOE, 1964, I. Smithsonian Contributions to Zoology 82:1–13. Figure 1.
- Bruce AJ. 1971b. *Periclimenes attenuatus* sp. nov. (Crustacea, Decapoda, Natantia, Pontoniinae), a new commensal shrimp from the Duke of York Islands. Pacific Science 25:533–544. Figures 1–5.
- Bruce AJ. 1972. Notes on Indo-Pacific Pontoniinae. XIX. *Allopontonia iaini* gen. nov., sp. nov., a new echinoid associate from Zanzibar (Decapoda, Caridea). Crustaceana 22:1–12. Figures 1–5.
- Bruce AJ. 1973a. Notes on some Indo-Pacific Pontoniinae. XXII. *Pliopontonia furtiva* gen. nov., sp. nov., a new shrimp associated with a coralliomorph zooantharian. Crustaceana 24:97–109. Figures 1–5, Plate 1.
- Bruce AJ. 1973b. Notes on some Indo-Pacific Pontoniinae. XXIV. *Dasycares zanzibarica* sp. nov., from the western Indian Ocean, with remarks on other species of *Dasycares* Kemp, 1922 (Decapoda, Natantia). Crustaceana 24:247–260. Figures 1–7.
- Bruce AJ. 1974a. *Coralliocaris viridis* sp. nov., a preliminary note (Decapoda Natantia, Pontoniinae). Crustaceana 26:222–224.
- Bruce AJ. 1974b. A report on a small collection of pontoniid shrimps from the Island of Farquhar. Crustaceana 27:189–203. Figures 1–8.
- Bruce AJ. 1975a. Notes on some Indo-Pacific Pontoniinae. XXV. Further observations upon *Periclimenes noverca* Kemp, with the designation of a new genus *Zenopontonia*, and some remarks upon *Periclimenes parasiticus* Borradaile. Crustaceana 28:275–285. Figures 1–3.
- Bruce AJ. 1975b. Notes on some Indo-Pacific Pontoniinae. XXVI. *Neoanchistus cardiodytes* gen. nov., sp. nov., a new mollusc-associated shrimp from Madagascar (Decapoda, Palaemonidae) Crustaceana 29:149–165. Figures 1–7.
- Bruce AJ. 1975c. Further observations on the Indo-West Pacific species of the genus *Palaemonella* Dana, 1852 (Decapoda Natantia, Pontoniinae). Crustaceana 29:169–185. Figures 1–7.
- Bruce AJ. 1976a. A report on some pontoniid shrimps collected from the Seychelles Islands by the F. R. V. Manihine, 1972, with a review of the Seychelles pontoniid shrimp fauna. Zoological Journal of the Linnean Society 59:89–153. Figures 1–30.
- Bruce AJ. 1976b. A synopsis of the pontoniid shrimp fauna of Central East Africa. Journal of the Marine Biological Association of India 16:462–490.
- Bruce AJ. 1976c. *Periclimenes soror* Nobili, a pontoniid shrimp new to the American fauna, with observations on its Indo-West Pacific distribution. Téthys 8:299–306. Figures 1–6.
- Bruce AJ. 1976d. Shrimps and prawns of coral reefs, with special reference to commensalism. In: Jones O, Endean R, editors. Biology and geology of coral. New York: Academic Press, Inc. Volume 3, Biology 2, 37–94. Figures 1–21, Table 1.
- Bruce AJ. 1977a. The possible identity of *Coralliocaris macrophthalmia* (H. Milne-Edwards, 1837) Decapoda Natantia, Pontoniinae. Crustaceana 32:203–205. Figure 1.
- Bruce AJ. 1977b. A report on a small collection of pontoniid shrimps from Queensland, Australia. Crustaceana 33:167–181. Figures 1–10.
- Bruce AJ. 1977c. Notes on some Indo-Pacific Pontoniinae, XXX. Some *Periclimenes* species from Madagascar (Decapoda Caridea). Crustaceana 33:265–274. Figures 1–5.
- Bruce AJ. 1977d. *Periclimenes kororensis* sp. nov., an unusual shrimp associate of the fungiid coral *Heliofungia actiniformis*. Micronesica 13:33–43. Figures 1–4.
- Bruce AJ. 1978a. A report on a collection of pontoniine shrimps from Madagascar and adjacent seas. Zoological Journal of the Linnean Society 62:205–290. Figures 1–44.

- Bruce AJ. 1978b. The re-examination of some pontoniine shrimp types first described by L. A. Borradaile. *Crustaceana* 34:251–268. Figures 1–9.
- Bruce AJ. 1979a. Records of some pontoniine shrimps from the South China Sea. *Cahiers de l'Indo-Pacifique* 1:215–248.
- Bruce AJ. 1979b. *Periclimes* *magnificus* sp. nov., a coelenterate associate from the Capricorn Islands (Decapoda, Palaemonidae). Notes on some Indo-Pacific Pontoniinae, XXXI. *Crustaceana Supplement* 5:195–208. Figures 1–6, Plate 1.
- Bruce AJ. 1980a. Some pontoniine shrimps from the Solomon Islands. *Micronesica* 16:261–269. Figures 1, 2.
- Bruce AJ. 1980b. On some pontoniine shrimps from Nounéa, New Caledonia. *Cahiers de l'Indo-Pacifique* 2:1–39. Figures 1–14.
- Bruce AJ. 1981a. Decapod Crustacea: Pontoniinae. In: Résultats des Campagnes MUSORSTOM, I: Philippines (18–28 Mars 1976). Mémoires ORSTOM 91:189–215. Figures 1–18.
- Bruce AJ. 1981b. Pontoniine shrimps from Viti Levu, Fijian Islands. *Micronesica* 17:77–95. Figures 1–11.
- Bruce AJ. 1982. The pontoniine shrimp fauna of Hong Kong. In: Morton BS, Tseng CK, editors, Proceedings of the First International Marine Biological Workshop: The Marine Flora and Fauna of Hong Kong and Southern China, Hong Kong, 1980. Hong Kong: Hong Kong University Press. p 233–284. Figures 1–26.
- Bruce AJ. 1983a. Additions to the marine fauna of the Northern Territory. 1. Decapod Crustacea: Caridea and Stenopodidea. The Beagle, Occasional Papers of the Northern Territory, Museum of Arts and Sciences 1:41–49.
- Bruce AJ. 1983b. Expédition Rumphius II (1975). Crustacés parasites, commensaux, etc. IX: Crustacés Décapodes (1ère partie: Natantia Pontoniinae). Bulletin du Muséum National d'Histoire Naturelle, Paris, Série 4 5A:871–902. Figures 1–10.
- Bruce AJ. 1983c. A note on the pontoniine shrimp fauna of La Réunion. *Bulletin of Marine Science* 33:165–166.
- Bruce AJ. 1983d. The pontoniine shrimps fauna of Australia. *Memoirs of the Australian Museum* 18:195–218.
- Bruce AJ. 1985a. Decapod Crustacea: Pontoniinae (MUSORSTOM II). In: Résultats des Campagnes MUSORSTOM, I et II. Philippines, Tome 2. Mémoires du Muséum National d'Histoire Naturelle 133:229–260. Figures 1–17.
- Bruce AJ. 1985b. Some caridean associates of scleractinian corals in the Ryukyu Islands. *Galaxea* 4:1–21. Figures 1–12, Plate 1.
- Bruce AJ. 1987. Notes on some Indo-Pacific Pontoniinae, XLIV. *Periclimes darwiniensis* sp. nov., from the Northern Territory, Australia (Decapoda, Caridea) *Crustaceana* 52:29–39. Figures 1–5.
- Bruce AJ. 1988a. A new palaemonid shrimp from the Zostera-beds of Moreton Bay, Queensland, Australia (Decapoda: Palaemonidae). The Beagle, Records of the Northern Territory Museum of Arts and Sciences 5:105–114. Figures 1–5.
- Bruce AJ. 1988b. The occurrence of *Palaemonetes atrinubes* Bray (Crustacea: Decapoda: Palaemonidae) on the Australian east coast. The Beagle, Records of the Northern Territory Museum of Arts and Sciences 5:115–117. Figure 1.
- Bruce AJ. 1988c. The shrimp fauna of a small tropical reef, the East Point Fish Reserve, Darwin. In: Larson HK, Michie MG, Hanley JR, editors. Darwin Harbour, proceedings of the workshop on Research and Management held in Darwin, 2–3 September, 1987, ANU North Australia Research Unit, Mangrove Monograph 4:226–245. Figure 1.
- Bruce AJ. 1989a. A report on some coral reef shrimps from the Philippine Islands. *Asian Marine Biology* 6:173–192. Figures 1–6.
- Bruce AJ. 1989b. Notes on some Indo-Pacific Pontoniinae, XLV. *Conchodytes maculatus* sp. nov., a new bivalve associate from the Australian Northwest Shelf. *Crustaceana* 56:182–192. Figures 1–6.
- Bruce AJ. 1990a. Crustacea Decapoda: deep-sea palaemonoid shrimps from New Caledonian waters. In: Crosnier A, editor. Résultats des Campagnes MUSORSTOM. Volume 6, Mémoires du Muséum National d'Histoire Naturelle, A, Zoologie 145:149–215. Figures 1–39.
- Bruce AJ. 1990b. *Periclimes franklini* sp. nov., a new deep-sea shrimp from the Coral Sea. The Beagle, Occasional Papers of the Northern Territory Museum of Arts and Sciences 7:55–64.
- Bruce AJ. 1990c. A new cnidarian-associated palaemonid shrimp from Port Essington, Cobourg Peninsula, Australia. *Indo-Malayan Zoology* 6(1989):229–243. Figures 1–8.
- Bruce AJ. 1991a. Crustacea Decapoda: further deep-sea palaemonid shrimps from New Caledonian waters. In: Crosnier A, editor. Résultats des Campagnes MUSORSTOM. Volume 9, Mémoires du Muséum National d'Histoire Naturelle, Série A, Zoologie 152:299–411. Figures 1–75.
- Bruce AJ. 1991b. Shallow-water palaemonoid shrimps from New Caledonia (Crustacea: Decapoda). In: Richer de Forges B, editor. Le benthos des fonds meubles des lagons de Nouvelle-Calédonie, 1. Études et Thèses. Paris: ORSTOM. 211–279. Figures 1–31.

- Bruce AJ. 1992. Two new species of *Periclimenes* (Crustacea: Decapoda: Palaemonidae) from Lizard Island, Queensland, with notes on some related taxa. *Records of the Australian Museum* 44:45–84. Figures 1–27.
- Bruce AJ. 1994. A synopsis of the Indo-West Pacific genera of the Pontoniinae (Crustacea: Decapoda: Palaemonidae). Königstein: Koeltz Scientific Books. 172 p.
- Bruce AJ. 1995. A re-examination of *Palaemonetes sinensis* (Sollaud, 1911) (Crustacea: Decapoda: Palaemonidae). The Beagle, Records of the Museums and Art Galleries of the Northern Territories 11:1–7. Figures 1–3.
- Bruce AJ. 1996. Crustacea Decapoda: palaemonid shrimps from the Indo-West Pacific region, mainly from New Caledonia. In: Crosnier A, editor. *Résultats des Campagnes MUSORSTOM, 15. Mémoires du Muséum National d'Histoire Naturelle* 168:197–267. Figures 1–31.
- Bruce AJ. 1998. New keys for the identification of Indo-West Pacific coral associated pontoniine shrimps, with observations on their ecology [Crustacea: Decapoda: Palaemonidae]. *Ophelia* 49:29–46.
- Bruce AJ. 2002a. *Leander manningi*, a new palaemonine shrimp from Western Australia (Crustacea, Decapoda, Palaemonidae), with a review of the Indo-West Pacific species of the genus *Leander* E. Desmarest, 1849. *Records of the Western Australian Museum* 21:71–81. Figures 1–6.
- Bruce AJ. 2002b. Notes on some Indo-Pacific Pontoniinae, XLVI. *Palaemonella foresti* sp. nov., a new pontoniine shrimp from Western Australia (Decapoda, Palaemonidae), with a review of the Indo-West Pacific species of the genus *Palaemonella* Dana, 1852. *Crustaceana* 75:277–298. Figures 1–4.
- Bruce AJ. 2003. The pontoniine shrimp fauna of Hong Kong and the South China Sea (Crustacea: Decapoda: Palaemonidae). In: Morton B, editor. *Perspectives on marine environment change in Hong Kong and southern China, 1977–2001. Proceedings of an international workshop reunion conference, Hong Kong, 21–26 October 2001, Hong Kong: Hong Kong University Press.* p 209–257. Table 1.
- Bruce AJ. 2004. A partial revision of the genus *Periclimenes* Costa, 1884 (Crustacea: Decapoda: Palaemonidae). *Zootaxa* 582:1–26.
- Bruce AJ. 2005. Pontoniine shrimps from the 2003 NORFANZ Expedition, 10 May–16 June [Crustacea: Decapoda: Palaemonidae]. *Zootaxa* 981:1–20. Figures 1–8.
- Bruce AJ, Coombes KE. 1995. The palaemonoid shrimp fauna (Crustacea: Decapoda: Caridea) of the Cobourg Peninsula, Northern Territory. The Beagle, Occasional Papers of the Northern Territory, Museum of Arts and Sciences 12:101–144. Figures 1–12.
- Bruce AJ, Okuno J, Li X. 2005. *Manipontonia* gen. nov., a new pontoniine shrimp genus for *Periclimenes psamathe* (De Man) [Crustacea: Decapoda: Palaemonidae]. *Zootaxa* 926:1–11. Figures 1–3.
- Bruce AJ, Svoboda A. 1984. A report on a small collection of coelenterate associated pontoniine shrimps from Cebu, Philippine Islands. *Asian Marine Biology* 1:87–99. Figures 1–7.
- Bruce AJ, Zmarzly DL. 1983. *Periclimenes pilipes*, new species, a crinoid associate from Eniwetak Atoll, Marshall Islands [Crustacea: Decapoda: Pontoniinae]. *Journal of Crustacean Biology* 3:644–654. Figures 1–6.
- Chace FA Jr, Bruce AJ. 1993. The caridean shrimps (Crustacea: Decapoda) of the *Albatross* Philippine Expedition 1907–1910, part 6: Superfamily Palaemonoidea. *Smithsonian Contributions to Zoology* 543:1–152. Figures 1–23.
- Dana JD. 1852. *Conspectus of the Crustacea of the exploring expedition under Capt. C. Wilkes U. S. N.* *Proceedings of the Academy of Natural Sciences of Philadelphia* 1852:10–28.
- Davie PJF. 2002. Crustacea: Malacostraca: Phyllocarida, Hoplocarida, Eucarida (part 1). In: Wells A, Houston WWK, editors. *Zoological catalogue of Australia. Volume 19.3A.* Melbourne: CRIRO Publishing. 551 p.
- De Grave S. 2000. Caridean shrimps (Crustacea, Decapoda) from Hansa Bay, Papua New Guinea: Palaemonidae and Gnathophyllidae. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Biologie* 70:119–148.
- De Man JG. 1888. Report on the podophthalmous Crustacea of the Mergui Archipelago collected for the Trustees of Indian Museum Calcutta by John Anderson F. R. S. Superintendent of the Museum. *Journal of the Linnean Society of London* 22:1–312. Plates 1–19.
- De Man JG. 1902. Die von Herrn Professor Kükenthal in Indischen Archipel gesammelten Dekapoden und Stomatopoden. In: Kükenthal W, editor. *Ergebnisse einer zoologischen Forschungsreise in den Molukken un Borneo, Abhandlungen der Senckenbergischen und Naturforschenden Gesellschaft* 25:467–929. Plates 19–27.
- Debelius H. 1999. Crustacea, guide of the world. Frankfurt: IKAN-Unterwasserarchiv. 321 p.
- Duris Z, Bruce AJ. 1995. A revision of the “*petithouarsi*” species-group of the genus *Periclimenes* Costa, 1844 (Crustacea: Decapoda: Palaemonidae). *Journal of Natural History* 29:619–671. Figures 1–22.
- Forskål P. 1775. *Descriptiones Animalium, Avium, Amphibiorum, Piscium, Insectorum, Vermium; quae in Itinere Orientali Observavit.* Haunia: Heineck et Faber. 164 p.
- Fransen CHJM. 1987. Notes on caridean shrimps of Easter Island with descriptions of three new species. *Zoologische Mededelingen, Leiden* 61:501–531. Figures 1–16.

- Fransen CHJM. 1989. Notes on caridean shrimps collected during the Snellius-II Expedition, I: associates of Anthozoa. *Netherlands Journal of Sea Research* 23:131–147. Figure 1–9.
- Fransen CHJM. 1994. Marine palaemonoid shrimps of the Netherlands Seychelles Expedition 1992–1993. *Zoologische Verhandelingen, Leiden* 297:85–152. Figures 1–112, Plates 1–4.
- Fujino T, Miyake S. 1969. On two new species of Palaemonid shrimps from Tanabe Bay, Kii Peninsula, Japan (Crustacea, Decapoda, Palaemonidae). *Publications of the Seto Marine Biological Laboratory* 17:143–154. Figures 1–5.
- Fujino T, Miyake S. 1970. Caridean and stenopodidean shrimps from the East China and the Yellow Seas (Crustacea, Decapoda, Natantia). *Journal of the Faculty of Agriculture, Kyushu University* 16:237–312. Figures 1–25.
- Gordon I. 1935. On new and imperfectly known species of Crustacea Macrura. *Journal of the Linnean Society of London* 39:307–351. Figures 1–27.
- Guérin-Ménéville FE. 1838. Crustacés, Arachnides et Insectes. In: Duperrey LJ, Voyage a l'author du monde, exécuté par Ordre du Roi, sur la Corvette de Sa Majesté, La Coquille, pendant les années 1822, 1823, 1824 et 1825. *Zoologie* 2 (no. 2, sect. 1): 1–47 (Crustacés), 48–319 (Arachnides et Insectes); plates (1–5 (Crustacés), 1–21 (Insectes). Paris: Arthus Bertrand.
- Guérin-Ménéville FE. 1855. Crustaceos. In: La Sagra: Historia Fisica Politica y Natural de la Isla de Cuba, Historia Natural. Volume 7 (atlas). Paris: Arthus Bertrand. 88 p, 3 plates.
- Hayashi K. 1986. Shrimps. In: Baba K, Hayashi K, Toriyama M, editors. Decapod crustaceans from continental shelf and slope around Japan. Tokyo: Tosho Printing Co. Ltd.. p 335. 22 text-figures, 176 photographs.
- Heller C. 1861. Synopsis der in rothen Meere vorkommenden Crustaceen. *Verhandlungen des Kaiserlich-königlichen Zoologisch-Botanischen Gesellschaft in Wien* 11:3–32.
- Henderson JR. 1893. A contribution to Indian carcinology. *Transactions of the Linnean Society of London, Series 2* 5:325–458. Plates 36–40.
- Hipeau-Jacquotte R. 1973. Étude des crevettes Pontoniinae (Palaemonidae) associées aux mollusques Pinnidae à Tuléar (Madagascar). 3. Morphologie externe et morphologie des pièces buccales. *Téthys Supplément* 1973:95–116.
- Hipeau-Jacquotte R. 1974. Étude des crevettes Pontoniinae (Palaemonidae) associées aux molluscs Pinnidae à Tuléar (Madagascar). 6. Comportement sexuel. *Téthys* 5:403–408. Figure 1.
- Holthuis LB. 1950. The Decapoda of the Siboga Expedition, part X: the Palaemonidae collected by the Siboga and Snellius Expeditions, with remarks on other species, part I: Subfamily Palaemoninae. *Siboga-Expeditie* 39a:1–268. Figures 1–52.
- Holthuis LB. 1951. The Subfamilies Euryrhynchinae and Pontoniinae. A general revision of the Palaemonidae (Crustacea Decapoda Natantia) of the Americas. II. *Allan Hancock Foundation Publication, Occasional Paper* 11:1–332. Plates 1–63.
- Holthuis LB. 1952a. The Decapoda of the Siboga Expedition, part XI: the Palaemonidae collected by the Siboga and Snellius Expeditions, with remarks on other species, part II: Subfamily Pontoniinae. *Siboga-Expeditie* 39a:1–254. Figures 1–110.
- Holthuis LB. 1952b. The subfamily Palaemoninae, part II. A general revision of the Palaemonidae (Crustacea Decapoda Natantia) of the Americas. *Allan Hancock Foundation Occasional Papers* 12:1–396. Figure 1, Plates 1–55.
- Holthuis LB. 1958. Contributions to the knowledge of the Red Sea, 8. Crustacea Decapoda from the northern Red Sea (Gulf of Aqaba and Sinai Peninsula). I. Macrura. *Bulletin of the Sea Fisheries Research Station (Haifa, Israel)* 17:1–40. Figures 1–15.
- Holthuis LB. 1959. Results of the re-examination of the type specimens of some species belonging to the subfamilies Pontoniinae and Palaemoninae (Crustacea Decapoda Macrura). *Zoologische Mededelingen, Leiden* 36:193–200. Figure 1.
- Holthuis LB. 1980. Shrimps and prawns of the world. An annotated catalogue of species of interest to fisheries. *FAO Fisheries Synopsis* 125:i–xvii, 1–231.
- Jacquotte R. 1963. Habitat electif des Pontoniinae commensales des Pinnidae de Tulear (Madagascar). *Recueil des Travaux de la Station Marine d'Endoume* 29:60–62. Figure 1.
- Kemp S. 1922. Notes on Crustacea Decapoda in the Indian Museum, XV: Pontoniinae. *Records of the Indian Museum* 24:113–288. Figures 1–105, Plates 3–9.
- Kemp S. 1925. Notes on Crustacea Decapoda in the Indian Museum, XVII: on various Caridea. *Records of the Indian Museum* 27:249–343. Figures 1–24.
- Kingsley JS. 1878. List of the North American Crustacea belonging to the sub-order Caridea. *Bulletin of the Essex Institute* 10:53–71.
- Kubo I. 1940a. A new shrimp, *Harpilius imperialis*. *Journal of the Imperial Fisheries Institute* 34:1–4. Figures 1–3.

- Kubo I. 1940b. Studies on Japanese palaemonoid shrimps, II: Pontoniinae. *Journal of the Imperial Fisheries Institute* 34:31–75. Figures 1–36.
- Kubo I. 1951. Some macrurous decapod Crustacea found in Japanese waters, with descriptions of four new species. *Journal of the Tokyo University of Fisheries* 38:259–289. Figures 1–16.
- Ledoyer M. 1984. Les Caridea (Crustacea: Decapoda) des herbiers de Phanérogames marines de Nouvelle-Calédonie (Région de Nouméa). *Zoologische Verhandelingen, Leiden* 211:1–58. Figures 1–21, Tables 1–5.
- Li X. 1996. The pontoniine shrimps from Nansha Islands, China. I. *Marine Fauna and Flora and Biogeography of the Nansha Islands and Neighbouring Waters* 2:222–233. Figures 1–9.
- Li X. 1997. Report on Gnathophyllidae and Pontoniinae (Decapoda, Palaemonoidea) shrimps from the Xisha Islands and adjacent waters, South China Sea. *Studia Marina Sinica (Qingdao)* 38:223–251. Figures 1–12.
- Li X. 1998. The pontoniine shrimps (Crustacea: Caridea: Palaemonoidea) from Nansha Islands, South China Sea. II. *Studies on Marine Fauna and Flora and Biogeography of the Nansha Islands and Neighbouring Waters* 3:218–226. Figures 1–9.
- Li X. 2000. Catalog of the genera and species of Pontoniinae Kingsley, 1878. Beijing: Xueyuan Press. 319. 408 figures.
- Li X. 2001. On some pontoniine shrimps (Crustacea: Caridea) from Hainan Island, South China Sea. In: Matsuura K, editor. *Marine fauna of the shallow waters around Hainan Island, South China Sea*. National Science Museum Monographs 21:75–86.
- Li X. 2004a. The pontoniine shrimps (Crustacea, Decapoda, Palaemonidae) from Anambas and Natuna Islands, Indonesia, collected by Anambas Expedition, 2002. *The Raffles Bulletin of Zoology Supplement* 11:67–72. Figures 1, 2.
- Li X. 2004b. Report on some pandalid and pontoniine shrimps from the Chinese Seas (Crustacea, Decapoda, Caridea). *Acta Zootaxonomica Sinica* 29:820–826. Figures 1–9.
- Li X, Bruce AJ, Manning RB. 2004. Some palaemonid shrimps (Crustacea: Decapoda) from Northern South China Sea, with descriptions of two new species. *The Raffles Bulletin of Zoology* 52:513–553. Figures 1–33.
- Li X, Liu JY. 2002. Report of pontoniine shrimps (Crustacea: Decapoda) collected by the joint Chinese–German marine biology expeditions to Hainan Island, South China Sea. I. *Anchistus, Conchodytes, Coralliocaris*. *Chinese Journal of Oceanology and Limnology* 20:371–377. Figures 1–4.
- Li X, Liu JY. 2003. Report of pontoniine shrimps (Crustacea: Decapoda) collected by the joint Chinese–German marine biology expeditions to Hainan Island, South China Sea. II. *Harpiliopsis, Ischnopontonia, Jocaste, Palaemonella, Periclimenaeus, Periclimenella*. *Chinese Journal of Oceanology and Limnology* 21:154–165. Figures 1–8.
- Li X, Liu JY. 2004. Report of pontoniine shrimps (Crustacea: Decapoda) collected by the joint Chinese–German marine biology expeditions to Hainan Island, South China Sea. III. *Periclimenes*. *Chinese Journal of Oceanology and Limnology* 22:89–100. Figures 1–8.
- Lucas H. 1846. Crustacés, Arachnides, Myriapodes et Hexapodes. *Exploration Scientifique de l'Algérie pendant les années 1840, 1841, 1842. Sciences physiques. Zoologie I, Histoire Naturelle des Animaux Articulés* 1:1–403. Plates 1–8.
- Marin IN, Britayev TA, Anker A. 2004. Pontoniine shrimps associated with cnidarians: new records and list of species from coastal waters of Viet Nam. *Arthropoda Selecta* 13:199–218. Figures 1–13.
- Markham JC. 1989. Three species of Isopoda Bopyridae new to the fauna of the Philippines. *The Beagle, Records of the Northern Territory Museum of Arts and Sciences* 6:141–148.
- Miers EJ. 1884. Crustacea. In: *Report of the zoological collections made in the Indo-Pacific Ocean during the voyage of H.M.S. "Alert" 1881–2*. London: British Museum (Natural History). p 178–322, 513–575. Plates 18–35, 46–52.
- Milne Edwards H. 1837. *Histoire naturelle des Crustacés, comprenant l'anatomie, la physiologie et la classification de ces animaux*. Volume 2. Paris: Roret. 532 p.
- Minemizu R, Okuno J, Takeda M. 2000. Marine decapod and stomatopod crustaceans mainly from Japan. Tokyo: Bun-Ichi Sogo Shuppan Co.. 344 p.
- Mitsuhashi M. 2000. A revision of the genus *Coralliocaris* Stimpson, 1860 (Crustacea: Decapoda: Palaemonidae) from Japan. *IOP Diving News* 11:2–7. Figures 1–11.
- Miyake S, Fujino T. 1968. Pontoniinid shrimps from the Palau Islands (Crustacea, Decapoda, Palaemonidae). *Journal of the Faculty of Agriculture, Kyushu University* 14:399–431. Figures 1–8.
- Monod T. 1976. Sur une nouvelle collection de Crustacés Décapodes de Nouméa (Nouvelle-Calédonie). *Cahiers du Pacifique* 19:133–152. Figures 1–82.
- Nobili G. 1901. Decapodi e Stomatopodi Eritrei del Museo Zoologico dell'Università di Napoli. *Annuario del Museo Zoologico della R. Università di Napoli, New Series* 1:1–20.

- Nobili G. 1904. Diagnoses préliminaires de vingt-huit espèces nouvelles de Stomatopodes et Décapodes Macroures de la mer Rouge. Bulletin de Muséum d'Histoire Naturelle, Paris 10:228–238.
- Nobili G. 1906a. Diagnoses préliminaires de Crustacés. Décapodes et Isopodes nouveaux recueillis par M. le Dr. G. Seurat aux Îles Touamotou. Bulletin du Muséum d'Histoire Naturelle 12:256–270.
- Nobili G. 1906b. Crustacés Décapodes et Stomatopodes: Mission J. Bonnier et Ch. Perez (Golfe Persique 1901). Bulletin Scientifique de la France et de la Belgique 40:13–159. Figures 1–3, Plates 2–7.
- Nobili G. 1907. Ricerche sui Crostacei della Polinesia. Decapodi, Stomatopodi, Anisopodi e Isopodi. Memorie della Reale Accademia della Scienze di Torino, Serie 2 57:351–430. Plates 1–3.
- Okuno J. 1999. *Palaemonella hachijo*, a new species of shrimp (Crustacea: Decapoda: Palaemonidae) from a submarine cave in southern Japan. Proceedings of the Biological Society of Washington 112:739–745. Figures 1–3.
- Okuno J. 2000. Additional specimens of *Palaemonella hachijo* Okuno, 1999 (Decapoda, Caridea, Palaemonidae). Crustaceana 73:1297–1299. Figure 1.
- Ortmann AE. 1890. Die Unterordnung Natantia Boas. Die Decapoden-Krebse des Strassburger Museums, mit besonderer Berücksichtigung der von Herrn Dr. Döderlein bei Japan und bei der Liu-Kiu-Inseln gesammelten und z. Z. in Strassburger Museum aufbewahrten Formen, I. Zoologische Jahrbucher Abteilung für Systematik, Geographie und Biologie der Thiere 5:437–542. Plates 36, 37.
- Patton WK. 1966. Decapod Crustacea commensal with Queensland branching corals. Crustaceana 10:271–295. Figures 1–3.
- Paulson O. 1875. Investigations on the Crustacea of the Red Sea with notes on Crustacea of the adjacent seas. Part I. Podophthalmata and Edriophthalmata (Cumacea). Kiev: SV Kul'zhenko, 144 p. 21 plates.
- Peters W. 1852. *Conchodytes*, eine neue in Muscheln lebende Gattung von Garneelen. Berichte über die zur Bekanntmachung geeigneten Verhandlungen der K. Preuss. Akademie der Wissenschaften zu Berlin 1852:588–595.
- Rathbun MJ. 1906. The Brachyura and Macrura of the Hawaiian Islands. Bulletin of the United States Fish Commission (1903) 23:827–930. Plates 1–24.
- Say T. 1817–18. An account of the Crustacea of the United States. Journal of the Academy of Natural Sciences of Philadelphia 1(1817):57–80, 97–101, 151–169, Plate 4; (1818):235–253, 313–319, 374–401, 423, 441, 445–458.
- Schenkel E. 1902. Beitrag zur Kenntnis der Dekapodenfauna von Celebes. Verhandlungen der Naturforschenden Gesellschaft in Basel 13:485–585. Plates 7–13.
- Stimpson W. 1860. Prodromus descriptionis animalium evertibratorum, quae in expeditione ad Oceanum Pacificum septentrionalem, a Republica Federata missa Cadwaladaro Ringgold et Johanne Rodgers Ducibus, observavit et descripsit. Proceedings of the Academy of Natural Sciences of Philadelphia 1860:22–48.
- Wicksten MK, Hendrickx ME. 1985. New records of caridean shrimps in the Gulf of California, Mexico. Proceedings of the Biological Society of Washington 98:571–573.
- Yokoya Y. 1936. Some rare and new species of decapod crustaceans found in the vicinity of the Misaki Marine Biological Station. Japanese Journal of Zoology 7:129–146. Figures 1–4.
- Zehntner L. 1894. Crustacés de l'Archipel Malais. Voyage de MM. M. Bedot et C. Pictet dans l'Archipel Malais. Revue Suisse de Zoologie et Annales du Musée d'Histoire Naturelle de Genève 2:135–214. Plates 7–9.

Appendix. Geographical distribution of presently reported species

	Comoro Islands	Madagascar	La Réunion	Île Europa	Philippines	Indonesia	Solomon Islands	Vanuatu	New Caledonia	Loyalty Islands	Fiji.	Tuvalu	Tonga
ANCHISTIOIDIDAE													
<i>Anchistioides willeyi</i>	+	+			+				+		+		
GNATHOPHYLLIDAE													
<i>Gnathophyllum americanum</i>									+	+			
PALAEMONIDAE													
PALAEMONINAE													
<i>Brachycarpus biunguiculatus</i>			+						+				
<i>Leander tenuicornis</i>									+	+			
<i>Macrobrachium australe</i>									+				
<i>Macrobrachium equidens</i>									+				
<i>Nematopalaemon tenuipes</i>		+				+							
<i>Palaemon debilis</i>				+					+				
<i>Palaemonetes atrimubes</i>									+				
<i>Urocaridella antonbruunii</i>			+						+				
<i>Urocaridella urocaridella</i>					+		+		+				
PONTONIINAE													
<i>Allopontonia iaini</i>									+				
<i>Altopontonia disparostris</i>												+	
<i>Amphipontonia kanak</i>									+				
<i>Anchistus custos</i>									+				
<i>Anchistus miersi</i>		+											
<i>Anchistus pectinis</i>									+				
<i>Climeniperæus truncoideus</i>					+								
<i>Conchodytes biunguiculatus</i>		+											
<i>Conchodytes maculatus</i>									+				
<i>Conchodytes meleagrinae</i>				+					+	+			
<i>Coralliocaris</i> sp.											+		
<i>Coralliocaris macrophthalma</i>									+				
<i>Coralliocaris superba</i>		+							+				
<i>Coralliocaris viridis</i>		+											
<i>Dasycaris ceratops</i>		+											
<i>Dasycaris symbiotes</i>									+				
<i>Dasycaris zanzibarica</i>		+											
<i>Exoclimenella denticulate</i>											+		
<i>Exoclimenella</i> cf. <i>denticulate</i>											+		
<i>Exoclimenella maldivensis</i>											+		
<i>Hamodactylus boschmai</i>									+				
<i>Hamodactylus noumeae</i>									+				
<i>Hamopontonia corallicola</i>					+								
<i>Harpiliopsis beaupresii</i>		+	+								+		
<i>Harpiliopsis depressa</i>									+				
<i>Harpiliopsis spinigera</i>									+				
<i>Harpilius lutescens</i>											+		
<i>Ischnopontonia lophos</i>		+											
<i>Jocaste japonica</i>											+		
<i>Jocaste lucina</i>											+		
<i>Kemponia agag</i>									+				
<i>Kemponia amymone</i>									+				

	Comoro Islands	Madagascar	La Réunion	Île Europa	Philippines	Indonesia	Solomon Islands	Vanuatu	New Caledonia	Loyalty Islands	Fiji	Tuvalu	Tonga
<i>Kemponia anacanthus</i>									+	+			
<i>Kemponia andamanensis</i>												+	
<i>Kemponia darwiniensis</i>									+				
<i>Kemponia elegans</i>												+	
<i>Kemponia ensifrons</i>									+				
<i>Kemponia grandis</i>									+	+			
<i>Kemponia kororensis</i>					+								
<i>Kemponia lacertae</i>										+			
<i>Kemponia nilandensis</i>			+						+				
<i>Kemponia seychellensis</i>									+				
<i>Kemponia cf. suvadiensis</i>									+				
<i>Kemponia tenuipes</i>									+	+	+		
<i>Manipontonia psamathe</i>		+							+				
Mesopontonia brevicarpalis			+										
<i>Mesopontonia gracilicarpus</i>									+				
<i>Metapontonia fungiacola</i>									+				
<i>Palaemonella crosnieri</i>										+			
<i>Palaemonella dolichodactylus</i>											+		
<i>Palaemonella hachijo</i>									+				
Palaemonella komaii											+		+
<i>Palaemonella potsi</i>					+				+	+			
<i>Palaemonella pusilla</i>									+	+			
<i>Palaemonella rotumana</i>									+	+			
<i>Palaemonella spinulata</i>									+	+			
<i>Paraclimenes franklinae</i>								+					+
<i>Paranchistus nobilii</i>									+				
<i>Paranchistus ornatus</i>		+											
<i>Periclimenella petithouarsi</i>		+											
<i>Periclimenella spinifera</i>									+	+			
<i>Periclimenes affinis</i>					+				+				
<i>Periclimenes alcocki</i>					+	+	+		+		+		+
<i>Periclimenes aleator</i>						+	+	+	+	+	+		
<i>Periclimenes amboinensis</i>					+				+	+			
<i>Periclimenes attenuatus</i>					+								
<i>Periclimenes brevicarpalis</i>		+							+	+			
<i>Periclimenes brevirostris</i>								+	+				
<i>Periclimenes calcaratus</i>					+								
<i>Periclimenes ceratophthalmus</i>									+	+			
<i>Periclimenes commensalis</i>					+				+				
<i>Periclimenes cristimanus</i>					+								
Periclimenes crosnieri						+							
<i>Periclimenes forcipulatus</i>							+						
Periclimenes forgesi									+				
<i>Periclimenes foveolatus</i>					+								
<i>Periclimenes hertwigi</i>					+	+	+		+				
<i>Periclimenes imperator</i>		+							+				
<i>Periclimenes incertus</i>		+					+		+				
<i>Periclimenes involens</i>			+										
<i>Periclimenes kemp</i> Bruce, 1969									+				
<i>Periclimenes laccadivensis</i>									+				
<i>Periclimenes lanipes</i>					+						+		

	Comoro Islands	Madagascar	La Réunion	Île Europa	Philippines	Indonesia	Solomon Islands	Vanuatu	New Caledonia	Loyalty Islands	Fiji.	Tuvalu	Tonga
<i>Periclimenes latipollex</i>									+				
<i>Periclimenes lepidus</i>		+							+				
<i>Periclimenes loyautensis</i>										+			
<i>Periclimenes magnificus</i>									+				
<i>Periclimenes macrophthalmus</i>											+		
<i>Periclimenes novaecaledoniae</i>									+				
<i>Periclimenes paralcocki</i>												+	
<i>Periclimenes paraleator</i>								+					
<i>Periclimenes pilipes</i>					+								
<i>Periclimenes platyrhynchus</i>									+				
<i>Periclimenes pseudalcocki</i>						+							
<i>Periclimenes soror</i>									+	+			
<i>Periclimenes tosaensis</i>											+		
<i>Periclimenes uniunguiculatus</i>	+								+				
<i>Periclimenes vaubani</i>									+				
<i>Periclimenes venustus</i>									+				
<i>Periclimenes</i> sp.			+										
<i>Philarius imperialis</i>									+	+			
<i>Platycaris latirostris</i>		+											
<i>Pliopontonia furtiva</i>									+				
<i>Pontoniopsis comanthi</i>					+				+				
<i>Thaumastocaris streptopus</i>									+				
<i>Zenopontonia noverca</i>		+							+				