



GABAergic signaling as therapeutic target for autism spectrum disorders

Giada Cellot¹ and Enrico Cherubini^{1,2*}

¹ Department of Neuroscience, Scuola Internazionale Superiore di Studi Avanzati, Trieste, Italy

² European Brain Research Institute, Rome, Italy

Edited by:

Yuri Bozzi, University of Trento, Italy

Reviewed by:

Yehezkel Ben-Ari, Institut National de la Santé et de la Recherche Médicale, France

Laura Cancedda, Istituto Italiano di Tecnologia, Italy

*Correspondence:

Enrico Cherubini, Department of Neuroscience, Scuola Internazionale Superiore di Studi Avanzati (SISSA), Via Bonomea 265, Trieste 34136, Italy
e-mail: cher@siissa.it

γ -Aminobutyric acid (GABA), the main inhibitory neurotransmitter in the adult brain, early in postnatal life exerts a depolarizing and excitatory action. This depends on accumulation of chloride inside the cell via the cation–chloride importer NKCC1, being the expression of the chloride exporter KCC2 very low at birth. The developmentally regulated expression of KCC2 results in extrusion of chloride with age and a shift of GABA from the depolarizing to the hyperpolarizing direction. The depolarizing action of GABA leads to intracellular calcium rise through voltage-dependent calcium channels and/or *N*-methyl-D-aspartate receptors. GABA-mediated calcium signals regulate a variety of developmental processes from cell proliferation migration, differentiation, synapse maturation, and neuronal wiring. Therefore, it is not surprising that some forms of neuro-developmental disorders such as autism spectrum disorders (ASDs) are associated with alterations of GABAergic signaling and impairment of the excitatory/inhibitory balance in selective neuronal circuits. In this review, we will discuss how changes of GABA_A-mediated neurotransmission affect several forms of ASDs including the Fragile X, the Angelman, and Rett syndromes. Then, we will describe various animal models of ASDs with GABAergic dysfunctions, highlighting their behavioral deficits and the possibility to rescue them by targeting selective components of the GABAergic synapse. In particular, we will discuss how in some cases, reverting the polarity of GABA responses from the depolarizing to the hyperpolarizing direction with the diuretic bumetanide, a selective blocker of NKCC1, may have beneficial effects on ASDs, thus opening new therapeutic perspectives for the treatment of these devastating disorders.

Keywords: autism spectrum disorders, GABA receptors, bumetanide, neuro-developmental disorders, excitatory inhibitory balance

Autism comprises a heterogeneous group of neuro-developmental disorders known as autism spectrum disorders (ASDs) characterized by deficits in verbal and non-verbal communication, social interaction, restricted interests, and stereotyped behavior (1). The incidence of ASDs (20–60/10000 children) has dramatically increased over the past decades because of the improvement of diagnostic criteria and increased attention of medical community (2). Complications arising from later age pregnancies and from excessive exposure of fetuses with a genetic vulnerable background to environmental factors (i.e., toxic agents) may also contribute to the higher incidence of these disorders in recent years.

In spite different etiologies, ASDs share overlapping symptoms, indicating common deficits in some neuro-developmental pathways. One of these involves the γ -aminobutyric acid (GABA)_A-mediated neurotransmission, known to play a crucial role in synaptic tuning and neuronal wiring in late pre and early postnatal days (3). Studies from animal models of ASDs indicate that a dysfunction in GABAergic signaling within particular neuronal circuits may account for most of the clinical symptoms found in autistic patients. The high co-morbidity of ASDs with epilepsy (30% of cases) further confirms this issue (4).

γ -Aminobutyric acid is the main inhibitory neurotransmitter in the adult mammalian brain. It inhibits neuronal firing by

activating two different classes of receptors, GABA_A and GABA_B. GABA_A receptors are integral ion channels while GABA_B receptors are coupled to ion channels via guanine nucleotide-binding proteins and second messengers. The opening of GABA_A receptors causes a net influx of chloride with consequent membrane hyperpolarization and reduction of cell firing. However, in particular conditions and during brain maturation the intracellular chloride concentration $[Cl^-]_i$ rises in such a way that the opening of anion channels by GABA produces a chloride efflux and a membrane depolarization that through the activation of a persistent non-inactivating sodium conductance (5) may reach the threshold for action potential generation (6, 7). Generally, low $[Cl^-]_i$ facilitates GABA-mediated inhibition, whereas high $[Cl^-]_i$ facilitates GABA-mediated excitation. The mechanisms underlying chloride accumulation inside immature neurons, start to be unveiled with a different efficacy of chloride co-transporters such as NKCC1 and KCC2, which import and export chloride, respectively. Before and immediately after birth, chloride accumulates inside the cell due to a reduced expression of the cation–chloride exporter KCC2. Later in development, the intracellular chloride concentration decreases thanks to the up-regulation of KCC2 (8–10). GABA-induced membrane depolarization facilitates calcium entry via voltage-dependent calcium channels and *N*-methyl-D-aspartate

(NMDA) receptors. Calcium rise leads to the activation of second messengers involved in a variety of developmental processes from cell migration and differentiation to synaptogenesis and circuit formation (11).

How GABA orchestrates these processes has been extensively reported (3, 12–14). GABAergic signals operate with multiple modalities at different developmental stages before glutamatergic ones (15). At the beginning, GABA works as a trophic factor, modulating neuronal migration and maturation (16). GABA receptors are expressed in neuronal progenitors before the establishment of synaptic contacts (17). At this stage, the receptors work as sensors for GABA present in the extracellular space after its release in a calcium- and SNARE-independent way from growth cones and astrocytes (18). The absence of an efficient uptake system enables this neurotransmitter to accumulate in the extracellular space and to reach a concentration sufficient to exert its depolarizing and excitatory effects on distal neurons. Blocking the depolarizing action *in utero* heavily affects migration and circuit formation (11, 19, 20).

At later developmental stages, when synapses are formed, the release of GABA and glutamate, generate a primitive form of network-driven oscillatory events known as giant depolarizing potentials (GDPs). GDPs are characterized by recurrent membrane depolarizations (lasting several hundred of milliseconds) that give rise to bursts of action potentials, separated by quiescent periods. This network activity thought to be the *in vitro* counterpart of “sharp waves” recorded in pups during immobility periods, sleep, and feeding (21), is reminiscent of the “trace discontinue” first described by Dreyfus-Brisac in the electroencephalogram of immature babies and characterized by intermittent bursts separated by periods of virtually complete suppression of activity (22).

In analogy with the synchronized activity generated in the disinhibited hippocampus by GABA_A receptor antagonists (23), GDPs emerge when a sufficient number of cells fire and the excitability of the network attains a certain threshold within a restricted temporal window (24). Although the entire hippocampus possesses the capacity to generate GDPs, for its extensive glutamatergic connections via recurrent collaterals, the CA3 area is particularly well equipped to generate synchronized activity. Furthermore, this area is able to initiate, upon membrane depolarization, intrinsic bursts which, by virtue of their spontaneous discharges and large spike output can drive other neurons to fire (25, 26). Burst firing is facilitated by a persistent slow sodium current (27) and by a tonic GABA_A-mediated conductance generated by the activation of extrasynaptic GABA_A receptors by “ambient” GABA whose depolarizing action would bring the membrane to the voltage window for activation of voltage-dependent sodium and calcium channels (28). Intrinsic bursting activity is boosted by the low expression of Kv7.2 and Kv7.3 channels responsible for the non-inactivating, low-threshold M current (I_M), which in adulthood controls spike after-depolarization and burst generation (29). The low density of I_M at birth contributes to produce intrinsic bursts that, in comparison with those observed in adults, are more robust, last longer and recur more regularly (25). GDPs-associated calcium transients act as coincident detector signals for enhancing synaptic efficacy at emerging GABAergic (30) and

glutamatergic synapses (31). Therefore, this early synchronized activity is fundamental for synaptic wiring and refinement of local neuronal circuits according to the Hebbian rule that “neurons that fire together wire together.”

γ -Aminobutyric acid is released from GABAergic interneurons that constitute a very heterogeneous group of cells, differentially classified according to their morphology, biophysical properties, molecular expression profile, and connectivity (32). These cells, mainly derived from the medial and caudal ganglionic eminences, undergo their final mitosis in these regions prior to their tangential migration into the cortical plate. The migration process, supposed to be calcium dependent, is regulated by a sequence of well-orchestrated processes involving guidance cues, neurotransmitter receptors (NMDA, GABA_A receptors) and voltage-dependent calcium channels (33, 34).

GABAergic interneurons not only exert a powerful control on network excitability but, in spite of their relatively low number (10–15% of the entire neuronal population), are able to synchronize a large number of principal cells giving rise to coherent oscillations, which support different behavioral states of the animals and high cognitive tasks (35).

Altogether, these observations point to GABA as one of the major players in the early assembly and formation of neuronal circuits in the developing brain. Therefore, it is not surprising that dysfunctions of GABAergic circuits have been implicated in various neuro-developmental and psychiatric disorders such as schizophrenia, autism, and epilepsy.

GABAergic DYSFUNCTIONS IN THE BRAIN OF ASD PATIENTS

The high frequency of epileptiform activity and the altered brain rhythms detected in the EEG of ASDs patients suggest a dysfunction of GABAergic transmission and an imbalance between excitation and inhibition (E/I) in local circuits involved in sensory, mnemonic, social, and emotional processes. However, as summarized in **Table 1**, more direct evidence in favor of a GABAergic dysfunction in ASDs derives from:

- a. Genetic observations
- b. *In vitro* analysis of post-mortem brain tissues
- c. *In vivo* studies on patients affected mainly by idiopathic forms of ASDs.

It is clear from the **Table 1** that most of cases are from juvenile and adult patients. This can be attributed to difficulties in obtaining post-mortem material from young children and to perform complex *in vivo* examinations such as positron emission tomography (PET) and single photon emission computed tomography (SPECT) in children of pediatric age. However, we cannot exclude that the same alterations are already present at early stages of development.

GENETIC OBSERVATIONS

The involvement of GABA_A receptors in ASDs was provided by genetic studies that have revealed submicroscopic abnormalities known as “copy-number variations” in chromosomal loci 15q11–q13, which contains a number of genes encoding for GABA_A receptor subunits (54). These loci can be affected either directly by

Table 1 | Alterations of GABAergic signaling in patients with idiopathic forms of autism and Rett syndrome.

Clinical phenotype	Alterations of GABAergic signaling	Age (years)	Reference
GENETIC OBSERVATIONS			
Idiopathic autism	Linkage disequilibrium in <i>GABRB3</i>	7.6 ± 6.2	(36, 37)
Idiopathic autism	Altered gene expression in interneurons	Not specified	(38)
IN VITRO ANALYSIS OF POST-MORTEM BRAIN TISSUES			
Idiopathic autism	Reduction in the density of GABA _A , GABA _B receptors and benzodiazepine binding sites in the anterior cingulate cortex	19/43	(39)
Idiopathic autism	Reduction in the density of GABA _B receptors cingulate cortex and fusiform gyrus	19/43	(40)
Idiopathic autism	Reduction in 3[H]flunitrazepam labeled benzodiazepines binding sites in the hippocampus	16/22	(41)
Idiopathic autism	Decreased number of GABAergic Purkinje cells in cerebellum	13/54	(42)
Idiopathic autism	Reduced level of GAD65 and GAD67 in Purkinje cells of cerebellar and parietal cortices	19/30	(43)
Idiopathic autism	Decreased GAD67 mRNA levels in cerebellar Purkinje cells	16/30	(44)
Idiopathic autism	Decreased GAD65 mRNA levels in cerebellar dentate nuclei	16/30	(45)
Idiopathic autism	Increased expression of GABAergic interneurons expressing calcium-binding proteins in the hippocampus	13/63	(46)
Rett syndrome	Disruption in the inhibitory architecture of the cell mini-columns	14.4 ± 4	(47)
IN VIVO STUDIES			
Idiopathic autism	Reduction of GABA concentration in the frontal lobe	2/12	(48)
Idiopathic autism	Reduced GABA levels in the perisylvian region of the left hemisphere	12.4 ± 5.2	(49)
Idiopathic autism	Reduced expression of GABA _A receptors in the superior and medial frontal cortex	7.3 ± 3.5	(50)
Rett syndrome	Reduced GABA _A receptor density in fronto-temporal cortex	27/41	(51)
Idiopathic autism	Significant reduction of α5 GABA _A receptor subunits in limbic areas	34/43	(52)
Rett syndrome	Reduced KCC2/NKCC1 ratio in the cerebrospinal fluid	0/19	(53)

Numbers refer to mean (usually ± SEM) or range values.

single point mutations or indirectly by epigenetic factors. Potential gene targets include *GABRB3*, *GABRA5*, and *GABRG3*, encoding for β3, α5, and γ3 subunits containing GABA_A receptors, respectively (36, 37). Systematic changes in GABA_A receptor subunit expression were found in the superior frontal cortex, parietal cortex, and cerebellum of autistic subjects. In addition, autism-related genes have been found to be expressed mainly in GABAergic interneurons (38).

IN VITRO ANALYSIS OF POST-MORTEM BRAIN TISSUES

Post-mortem analysis on brain tissues from ASD patients as well as genetic and *in vivo* studies have largely contributed to unveil the impact of GABAergic signaling in these disorders. Thus, as compared to controls, a significant reduction in the density of GABA_A, GABA_B receptors, and benzodiazepine binding sites was detected in the supra and infragranular layers of the anterior cingulate cortex (known to participate in a variety of processes including socio-emotional behavior and other associative functions via pre-frontal cortex connectivity) in brain samples from autistic subjects (39, 40). A reduction of ³[H]muscimol labeled GABA_A receptors and 3[H]flunitrazepam labeled benzodiazepines binding sites was found also in the hippocampus (41, 55). Neuropathological studies

from the cerebellum of individuals with ASDs have demonstrated that GABAergic Purkinje cells are particularly vulnerable since their number appears considerably reduced respect to controls (42, 56, 57). This effect was found to be associated with reduced levels of mRNA encoding for glutamic acid decarboxylase (GAD) 65 and 67, rate limiting enzymes responsible for the conversion of glutamate to GABA (43–45, 58). In contrast with Purkinje cells, enhanced levels of mRNA for GAD67 were found in stellate cells, a subtype of interneuron innervating Purkinje cells (43). Interestingly, pathological studies of brains of individual affected by autism demonstrated an increased expression of GABAergic interneurons expressing calcium-binding proteins such as calbindin-, calretinin-, and parvalbumin in the hippocampal formation (46). Since these interneurons are capable of buffering calcium, the intracellular messenger that controls several transduction pathways, an alteration of calcium signaling may have dramatic consequences on neuronal functions and dynamics.

Abnormalities in micro-columnar organization of prefrontal cortex in brain tissues from autistic individuals, including two with Angelman syndrome, have been also detected (59, 60). The narrower size of mini-columns in autistic patients respect to controls may reflect defects in GABAergic fibers within and

between cortical mini-columns due to reductions in the neuropil, which separates adjacent mini-columns (47). This may alter local connectivity and lateral inhibition.

IN VIVO STUDIES ON PATIENTS AFFECTED MAINLY BY IDIOPATHIC FORMS OF ASDs

In vivo studies from ASD patients are rather limited due to the difficulty of measuring GABAergic function *in vivo*. Using proton magnetic resonance spectroscopy, Harada et al. (48) reported a reduction of GABA concentration in the frontal lobe of children with ASDs respect to controls. However, in this study ASD patients were sedated with triclofos a compound that, by potentiating GABA action, may have biased the results. Using the same technique, Rojas et al. (49) have reported reduced GABA levels in the perisylvian region of the left hemisphere of autistic patients, further supporting the involvement of GABAergic neurotransmission in ASDs. Using SPECT and ^{123}I -iomazenil, a selective GABA_A-benzodiazepine ligand, a reduced expression of GABA_A receptors in the superior and medial frontal cortex of autistic patients was detected (50). Using the same technique, a reduced GABA_A receptor density was observed in three adult females affected by the Rett syndrome (51). In addition, a pilot study using PET and [^{11}C]Ro15-4513 to measure the expression levels of $\alpha 5$ GABA_A receptor subunits (localized mainly to extrasynaptic regions where they mediate tonic inhibition), has demonstrated a significant reduction of these subunits in limbic areas of ASD patients respect to controls (52).

Indirect evidence for deficits of GABAergic transmission in ASDs was provided by a recent magneto-encephalographic study that revealed impaired gamma-band activity in selective brain areas of adults with autism during perceptual visual processing (61). Gamma-band oscillations known to critically depend on negative feedback inhibition of principal cells by GABAergic interneurons, including parvalbumin-positive ones (62), are crucial for integrating multisensory information into a coherent representation (63). Although no data from young ASD patients are available, the possibility that such impairment may result from aberrant pre and perinatal development cannot be excluded.

Although informative, these studies referred mainly to idiopathic forms of ASDs and failed to address whether particular types of autistic syndromes, including monogenic ones were similarly affected. In this perspective, it is worth noting that a significant reduction of the cation-chloride importer KCC2 in the cerebrospinal fluid of Rett syndrome patients as compared to controls, with a consequent reduction in the KCC2/NKCC1 ratio was found (53). In accord with the notion that the methyl-CpG-binding protein 2 (MeCP2), encoded by the X-linked *Mecp2* gene, is highly expressed in GABAergic interneurons where it regulates their function these data suggest that a GABA dysfunction underlies the pathophysiology of the Rett syndrome.

Altogether, these findings indicate that alterations of GABAergic signaling in selective microcircuits of well-defined brain areas exert a fundamental role in the pathogenesis of ASDs.


GABAergic DYSFUNCTIONS IN ANIMAL MODELS OF ASDs

Over the last decade several animal models of ASDs have been generated, in order to identify the molecular and cellular mechanisms

underlying these disorders and to develop new therapeutic tools. To this aim, genetic defects detected in patients with ASDs have been introduced in the genome of mice. Animal models have been produced not only by manipulating candidate genes but also environmental factors or drugs such as valproic acid (VPA), an antiepileptic known to be a risk factor for autism in offspring of mothers treated during pregnancy with this drug (64). In accordance with human observations, animal studies have revealed important dysfunctions in GABAergic signaling occurring at different locations of the GABAergic synapse (Figure 1).

Table 2 summarizes alterations of GABAergic transmission observed in the most commonly used animal models of ASDs.

The *Fragile X* syndrome is a monogenetic disorder caused by mutations of the *FMR1* gene, located in chromosome X (Xq27.3) and encoding for the *Fragile X* mental retardation protein (FMRP), which is involved in the translation of a variety of mRNAs. Gene mutations do not allow the expression of the FMRP protein, determining a phenotype characterized by intellectual disabilities associated with language deficits, hyperactivity, autistic behavior, and seizures (85). Mice lacking the *FMR1* gene (Fmr1 KO mice) show an increased network excitability associated with an E/I imbalance in cortical circuits involving specific types of interneurons (65, 67, 86, 87). Thus, in the somatosensory cortex, the reduced activity of parvalbumin-positive cells (66) may contribute to enhance cell excitability and to affect gamma frequency oscillations, thought to be involved in high cognitive functions (86). The enhanced coupling of group I metabotropic glutamate receptors (mGluR) signaling and cannabinoid receptors mobilization may further enhance suppression of inhibition (88). Other factors contributing to enhance cell excitability include: (i) a reduced expression of GABA_A receptors subunits in the cortex. This may represent an evolutionary conserved hallmark of the *Fragile X* syndrome since similar reduction has been detected in the *Fragile X* fruit fly model (89). (ii) A down regulation of mRNA encoding for GAD65 and GAD67 in presynaptic terminals of the cortex and cerebellum; (iii) a reduction of mRNA encoding for gephyrin, the scaffold molecule responsible for glycine and GABA_A receptor clustering and stabilization at post-synaptic sites; (iv) reduced phasic and tonic GABA_A-mediated conductances in the amygdala associated with a down regulation of GABA_A receptor subunits (65, 68, 89); (v) a down regulation of GABA_A-mediated tonic inhibition associated with a reduced expression of $\alpha 5$ and δ GABA_A receptors subunits in subicular neurons (69). In contrast, electrophysiological recordings from the striatum of Fmr1 KO mice have revealed a selective increase in basal inhibitory neurotransmission, caused by the enhanced probability of GABA release (90). Overall these data suggest that in this animal model modifications of the GABAergic function are region-specific. Interestingly, a recent study from the hippocampus of mice carrying the *Fragile X* mutation, has revealed the loss of oxytocin-mediated GABA_A-mediated inhibition during the delivery process (70). In rats, oxytocin, the maternal hormone involved in delivery, has been shown to cause an abrupt shift of GABA from the depolarizing to the hyperpolarizing direction, thus exerting a neuroprotective and analgesic action on newborns (91). Unlike wild-type animals, the depolarizing action of GABA, apparently caused by the reduced expression of KCC2 with consequent high $[\text{Cl}^-]_i$, persists in juvenile life. The



increased excitatory drive to principal cells boosts network activity in the hippocampus, suggesting that an altered E/I balance may be caused not only by a decreased GABA_A-mediated inhibition but also by an enhanced GABA_A-mediated excitation (70).

Another neuro-developmental disorder showing high comorbidity with autism is the *Rett syndrome*. This is caused by mutations in the X-linked *Mecp2* gene that codes for the transcriptional factor MeCP2, highly expressed in GABAergic neurons (92). This disorder affects mainly girls who develop normally until the age of 6/18 months. Later, they develop cognitive deficits, loss of speech, motor abnormalities, respiratory dysrhythmias, stereotyped behavior, and seizures leading sometimes to premature death (93). Mice lacking MeCP2 or engineered to express an allele mimicking some mutation present in the *Rett syndrome* exhibit neurological symptoms reminiscent of those found in patients affected by the *Rett syndrome*. Similarly to Fmr1 KO mice, these animals exhibit alterations of the E/I balance (71), associated with a decrease in somatic GABA content and low levels of mRNA encoding for GAD65 and GAD67 in cortical and striatal neurons (75). While in the thalamus, the *Mecp2* gene appears to differentially regulate the development of GABAergic synapses in excitatory and inhibitory neurons (94), in the brain stem, the E/I imbalance is due to the depression of GABAergic transmission originating at both pre- and post-synaptic levels (72). A reduced GABAergic inhibition is present also in the locus coeruleus (73). In the hippocampus of *Mecp2* KO mice, a significant reduction in the quantal size of miniature inhibitory post-synaptic currents (mIPSCs) has been detected. This would account for the impairment of long-term

potentiation (LTP) induced at CA3–CA1 synapses by theta burst stimulation (74).

An E/I imbalance has been found also in individuals with *Tuberous sclerosis*, a genetic multisystem disorder characterized by wide spread hamartomas in several organs, including brain, heart, skin, eyes, kidney, lung, and liver (95). Tuberous sclerosis patients exhibit a variety of neurological disorders including mental retardation, autism-like disorders, and epilepsy. The affected genes are *Tsc1* and *Tsc2* encoding hamartin and tuberin, respectively. The hamartin–tuberin complex inhibits the mammalian target of rapamycin pathway that controls cell growth and proliferation (95). Immunocytochemical and western blots experiments have demonstrated that this disorder is associated with a decrease of $\alpha 1$ GABA_A receptor subunits and reduced and enhanced levels of KCC2 and NKCC1, respectively, in tubers. Changes in the expression of KCC2 and NKCC1 account for the excitatory action of GABA revealed with patch clamp in slices from *Tuberous sclerosis* tubers (96). However, electrophysiological data are still preliminary and should be taken with caution since they refer only to experiments from cortical slices obtained from tubers of a single patient. The *Angelman* or the closely related *Prader–Willi* syndromes (depending from which parent the deletion has been inherited), characterized by mental retardation, autistic behavior, and seizures, are determined by the loss of function of the *Ube3a* gene encoding for the ubiquitin E3 ligase or *Gabrb3*, *Gabra5*, and *Gabrg3* genes, encoding, respectively for $\beta 3$, $\alpha 5$, and $\gamma 3$ GABA_A receptor subunits localized in the same chromosome region 15q11–q13. Ube3a-deficient mice exhibit a reduced tonic

Table 2 | Alterations of GABAergic signaling in animal models of ASDs (E, embryonic day; P, postnatal day).

Animal models	Alterations of GABAergic signaling	Age	Reference
Fmr1 KO (<i>X Fragile</i>)	Reduced GABA release in the amygdala	P20–30	(65)
	Reduced number of parvalbumin-positive interneurons	Adult	(66)
	Altered E/I balance	P14–P30	(67)
	Decreased expression of GAD67 and GABA _A receptor subunits	Adult	(68)
	Down regulation of GABA _A -mediated tonic inhibition	Adult	(69)
	Persistent depolarizing effect of GABA in juvenile animals	E20–P30	(70)
MECP2 KO (Rett syndrome)	Altered E/I balance	P14–P35	(71)
	Depressed GABAergic synaptic transmission	P7	(72)
		P14–P28	(73)
		Adult	(74)
	Reduced expression of GAD65 and GAD 67	Adult	(75)
Ube3a-deficient mouse (Angelman syndrome)	Reduced GABA _A -mediated tonic inhibition	P25–P28 or adult	(76)
GABRB3 KO (Angelman syndrome)	Decreased expression of GABA _A receptors	Adult	(77)
<i>Scn1a</i> ^{+/-} mice (Dravet's syndrome)	Altered E/I balance	P21–P30	(78)
NL3 ^{R451C} KI	Increased GABAergic neurotransmission; increased VGAT and gephyrin expression	P13–P16	(79)
	Increased frequency of GDPs	P4–P35	(14)
	Circuit specific changes in GABAergic signaling in the hippocampus and in the cortex	P21–P35	(80)
		P9–P15	(81)
	Decreased number of PV ⁺ interneurons	P21–P35	(82)
En2 KO	Reduction of GABAergic markers during development; reduced number of GABAergic interneurons	Adult	(83)
VPA model	Decreased GABA _A -mediated neurotransmission	P23–P45	(84)
	Persistent depolarizing effect of GABA in juvenile animals	E20–P30	(70)

GABA-mediated inhibition in the cerebellum, caused by a GAT-1 dependent decrease of GABA concentration in the extracellular space (76). Moreover, β 3-deficient mice, which have a phenotype similar to some forms of *Angelman* syndrome, exhibit a reduced expression of GABA_A receptors in selective brain regions as determined by quantitative autoradiography (77).

The *Dravet's* syndrome is caused by a haploinsufficiency of the *SCN1A* gene encoding for voltage-gated sodium channel Nav1.1. Children affected by this disorder show intractable seizures, cognitive deficit, and autism spectrum behaviors. GABA plays a pivotal role also in this disorder as demonstrated by the reduced activity of Nav1.1 channels in forebrain GABAergic interneurons of *Scn1a*^{+/-} mice. These animals exhibited an E/I imbalance resulting from a decreased frequency of spontaneous inhibitory post-synaptic currents and an increased frequency of spontaneous excitatory post-synaptic currents in the hippocampus and prefrontal cortex. The impairment of GABAergic neurotransmission is associated with behavioral and cognitive deficits similar to those found in patients affected by the *Dravet's* syndrome (78).

Although rare, single mutations of genes encoding for adhesion molecules of the Neuroligin (NL) family found in individuals with autism have been introduced in mice. One of these, the R451C mutation of the *Nlgn3* gene encoding for NL3, has been found

in a family with children affected by ASDs (97). NLs are adhesion post-synaptic proteins that, by binding to their presynaptic partners, neuroligins, functionally couple the post-synaptic densities with the transmitter release machinery, thus contributing to synapses stabilization (98). Mice carrying the NL3 R451C mutation (NL3^{R451C} knock-in mice) show modifications of GABAergic signaling associated with behavioral deficits reminiscent of those found in autistic children (79, 99). These mice exhibit an increased frequency of mIPSCs in the somatosensory cortex (79) and in the CA3 region of the hippocampus (100) where they contribute to boost GDPs activity. A more detailed analysis of GABAergic microcircuits in the hippocampus has unveiled an increased GABA release at synapses between cholecystokinin (CCK)-positive endocannabinoids-sensitive interneurons and pyramidal cells and a decreased GABA release at synapses between parvalbumin-positive basket cells and principal cells (80). The similar phenotype found at CCK-pyramidal cell synapses in NL3^{R451C} knock-in mice and in NL3 KO mice suggests a loss of function consisting in the loss of tonic endocannabinoid signaling at these connections. It is worth noting that NL3^{R451C} knock-in mice present an asymmetric reduction of parvalbumin-positive basket cells across the two hemispheres (82).

Similarly to Földy et al. (80), a reduced probability of GABA release has been found in parvalbumin-positive basket cell–spiny

neuron synapses in layer IV somatosensory barrel cortex of juvenile NL3^{R451C} knock in mice. Such deficit determines an alteration of the E/I balance in this cortical layer together with a modification of the temporal window for integration of sensory inputs in principal cells (81). The altered sensory representations may underline deficits in coherent percepts of autistic children.

Mutations in the *En2* gene, coding for the homeobox-containing transcription factor engrailed-2 (EN2), involved in patterning and neuronal differentiation of the midbrain/hindbrain region, have been associated with ASDs. *En2* KO mice have been proposed as a model for ASDs due to their behavioral abnormalities similar to those observed in individuals with ASDs (101). The *En2* gene is also involved in the development or maintenance of GABAergic interneurons as demonstrated by the selective loss of parvalbumin, somatostatin, and neuropeptide Y positive interneurons in the cortex and hippocampus of *En2* KO mice (83). This effect appears to be region-specific since different subpopulations of interneurons are affected in posterior brain areas (102). Deficits in GABAergic neurotransmission account for the higher susceptibility to seizures of *En2* KO mice respect to controls (103).

Valproic acid is a histone deacetylase inhibitor, widely used to cure epilepsy and bipolar disorders. VPA is also a potent teratogen since children exposed *in utero* to this drug have a much higher risk of developing an autistic-type behavior than normal children (64). Pups exposed to VPA *in utero* show neuro-developmental abnormalities and behavioral deficits similar to ASDs (104). As in other animal models of ASDs, the autistic phenotype is associated to an altered E/I balance due to a decreased GABAergic signaling, which affects both pre- and post-synaptic sites, leaving the extrasynaptic transmission unaffected (84). Interestingly, like the NL3 model, the VPA model exhibits an asymmetric reduction of parvalbumin-positive cells across the two hemispheres (82). Similarly to *FRM1* KO mice, also in this animal model a loss of oxytocin-mediated GABA_A-mediated inhibition occurs during the transition from fetal to postnatal life (70). Also here as in *Fmr1* KO mice, the depolarizing action of GABA in the hippocampus persists in juvenile life. The excitatory GABAergic drive to principal cells leads to an increased network activity in a broad spectrum of frequencies including gamma oscillations.

The different animal models mentioned here have in common dysfunctions of GABAergic signaling leading to alterations of the E/I balance in selective brain circuits. These alterations can be rescued by selective tools that regulate GABA_A-mediated synaptic transmission.

THERAPEUTIC INTERVENTIONS TO RESCUE GABAergic DYSFUNCTIONS IN ASDs

Figure 1 shows a GABAergic synapse with its different constituents.

As summarized in **Figure 1A**, GABA released from a presynaptic terminal binds to post-synaptic GABA_A receptors localized on precise apposition to presynaptic release sites. It binds also to presynaptic and post-synaptic GABA_A and GABA_B receptors [for a review see Ref. (105)]. Post-synaptic GABA_B receptors are localized mainly at perisynaptic sites. Post-synaptic GABA_A receptors are maintained in the right place by gephyrin, a scaffold protein which, by interacting with cytoskeletal anchoring elements,

contributes to regulate receptor trafficking in and out of the synapses (106). Before being released in a calcium-dependent way, GABA is synthesized by GAD65 and GAD67 and it is stored in presynaptic vesicles by the vesicular transporter VGAT, which uses the electrochemical gradient for H⁺ to shuffle and pack GABA into synaptic vesicles (107). The probability of GABA release is under control of presynaptic receptors including GABA_A and GABA_B receptors. Adhesion molecules of the neuroligin–neurexin families ensure the cross-talk between post and presynaptic elements of the synapses (98). The figure also shows that the presynaptic terminal contains sodium channels, since, as already mentioned, a selective mutation of these channels in GABAergic interneurons of the forebrain is responsible for the Dravet's syndrome. After being released, GABA is taken up into nerve terminals and astrocytes by GABA transporters (GATs) localized on presynaptic nerve terminals and astrocytes.

Figure 1B shows the direction of chloride fluxes through post-synaptic GABA_A receptors in immature and mature neurons. The influx or efflux of chloride in and out of the cells is mainly dictated by the activity of two developmentally regulated cation–chloride co-transporters, the NKCC1 and KCC2, which pump chloride inside and outside of neurons, respectively (8). Therefore, different components of the synapses can be selectively targeted by agents that can rescue their functions.

Due to the co-morbidity of ASDs with epilepsy, anticonvulsants are widely used for the symptomatic treatment of these disorders (108). Some of these drugs have also an anti-anxiety effect and may indirectly increase GABA_A neurotransmission by promoting the synthesis of GABA or by inhibiting their reuptake or breakdown. Interestingly, treatment with vigabatrin, which blocks GABA catabolism by inhibiting GABA transaminase, is able to control seizures and to improve the autistic behavior of children affected by tuberous sclerosis (109).

γ-Aminobutyric acid agonists have proved to be effective in normalizing the E/I imbalance in animal models of Autism: they can either directly enhance inhibition or indirectly reduce excitation. Thus, in BTBR mice [an animal model of idiopathic autism, Ref. (110)] or in *Scn1a*^{+/-} mice [a monogenic model of ASDs, Ref. (78)], in which a reduced GABA_A-mediated inhibition occurs, the treatment with low non-sedative, non-anxiolytic doses of benzodiazepines, or clonazepam, known to enhance GABAergic signaling *via* allosteric modulation of post-synaptic GABA_A receptors, leads to an improvement of social and cognitive deficits. Interestingly, in both BTBR and *Scn1a*^{+/-} mice, the up-regulation of α2 and/or α3 containing GABA_A receptors subunits by L-838,417, a selective positive allosteric modulator of these subunits, which does not induce tolerance, is able to mimic the effects of benzodiazepines, thus providing an attractive tool to enhance GABAergic neurotransmission and to improve behavioral deficits in ASDs (110). Consistent with this view, clinical trials using α2/α3 selective positive allosteric modulators of GABA_A receptors have been developed with AstraZeneca and the National Institutes of Health (<http://clinicaltrials.gov/show/NCT01966679>).

Animal models of the Rett syndrome are currently used in translational studies for the preclinical evaluation of new therapeutic trials (111). Genetic studies have provided evidence that neuronal dysfunctions and autistic symptoms can be reversed

upon restoration of MeCP2 protein in mice in which the *Mecp2* gene has been silenced using a *lox-Stop-lox* cassette (112, 113), suggesting that in this animal model, the neuronal connectivity is not altered and neurons and glia are not permanently damaged by the *Mecp2* loss. Pharmacological studies have demonstrated amelioration of autistic symptoms with IGF-1 (114) or BDNF (115). However, since most of autistic features of the Rett syndrome can be recapitulated by deleting the *Mecp2* gene in GABA releasing neurons (75), these are expected to be rescued by drugs that enhance GABAergic neurotransmission.

Treatment of *Fmr1* KO mice (an animal model of *Fragile X* syndrome), exhibiting an hyper-excitability and a GABAergic dysfunction in the basolateral nucleus of the amygdala, with gaboxadol (THIP), a selective agonist at δ subunits containing perisynaptic or extrasynaptic GABA_A receptors, results in an increased GABA_A-mediated tonic conductance and in beneficial effects on learning deficits and behavioral disturbances linked to this disorder. This supports the hypothesis that tonic inhibition is a putative target for the treatment of *Fragile X* syndrome (65, 116). In addition, consistent with the notion that mGluR are important regulators of protein synthesis, which is translationally repressed by the FMR protein, partial inhibition of mGluR5 or inhibition of excessive glutamate release by the GABA_B receptor agonist arbaclofen in *Fmr1* KO mice have led to promising results (117, 118). Although the selective activation of GABA_B receptors with arbaclofen has the potential to improve social function and behavior in patients with *Fragile X* syndrome (119), its use in clinical trials is still under debate (120).

It is worth mentioning that in some children affected by ASDs, allosteric modulators of GABA_A receptors such as benzodiazepines have a paradoxical effect, increasing anxiety, and aggression (121). Therefore, restoring low $[Cl^-]_i$ and the inhibitory action of GABA, using the selective blocker of the NKCC1 chloride importer bumetanide, may have beneficial effects. In an elegant study, Tyzio et al. (70) have convincingly demonstrated that, treating pregnant VPA rats and *Fragile X* mice shortly before delivery with bumetanide, suppresses in both animal models the excitatory action of GABA and prevents the autistic-like behavior in offspring. Bumetanide may not only reverse GABA action from the depolarizing to the hyperpolarizing direction, but it may also reduce cell excitability by an ephaptic type of mechanism involving regulation of the cell volume and the extracellular space (122). Animal data validate previous findings by the same group, showing amelioration of autistic symptoms in children treated for 3 months with bumetanide (123, 124). Although these studies should be expanded to larger multicenter trials with more restricted inclusion and exclusion criteria and more extended investigations on the dose/response action of the diuretic as well as persistence of action after drug's withdrawal, the significant improvement of the autistic behavior associated with a statistically significant amelioration of childhood autism rating scale (CARS) and clinical global impressions (CGI) scores in the absence of clear side effects make this diuretic a very promising drug to cure ASDs. In a parallel study, bumetanide was shown to improve accuracy in facial emotional labeling and to increase brain activation in areas involved in social and emotional perception (125).

CONCLUSION

Several lines of evidences suggest that ASDs are neurodevelopmental disorders characterized by a clear E/I imbalance in selective neuronal circuits. Such disequilibrium appears to be mainly related to heterogeneous defects of GABAergic signaling in different brain structures. This has paved the way toward the development of new drugs for the cure of these devastating disorders. In particular, several studies have shown that drugs acting on GABAergic synapses are able to rescue behavioral deficits in animal models of autism and to ameliorate at least some of the symptoms observed in ASD patients.

In order to successfully translate therapeutic approaches from animal to humans, it is necessary to develop animal models of ASDs that faithfully trace the behavioral alterations detected in ASD patients. In addition, the effects of drugs should be validated in humans in large scale clinical trials with accurate controls.

Since ASDs are developmental disorders, the early pharmacological intervention, guaranteed by an early diagnosis, is essential. This means that very young children will receive drugs, whose side effects have to be carefully considered. For instance, many drugs acting on GABAergic synapses can generate addiction or give rise to paradoxical reactions, as brain circuits are still immature and GABA may still exert a depolarizing and excitatory action which can be prolonged at late stages of development. Only accurate studies and the use of well suited animal models can help to design pharmacological tools with minimal risks.

AUTHOR CONTRIBUTIONS

Giada Cellot and Enrico Cherubini wrote the paper.

ACKNOWLEDGMENTS

This study was supported by a grant from Telethon (GGP11043) and from Ministero dell'Istruzione, dell'Università e della Ricerca (PRIN 2011) to Enrico Cherubini. The authors are particularly grateful to Dr. R. Pizzarelli who contributed to some of the original work on an animal model of ASDs, as well as all members of the laboratory for useful discussions.

REFERENCES

1. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. Arlington, VA: American Psychiatric Association (2000).
2. Levy SE, Mandell DS, Schultz R. Autism. *Lancet* (2009) **374**:1627–38. doi:10.1016/S0140-6736(09)61376-3
3. Ben-Ari Y, Woodin MA, Sernagor E, Cancedda L, Vinay L, Rivera C, et al. Refuting the challenges of the developmental shift of polarity of GABA actions: GABA more exciting than ever! *Front Cell Neurosci* (2012) **6**:35. doi:10.3389/fncel.2012.00035
4. Frye RE, Rossignol D, Casanova MF, Brown GL, Martin V, Edelson S, et al. A review of traditional and novel treatments for seizures in autism spectrum disorder: findings from a systematic review and expert panel. *Front Public Health* (2013) **1**:31. doi:10.3389/fpubh.2013.00031
5. Valeeva G, Abdullin A, Tyzio R, Skorinkin A, Nikolski E, Ben-Ari Y, et al. Temporal coding at the immature depolarizing GABAergic synapse. *Front Cell Neurosci* (2010) **14**:4. doi:10.3389/fncel.2010.00017
6. Ben-Ari Y, Cherubini E, Corradetti R, Gaiarsa JL. Giant synaptic potentials in immature rat CA3 hippocampal neurones. *J Physiol* (1989) **416**:303–25.
7. Cherubini E, Gaiarsa JL, Ben-Ari Y. Review GABA: an excitatory transmitter in early postnatal life. *Trends Neurosci* (1991) **14**:515–9. doi:10.1016/0166-2236(91)90003-D
8. Rivera C, Voipio J, Payne JA, Ruusuvoori E, Lahtinen H, Lamsa K, et al. The K⁺/Cl⁻ co-transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. *Nature* (1999) **397**:251–5. doi:10.1038/16697

9. Yamada J, Okabe A, Toyoda H, Kilb W, Luhmann HJ, Fukuda A. Cl-uptake promoting depolarizing GABA actions in immature rat neocortical neurones is mediated by NKCC1. *J Physiol* (2004) **557**:829–41. doi:10.1113/jphysiol.2004.062471
10. Dzhalal VI, Talos DM, Sdrulla DA, Brumback AC, Mathews GC, Benke TA, et al. NKCC1 transporter facilitates seizures in the developing brain. *Nat Med* (2005) **11**:1205–13. doi:10.1038/nm1301
11. Wang DD, Kriegstein AR. Defining the role of GABA in cortical development. *J Physiol* (2009) **587**:1873–9. doi:10.1113/jphysiol.2008.167635
12. Ben-Ari Y, Gaiarsa JL, Tyzio R, Khazipov R. GABA: a pioneer transmitter that excites immature neurons and generates primitive oscillations. *Physiol Rev* (2007) **87**:1215–84. doi:10.1152/physrev.00017.2006
13. Cherubini E, Griguoli M, Safulina V, Lagostena L. The depolarizing action of GABA controls early network activity in the developing hippocampus. *Mol Neurobiol* (2011) **43**:97–106. doi:10.1007/s12035-010-8147-z
14. Pizzarelli R, Cherubini E. Alterations of GABAergic signaling in autism spectrum disorders. *Neural Plast* (2011) **2011**:297153. doi:10.1155/2011/297153
15. Tyzio R, Represa A, Jorquera I, Ben-Ari Y, Gozlan H, Aniksztejn L. The establishment of GABAergic and glutamatergic synapses on CA1 pyramidal neurons is sequential and correlates with the development of the apical dendrite. *J Neurosci* (1999) **19**:10372–82.
16. Behar TN, Schaffner AE, Scott CA, O'Connell C, Barker JL. Differential response of cortical plate and ventricular zone cells to GABA as a migration stimulus. *J Neurosci* (1998) **18**:6378–87.
17. Nguyen L, Rigo JM, Rocher V, Belachew S, Malgrange B, Rogister B, et al. Neurotransmitters as early signals for central nervous system development. *Cell Tissue Res* (2001) **305**:187–202. doi:10.1007/s004410000343
18. Demarque M, Represa A, Becq H, Khalilov I, Ben-Ari Y, Aniksztejn L. Paracrine intercellular communication by a Ca²⁺- and SNARE-independent release of GABA and glutamate prior to synapse formation. *Neuron* (2002) **36**:1051–61. doi:10.1016/S0896-6273(02)01053-X
19. Manent JB, Demarque M, Jorquera I, Pellegrino C, Ben-Ari Y, Aniksztejn L, et al. A noncanonical release of GABA and glutamate modulates neuronal migration. *J Neurosci* (2005) **25**:4755–65. doi:10.1523/JNEUROSCI.0553-05.2005
20. Cancedda L, Fiumelli H, Chen K, Poo MM. Excitatory GABA action is essential for morphological maturation of cortical neurons in vivo. *J Neurosci* (2007) **27**:5224–35. doi:10.1523/JNEUROSCI.5169-06.2007
21. Leinekugel X, Khazipov R, Cannon R, Hirase H, Ben-Ari Y, Buzsáki G. Correlated bursts of activity in the neonatal hippocampus in vivo. *Science* (2002) **296**:2049–52. doi:10.1126/science.1071111
22. Stockard-Pope J, Werner SS, Bickford RG. *Atlas of Neonatal Electroencephalography*. New York, NY: Raven Press (1992).
23. De la Prida LM, Huberfeld G, Cohen I, Miles R. Threshold behavior in the initiation of hippocampal population bursts. *Neuron* (2006) **49**:131–42. doi:10.1016/j.neuron.2005.10.034
24. De la Prida L, Sanchez-Andres JV. Heterogeneous populations of cells mediate spontaneous synchronous bursting in the developing hippocampus through a frequency-dependent mechanism. *Neuroscience* (2000) **97**:227–41. doi:10.1016/S0306-4522(00)00029-4
25. Safulina V, Zacchi P, Tagliatela M, Yaari Y, Cherubini E. Low expression of Kv7/M channels facilitates intrinsic and network bursting in the developing rat hippocampus. *J Physiol* (2008) **586**:5437–53. doi:10.1113/jphysiol.2008.156257
26. Sipilä ST, Huttu K, Soltesz I, Voipio J, Kaila K. Depolarizing GABA acts on intrinsically bursting pyramidal neurons to drive giant depolarizing potentials in the immature hippocampus. *J Neurosci* (2005) **25**:5280–9. doi:10.1523/JNEUROSCI.0378-05.2005
27. Sipilä ST, Huttu K, Voipio J, Kaila K. Intrinsic bursting of immature CA3 pyramidal neurons and consequent giant depolarizing potentials are driven by a persistent Na⁺ current and terminated by a slow Ca²⁺-activated K⁺ current. *Eur J Neurosci* (2006) **23**:2330–8. doi:10.1111/j.1460-9568.2006.04757.x
28. Marchionni I, Omrani A, Cherubini E. In the developing rat hippocampus a tonic GABA-mediated conductance selectively enhances the glutamatergic drive of principal cells. *J Physiol* (2007) **581**:515–28. doi:10.1113/jphysiol.2006.125609
29. Yue C, Yaari Y. KCNQ/M channels control spike after depolarization and burst generation in hippocampal neurons. *J Neurosci* (2004) **24**:4614–24. doi:10.1523/JNEUROSCI.0765-04.2004
30. Kasyanov AM, Safulina V, Voronin LL, Cherubini E. GABA-mediated giant depolarizing potentials as coincidence detectors for enhancing synaptic efficacy in the developing hippocampus. *Proc Natl Acad Sci U S A* (2004) **101**:3967–72. doi:10.1073/pnas.0305974101
31. Mohajerani MH, Sivakumaran S, Zacchi P, Aguilera P, Cherubini E. Correlated network activity enhances synaptic efficacy via BDNF and the ERK pathway at immature CA3 CA1 connections in the hippocampus. *Proc Natl Acad Sci U S A* (2007) **104**:13176–81. doi:10.1073/pnas.0704533104
32. Freund TF, Buzsáki G. Interneurons of the hippocampus. *Hippocampus* (1996) **6**:347–470. doi:10.1002/(SICI)1098-1063(1996)6:4<347::AID-HIPO1>3.0.CO;2-I
33. Komuro H, Rakic P. Orchestration of neuronal migration by activity of ion channels, neurotransmitter receptors, and intracellular Ca²⁺ fluctuations. *J Neurobiol* (1998) **37**:110–30. doi:10.1002/(SICI)1097-4695(199810)37:1<110::AID-NEU9>3.3.CO;2-Q
34. Spitzer NC. Activity-dependent neuronal differentiation prior to synapse formation: the functions of calcium transients. *J Physiol Paris* (2002) **96**:73–80. doi:10.1016/S0928-4257(01)00082-1
35. Klausberger T, Somogyi P. Neuronal diversity and temporal dynamics: the unity of hippocampal circuit operations. *Science* (2008) **321**:53–7. doi:10.1126/science.1149381
36. Cook EH, Courchesne RY, Cox NJ, Lord C, Gonen D, Guter SJ, et al. Linkage-disequilibrium mapping of autistic disorder, with 15q11-13 markers. *Am J Hum Genet* (1998) **62**:1077–83. doi:10.1086/301832
37. Shao Y, Cuccaro ML, Hauser ER, Raiford KL, Menold MM, Wolpert CM, et al. Fine mapping of autistic disorder to chromosome 15q11-q13 by use of phenotypic subtypes. *Am J Hum Genet* (2003) **72**:539–48. doi:10.1086/367846
38. Xu X, Wells AB, O'Brien DR, Nehorai A, Dougherty JD. Cell type-specific expression analysis to identify putative cellular mechanisms for neurogenetic disorders. *J Neurosci* (2014) **34**:1420–31. doi:10.1523/JNEUROSCI.4488-13.2014
39. Oblak A, Gibbs TT, Blatt GJ. Decreased GABA receptors and benzodiazepine binding sites in the anterior cingulate cortex in autism. *Autism Res* (2009) **2**:205–19. doi:10.1002/aur.88
40. Oblak A, Gibbs T, Blatt GJ. Decreased GABA receptors in the cingulate cortex and fusiform gyrus in autism. *J Neurochem* (2010) **114**:1414–23. doi:10.1111/j.1471-4159.2010.06858.x
41. Guptill JT, Booker AB, Gibbs TT, Kemper TL, Bauman ML, Blatt GJ. [3H]-flunitrazepam-labeled benzodiazepine binding sites in the hippocampal formation in autism: a multiple concentration autoradiographic study. *J Autism Dev Disord* (2007) **37**:911–20. doi:10.1007/s10803-006-0226-7
42. Whitney ER, Kemper TL, Bauman ML, Rosene DL, Blatt GJ. Cerebellar Purkinje cells are reduced in a subpopulation of autistic brains: a cerebellar experiment using calbindin-D28k. *Cerebellum* (2008) **7**:406–16. doi:10.1007/s12311-008-0043-y
43. Fatemi SH, Halt AR, Sary JM, Kanodia R, Schulz SC, Realmuto GR. Glutamic acid decarboxylase 65 and 67 kDa proteins are reduced in autistic parietal and cerebellar cortices. *Biol Psychiatry* (2002) **52**:805–10. doi:10.1016/S0006-3223(02)01430-0
44. Yip J, Soghomonian JJ, Blatt GJ. Decreased GAD67 mRNA levels in cerebellar Purkinje cells in autism: pathophysiological implications. *Acta Neuropathol* (2007) **113**:559–68. doi:10.1007/s00401-006-0176-3
45. Yip J, Soghomonian JJ, Blatt GJ. Decreased GAD65 mRNA levels in select subpopulations of neurons in the cerebellar dentate nuclei in autism: an in situ hybridization study. *Autism Res* (2009) **2**:50–9. doi:10.1002/aur.62
46. Lawrence YL, Kemper TL, Bauman ML, Blatt GJ. Parvalbumin, calbindin, and calretinin immunoreactive interneuron density in autism. *Acta Neurol Scand* (2010) **121**:99–108. doi:10.1111/j.1600-0404.2009.01234.x
47. Casanova MF, Buxhoeveden DP, Gomez J. Disruption of the inhibitory architecture of the cell minicolumn: implications for autism. *Neuroscientist* (2003) **9**:496–507. doi:10.1177/1073858403253552
48. Harada M, Taki MM, Nose A, Kubo H, Mori K, Nishitani H, et al. Non-invasive evaluation of the GABAergic/glutamatergic system in autistic patients observed by MEGA-editing proton MR spectroscopy using a clinical 3 Tesla instrument. *J Autism Dev Disord* (2011) **41**:447–54. doi:10.1007/s10803-010-1065-0
49. Rojas DC, Singel D, Steinmetz S, Hepburn S, Brown MS. Decreased left perisylvian GABA concentration in children with autism and unaffected siblings. *Neuroimage* (2014) **86**:28–34. doi:10.1016/j.neuroimage.2013.01.045

50. Mori T, Mori K, Fujii E, Toda Y, Miyazaki M, Harada M, et al. Evaluation of the GABAergic nervous system in autistic brain: (123I)-iomazenil SPECT study. *Brain Dev* (2012) **34**:648–54. doi:10.1016/j.braindev.2011.10.007
51. Yamashita Y, Matsushita T, Ishibashi M, Kimura A, Onishi Y, Yonekura Y, et al. Decrease in benzodiazepine receptor binding in the brains of adult patients with Rett syndrome. *J Neurol Sci* (1998) **154**:146–50. doi:10.1016/S0022-510X(97)00223-2
52. Mendez MA, Horder J, Myers J, Coghlan S, Stokes P, Erritzoe D, et al. The brain GABA-benzodiazepine receptor alpha-5 subtype in autism spectrum disorder: a pilot [(11C)Ro15-4513 positron emission tomography study. *Neuropharmacology* (2013) **68**:195–201. doi:10.1016/j.neuropharm.2012.04.008
53. Duarte ST, Armstrong J, Roche A, Ortez C, Pérez A, O'Callaghan Mdel M, et al. Abnormal expression of cerebrospinal fluid cation chloride cotransporters in patients with Rett syndrome. *PLoS One* (2013) **8**:e68851. doi:10.1371/journal.pone.0068851
54. Coghlan S, Horder J, Inkster B, Mendez MA, Murphy DG, Nutt DJ. GABA system dysfunction in autism and related disorders: from synapse to symptoms. *Neurosci Biobehav Rev* (2012) **36**:2044–55. doi:10.1016/j.neubiorev.2012.07.005
55. Blatt GJ, Fitzgerald CM, Guptill JT, Booker AB, Kemper TL, Bauman ML. Density and distribution of hippocampal neurotransmitter receptors in autism: an autoradiographic study. *J Autism Dev Disord* (2001) **31**:537–43. doi:10.1023/A:1013238809666
56. Williams RS, Hauser SL, Purpura DP, DeLong GR, Swisher CN. Autism and mental retardation: neuropathologic studies performed in four retarded persons with autistic behavior. *Arch Neurol* (1980) **37**:749–53. doi:10.1001/archneur.1980.00500610029003
57. Bauman ML, Kemper TL. Histoanatomic observations of the brain in early infantile autism. *Neurology* (1985) **35**:866–74. doi:10.1212/WNL.35.6.866
58. Yip J, Soghomonian JJ, Blatt GJ. Increased GAD67 mRNA levels in cerebellar interneurons in autism: implications for Purkinje cell dysfunction. *J Neurosci Res* (2008) **86**:525–30. doi:10.1002/jnr.21520
59. Casanova MF, Buxhoeveden DP, Switala AE, Roy E. Minicolumnar pathology in autism. *Neurology* (2002) **58**:428–32. doi:10.1212/WNL.58.3.428
60. Casanova MF, Buxhoeveden DP, Switala AE, Roy E. Asperger's syndrome and cortical neuropathology. *J Child Neurol* (2002) **17**:142–5. doi:10.1177/088307380201700211
61. Sun L, Grützner C, Bölte S, Wibrall M, Tozman T, Schlitt S, et al. Impaired gamma-band activity during perceptual organization in adults with autism spectrum disorders: evidence for dysfunctional network activity in fronto-posterior cortices. *J Neurosci* (2012) **32**:9563–73. doi:10.1523/JNEUROSCI.1073-12.2012
62. Sohal VS, Zhang F, Yizhar O, Deisseroth K. Parvalbumin neurons and gamma rhythms enhance cortical circuit performance. *Nature* (2009) **459**:698–702. doi:10.1038/nature07991
63. Uhlhaas PJ, Singer W. Neural synchrony in brain disorders: relevance for cognitive dysfunctions and pathophysiology. *Neuron* (2006) **52**:155–68. doi:10.1016/j.neuron.2006.09.020
64. Meador KJ, Loring DW. Risks of in utero exposure to valproate. *JAMA* (2013) **309**(16):1730–1. doi:10.1001/jama.2013.4001
65. Olmos-Serrano JL, Paluszkiwicz SM, Martin BS, Kaufmann WE, Corbin JG, Huntsman MM. Defective GABAergic neurotransmission and pharmacological rescue of neuronal hyperexcitability in the amygdala in a mouse model of Fragile X syndrome. *J Neurosci* (2010) **30**:9929–38. doi:10.1523/JNEUROSCI.1714-10.2010
66. Selby L, Zhang C, Sun QQ. Major defects in neocortical GABAergic inhibitory circuits in mice lacking the Fragile X mental retardation protein. *Neurosci Lett* (2007) **412**:227–32. doi:10.1016/j.neulet.2006.11.062
67. Gibson JR, Bartley AF, Hays SA, Huber KM. Imbalance of neocortical excitation and inhibition and altered UP states reflect network hyperexcitability in the mouse model of Fragile X syndrome. *J Neurophysiol* (2008) **100**:2615–26. doi:10.1152/jn.90752.2008
68. Gantois I, Vandesompele J, Speleman F, Reyniers E, D'Hooge R, Severijnen LA, et al. Expression profiling suggests underexpression of the GABA(A) receptor subunit delta in the Fragile X knockout mouse model. *Neurobiol Dis* (2006) **21**:346–57. doi:10.1016/j.nbd.2005.07.017
69. Curia G, Papouin T, Séguéla P, Avoli M. Downregulation of tonic GABAergic inhibition in a mouse model of Fragile X syndrome. *Cereb Cortex* (2009) **19**:1515–20. doi:10.1093/cercor/bhn159
70. Tyzio R, Nardou R, Ferrari DC, Tsintsadze T, Shahrokhi A, Eftekhari S, et al. Oxytocin-mediated GABA inhibition during delivery attenuates autism pathogenesis in rodent offspring. *Science* (2014) **343**:675–9. doi:10.1126/science.1247190
71. Dani VS, Chang Q, Maffei A, Turrigiano GG, Jaenisch R, Nelson SB. Reduced cortical activity due to a shift in the balance between excitation and inhibition in a mouse model of Rett syndrome. *Proc Natl Acad Sci U S A* (2005) **102**:12560–5. doi:10.1073/pnas.0506071102
72. Medrihan L, Tantalaki E, Aramuni G, Sargsyan V, Dudanova I, Missler M, et al. Early defects of GABAergic synapses in the brain stem of a MeCP2 mouse model of Rett syndrome. *J Neurophysiol* (2008) **99**:112–21. doi:10.1152/jn.00826.2007
73. Jin X, Cui N, Zhong W, Jin XT, Jiang C. GABAergic synaptic inputs of locus coeruleus neurons in wild-type and Mecp2-null mice. *Am J Physiol Cell Physiol* (2013) **304**:844–57. doi:10.1152/ajpcell.00399.2012
74. Zhang L, He J, Jugloff DG, Eubanks JH. The MeCP2-null mouse hippocampus displays altered basal inhibitory rhythms and is prone to hyperexcitability. *Hippocampus* (2008) **18**:294–309. doi:10.1002/hipo.20389
75. Chao HT, Chen H, Samaco RC, Xue M, Chahrour M, Yoo J, et al. Dysfunction in GABA signalling mediates autism-like stereotypies and Rett syndrome phenotypes. *Nature* (2010) **468**:263–9. doi:10.1038/nature09582
76. Egawa K, Kitagawa K, Inoue K, Takayama M, Takayama C, Saitoh S, et al. Decreased tonic inhibition in cerebellar granule cells causes motor dysfunction in a mouse model of Angelman syndrome. *Sci Transl Med* (2012) **4**:163ra157. doi:10.1126/scitranslmed.3004655
77. Sinkkonen ST, Homanics GE, Korpi ER. Mouse models of Angelman syndrome, a neurodevelopmental disorder, display different brain regional GABA(A) receptor alterations. *Neurosci Lett* (2003) **340**:205–8. doi:10.1016/S0304-3940(03)00123-X
78. Han S, Tai C, Westenbroek RE, Yu FH, Cheah CS, Potter GB, et al. Autistic-like behaviour in Scn1a^{+/-} mice and rescue by enhanced GABA-mediated neurotransmission. *Nature* (2012) **489**:385–90. doi:10.1038/nature11356
79. Tabuchi K, Blundell J, Etherton MR, Hammer RE, Liu X, Powell CM, et al. A neuroligin-3 mutation implicated in autism increases inhibitory synaptic transmission in mice. *Science* (2007) **318**:71–6. doi:10.1126/science.1146221
80. Földy C, Malenka RC, Südhof TC. Autism-associated neuroligin-3 mutations commonly disrupt tonic endocannabinoid signaling. *Neuron* (2013) **78**:498–509. doi:10.1016/j.neuron.2013.02.036
81. Cellot G, Cherubini E. Reduced inhibitory gate in the barrel cortex of neuroligin3^{R451C} knock-in mice, an animal model of autism spectrum disorders. *Physiol Rep* (2014) **2**:e12070. doi:10.14814/phy2.12077
82. Gogolla N, Leblanc JJ, Quast KB, Südhof TC, Fagiolini M, Hensch TK. Common circuit defect of excitatory-inhibitory balance in mouse models of autism. *J Neurodev Disord* (2009) **1**:172–81. doi:10.1007/s11689-009-9023-x
83. Sgadò P, Genovesi S, Kalinovsky A, Zunino G, Macchi F, Allegra M, et al. Loss of GABAergic neurons in the hippocampus and cerebral cortex of engrailed-2 null mutant mice: implications for autism spectrum disorders. *Exp Neurol* (2013) **247**:496–505. doi:10.1016/j.expneurol.2013.01.021
84. Banerjee A, García-Oscos F, Roychowdhury S, Galindo LC, Hall S, Kilgard MP, et al. Impairment of cortical GABAergic synaptic transmission in an environmental rat model of autism. *Int J Neuropsychopharmacol* (2013) **16**:1309–18. doi:10.1017/S1461145712001216
85. Oostra BA, Chiurazzi P. The Fragile X gene and its function. *Clin Genet* (2001) **60**:399–408. doi:10.1034/j.1399-0004.2001.600601.x
86. Paluszkiwicz SM, Martin BS, Huntsman MM. Fragile X syndrome: the GABAergic system and circuit dysfunction. *Dev Neurosci* (2011) **33**:349–64. doi:10.1159/000329420
87. Gonçalves JT, Anstey JE, Golshani P, Portera-Cailliau C. Circuit level defects in the developing neocortex of Fragile X mice. *Nat Neurosci* (2013) **16**:903–9. doi:10.1038/nn.3415
88. Zhang L, Alger BE. Enhanced endocannabinoid signaling elevates neuronal excitability in Fragile X syndrome. *J Neurosci* (2010) **30**:5724–9. doi:10.1523/JNEUROSCI.0795-10.2010

89. D'Hulst C, Heulens I, Brouwer JR, Willemsen R, De Geest N, Reeve SP, et al. Expression of the GABAergic system in animal models for Fragile X syndrome and Fragile X associated tremor/ataxia syndrome (FXTAS). *Brain Res* (2009) **1253**:176–83. doi:10.1016/j.brainres.2008.11.075
90. Centonze D, Rossi S, Mercaldo V, Napoli I, Ciotti MT, De Chiara V, et al. Abnormal striatal GABA transmission in the mouse model for the Fragile X syndrome. *Biol Psychiatry* (2008) **63**:963–73. doi:10.1016/j.biopsych.2007.09.008
91. Tyzio R, Cossart R, Khalilov I, Minlebaev M, Hübner CA, Represa A, et al. Maternal oxytocin triggers a transient inhibitory switch in GABA signaling in the fetal brain during delivery. *Science* (2006) **314**:1788–92. doi:10.1126/science.1133212
92. Akbarian S, Chen RZ, Gribnau J, Rasmussen TP, Fong H, Jaenisch R, et al. Expression pattern of the Rett syndrome gene MeCP2 in primate prefrontal cortex. *Neurobiol Dis* (2001) **8**:784–91. doi:10.1006/nbdi.2001.0420
93. Amir RE, Van den Veyver IB, Wan M, Tran CQ, Francke U, Zoghbi HY. Rett syndrome is caused by mutations in X-linked MECP2, encoding methyl-CpG-binding protein 2. *Nat Genet* (1999) **23**:185–8. doi:10.1038/13810
94. Zhang ZW, Zak JD, Liu H. MeCP2 is required for normal development of GABAergic circuits in the thalamus. *J Neurophysiol* (2010) **103**:2470–81. doi:10.1152/jn.00601.2009
95. Curatolo P, Bombardieri R, Jozwiak S. Tuberous sclerosis. *Lancet* (2008) **372**:657–68. doi:10.1016/S0140-6736(08)61279-9
96. Talos DM, Sun H, Kosaras B, Joseph A, Folkert RD, Poduri A, et al. Altered inhibition in tuberous sclerosis and type IIb cortical dysplasia. *Ann Neurol* (2012) **71**:539–51. doi:10.1002/ana.22696
97. Jamain S, Quach H, Betancur C, Råstam M, Colinaux C, Gillberg IC, et al. Mutations of the X-linked genes encoding neuroligins NLGN3 and NLGN4 are associated with autism. *Nat Genet* (2003) **34**:27–9. doi:10.1038/ng1136
98. Südhof TC. Neuroligins and neuroligins link synaptic function to cognitive disease. *Nature* (2008) **455**:903–11. doi:10.1038/nature07456
99. Etherton M, Földy C, Sharma M, Tabuchi K, Liu X, Shamloo M, et al. Autism-linked neuroligin-3 R451C mutation differentially alters hippocampal and cortical synaptic function. *Proc Natl Acad Sci U S A* (2011) **108**:13764–9. doi:10.1073/pnas.1111093108
100. Pizzarelli R, Cherubini E. Developmental regulation of GABAergic signalling in the hippocampus of neuroligin 3 R451C knock-in mice: an animal model of autism. *Front Cell Neurosci* (2013) **4**(7):85. doi:10.3389/fncel.2013.00085
101. Cheh MA, Millonig JH, Roselli LM, Ming X, Jacobsen E, Kamdar S, et al. En2 knockout mice display neurobehavioral and neurochemical alterations relevant to autism spectrum disorder. *Brain Res* (2006) **1116**:166–76. doi:10.1016/j.brainres.2006.07.086
102. Sudarov A, Joyner AL. Cerebellum morphogenesis: the foliation pattern is orchestrated by multi-cellular anchoring centers. *Neural Dev* (2007) **3**(2):26. doi:10.1186/1749-8104-2-26
103. Tripathi PP, Sgado P, Scali M, Viaggi C, Casarosa S, Simon HH, et al. Increased susceptibility to kainic acid-induced seizures in engrailed-2 knockout mice. *Neuroscience* (2009) **159**:842–9. doi:10.1016/j.neuroscience.2009.01.007
104. Wagner GC, Reuhl KR, Cheh M, McRae P, Halladay AK. A new neurobehavioral model of autism in mice: pre- and postnatal exposure to sodium valproate. *J Autism Dev Disord* (2006) **36**:779–93. doi:10.1007/s10803-006-0117-y
105. Cherubini E. *Phasic GABA_A-Mediated Inhibition: Jasper's Basic Mechanisms of the Epilepsies*. Bethesda, MD: National Center for Biotechnology Information (2012).
106. Kneussel M, Betz H. Receptors, gephyrin and gephyrin-associated proteins: novel insights into the assembly of inhibitory postsynaptic membrane specializations. *J Physiol* (2000) **1**:1–9. doi:10.1111/j.1469-7793.2000.t01-4-00001.x
107. Buddhala C, Hsu CC, Wu JY. A novel mechanism for GABA synthesis and packaging into synaptic vesicles. *Neurochem Int* (2009) **55**:9–12. doi:10.1016/j.neuint.2009.01.020
108. Di Martino A, Tuchman RE. Antiepileptic drugs: affective use in autism spectrum disorders. *Pediatr Neurol* (2001) **25**:199–207. doi:10.1016/S0887-8994(01)00276-4
109. Jambaqué I, Chiron C, Dumas C, Mumford J, Dulac O. Mental and behavioural outcome of infantile epilepsy treated by vigabatrin in tuberous sclerosis patients. *Epilepsy Res* (2000) **38**:151–60. doi:10.1016/S0920-1211(99)00082-0
110. Han S, Tai C, Jones CJ, Scheuer T, Catterall WA. Enhancement of inhibitory neurotransmission by GABA_A receptors having $\alpha 2,3$ -subunits ameliorates behavioral deficits in a mouse model of autism. *Neuron* (2014) **81**:1282–9. doi:10.1016/j.neuron.2014.01.016
111. Katz DM, Berger-Sweeney JE, Eubanks JH, Justice MJ, Neul JL, Pozzo-Miller L, et al. Preclinical research in Rett syndrome: setting the foundation for translational success. *Dis Model Mech* (2012) **5**:733–45. doi:10.1242/dmm.011007
112. Guy J, Gan J, Selfridge J, Cobb S, Bird A. Reversal of neurological defects in a mouse model of Rett syndrome. *Science* (2007) **315**:1143–7. doi:10.1126/science.1138389
113. Giacometti E, Luikenhuis S, Beard C, Jaenisch R. Partial rescue of MeCP2 deficiency by postnatal activation of MeCP2. *Proc Natl Acad Sci U S A* (2007) **104**:1931–6. doi:10.1073/pnas.0610593104
114. Tropea D, Giacometti E, Wilson NR, Beard C, McCurry C, Fu DD, et al. Partial reversal of Rett syndrome-like symptoms in MeCP2 mutant mice. *Proc Natl Acad Sci U S A* (2009) **106**:2029–34. doi:10.1073/pnas.0812394106
115. Kline DD, Ogier M, Kunze DL, Katz DM. Exogenous brain-derived neurotrophic factor rescues synaptic dysfunction in Mecp2-null mice. *J Neurosci* (2010) **30**:5303–10. doi:10.1523/JNEUROSCI.5503-09.2010
116. Olmos-Serrano JL, Corbin JG, Burns MP. The GABA(A) receptor agonist THIP ameliorates specific behavioral deficits in the mouse model of Fragile X syndrome. *Dev Neurosci* (2011) **33**:395–403. doi:10.1159/00032884
117. Wang LW, Berry-Kravis E, Hagerman RJ. Fragile X: leading the way for targeted treatments in autism. *Neurotherapeutics* (2010) **7**:264–74. doi:10.1016/j.nurt.2010.05.005
118. Zoghbi HY, Bear MF. Synaptic dysfunction in neurodevelopmental disorders associated with autism and intellectual disabilities. *Cold Spring Harb Perspect Biol* (2012) **4**:a009886. doi:10.1101/cshperspect.a009886
119. Berry-Kravis EM, Hessel D, Rathmell B, Zarevics P, Cherubini M, Walton-Bowen K, et al. Effects of STX209 (arbaclofen) on neurobehavioral function in children and adults with Fragile X syndrome: a randomized, controlled, phase 2 trial. *Sci Transl Med* (2012) **4**:152ra127. doi:10.1126/scitranslmed.3004214
120. Oberman LM. mGluR antagonists and GABA agonists as novel pharmacological agents for the treatment of autism spectrum disorders. *Expert Opin Investig Drugs* (2012) **21**(12):1819–25. doi:10.1517/13543784.2012.729819
121. Marrosu F, Marrosu G, Rachel MG, Biggio G. Paradoxical reactions elicited by diazepam in children with classic autism. *Funct Neurol* (1987) **2**:355–61.
122. Hochman DW. The extracellular space and epileptic activity in the adult brain: explaining the antiepileptic effects of furosemide and bumetanide. *Epilepsia* (2012) **1**:18–25. doi:10.1111/j.1528-1167.2012.03471.x
123. Lemonnier E, Ben-Ari Y. The diuretic bumetanide decreases autistic behaviour in five infants treated during 3 months with no side effects. *Acta Paediatr* (2010) **99**:1885–8. doi:10.1111/j.1651-2227.2010.01933.x
124. Lemonnier E, Degrez C, Phelep M, Tyzio R, Josse F, Grandgeorge M, et al. A randomised controlled trial of bumetanide in the treatment of autism in children. *Transl Psychiatry* (2012) **2**:e202. doi:10.1038/tp.2012.124
125. Hadjikhani N, Zürcher NR, Rogier O, Ruest T, Hippolyte L, Ben-Ari Y, et al. Improving emotional face perception in autism with diuretic bumetanide: a proof-of-concept behavioral and functional brain imaging pilot study. *Autism* (2013). doi:10.1177/1362361313514141

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 19 May 2014; accepted: 23 June 2014; published online: 08 July 2014.

Citation: Cellot G and Cherubini E (2014) GABAergic signaling as therapeutic target for autism spectrum disorders. *Front. Pediatr.* 2:70. doi: 10.3389/fped.2014.00070

This article was submitted to *Child and Neurodevelopmental Psychiatry*, a section of the journal *Frontiers in Pediatrics*.

Copyright © 2014 Cellot and Cherubini. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.