

# UC Santa Cruz

## UC Santa Cruz Previously Published Works

### Title

GAP PHENOMENA AND CURVATURE ESTIMATES FOR CONFORMALLY COMPACT EINSTEIN MANIFOLDS

### Permalink

<https://escholarship.org/uc/item/6tw8n95x>

### Journal

TRANSACTIONS OF THE AMERICAN MATHEMATICAL SOCIETY, 369(6)

### ISSN

0002-9947

### Authors

Li, Gang  
Qing, Jie  
Shi, Yuguang

### Publication Date

2017

### DOI

10.1090/tran/6925

Peer reviewed

# GAP PHENOMENA AND CURVATURE ESTIMATES FOR CONFORMALLY COMPACT EINSTEIN MANIFOLDS

GANG LI<sup>†</sup>, JIE QING<sup>\*</sup> AND YUGUANG SHI<sup>‡</sup>

**ABSTRACT.** In this paper we first use the result in [12] to remove the assumption of the  $L^2$  boundedness of Weyl curvature in the gap theorem in [9] and then obtain a gap theorem for a class of conformally compact Einstein manifolds with very large renormalized volume. We also use the blow-up method to derive curvature estimates for conformally compact Einstein manifolds with large renormalized volume. The second part of this paper is on conformally compact Einstein manifolds with conformal infinities of large Yamabe constants. Based on the idea in [15] we manage to give the complete proof of the relative volume inequality (1.9) on conformally compact Einstein manifolds. Therefore we obtain the complete proof of the rigidity theorem for conformally compact Einstein manifolds in general dimensions with no spin structure assumption (cf. [29, 15]) as well as the new curvature pinch estimates for conformally compact Einstein manifolds with conformal infinities of very large Yamabe constant. We also derive the curvature estimates for conformally compact Einstein manifolds with conformal infinities of large Yamabe constant.

## 1. INTRODUCTION

The study of conformally compact Einstein manifolds is fundamental in establishing mathematical theory of the so-called AdS/CFT correspondence proposed in the theory of quantum gravity in theoretic physics. It is well understood that there is the rigidity phenomenon for conformally compact Einstein manifolds, that is, a conformally compact Einstein manifold whose conformal infinity is the conformal round sphere has to be the hyperbolic space [3, 29]. On the other hand, in [18, 6], it was shown that for each conformal sphere that is sufficiently close to the conformal round sphere there exists a conformally compact Einstein metric on the ball whose conformal infinity is the given conformal sphere. Those conformally compact Einstein metrics constructed in [18, 6] are automatically close to the hyperbolic space in some appropriate way. In an attempt to understand if those conformally compact Einstein metrics given in [18, 6] are the unique ones, in this paper, we describe some gap phenomena in terms of renormalized volumes and derive curvature estimates when either the renormalized volume is close to that of hyperbolic space or the Yamabe constant of the conformal infinity is close to that of conformal round sphere.

---

2010 *Mathematics Subject Classification.* Primary 53C25; Secondary 58J05.

<sup>†</sup> Research supported by China Postdoctoral Science Foundation Grant 2014M550540.

<sup>\*</sup> Research supported by NSF grant DMS-1303543.

<sup>‡</sup> Research supported by NSF grant of China 10990013.

The gap theorem in this paper for renormalized volumes of conformally compact Einstein 4-manifolds grows out of the gap theorem in [9] for closed Bach flat 4-manifolds. As a consequence of recent remarkable work of Cheeger and Naber [12] we first remove the dependence of the  $L^2$  of the Weyl curvature from the gap theorem in [9] and obtain the following:

**Theorem 1.1.** *There exists a positive small number  $\epsilon$  such that a closed Bach flat 4-manifold  $(M, g)$  of positive Yamabe type has to be conformally equivalent to the round 4-sphere, if*

$$\int_M \sigma_2(A[g])dV_g \geq (1 - \epsilon)16\pi^2,$$

where  $\sigma_2(A[g])$  is the second symmetric function of the eigenvalues of the Schouten curvature tensor  $A[g] = \frac{1}{n-2}(\text{Ric} - \frac{R}{2(n-1)}g)$  of the metric  $g$ .

Our proof of Theorem 1.1 is slightly different from that in [9]. Our approach instead relies on the control of the Yamabe constant. The Yamabe constant is defined as follows:

$$(1.1) \quad Y(M, g) = \inf \left\{ \frac{\int_M \left( \frac{4(n-1)}{n-2} |\nabla u|^2 + R[g]u^2 \right) dv[g]}{\left( \int_M u^{\frac{2n}{n-2}} dv[g] \right)^{\frac{n-2}{n}}} : u \in C_c^\infty(M) \setminus \{0\} \right\},$$

where  $C_c^\infty(M)$  is the space of smooth functions with compact support in  $M$ . It is easily seen that  $Y(M, g)$  is a conformal invariant when  $(M, g)$  is compact with no boundary, in which case we denote it by  $Y(M, [g])$  instead. On the other hand, on a complete non-compact manifold,  $Y(M, g)$  is rather a local Yamabe constant. For instance

$$Y(\mathbb{R}^n, g_E) = Y(\mathbb{H}^n, g_{\mathbb{H}}) = Y(\mathbb{S}^n, [g_{\mathbb{S}}]).$$

The Yamabe constant is as convenient to use as the Euclidean volume growth bound when rescaling metrics. Particularly we observe the following fact:

**Lemma 1.2.** *Suppose that  $(X^4, g)$  is a complete non-compact Ricci flat manifold. And suppose that*

$$Y(M, g) > 2^{-\frac{1}{2}} Y(\mathbb{R}^4, g_E).$$

*Then  $(X^4, g)$  is isometric to the Euclidean 4-space  $(\mathbb{R}^4, g_E)$ .*

Lemma 1.2 is the special case of Lemma 3.4 when  $n = 4$ . The proof of this fact is rather straightforward using the end analysis based on [1, 5] and the simple consequence Lemma 3.3 from the new remarkable work in [12] in dimension 4.

To see the control of Yamabe constant from the integral of  $\sigma_2(A)$ , with the Yamabe metric  $g_Y$  on a compact manifold, we have

$$(1.2) \quad \int_M (\sigma_2(A)dv)[g_Y] = \int_M \left( \frac{1}{24}R^2 - \frac{1}{2}|\text{Ric}^\circ|^2 \right) dv[g_Y] \leq \frac{1}{24}(Y(M, [g]))^2.$$

It is known from [17] that, on a conformally compact Einstein 4-manifold  $(X^4, g^+)$  with the conformal infinity  $(\partial X, [\hat{g}])$ ,

$$(1.3) \quad \text{Vol}(\{x > \epsilon\}) = \frac{1}{3}\text{Vol}(\partial X, \hat{g})\epsilon^{-3} - \frac{1}{8} \int_{\partial X} (Rdv)[\hat{g}]\epsilon^{-1} + V(X^4, g^+) + o(1)$$

where  $R[\hat{g}]$  is the scalar curvature of the metric  $\hat{g}$  and  $x$  is the geodesic defining function associated with a representative  $\hat{g}$  of the conformal infinity. It turns out that  $V(X^4, g^+)$  in (1.3) is independent of representatives and is called the renormalized volume [20, 17]. The expansion (1.3) uses the expansion of the Einstein metric  $g^+$  given in [16]

$$(1.4) \quad g^+ = x^{-2}(dx^2 + g_x) = x^{-2}(dx^2 + \hat{g} + g^{(2)}x^2 + g^{(3)}x^3 + o(x^3))$$

where  $g^{(2)}$  is a curvature tensor of  $\hat{g}$  while  $g^{(3)}$  is non-local. Moreover, in [2] (see also [10]), it is shown that

$$(1.5) \quad 8\pi^2\chi(X) = \frac{1}{4} \int_X (|W|^2 dv)[g^+] + 6V(X^4, g^+)$$

for a conformally compact Einstein 4-manifold  $(X^4, g^+)$ . Consequently, one knows

$$(1.6) \quad \int_{X^4} (\sigma_2(A)dv)[\bar{g}] = 6V(X^4, g^+),$$

where  $\bar{g} = x^2g^+$  is a conformal compactification. Therefore, given a conformally compact Einstein manifold, one may control the Yamabe constant for compactifications from the renormalized volume. Following the idea in [11], one may consider the doubling of the compactified manifold from a conformally compact Einstein manifold and obtain the corresponding gap theorem.

**Theorem 1.3.** *There exists a small positive number  $\epsilon$  such that a conformally compact Einstein 4-manifold  $(X^4, g^+)$  with the conformal infinity of positive Yamabe type has to be isometric to the hyperbolic space, if its renormalized volume satisfies*

$$(1.7) \quad V(X^4, g^+) \geq (1 - \epsilon)\frac{4\pi^2}{3} = (1 - \epsilon)V(\mathbb{H}^4, g_{\mathbb{H}})$$

and the non-local term  $g^{(3)}$  in (1.4) for  $g^+$  vanishes.

Theorem 1.3 indicates that the hyperbolic space is the only ‘‘critical point’’ of the renormalized volume among all conformally compact Einstein manifolds that satisfy (1.7), since the Euler-Lagrangian equation for the renormalized volume is  $g^{(3)} = 0$  by the calculation made in [2]. On the other hand, Theorem 1.3 clearly does not hold if one drops the assumption  $g^{(3)} = 0$ . In fact the conformally compact Einstein metrics constructed in [18, 6] provide plenty of examples of conformally compact Einstein metrics that satisfy (1.7) for arbitrarily small  $\epsilon$  and with conformal infinities of positive Yamabe type. Nevertheless, one obtains the following curvature estimate for conformally compact Einstein metrics when the renormalized volume is large enough.

**Theorem 1.4.** *For any  $0 < \epsilon < \frac{1}{2}$ , there is a positive constant  $C$  such that, on a conformally compact Einstein 4-manifold  $(X, g^+)$ ,*

$$\|Rm\|[g^+] \leq C,$$

where  $Rm[g^+]$  is the Riemann curvature tensor of the metric  $g^+$ , provided that the conformal infinity is of positive Yamabe type and (1.7) holds.

Next we focus our attention to those conformally compact Einstein manifolds whose conformal infinities have large Yamabe constants. It is a very original idea in [15] to use the Yamabe constant of the conformal infinity to control the relative volume growth of geodesic balls in

conformally compact Einstein manifolds. We recognize the important contribution of [15], but are compelled to present a complete and correct proof of the following:

**Theorem 1.5.** *Suppose that  $(X^n, g^+)$  is an AH manifold of  $C^3$  regularity and with the conformal infinity of positive Yamabe type. Let  $p \in X^n$  be a fixed point. Assume*

$$(1.8) \quad Ric[g^+] \geq -(n-1)g^+ \text{ and } R[g^+] + n(n-1) = o(e^{-2t})$$

for the distance function  $t$  from  $p$ . Then

$$(1.9) \quad \left( \frac{Y(\partial X, [\hat{g}])}{Y(\mathbb{S}^{n-1}, [g_{\mathbb{S}}])} \right)^{\frac{n-1}{2}} \leq \frac{Vol(\partial B_{g^+}(p, t))}{Vol(\partial B_{g_{\mathbb{H}}}(0, t))} \leq \frac{Vol(B_{g^+}(p, t))}{Vol(B_{g_{\mathbb{H}}}(0, t))} \leq 1,$$

where  $B_{g^+}(p, t)$  and  $B_{g_{\mathbb{H}}}(0, t)$  are geodesic balls.

The  $C^3$  regularity is used to construct the geodesic defining function  $x$  and  $C^2$  conformal compactification  $\bar{g} = x^2 g^+$ , for each given representative  $\hat{g}$ . To start the proof we first need to clear a technical issue.

**Lemma 1.6.** *Suppose that  $(X^n, g^+)$  is AH of  $C^2$  regularity and that  $x$  is a defining function. Assume*

$$(1.10) \quad Ric[g^+] \geq -(n-1)g^+ \text{ and } R[g^+] + n(n-1) = o(x)$$

Then there is a constant  $C_0$  such that

$$(1.11) \quad |K[g^+] + 1| \leq C_0 x^2$$

for any sectional curvature  $K$ .

It takes substantial arguments to finish the proof of Theorem 1.5 based on the idea presented in [15]. The first issue is that (4.8) (cf. (3.17) in [15]) may not be available as claimed in [15], since the distance function  $t$  is only Lipschitz in general. We devote Section 5 to solve this issue by the careful study of cut loci based on [27, 22]. The second issue is that the estimate (4.1) (cf. (2.3) in [15]) is not known to hold without assuming the convexity of the geodesic spheres (cf. [30, 21]). We devote Section 6 to derive the total scalar curvature estimate (4.10) without (4.1). Our argument in Section 6 uses more delicate analysis of the Riccati equations on AH manifolds and volume estimates along geodesics where the mean curvature of the geodesic sphere is small.

As argued in [15], Theorem 1.5 implies the rigidity of conformally compact Einstein manifolds for any dimension, which are the cases where the conformal infinity is exactly the round spheres (cf. [26, 3, 24, 29, 32, 14, 7, 30, 15]).

**Theorem 1.7.** *Suppose that  $(X^n, g^+)$  is AH of  $C^3$  regularity and that (1.8) holds. Then  $(X^n, g^+)$  is isometric to the hyperbolic space  $(\mathbb{H}^n, g_{\mathbb{H}})$ , provided that the conformal infinity  $(\partial X, [\hat{g}])$  is the round conformal sphere.*

Notice that, when the conformal infinity is the round sphere, a conformally compact Einstein metric  $g^+$  is always smooth according to [13], provided that it is at least of  $C^2$  regularity. Therefore Theorem 1.7 does cover the most general rigidity theorem for conformally compact Einstein manifolds whose conformal infinity is the round sphere in any dimension. In this paper we also deduce from the relative volume growth estimates (1.9) the following interesting curvature pinch estimates.

**Theorem 1.8.** *For any  $\epsilon > 0$ , there exists  $\delta > 0$ , for any conformally compact Einstein manifold  $(X^n, g^+)$  ( $n \geq 4$ ), one gets*

$$(1.12) \quad |K[g^+] + 1| \leq \epsilon,$$

for all sectional curvature  $K$  of  $g^+$ , provided that

$$Y(\partial X, [\hat{g}]) \geq (1 - \delta)Y(S^{n-1}, [g_{\mathbb{S}}]).$$

Particularly, any conformally compact Einstein manifold with its conformal infinity of Yamabe constant sufficiently close to that of the round sphere is necessarily negatively curved.

This result is even more interesting because it gives the curvature pinch estimate which only relies on the Yamabe constant of the conformal infinity. Particularly, from Theorem 1.8 we now know that any conformally compact Einstein manifold whose conformal infinity is prescribed as a conformal sphere that is sufficiently close to the round conformal sphere is negatively curved, which was only known to be true for those conformally compact Einstein manifolds constructed in [18, 6].

As a consequence of the proof of Theorem 1.8 we also get:

**Corollary 1.9.** *For  $L > 0$ ,  $\tau > \frac{1}{2}$  and  $n \geq 4$ , there is a number  $C$  such that, on any conformally compact Einstein manifold  $(X^n, g^+)$ ,*

$$|W|[g^+] \leq C$$

provided that

$$Y(\partial X, [\hat{g}]) \geq \tau^{\frac{2}{n-1}} Y(\mathbb{S}^n, [g_{\mathbb{S}}])$$

and

$$\int_{X^n} (|W|^{\frac{n}{2}} dv)[g^+] \leq L$$

when  $n \geq 5$ .

The organization of this paper is as follows: in Section 2, we will introduce some basics about AH manifolds and conformally compact Einstein manifolds. Particularly we will prove Lemma 1.6. In Section 3 we use the renormalized volume to control the Yamabe constant of the conformally compact Einstein 4-manifolds and prove Theorem 1.3 of the gap phenomenon and Theorem 1.4 of curvature estimates. In Section 4 We will first sketch a proof of Theorem 1.5 based on the idea in [15]. We will identify the incompleteness and incorrectness of the arguments in [15] in the sketch of the proof of Theorem 1.5. Then we will use Theorem 1.5 to obtain Theorem 1.8 and Corollary 1.9. In Section 5 and Section 6 we will resolve the gaps identified in Section 4 and present the complete and correct proof of Theorem 1.5 with details.

**Acknowledgment** The authors are very grateful to Beijing International Center for Mathematical Research (BICMR) for the hospitality. Most of research in this paper was conducted during the summer when the authors visit BICMR. The first author would like to thank Professor Matthew Gursky for his interest of the problem and encouragement

## 2. PRELIMINARIES

Let us recall some basics about AH manifolds and conformally compact Einstein manifolds. First we use the following definition for conformally compact Einstein manifolds.

**Definition 2.1.** *Suppose that  $X^n$  is the interior of a smooth compact manifold  $\overline{X^n}$  with boundary  $\partial X^{n-1}$ . A Riemannian metric  $g^+$  on  $X^n$  is said to be conformally compact of  $C^{k,\alpha}$  regularity if, for a smooth defining function  $x$  for the boundary  $\partial X^{n-1}$  in  $\overline{X^n}$ ,  $\bar{g} = x^2 g^+$  can be extended to a  $C^{k,\alpha}$  Riemannian metric on  $\overline{X^n}$ . If, in addition,  $|dx|_{x^2 g^+}|_{x=0} = 1$ , then we say  $(X^n, g^+)$  is asymptotically hyperbolic (AH in short) of  $C^{k,\alpha}$  regularity. And if, in addition,  $g^+$  is at least of  $C^2$  regularity and Einstein, that is,*

$$(2.1) \quad \text{Ric}[g^+] = -(n-1)g^+,$$

then we say  $(X^n, g^+)$  is a conformally compact Einstein manifold.

A smooth defining function  $x$  for the boundary  $\partial X$  in a smooth manifold  $\overline{X^n}$  is a smooth nonnegative function from  $\overline{X^n}$  such that

- $x > 0$  in the interior  $X^n$ ;
- $x = 0$  on the boundary  $\partial X^{n-1}$ ;
- $dx \neq 0$  on the boundary  $\partial X^{n-1}$ .

The compactification  $\bar{g}$  induces a metric  $\hat{g}$  on the boundary  $\partial X^{n-1}$  and changes conformally when the defining function  $x$  varies. Hence a conformally compact metric  $g^+$  always induces a conformal structure  $[\hat{g}]$  on the boundary  $\partial X^{n-1}$ . The conformal manifold  $(\partial X^{n-1}, [\hat{g}])$  is called the conformal infinity of the conformally compact manifold  $(X^n, g^+)$ .

Before we recall basic properties of conformally compact Einstein manifolds, we give a proof of Lemma 1.6 based on the proof of Lemma 3.1 in [13].

*Proof of Lemma 1.6* The first step completely follows the proof of Lemma 3.1 in [13] and concludes that, there is a coordinate at infinity (up to a  $C^3$  collar diffeomorphism in the language in [13]) such that, for a defining function  $x$ ,

$$(2.2) \quad \bar{g} = x^2 g^+ = dx^2 + \hat{g} + g^{(1)}x + O(x^2) \in C^2(\overline{X^n})$$

for some symmetric 2-tensor  $g^{(1)}$  and a representative  $\hat{g} = x^2 g^+|_{\Gamma \partial X}$  on  $\partial X$ .

We then calculate the transforms of Riemann curvature, Ricci curvature and scalar curvature based on  $g^+ = x^{-2} \bar{g}$ .

$$R_{ijkl}[g^+] = -(g_{ik}^+ g_{jl}^+ - g_{jk}^+ g_{il}^+) + x^{-1} (g_{ik}^+ \nabla_j^{\bar{g}} \nabla_l^{\bar{g}} x + g_{jl}^+ \nabla_i^{\bar{g}} \nabla_k^{\bar{g}} x - g_{il}^+ \nabla_j^{\bar{g}} \nabla_k^{\bar{g}} x - g_{jk}^+ \nabla_i^{\bar{g}} \nabla_l^{\bar{g}} x) + O(x^2),$$

$$R_{ik}[g^+] = -(n-1)g^+ + x^{-1} ((n-2) \nabla_i^{\bar{g}} \nabla_k^{\bar{g}} x + \Delta[\bar{g}] x \bar{g}_{ik}) + O(x^2),$$

and

$$R[g^+] = -n(n-1) + 2x(n-1) \Delta[\bar{g}] x + O(x^2).$$

In the same time we can calculate

$$\nabla_i^{\bar{g}} \nabla_k^{\bar{g}} x = \frac{1}{2} \partial_x \bar{g}_{ik} + O(x).$$

Therefore the condition (1.10) are translated to

$$(n-2)\partial_x \bar{g}_{ik} + \bar{g}^{jl}\partial_x \bar{g}_{jl}\bar{g}^{ik} \geq 0 \text{ and } \bar{g}^{jl}\partial_x g_{jl} = 0 \text{ at } x=0,$$

which implies  $g^{(1)} = 0$  and thus (1.11).  $\square$

The fundamental properties of conformally compact Einstein 4-manifolds that are useful to us are summarized in the following:

**Lemma 2.2.** ([16, 17, 13, 25]) *Let  $(X^4, g^+)$  be a conformally compact Einstein manifold and  $x$  be the geodesic defining function associated with a representative  $\hat{g}$  of the conformal infinity  $(\partial X^3, [\hat{g}])$ . In a neighborhood of the infinity*

$$(2.3) \quad g = x^{-2}\tilde{g} = x^{-2}(dx^2 + g_x)$$

with the expansion

$$(2.4) \quad g_x = \hat{g} + g^{(2)}x^2 + g^{(3)}x^3 + \sum_{k=4}^m g^{(k)}x^k + o(x^m)$$

for any  $m \geq 4$ , where  $g^{(k)}$  is a symmetric  $(0, 2)$  tensor on  $\partial X$  for all  $k$  and  $g^{(3)}$  is the so-called non-local term. Moreover, the expansion (2.4) only has even power terms, when  $g^{(3)}$  vanishes.

As a consequence of the expansion (2.4) one gets the following volume expansion:

**Lemma 2.3.** ([20, 17]) *Let  $(X^4, g^+)$  be a conformally compact Einstein manifold and  $x$  be the geodesic defining function associated with a representative  $\hat{g}$  of the conformal infinity  $(\partial X^3, [\hat{g}])$ . One has*

$$\text{Vol}(\{x > \epsilon\}) = \frac{1}{3}\text{Vol}(\partial X, \hat{g})\epsilon^{-3} - \frac{1}{8}\int_{\partial X} (Rdv)[\hat{g}]\epsilon^{-1} + V(X^4, g^+) + o(1).$$

More importantly  $V(X^4, g^+)$  is independent of the choice of representative  $\hat{g}$  and is called the renormalized volume.

To appreciate the global invariant  $V(X^4, g^+)$  of a conformally compact Einstein 4-manifold  $(X^4, g^+)$  we recall the Gauss-Bonnet formula observed in [2, 11].

**Lemma 2.4.** *Let  $(X^4, g^+)$  be a conformally compact Einstein 4-manifold. Then*

$$(2.5) \quad 8\pi^2\chi(X) = \frac{1}{4}\int_X (|W|^2 dv)[g^+] + 6V(X^4, g^+)$$

Comparing the Gauss-Bonnet formula (2.5) with the Gauss-Bonnet formula for compact 4-manifold  $(X, \bar{g})$  with totally geodesic boundary:

$$(2.6) \quad 8\pi^2\chi(X) = \frac{1}{4}\int_X (|W|^2 dv)[\bar{g}] + \int_X (\sigma_2(A)dv)[\bar{g}],$$

we arrive at

$$(2.7) \quad \int_X (\sigma_2(A)dv)[\bar{g}] = 6V(X^4, g^+),$$

for any compactification  $\bar{g} = x^2g^+$ .



To discuss the conformal gap theorem in [9] we recall the definition of Bach curvature tensor and  $\epsilon$ -regularity theorem for Bach flat Yamabe metrics. On 4-manifolds the Bach curvature tensor is a symmetric 2-tensor

$$(2.8) \quad B_{ij} = W_{kijl}{}^{lk} + \frac{1}{2}R^{kl}W_{kijl}.$$

The  $\epsilon$ -regularity theorem has been established in [31] as follows:

**Lemma 2.5.** ([31]) *Suppose that  $(M^4, g)$  is a Bach flat 4-manifold with Yamabe constant  $Y > 0$  and  $g$  is a Yamabe metric. Then there exist positive numbers  $\tau_k$  and  $C_k$  depending on  $Y$  such that, for each geodesic ball  $B_{2r}(p)$  centered at  $p \in M$ , if*

$$\int_{B_{2r}(p)} |Rm|^2 dv \leq \tau_k,$$

then

$$(2.9) \quad \sup_{B_r(p)} |\nabla^k Rm| \leq \frac{C_k}{r^{2+k}} \left( \int_{B_{2r}(p)} |Rm|^2 dv \right)^{\frac{1}{2}}.$$

### 3. CONFORMALLY COMPACT EINSTEIN MANIFOLDS WITH LARGE RENORMALIZED VOLUMES

In this section we use Cheeger and Naber's result in [12] in dimension 4 to drop the  $L^2$  boundedness condition in the conformal gap theorem in [9] and prove a version of conformal gap theorem of renormalized volumes on conformally compact Einstein manifolds. First we state Cheeger and Naber's result.

**Theorem 3.1.** (Theorem 1.5 [12]) *There exists  $C = C(v)$  such that, if  $M^4$  satisfies  $|\text{Ric}_{M^4}| \leq 3$  and  $\text{Vol}(B_1(p)) > v > 0$ , then*

$$(3.1) \quad \int_{B_1(p)} |Rm|^2 dV \leq C,$$

where  $B_1(p)$  is a geodesic ball on  $(M^4, g)$ .

We then recall the following theorem on the end analysis in [1, 5].

**Theorem 3.2.** ([1, 5]) *Let  $(M^n, g)$  ( $n \geq 4$ ) be a complete noncompact Ricci flat Riemannian manifold satisfying that*

$$(3.2) \quad \text{Vol}(B_r) \geq Cr^n \text{ for all } r > 0,$$

$$(3.3) \quad \int_M (|Rm|^{\frac{n}{2}} dV)[g] < \infty.$$

Then given  $q \in M$ , there is an  $R_0 > 0$  such that

- $M \setminus B_{R_0}(q)$  is diffeomorphic to the cone  $(R_0, \infty) \times S^{n-1}/\Gamma$ ;
- $g = g_F + O(r^{-2})$ , where  $g_F$  is the flat metric on the cone.

Moreover, if  $M$  is simply connected at infinity, then  $(M^n, g)$  is isometric to the Euclidean space.

It turns out, as a straightforward consequence of Theorem 3.1 of Cheeger and Naber, one can drop the assumption (3.3) in Theorem 3.2 in dimension 4.

**Lemma 3.3.** *Let  $(M^4, g)$  be a complete noncompact Ricci flat Riemannian 4-manifold satisfying the Euclidean volume growth assumption (3.2). Then, given  $q \in M$ , there is an  $R_0 > 0$  such that*

- $M \setminus B_{R_0}(q)$  is diffeomorphic to the cone  $(R_0, \infty) \times S^3/\Gamma$ ;
- $g = g_F + O(r^{-2})$ , where  $g_F$  is the flat metric on the cone.

Moreover, if  $M$  is simply connected at infinity, then  $(M^4, g)$  is isometric to the Euclidean 4-space.

*Proof.* In the light of Theorem 3.2 ([1, 5]) it suffices to show that, there is a constant  $C$  such that

$$\int_{B_R} (|Rm|^2 dv)[g] \leq C$$

for any  $R > 0$ . For any fixed  $R > 0$ , we consider the metric  $g_R = R^{-2}g$ . It is easily verified that Theorem 3.1 is applicable to  $g_R$ . Hence the proof is complete.  $\square$

Theorem 1.1 is an improved version of the conformal gap theorem in [9] based on the above Lemma 3.3. But we will present a complete proof of Theorem 1.1 that is slightly different from that in [9]. Our approach relies on the control of the Yamabe constant via (1.2) and the following observation that describes the influence of the end structure from the Yamabe constant similar to that of the lower bound of the Euclidean volume growth.

**Lemma 3.4.** *Let  $(M^n, g)$  ( $n \geq 4$ ) be a complete noncompact Ricci flat Riemannian manifold. Assume (3.3) when  $n \geq 5$ . Then  $(M^n, g)$  is isometric to the Euclidean space  $\mathbb{R}^n$ , provided that*

$$(3.4) \quad Y(M, g) \geq \tau Y(\mathbb{R}^n, g_E)$$

for some  $\tau > 2^{-\frac{2}{n}}$ .

*Proof.* First of all, according to Lemma 3.2 in [19], one indeed has a lower bound for the Euclidean volume growth from the lower bound of the Yamabe constant. Then from our Lemma 3.3 and the arguments in [1, 5] it is known that the tangent cone  $(M_\infty^n, g_\infty)$  at infinity for the Ricci flat manifold  $(M^n, g)$  is a flat cone  $\mathbb{R}^n/\Gamma$ . Because the blow-down: the rescaled manifolds  $(M^n, \lambda_i^2 g, p)$ , as  $\lambda_i \rightarrow 0$  with a fixed point  $p \in M^n$ , converges to the tangent cone  $(M_\infty, g_\infty, p_\infty)$  at infinity in Cheeger-Gromov topology away from the singular point  $p_\infty$  (cf. [1, 5]). It is rather easily seen from (1.1) that the Yamabe quotient of the blow-down sequence converges to that of the tangent cone at infinity for any smooth function with compact support away from the vertex of the cone. For a smooth function with compact support in general, we simply use cut-off functions to modify it as follows. For any small  $s > 0$ , we consider cut-off functions:

$$\begin{cases} \phi = 0 & \text{when } r \leq s \\ \phi = 1 & \text{when } r \geq 2s \\ \phi \in C^\infty \end{cases}$$

We may require that  $|\nabla\phi| \leq Cs^{-1}$  for some constant  $C$ . Then, for a function  $u$  with compact support, which is smooth away from the vertex and Lipschitz across the vertex, we calculate

$$\begin{aligned} \int_{\mathbb{R}^n/\Gamma} |\nabla u|^2 &= \int_{\mathbb{R}^n/\Gamma} |\nabla(\phi u)|^2 + \int_{\mathbb{R}^n/\Gamma} (2\nabla(\phi u) \cdot \nabla((1-\phi)u) + |\nabla((1-\phi)u)|^2) \\ &= \int_{\mathbb{R}^n/\Gamma} |\nabla(\phi u)|^2 + O(s^{n-2}) \end{aligned}$$

and

$$\int_{\mathbb{R}^n/\Gamma} u^{\frac{2n}{n-2}} = \int_{\mathbb{R}^n/\Gamma} (\phi u)^{\frac{2n}{n-2}} + O(s^n)$$

as  $s \rightarrow 0$ . Therefore, due to the scaling invariance of the Yamabe constant, we also know that

$$Y(M_\infty^n, g_\infty) \geq \liminf Y(M, \lambda_i g) \geq \tau Y(\mathbb{R}^n, g_E)$$

for some  $\tau > 2^{-\frac{2}{n}}$  from our assumption (3.4). Suppose that

$$D_\Gamma : \mathbb{R}^n \rightarrow \mathbb{R}^n/\Gamma$$

is the desingularization of the cone metric. Then it is easily seen that

$$(3.5) \quad \frac{\int_{\mathbb{R}^n} \frac{4(n-1)}{n-2} (|\nabla u \circ D_\Gamma|^2 dv)[g_E]}{(\int_{\mathbb{R}^n} (u \circ D_\Gamma)^{\frac{2n}{n-2}} dv[g_E])^{\frac{n-2}{n}}} = 2^{\frac{2}{n}} \frac{\int_{M_\infty^n} \frac{4(n-1)}{n-2} (|\nabla u|^2 dv)[g_\infty]}{(\int_{M_\infty^n} u^{\frac{2n}{n-2}} dv[g_\infty])^{\frac{n-2}{n}}}.$$

Hence, when using appropriately modifications of the standard function

$$u \circ D_\Gamma = \left( \frac{2}{1 + |x|^2} \right)^{\frac{n-2}{2}}$$

in (3.5), one easily gets

$$Y(M_\infty^n, g_\infty) \leq |\Gamma|^{-\frac{2}{n}} Y(\mathbb{R}^n, g_E),$$

which, comparing to (3.4), forces the group  $\Gamma$  to be trivial and the Ricci flat manifold  $(M^n, g)$  to be isometric to the Euclidean space  $(\mathbb{R}^n, g_E)$  according Lemma 3.3 ([12]) in dimension 4 and Theorem 3.2 ([1, 5]) in general dimensions. So the proof is complete.  $\square$

Now we are ready to give the proof of our Theorem 1.1:

*Proof of Theorem 1.1:* If not, then there exists a sequence of Bach flat 4-manifolds  $(M_j^4, (g_Y)_j)$ , that are not conformally equivalent to the round 4-sphere, where  $(g_Y)_j$  is the Yamabe metric with  $R[(g_Y)_j] = 12$  and

$$\int_M (\sigma_2(A) dv)[(g_Y)_j] = (1 - \epsilon_j)^2 16\pi^2 \rightarrow 16\pi^2$$

as  $j \rightarrow \infty$ . Hence, in the light of (1.2), we have

$$(3.6) \quad (1 - \epsilon_j) Y(\mathbb{S}^4, [g_\mathbb{S}]) \leq Y(M_j^4, [(g_Y)_j]) \leq Y(\mathbb{S}^4, [g_\mathbb{S}])$$

and

$$(3.7) \quad \int_{M_j} (|\mathring{\text{Ric}}|^2 dv)[(g_Y)_j] \leq \frac{1}{12} \epsilon_j \rightarrow 0.$$

Here we use the fact that, on the round sphere  $(\mathbb{S}^4, g_\mathbb{S})$ ,

$$\int_{\mathbb{S}^4} (\sigma_2(A) dv)[g_\mathbb{S}] = \frac{1}{24} Y(\mathbb{S}^4, [g_\mathbb{S}])^2.$$

From the argument in [9], to get a contradiction it suffices to show that  $M_j^4$  is diffeomorphic to  $\mathbb{S}^4$  for a subsequence of  $j$ . More precisely, here we use the following interesting result from [9]:

**Lemma 3.5.** ([9]) *There is  $\epsilon > 0$  such that a Bach flat metric  $g$  on 4-sphere  $\mathbb{S}^4$  is conformal to the standard round metric  $g_{\mathbb{S}}$ , provided that*

$$\int_{\mathbb{S}^4} (|W|^2 dv)[g] \leq \epsilon.$$

The rest of the proof of Theorem 1.1 follows from a more or less standard rescaling argument based on our Lemma 3.4. We first derive a contradiction if there was curvature blow-up. Again, we use Lemma 3.2 in [19] to get the uniform lower bound on the Euclidean volume growth for such sequence of manifolds. Then, one stands at the point of curvature blow-up, that is,  $p_j \in M_j$  such that

$$\lambda_j = |Rm|(p_j)[(g_Y)_j] = \max_{M_j} |Rm|[(g_Y)_j] \rightarrow \infty$$

and considers the sequence of pointed Riemannian manifold  $(M_j, g_j, p_j)$  with the rescaled metric  $g_j = \lambda_j^2 (g_Y)_j$ . Therefore, according to the curvature estimates established, for example, in [31], one derives a subsequence that converges to complete non-compact manifold  $(M_\infty, g_\infty, p_\infty)$  in Cheeger-Gromov topology. As the consequences of (3.6) and (3.7) one knows that

- $Y(M_\infty, g_\infty) = Y(\mathbb{S}^n, [g_{\mathbb{S}}]) = Y(\mathbb{R}^n, g_E)$  and
- $\text{Ric}[g_\infty] = 0$  and  $R[g_\infty] = 0$ .

Here we use the argument similar to that in the proof in Lemma 3.4 to derive the equality  $Y(M_\infty, g_\infty) = Y(\mathbb{R}^n, g_E)$ . Therefore  $(M_\infty, g_\infty)$  is isometric to the Euclidean 4-space according to Lemma 3.4 in the same time  $|Rm|(p_\infty)[g_\infty] = 1$ , which is a contradiction. On the other hand, if there is no curvature blow-up for the sequence  $(M_j, (g_Y)_j)$ , then, the same compactness argument implies that there would be a subsequence that converges to the round sphere in Cheeger-Gromov topology, which is impossible due to Lemma 3.5. Thus the proof of Theorem 1.1 is complete.  $\square$

Next we use the facts collected in the section of preliminaries to give a proof of Theorem 1.3:

*Proof of Theorem 1.3:* Since  $g^{(3)} = 0$ , from the expansion (2.4), the doubling

$$(X_D, \tilde{g}) = (\bar{X} \bigcup \bar{X}, \tilde{g})$$

is a smooth Bach flat 4-manifold (for more details about the doubling please see [10], which uses [8]). In fact, it is also shown in [10] that the doubling  $(X_D, \tilde{g})$  is of positive Yamabe type from the assumption that the conformal infinity  $(\partial X^3, [\hat{g}])$  is of positive Yamabe type. In the mean time we recall from (2.7) that

$$\int_{X_D} (\sigma_2(A) dv)[\tilde{g}] = 2 \int_X (\sigma_2(A) dv)[\tilde{g}] = 12V(X^4, g^+)$$

to conclude that

$$\int_{X_D} (\sigma_2(A) dv)[\tilde{g}] \geq (1 - \epsilon) 16\pi^2$$

by the assumption (1.7). Now one applies Theorem 1.1 to the doubling  $(X_D, \tilde{g})$  and derives that  $(X_D, \tilde{g})$  is conformally equivalent to the round 4-sphere when  $\epsilon$  is sufficiently small. Particularly one obtains that  $(X^4, g^+)$  is a simply connected Riemannian manifold of constant curvature  $-1$  (cf. [10]) when  $\epsilon$  is sufficiently small, which completes the proof.  $\square$

To derive Theorem 1.4 we carry the above rescaling scheme with Lemma 3.4 on conformally compact Einstein manifolds. In fact we are able to derive the curvature bound for conformally compact Einstein manifolds in general dimensions, to which Theorem 1.4 is a corollary.

**Theorem 3.6.** *For a constant  $B$  and a constant  $\tau > 2^{-\frac{2}{n}}$ , there exists a constant  $C = C(n, \tau, B) > 0$  such that*

$$(3.8) \quad |Rm| \leq C,$$

for any conformally compact Einstein manifold  $(X^n, g^+)$  ( $n \geq 4$ ) with

- $Y(X^n, g^+) \geq \tau Y(\mathbb{H}^n, g_{\mathbb{H}})$  and
- $\int_{X^n} (|W|^{\frac{n}{2}} dv)[g^+] \leq B$  when  $n \geq 5$ .

*Proof.* Suppose otherwise that there is a sequence of conformally compact Einstein manifolds  $(X_j^n, g_j^+)$  satisfying the assumptions in the theorem with curvature blowing up. Since conformally compact Einstein manifolds are always asymptotically hyperbolic, we may extract a sequence of points  $p_j \in X_j^n$  such that

$$\lambda_j = |Rm|(p_j)[g_j^+] = \max_{X_j^n} |Rm|[g_j^+] \rightarrow \infty,$$

and consider the pointed rescaled manifolds  $(X_j^n, \lambda_j g_j^+, p_j)$ . It is then easily seen that there is a subsequence  $(X_j^n, g_j^+, p_j)$  converges in Cheeger-Gromov topology to a complete non-compact Ricci flat manifold  $(X_\infty^n, g_\infty, p_\infty)$  with

- $Y(X_\infty^n, g_\infty) \geq \tau Y(\mathbb{H}^n, g_{\mathbb{H}}) = \tau Y(\mathbb{R}^n, g_E)$ ;
- $|Rm|(p_\infty)[g_\infty] = 1$ ;
- $\int_{X_\infty^n} (|W|^{\frac{n}{2}} dv)[g_\infty] \leq B$  when  $n \geq 5$ ,

which is a contradiction in the light of Lemma 3.4. □

**Corollary 3.7.** *For  $\epsilon < \frac{1}{2}$ , there exists a number  $C > 0$  such that*

$$|Rm|[g^+] \leq C,$$

for any conformally compact 4-manifold  $(X^4, g^+)$  with conformal infinity of positive Yamabe type and

$$V(X^4, g^+) \geq (1 - \epsilon) \frac{4\pi^2}{3} = (1 - \epsilon) V(\mathbb{H}^4, g_{\mathbb{H}}).$$

*Proof.* It suffices to verify that

$$(3.9) \quad Y(X^4, g^+) \geq (1 - \epsilon)^{\frac{1}{2}} Y(\mathbb{H}^4, g_{\mathbb{H}}).$$

From (3.6) in the proof of Lemma 3.4 we know

$$Y(X_D, \tilde{g}) \geq (1 - \epsilon)^{\frac{1}{2}} Y(\mathbb{H}^4, g_{\mathbb{H}}),$$

which implies (3.9) by the conformal invariance of the Yamabe constant. □

Clearly, the local Yamabe constant  $Y(X^4, g^+)$  for a conformally compact Einstein 4-manifold approaches that of the hyperbolic space as the renormalized volume  $V(X^4, g^+)$  approaches that of the hyperbolic space. Before we end this section we state an easy observation based on the construction of Aubin on the impact to local Yamabe constant from local geometry in higher dimensions (cf. Paragraph 6.10 in [4, 23]).

**Proposition 3.8.** *For any  $\epsilon > 0$ , there is  $\delta > 0$  such that, for any section curvature  $K$  at any point on any conformally compact Einstein manifold  $(X^n, g^+)$  ( $n \geq 6$ ), one has*

$$|K + 1| \leq \epsilon,$$

*provided that*

$$Y(X^n, g^+) \geq (1 - \delta)Y(\mathbb{H}^n, g_{\mathbb{H}}).$$

*Proof.* Assume otherwise, for some  $\epsilon_0 > 0$ , there is a sequence of conformally compact Einstein manifolds  $(X_j^n, g_j^+)$  and a sequence  $\delta_j \rightarrow 0$  such that

$$Y(X_j^n, g_j^+) \geq (1 - \delta_j)Y(\mathbb{H}^n, g_{\mathbb{H}}),$$

but

$$|K + 1| > \epsilon_0,$$

for a sectional curvature  $K$  at some point on  $X_j^n$ . We are going to derive contradictions in two steps. First, if there is a subsequence  $(X_j^n, g_j^+)$  whose sectional curvature is not bounded as  $j \rightarrow \infty$  (for convenience, we continue to use the same index  $j$ ), then we may rescale the metrics

$$\tilde{g}_j = \lambda_j^2 g_j^+$$

for  $\lambda_j = |W|(p_j)[g_j^+] = \max_{X_j^n} |W|[g_j^+]$ . Then, due to the curvature estimates for Einstein manifolds, it is easily seen that, at least for a subsequence,  $(X_j^n, \tilde{g}_j, p_j)$  converges to a complete non-compact Ricci flat manifold  $(X_\infty^n, g_\infty, p_\infty)$  in Cheeger-Gromov topology. But  $|W|(p_\infty)[g_\infty] = 1$  contradicts with the fact that  $Y(X_\infty^n, g_\infty) = Y(\mathbb{R}^n, g_E)$  in the light of the estimate of Yamabe constant in Paragraph 6.10 in [4] (cf. also [23]).

Secondly, if there is no curvature blowup, then one may extract a subsequence such that

$$|W|(p_j)[g_j^+] \rightarrow w > 0$$

and  $(X_j^n, g_j^+, p_j)$  converges to a complete non-compact Einstein manifold  $(X_\infty^n, g_\infty^+, p_\infty)$  in Cheeger-Gromov topology. But, again,  $|W|(p_\infty)[g_\infty^+] = w > 0$  contradicts with the fact that  $Y(X_\infty^n, g_\infty^+) = Y(\mathbb{H}^n, g_{\mathbb{H}})$  in the light of the estimate of Yamabe constant in Paragraph 6.10 in [4](cf. also [23]).  $\square$

One wonders whether Proposition 3.8 still holds in dimension 4, which would be much more significant.

#### 4. CONFORMALLY COMPACT EINSTEIN MANIFOLDS WHOSE CONFORMAL INFINITIES HAVE LARGE YAMABE CONSTANTS

In this section we will first present the idea in [15] to establish the relative volume growth bounds (1.9). We recognize the contribution from [15] but are compelled to give self-contained arguments for a complete, vigorous and correct proof of (1.9) to the best of our knowledge. We will point out what are not clear and not correct in [15] and finish filling those gaps in the subsequent sections. Then we will carry out the rescaling argument to derive the curvature estimates in Theorem 1.8 in two steps similar to that in the proof of Proposition 3.8.

We recall that a Riemannian manifold  $(X^n, g^+)$  is said to be AH (short for asymptotically hyperbolic) if it is conformally compact and its curvature goes to  $-1$  at the infinity. Obviously

a conformally compact Einstein manifold is always AH. Let us consider the distance function to a given point  $p_0 \in X^n$ :

$$t = \text{dist}(\cdot, p_0)$$

and the geodesic sphere  $\Gamma_t = \{p \in X^n : \text{dist}(p, p_0) = t\}$ . The important initial step is the estimate (2.3) in Lemma 2.1 of [15], which is

$$(4.1) \quad \nabla_g^2 t(v, v) = 1 + O(e^{-\beta t}),$$

where  $v$  is any unit vector perpendicular to  $\nabla t$  and  $\beta$  is any positive number less than 2. It is not clear to us how Section 6.2 in [28] is applied in the proof (4.1) in [15]. In fact the estimate (4.1) does not seem to be correct without the convexity of geodesic spheres (cf. [30, 21]).

It is observed in [15] that one may employ the Bishop-Gromov relative volume comparison theorem and get

$$\frac{\text{Vol}(\Gamma_t, g^+)}{\text{Vol}(\Gamma_t, g_{\mathbb{H}})} \leq \frac{\text{Vol}(B(t, p_0), g^+)}{\text{Vol}(B(t, 0), g_{\mathbb{H}})} \leq 1$$

for all  $t > 0$  on a conformally compact Einstein manifold. Hence the real issue for (1.9) is the lower bound and the key is to establish the relative volume lower bound by the limit

$$\lim_{t \rightarrow \infty} \frac{\text{Vol}(\Gamma_t, g^+)}{\text{Vol}(\Gamma_t, g_{\mathbb{H}})}.$$

It is very original in [15] to realize that one may use the Yamabe quotient to bound the relative volume  $\frac{\text{Vol}(\Gamma_t, g^+)}{\text{Vol}(\Gamma_t, g_{\mathbb{H}})}$  from below. Suppose that  $(X^n, g^+)$  is an AH manifold and that  $x$  is the geodesic defining function associated with a representative  $\hat{g}$  of the conformal infinity  $(\partial X^{n-1}, [\hat{g}])$ . Let

$$(4.2) \quad r = -\log \frac{x}{2}$$

and  $\Sigma_r = \{p \in X^n : r(p) = r\}$  be the level set of the geodesic defining function. We would like to mention that the inequality (2.2) in Lemma 2.1 of [15] is a well known fact about AH manifolds of  $C^3$  regularity, which is

$$(4.3) \quad Ddr(v, v) = 1 + O(e^{-2r}),$$

where  $v$  is any unit vector perpendicular to  $\nabla r$ , provided that (1.11) holds. Let  $\bar{g} = x^2 g^+$  be the conformal compactification from the defining function and let

$$\bar{g}_t = \bar{g}|_{\Gamma_t} \text{ and } \bar{g}_r = \bar{g}|_{\Sigma_r}.$$

Also let  $\tilde{g} = 4e^{-2t} g^+ = \psi^{\frac{4}{n-3}} \bar{g}$  be the conformal compactification from the distance function and let

$$\tilde{g}_t = \tilde{g}|_{\Gamma_t} \text{ and } \tilde{g}_r = \tilde{g}|_{\Sigma_r},$$

where  $\psi = e^{\frac{n-3}{2}(r-t)}$ . First of all it is easily seen that  $u = r - t$  is bounded on  $X^n$  by the triangle inequality for distance functions. It is then observed in Lemma 3.1 of [15] that  $|\nabla u|[\bar{g}]$  is bounded. For the convenience of readers we present a complete proof and an easy fix for a gap in [15].

**Lemma 4.1.** *Suppose that  $(X^n, g^+)$  is an AH manifold of  $C^3$  regularity and (1.11) holds. Then there is a constant  $C$  such that*

$$(4.4) \quad |du|[\tilde{g}] \leq C,$$

when  $r$  is large enough.

*Proof.* It suffices to show that

$$(4.5) \quad g^+(\nabla t, \nabla r) = 1 + O(e^{-2t}).$$

First, as noticed in the proof of Lemma 3.1 in [15], if let  $\phi = g^+(\nabla t, \nabla r)$ , then

$$(4.6) \quad \frac{d}{dt}\phi = \nabla^2 r(\nabla t, \nabla t) = (1 - \phi^2)\nabla^2 r(v, v),$$

where  $\nabla t = \sqrt{1 - \phi^2}v + \phi\nabla r$ . We want to point out that it is not enough to derive (4.5) just from (4.6) and (4.3). One needs the next lemma, which turns out to an easy fact but not a consequence of (4.5) as presented in the proof of Lemma 4.1 of [15].  $\square$

**Lemma 4.2.** *Suppose that  $(X^n, g^+)$  is AH of  $C^3$  regularity and (1.11) holds. Let  $x$  be the geodesic defining function associated with a representative  $\hat{g}$  of the conformal infinity  $(\partial X, [\hat{g}])$ . Let  $p_0 \in X^n$  is a fixed point and  $t$  be the distance to  $p_0$  in  $g^+$ . And let  $r$  be given in (4.2). Then*

$$g^+(\nabla r, \nabla t) > 0$$

at the point  $t$  is smooth and  $r$  is sufficiently large.

*Proof.* It is known (cf. (2.2) in Lemma 2.1 of [15]) that, there is  $r_0 > 0$  such that, for  $r > r_0$ ,

$$\nabla^2 r(v, v) > \frac{1}{2}$$

for all unit vector  $v$  that is perpendicular to  $\nabla r$ . Then we claim that

$$\phi = g^+(\nabla t, \nabla r) > 0$$

for any point  $p$  where  $t$  is smooth and  $r > r_0$ . To see this, one considers the minimal geodesic  $\gamma$  that connects  $p$  to  $p_0$  and realizes the distance. Then, in the light of (4.6), it is not hard to see that  $\phi > 0$  from the time  $t_0$  when the geodesic  $\gamma$  exits from  $\Sigma_{r_0}$ . Because  $\phi \leq 1$  for all  $t$  and  $\phi(t_0) \geq 0$ .  $\square$

By Lemma 4.2, for  $r$  large, a geodesic from  $p_0$  can touch  $\Sigma_r$  at most once till the time it hits the cut locus of  $p_0$ . Hence by (4.4) the metric  $\tilde{g} = \psi^{\frac{4}{n-3}}\bar{g}$  extends to  $\bar{X}^n$  with Lipschitz regularity up to the boundary. In fact, as a consequence of the estimate (4.3) and Lemma 4.1 as observed in Corollary 2.2 of [15], one has the following:

**Corollary 4.3.** *Suppose that  $(X^n, g^+)$  is AH of  $C^3$  regularity and (1.11) holds. Then the second fundamental form of the level set  $\Sigma_r$  in  $(X^n, \tilde{g})$  converges to zero as  $r \rightarrow \infty$ .*

Corollary 4.3 is used in the calculation of the scalar curvature  $R[\tilde{g}_r]$  for  $(\Sigma_r, \tilde{g}_r)$  (cf. (5.5) in [15]). The following is one of the key steps in [15]. We will present the idea from [15] and point out what are not clear and not correct. We will finish filling those gaps in the subsequent sections.



**Lemma 4.4.** *Suppose that  $(X^n, g^+)$  is an AH manifold of  $C^3$  regularity and (1.8) holds, with its conformal infinity  $(\partial X^{n-1}, [\hat{g}])$  of positive Yamabe type. Then*

$$\left(\frac{Y(\partial X, [\hat{g}])}{(n-2)(n-1)}\right)^{\frac{n-1}{2}} \leq \text{Vol}(\partial X, \tilde{g}_0) = \lim_{t \rightarrow \infty} \text{Vol}(\Gamma_t, \tilde{g}_t),$$

where  $\tilde{g}_0$  is the continuous extension of  $\tilde{g}$  to the boundary.

*Proof.* We are recapturing the proof given in [15] in the way that the use of the estimate (4.1) and the places where more vigorous arguments are required are explicitly identified. The full proof of this lemma will be completed in the subsequent sections.

The first step is to derive (3.8) in [15]. Using the Lipschitz extension of  $\psi$  based on Lemma 4.1, we have

$$(4.7) \quad \frac{\int_{\partial X} \left( \frac{4(n-2)}{n-3} |\nabla \psi|^2 + R\psi^2 \right) dv [\hat{g}]}{\left( \int_{\partial X} \psi^{\frac{2(n-1)}{n-3}} dv [\hat{g}] \right)^{\frac{n-3}{n-1}}} \leq \liminf_{r \rightarrow \infty} \frac{\int_{\Sigma_r} \left( \frac{4(n-2)}{n-3} |\nabla \psi|^2 + R\psi^2 \right) dv [\tilde{g}_r]}{\left( \int_{\Sigma_r} \psi^{\frac{2(n-1)}{n-3}} dv [\tilde{g}_r] \right)^{\frac{n-3}{n-1}}}.$$

On the other hand, we remark that, though

$$(4.8) \quad \int_{\Sigma_r} \left( \frac{4(n-2)}{n-3} |\nabla \psi|^2 + R\psi^2 \right) dv [\tilde{g}_r] = \int_{\Sigma_r} (Rdv) [\tilde{g}_r]$$

easily holds when  $\psi$  is smooth, (4.8) (cf. (3.7) in [15]) may not be correct when  $\psi$  is only known to be Lipschitz. Our approach is to use the deep understanding of the structure of cut loci to overcome the challenge based on [27, 22]. We will deal with this issue and complete this first step in Section 5 (cf. Theorem 5.4).

The second step is to obtain the pointwise scalar curvature estimate

$$(4.9) \quad R[\tilde{g}_r] \leq (n-1)(n-2) + o(1)$$

(cf. (5.1) in [15]). The proof of Lemma 5.1 in [15] uses Corollary 4.3 and the Laplacian comparison theorem. More importantly it uses the estimate (4.1) in calculating

$$\nabla^2 t(\nabla r, \nabla r) = 1 - (g^+(\nabla r, \nabla r))^2 + O(e^{-3t}),$$

which indeed would imply (4.9) and

$$(4.10) \quad \liminf_{r \rightarrow \infty} \frac{\int_{\Sigma_r} (Rdv) [\tilde{g}_r]}{\left( \int_{\Sigma_r} dv [\tilde{g}_r] \right)^{\frac{n-3}{n-1}}} \leq (n-1)(n-2) \lim_{r \rightarrow \infty} \text{Vol}(\Sigma_r, \tilde{g})^{\frac{2}{n-1}} = (n-1)(n-2) \text{Vol}(\partial X, \tilde{g}_0)^{\frac{2}{n-1}}.$$

We will present a proof of (4.10) without assuming the estimate (4.1) at each smooth point of  $t$  on  $\Sigma_r$ . Our key idea is to show the part of  $\Sigma_r$  where (4.1) does not hold has arbitrarily small volume. We will present a complete and correct proof the second step in Section 6 (cf. Theorem 6.8).

The last step is to show that

$$(4.11) \quad \lim_{t \rightarrow \infty} \text{Vol}(\Gamma_t, \tilde{g}_t) = \text{Vol}(\partial X, \tilde{g}_0).$$

To do so, similar to the discussions in Section 6 of [15] that is based on Lemma 4.1, one considers the geodesic sphere  $\Gamma_t$  as a Lipschitz graph over  $\partial X$  in  $(X^n, \tilde{g})$ , where

$$x(t) \rightarrow 0 \text{ as } t \rightarrow \infty$$

in  $W^{1,p}$  topology for any  $p \in [1, \infty)$ . It is then easily seen that (4.11) holds, as shown in [15].  $\square$

Consequently, from the idea in [15], for conformally compact Einstein manifolds, we have the lower bound of the relative volume growth.

*Proof of Theorem 1.5* One only needs to realize that, as calculated in [15], the following:

$$(4.12) \quad \frac{\text{Vol}(\Gamma_t, g^+)}{\text{Vol}(\Gamma_t, g_{\mathbb{H}})} = \frac{\text{Vol}(\Gamma_t, \tilde{g}_t) \left(\frac{e^t}{2}\right)^{n-1}}{\omega_{n-1} \sinh^{n-1} t} = \frac{\text{Vol}(\Gamma_t, \tilde{g}_t)}{\omega_{n-1}} + o(1)$$

as  $t \rightarrow \infty$ .  $\square$

Now we are ready to prove Theorem 1.8.

*Proof of Theorem 1.8.* First we want to show that, there are constant  $\delta_0 > 0$  and  $C$  such that

$$|W|[g^+] \leq C$$

for any conformally compact Einstein manifolds that satisfy the assumptions in Theorem 1.8 for  $0 < \delta \leq \delta_0$ . Assume otherwise there is a sequence of conformally compact Einstein manifolds  $(X_j^n, g_j^+)$  such that

$$|W|[g_j^+] \rightarrow \infty \text{ and } Y(\partial X_j, [\hat{g}_j]) \rightarrow Y(\mathbb{S}, [g_{\mathbb{S}}])$$

as  $j \rightarrow \infty$ . By Theorem 1.5, we know that

$$(4.13) \quad \left(\frac{Y(\partial X_j, [\hat{g}_j])}{Y(\mathbb{S}, [g_{\mathbb{S}}])}\right)^{\frac{n-1}{2}} \leq \frac{\text{Vol}(\Gamma_t, g_j^+)}{\text{Vol}(\Gamma_t, g_{\mathbb{H}})} \leq \frac{\text{Vol}(B(p_j, t), g_j^+)}{\text{Vol}(B(0, t), g_{\mathbb{H}})} \leq 1$$

for  $t > 0$ . Since  $|W|[g_j^+](p) \rightarrow 0$  as  $p \rightarrow \infty$  on each conformally compact Einstein manifold  $(X_j^n, g_j^+)$ , there exists a point  $p_j \in X_j^n$  so that

$$\tau_j = |W|[g_j^+](p_j) = \max_{p \in X_j^n} |W|[g_j^+](p) \rightarrow \infty$$

as  $j \rightarrow \infty$ . We then consider the rescaled metric  $g_j = \tau_j g_j^+$  on the pointed manifold  $(X_j^n, p_j)$ . From (4.13), one may conclude that the sequence of pointed Einstein manifolds  $(X_j^n, \tau_j g_j^+, p_j)$  converges to a Ricci flat manifold  $(X_\infty^n, g_\infty, p_\infty)$  in Cheeger-Gromov topology. In particular, one gets, again from (4.13),

$$\text{Vol}(\Gamma_t, g_\infty) = \text{Vol}(\Gamma_t, g_E)$$

for all  $t > 0$ , which implies that  $(X_\infty^n, g_\infty)$  is isometric to the Euclidean space  $(\mathbb{R}^n, g_E)$  and hence contradicts with  $|W|(p_\infty)[g_\infty] = 1$ .

To finish the proof of Theorem 1.8 we assume again otherwise, there are  $\epsilon_0 > 0$  and a sequence of conformally compact Einstein manifolds  $(X_j^n, g_j^+)$  such that

$$|W|(p_j)[g_j^+] \geq \epsilon_0 \text{ and } Y(\partial X, [\hat{g}_j]) \rightarrow Y(\mathbb{S}^n, [g_{\mathbb{S}}])$$

as  $j \rightarrow \infty$ . According to the above uniform bound for the curvature for such a sequence, we may extract a subsequence of pointed Einstein manifolds  $(X_j^n, g_j^+, p_j)$  with  $|W|(p_j)[g_j^+] \geq \epsilon_0 > 0$ , which converges to an Einstein manifold  $(X_\infty^n, g_\infty, p_\infty)$  in Cheeger-Gromov topology. Then the exact same argument at the end of Section 7 in [15] produces a contradiction from  $|W|(p_\infty)[g_\infty] \geq \epsilon_0 > 0$ . So the proof is complete.  $\square$

It is obvious that the argument in the first step in the above proof of Theorem 1.8 implies Corollary 1.9, in the same spirit as in the proof of Lemma 3.4.

## 5. NORMAL CUT LOCI

In this section we focus on the issue in the first step of the proof of Lemma 4.4. First  $\psi$  is smooth away from the cut loci of the point  $p$  in a conformally compact Einstein manifold  $(X^n, g^+)$ . Hence it is necessary to understand the fine structure of the set of cut loci and the behavior of the distance function near the cut loci in order to study (4.8).

One might think, for a fixed  $p$  in a complete and non-compact manifold  $(M^n, g)$ , the cut loci  $C_p$  may stay in a compact set. In fact, to the contrary, any component of  $C_p$  extends to the infinity, unless  $p$  is a pole. This is because  $M^n \setminus C_p$  is always diffeomorphic to the Euclidean space  $\mathbb{R}^n$ .

On a complete Riemannian manifold  $(M^n, g)$  with a fixed point  $p \in M^n$ , the set  $C_p$  of cut loci consists of the set  $Q_p$  of conjugate points and the set  $A_p$  of non-conjugate cut loci. Among the points in  $A_p$  we call those from which there are exactly two minimal geodesics connecting to  $p$  and realizing the distance to  $p$  in  $(M^n, g)$  the normal cut loci, according to [27, 22]. We denote the set of all normal cut loci by  $N_p$  and the rest of non-conjugate cut loci by  $L_p$ . In those notations we have

$$C_p = Q_p \cup L_p \cup N_p.$$

We recall from [27, 22] the following facts about the cut loci on Riemannian manifolds in general.

**Lemma 5.1.** ([27, 22]) *Suppose that  $(M^n, g)$  is a complete Riemannian manifold and that  $p \in M^n$ . Then*

- *The closed set  $Q_p \cup L_p$  is of Hausdorff dimension no more than  $n - 2$ .*
- *The set  $N_p$  of normal cut loci consists of possibly countably many disjoint smooth hypersurfaces in  $M^n$ .*
- *Moreover, at each normal cut locus  $q \in N_p$ , there is a small open neighborhood  $U$  of  $q$  such that  $U \cap C_p = U \cap N_p$  is a piece of smooth hypersurface in  $M^n$ .*

In our cases, on a conformally compact Einstein manifold  $(X^n, g^+)$  with a given point  $p$ , we are concerned with the set

$$\gamma_r = \Sigma_r \cap C_p = (\Sigma_r \cap (Q_p \cup L_p)) \cup (\Sigma_r \cap N_p) = \gamma_r^{QL} \cup \gamma_r^N,$$

where  $\psi$  is not smooth as a Lipschitz function on  $\Sigma_r$ . Before we move to look closely on (4.8) we would like to mention some more facts about the distance function  $t$  and the geodesic spheres in our context.

**Lemma 5.2.** *Suppose that  $(X^n, g^+)$  is AH of  $C^3$  regularity and (1.11) holds. Then*

- When  $t$  is sufficiently large, the geodesic sphere  $\Gamma_t$  is a Lipschitz graph over  $\partial X$ .
- When  $t$  is sufficiently large, the outward angle of the corner at the normal cut locus on  $\Gamma_t$  is always less than  $\pi$ .

*Proof.* The first statement can be proven using the same argument as in Section 4 of [15] (cf. Lemma 4.1 and Lemma 4.2 in the previous section).

From [27, 22] we know that the singularities for the geodesic sphere at normal cut loci are corners that are, at least locally, the meet of two smooth hypersurfaces. To see the outward angle of such corner at each normal cut locus is always less than  $\pi$ , let us assume otherwise. Let  $\gamma$  be a (distance realizing) minimal geodesic from the fixed  $p$  to a normal cut locus  $q \in \Gamma_t$  where the inward angle is less than  $\pi$ . We may push toward the geodesic sphere  $\Gamma_t$  from inside a small geodesic ball centered along  $\gamma$ . Clearly the geodesic ball will definitely touch the geodesic sphere  $\Gamma_t$  at some point  $\bar{q} \in \Gamma_t$  before it reaches the corner  $q \in \Gamma_t$ . This yields a contradiction because the distance from  $p$  to the point  $\bar{q} \in \Gamma_t$  would be definitely less than  $t$ .  $\square$

By compactness of  $\gamma_r^{QL}$ , there are finitely many points  $p_i \in \gamma_r^{QL}$  so that  $B_r^\epsilon = \bigcup_i B_\epsilon(p_i)$  covers  $\gamma_r^{QL}$ . Then  $B_r^\epsilon$  together with  $\gamma_r^N$  covers the set  $\gamma_r$ . Then we perform the integral by parts to calculate the Yamabe functional for  $\psi$  as follows:

$$(5.1) \quad \int_{\Sigma_r \setminus (B_r^\epsilon \cup \gamma_r^N)} \left( \frac{4(n-2)}{n-3} |\nabla \psi|^2 + R\psi^2 \right) dv [\bar{g}_r] = \int_{\Sigma_r \setminus (B_r^\epsilon \cup \gamma_r^N)} (Rdv) [\bar{g}_r] \\ + \oint_{\partial(\Sigma_r \setminus (B_r^\epsilon \cup \gamma_r^N))} \frac{4(n-2)}{n-3} \psi \left( \frac{\partial}{\partial n} \psi d\sigma \right) [\bar{g}_r]$$

In the light of Lemma 5.1, for almost every  $r$  as  $r \rightarrow \infty$  and almost every  $\epsilon$  as  $\epsilon \rightarrow 0$ , one may assume  $\gamma_r^{QL}$  is of Hausdorff dimension no more than  $n-3$  and  $\gamma_r^N \setminus B_r^\epsilon$  is a union of finitely many disjoint hypersurfaces in  $\Sigma_r$ . Hence

$$\oint_{\partial(\Sigma_r \setminus (B_r^\epsilon \cup \gamma_r^N))} \psi \left( \frac{\partial}{\partial n} \psi d\sigma \right) [\bar{g}_r] = \oint_{\partial B_r^\epsilon} \psi \left( \frac{\partial}{\partial n} \psi d\sigma \right) [\bar{g}_r] + \oint_{\gamma_r^N \setminus B_r^\epsilon} \psi \left( \left( \frac{\partial}{\partial n^+} \psi + \frac{\partial}{\partial n^-} \psi \right) d\sigma \right) [\bar{g}_r]$$

where  $n^+$  and  $n^-$  are the two outward normal directions to  $\gamma_r^N$  from the inside of  $\Sigma_r \setminus (B_r^\epsilon \cup \gamma_r^N)$ . It is not hard to see that

$$(5.2) \quad \oint_{\partial B_r^\epsilon} \psi \left( \frac{\partial}{\partial n} \psi d\sigma \right) [\bar{g}_r] \rightarrow 0$$

as  $\epsilon \rightarrow 0$ . Because,  $\gamma_r^{QL}$  is compact and of Hausdorff dimension no more than  $n-3$ ; and  $\psi$  is uniformly Lipschitz on  $\Sigma_r$  (cf. (4.4) in Lemma 4.1). Recall that  $\psi = e^{\frac{n-3}{2}(r-t)}$  at least for almost every  $r$ . Therefore

$$\frac{\partial}{\partial n^\pm} \psi = -\frac{n-3}{2} \psi \frac{\partial t}{\partial n^\pm} = -\frac{n-3}{2} \psi (\nabla t)^\pm \cdot n^\pm$$

and

$$(5.3) \quad \frac{\partial}{\partial n^+} \psi + \frac{\partial}{\partial n^-} \psi = -\frac{n-3}{2} \psi \left( \frac{\partial t}{\partial n^+} + \frac{\partial t}{\partial n^-} \right) = -\frac{n-3}{2} \psi ((\nabla t)^+ - (\nabla t)^-) \cdot n^+,$$


where  $(\nabla t)^\pm$  is the gradient of the distance function  $t$  with respect to the metric  $\bar{g}$  from either side of the corner  $\gamma_r^N$ .

**Lemma 5.3.** For almost every  $r$ , when  $\gamma_r^N$  is union of disjoint hypersurfaces in  $\Sigma_r$ ,

$$(5.4) \quad \frac{\partial t}{\partial n^+} + \frac{\partial t}{\partial n^-} \geq 0$$

at each point on  $\gamma_r^N$ .

*Proof.* Given a point  $q \in \gamma_r^N$ , let us consider the plane  $P$  spanned by  $(\nabla t)^+$  and  $(\nabla t)^-$ . And let  $n^t$  be the unit direction of the projection of  $n^+$  to the plane  $P$ . One notices that, from (4.5), the angle between  $(\nabla t)^+$  and  $(\nabla t)^-$  is arbitrarily small as well as the angle between  $(\nabla t)^\pm$  and  $n^t$  is arbitrarily close to  $\frac{\pi}{2}$ , as  $r \rightarrow \infty$ . In the light of (5.3), to verify (5.4) is to verify that the angle from  $(\nabla t)^-$  to  $n^t$  is not smaller than the one from  $(\nabla t)^+$  to  $n^t$ , since  $\|(\nabla t)^+\| = \|(\nabla t)^-\|$ .



This turns out to be true because the outward angle of the corner at any normal cut locus on the geodesic sphere is always less than  $\pi$  according to Lemma 5.2.  $\square$

From the proof of Lemma 5.3, we can observe that for  $r > 0$  large,  $N_p$  intersects with  $\Sigma_r$  transversely, so that for  $q \in N_p$ , there exists  $v \in T_q N_p$  so that the angle between  $v$  and  $\nabla_g r(q)$  is bounded by  $Ce^{-r}$  with a uniform constant  $C > 0$ . Therefore,  $\mathcal{H}^{n-1}(\Sigma_r \cap C_p) = 0$ . Similarly,  $\mathcal{H}^{n-1}(\Gamma_t \cap C_p) = 0$  for  $t > 0$  large. To summarize what we have so far in this section we state the following proposition:

**Theorem 5.4.** Suppose that  $(X^n, g^+)$  is an AH manifold of  $C^3$  regularity and (1.11) holds. Let  $p$  be a fixed point on  $X^n$ . Let  $t$  be the distance function to  $p$  on  $(X^n, g^+)$  and let  $r = -\log \frac{x}{2}$ , where  $x$  is the geodesic defining function associated with a representative  $\hat{g}$  of the conformal infinity  $(\partial X, [\hat{g}])$ . Then for almost all large  $r > 0$  so that  $\mathcal{H}^{n-2}(\gamma_r^{QL}) = 0$ , it holds that

$$(5.5) \quad \int_{\Sigma_r} \left( \frac{4(n-2)}{n-3} |\nabla \psi|^2 + R\psi^2 \right) dv [\tilde{g}_r] \leq \int_{\Sigma_r \setminus (B_r^\epsilon \cup \gamma_r^N)} (Rdv) [\tilde{g}_r] + o_\epsilon(1),$$

where  $\psi = e^{\frac{n-3}{2}(r-t)}$  and  $o_\epsilon(1) \rightarrow 0$  as  $\epsilon \rightarrow 0$ .

## 6. ESTIMATES OF THE TOTAL SCALAR CURVATURE

In this section we focus on the issue in the second step of the proof of Lemma 4.4. Let us first be very clear on how (4.1) is used in the argument in [15] and what one can hope to get for an upper bound for the scalar curvature  $R[\tilde{g}_r]$ . Recall from [15], to estimate the scalar curvature  $R[\tilde{g}_r]$ , one starts with (5.5) in [15],

$$R[\tilde{g}_r] = R[\tilde{g}] - 2\text{Ric}[\tilde{g}](N, N) + o(1)$$

and

$$\text{Ric}[\tilde{g}](N, N) = \frac{e^{2t}}{4}(\text{Ric}[g^+](\nabla r, \nabla r) + \Delta t + (n-2)(1 - (g^+(\nabla r, \nabla t))^2)(\nabla^2 t(v, v) - 1))$$

where  $N = \frac{1}{2}e^t \nabla r$  and

$$\nabla r = g^+(\nabla r, \nabla t)\nabla t + \sqrt{1 - (g^+(\nabla r, \nabla t))^2}v$$

for some unit vector  $v \perp \nabla t$ . Hence, following the calculation in [15] and assuming

$$(6.1) \quad \text{Ric}[g^+] \geq -(n-1)g^+ \text{ and } R[g^+] = -n(n-1) + o(e^{-2t}),$$

one arrives at

$$(6.2) \quad R[\tilde{g}_r] \leq (n-1)(n-2) + \frac{n-2}{2}e^{2t}(1 - (g^+(\nabla r, \nabla t))^2)(1 - \nabla^2 t(v, v)) + o(1),$$

which implies (4.9) whenever (4.1) is available. In fact it is clear that, any lower bound of the principal curvature would yield an upper bound for the scalar curvature  $R[\tilde{g}_r]$  from (6.2) and (4.5). To our best knowledge, one does not have any a priori lower bound of the principal curvature, though one indeed can manage to get the desired upper bound for the principal curvature, by the nature of the Riccati equations in general (cf. [30, 21]). What we observe is that, smaller the principal curvature is; smaller the mean curvature is; and therefore smaller the surface volume element is, at any smooth point of the distance function. Thus we will still be able to obtain the desired estimate for the total scalar curvature

$$(6.3) \quad \int_{\Sigma_r \setminus C_{p_0}} (Rdv)[\tilde{g}_r] \leq (n-1)(n-2) \int_{\Sigma_r} dv[\tilde{g}_r] + o(1)$$

where  $o(1) \rightarrow 0$  as  $r \rightarrow \infty$ . We will organize this section into three subsections.

**6.1. Curvature estimates based on the Riccati equations.** In this section we would like to derive the curvature estimates based on the Riccati equations on AH manifolds. Let us start with the Riccati equation for the shape operator of the geodesic spheres along a geodesic  $\gamma(t)$  (cf. for example, (2.1) in [15] or [30, 21, 28])

$$(6.4) \quad \nabla_{\nabla t} S + S^2 = -(R_{\nabla t})[g^+]$$

where  $S = \nabla^2 t$  is the shape operator and  $(R_{\nabla t})[g^+](v) = (R(v, \nabla t)\nabla t)[g^+]$ . Particularly, one has

$$(6.5) \quad 1 - C_0 e^{-2t} \leq \mu'(t) + \mu^2(t) \leq 1 + C_0 e^{-2t}$$

for the principal curvature  $\mu$ , on an AH manifold with (1.11) (cf. (2.6) in [15]). We will use  $\mu_m$  and  $\mu_M$  to stand for the smallest and the biggest principal curvature respectively. The step following (2.6) in the proof of (2.3) in [15] does not seem to be correct to us. In the rest of this subsection we will present a careful study of the Riccati equations and derive the upper bounds

and lower bounds of the principal curvature that hold on AH manifolds.

The first is a general fact for complete Riemannian manifolds.

**Lemma 6.1.** *Suppose that  $(M^n, g)$  is a complete Riemannian manifold and  $p \in M^n$  is a fixed point. For any given  $t_0 > 0$ , there is a constant  $C$  such that*

$$(6.6) \quad \mu_M(q) \leq C$$

for all  $q \in \Gamma_{t_0} \setminus C_p$ , where  $\Gamma_{t_0}$  is the geodesic sphere of radius  $t_0$  and  $C_p$  is the cut loci of  $p$  in  $(M^n, g)$ .

*Proof.* Suppose otherwise that there is a sequence  $q_k \in \Gamma_{t_0} \setminus C_p$  such that

$$\mu_M(q_k) \rightarrow \infty.$$

Without loss of generality one may assume that  $q_k \rightarrow q_0 \in \Gamma_{t_0}$  and the geodesic  $\gamma_k(t)$  that realizes the distance and connects  $q_k$  and  $p$  converges to a minimizing geodesic  $\gamma(t)$  from  $q_0$  to  $p$ . Hence  $q_0$  is the first cut locus of  $p$  along  $\gamma$ . Set

$$\mu = \mu_M(\gamma(\frac{1}{2}t_0)).$$

Since the set  $C_p$  of cut loci of  $p$  is closed, there is an open neighborhood  $U$  of  $\gamma(\frac{1}{2}t_0)$  where  $t$  is smooth. One may assume that

$$\mu_M(q) \leq 2\mu$$

for all  $q \in U$  (take a smaller set if necessary). Particularly

$$\gamma_k(\frac{1}{2}t_0) \in U \text{ and } \mu_M(\gamma_k(\frac{1}{2}t_0)) \leq 2\mu$$

for all sufficiently large  $k$ , which is easily seen to contradict with the fact that

$$\mu_M(q_k) = \mu_M(\gamma_k(t_0)) \rightarrow \infty$$

in the light of the Riccati equation (6.4). So the proof is complete.  $\square$

Consequently we have the sharp upper bound for the principal curvature of geodesic spheres in AH manifolds.

**Corollary 6.2.** *Suppose that  $(X^n, g^+)$  is AH and  $p \in X^n$  is a fixed point. And suppose that the curvature condition (1.11) holds on  $(X^n, g^+)$ . Then, for  $t_0 > 0$ , there exists a constant  $C$  such that*

$$(6.7) \quad \mu_M(q) \leq 1 + C(t+1)e^{-2t}$$

for all  $q \in \Gamma_t \setminus C_p$  and all  $t \geq t_0$ .

*Proof.* From (6.5) one has

$$\mu'_M + \mu_M^2 \leq 1 + C_0 e^{-2t}.$$

Then one may consider  $z = \mu_M - 1$  along the minimizing geodesic  $\gamma(s)$  from  $p$  to  $q \in \Gamma_t$  and the equation (6.5) for  $z$

$$z' + 2z \leq C_0 e^{-2s} - z^2 \leq C_0 e^{-2s},$$

which easily implies (6.7). We remark that the constant  $C$  may vary in different places.  $\square$

Roughly speaking, the reason one has the sharp upper bound (6.7) is because 1 is a sink for the equation

$$(6.8) \quad \mu' + \mu^2 = 1.$$

On the other hand,  $-1$  is a source for the equation (6.8), one could not expect a lower bound in general. In fact the best one can say about the lower bound for the principal curvature is as follows:

**Proposition 6.3.** *Suppose that  $(X^n, g^+)$  is AH and  $p \in X^n$  is a fixed point. And suppose that the curvature condition (1.11) holds on  $(X^n, g^+)$ . Then*

$$\mu_m(q) \geq -\sqrt{1 + C_0 e^{-2t}}$$

for  $q \in \Gamma_t$  and  $q$  is on a minimizing geodesic ray that runs from  $p$  to the infinity without intersecting the set  $C_p$  of cut loci of  $p$ .

*Proof.* Suppose that  $\gamma$  is a minimizing geodesic that runs from  $p$  to the infinity without intersecting the set  $C_p$  of cut loci of  $p$ . Assume otherwise

$$\mu_m(\gamma(t)) < -\sqrt{1 + C_0 e^{-2t}}.$$

Then, based on (6.5), it is not hard to show that  $\mu_m(\gamma(s))$  goes to  $-\infty$  in a finite time after  $t$ , which is a contradiction.  $\square$

**6.2. Volume estimates where the mean curvature is not big enough.** In this section we are concerned with the set

$$U_r^\delta = \{q \in \Sigma_r \setminus C_p : H(q) < (n-1)(1-\delta)\}$$

where  $H = \Delta t$ . We would like to show that

$$\int_{U_r^\delta} dv[\tilde{g}_r] \rightarrow 0$$

as  $r \rightarrow \infty$ .

We first notice from (4.5) in the proof of Lemma 4.1 that

$$(6.9) \quad \|dv[g^+](\nabla t) - dv[g^+](\nabla r)\|_{g^+} \leq C e^{-t}$$

and

$$(6.10) \quad \left| \frac{2^{n-1} e^{-(n-1)t} dv[g^+](\nabla r)}{2^{n-1} e^{-(n-1)t} dv[g^+](\nabla t)|_{(\nabla r)^\perp}} - 1 \right| = \left| \frac{dv[g^+](\nabla r)}{dv[g^+](\nabla t)|_{(\nabla r)^\perp}} - 1 \right| \leq C e^{-2t},$$

where

$$(6.11) \quad dv[\tilde{g}_r] = 2^{n-1} e^{-(n-1)t} dv[g^+](\nabla r)$$

and  $(\nabla r)^\perp$  stands for the hyperplane perpendicular to  $\nabla r$ . This gives us the clue how the mean curvature  $H = \Delta t$  is related to the volume element  $dv[\tilde{g}_r]$  on  $\Sigma_r$ . We now focus on the behavior of the  $(n-1)$ -form  $dv[g^+](\nabla t)$  along geodesics  $\gamma$  from  $p$ .



To be more clearer we set a parallel orthonormal frame  $\{\omega_1, \omega_2 \cdots, \omega_{n-1}\}$  along a geodesic  $\gamma$  for the dual of the subspace perpendicular to the geodesic  $\gamma$ . Then we may write

$$dv[g^+](\nabla t) = \mathcal{J}\omega_1 \cdot \omega_2 \cdots \omega_{n-1}.$$

Now we recall from the first variation formula in Riemannian geometry that

$$(6.12) \quad \frac{d\mathcal{J}}{dt} = H\mathcal{J}.$$

To state the key observation for the volume estimate, for each small  $\delta$ , we let  $t_\delta$  be a fixed large number such that

$$(n-2)C(t+1)e^{-2t} \leq \frac{\delta}{4} \text{ and } 1 - C_0e^{-2t} \geq (1 - \frac{1}{2}\delta)^2$$

for all  $t \geq t_\delta$ , where  $C$  is the constant in (6.7) and  $C_0$  is the constant in (1.11).

**Lemma 6.4.** *Suppose that  $(X^n, g^+)$  ( $n \geq 4$ ) is AH of  $C^3$  regularity and  $p \in X^n$  is a fixed point. And suppose that the curvature condition (1.11) holds. Let  $q \in U_r^\delta$  and  $\gamma(s)$  be the minimizing geodesic from  $p$  to  $q = \gamma(t)$  for  $t > t_\delta$ . Then*

$$(6.13) \quad H(\gamma(s)) \leq (n-1)\left(1 - \frac{\delta}{4(n-1)}\right)$$

for all  $s \in (t_\delta, t)$ .

*Proof.* Assume otherwise, for some  $s \in (t_\delta, t)$  such that

$$H(\gamma(s)) > (n-1)\left(1 - \frac{\delta}{4(n-1)}\right).$$

Using the sharp upper bound (6.7) we conclude that

$$(6.14) \quad \mu_m(\gamma(s)) > 1 - \frac{\delta}{2}.$$

It is then easily seen that (6.14) implies

$$\mu_m(q) = \mu_m(\gamma(t)) \geq 1 - \frac{3}{4}\delta > 1 - \delta.$$

Because, from (6.5), one has

$$\mu'_m + \mu_m^2 \geq 1 - C_0e^{-2t} \geq (1 - \frac{1}{2}\delta)^2$$

in  $(t_\delta, t)$ , which means  $\mu'_m$  is positive whenever  $\mu_m \in (0, 1 - \frac{1}{2}\delta)$ . □

Consequently, from (6.11), (6.10), and (6.12), considering  $U_r^\delta$  as a graph on a subset of  $\Gamma_{t_\delta}$  induced by exponential map at  $p$ , we have

**Proposition 6.5.** *Suppose that  $(X^n, g^+)$  is an AH manifold of  $C^3$  regularity and  $p \in X^n$  is a fixed point. And suppose that the curvature condition (1.11) holds. Then*

$$(6.15) \quad \mathcal{J}(q) \leq \mathcal{J}(\gamma(t_\delta))e^{(n-1)(1-\frac{\delta}{4(n-1)})(t-t_\delta)}$$

for  $q \in \Gamma_t \setminus C_p$  with  $H < (n-1)(1-\delta)$  and  $\gamma$  is the minimizing geodesic from  $p$  to  $q$ . Thus, for each fixed small  $\delta > 0$ ,

$$(6.16) \quad \int_{U_r^\delta} dv[\tilde{g}_r] \rightarrow 0$$

as  $r \rightarrow \infty$ .

**6.3. Volume estimates where the mean curvature is very negative.** If there were a lower bound for the mean curvature, then (6.2) and (6.7) would yield a upper bound the scalar curvature  $R[\tilde{g}_r]$  and one would have by now completed the proof of (6.3). Therefore we need to estimate the lower bound of  $Hdv[g^+](\nabla t)$  at least when the mean curvature  $H$  is very negative. First of all one easily derives from the first variational formula (6.12) that, along a geodesic  $\gamma(s)$ ,

$$(6.17) \quad \frac{d}{ds}(Hdv[g^+](\nabla t)) = (H' + H^2)dv[g^+](\nabla t).$$

The key observation for this subsection is the following more detailed statement of Proposition 6.3 about the Riccati equation:

**Lemma 6.6.** *Suppose that  $(X^n, g^+)$  is AH and  $p \in X^n$  is a fixed point. And suppose that the curvature condition (1.11) holds. Then, for any  $t_0 > 0$ , there is a constant  $C > 0$ , for  $q \in \Gamma_t \setminus C_p$  and  $\gamma$  being the minimizing geodesic from  $p$  to  $q$ , one always has*

$$(6.18) \quad \mu_m(\gamma(t-1)) \geq -C,$$

provided that  $t > t_0 + 1$ .

*Proof.* To start, one considers the Riccati equation (6.5) along the geodesic  $\gamma$  when  $\mu(s_0) < -\sqrt{1 + C_0 e^{-2s_0}}$  with  $s_0 = t - 1 > t_0$ . Recall

$$\mu' + \mu^2 \leq 1 + C_0 e^{-2s} \leq 1 + C_0 e^{-2s_0}$$

for  $s > s_0$ , which implies

$$\mu(s) \leq -a \frac{1 + \chi e^{2a(s-s_0)}}{1 - \chi e^{2a(s-s_0)}}$$

where  $a = \sqrt{1 + C_0 e^{-2s_0}}$  and  $\chi = -\frac{a+\mu(s_0)}{a-\mu(s_0)} \in (0, 1)$ . Therefore it is clear that  $\mu$  reaches  $-\infty$  at

$$s_1 = t - 1 - \frac{1}{2a} \log \chi$$

and

$$\lim_{\mu(t-1) \rightarrow -a} \chi = 0 \text{ and } \lim_{\mu(t-1) \rightarrow -\infty} \chi = 1.$$

Thus

$$-\frac{1}{2a} \log \chi < 1$$

when  $\mu(t-1)$  is sufficient negative, which implies, there is a constant  $C = C(t_0)$  such that

$$\mu_m(\gamma(t-1)) > -C$$

for  $\gamma(t) \in \Gamma_t \setminus C_p$  and  $t > t_0 + 1$ . □

Now we are ready to deal with the volume estimate at places the mean curvature is very negative.

**Proposition 6.7.** *Suppose that  $(X^n, g^+)$  is AH and  $p \in X^n$  is a fixed point. And suppose that the curvature condition (1.11) holds. Then there is a constant  $C > 0$  such that, for  $q \in \Gamma_t \setminus C_p$  with  $H(q) \leq -2(n-1)$  and  $\gamma$  being the minimizing geodesic from  $p$  to  $q$ ,*

$$(6.19) \quad H\mathcal{J}(\gamma(t)) \geq -Ce^{(n-1)(1-\frac{\delta}{4(n-1)})t},$$

provided that  $t > t_\delta + 1$ .

*Proof.* Using Riccati equation and (6.17), we end at

$$\frac{d}{ds}(H\mathcal{J}) = (-\text{Ric}[g^+](\nabla t, \nabla t) - |\nabla^2 t|^2 + H^2)\mathcal{J}$$

for  $s \in (t-1, t) \subset (t_\delta, t)$ . Hence we are looking for lower bound for  $(\Delta t)^2 - |\nabla^2 t|^2$ . Let

$$\{\mu_i : i = 1, \dots, n-1\}$$

be the eigenvalues of  $\nabla^2 t$ . We may calculate

$$(\Delta t)^2 - |\nabla^2 t|^2 = \sum_{i \neq j} \mu_i \mu_j \geq \sum_{\mu_i \mu_j < 0} \mu_i \mu_j \geq 2(n-1)^2 \mu_m \geq 2(n-1)^2 H - C$$

for some constant  $C > 0$ , where the last inequality uses the sharp upper bound (6.7). Notice also that  $H \leq -2(n-1)$  yields that  $\mu_m \leq -2$ , which guarantees that there is always some negative eigenvalues. Therefore by (1.11), we obtain

$$(6.20) \quad \frac{d}{ds}(H\mathcal{J}) \geq 2(n-1)^2 H\mathcal{J} - C\mathcal{J}$$

for some constant  $C > 0$ , which can be rewritten as follows:

$$\frac{d}{ds}(e^{-2(n-1)^2 s} H\mathcal{J}) \geq -Ce^{-2(n-1)^2 s} \mathcal{J}$$

for  $s \in (t-1, t)$  Thus

$$(6.21) \quad H\mathcal{J}(\gamma(t)) \geq e^{2(n-1)^2} H\mathcal{J}(\gamma(t-1)) - Ce^{2(n-1)^2 t} \int_{t-1}^t e^{-2(n-1)^2 s} \mathcal{J}(\gamma(s)) ds.$$

Now, in the light of Lemma 6.6, (6.19) is proven if (6.15) in Proposition 6.5 is applicable to all points  $\gamma(s) : s \in [t-1, t]$ . To see (6.15) holds at each point  $\gamma(s) : s \in [t-1, t]$ , one uses that  $\mu_m(\gamma(t)) < -2$  and the Riccati equation (6.5) for  $t-1 > t_\delta$ . Because one gets that  $\mu_m(\gamma(s)) < -1 + C_0 e^{-2t_\delta}$  and hence  $H(\gamma(s)) \leq (n-1)(1-\delta)$  for all  $s \in [t-1, t]$ , at least when  $\delta$  is small enough.  $\square$

Combining Proposition 6.5 and Proposition 6.7 we thus have obtained

**Theorem 6.8.** *Suppose that  $(X^n, g^+)$  is AH of  $C^3$  regularity and  $p \in X^n$  is a fixed point. And suppose that (6.1) holds. Then for almost all large  $r > 0$  so that  $\mathcal{H}^{n-2}(\gamma_r^{QL}) = 0$ , it holds that*

$$(6.22) \quad \int_{\Sigma_r \setminus (B_\epsilon^r \cup N_p)} (Rdv)[\tilde{g}_r] \leq (n-1)(n-2) \int_{\Sigma_r} dv[\tilde{g}_r] + o_r(1),$$

where  $o_r(1)$  is independent of  $\epsilon > 0$  and  $o_r(1) \rightarrow 0$  as  $r \rightarrow \infty$ .

*Proof.* At any point  $q \in \Sigma_r \setminus C_p$ , by (6.10),

$$\left| \frac{Hdv[g^+](\nabla r)}{Hdv[g^+](\nabla t)|_{(\nabla r)^\perp}} - 1 \right| \leq Ce^{-2t}.$$

Since  $|t - r|$  is uniformly controlled, for  $q \in \Sigma_r \setminus C_p$  so that  $H \leq -2(n - 1)$  we get the decay of  $Hdv[g^+](\nabla r)$  by Proposition 6.7.

For any  $\delta > 0$ , for  $r > 0$  large as in the statement of the theorem, considering  $\Sigma_r$  as a graph of  $\Gamma_{t_\delta}$  induced by the exponential map at  $p$ , we have the splitting of the integration,

$$\begin{aligned} & \int_{\Sigma_r \setminus (B_\epsilon^r \cup N_p)} (RdV)[\tilde{g}_r] \\ &= \int_{\{q \in \Sigma_r \setminus (B_\epsilon^r \cup N_p) : H(q) > (n-1)(1-\delta)\}} (RdV)[\tilde{g}_r] + \int_{\{q \in \Sigma_r \setminus (B_\epsilon^r \cup N_p) : -2(n-1) < H \leq (n-1)(1-\delta)\}} (RdV)[\tilde{g}_r] \\ &+ \int_{\{q \in \Sigma_r \setminus (B_\epsilon^r \cup N_p) : H \leq -2(n-1)\}} (RdV)[\tilde{g}_r]. \end{aligned}$$

Note that  $Ddt(v, v) \geq \mu_m \geq H - C$ . By Proposition 6.5 and Proposition 6.7 combining with (6.11) and (6.12) and arbitrary choice of  $\delta > 0$ , we completes the proof of (6.23).  $\square$

#### REFERENCES

- [1] M. Anderson, *Ricci curvature bounds and Einstein metrics on compact manifolds*, J. Amer. Math. Soc. **2** (1989), no. 3, 455-490.
- [2] M. Anderson,  *$L^2$  curvature and volume renormalization of the AHE metrics on 4-manifolds*, Math. Res. Lett. **8** (2001), no. 3, 171-188.
- [3] L. Andersson and M. Dahl, *Scalar curvature rigidity for asymptotically hyperbolic manifolds*, Ann. Global Anal. and Geo. **16** (1998) 1-27.
- [4] T. Aubin, *Nonlinear Analysis on Manifolds, Monge-Ampère Equations*, Grundlehren der mathematischen Wissenschaften **252**, 1982 Springer-Verlag New York.
- [5] S. Bando, A. Kasue, H. Nakajima, *On a construction of coordinates at infinity on manifolds with fast curvature decay and maximal volume growth*, Invent. math. **97** (1989), 313-349.
- [6] O. Biquard, *Asymptotically symmetric Einstein metrics*, Translated from the 2000 French original by Stephen S. Wilson. SMF/AMS Texts and Monographs, 13. American Mathematical Society, Providence, RI; Societe Mathematique de France, Paris, (2006), vi+105 pp.
- [7] V. Bonini, P. Miao, J. Qing, *Ricci curvature rigidity for weakly asymptotically hyperbolic manifolds*, Comm. Anal. Geom. **14** (3) (2006) 603 - 612.
- [8] A. Chang, M. Gursky, P. Yang, *A conformally invariant sphere theorem in four dimensions*, Publ. Math. Inst. Hautes tudes Sci. **98** (2003), 105-143.
- [9] A. Chang, J. Qing, P. Yang, *On A Conformal Gap and Finiteness Theorem for A Class of Four-manifolds*, GAFA, **17** (2007), 404-434.
- [10] A. Chang, J. Qing, P. Yang, *Renormalized volumes for conformally compact Einstein manifolds*. (Russian) Sovrem. Mat. Fundam. Napravl. **17** (2006), 129 - 142; translation in J. Math. Sci. (N. Y.) **149** (2008), no. 6, 1755 - 1769.
- [11] A. Chang, J. Qing, P. Yang, *On the topology of conformally compact Einstein 4-manifolds*, Noncompact problems at the intersection of geometry, analysis, and topology, 49C61, Contemp. Math., 350, Amer. Math. Soc., Providence, RI, 2004.
- [12] J. Cheeger, A. Naber, *Regularity of Einstein Manifolds and The Codimension 4 Conjecture*, arXiv:1406.6534v1, preprint.

- [13] P.T. Chruściel, E. Delay, J.M. Lee, D.N. Skinner, *Boundary Regularity of Conformally Compact Einstein Metrics*, J. Differential Geom. **69** (2005), no. 1, 111-136.
- [14] Piotr Chruściel and M. Herzlich, *The mass of asymptotically hyperbolic Riemannian manifolds*, Pacific J. Math. **212** (2003), no. 2, 231- 264..
- [15] S. Dutta and M. Javaheri, *Rigidity of Conformally Compact Manifolds with The Round Sphere as Conformal Infinity*, Adv. Math., **224**, (2010), 525-538.
- [16] C. Fefferman and C.R. Graham, *Conformal invariants*, in Elie Cartan et les Mathematiques dAujourd'hui, Asterisque (1985), 95-116.
- [17] R. Graham, *Volume and Area Renormalizations for Conformally Compact Einstein Metrics*, Rend. Circ. Mat. Palermo. Ser. II Suppl. **63** (2000) 31 - 42, arXiv:math/9909042v1.
- [18] R. Graham, J. Lee, *Einstein metrics with prescribed conformal infinity on the ball*, Adv. Math. **87** (1991), no. 2, 186-225.
- [19] E. Hebey, *Sobolev spaces on Riemannian manifolds*, Lect. Notes Math., Vol. **1635**. Berlin: Springer 1996.
- [20] M. Henningson and K. Skenderis, *The holographic Weyl anomaly*, J. High Ener. Phys. **07** (1998), 1 - 23, hep-th/9806087; Holography and the Weyl anomaly, hep-th/9812032.
- [21] X. Hu, J. Qing, Y. Shi, *Regularity and rigidity of asymptotically hyperbolic manifolds*, Adv. Math., **230**, (2012), 2332-2363.
- [22] J. Itoh, M. Tanaka, *The dimension of a cut locus on a smooth Riemannian manifold*, Tohoku Math. J. (2), **50**, (1998), 571-575.
- [23] J. M. Lee, T. H. Parker, *The Yamabe Problem*, Bulletin of Amer. Math. Soc. **17** (1987), no. 1.
- [24] M.C. Leung, *Pinching theorem on asymptotically hyperbolic spaces*, Int. J. Math. 4 (**5**) (1993) 841- 857.
- [25] R. Mazzeo, *Elliptic Theory of Differential Edge Operators I*, Comm. Partial Differential Equations **16**(10) (1991), 1615 - 1664.
- [26] M. Min-Oo, *Scalar curvature rigidity of asymptotically hyperbolic spin manifolds*, Math. Ann. 285 (1989) 527 - 539.
- [27] V. Ozols, *Cut Loci in Riemannian Manifolds*, Tôhoku Math. J. **26** (1974), 219 - 227.
- [28] P. Petersen, *Riemannian Geometry*, Graduate Texts in Mathematics **171**, 1998 Springer-Verlag New York.
- [29] J. Qing, *On the rigidity for conformally compact Einstein manifolds*, Int. Math. Res. Not. (2003), no. 21, 1141-1153.
- [30] Y. Shi, G. Tian, *Rigidity of Asymptotically Hyperbolic Manifolds*, Commun.Math.Phys, **259**, (2005) 545-559.
- [31] G. Tian, J. Viaclovsky, *Bach-flat asymptotically locally Euclidean metrics*, Invent. Math. **160**(10) (2005), no. 2, 357 - 415.
- [32] X. D. Wang, *The mass of asymptotically hyperbolic manifolds*, J. Diff. Geom. 57 (**2**) (2001), 273 - 299.

GANG LI, BEIJING INTERNATIONAL CENTER FOR MATHEMATICAL RESEARCH, PEKING UNIVERSITY, BEIJING, CHINA  
*E-mail address:* runxing3@gmail.com

JIE QING, DEPARTMENT OF MATHEMATICS, UC SANTA CRUZ, CALIFORNIA, USA  
*E-mail address:* qing@ucsc.edu

YUGUANG SHI, KEY LABORATORY OF PURE AND APPLIED MATHEMATICS, SCHOOL OF MATHEMATICS SCIENCE, PEKING UNIVERSITY, BEIJING, 100871, P.R. CHINA.  
*E-mail address:* ygshi@math.pku.edu.cn