

# GC-MS analysis of bioactive compounds from the whole plant ethanolic extract of *Evolvulus alsinoides* (L.) L

Duraisamy Gomathi · Manokaran Kalaiselvi ·  
Ganesan Ravikumar · Kanakasabapathi Devaki ·  
Chandrasekar Uma

Revised: 27 May 2013 / Accepted: 8 July 2013 / Published online: 24 July 2013  
© Association of Food Scientists & Technologists (India) 2013

**Abstract** Medicinal plants are at great interest to the researcher in the field of biotechnology, as most of the drug industries depend in medicinal plants for the production of pharmaceutical compounds. Plants are the traditional sources for many chemicals used as pharmaceutical biochemicals, fragrances, food colours and flavours in different countries especially in India. Most herbal medicines and their derivative products were often prepared from crude plant extracts, which comprise a complex mixture of different phytochemical constituents (plant secondary metabolites). The chemical features of these constituents differ considerably among different species. GC-MS method used for the analysis of the obtained extracts can be an interesting tool for testing the amount of some active principles in herbs used in cosmetic, drugs, pharmaceutical or food industry. The aim of this study was to carry out for identification of bioactive compounds from the whole plant ethanolic extract of *Evolvulus alsinoides* by Gas chromatography and Mass spectroscopy (GC-MS). GCMS analysis of ethanolic extract was done by standard protocol using the equipment Thermo GC-Trace Ultra Version: 5.0, Thermo MS DSQ II. The GC-MS analysis revealed the presence of various compounds like piperine, octodecanoic acids, hexadecanoic acid and squalene in the ethanolic extract of *Evolvulus alsinoides*. Hence, the *Evolvulus alsinoides* may have chemopreventive, anticancer, anti-microbial activity, antioxidant and antidiabetic activity due to the presence of secondary metabolites in the ethanolic extract. Due to the presence of esters which can be used as a flavoring agent in food industries. These findings support the traditional use of *Evolvulus alsinoides* in various

disorders. Further studies are needed to isolate active principle of the extract as well as to elucidate their exact mechanism of action in various disorders.

**Keywords** *Evolvulus alsinoides* · Secondary metabolites · GC-MS analysis · Whole plant

## Introduction

Free radicals play a crucial role in the development of tissue damage in pathological events. Antioxidants are chemical compounds which have the ability to quench the free radicals and thereby it prevents the human body against various diseases. Plants are the rich sources of antioxidants which contain secondary metabolites such as phenolic and flavonoid compounds commonly which act as antioxidants with redox and metal chelating properties (Karimi and Jaafar 2011). Antioxidants are chemical compounds extremely useful to humans, which has the ability to reduce free radicals and/or to decrease their rate of production and lipid peroxidation in human bodies that cause various human diseases and aging (Galati and Brien 2004). The active principles of many drugs found in plants are secondary metabolites (Cragg and Newman 2005). Plants provide us with rich sources of natural antioxidants (Biswas et al. 2005) and so the phytochemical investigation on the extract for their main phytochemicals is very vital. Extraction is the main step for the recovery and isolation of bioactive phytochemicals from plant materials, before component analysis (Karimi and Jaafar 2011). Hence the medicinal plants are important sources of medicines and presently about 25 % of pharmaceutical prescriptions (Uma et al. 2009). According to the world health organization, the traditional medicine system will continue to play an essential role in health care system since over 80 % of the population in the third world country relies on the use of traditional medicine. Medicinal plants are

D. Gomathi · M. Kalaiselvi · G. Ravikumar · K. Devaki  
Department of Biochemistry, Karpagam University,  
Coimbatore 641 021, India

C. Uma (✉)  
Hawasaa University, Hawasaa, Ethiopia  
e-mail: umachandrasekaran29@gmail.com




and Xu 1988; Shi et al. 2005). Most of the volatile components are long chain unsaturated fatty acids which play very important roles in biological systems because they are structural elements of many valuable compounds as well as being important sources of energy (Mu et al. 2001). In recent years, increasing research has been carried out on fatty acids and the results obtained show that they possess significant sedative and hypnotic effects (Zhang et al. 1995). Hence, the present study was aimed to evaluate the chemical constituents in the whole plant ethanolic extract of *Evolvulus alsinoides* (L.) L.

## Materials and methods

### Collection and identification of plant material

*Evolvulus alsinoides* (L.) L. used for the investigation was obtained from Coimbatore District, Tamilnadu, India. The plant was authenticated by Dr. P. Satyanarayana, Botanical Survey of India, Tamil Nadu Agricultural University (TNAU) Campus, Coimbatore. The voucher number is BSI/SRC/5/23/2011-12/Tech.-514. Fresh whole plant material of *Evolvulus*



**Fig. 2** Mass spectra of identified compound from ethanolic extract of *Evolvulus alsinoides*




Fig. 2 (continued)

*alsinoides* was washed under running tap water, air dried and powdered in electric blender.

Sample extraction

A 100 g sample of dried plant powder was extracted in 500 ml of ethanol in an orbital shaker for 72 h. Repeated extraction was done with the same solvent until a clear colorless solvent was obtained. Obtained extract was evaporated to dryness and stored at 4 °C in an airtight container for further use.

Gas chromatography-mass spectroscopic analysis of *Evolvulus alsinoides*

GC-MS analysis of the whole plant extract of *Evolvulus alsinoides* was performed using the equipment Thermo GC-Trace Ultra Version: 5.0, Thermo MS DSQ II. The equipment has a DB 35 – MS Capillary Standard non-polar column with dimensions of 30 mm×0.25 mm ID×0.25 μm film. The carrier gas used is Helium with at low of 1.0 ml/min. The injector was operated at 250 °C and the oven temperature was

programmed as follows: 60 °C for 15 min, then gradually increased to 280 °C at 3 min. The identification of components was based on Willey and NIST libraries as well as comparison of their retention indices. The constituents were identified after comparison with those available in the computer library (NIST and Willey) attached to the GC-MS instrument and the results obtained have been tabulated.

## Result and discussion

Medicinal plants are the resources of new drugs. Many of the modern medicines are produced indirectly from the medicinal plants. They have contributed many ingredients to fight against various diseases and illness. The analysis and extraction of plant material play an important role in the development, modernization and quality control of herbal formulations. Studying of medicinal plants also facilitates to comprehend plant toxicity and also helps to protect human and animals from natural poisons. Hence the present study was undertaken to find out the bioactive compounds present in the ethanolic extract of *Evolvulus alsinoides* by using Gas chromatography and Mass spectroscopy. The active principles with their retention time (RT), molecular formula, molecular weight (MW), concentration (peak area %) are presented in Table 1 and Fig. 1 which shows the presence of 16 bioactive phytochemical compounds in the ethanolic extract of *Evolvulus alsinoides*. The mass spectra of identified compounds from *Evolvulus alsinoides* were presented in Fig. 2.

Among the identified phytochemicals, tetradecanoic acid, squalene, piperine and n Hexadecanoic acid have the property of antioxidant, 5- $\alpha$ -reductase inhibitor, antifibrinolytic, hemolytic and antimicrobial activities (Bodoprost and Rosemeyer 2007; Abirami and Rajendran 2011; Kala et al. 2011). Octadecadienoic acid (Z, Z) have the property of anti-inflammatory, hypocholesterolemic and antiarthritic activity which was reported by the earlier workers (Rani et al. 2009; Ponnamma and Manjunath 2012; Uma et al. 2009). Naphthalene also having good antimicrobial activity (Uma et al. 2009)

The squalene (triterpene) is a phenolic compound and that the terpenes are found in latex and resins of some plants and physiological function of these compounds are generally believed to be a chemical in defense against certain pathogens causing human and animal diseases (Scortichini and Rossi 1991). Their activity is a function of the lipophilic properties of the constituent terpenes, the properties of their functional groups and their aqueous solubility (Mahato and Sen 1997; Ezhilan and Neelamegam 2012). It possesses anti microbial activity, chemopreventive activity against colon carcinogenesis (Rao et al. 1998). In addition to antimicrobial activity, the squalene was also reported to have anticancer, antioxidant, chemopreventive, gastroprotective and hepatoprotective

effects, pesticide, anti-tumor and sunscreen properties (Sunitha et al. 2001; Ukiva et al. 2002; Katerere et al. 2003).

Piperine is used all over the world for various illnesses. Some health benefits that can be gained from using piperine may be helpful in reducing inflammation, improving digestion, the reduction of pain and the relief of Asthma. It has also been extensively evaluated antidepressant (Li et al. 2007), anticonvulsant (Wang et al. 1999), antioxidant (Rauscher et al. 2000; Prakash and Srinivasan 2010), antimutagenic (El Hamass et al. 2003), hepatoprotective (Matsuda et al. 2008) endocrine and several other activities (Vijayakumar and Nalini 2006; Atal et al. 2012).

Esters are important organic compounds with increasing number of commercial applications (Foresti et al. 2005). These compounds are largely used in fragrances, cosmetics detergents, flavors and pharmaceuticals. Esters (ethyl oleates) may also be used as plasticizers and lubricants; biological additives and hydraulic fluids (Hazarika et al. 2002). The use of ethyl oleate in commercial applications has been hampered due to the low amounts that can be recovered from natural sources (Radzi et al. 2006; Martiinez-Ruiz et al. 2008). Due to the presence of above mentioned compounds in the whole plant ethanolic extract of *Evolvulus alsinoides* may be used in various pharmaceutical and industrial applications.

## Conclusion

Medicinal plants, which form the backbone of traditional medicine, in the last few decades have been the subject for very intense pharmacological studies, this has been brought about by the acknowledgement of the value of medicinal plants as potential sources of new compounds of therapeutic value and as sources of lead compounds in drug development. Thus, the identification of bioactive compound in *Evolvulus alsinoides* was done by GC-MS analysis which shows the presence of 16 compounds. Among the identified compounds, octadecanoic acid, n-hexadecanoic acid, squalene and piperine have the role in antioxidant, antimicrobial, anticancer, antidiabetic and antiinflammatory effects. From this study it can be concluded that the *Evolvulus alsinoides* may serve as a new potential source of medicines due to the presence of these phytochemicals and bioactive compounds.

**Acknowledgment** We are thankful to our Chancellor, Advisor, Vice Chancellor and Registrar of Karpagam University for providing facilities and encouragement.

## References

- Abirami P, Rajendran A (2011) GC-MS determination of bioactive compounds of *Indigofera aspalathoides*. J Nat Prod Plant Resour 1(4):126–130

- Atal S, Agrawal PR, Vyas S, Phadnis P, Rai N (2012) Evaluation of the effect of piperine per se on blood glucose level in alloxan-induced diabetic mice. *Acta Pol Pharm-Drug Res* 69(5):965–969
- Auddy B, Ferreira M, Blasina F, Lafon L, Arredondo F, Dajas F, Tripathi PC et al (2003) Screening of antioxidant activity of some three Indian medicinal plants traditionally used for the management of neurodegenerative diseases. *J Ethnopharmacol* 84(2–3):131–138
- Austin DF (2008) *Evolvulus alsinoides* (Convolvulaceae): an American herb in the old World. *J Ethnopharmacol* 117:185–198
- Biswas S, Bhattacharyya J, Dutta AG (2005) Oxidant induced injury of erythrocyte-role of green tea leaf and ascorbic acid. *Mol Cell Biochem* 276:205–210
- Cragg GM, Newman DJ (2005) A continuing source of novel drug leads. *Pure Appl Chem* 77:7–24
- Daniel FA (2008) Review of *Evolvulus alsinoides* (Convolvulaceae): an American herb in the old world. *J Ethnopharmacol* 117:185–198
- Deng SX, Wang DC, Wang MD et al (1964) The cardiac effect of *Periploca sepium*. *Acta Pharm Sin* 11:75
- El Hamass R, Idaomar M, Alonso-Moraga M, Serrano AM (2003) Antimutagenic properties of bell and black peppers. *Food Chem Toxicol* 41(1):41–47
- Ezhilan BP, Neelamegam R (2012) GC-MS analysis of phytochemicals in the ethanol extract of *Polygonum chinense* L. *Pharmacogn Res* 4(1):11–14
- Foresti ML, Errazu A, Ferreira M (2005) Effect of several reaction parameters in the solvent-free ethyl oleate synthesis using *Candida rugosa* lipase immobilised on polypropylene. *Biochem Eng J* 25(1):69–77
- Galati G, Brien PJ (2004) Potential toxicity of flavonoids and other dietary phenolics: significance for their chemopreventive and anticancer properties. *Free Radic Biol Med* 37:287–303
- Hazarika S, Goswami N, Dutta N, Hazarika A (2002) Ethyl oleate synthesis by porcine pancreatic lipase in organic solvent. *Chem Eng J* 85(1):61–68
- Hedeji I, Xu JP (1988) Pregnane glycoside from an antitumour fraction of *Periploca sepium*. *Phytochemistry* 27:1173
- Kala SMJ, Balasubramanian T, Soris PT, Mohan VR (2011) GC-MS determination of bioactive components of *Eugenia singampattiana* Bedd. *Int J ChemTech Res* 3(3):1534–1537
- Karimi E, Jaafar HZE (2011) HPLC and GC-MS determination of bioactive compounds in microwave obtained extracts of three varieties of *Labisia pumila* benth. *Molecules* 16:6791–6805
- Katerere FR, Grev AI, Nash RJ, Waigh RD (2003) Antimicrobial activity of pentacyclic triterpenes isolated from African combretaceae. *Phytochemistry* 63:81–88
- Li S, Wang C, Wang M, Li W, Matsumoto K, Tang Y (2007) Antidepressant like effects of piperine in chronic mild stress treated mice and its possible mechanisms. *Life Sci* 80(15):1373–1381
- Madhavan V, Yoganasimhan SN, Gurudeva MR (2008) Pharmacognostical studies on Sankhapushphi (*Convolvulus microphyllus* Sieb. Ex Spreng. and *Evolvulus alsinoides* (L.) L. *Indian J Trad Knowl* 7(4):529–541
- Mahato SB, Sen S (1997) Advances in triterpenoid research-1990–1994. *Phytochemistry* 44:115–123
- Martínez-Ruiz A, García HS, Saucedo-Castaneda G, Favela-Torres E (2008) Organic phase synthesis of ethyl oleate using lipases produced by solid-state fermentation. *Appl Biochem Biotechnol* 151(2–3):393–401
- Matsuda H, Ninomiya K, Morikawa T, Yasuda D, Yamaguchi I, Yoshikawa M (2008) Protective effects of amide constituents from the fruit of *Piper chaba* on D-galactosamine/TNF-alpha-induced cell death in mouse hepatocytes. *Bioorg Med Chem Lett* 18(6):2038–2042
- Mu YM, Yanass T et al (2001) Saturated FFAs, palmitic acid and stearic acid, induced apoptosis in human granulose cells. *J Endocrinol* 142:3590–3597
- Ponnamma SU, Manjunath K (2012) GC-MS Analysis of phytochemicals in the methanolic extract of *Justicia wynaadensis* (nees) T. anders. *Int J Pharm Bio Sci* 3(3):570–576
- Prakash UN, Srinivasan K (2010) Gastrointestinal protective effect of dietary spices during ethanol-induced oxidant stress in experimental rats. *Appl Physiol Nutr Metab* 35(2):134–141
- Radzi SM, Basri M, Salleh AB, Ariff A, Mohamad R, Rahman MBAR, Rahman RNZRA (2006) Optimization study of large-scale enzymatic synthesis of ethyl oleate, a liquid wax ester, by response surface methodology. *J Chem Technol Biotechnol* 81:374–380
- Rani LS, Mohan VR, Regini GS, Kalidass C (2009) GC-MS analysis of ethanolic extract of *Pothos scandens* leaf. *J Herb Medi Toxicol* 3:159–160
- Rao CV, Newmark HL, Reddy BS (1998) Chemopreventive effect of squalene on colon cancer. *Carcinogenesis* 19:287–297
- Rauscher FM, Sanders RA, Watkins JB (2000) Effects of piperine on antioxidant pathways in tissues from normal and streptozotocin-induced diabetic rats. *J Biochem Mol Toxicol* 14:329–334
- Rosemeyer J, Bodoprost H (2007) Analysis of phenacyl ester derivatives of fatty acids from human skin surface sebum by reversed-phase HPLC: chromatographic mobility as a function of physico-chemical properties. *Int J Mol Sci* 8:1111–1124
- Scortichini M, Rossi PM (1991) Preliminary *in vitro* evaluation of antimicrobial activity of terpenes and terpenoids towards *Erwinia amylovora* (Burrill). *J Appl Bact* 71:109–112
- Sethiya NK, Nahata A, Mishra SH, Dixit VK (2009) An update on shankpushpi, a cognition-boosting ayurvedia medicine. *J Chin Integr Med* 7(11):1001–1022
- Sharma PV (1983) *Dravyaguna vijinana*, (in Hindi) (Showkamba, Varanasi). Pp:5
- Shi QH, Ma YM et al (2005) Preliminary study on chemical composition and insecticidal activity of root bark from *Periploca sepium*. *Acta Agric Boreali-occidentalis Sin* 14:141–144
- Sunitha S, Nagaraj M, Varalakshmi P (2001) Hepatoprotective effect of lupeol and lupeol linoleate on tissue antioxidant defence system in cadmium-induced hepatotoxicity in rats. *Fitoterapia* 72(5):516–523
- Ukiva M, Akihisa T, Tokuda H, Suzuki H, Mukainaka T, Ichiishi E (2002) Antitumor promoting effects and cytotoxic activity against human cancer lines of triterpene diols and triols from edible chrysanthemum flowers. *Cancer Lett* 177:7–12
- Uma B, Prabhakar K, Rajendran S, Sarayu LY (2009) Studies on GC/MS spectroscopic analysis of some bioactive antimicrobial compounds from *Cinnamomum zeylanicum*. *J Med Plants* 8(31):125–131
- Vijayakumar RS, Nalini N (2006) Piperin, an active principle from *Piper nigrum*, modulates hormonal and apo lipoprotein profiles in hyperlipidemic rats. *J Basic Clin Physiol Pharmacol* 17(2):71–86
- Vuorela P, Leinonen M, Saikku P, Tammela P, Rouha JP, Wennberg T, Vuorela H (2004) *Curr Med Chem* 11:1375–1389
- Wang L, Zhao D, Zhang Z, Zuo C, Zhang Y, Pei YQ, Lo YQ (1999) Trial of antiepilepsirine (AES) in children with epilepsy. *Brain Dev* 21(1):36–40
- Zhang QX, Wang QL, Han JH (1995) Effect of fatty oil in *Periploca sepium* on neural system in mice. *Xi'an Med Univ* 16:43–44