

Supporting information

GC-MS Metabolite profile and identification of unusual homologous cannabinoids in high potency *Cannabis sativa*

Josep Basas-Jaumandreu and F. Xavier C. de las Heras*

¹*Department of Mining, Industrial and ICT – Escola Politècnica Superior d'Enginyeria de Manresa*

Universitat Politècnica de Catalunya (UPC-BarcelonaTech)


Av. Bases de Manresa 61-73

E-08242 Manresa, Catalonia, Spain

*Corresponding author. Telf.: + 34 938 777 236; Fax: + 34 938 777 202; E-mail address: Fxavier.de.las.heras@upc.edu; Orcid 0000-0002-9865-5878; ResearcherID N-9657-2013; Scopus Author ID 7004247478.

Supplementary Data Contents:

- 1S.** Mass spectrum of the pentyl cannabinolic acid (C₅-CBNA) as its TMS derivative identified in the seeds of *C. sativa*.....2
- 2S.** Mass chromatogram *m/z* 303 corresponding to the four isomers (a-d) of cannabinochromene-C₅ CBC (CBC-C₅) identified in hemp seeds and the corresponding mass spectra. The structure of main ions is also showed.....3
- 3S.** Mass chromatogram *m/z* 419 corresponding to the four isomers (a-d) of cannabichromenic acid (CBCA (A)-C₅) identified in hemp seeds and the corresponding mass spectra of the TMS derivative. The meaning of main ions is also showed.....4
- 4S.** Mass chromatograms *m/z* 337 (a) for the identification of cannabinerol (CBNR) and cannabigerol (CBG) and *m/z* 561 (b) in order to represent cannabinerolic (CBNRA) and cannabigerolic (CBGA) acids present in the seeds of hemp (*C. sativa*).....5
- 5S.** Mass spectra of the trimethylsilylated derivative cannabinoid isomers of (*Z*)-cannabinerol (a) and (*E*)-cannabigerol (b) the latter previously reported [31] also as TMS derivative and identified in the total extract of the seeds of *C. sativa*.....6



1S

