



Gene-Centric Meta-Analysis of Lipid Traits in African, East Asian and Hispanic Populations

Citation

Elbers, Clara C., Yiran Guo, Vinicius Tragante, Erik P. A. van Iperen, Matthew B. Lanktree, Berta Almoguera Castillo, Fang Chen, et al. 2012. Gene-centric meta-analysis of lipid traits in African, East Asian and Hispanic populations. PLoS ONE 7(12): e50198.

Published Version

doi:10.1371/journal.pone.0050198

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:10582091>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Gene-Centric Meta-Analysis of Lipid Traits in African, East Asian and Hispanic Populations

Clara C. Elbers^{1,2,3,9}, Yiran Guo^{4,5,9}, Vinicius Tragante^{2,6}, Erik P. A. van Iperen⁷, Matthew B. Lanktree⁸, Berta Almoguera Castillo⁴, Fang Chen⁹, Lisa R. Yanek¹⁰, Mary K. Wojczynski^{11,18}, Yun R. Li⁴, Bart Ferwerda¹, Christie M. Ballantyne¹², Sarah G. Buxbaum^{13,61}, Yii-Der Ida Chen¹⁴, Wei-Min Chen^{9,15}, L. Adrienne Cupples^{16,17}, Mary Cushman⁵⁴, Yanan Duan¹⁸, David Duggan¹⁹, Michele K. Evans⁵⁹, Jyotika K. Fernandes⁵⁵, Myriam Fornage²⁰, Melissa Garcia²¹, W. Timothy Garvey⁵⁶, Nicole Glazer¹⁶, Felicia Gomez¹, Tamara B. Harris²¹, Indrani Halder⁶⁰, Virginia J. Howard⁵⁷, Margaux F. Keller⁴⁹, M. Ilyas Kamboh²⁸, Charles Kooperberg²³, Stephen B. Kritchevsky^{24,25}, Andrea LaCroix²³, Kiang Liu²⁶, Yongmei Liu²⁷, Kiran Musunuru^{29,30,31}, Anne B. Newman³², N. Charlotte Onland-Moret^{2,3}, Jose Ordovas³³, Inga Peter³⁴, Wendy Post³⁵, Susan Redline³⁶, Steven E. Reis²², Richa Saxena^{37,38}, Pamela J. Schreiner³⁹, Kelly A. Volcik⁴⁰, Xingbin Wang²⁸, Salim Yusuf⁴¹, Alan B. Zonderland⁵⁸, Sonia S. Anand⁴¹, Diane M. Becker¹⁰, Bruce Psaty^{42,43}, Daniel J. Rader⁴⁴, Alex P. Reiner²³, Stephen S. Rich⁹, Jerome I. Rotter¹⁴, Michèle M. Sale^{9,45,46}, Michael Y. Tsai⁴⁷, Ingrid B. Borecki¹⁸, Robert A. Hegele⁴⁸, Sekar Kathiresan^{29,30,31}, Michael A. Nalls⁴⁹, Herman A. Taylor, Jr.⁵⁰, Hakon Hakonarson⁴, Suthesh Sivapalaratnam⁵¹, Folkert W. Asselbergs^{3,2,6}, Fotios Drenos⁵², James G. Wilson⁵³, Brendan J. Keating^{4*}

1 Department of Genetics, University of Pennsylvania, School of Medicine, Philadelphia, Pennsylvania, United States of America, **2** Department of Medical Genetics, Biomedical Genetics, University Medical Center, Utrecht, The Netherlands, **3** Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, The Netherlands, **4** Center for Applied Genomics, Abramson Research Center, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, United States of America, **5** BGI-Shenzhen, Shenzhen, People's Republic of China, **6** Department of Cardiology, Division Heart and Lungs, University Medical Center Utrecht, Utrecht, The Netherlands, **7** Department of Clinical Epidemiology, Biostatistics and Bioinformatics, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands, **8** Departments of Medicine and Biochemistry, Schulich School of Medicine and Dentistry, University of Western Ontario, London, Ontario, Canada, **9** Center for Public Health Genomics, University of Virginia, Charlottesville, Virginia, United States of America, **10** GeneSTAR Research Program, Division of General Internal Medicine, Johns Hopkins School of Medicine, Baltimore, Maryland, United States of America, **11** Department of Biostatistics, University of Alabama at Birmingham, Birmingham, Alabama, United States of America, **12** Baylor College of Medicine, Houston, Texas, United States of America, **13** Jackson Heart Study, Jackson State University, Jackson, Mississippi, United States of America, **14** Medical Genetics Institute, Cedars-Sinai Medical Center, Los Angeles, California, United States of America, **15** Department of Public Health Sciences, University of Virginia, Charlottesville, Virginia, United States of America, **16** Boston University, Boston, Massachusetts, United States of America, **17** The National Heart, Lung, Blood Institute's Framingham Heart Study, Framingham, Massachusetts, United States of America, **18** Division of Statistical Genomics and Department of Genetics Washington University School of Medicine, St. Louis, Missouri, United States of America, **19** Translational Genomics Research Institute, Phoenix, Arizona, United States of America, **20** The University of Texas Health Science Center at Houston, Houston, Texas, United States of America, **21** Laboratory for Epidemiology, Demography, and Biometry, National Institute on Aging, National Institutes of Health, Bethesda, Maryland, United States of America, **22** School of Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America, **23** Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, Washington, United States of America, **24** Department of Preventive Medicine, University of Tennessee Health Science Center, Memphis, Tennessee, United States of America, **25** Sticht Center on Aging, Wake Forest University, Winston-Salem, North Carolina, United States of America, **26** Department of Preventive Medicine, Northwestern University Feinberg School of Medicine, Chicago, Illinois, United States of America, **27** Department of Epidemiology and Prevention, Division of Public Health Sciences, Wake Forest University, Winston-Salem, North Carolina, United States of America, **28** Department of Human Genetics, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America, **29** Broad Institute, Cambridge, Massachusetts, United States of America, **30** Harvard Medical School, Boston, Massachusetts, United States of America, **31** Massachusetts General Hospital, Boston, Massachusetts, United States of America, **32** Department of Epidemiology, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America, **33** JM-USDA Human Nutrition Research Center on Aging at Tufts University, Boston, Massachusetts, United States of America, **34** Department of Genetics and Genomic Sciences, Mount Sinai School of Medicine, New York, New York, United States of America, **35** Division of Cardiology, Department of Medicine, Johns Hopkins University, Baltimore, Maryland, United States of America, **36** Brigham and Women's Hospital and Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, United States of America, **37** Center for Human Genetic Research, Massachusetts General Hospital, Boston, Massachusetts, United States of America, **38** Program in Medical and Population Genetics, Broad Institute, Cambridge, Massachusetts, United States of America, **39** School of Public Health, University of Minnesota, Minneapolis, Minnesota, United States of America, **40** Division of Epidemiology, Human Genetics and Environmental Sciences, Human Genetics Center, School of Public Health, University of Texas Health Science Center, Houston, Texas, United States of America, **41** Population Health Research Institute, Hamilton Health Sciences, McMaster University, Hamilton, Ontario, Canada, **42** Cardiovascular Health Research Unit, Departments of Medicine, Epidemiology, and Health Services, University of Washington, Seattle, Washington, United States of America, **43** Group Health Research Institute, Group Health Cooperative, Seattle, Washington, United States of America, **44** Cardiovascular Institute, the Perelman School of Medicine at the University of Pennsylvania, Philadelphia, Pennsylvania, United States of America, **45** Department of Biochemistry and Molecular Genetics, University of Virginia, Charlottesville, Virginia, United States of America, **46** Department of Medicine, University of Virginia, Charlottesville, Virginia, United States of America, **47** Department of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis, Minnesota, United States of America, **48** Robarts Research Institute, University of Western Ontario, London, Ontario, Canada, **49** Laboratory of Neurogenetics, Intramural Research Program, National Institute on Aging, National Institutes of Health, Bethesda, Maryland, United States of America, **50** Jackson State University, Tougaloo College, and the University of Mississippi Medical Center, Jackson, Mississippi, United States of America, **51** Department of Vascular Medicine, Academic Medical Center, Amsterdam, The Netherlands, **52** Centre for Cardiovascular Genetics, Institute of Cardiovascular Science, Faculty of Population Health Sciences, University College London, London, United Kingdom, **53** Department of Physiology and Biophysics, University of Mississippi Medical Center, Jackson, Mississippi, United States of America, **54** Department of Medicine, Thrombosis and Hemostasis Program, University of Vermont, Burlington, Vermont, United States of America, **55** Division of Endocrinology, Diabetes and Medical Genetics, College of Medicine, Medical University of South Carolina, Charleston, SC United States of America, **56** Department of Nutrition Sciences, University of Alabama at

Birmingham, Birmingham, Alabama, United States of America, **57** Department of Epidemiology, University of Alabama at Birmingham, Birmingham, Alabama, United States of America, **58** Laboratory of Personality and Cognition, National Institute on Aging, National Institutes of Health, Bethesda, Maryland, United States of America, **59** Health Disparities Unit, National Institute on Aging, National Institutes of Health, Baltimore, Maryland, United States of America, **60** Heart and Vascular Institute, School of Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America, **61** School of Health Sciences, Department of Epidemiology and Biostatistics, Jackson State University, Jackson, Mississippi, United States of America

Abstract

Meta-analyses of European populations has successfully identified genetic variants in over 100 loci associated with lipid levels, but our knowledge in other ethnicities remains limited. To address this, we performed dense genotyping of ~2,000 candidate genes in 7,657 African Americans, 1,315 Hispanics and 841 East Asians, using the IBC array, a custom ~50,000 SNP genotyping array. Meta-analyses confirmed 16 lipid loci previously established in European populations at genome-wide significance level, and found multiple independent association signals within these lipid loci. Initial discovery and *in silico* follow-up in 7,000 additional African American samples, confirmed two novel loci: rs5030359 within *ICAM1* is associated with total cholesterol (TC) and low-density lipoprotein cholesterol (LDL-C) ($p = 8.8 \times 10^{-7}$ and $p = 1.5 \times 10^{-6}$ respectively) and a nonsense mutation rs3211938 within *CD36* is associated with high-density lipoprotein cholesterol (HDL-C) levels ($p = 13.5 \times 10^{-12}$). The rs3211938-G allele, which is nearly absent in European and Asian populations, has been previously found to be associated with *CD36* deficiency and shows a signature of selection in Africans and African Americans. Finally, we have evaluated the effect of SNPs established in European populations on lipid levels in multi-ethnic populations and show that most known lipid association signals span across ethnicities. However, differences between populations, especially differences in allele frequency, can be leveraged to identify novel signals, as shown by the discovery of *ICAM1* and *CD36* in the current report.

Citation: Elbers CC, Guo Y, Tragante V, van Iperen EPA, Lanktree MB, et al. (2012) Gene-Centric Meta-Analysis of Lipid Traits in African, East Asian and Hispanic Populations. *PLoS ONE* 7(12): e50198. doi:10.1371/journal.pone.0050198

Editor: Yan Gong, College of Pharmacy, University of Florida, United States of America

Received: July 6, 2012; **Accepted:** October 22, 2012; **Published:** December 7, 2012

Copyright: © 2012 Elbers et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The CARE Consortium wishes to acknowledge the support of the National Heart, Lung, and Blood Institute and the contributions of the research institutions, study investigators, field staff and study participants in creating this resource for biomedical research. The following nine parent studies have contributed parent study data, ancillary study data, and DNA samples through the Massachusetts Institute of Technology - Broad Institute (N01-HC-65226) to create this genotype/phenotype database for wide dissemination to the biomedical research community: the Atherosclerosis Risk in Communities (ARIC) study, the Cardiovascular Health Study (CHS), the Cleveland Family Study (CFS), the Cooperative Study of Sickle Cell Disease (CSSCD), the Coronary Artery Risk Development in Young Adults (CARDIA) study, the Framingham Heart Study (FHS), the Jackson Heart Study (JHS), the Multi-Ethnic Study of Atherosclerosis (MESA), and the Sleep Heart Health Study (SHHS). C.C.E is supported by a Rubicon grant from the Netherlands Organization for Scientific Research (NWO). F.W.A. is supported by a clinical fellowship from the Netherlands Organisation for Health Research and Development (ZonMw grant 90700342). Additional funding information can be found in the Supporting Information S1. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: bkeating@mail.med.upenn.edu

☞ These authors contributed equally to this work.

Introduction

Plasma levels of circulating total cholesterol (TC), low-density lipoprotein (LDL-C), high-density lipoprotein (HDL-C) and triglycerides (TG) are associated with coronary artery disease (CAD) and are targets for therapeutic intervention [1]. Multiple environmental and genetic factors influence these plasma lipid levels, with heritability estimated to range from 0.28 to 0.78 in twin and family studies [2]. To date, >100 lipid-associated loci have been described, using studies mainly based on individuals of European ancestry [3]. Together, known variants affecting plasma lipid levels explain 10–12% of the total variance and 25–30% of the genetic variance [3] indicating that other loci and independent signals in established loci are likely to additionally contribute to the trait.

Lipid levels have been demonstrated to vary between ethnic groups [4]. Africans and East Asians have higher levels of HDL-C and lower levels of TG compared to Europeans [5] though the underlying mechanisms of these ethnic differences remain unknown. Genetic contributors to lipid concentrations are less well understood in non-European populations partly due to less well-powered genetic studies being attempted to date and most

genotyping platforms are designed to have optimal coverage in European studies. An important first step towards understanding genetic risk across populations is to establish whether plasma lipid associated loci, identified in Europeans, span across multiple ethnicities or are population-specific. In a recent analysis, most of these known lipid loci had the same direction of association in different ethnic groups as in Europeans, despite presumed differences in linkage disequilibrium (LD) between marker and causal variants in each population [6]. Using regional LD in different ethnicities can help to refine association signals and to distinguish causal variants from correlated markers [7]. Furthermore, independent association signals in established lipid loci in one ethnicity may be useful to highlight causal signal(s) in other ethnicities.

The ITMAT-Broad-CARE (IBC) array (also referred to as the CardioChip or HumanCVD Beadchip [Illumina]) was specifically designed to densely tag ~2000 genes with known or potential roles in lipid and cardiovascular traits using ~50,000 single nucleotide polymorphisms (SNPs) [8]. Sequencing data from European, African American and Yoruba individuals was included for SNP selection in IBC array development. The IBC array drew upon knowledge of lipid metabolism and cardiovascular physiology, as

well as early GWAS and sequencing studies to target efforts towards regions with higher *a priori* evidence of association, reducing cost per sample, and improving efficiency of replication studies. The IBC array has been successfully used for multiple cardiovascular-related phenotypes [9,10,11,12]. Results are reported elsewhere for the association of lipid phenotypes in European-derived cohorts with variants on the IBC array [13].

In this study we set out to discover novel lipid loci, fine map signals to identify causal genes at implicated loci, and gain a greater understanding of the genetic architecture of lipid traits across ethnicities. Here, we have used the IBC array to examine association results for TC, LDL-C, HDL-C and TG across seven non-European study populations, including African Americans (n = 7,657), Hispanics (n = 1,315) and East Asians (n = 841). Using conditional analyses, we sought to identify independent signals from within associated loci. Finally, we assessed the direction of effect in non-Europeans of new and established loci found in European-derived populations, and tested a composite risk score of known loci across ethnicities.

Materials and Methods

Ethics statement

All participants in each of the cohorts gave informed written consent. The Institutional Review Boards (IRBs) of each CARE cohort (i.e., the IRBs for each cohort's field centers, coordinating center, and laboratory center) have reviewed and approved the cohort's interaction with CARE. The study described in this manuscript was approved by the Committee on the Use of Humans as Experimental Subjects (COUHES) of the Massachusetts Institute of Technology.

Participating studies

Data from African-American, Hispanic and East Asian participants from seven cohorts were included for this study (Figure 1). Participants were ≥ 21 years of age. All seven studies contributed individual-level genotypes and phenotypes. Features of the included cohorts are presented in Table S1 and summary statistics are listed in Table S2. Six replication studies were used comprising African American individuals.

Phenotype definitions

Lipid phenotypes were taken from baseline or first measurements for all fasting individuals. All measurements were converted to mmol/L, with TC and HDL-C measurements converted from mg/dL by dividing by 38.67, and TG measurements converted from mg/dL by dividing by 88.57. TG values were $\log(10)$ -transformed as TG values were not normally distributed. LDL-C was calculated according to Friedewald's formula $L \sim C - H - kT$ where C is total cholesterol, H is HDL-C, L is LDL-C, T is TG and k is 0.45 for mmol/L (or 0.20 if measured in mg/dl) [14]. If TG values were >4.51 mmol/L (>400 mg/dL), then LDL-C was treated as a missing value.

Genotyping and quality control

Genotyping in each participating cohort was performed using the IBC array [8]. SNPs were clustered into genotypes using the Illumina Genomestudio software and were subjected to quality control filters at the sample and SNP level, separately within each cohort. Samples were excluded for individual call rates $<90\%$, gender mismatch, and duplicate discordance. SNPs were removed for call rates $<95\%$ or Hardy-Weinberg equilibrium (HWE) $p < 10^{-7}$. Due to low frequency SNPs included in the design, and the aim to capture low frequency variants of large effect across the

combined dataset, we filtered only on minor allele frequency (MAF) < 0.005 .

Statistical analyses

Evaluation of population stratification. Self-reported ethnicity was verified by multidimensional scaling analysis of identity-by-state distances as implemented in PLINK [15], including HapMap panels as reference standards. After pruning of SNPs in linkage disequilibrium ($r^2 > 0.3$), Eigenstrat was used to compute principal components within each ethnic group separately for use as covariates in the regression analyses [16].

Association testing. Association analysis was performed in each study using an additive genetic model with one degree of freedom. Gender stratified analyses were performed using three multivariate models: Model 1, including 10 principal components (PCs); Model 2, including 10 PCs, age, and lipid medication; and Model 3, including 10 PCs, age, lipid medication, type 2 diabetes (T2D), smoking and BMI. The genomic control inflation factor, lambda, was calculated for each cohort and used for within-study correction before meta-analysis. Genomic control inflation factors (λ) ranged from 1.00 to 1.054.

Meta-analyses within each ethnic group were performed by two independent analysts using a fixed-effect inverse-variance approach in two different software packages: MANTEL (www.broadinstitute.org/~debakker/mantel.html) and METAL [17]. Results were highly concordant, reflecting a robust data analyses pipeline. Additionally, the directions of effect of lead SNPs from previously identified loci from the European IBC array meta-analysis [13] were evaluated for consistency in African Americans, Hispanics and Asians. To gauge an appropriate significance threshold, data from the Candidate gene Association Resource (CARE) IBC array studies [18] which is available on dbGAP (www.ncbi.nlm.nih.gov/gap) were employed and it was determined that after accounting for LD, the effective number of independent tests was $\sim 26,500$ for African Americans, $\sim 23,500$ for Hispanics, and $\sim 15,500$ for East Asians. This produces experimental or 'array-wide' statistical thresholds of $p = 1.9 \times 10^{-6}$, $p = 2.1 \times 10^{-6}$ and $p = 3.2 \times 10^{-6}$, respectively, to maintain a false positive rate of 5% for each of the three ethnic groups. While we have adopted these 'array-wide' statistical thresholds for this study, we also highlight loci associated at a more conventional genome-wide significance threshold of $p < 5.0 \times 10^{-8}$.

Additionally, the F^2 statistic was calculated to quantify the proportion of total variation due to heterogeneity, as described previously [19].

Conditional Analyses. Loci harboring evidence for association of $P < 1 \times 10^{-5}$ in African Americans were examined for the presence of multiple, independent signals via conditional analyses in PLINK [15]. A term was added to the regression model including the lead SNP as a covariate, and SNPs within a ± 500 kb region were evaluated for significance. A locus-specific Bonferroni correction, as employed in previous IBC studies [20], was applied to determine significance of independent signals within candidate genes genotyped at each locus. On average, the windows contained 195.2 (± 107.0) variants with a range between 12 for *ACADL* and 359 for *PCSK9*. Because of limited power due to low sample size, we did not perform conditional analyses in Hispanics and East Asians.

Genetic Risk Score Analyses and direction of effect. Within each ethnic group, we generated a genetic risk score using 28 SNPs for TC, 20 SNPs for LDL-C, 24 SNPs for HDL-C, and 21 SNPs for TG that had been found to be array-wide significant ($p = 2.6 \times 10^{-6}$) in the European-ancestry IBC meta-analysis [21] (Table S3), weighted by the beta as described




Figure 1. Schematic design of study for the multi-ethnic IBC-Lipid association meta-analysis. The workflow includes primary analyses and secondary analyses. Details can be found in the text. doi:10.1371/journal.pone.0050198.g001

previously [22,23]. To account for missing data we adjusted the values for the number of genotyped risk alleles per individual. We evaluated for each ethnic group the contribution of the weighted genetic risk score to TC, HDL-C, LDL-C and TG in linear regression models adjusting for 10 PCs. Additionally, we compared the relative betas across quartiles of risk by linear regression. These loci were additionally investigated to study direction of effect across ethnicities.

Replication

In order to confirm putative novel loci, we replicated previously undetected lipid signals ($p < 1.0 \times 10^{-5}$) in 7,000 African American individuals from six replication cohorts and in 61,636 samples from the European-ancestry IBC meta-analysis [21]. Recent power analyses suggest that large-scale multi-ethnic association studies may have greater statistical power to detect causal alleles because of random genetic drift elevating global risk variants to higher allele frequency in some populations [24]. All but one replication study provided summary results of SNPs that were genotyped on platforms other than the IBC array, or imputed using 1000 Genomes data. Features of the replication datasets included in this meta-analysis are described in Table S1.

Results

Meta-analyses of African, Hispanic and East Asian populations

Meta-analyses of IBC array association results for plasma TC, LDL-C, HDL-C and TG levels in five African American studies ($n = 7,657$), two Hispanic studies ($n = 1,315$) and three East Asian studies ($n = 841$) were performed independently. Results of different association models did not differ substantially. Therefore, results of model 1, an additive model with 10 PCs as covariates, are presented in the main text (Table 1) and results of other models are presented in the supplements (Table S4). After fixed-effect inverse-variance meta-analysis, we found that 23, five and two loci in African Americans, Hispanics and East Asian samples respectively, were significantly associated with a lipid trait at their respective array-wide significance thresholds, with twelve, three and one loci respectively surpassing the traditional genome-wide significance threshold (see Table 1; Figure 1). Two of these loci, intercellular adhesion molecule 1 (*ICAM1*) and CD36 molecule thrombospondin receptor (*CD36*), have not previously been reported to be associated with a lipid trait in a large-scale genomic study (Figure 2).

African Americans. We found five independent loci that were associated with TC at the genome-wide significance threshold. Four of these signals were SNPs lying within previously described loci: *LDLR* (rs6511720, $p = 1.4 \times 10^{-13}$); *CELSR2*





Figure 2. Regional plots for novel lipid loci with array-wide significant regions in IBC meta-analysis of African ancestry. A. *CD36* region, **B.** *ICAM1* region. Loci are shown as the lead SNP with a flanking region depicting the candidate gene and nearby genes included on the array. The purple diamond represents the lead SNP in the IBC meta-analysis and the dots represent the surrounding SNPs, with the different colors showing the LD relationship with the lead SNP based on YRI HapMap II information. $-\log_{10}$ p-values for association with HDL-C (for *CD36*) and TC (for *ICAM1*) are shown for each SNP (left-hand axis). Recombination rates in YRI HapMap II is shown in blue traces (right-hand axis). doi:10.1371/journal.pone.0050198.g002

(rs12740374, $p = 4.4 \times 10^{-13}$); *APOE* (rs389261, $p = 2.1 \times 10^{-11}$) and *PCSK9* (rs11806638, $p = 2.00 \times 10^{-9}$), while one signal was a novel SNP within *ICAM1* (rs5030359, $p = 5.2 \times 10^{-9}$). Three SNPs in the previously known loci, *CELSR2* (rs12743074, $p = 1.9 \times 10^{-17}$), *APOE* (rs389261, $p = 1.0 \times 10^{-12}$) and *PCSK9* (rs11800231, $p = 1.0 \times 10^{-10}$) reached genome-wide significance for association with LDL-C. We also identified a novel signal within *ICAM1* (rs5030359, $p = 1.1 \times 10^{-7}$) that is associated with LDL-C in African Americans at array-wide significance. Genome-wide significant association with HDL-C was observed for three SNPs in previously identified loci within *CETP* (rs17231520 $p = 2.0 \times 10^{-46}$), *LPL* (rs13702 $p = 1.3 \times 10^{-9}$) and *LIPC* (rs2070895 $p = 4.2 \times 10^{-8}$). Of the array-wide significant loci, rs3211938 within *CD36* ($p = 3.1 \times 10^{-7}$) has been previously described to be associated with HDL-C in a candidate gene study of 2,020 African Americans [25] but had not previously been identified in a large-scale genomic study. For TG, we identified one association signal, rs12721054, within the previously reported *APOE* locus with TG with at genome-wide significance ($p = 1.0 \times 10^{-21}$).

Hispanics. Genome-wide significant association with HDL-C was observed for two SNPs in previously identified loci within *CETP* (rs3764261, $p = 3.4 \times 10^{-11}$) and *LIPC* (rs8034802, $p = 1.8 \times 10^{-8}$). For TG, we identified one genome-wide signal within the previously reported *APOA5* locus (rs10750097, $p = 2.1 \times 10^{-12}$). Genome-wide significant association for TC and LDL-C was not observed in our Hispanic populations.

East Asians. In East Asians, the rs662799 variant within *ζNF259/APOA5* was significantly associated with TG ($p = 1.6 \times 10^{-13}$). The opposite allele of the same SNP was study-wide significantly associated with HDL-C. Genome-wide or study-wide significant genetic association was not observed for LDL-C or TC in our East Asian populations.

Independent signals within single genetic loci in African Americans

The current investigation using the IBC array included rare SNPs at candidate loci collected in sequencing data from Europeans and Africans and dense genotyping, which can potentially be used to identify independent signals for lipids within genes at known or novel loci. We repeated association studies conditioning on the lead SNP in 23 loci with $P < 1.0 \times 10^{-5}$. After Bonferroni correction for the number of SNPs at each candidate gene locus, we found independent lipid signals at the *LDLR*, *APOE*, *PCSK9* and *APOB* loci for TC, at the *APOE*, *PCSK9*, *LDLR*, and *APOB* loci for LDL-C, at the *APOC1/APOE*, and *LPL* loci for TG and at the *CETP*, *LPL*, *CD36* and the *TRADD/LCAT* for HDL-C (Table 2).

Three loci harbored two independent signals at genome-wide significance. The alleles rs6511720-G (risk allele frequency [RAF] = 0.86) and rs17242787-T (RAF = 0.98) within the *LDLR* gene showed association with TC with a p-value of 1.04×10^{-13} and 4.7×10^{-9} respectively in the original analyses. After conditioning on rs6511720-G, the p value for rs17242787-T remained significant ($p = 2.4 \times 10^{-10}$). Also for LDL-C, we found two independent genome-wide significant signals within the *APOE* locus: rs389261-A (RAF = 0.25) and rs283813-T (RAF = 0.67). Furthermore, the SNPs rs17231520-A (RAF = 0.07) and

rs4783961-A (RAF = 0.44) within the *CETP* gene were both strongly associated with HDL-C and after conditioning on the lead signal, the secondary signal remained significant with $p = 2.8 \times 10^{-20}$. Interestingly, the newly identified *CD36* locus also harbored two independent signals, with the second signal showing association with locus-wide significance. The r^2 between the two SNPs in HapMap-YRI was 0.118.

Replication

In order to confirm putative novel signals, we carried out *in silico* follow-up of ten SNPs within novel loci and previously unreported SNPs within known lipid-associated loci ($P < 1.0 \times 10^{-5}$) in six African American studies, comprising together 7,000 samples. Only HeartSCORE was genotyped using the IBC array and provided association results for all SNPs. All other replication studies contributed association results for up to seven genotyped and imputed SNPs. Imputed SNPs were only included in the study when passing the 95% confidence threshold. Combined meta-analysis of the discovery and replication studies led to genome-wide significant signals at the *CD36* locus ($p = 13.5 \times 10^{-12}$; Table 3) for association with HDL-C. A signal within *ACADL* was not significant after meta-analysis of the discovery and replication studies. However, the direction of effect was consistent with our discovery dataset in three of six studies, so it is possible that the signal has a weak effect and the locus is undetectable due to limited statistical power. Also, previously unidentified signals in known lipid loci showed genome-wide significant association in the combined discovery and replication meta-analysis: rs11806638 within *PCSK9* was found to be associated with TC; rs389261 within *APOE* was associated with LDL-C levels; rs17231520 within the *CETP* locus and rs35673026 within the *LCAT* locus were found to be associated with HDL-C; and rs12721054 within *APOE* was associated with TG levels (Table 3).

Comparison of lipid loci in African Americans to Europeans

Utilizing the results of each of the meta-analyses from the three available ethnicities, we sought to refine localization of known lipid signals or reveal novel independent signals within known loci based upon differential LD (see Table 1). The dense genotyping within each locus on the IBC array enabled detailed comparisons of loci that harbored array-wide significant SNPs in Africans Americans, Hispanics and East Asians as well as in the IBC meta-analysis of up to 61,636 individuals of Europeans ancestry [21] (see Table 1 and Table S3).

The strongest signal for HDL-C in African Americans is rs17231520 within *CETP* ($p = 2.0 \times 10^{-46}$; Table 1). This SNP is associated with HDL-C in the same direction in Europeans with $p = 3.3 \times 10^{-4}$. However, in Europeans there is less power to detect this signal at array-wide significance, as the MAF in Europeans is only 0.2% (versus 7% in African Americans) and was screened out in many European studies for the IBC meta-analysis. Furthermore, rarer variants are often not correctly clustered optimally during QC, making them less likely to pass the standard quality control (including genotyping threshold or HWE check). This is also observed for the most strongly associated SNPs within *CD36*

Table 1. Associated loci with lipid traits in individuals of African, East Asian and Hispanic ancestry.

African-Americans											Results IBC Europeans					
Trait	Chr	BP	Candidate Gene	SNP	Risk allele/RAF	Beta (SE)	P	% I2	RAF	Beta (SE)	P	% I2	RAF	Beta (SE)	P	% I2
TC	1	55290748	PCSK9	rs11806638	C	0.68	1.99×10 ⁻⁹	8	0.94	0.01 (0.013)	0.28	0				0
TC	1	109619113	CELSR2	rs12740374	G	0.75	4.43×10 ⁻¹³	0	0.76	0.13 (0.008)	1.79×10 ⁻⁶³	34.5				34.5
TC	2	21119200	APOB	rs12720826	T	0.87	7.84×10 ⁻⁸	0	0.999	NA	NA	NA				NA
TC	2	210769915	ACADL	rs6739874	T	0.05	4.65×10 ⁻⁶	0	0.18	NA	NA	NA				NA
TC	19	10249462	ICAM1	rs5030359	G	0.99	5.22×10 ⁻⁹	0	0.998	NA	NA	NA				NA
TC	19	11063306	LDLR	rs6511720	G	0.86	1.39×10 ⁻¹³	26.2	0.88	0.17 (0.009)	1.81×10 ⁻⁷³	24.3				24.3
TC	19	50112183	APOE	rs389261	A	0.25	2.07×10 ⁻¹¹	0	0.001	NA	NA	NA				NA
LDL-C	1	55290528	PCSK9	rs11800231	G	0.83	1.02×10 ⁻¹⁰	37.3	0.96	0.009 (0.015)	0.60	0				0
LDL-C	1	109619113	CELSR2	rs12740374	G	0.75	1.92×10 ⁻¹⁷	0.8	0.75	0.12 (0.007)	2.62×10 ⁻⁶⁶	41.6				41.6
LDL-C	2	21141826	APOB	rs562338	G	0.4	6.54×10 ⁻⁸	0	0.82	0.12 (0.008)	7.38×10 ⁻⁵¹	0				0
LDL-C	17	61641042	APOH	rs1801689	C	0.007	4.70×10 ⁻⁷	25.5	0.03	0.11 (0.016)	8.57×10 ⁻¹²	0				0
LDL-C	19	10249462	ICAM1	rs5030359	G	0.99	1.09×10 ⁻⁷	0	0.998	NA	NA	NA				NA
LDL-C	19	11059187	LDLR	rs17248720	C	0.73	7.13×10 ⁻¹⁰	0	0.88	0.16 (0.009)	8.81×10 ⁻⁷²	37.5				37.5
LDL-C	19	50112183	APOE	rs389261	A	0.25	1.03×10 ⁻¹²	0	0.001	NA	NA	NA				NA
HDL-C	7	80138385	CD36	rs3211938	G	0.09	3.09×10 ⁻⁷	28.3	0.0005	NA	NA	NA				NA
HDL-C	8	19868772	LPL	rs13702	C	0.51	1.34×10 ⁻⁹	46.8	0.3	0.04 (0.002)	3.57×10 ⁻⁷⁴	24				24
HDL-C	15	56511231	LIPC	rs2070895	A	0.51	4.16×10 ⁻⁸	0	0.21	0.04 (0.002)	9.76×10 ⁻⁵⁸	41.2				41.2
HDL-C	16	5553328	CETP	rs17231520	A	0.07	2.03×10 ⁻⁴⁶	14.5	0.002	0.16 (0.044)	3.28×10 ⁻⁰⁴	0				0
HDL-C	16	66534352	LCAT	rs35673026	T	0.004	2.40×10 ⁻⁶	0	0.001	NA	NA	NA				NA
TG	2	27584444	GCKR	rs1260326	T	0.15	5.54×10 ⁻⁷	13.7	0.41	0.06 (0.003)	1.56×10 ⁻⁸³	32.6				32.6
TG	8	19864004	LPL	rs328	C	0.93	1.74×10 ⁻⁷	19.9	0.9	0.08 (0.005)	5.73×10 ⁻¹⁶	38.1				38.1
TG	11	116170289	APOA5	rs9804646	C	0.65	2.55×10 ⁻⁶	2.3	0.92	0.02 (0.005)	1.57×10 ⁻⁷	11.4				11.4
TG	19	50114427	APOE	rs12721054	A	0.89	1.01×10 ⁻²¹	2.7	0.998	NA	NA	NA				NA
East Asians											Results IBC Europeans					
Trait	Chr	BP	Candidate Gene	SNP	Risk allele/RAF	Beta (SE)	P	% I2	RAF	Beta (SE)	P	% I2	RAF	Beta (SE)	P	% I2
HDL-C	11	116168917	ZNF259/APOA5	rs662799	A	0.71	8.91×10 ⁻⁷	0	0.93	0.03 (0.004)	1.42×10 ⁻¹⁸	28.3				28.3
TG	11	116168917	ZNF259/APOA5	rs662799	G	0.29	1.57×10 ⁻¹³	28.4	0.07	0.1102 (0.0058)	5.93×10 ⁻⁸⁹	58				58
Hispanics											Results IBC Europeans					
Trait	Chr	BP	Candidate Gene	SNP	Risk allele/RAF	Beta (SE)	P	% I2	RAF	Beta (SE)	P	% I2	RAF	Beta (SE)	P	% I2
TC	1	109619113	CELSR2	rs12740374	G	0.78	7.88×10 ⁻⁷	41.5	0.76	0.13 (0.008)	1.79×10 ⁻⁶³	34.5				34.5
TC	12	130896484	MMP17	rs10902456	G	0.73	6.96×10 ⁻⁶	0	0.69	-0.005 (0.006)	0.40	0				0
LDL-C	1	109619113	CELSR2	rs12740374	G	0.78	1.10×10 ⁻⁷	25.8	0.76	0.13 (0.008)	2.62×10 ⁻⁶⁶	34.5				34.5
LDL-C	19	47603609	LPIE	rs34052647	G	0.96	9.65×10 ⁻⁶	0	0.999	NA	NA	NA				NA

Table 1. Cont.

Hispanics		Results IBC Europeans											
Trait	Chr	BP	Candidate Gene	SNP	Risk allele	RAF	Beta (SE)	P	% I2	RAF	Beta (SE)	P	% I2
HDL-C	1	202392868	REN	rs11571087	T	0.01	0.28 (0.060)	2.58 × 10 ⁻⁶	53	0.03	0.006 (0.006)	0.89	13.1
HDL-C	3	149905771	AGTR1	rs12721308	G	0.01	0.27 (0.060)	4.95 × 10 ⁻⁶	0	0.001	NA	NA	NA
HDL-C	15	56512084	LIPC	rs8034802	A	0.48	0.07 (0.012)	1.82 × 10 ⁻⁸	2.5	0.28	0.03 (0.002)	4.86 × 10 ⁻⁴³	18.2
HDL-C	16	55550825	CETP	rs3764261	A	0.31	0.08 (0.012)	3.42 × 10 ⁻¹¹	63.2	0.31	0.07 (0.002)	8.45 × 10 ⁻²⁷⁰	65.3
TG	11	116169250	APOA5	rs10750097	G	0.4	0.14 (0.020)	2.14 × 10 ⁻¹²	0	0.21	0.07 (0.004)	4.16 × 10 ⁻⁹³	52.9

Chr chromosome, BP base pair, RAF risk allele frequency, SE standard error.
doi:10.1371/journal.pone.0050198.t001

(rs3211938) and *LCAT* (rs35673026) for HDL-C in African-Americans, as they show the same direction of effect in Europeans, but do not reach significance, given low MAF and absence in the majority of European studies for IBC meta-analysis. For two loci, *LIPC* and *LPL*, the strongest associated SNP in African Americans for HDL-C was the same or among the most highly associated SNPs in Europeans. Also, for the LDL-C-associated loci *CELSR2*, *APOB*, *APOH* and *LDLR*, the strongest signals in African Americans did overlap or represented similar signals that were highly associated with LDL-C in Europeans. The newly identified SNP for LDL-C, rs5030359 within *ICAMI*, has an observed MAF of 0.8% in African Americans and 0.2% in Europeans. In Europeans, this SNP is not associated with LDL-C ($p = 0.3231$), but the SNP is only present in very few European studies that are included in the IBC meta-analysis. The most associated signals within *PCSK9* and *APOE* in African Americans are different, independent signals compared to the most associated SNPs within these loci in Europeans. Again, both signals are common in African Americans and have very low frequencies in Europeans: MAF for SNPs in *PCSK9* and *APOE* are 17% and 25% in African Americans and 0.5% and 0.1% in Europeans respectively.

Among the array-wide statistically significant loci that were associated with TG in African Americans, three SNPs within *GCKR*, *LPL* and *APOA5* were the same as or amongst the most highly associated SNPs in Europeans. SNP rs12721054 in *APOE* appeared to be a novel independent signal for TG in African Americans. This SNP showed an opposite effect in European-derived cohorts, although it was observed rarely in the meta-analysis of European populations (MAF = 0.2%) [13].

For TC, we observed the same pattern as for other lipid traits. The strongest associated SNPs within loci associated with TC overlapped with the same signals in Europeans (SNPs within *CELSR2*, *APOB*, *LDLR* and *APOE*), or were independent signals in African Americans that could not be replicated in Europeans because of low frequency (*PCSK9*, *ACADL* and *ICAMI*).

Direction of effect concordance with lead SNPs identified in European populations

Direction of effect across different ethnicities was studied for 28 previously established TC risk loci, 20 LDL-C loci, 24 HDL-C loci, and 21 TG associated loci. Not all SNPs passed the initial quality control, so number of investigated SNPs differed by trait and ethnicity (Table S3).

Concordance in direction of effect was observed for 21/27 ($p = 0.033$), 15/20 ($p = 0.102$), 16/23 ($p = 0.176$) and 19/21 ($p = 0.004$) association signals for TC, LDL-C, HDL-C and TG, respectively, between Europeans and African Americans; 23/28 ($p = 0.011$), 16/20 ($p = 0.047$), 21/23 ($p = 0.002$) and 19/21 ($p = 0.004$) SNPs were concordant in direction of effect for TC, LDL-C, HDL-C and TG respectively between Europeans and Hispanics. Finally, 17/24 SNPs for TC ($p = 0.140$), 11/16 SNPs for LDL-C ($p = 0.279$), 16/29 SNPs for HDL-C ($p = 0.196$) and 17/21 ($p = 0.035$) SNPs for TG were concordant between Europeans and East Asians (Table S3).

Genetic risk score analysis

To study whether we could find elevated lipid levels in multi-ethnic samples with cumulative numbers of risk alleles that were previously found to be associated in Europeans, we evaluated the contribution of the weighted genetic risk score for lipids in linear regression models adjusting for 10 PCs and compared the relative beta's ratios across quartiles of risk. We demonstrated a significant per quartile risk effect in African-Americans (ranging from $p < 10^{-10}$ for TG to $p < 10^{-33}$ for HDL-C), Hispanics (ranging

Table 2. Loci with significant evidence of independent lipid association signals.

Trait	Gene	SNP	Chr	BP	Risk Allele	RAF	Original signal		Second signal		Third signal		r2 with lead SNP	
							Beta (SE)	P	Beta (SE)	P	Beta (SE)	P		
TC	LDLR	rs6511720	19	11063306	G	0.86	0.18 (0.024)	1.39×10 ⁻¹³						
		rs17242787	19	11063460	T	0.98	0.35 (0.059)	4.67×10 ⁻⁹	0.37 (0.059)	2.44×10 ⁻¹⁰		0.004		
	APOE	rs389261	19	50112183	A	0.25	0.13 (0.0120)	2.07×10 ⁻¹¹						
		rs283813	19	50081014	T	0.67	0.09 (0.018)	3.60×10 ⁻⁷	0.09 (0.018)	1.30×10 ⁻⁶		0.001		
		rs12721054	19	50114427	A	0.88	0.15 (0.027)	4.75×10 ⁻⁸		0.10 (0.027)	1.85×10 ⁻⁴		0.025	
	PCSK9	rs11806638	1	55290748	C	0.68	0.11 (0.018)	1.99×10 ⁻⁹						
		rs505151	1	55301775	G	0.24	0.11 (0.019)	7.50×10 ⁻⁹	0.09 (0.0120)	9.78×10 ⁻⁶		0.085		
	APOB	rs12720826	2	21119200	T	0.87	0.13 (0.025)	7.84×10 ⁻⁸						
		rs562338	2	21141826	G	0.4	0.09 (0.017)	2.00×10 ⁻⁷	0.07 (0.018)	1.20×10 ⁻⁴		0.054		
	LDL-C	APOE	rs389261	19	50112183	A	0.25	0.14 (0.020)	1.03×10 ⁻¹²					
rs283813			19	50081014	T	0.67	0.12 (0.018)	7.27×10 ⁻¹²	0.12 (0.018)	3.30×10 ⁻¹¹		0.001		
		rs166907	19	50078695	G	0.12	0.08 (0.026)	2.92×10 ⁻³		0.16 (0.031)	1.96×10 ⁻⁷		0.001	
PCSK9		rs11800231	1	55290528	G	0.82	0.14 (0.022)	1.02×10 ⁻¹⁰						
		rs505151	1	55301775	G	0.24	0.12 (0.019)	1.91×10 ⁻¹⁰	0.10 (0.020)	2.15×10 ⁻⁷		0.12		
		rs1165287	1	55292800	A	0.25	0.09 (0.02)	7.42×10 ⁻⁶		0.08 (0.021)	1.17×10 ⁻⁴		0.08	
LDLR		rs17248720	19	11059187	C	0.73	0.11 (0.019)	7.13×10 ⁻¹⁰						
		rs6511720	19	11063306	T	0.86	0.19 (0.024)	7.91×10 ⁻¹⁵	0.15 (0.030)	2.93×10 ⁻⁷		0.251		
		rs17242787	19	11063460	T	0.98	0.31 (0.06)	1.76×10 ⁻⁷		0.33 (0.062)	8.12×10 ⁻⁸		0.071	
APOB		rs562338	2	21119200	G	0.4	0.09 (0.017)	6.54×10 ⁻⁸						
		rs12720826	2	21119200	T	0.87	0.13 (0.025)	7.84×10 ⁻⁸	0.11 (0.026)	1.30×10 ⁻⁴		0.054		
HDL-C		CETP	rs17231520	16	55553328	A	0.07	0.19 (0.013)	2.03×10 ⁻⁴⁶					
			rs4783961	16	55552395	A	0.44	0.09 (0.007)	6.08×10 ⁻⁴⁰	0.06 (0.007)	2.83×10 ⁻²⁰		0.165	
			rs7499892	16	55564091	C	0.62	0.07 (0.007)	7.27×10 ⁻²⁴		0.04 (0.006)	6.40×10 ⁻⁹		0.078
	LPL	rs13702	8	19868772	C	0.51	0.04 (0.007)	1.34×10 ⁻⁹						
		rs3289	8	19867472	T	0.93	0.07 (0.013)	5.07×10 ⁻⁸	0.06 (0.013)	2.70×10 ⁻⁵		0.047		
	CD36	rs3211938	7	80138385	G	0.09	0.06 (0.012)	3.09×10 ⁻⁷						
		rs3211849	7	80121259	G	0.54	0.02 (0.007)	5.50×10 ⁻³	0.03 (0.007)	1.33×10 ⁻⁵		0.118		
	TRADD/ LCAT	rs35673026	16	66534352	T	0.004	0.28 (0.059)	2.40×10 ⁻⁶						
		rs2233455	16	65765434	T	0.29	0.03 (0.007)	3.72×10 ⁻⁶	0.03 (0.007)	2.78×10 ⁻⁶		0		
	TG	APOC1/ APOE	rs12721054	19	50114427	A	0.89	0.12 (0.013)	1.01×10 ⁻²¹					
rs7258987			19	50124360	T	0.03	0.11 (0.024)	2.14×10 ⁻⁶	0.11 (0.024)	3.42×10 ⁻⁶		0.003		
LPL		rs328	8	19864004	C	0.93	0.08 (0.016)	1.74×10 ⁻⁷						
		rs3289	8	19867472	G	0.07	0.08 (0.016)	2.82×10 ⁻⁶	0.07 (0.016)	1.52×10 ⁻⁵		0.003		

Chr chromosome, BP base pair, RAF risk allele frequency, SE standard error.
doi:10.1371/journal.pone.0050198.t002

from $p < 10^{-1}$ for LDL-C to $p < 10^{-23}$ for TC) and East Asians (ranging from $p < 0.02$ for HDL-C to $p < 10^{-6}$ for TG) (see Table 4). Quartiles based on weighted risk alleles and lipid level distribution for each ethnicity is shown in Figure S1.

Discussion

The current study reports a meta-analysis of lipid association studies in African Americans, Hispanics and East Asians using the IBC array, and has identified two novel loci associated with TC and LDL-C levels (rs5030359 in *ICAM1*) and HDL-C levels (rs3211938 in *CD36*) in African Americans. Additionally, we have uncovered multiple independent association signals within estab-

lished lipid loci, demonstrating the value of dense SNP genotyping to uncover genetic variation associated with lipid levels. Furthermore, we have evaluated the impact of established SNPs, previously associated with lipids in Europeans populations, on lipid levels in three additional populations, showing that many known association signals for lipids span across ethnicities.

CD36

This study shows association between the nonsense coding variant rs3211938-G in *CD36* and HDL-C levels at conventional genome-wide significance for African Americans ($p < 5 \times 10^{-9}$). This SNP has previously been reported to be associated with

Table 3. Replication results of nine signals in 7,000 African Americans.

Novel loci		Discovery Set (N = 7,657)				Replication Set (N = 7,000)				Combined Set (N = 14,657)			
Trait	Candidate Gene	SNP	Risk allele/RAF	Beta (SE)	P	Beta (SE)	P	Beta (SE)	P	Beta (SE)	P		
TC	ACADL	rs6739874	T	0.05 0.17 (0.038)	4.65 × 10 ⁻⁶	0.01 (0.043)	0.86	0.11 (0.029)	1.72 × 10 ⁻⁴				
TC	ICAM1	rs5030359	G	0.99 0.57 (0.098)	5.22 × 10 ⁻⁹	0.48 (0.295)	0.10	0.44 (0.095)	4.33 × 10 ⁻⁶				
LDL-C	ICAM1	rs5030359	G	0.99 0.51 (0.096)	1.09 × 10 ⁻⁷	0.47 (0.264)	0.08	0.40 (0.089)	8.74 × 10 ⁻⁶				
HDL-C	CD36	rs3211938	G	0.09 0.06 (0.012)	3.09 × 10 ⁻⁷	0.06 (0.013)	3.12 × 10 ⁻⁶	0.06 (0.009)	3.49 × 10 ⁻¹²				
unidentified signals in previously known loci													
Trait	Candidate Gene	SNP	Risk allele/RAF	Beta (SE)	P	Beta (SE)	P	Beta (SE)	P	Beta (SE)	P		
TC	PCSK9	rs11806638	C	0.68 0.11 (0.018)	1.99 × 10 ⁻⁹	0.15 (0.029)	3.78 × 10 ⁻⁷	0.12 (0.016)	2.38 × 10 ⁻¹⁴				
LDL-C	APOE	rs389261	A	0.25 0.14 (0.020)	1.03 × 10 ⁻¹²	0.13 (0.028)	4.43 × 10 ⁻⁶	0.14 (0.016)	1.70 × 10 ⁻¹⁷				
HDL-C	CETP	rs17231520	A	0.07 0.19 (0.013)	2.03 × 10 ⁻⁴⁶	0.18 (0.021)	2.35 × 10 ⁻¹⁷	0.18 (0.011)	1.70 × 10 ⁻⁶²				
HDL-C	LCAT	rs35673026	T	0.005 0.28 (0.059)	2.40 × 10 ⁻⁶	0.05 (0.12)	0.71	0.24 (0.053)	9.06 × 10 ⁻⁶				
TG	APOE	rs12721054	A	0.89 0.12 (0.013)	1.01 × 10 ⁻²¹	0.05 (0.008)	4.71 × 10 ⁻¹³	0.07 (0.007)	1.31 × 10 ⁻²⁸				

RAF risk allele frequency, SE standard error.
doi:10.1371/journal.pone.0050198.t003

Table 4. Risk score analysis of lipid profile in multiethnic populations, using weighted score of known lipid SNPs.

African Americans				
Trait	TC	LDL-C	HDL-C	TG
Beta (SE)	0.12 (0.011)	0.08 (0.01)	0.05 (0.004)	0.02 (0.004)
P	7.57×10^{-23}	5.45×10^{-15}	3.13×10^{-33}	4.83×10^{-10}
Quartiles of Risk Alleles				
Q1 Beta (SE)	ref	ref	ref	ref
Q2 BETA (SE)	0.12 (0.034)	0.12 (0.033)	0.05 (0.014)	0.06 (0.017)
P	2.99×10^{-04}	3.07×10^{-04}	1.23×10^{-04}	8.04×10^{-04}
Q3 BETA (SE)	0.25 (0.035)	0.16 (0.034)	0.10 (0.014)	0.05 (0.017)
P	1.88×10^{-12}	1.88×10^{-06}	2.45×10^{-12}	2.62×10^{-03}
Q4 BETA (SE)	0.33 (0.036)	0.26 (0.034)	0.16 (0.013)	0.12 (0.017)
P	1.98×10^{-20}	7.87×10^{-15}	6.73×10^{-32}	3.37×10^{-13}
Hispanics				
Trait	TC	LDL-C	HDL-C	TG
Beta (SE)	0.17 (0.017)	0.08 (0.015)	0.07 (0.009)	0.07 (0.009)
P	2.16×10^{-23}	2.07×10^{-07}	4.43×10^{-14}	4.43×10^{-14}
Quartiles of Risk Alleles				
Q1 Beta (SE)	ref	ref	ref	ref
Q2 BETA (SE)	0.09 (0.063)	0.06 (0.046)	0.06 (0.019)	0.05 (0.036)
P	0.15	0.19	0.005	0.15
Q3 BETA (SE)	0.26 (0.063)	0.09 (0.048)	0.08 (0.02)	0.02 (0.033)
P	2.78×10^{-05}	0.06	9.65×10^{-05}	0.56
Q4 BETA (SE)	0.47 (0.056)	0.24 (0.047)	0.166	0.18 (0.034)
P	6.42×10^{-17}	2.65×10^{-07}	3.42×10^{-16}	2.68×10^{-07}
East Asians				
Trait	TC	LDL-C	HDL-C	TG
Beta (SE)	0.0642 (0.0288)	0.0137 (0.0272)	0.105 (0.044)	0.088 (0.0186)
P	0.03	0.62	0.02	2.29×10^{-06}
Quartiles of Risk Alleles				
Q1 Beta (SE)	ref	ref	ref	ref
Q2 BETA (SE)	0.32 (0.097)	-0.03 (0.084)	0.09 (0.039)	0.08 (0.059)
P	8.37×10^{-04}	0.74	0.02	0.17
Q3 BETA (SE)	0.22 (0.094)	-0.08 (0.092)	0.13 (0.038)	0.10 (0.063)
P	0.02	0.37	6.82×10^{-04}	0.1
Q4 BETA (SE)	0.25 (0.092)	0.05 (0.085)	0.19 (0.039)	0.27 (0.058)
P	0.007	0.53	2.27×10^{-06}	2.39×10^{-06}

ref reference group, SE standard error.
doi:10.1371/journal.pone.0050198.t004

increased HDL-C levels ($p = 0.00018$), decreased TG levels ($p = 0.0059$) and protection against metabolic syndrome ($p = 0.0012$) in a candidate gene study including 2,020 African Americans that did not overlap with samples in our meta-analyses [25]. Also, a variant within *CD36* was associated with LDL levels in two small studies [26,27]. The *CD36* finding is present in an accompanying paper [52] from the wider NHLBI CARE lipid studies which essentially uses the same discovery cohorts for African Americans that we present here although our analysis differs in that (a) it screened out related individuals (b) it takes additional covariates into account through the use of the three

multivariate models and (c) our analysis filtered more stringently on R^2 and (d) we replicated these findings in additional studies.

CD36, which is present on gustatory, olfactory and intestinal epithelial cells, is involved in the orosensory perception of fatty acids [28,29]. Also, lipid ingestion affects lingual *CD36* expression in mice [30]. Therefore, *CD36* may influence fat intake, and hence, serum lipid levels. SNPs within *CD36*, other than the one we found in this study, were linked to obesity in a case-control study [31]. However, this finding could not be replicated in a larger cohort [32]. In mouse models, *CD36* deficiency impairs intestinal lipid secretion and results in hypertriglyceridemia [33]

and others show that CD36 deficiency rescues lipotoxic cardiomyopathy [34].

CD36 is an integral membrane protein found on the surface of many cell types and binds many ligands including oxidized lipid proteins [35,36], long-chain fatty acids [37] and erythrocytes that are parasitized with the malaria parasite *Plasmodium falciparum* [38]. The rs3211938-G variant is nearly absent in Europeans and Asians and shows a signature of selection in African Americans and some African populations [39,40]. Additionally, rs3211938-G has been shown in previous studies to be associated with CD36 deficiency and with susceptibility to malaria, although this has not been confirmed in other studies [41,42].

ICAM1

The rs5030359 variant in *ICAM1*, is observed in this study to be associated with TC and LDL-C at conventional genome-wide significance. *ICAM1* encodes a cell surface glycoprotein that is typically expressed on endothelial cells and cells of the immune system [43]. However, rs5030359 maps to a gene-dense region (Figure 2b), so it cannot be excluded that there is another gene underlying the signal. The rs5030359 variant is ~800 kb downstream of a previously identified lipids signal within the *LDLR* region, but conditional analyses showed that the two loci are independent. Using fine-mapping in non-African populations to point to the most likely gene underlying the signal, is not possible as the SNP is very rare in Europeans, with a MAF of 0.002, and absent in our Hispanic and East Asian populations. Previously, common variants within *ICAM1* were found to be associated with soluble ICAM1 (sICAM1) concentrations in Europeans [44,45]. sICAM1 has been associated with several common diseases such as diabetes, heart disease, stroke, and malaria [46,47]. sICAM1 levels were associated with progression of carotid intima media thickness in young adults [48,49] and in asymptomatic dyslipidaemia subjects [50]. Additionally, sICAM1 levels were found to be higher in Europeans than in Africans [49].

Differences in signals within lipid loci in multiple ethnicities

We were able to use the dense SNP genotyping in loci on the IBC array to analyze and compare lipid-associated loci, particularly between African Americans and Europeans. Our analyses showed multiple examples of signals that were associated with lipid levels in one ethnicity but not another (Table 1).

First, some of the strongest associated SNPs in one ethnicity may be rare or absent in other ethnicities. This is a well-established phenomenon, e.g., truncation mutations in *PCSK9* that are of low frequency in African Americans and absent in individuals of European origin, that result in a robust reduction in LDL-C levels and coronary heart disease risk [51,52]. In this study we find that the majority of the observed discrepancies across ethnicities in association of SNPs with lipid traits can be attributed to differences in allele frequency. For example, rs3211938 in *CD36* is much more highly associated with HDL-C in African Americans ($p = 1.8 \times 10^{-11}$) than in Europeans ($p = 0.08$) with a large discrepancy in RAFs (7% vs. 0.2%).

In other loci, the strongest associated polymorphisms varied across populations, for example in the *BUD13/ZN259/APOA5* region (Table S3, Figure S2). In theory these regions could be excellent candidates for fine-mapping, but our efforts and association results could not narrow down the loci. When conducting meta-analyses across multiple ethnicities we observed that the stronger p-value association typically tracked with the higher heterogeneity I^2 values (Figure S3). This high I^2 suggests high heterogeneity, but it could also be the effect of low sample

sizes of the combined cohorts (especially for Hispanics and East Asians).

One limitation of this study is the sample size available particularly the Hispanic and the East Asian available samples and this obviously limited our ability to find new signals in these populations and to replicate many previously established lipid signals. Also, not all previously described signals for lipids were present on the IBC array, as the array was designed to densely cover genes regions, rather than the whole genome. However, using this approach we did find signals for lipids that remained uncovered using the genome-wide association approach, as both rs5030359 within *ICAM1* and rs3211938 within *CD36* were not present on conventional genome-wide arrays.

In conclusion, we performed dense genotyping of ~2,000 candidate genes in 7,657 African Americans, 1,315 Hispanics and 841 East Asians using IBC 50K SNP genotyping array and we found and confirmed two novel signals for lipids by replication in 7,000 African Americans. Additionally we evaluated the effect of SNPs established in European populations on lipid levels in multi-ethnic populations and show that most known lipid association signals span across ethnicities. However, differences between populations, especially differences in allele frequency, can be leveraged to identify novel signals.

Supporting Information

Figure S1 Quartiles based on the number of weighted risk alleles and lipid level distribution.
(TIF)

Figure S2 Association results of the *BUD13/ZN259/APOA5* regions with TG in multiple ethnicities. Beta's of SNPs are shown in the *BUD13/ZN259/APOA5* region from association results in each ethnicity separately. $-\log P$ -value and I^2 are from multi-ethnic meta-analyses. Squares mark the strongest association signals per ethnicity. The three independent signals in Europeans are depicted in green, the top signal in African Americans is shown in blue and Hispanics and East Asian meta-analyses results are in red and yellow respectively.
(PDF)

Figure S3 Correlation between $-\log P$ -value and I^2 in the *BUD13/ZN259/APOA5* region for TG association results.
(TIF)

Supporting Information S1 Supplementary acknowledgments.
(DOCX)

Table S1 Characteristics of studies contributing to the multi-ethnic IBC lipids meta-analyses.
(XLSX)

Table S2 Summary statistics for covariates in participating studies providing individual-level data.
(XLSX)

Table S3 Association results for known lipids loci in Europeans, African-Americans, Hispanics and East Asians.
(XLSX)

Table S4 Loci associated with lipid traits in individuals of African American, Hispanic and East Asian origin (model 2, model 3).
(XLSX)

Author Contributions

Conceived and designed the experiments: CCE FWA FD JGW BJK. Performed the experiments: VT EPAVI MBL BAC FC LRY MKW YRL

BF CMB YDIC WMC LAC YD DD MF MG NG FG TBH CK SBK AL KL YL KM ABN NCOM JO IP WP RS PJS KAV SY SSA DMB BP APR SSR JIR MMS MYT IBB RAH SK MAN DJR HH SS BJK SGB HAT SER SR MC MKE JKF WTG IH VJH MFK MIK XW ABZ.

Analyzed the data: CCE YG VT BJK. Contributed reagents/materials/analysis tools: FD FWA FD JGW BJK. Wrote the paper: CCE VT FD FWA FD JGW BJK.

References

- Arsenault BJ, Boekholdt SM, Kastelein JJ (2011) Lipid parameters for measuring risk of cardiovascular disease. *Nat Rev Cardiol* 8: 197–206.
- Heller DA, de Faire U, Pedersen NL, Dahlen G, McClearn GE (1993) Genetic and environmental influences on serum lipid levels in twins. *N Engl J Med* 328: 1150–1156.
- Teslovich TM, Musunuru K, Smith AV, Edmondson AC, Stylianou IM, et al. (2010) Biological, clinical and population relevance of 95 loci for blood lipids. *Nature* 466: 707–713.
- Ford ES, Giles WH, Dietz WH (2002) Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA* 287: 356–359.
- Park YW, Zhu S, Palaniappan L, Heshka S, Carnethon MR, et al. (2003) The metabolic syndrome: prevalence and associated risk factor findings in the US population from the Third National Health and Nutrition Examination Survey, 1988–1994. *Arch Intern Med* 163: 427–436.
- Lanktree MB, Anand SS, Yusuf S, Hegele RA (2009) Replication of genetic associations with plasma lipoprotein traits in a multiethnic sample. *J Lipid Res* 50: 1487–1496.
- Lettre G, Palmer CD, Young T, Ejebe KG, Allayee H, et al. (2011) Genome-wide association study of coronary heart disease and its risk factors in 8,090 African Americans: the NHLBI CARE Project. *PLoS Genet* 7: e1001300.
- Keating BJ, Tischfield S, Murray SS, Bhargava T, Price TS, et al. (2008) Concept, design and implementation of a cardiovascular gene-centric 50 k SNP array for large-scale genomic association studies. *PLoS One* 3: e3583.
- Consortium IKC (2011) Large-scale gene-centric analysis identifies novel variants for coronary artery disease. *PLoS Genet* 7: e1002260.
- Fox ER, Young JH, Li Y, Dreisbach AW, Keating BJ, et al. (2011) Association of genetic variation with systolic and diastolic blood pressure among African Americans: the Candidate Gene Association Resource study. *Hum Mol Genet* 20: 2273–2284.
- Clarke R, Peden JF, Hopewell JC, Kyriakou T, Goel A, et al. (2009) Genetic variants associated with Lp(a) lipoprotein level and coronary disease. *N Engl J Med* 361: 2518–2528.
- Saxena R, Elbers CC, Guo Y, Peter I, Gaunt TR, et al. (2012) Large-scale gene-centric meta-analysis across 39 studies identifies type 2 diabetes loci. *Am J Hum Genet* 90: 410–425.
- Asselbergs FW, Guo Y, van Iperen EP, Sivapalaratnam S, Tragante V, et al. (2012) Large-Scale Gene-Centric Meta-analysis across 32 Studies Identifies Multiple Lipid Loci. *Am J Hum Genet*.
- Friedewald WT, Levy RI, Fredrickson DS (1972) Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 18: 499–502.
- Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, et al. (2007) PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 81: 559–575.
- Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, et al. (2006) Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet* 38: 904–909.
- Willer CJ, Li Y, Abecasis GR (2010) METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 26: 2190–2191.
- Musunuru K, Lettre G, Young T, Farlow DN, Pirruccello JP, et al. (2010) Candidate gene association resource (CARE): design, methods, and proof of concept. *Circ Cardiovasc Genet* 3: 267–275.
- Higgins JP, Thompson SG (2002) Quantifying heterogeneity in a meta-analysis. *Stat Med* 21: 1539–1558.
- Saxena R, Elbers CC, Guo Y, Peter I, Gaunt TR, et al. (2012) Large-Scale Gene-Centric Meta-Analysis across 39 studies Identifies Type 2 Diabetes Loci. *Am J Hum Genet*.
- Asselbergs FWG, Sivapalaratnam S, van Iperen EP, de O VT, Lanktree MB, et al. (2012) Twenty-three unreported genetic associations with lipid phenotypes: a dense gene-centric meta-analysis in 66,240 individuals across 32 studies. submitted to *AJHG*.
- Cornelis MC, Qi L, Zhang C, Kraft P, Manson J, et al. (2009) Joint effects of common genetic variants on the risk for type 2 diabetes in U.S. men and women of European ancestry. *Ann Intern Med* 150: 541–550.
- Waters KM, Stram DO, Hassanein MT, Le Marchand L, Wilkens LR, et al. (2010) Consistent association of type 2 diabetes risk variants found in Europeans in diverse racial and ethnic groups. *PLoS Genet* 6.
- Pulit SL, Voight BF, de Bakker PI (2010) Multiethnic genetic association studies improve power for locus discovery. *PLoS One* 5: e12600.
- Love-Gregory L, Sherva R, Sun L, Wasson J, Schappe T, et al. (2008) Variants in the CD36 gene associate with the metabolic syndrome and high-density lipoprotein cholesterol. *Hum Mol Genet* 17: 1695–1704.
- Morii T, Ohno Y, Kato N, Hirose H, Kawabe H, et al. (2009) CD36 single nucleotide polymorphism is associated with variation in low-density lipoprotein-cholesterol in young Japanese men. *Biomarkers* 14: 207–212.
- Goyenechea E, Collins LJ, Parra D, Liu G, Snieder H, et al. (2008) CD36 gene promoter polymorphisms are associated with low density lipoprotein-cholesterol in normal twins and after a low-calorie diet in obese subjects. *Twin Res Hum Genet* 11: 621–628.
- Gaillard D, Laugerette F, Darcel N, El-Yassimi A, Passilly-Degrace P, et al. (2008) The gustatory pathway is involved in CD36-mediated orosensory perception of long-chain fatty acids in the mouse. *FASEB J* 22: 1458–1468.
- Laugerette F, Passilly-Degrace P, Patris B, Niot I, Febbraio M, et al. (2005) CD36 involvement in orosensory detection of dietary lipids, spontaneous fat preference, and digestive secretions. *J Clin Invest* 115: 3177–3184.
- Martin C, Passilly-Degrace P, Gaillard D, Merlin JF, Chevrot M, et al. (2011) The lipid-sensor candidates CD36 and GPR120 are differentially regulated by dietary lipids in mouse taste buds: impact on spontaneous fat preference. *PLoS One* 6: e24014.
- Bokor S, Legry V, Meirhaeghe A, Ruiz JR, Mauro B, et al. (2010) Single-nucleotide polymorphism of CD36 locus and obesity in European adolescents. *Obesity (Silver Spring)* 18: 1398–1403.
- Choquet H, Labrune Y, De Graeve F, Hinney A, Hebebrand J, et al. (2011) Lack of association of CD36 SNPs with early onset obesity: a meta-analysis in 9,973 European subjects. *Obesity (Silver Spring)* 19: 833–839.
- Drover VA, Ajmal M, Nassir F, Davidson NO, Nauli AM, et al. (2005) CD36 deficiency impairs intestinal lipid secretion and clearance of chylomicrons from the blood. *J Clin Invest* 115: 1290–1297.
- Yang J, Sambandam N, Han X, Gross RW, Courtois M, et al. (2007) CD36 deficiency rescues lipotoxic cardiomyopathy. *Circ Res* 100: 1208–1217.
- Endemann G, Stanton LW, Madden KS, Bryant CM, White RT, et al. (1993) CD36 is a receptor for oxidized low density lipoprotein. *J Biol Chem* 268: 11811–11816.
- Calvo D, Gomez-Coronado D, Suarez Y, Lasuncion MA, Vega MA (1998) Human CD36 is a high affinity receptor for the native lipoproteins HDL, LDL, and VLDL. *J Lipid Res* 39: 777–788.
- Abumrad NA, el-Maghrabi MR, Amri EZ, Lopez E, Grimaldi PA (1993) Cloning of a rat adipocyte membrane protein implicated in binding or transport of long-chain fatty acids that is induced during preadipocyte differentiation. Homology with human CD36. *J Biol Chem* 268: 17665–17668.
- Oquendo P, Hundt E, Lawler J, Seed B (1989) CD36 directly mediates cytoadherence of *Plasmodium falciparum* parasitized erythrocytes. *Cell* 58: 95–101.
- Bhatia G, Patterson N, Pasaniuc B, Zaitlen N, Genovese G, et al. (2011) Genome-wide comparison of African-ancestry populations from CARE and other cohorts reveals signals of natural selection. *Am J Hum Genet* 89: 368–381.
- Ayodo G, Price AL, Keinan A, Ajwang A, Otieno MF, et al. (2007) Combining evidence of natural selection with association analysis increases power to detect malaria-resistance variants. *Am J Hum Genet* 81: 234–242.
- Aitman TJ, Cooper LD, Norsworthy PJ, Wahid FN, Gray JK, et al. (2000) Malaria susceptibility and CD36 mutation. *Nature* 405: 1015–1016.
- Fry AE, Ghansa A, Small KS, Palma A, Auburn S, et al. (2009) Positive selection of a CD36 nonsense variant in sub-Saharan Africa, but no association with severe malaria phenotypes. *Hum Mol Genet* 18: 2683–2692.
- van de Stolpe A, van der Saag PT (1996) Intercellular adhesion molecule-1. *J Mol Med (Berl)* 74: 13–33.
- Pare G, Chasman DI, Kellogg M, Zec RY, Rifai N, et al. (2008) Novel association of ABO histo-blood group antigen with soluble ICAM-1: results of a genome-wide association study of 6,578 women. *PLoS Genet* 4: e1000118.
- Bielinski SJ, Reiner AP, Nickerson D, Carlson C, Bailey KR, et al. (2011) Polymorphisms in the ICAM1 gene predict circulating soluble intercellular adhesion molecule-1 (sICAM-1). *Atherosclerosis* 216: 390–394.
- Ridker PM, Hennekens CH, Roitman-Johnson B, Stampfer MJ, Allen J (1998) Plasma concentration of soluble intercellular adhesion molecule 1 and risks of future myocardial infarction in apparently healthy men. *Lancet* 351: 88–92.
- Song Y, Manson JE, Tinker L, Rifai N, Cook NR, et al. (2007) Circulating levels of endothelial adhesion molecules and risk of diabetes in an ethnically diverse cohort of women. *Diabetes* 56: 1898–1904.
- Gross MD, Bielinski SJ, Suarez-Lopez JR, Reiner AP, Bailey K, et al. (2012) Circulating soluble intercellular adhesion molecule 1 and subclinical atherosclerosis: the Coronary Artery Risk Development in Young Adults Study. *Clin Chem* 58: 411–420.
- Nguyen QM, Srinivasan SR, Xu JH, Chen W, Berenson GS (2010) Distribution and cardiovascular risk correlates of plasma soluble intercellular adhesion molecule-1 levels in asymptomatic young adults from a biracial community: the Bogalusa Heart Study. *Ann Epidemiol* 20: 53–59.
- Karasek D, Vaverkova H, Frysak Z, Halenka M, Jackuliakova D, et al. (2011) Soluble intercellular cell adhesion molecule-1 and vascular cell adhesion molecule-1 in asymptomatic dyslipidemic subjects. *Int Angiol* 30: 441–450.

51. Cohen J, Pertsemlidis A, Kotowski IK, Graham R, Garcia CK, et al. (2005) Low LDL cholesterol in individuals of African descent resulting from frequent nonsense mutations in PCSK9. *Nat Genet* 37: 161–165.
52. Cohen JC, Boerwinkle E, Mosley TH Jr, Hobbs HH (2006) Sequence variations in PCSK9, low LDL, and protection against coronary heart disease. *N Engl J Med* 354: 1264–1272.