

Gene lineages and eastern North American palaeodrainage basins: phylogeography and speciation in salamanders of the *Eurycea bislineata* species complex

KENNETH H. KOZAK, RUSSELL A. BLAINE and ALLAN LARSON

Department of Biology, Washington University, Saint Louis, MO 63130, USA

Abstract

Contemporary North American drainage basins are composites of formerly isolated drainages, suggesting that fragmentation and fusion of palaeodrainage systems may have been an important factor generating current patterns of genetic and species diversity in stream-associated organisms. Here, we combine traditional molecular-phylogenetic, multiple-regression, nested clade, and molecular-demographic analyses to investigate the relationship between phylogeographic variation and the hydrogeological history of eastern North American drainage basins in semiaquatic plethodontid salamanders of the *Eurycea bislineata* species complex. Four hundred forty-two sequences representing 1108 aligned bases from the mitochondrial genome are reported for the five formally recognized species of the *E. bislineata* complex and three outgroup taxa. Within the ingroup, 270 haplotypes are recovered from 144 sampling locations. Geographic patterns of mtDNA-haplotype coalescence identify 13 putatively independent population-level lineages, suggesting that the current taxonomy of the group underestimates species-level diversity. Spatial and temporal patterns of phylogeographic divergence are strongly associated with historical rather than modern drainage connections, indicating that shifts in major drainage patterns played a pivotal role in the allopatric fragmentation of populations and build-up of lineage diversity in these stream-associated salamanders. More generally, our molecular genetic results corroborate geological and faunistic evidence suggesting that palaeodrainage connections altered by glacial advances and headwater erosion occurring between the mid-Miocene and Pleistocene epochs explain regional patterns of biodiversity in eastern North American streams.

Keywords: Amphibia, Appalachian Mountains, biogeography, drainage history, Plethodontidae, stream capture, vicariance

Received 28 June 2005; revision accepted 24 August 2005

Introduction

A continuing challenge in evolutionary biology is to understand population processes that lead to the formation of new evolutionary lineages. When combined with geographic patterns of genetic variation predicted independently from geological (Durand *et al.* 1999; Waters *et al.* 2001; Johnson 2002) and palaeoclimatic sources (Masta 2000; Knowles 2001; Carstens *et al.* 2004), the genealogical history and spatial distributions of gene lineages provide a powerful tool for

understanding historical and contemporary factors that influence the genetic compositions of populations, and ultimately, speciation (Hewitt 2001; Templeton 2001).

In eastern North America, populations of stream-dwelling organisms have experienced dramatic geological changes in the major drainage patterns that structure their geographic patterns of gene exchange, setting the stage for divergence and speciation. Drainage patterns west of the Eastern Continental Divide (ECD) (Fig. 1A), a major physiographic feature that separates eastern-flowing drainages of the Atlantic Slope from those that ultimately drain into the Gulf of Mexico, were profoundly different from current ones (Fig. 1B). As a result of Pleistocene glacial advances, flow patterns of rivers were reversed to the south, resulting in fusion of formerly isolated drainage systems and the origin

Correspondence: Kenneth H. Kozak, Present address: Department of Ecology & Evolution, Stony Brook University, Stony Brook, NY 11794-5245, USA. Fax: 631-632-7626; E-mail: kozak@life.bio.sunysb.edu




Fig. 1 Major Pliocene (A) and contemporary (B) drainage basins in eastern North America. Tributaries of the present-day Ohio River were organized into three major drainage basins separated by dispersal barriers: the Laurentian, Teays, and Old-Ohio (Burr & Page 1986; Hocutt *et al.* 1986; Mayden 1988). The Teays River basin included the Kanawha, Licking, and Kentucky drainage systems; the Old-Ohio system included the Wabash and Green drainage systems. Glacial advances during the Pleistocene diverted tributaries of the Teays system southward to form the modern Ohio River system. Portions of the Teays were diverted into the Upper Wabash system of the Old-Ohio basin. In addition, the formerly north-flowing Old Upper Allegheny River of the Laurentian system reversed to flow south and was integrated into the Teays River basin. Similarly, the present-day Tennessee River basin is a hypothesized composite of formerly separate palaeodrainage basins that had independent southward courses to the Gulf of Mexico (Thornbury 1965; Mayden 1988; but see Starnes & Etnier 1986). Headwaters of Atlantic Slope drainages are also composites of formerly isolated rivers. Geological evidence and freshwater-fish zoogeographic patterns indicate that Pleistocene stream captures diverted the headwaters of the New and Upper Tennessee systems into the James–Roanoke and Santee drainages of the Atlantic Slope, respectively (Ross 1969, 1971; Hocutt *et al.* 1986; Starnes & Etnier 1986).

of the modern Ohio River basin. Likewise, the contemporary Tennessee River basin is a hypothesized composite of two major drainage basins that had independent southward courses to the Gulf of Mexico prior to the close of the Pliocene epoch.

Similarly, as a result of stream capture, headwaters of Atlantic Slope drainages are composites of formerly separate drainages. Stream capture occurs when differential erosion at the headwaters of a drainage divide transfers part of a stream from one drainage system to another, facilitating biotic exchange between previously isolated drainage basins (Burr & Page 1986). This process permits dispersal between formerly isolated drainages and ultimately may lead to speciation if the new drainage divide continues to impede gene exchange among drainage systems. Both hydrogeology and zoogeographic patterns in freshwater fishes indicate that stream captures diverted the headwaters of the New

and Upper Tennessee systems into the James–Roanoke and Santee drainages of the Atlantic Slope, respectively (Ross 1969, 1971; Hocutt *et al.* 1986; Starnes & Etnier 1986).

The complex history of drainage-basin fragmentation and fusion in eastern North America generates a priori hypotheses for the phylogeographic structuring of populations throughout their evolutionary histories. First, the hypothesis that the Upper and Lower portions of the Tennessee River had independent outlets to the Gulf of Mexico predicts that haplotypes sampled from the present-day Tennessee River will not form a clade. Instead, an ancient pre-Pleistocene phylogeographic break is expected among haplotypes sampled from the Upper and Lower Tennessee River basins. Haplotypes sampled from the Lower Tennessee River are expected to group with those from the Cumberland drainage, while those from the Upper Tennessee River are predicted to group with those from Gulf drainage systems. Second, the

hypothesis that stream capture played an important role in the movement of populations across the ECD predicts that haplotypes in translocated rivers should retain phylogenetic affinity with those of their ancestral drainage. Specifically, haplotypes sampled from the James–Roanoke and Santee drainages on the Atlantic Slope should be derived from ancestral haplotypes distributed in the New and Upper Tennessee systems, respectively. As a corollary, given the hypothesized Pleistocene timing of these stream captures (Ross 1969, 1971), genetic signatures of range expansion across the ECD are also predicted. Integration of traditional molecular-phylogenetic methods and population-genetic approaches based on coalescent theory permits rigorous tests of these phylogeographic predictions.

Recent applications of mitochondrial DNA haplotype phylogenies in North American freshwater fishes demonstrate the importance of historical drainage connections in shaping geographic patterns of genetic variation (Strange & Burr 1997; Kreiser *et al.* 2001; Near *et al.* 2001; Hardy *et al.* 2002; Berendzen *et al.* 2003). However, there have been surprisingly few tests of drainage evolution as a determinant of biotic diversity in other codistributed stream-associated organisms (Routman *et al.* 1994; Voss *et al.* 1995). Here, we examine the relationship between the geological history of eastern North American drainage basins and geographic patterns of mtDNA sequence variation in semiaquatic plethodontid salamanders of the *Eurycea bislineata* (two-lined salamanders) species complex.

Populations of two-lined salamanders are common inhabitants of eastern North American streams. Several aspects of their life history suggest that gene flow occurs primarily through stream corridors and therefore predict phylogeographic structuring that reflects the hydrogeological history of drainage basins. Both larvae and adults show seasonal upstream and downstream movements (Bruce 1986). In some populations, adults engage in seasonal terrestrial movements; however, they are highly philopatric and return to their natal streams to overwinter and to breed (MacCulloch & Bider 1975). Current taxonomy of the group recognizes five species distributed across the Interior Lowlands, Appalachian Highlands, Piedmont, and Coastal Plain of eastern North America (Sever *et al.* 1976; Jacobs 1987; Kozak & Lannoo 2005), making it an ideal study system for investigating the biological significance of geological interactions among the region's major drainage basins. Furthermore, despite overall morphological stasis, the *E. bislineata* complex shows evidence of ancient geographic genetic fragmentation dating to as early as the Miocene (Jacobs 1987).

If gene exchange among populations is influenced by the interconnectivity of streams, phylogenetic relationships among gene lineages and processes generating population structure within those lineages should reflect historical connections among drainage systems. In this study, we use a combination of traditional phylogenetic, matrix-

correspondence, nested clade, and population-demographic analyses to test a priori genetic predictions of historical fragmentation and population structure derived from eastern North America's dynamic hydrologic history.

Materials and methods

Population sampling, DNA sequencing, and alignment

We obtained mtDNA sequence data from 438 salamanders sampled from 144 localities across the range of the *Eurycea bislineata* complex (Fig. 2). Sample sizes range from 1 to 10 individuals per location (mean = 3). Samples of *Eurycea guttolineata*, *Eurycea longicauda*, and *Eurycea quadridigitata* were used as outgroups for phylogenetic analyses. Sample sizes and sampling locations are listed in Table S1 (Supplementary material).

DNA extraction, amplification, and sequencing were performed as in Kozak *et al.* (2005). Amplification and sequencing of the mitochondrial ND2 gene and the adjacent tRNA^{Trp} gene were conducted using primers L4437 (5'-AAGCTTTCGGGCCCATACC-3') and H5934 (5'-AGRGTGCCAATGCTTTGTGRTT-3') and primers L4882 (5'-TGACAAAACTAGCC-3') and H5617a (5'-AAAATRTCTGRGTTGCATTAG-3') were used as internal sequencing primers (Macey *et al.* 1997). The protein-coding ND2 gene was translated to amino acids using MACCLADE (Maddison & Maddison 1992) to check for premature stop codons. The tRNA sequence was aligned manually based on models of secondary structure (Macey & Verma 1997). Alignment was straightforward and unambiguous. Length variation consisted of a single base-pair insertion/deletion in the D-loop of the tRNA^{Trp} gene in some individuals.

Phylogenetic analyses

Hierarchical phylogenetic relationships among haplotypes were estimated using Bayesian and parsimony optimality criteria. MR MODELTEST 1.1b was used to select the model of nucleotide substitution that best fit the mtDNA sequence data (www.ebc.uu.se/systzoo/staff/nylander.html). Bayesian phylogenetic analyses were then implemented in MRBAYES 3.01 (Huelsenbeck & Ronquist 2001) using a GTR + I + Γ model of evolution. Four incrementally heated Markov chains were run for 5×10^6 generations, sampling every 5000 generations for a total of 1000 samples. Flat priors were used for all substitution-parameter estimates, and random trees were used to begin each Markov chain. To ensure that the Markov chains reached a stable equilibrium, ln-likelihood values for sampling points were plotted against generation time. Equilibrium sample points were used to generate a 50% majority-rule consensus tree, where the percentage of samples that recover a particular node




Fig. 2 Geographic distribution of *Eurycea bislineata*-complex sampling locations included in this study. Open (A–E) and shaded (F–M) symbols designate the major clades (northern and southern, respectively) to which haplotypes sampled from that population belong. The mixed circle for location 136 denotes sympatric occurrence of haplotypes from the northern and southern clades. Symbols delimit the geographic distributions of the major haplotype clades denoted in Fig. 4. Lineage A: Potomac and Lower Chesapeake drainages; Lineage B: Potomac, Kanawha, Lower Chesapeake, Monongahela, Lower Hudson, Upper Susquehanna, Oswego, Lower Connecticut, Saco, and Upper Connecticut drainages; Lineage C: Cumberland and Lower Tennessee drainages; Lineage D: Green, Wabash and Lower-Ohio, Kentucky, Licking and Little Miami drainages; Lineage E: New, Greenbriar, Middle-Ohio, Roanoke, James, Santee, Shenandoah, and Upper Tennessee (Holston) drainages; Lineage F: Upper Tennessee (Emory) and South Fork Cumberland drainages; Lineage G: Upper Tennessee (Little Tennessee) drainage; Lineage H: Cahaba drainage; Lineage I: Santee and Roanoke drainages; Lineage J: Upper Tennessee (Nolichucky and French Broad), Santee, Enoree, Cape Fear, and Neuse drainages; Lineage K: Pearl drainage; Lineage L: Pascagoula, Middle-Tennessee-Elk, Apalachicola, Florida-Panhandle Coastal, Ochlockonee, Escambia, Alabama, Altamaha, and Savannah drainages; Lineage M: Upper Tennessee (Hiwassee, Little Tennessee, Tuckasegee), Middle Tennessee-Elk, Coosa-Tallapoosa, Savannah, and Santee drainages.

represents the posterior probability of that clade. To verify that analyses were not trapped on local optima, five replicate searches were conducted. The independent analyses were considered to have converged on the optimal joint posterior distribution if similar ln-likelihood scores and parameter estimates were achieved (Huelsenbeck & Bollback 2001).

PAUP* 4.0b10 (Swofford 2002) was used to reconstruct phylogenetic relationships among haplotypes using maximum parsimony and 100 heuristic searches with random addition of sequences and tree-bisection–reconnection (TBR) branch swapping. To facilitate computational efficiency, parsimony analyses were conducted with two representative haplotypes from each of the major haplotype clades resolved in the Bayesian phylogenetic analyses (see Results). To assess support for individual nodes, 100 nonparametric bootstrap pseudoreplicates with 10 random taxon-addition-sequence replicates per bootstrap pseudoreplicate were performed. Decay indices (Bremer 1994) were calculated for each node using TREEROT version 2a (Sorenson 1999).

To delineate boundaries of putatively independent evolutionary lineages, we employed the tree-based method of Wiens & Penkrot (2002), which uses geographic patterns of coalescence among DNA haplotypes to test for gene exchange among populations of a focal species and one or more closely related species. Briefly, discordance between the phylogenetic relatedness of haplotypes and their geographic proximity provides evidence for gene flow among populations of the focal species (Slatkin & Maddison 1989). Deep phylogenetic splits among haplotype clades that replace each other geographically delimit putatively independent evolutionary lineages concealed by previous taxonomy. Incomplete sorting of ancestral polymorphisms following lineage splitting may also lead to discordance between gene trees and geography, in which case this procedure would lead to a conservative estimate of the number of putatively independent lineages.

To obtain estimates of divergence times, we calculated the mean pairwise sequence divergence between lineages and employed an expected rate of 1.3% sequence divergence per million years calibrated from homologous sequences in salamandrid salamanders (Weisrock *et al.* 2001) and bufonid frogs (Macey *et al.* 1998). Mean uncorrected *p*-distances and Tamura–Nei corrected distances (Tamura & Nei 1993) among major lineages were calculated using MEGA version 2 (Kumar *et al.* 2001). To test for rate heterogeneity among lineages, we performed relative-rate tests using the program RRTREE (Robinson-Rechavi & Huchon 2000).

Drainage-basin history and phylogeographic divergence

The hypothesis that interconnectivity of drainage systems influences patterns of phylogeographic structure predicts that salamander populations residing in different drainage

basins should display greater genetic divergence than those distributed within the same one. Given the ancient evolutionary history of the *E. bislineata* complex suggested by allozyme-based genetic distances (Jacobs 1987) and mtDNA sequence divergence, phylogeographic patterns may reflect historical as well as contemporary drainage connections. We used partial matrix correlation tests (PMCT) to evaluate congruence between patterns of phylogenetic divergence and drainage-basin connections (Smouse *et al.* 1986; Legendre *et al.* 1994; Thorpe *et al.* 1996).

PMCTs measure the correlation between a response matrix and multiple independent matrices simultaneously (Fig. 3). Simultaneous consideration of multiple matrices is critical because putative causes of phylogeographic divergence may be either correlated or additive in their effects. For example, as the geographic distance separating populations increases, so does the possibility that the populations being compared are distributed among different drainages (i.e. the geographic distance separating populations and the spatial arrangement of drainage basins are not independent). In addition, under isolation by distance the geographic distance separating populations is a significant predictor of their genetic divergence (Slatkin 1993). Therefore, to test the hypothesis that patterns of phylogenetic divergence are influenced by the interconnectivity of stream drainages, it is necessary to determine whether the degree of genetic divergence among populations residing in different drainage basins is greater than expected from geographic distance alone. Finally, drainage connections from different geological epochs are not independent because while some drainage connections were modified, others remained static, making it necessary to estimate the contributions of historical and contemporary drainage patterns relative to each other.

A phylogenetic-divergence matrix measuring the patristic distances separating haplotypes along the Bayesian consensus phylogram was designated the response matrix; the independent matrices included a matrix of great-circle distances between all pairs of sampling locations to control for the spatial effects mentioned above, binary design matrices corresponding to Pliocene, Pleistocene, and contemporary drainage connections, and the current species-level taxonomy. For the drainage-connection matrices, each element was set to zero if the haplotypes being compared were in the same drainage basin for the time period under consideration, or one if they were in different drainage basins. Similarly, haplotypes sampled from the same or different formally recognized species were assigned a value of zero or one, respectively. Each matrix was normalized (zero mean, unit variance). Because elements within individual matrices are not independent, significance of the partial regression coefficients was assessed by randomly permuting the order of values in the phylogenetic-divergence matrix 9999 times using the software package PERMUTE! (Casgrain 2001). If




Fig. 3 Hypothetical example outlining the basic procedure used to conduct partial matrix correlation tests. A map of haplotype sampling locations is converted to matrices corresponding to the pairwise geographic distance separating sampling locations and drainage connections from different geological time periods. For the drainage-connection matrices, each element was set to zero if the haplotype sampling locations being compared were in the same drainage basin for the time period under consideration, or one if they were in different drainage basins. A partial correlation between each matrix and the haplotype patristic distance matrix is calculated and its statistical significance is evaluated via permutation of the latter matrix. Partial correlations are proportional to the amount of variation that variable explains in the haplotype phylogeny after controlling for its nonindependence with the other independent matrices considered.

fewer than 5% of the randomizations had partial regression coefficients greater than, or equal to those of the real data set, the results were considered significant. Figure 3 illustrates a hypothetical example outlining the basic procedure used to conduct the PMCTs.

Molecular demographic and nested clade analyses

A potential limitation of using contemporary distributions of lineages to assess the geographic context of lineage splitting is that ranges may shift over time (Losos & Glor 2003). To assess the geographic stability of lineages, we used Tajima's *D*-test (Tajima 1989) as implemented in ARLEQUIN version 2.0 (Schneider *et al.* 2000) to test for population-genetic equilibrium within major population-level lineages. Assuming neutrality of molecular variation, this test can be used to detect increases in population size that accompany recent range movements. To test whether patterns of mtDNA variation are consistent with the hypothesis of neutrality, we employed the MK test (McDonald & Kreitman 1991) in DNASP (Rozas *et al.* 2003), using haplotypic variation within and among the major lineages resolved in the hierarchical phylogenetic analyses to compare the ratio of nonsynony-

mous and synonymous substitutions within and among species, respectively,

The hypothesis that dispersal across the ECD was facilitated by headwater stream capture predicts (i) genetic evidence of range expansion across the ECD, and (ii) haplotypes distributed in James–Roanoke and Santee drainages of the Atlantic slope should be derived from those in the New and Upper Tennessee drainages, respectively. These predictions are ideally suited for testing with NCA, which uses the geographic distributions of ancestral (interior) haplotypes relative to younger (tip) ones to draw inferences about processes that have shaped spatial patterns of genetic variation. We used statistical parsimony as implemented in TCS version 1.18 (Clement *et al.* 2000) to link haplotypes into a minimum-connecting network depicting their genealogical relationships. The procedures of Templeton *et al.* (1987) and Templeton & Sing (1993) were then used to impose a hierarchical nesting structure on haplotype networks. In cases where it was possible to use a maximum-parsimony tree to link haplotype networks that could not be connected at the 95% level of confidence using statistical parsimony, we grouped individual networks as sister clades at equal nesting levels (Templeton *et al.* 1987) and

used outgroup rooting to determine the tip/interior status of the nested groups. GEODIS version 2.0 (Posada *et al.* 2000) was used to calculate (i) the clade distance D_c , which measures the average distance of haplotypes in a group from its geographic centre, (ii) the nested clade distance D_n , which measures how far a haplotype group is from the geographic centre of other groups with which it is nested, and (iii) the average D_c and D_n separating interior and tip groupings of haplotypes. To test whether the geographic distributions of haplotypes were more widespread or restricted than expected by chance, we used a categorical permutation contingency analysis. The most likely historical and recurrent processes responsible for statistically significant patterns of phylogeographic variation were inferred using the revised inference key (Templeton 2004).

Results

Hierarchical phylogenetic analyses

Four hundred forty-two sequences representing 1108 aligned bases are reported for the five formally recognized species of the *Eurycea bislineata* complex and three outgroup taxa. Within the ingroup, 270 haplotypes are recovered from the 144 sampling locations (Table S2). Absence of premature stop codons in the protein-coding ND2 gene region, functional stability of the tRNA^{Trp} gene, and strong bias against guanine on the light strand indicate that the DNA sequences are from the mitochondrial genome and not nuclear-integrated copies of mitochondrial genes (Zhang & Hewitt 1996). Likelihood-ratio tests favour the GTR + I + Γ model of nucleotide substitution. Including the three outgroup taxa, the haplotype data set contains 632 variable sites of which 509 are parsimony informative (474 within the ingroup). Bayesian phylogenetic analysis using the GTR + I + Γ model of evolution produces a 50% majority-rule consensus tree with a mean ln-likelihood of -14382.84 (SD = 22.93) following a 'burn in' of 500 000 generations (Fig. 4). Parsimony analysis results in a single tree of 1301 steps. Because both analyses produced highly congruent estimates of phylogenetic relationship among the major haplotype clades, only the Bayesian consensus phylogram is presented, with posterior probabilities and nonparametric bootstrap values from the parsimony analyses included for shared branches.

Eurycea bislineata-complex haplotypes form a well-supported monophyletic group to the exclusion of *Eurycea guttolineata*, *Eurycea longicauda*, and *Eurycea quadridigitata*. Tree-based lineage delimitation identifies 13 phylogenetically and geographically distinct haplotype groups diagnosing putatively independent evolutionary lineages (Fig. 4). In some cases it is possible to delimit additional such haplotype groupings; however, we have conservatively focused on the most inclusive groups of haplotypes that are

concordant with geography to delimit population-level lineages. A high ratio of between vs. within-clade molecular divergence indicates a deep history of geographic genetic fragmentation among these lineages: Tamura–Nei distances among lineages average 13.5% (uncorrected average = 11.5%), whereas the mean within-lineage divergence is 1.5% (Table 1). Based on these estimates, the vast majority of divergence events between inferred sister lineages are estimated to have occurred during the late Pliocene to mid-Miocene (~4–11 Myr), with a single lineage-splitting event dating to the Pleistocene (A vs. B; ~1.5 Myr). Relative-rate tests suggest that lineage-specific rate heterogeneity is not prevalent in the haplotype data set; only a single lineage (lineage K, $P < 0.02$) has accumulated nucleotide substitution at a significantly different rate.

The deepest phylogenetic divergence in the complex separates haplotypes into strongly supported northern and southern clades. Northern clade haplotypes are distributed primarily among drainages in the Lower Tennessee, former Old-Ohio, former Teays, mid-Atlantic slope, and those in the northeast that were inundated by Pleistocene glaciations. Southern clade haplotypes are distributed primarily in the Gulf Coastal Plain, southern Atlantic slope, and Upper Tennessee drainage systems. Haplotypes from the northern and southern clades occur sympatrically in the headwaters of the Santee drainage system in the southern Appalachian Highlands.

In general, major haplotype clades in the *E. bislineata* complex exhibit greater concordance with geography than they do with current taxonomy. Three clades correspond to formally recognized species (A + B: *E. bislineata*; G: *Eurycea junaluska*; H: *Eurycea aquatica*). However, haplotypes sampled from *Eurycea cirrigera* and *Eurycea wilderae* do not form monophyletic groups of haplotypes. Instead, they span the basal north–south split in the phylogeny, interdigitating with each other and the remaining taxa of the complex.

Drainage-basin history and phylogeographic divergence

Multiple regression of the five matrices on the haplotype phylogeny was highly significant (Table 2). The partial regression coefficients represent the slope for that variable when all other variables are held constant. Significant partial regression coefficients for geographic distance, Tertiary, Pleistocene, and contemporary drainage connections, and the current alpha taxonomy indicate that each of these factors explains phylogeographic structure in the haplotype phylogeny. Although phylogenetic divergence is strongly correlated with geography, historical drainage-basin connections explain more variation in the structure of the phylogeny than expected from the geographic distance separating haplotype sampling locations. Of the matrices considered, Pliocene drainage-basin connections explain the greatest amount of variation in the phylogeny. Interestingly,




Fig. 4 Bayesian 50% majority-rule consensus phylogram for the 270 *Eurycea bislineata*-complex haplotypes and three outgroup taxa. Posterior probabilities based on 4500 post burn-in trees (which had a mean ln-likelihood of -14 382.84; SD = 22.93) are shown above the branches; nonparametric bootstrap proportions, followed by decay indices for the parsimony analysis are below. Haplotypes in bold are included in the parsimony analyses. Clades for which nested clade analysis (NCA) was performed are shaded grey. For illustrative purposes, branch-support values are omitted from short near-terminal branches within major population lineages. Haplotypes are labelled with either the outgroup species name, or designation listed in Table S2. Locations from which haplotypes were sampled (Table S1; Fig. 2) are given in parentheses. Geographic distributions for major population lineages are depicted using symbols as in Fig. 2. (A) The northern clade containing major population lineages A-E; (B) The southern clade containing major population lineages F-M. Symbols for population lineages A-M follow Fig. 2.




Fig. 4 Continued

Table 1 Matrix of haplotypic divergences within and between 13 population-level lineages identified through phylogeographic analysis. Average Tamura–Nei-corrected distances between lineages are below the diagonal; average uncorrected proportion of sites differing between lineages are above the diagonal; mean Tamura–Nei distances among haplotypes within inferred population-level lineages are shown in boldface on the diagonal

	A	B	C	D	E	F	G	H	I	J	K	L	M
A	0.005	0.022	0.108	0.106	0.111	0.127	0.136	0.134	0.136	0.133	0.146	0.129	0.127
B	0.022	0.002	0.10	0.099	0.105	0.125	0.131	0.129	0.126	0.127	0.138	0.124	0.121
C	0.121	0.111	0.037	0.128	0.123	0.147	0.149	0.150	0.152	0.154	0.153	0.147	0.144
D	0.118	0.109	0.146	0.007	0.065	0.139	0.146	0.148	0.147	0.141	0.145	0.142	0.138
E	0.125	0.117	0.139	0.069	0.028	0.141	0.145	0.140	0.140	0.137	0.143	0.134	0.128
F	0.145	0.143	0.173	0.161	0.164	0.006	0.089	0.081	0.103	0.093	0.110	0.099	0.093
G	0.156	0.149	0.174	0.168	0.168	0.097	0.002	0.089	0.114	0.110	0.119	0.110	0.101
H	0.155	0.149	0.176	0.173	0.163	0.088	0.098	0.001	0.106	0.105	0.126	0.099	0.098
I	0.156	0.143	0.178	0.170	0.162	0.116	0.128	0.119	0.007	0.079	0.125	0.096	0.089
J	0.152	0.144	0.181	0.162	0.158	0.102	0.123	0.117	0.085	0.013	0.110	0.090	0.083
K	0.170	0.161	0.179	0.167	0.165	0.123	0.134	0.144	0.143	0.122	0.004	0.094	0.087
L	0.148	0.143	0.173	0.165	0.154	0.110	0.124	0.111	0.107	0.098	0.104	0.033	0.051
M	0.146	0.139	0.168	0.159	0.146	0.103	0.112	0.109	0.097	0.090	0.095	0.054	0.031

Table 2 Partial matrix-correlation tests of geographic distance, historical and contemporary drainage–basin connections, and taxonomic species categories on the haplotype phylogeny

Matrix	b^*	t -statistic†	P value‡
Geographic distance	0.157	33.27	0.0001
Pliocene drainage basins	0.475	95.55	0.0001
Pleistocene drainage basins	0.237	30.03	0.0001
Contemporary drainage basins	−0.203	−24.74	0.0001
Species-level taxonomy	0.084	19.07	0.0011

*Partial regression coefficient; †Student's t ; ‡Significance of the partial-regression coefficients and associated t -statistics determined by 9999 randomizations of the haplotype patristic-distance matrix.

contemporary drainage connections are negatively correlated with the patristic distance separating haplotypes, which results from some very divergent haplotype clades occupying the same present-day drainage systems. This phylogenetic pattern is expected when modern drainage basins are composites of previously isolated stream systems. Three formally recognized species in the complex correspond to monophyletic groups of haplotypes (*E. bislineata*, *E. aquatica*, and *E. junaluska*), which results in a weak but significant correlation between phylogenetic structure and current species-level taxonomy.

Molecular-demographic and nested clade analyses

Genetic signatures of geographically extensive population expansions are detected in two lineages (B and D) that were closest to the receding Wisconsin Ice Sheet (Table 3). The range of lineage B encompasses northeastern drainage

Table 3 Tajima's D -test for recent population expansion. Estimates of the population parameter θ ($2N_e\mu$) based on the number of pairwise nucleotide differences (θ_π) and segregating sites (θ_s) for haplotype clades identified in hierarchical phylogenetic analyses (Fig. 4). n is the number individuals sampled in each lineage or subclade. Standard errors of θ estimates are shown in parentheses. Significantly negative Tajima's D values (shown in boldface) indicate an excess of young mutations and support the hypothesis of recent population expansion. Lineages F, G, H, and I are excluded from analysis due to the small number of individuals included ($n < 10$)

Lineage	n	θ_π	θ_s	Tajima's D	P value
A	10	4.78 (2.88)	4.95 (2.32)	−0.16	0.475
B	42	1.03 (0.78)	4.18 (1.52)	−2.43	0.001
C	35	33.70 (16.77)	30.60 (9.39)	−0.34	0.698
D	48	6.29 (3.27)	11.72 (3.59)	−1.61	0.025
E	64	27.14 (13.35)	33.20 (9.02)	−0.67	0.288
J	58	13.54 (6.54)	22.68 (6.28)	−1.47	0.041
L	60	35.80 (17.52)	41.17 (11.23)	−0.46	0.364
M	90	32.03 (15.63)	35.89 (9.14)	−0.38	0.391

systems that were covered by the Wisconsin Ice Sheet during the last glacial maximum. Similarly, lineage D is distributed across tributaries of the former Old-Ohio River and a single tributary of the ancient Teays River basin (Kentucky River), which were impounded during the last glacial maximum. In contrast, genetic evidence for long-term persistence of lineages and subclades is detected across the Lower Tennessee and Cumberland drainage basins (lineage C), tributaries of the former Teays River basin (lineage E), Gulf Coastal Plain (lineage L), and large portions of the Upper Tennessee River basin (lineage M).




Fig. 5 Graphical summary of the nested haplotype structure and nested clade analysis (NCA) results for subclade 6-1 of lineage E. Individual haplotypes are listed across the top, with increasingly more inclusive nested groups extending to the bottom. Interior haplotypes/nested groups are depicted in bold italics. Significant D_c , D_n , and I-T values are reported. Distances that are significantly small or large are indicated with S or L, respectively. The path taken through the inference key of Templeton (2004) is shown; RGF, restricted gene flow; CRE, contiguous range expansion; FR, fragmentation.

NCA of lineages E and J reveals phylogeographic patterns consistent with the predictions of the stream-capture hypothesis. The interior clade within a portion of lineage E that spans the Ohio River basin–Atlantic slope divide (subclade 5-1) has a significantly restricted geographic distribution, indicating a contiguous range expansion from the Upper New River drainage system onto the Atlantic Slope (Fig. 5). Similarly, for lineage J, a significantly small interior-tip clade distance suggests that the oldest event shaping geographic patterns of genetic variation in this lineage was a contiguous range expansion onto the Atlantic Coastal Plain from the headwaters of the Nolichucky and Upper Catawba drainage systems (Fig. 6). Fragmentation is inferred between populations in the Neuse + Cape Fear drainages and the remaining samples, which are distributed across tributaries of the Santee, Pee Dee, and Upper Tennessee drainage systems. The genetic signal of range expansion and long-distance colonization from the headwaters of the Upper Tennessee and Upper Catawba is also reflected in less inclusive nested groups of haplotypes (4–8 and 3–20).

The MK test fails to reject the hypothesis of selective neutrality for the ND2 gene, suggesting that our inferences of population history and structure are not likely to be confounded by natural selection acting on molecular variation.

Discussion

Long-term persistence of lineages and population structure

Populations of the *Eurycea bislineata* complex are distributed across all of eastern North America's major drainage basins, including those in the Ohio River basin and northeast that were strongly affected by Pleistocene glacial advances. Our phylogeographic analysis identifies 13 geographically circumscribed population-level lineages in the group. Molecular divergences between sister lineages indicate that the vast majority of lineages originated through ancient population fragmentation events whose genetic signatures were not erased by Pleistocene drainage modifications (Hocutt *et al.* 1986; Mayden 1988) and climatic changes (Watts 1980; Webb & Bartlein 1992; Webb *et al.* 1995).

Geographically extensive range expansions are detected for the two lineages (B and D) closest to the receding Wisconsin Ice Sheet, whereas lineages to the south have more restricted geographic ranges that show evidence of long-term population stability (Fig. 2; Table 2). This spatial and temporal pattern of genetic variation matches the leading-edge model of population expansion where lineages that are closest to receding glaciers undergo exponential population




Fig. 6 Graphical summary of the nested haplotype structure and nested clade analysis (NCA) results for lineage J as in Fig. 5; LDC, long-distance colonization.

growth as they colonize recently created habitats, and lineages to the south form barriers to the geographic expansion of neighbouring lineages due to demographic and competitive effects, permitting long-term persistence of their population-genetic structure (Hewitt 1996; Zamudio & Savage 2003). These results, coupled with the extremely low vagility of two-lined salamanders, suggest that many of the lineages have sufficient geographic stability to preserve the phylogenetic signature of ancient hydrological connections and the geographic context of lineage splitting.

Drainage-basin history and phylogeography

Geomorphological evidence demonstrates that pre-Pleistocene drainage patterns in eastern North America differed drastically from current hydrological connections. Because of dispersal following the fusion of previously isolated drainages, some lineages of the *E. bislineata* complex no longer exhibit endemism in drainage basins. However, a strong association between historical drainage connections and phylogeographic divergence provides compelling support for the hypothesis that drainage evolution shaped

geographic patterns of genetic diversity across eastern North America (Table 2).

A striking feature of the *E. bislineata* complex's evolutionary history is the strong congruence of phylogeographic patterns with historical, rather than contemporary drainage patterns. The deep phylogeographic split between *E. bislineata*-complex haplotypes sampled from the Lower Tennessee + Cumberland and the Upper Tennessee River drainages is predicted by the hypothesis that the modern Tennessee River drainage formed by fusion of separate palaeodrainage basins that had independent outlets to the Gulf of Mexico until the close of the Pliocene (Figs 1, 2 and 4). Further support for historically separate outlets of the Upper and Lower Tennessee Rivers comes from the phylogenetic relationships of lineages G, H, L and M (Fig. 4). Lineage M contains primarily haplotypes from Upper Tennessee River drainages and is phylogenetically nested inside a clade with an inferred origin in the Gulf drainage basin. Similarly, *Eurycea junaluska* (lineage G: Upper Tennessee) and *Eurycea aquatica* (lineage H: Mobile basin) exhibit a sister-group relationship. The estimated times of divergence between lineages L and M (5%; ~4 Myr) and G and H (9%; ~7 Myr) are temporally

congruent with a Miocene to Pliocene formation of the modern Tennessee River basin. Together, these phylogeographic patterns provide strong support for vicariance associated with fragmentation of an ancestral drainage system connecting the Upper Tennessee River and Mobile basin.

Discovery of a deep phylogeographic split separating two distinct evolutionary lineages in the modern Ohio River basin also shows an influence of palaeodrainage basins on current geographic patterns of genetic diversity. The geographic transition between lineages D and E is broadly concordant with the pre-Pleistocene boundary between the Old-Ohio and Teays systems (Figs 1 and 2), suggesting that lineage splitting accompanied geographic isolation imposed by the divide separating these palaeodrainage basins. Furthermore, the divergence for these lineages appears to pre-date the Pleistocene integration of the modern Ohio River basin (6.5%; ~5 Myr). The geographic origins of these lineages, however, require further investigation. Although lineage E haplotypes are clearly associated with drainages of the former Teays River basin, the ancestral location of lineage D haplotypes is ambiguous. Additional sampling and coalescent simulations are needed to determine whether the latter lineage expanded from an Old Ohio, or Teays refugium (e.g. Carstens *et al.* 2005). Regardless of the geographic context of lineage splitting, these phylogeographic patterns contrast with those of codistributed stream fishes where pre-Pleistocene genetic structure was erased by glaciation and followed by rapid population expansion across most of the modern Ohio River basin (Near *et al.* 2001; Berendzen *et al.* 2003).

Secondary contacts between deeply divergent mitochondrial lineages coincide with geographic locations of major drainage divides, a pattern that reinforces the role of drainage basins in generating contemporary geographic patterns of genetic variation. For example, sympatry between lineages that span the basal north–south split in the phylogeny closely parallels the juxtaposition of the New, Tennessee, and Atlantic Slope divides in the southern Appalachian Highlands. Additional such zones of sympatry between lineages occur along the Ohio–Cumberland and Tennessee River–Gulf divides.

Despite its potential as a contemporary isolating mechanism, the ECD has migrated westward as Atlantic Slope drainage systems eroded through the Blue Ridge, fusing with and reversing the flow of western headwater drainage systems. Geological evidence indicates headwater capture of the New by the James and Roanoke Rivers (Ross 1969; Hocutt *et al.* 1986), and tributaries of the Upper Tennessee by the Santee and Savannah River drainages (Ross 1971; Starnes & Etnier 1986). Freshwater-fish distributions also indicate multiple faunal invasions of the Atlantic Slope from drainages west of the ECD facilitated by these stream captures (Hocutt *et al.* 1986; Starnes & Etnier 1986).

Geographic genetic variation in the *E. bislineata* complex supports a role for colonization of the Atlantic Slope associated with headwater stream capture. Haplotypes sampled from the Kanawha–New River drainage span the deepest phylogenetic divergence within lineage E, suggesting a Teays/Ohio River basin origin for this lineage with subsequent dispersal to the James, Roanoke, and Santee drainages on the Atlantic Slope, and Holston drainage in the Upper Tennessee River basin. Geological evidence also supports the New–Roanoke–James and New–Holston exchanges (Ross 1969; Hocutt *et al.* 1986). The haplotype phylogeny identifies two dispersal events to the Atlantic Slope in the vicinity of the drainage divide separating the Kanawha + New and Roanoke + James systems. Extensive geographic sampling for nested group 6-1 of lineage E reveals a range expansion from the headwaters of the New and Roanoke Rivers into the James River basin, the genetic pattern predicted by an eastward dispersal event facilitated by a series of hypothesized Pleistocene stream captures between these drainage systems. A recent fish phylogeographic study also demonstrates similar dispersal between the New and Roanoke drainage systems (Berendzen *et al.* 2003), suggesting that stream capture played a key role in shaping current geographic patterns of genetic and species diversity along the Atlantic Slope.

Within the southern clade, lineage J exhibits recent range expansion from the Nolichucky (Upper Tennessee) and headwaters of the upper Catawba (Santee) onto the Atlantic Coastal Plain. This phylogeographic pattern is consistent with the hypothesis that population expansion across the ECD was associated with a geologically inferred stream capture between these drainage systems (Ross 1971), and it provides strong support for the hypothesis that piracy of the upper Tennessee by headwaters of the Santee drainage system created dispersal corridors across the ECD for stream-dependent faunas (Starnes & Etnier 1986).

Gene trees and species limits

An important strength of using mtDNA haplotype phylogenies to diagnose independent evolutionary lineages is that mitochondrial genes have an effective population size one quarter that of nuclear genes (Wiens & Penkrot 2002). As a result, species become reciprocally monophyletic for mtDNA haplotypes more quickly than for those of neutral nuclear genes (Moore 1995; but see Hoelzer 1997). The ancient history of phylogeographic fragmentation uncovered within the *E. bislineata* complex and its strong congruence with historical drainage connections suggest that the current taxonomy underestimates species-level diversity of the group and obscures its biogeographic history. Moreover, few haplotypes are shared among sampling locations, indicating that even at local spatial scales, *E. bislineata*-complex populations are highly genetically structured, a

result suggesting that ongoing gene flow is not likely to be a homogenizing force where the geographic ranges of major population-level lineages abut one another.

Eurycea aquatica and *E. junaluska* each form monophyletic groups of haplotypes and are morphologically diagnosable from other sympatric lineages of the complex (Rose & Bush 1963; Sever *et al.* 1976; Ryan 1997). Thus, in contrast with earlier work that synonymized these taxa with *Eurycea cirrigera* (Jacobs 1987), our study provides strong support for their recognition as species under the phylogenetic, evolutionary, and biological species concepts (de Queiroz 1998).

Eurycea bislineata, *E. cirrigera*, and *Eurycea wilderae* each include deeply divergent haplotype clades that replace each other geographically. The distributions of lineages A, B, D, E, L, and M are broadly congruent with geographic limits of cryptic, allozyme-based genetic groupings (Jacobs 1987), supporting the hypothesis that these haplotype clades diagnose cryptic, independent evolutionary lineages. Moreover, a combination of genetic, ecological, and sexual-isolation analyses across contact zones between lineages L and M (Camp *et al.* 2000; Kozak & Montanucci 2001; Kozak 2003) and B and D (Guttman & Karlin 1986) provide independent evidence for a lack of genetic and/or ecological exchangeability (Templeton 2001) between these deeply divergent, parapatric haplotype clades. Lineages C, F, I, and K lack independent confirmation of their phylogenetic distinctness.

One potential drawback of using mtDNA to delimit species is that it may be particularly susceptible to introgression across species boundaries, leading to discordance between gene trees and species trees (Good *et al.* 2003; Glor *et al.* 2004; Morando *et al.* 2004; Weisrock *et al.* 2005). The presence of haplotypes from both *E. cirrigera* and *E. wilderae* in lineages E and J in a region of sympatry with lineage M in the southern Appalachian Highlands may indicate mtDNA introgression across the boundaries of distinct evolutionary lineages. Such geographically structured discordance is not expected for retention of ancestral polymorphisms (Good *et al.* 2003). Furthermore, the close association of these lineages with the New, Tennessee, and Atlantic Slope drainage divides is suggestive of hybridization along the margins of historical barriers to gene exchange. Alternatively, the colour morphology used to diagnose *E. wilderae* may have evolved repeatedly during the evolutionary history of the *E. bislineata* complex, with the haplotype phylogeny providing a more accurate picture of the geographic limits of independent evolutionary lineages. Supporting the latter hypothesis, geographic patterns of genetic variation in allozymes also demonstrate that *E. wilderae* is nonmonophyletic and reveal an ancient genetic break separating populations in the New and Tennessee River drainages (Jacobs 1987). Further fine-scale population-genetic studies that employ multiple unlinked molecular markers are needed to discriminate these alternatives.

Conclusions

The geological history of stream drainage basins provides a wealth of a priori hypotheses for the geographic fragmentation of populations of stream-dependent organisms that are ideally suited for testing with phylogeographic methodologies. Here, we demonstrate that salamanders of the *E. bislineata* complex exhibit a complex phylogeographic history that preserves the genetic signatures of palaeo-drainage connections that were altered by glacial advances and headwater erosion. Our molecular genetic results reinforce geological and faunistic evidence for composite origins of major eastern North American drainage systems. More generally, our study provides a general methodological framework for combining independent biological and geological information to formulate and to test hypotheses of the complex interaction between historical and contemporary factors that generate and maintain patterns of biodiversity in stream-dependent organisms. The generality of the phylogeographic hypotheses tested in this study should be investigated using comparable analyses in other codistributed stream-dependent species.

Acknowledgements

We thank R. Bonett, C. Camp, R. Highton, D. Huestis, J. Marshall, K. Zamudio, W. Savage, W. Van Devender, and the LSU Collection of Genetic Resources for donating tissue samples for genetic analysis. R. Bonett, M. Gifford, and three anonymous reviewers provided helpful comments on the manuscript. The American Society of Ichthyologists and Herpetologist's Gaige Fund, Highlands Biological Station's Bruce Family and Charles Ash Scholarships, and a National Science Foundation Doctoral Dissertation Improvement Grant DEB-0206739 generously supported this research.

Supplementary material

The supplementary material is available from <http://www.blackwellpublishing.com/products/journals/suppmat/MEC/MEC2757/MEC2757sm.htm>

Table S1 Geographic sampling information for the salamanders used in this study. Population numbers correspond to Fig. 2.

Table S2 List of *Eurycea bislineata* complex haplotypes sequenced in this study, the localities in which they were found, field numbers of specimens from which they were sampled, and GenBank Accession numbers. Haplotypes are named according to the taxonomic species from which they were sampled. Populations are numbered as in Table S1, and Figs 2 and 4, with the number of individuals sampled for each haplotype shown in parentheses. KHK, Kenneth H. Kozak field series; RMB, Ronald M. Bonett field series; WKS, Wesley K. Savage field series; H, Louisiana State University Museum of Natural History Collection of Genetic Resources; numbers without a prefix indicate Richard Highton field numbers. Voucher specimens from KHK's field series will be deposited in the LSU Museum of Natural History following a formal taxonomic revision of the complex.

References

- Berendzen PB, Simons AM, Wood RM (2003) Phylogeography of the northern hogsucker, *Hypentelium nigricans* (Teleostei: Cypriniformes): genetic evidence for the existence of the ancient Teays River. *Journal of Biogeography*, **30**, 1139–1152.
- Bremer K (1994) Branch support and tree stability. *Cladistics*, **10**, 295–304.
- Bruce RC (1986) Upstream and downstream movements of *Eurycea bislineata* and other salamanders in a southern Appalachian stream. *Herpetologica*, **42**, 149–155.
- Burr BM, Page LM (1986) Zoogeography of fishes of the Lower Ohio–Upper Mississippi Basin. In: *The Zoogeography of North American Freshwater Fishes* (eds Hocutt CH, Wiley EO), pp. 290–324. John Wiley & Sons, New York.
- Camp CD, Marshall JL, Landau KR, Austin RM Jr, Tilley SG (2000) Sympatric occurrence of two species of the two-lined salamander (*Eurycea bislineata*) complex. *Copeia*, **2000**, 572–578.
- Carstens BC, Stevenson AL, Degenhardt JD, Sullivan J (2004) Testing nested phylogenetic and phylogeographic hypotheses in the *Plethodon vandykei* species group. *Systematic Biology*, **53**, 781–792.
- Carstens BC, Degenhardt JD, Stevenson AL, Sullivan J (2005) Accounting for coalescent stochasticity in testing phylogeographical hypotheses: modeling Pleistocene population structure in the Idaho giant salamander *Dicamptodon aterrimus*. *Molecular Ecology*, **14**, 255–265.
- Casgrain P (2001) *PERMUTE!*, Version 3.4 alpha. University of Montréal, Montréal, Québec.
- Clement MD, Posada D, Crandall KA (2000) rcs: a computer program to estimate gene genealogies. *Molecular Ecology*, **9**, 1657–1659.
- Durand JD, Templeton AR, Guinand B, Imsiridou A, Bouvet Y (1999) Nested clade and phylogeographic analyses of the chub, *Leuciscus cephalus* (Teleostei, Cyprinidae), in Greece: implications for Balkan Peninsula biogeography. *Molecular Phylogenetics and Evolution*, **13**, 566–580.
- Glor RE, Gifford ME, Larson A *et al.* (2004) Partial island submergence and speciation in an adaptive radiation: a multi-locus analysis of the Cuban green anoles. *Proceedings of the Royal Society of London. Series B, Biological Sciences*, **271**, 2257–2265.
- Good JM, Demboski JR, Nagorsen DW, Sullivan J (2003) Phylogeography and introgressive hybridization: chipmunks (genus *Tamias*) in the northern Rocky Mountains. *Evolution*, **57**, 1900–1916.
- Guttman SI, Karlin AA (1986) Hybridization of cryptic species of two-lined salamanders (*Eurycea bislineata* complex). *Copeia*, **1986**, 96–108.
- Hardy ME, Grady JM, Routman E (2002) Intraspecific phylogeography of the slender madtom: the complex evolutionary history of the Central Highlands of the United States. *Molecular Ecology*, **11**, 2393–2403.
- Hewitt GM (1996) Some genetic consequences of ice ages, and their role in divergence and speciation. *Biological Journal of the Linnean Society*, **58**, 247–276.
- Hewitt GM (2001) Speciation, hybrid zones and phylogeography – or seeing genes in space and time. *Molecular Ecology*, **10**, 537–549.
- Hocutt CH, Jenkins RE, Stauffer JR Jr (1986) Zoogeography of the fishes of the central Appalachians and central Atlantic Coastal Plain. In: *The Zoogeography of North American Freshwater Fishes* (eds Hocutt CH, Wiley EO), pp. 161–211. John Wiley & Sons, New York.
- Hoelzer GA (1997) Inferring phylogenies from mtDNA variation: mitochondrial-gene trees versus nuclear-gene trees revisited. *Evolution*, **51**, 622–626.
- Huelsenbeck JP, Bollback JP (2001) Empirical and hierarchical Bayesian estimation of ancestral states. *Systematic Biology*, **50**, 351–366.
- Huelsenbeck JP, Ronquist F (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics*, **17**, 754–755.
- Jacobs JF (1987) A preliminary investigation of geographic genetic variation and systematics of the two-lined salamander, *Eurycea bislineata* (Green). *Herpetologica*, **43**, 423–446.
- Johnson JB (2002) Evolution after the flood: phylogeography of the desert fish Utah chub. *Evolution*, **56**, 948–960.
- Knowles LL (2001) Did the Pleistocene glaciations promote divergence? Tests of explicit refugial models in montane grasshoppers. *Molecular Ecology*, **10**, 691–701.
- Kozak KH (2003) Sexual isolation and courtship behavior in salamanders of the *Eurycea bislineata* species complex, with comments on the evolution of the mental gland and pheromone delivery behavior in the Plethodontidae. *Southeastern Naturalist*, **2**, 281–292.
- Kozak KH, Lannoo MJ (2005) *Eurycea aquatica*. In: *Amphibian Declines: the Conservation Status of United States Species* (ed. Lannoo MJ), pp. 734–735. University of California Press, Berkeley, California.
- Kozak KH, Montanucci RR (2001) Genetic variation across a contact zone between montane and lowland forms of the two-lined salamander (*Eurycea bislineata*) species complex: a test of species limits. *Copeia*, **2001**, 25–34.
- Kozak KH, Larson A, Bonett RM, Harmon LJ (2005) Phylogenetic analysis of ecomorphological divergence, community structure, and diversification rates in dusky salamander (Plethodontidae: *Desmognathus*). *Evolution*, **59**, 2000.
- Kreiser BR, Mitton JB, Woodling JD (2001) Phylogeography of the plains killifish, *Fundulus zebrinus*. *Evolution*, **55**, 339–350.
- Kumar S, Tamura K, Jakobsen IB, Nei M (2001) MEGA2: Molecular evolutionary genetic analysis software. Arizona State University, Tempe, Arizona.
- Legendre P, Lapointe F-J, Casgrain P (1994) Modeling brain evolution from behavior: a permutational regression approach. *Evolution*, **48**, 1487–1499.
- Losos JB, Glor RE (2003) Phylogenetic comparative methods and the geography of speciation. *Trends in Ecology & Evolution*, **18**, 220–227.
- MacCulloch RD, Bider JR (1975) Phenology, migrations, circadian rhythm and the effect of precipitation on the activity of *Eurycea b. bislineata* in Quebec. *Herpetologica*, **31**, 433–439.
- Macey JR, Verma A (1997) Homology in phylogenetic analysis: alignment of transfer RNA genes and the phylogenetic position of snakes. *Molecular Phylogenetics and Evolution*, **7**, 272–279.
- Macey JR, Larson A, Ananjeva NB, Fang ZL, Papenfuss TJ (1997) Two novel gene orders and the role of light-strand replication in rearrangement of the vertebrate mitochondrial genome. *Molecular Biology and Evolution*, **14**, 91–104.
- Macey JR, Schulte JA, Larson A *et al.* (1998) Phylogenetic relationships of toads in the *Bufo bufo* species group from the eastern escarpment of the Tibetan Plateau: a case of vicariance and dispersal. *Molecular Phylogenetics and Evolution*, **9**, 80–87.
- Maddison WP, Maddison DR (1992) *MACCLADE: analysis of phylogeny*

- and character evolution, version 3.0. Sinauer Associates, Sunderland, Massachusetts.
- Masta SE (2000) Phylogeography of the jumping spider *Habronattus pugillis* (Aranae: Salticidae): recent vicariance of sky island populations? *Evolution*, **54**, 1699–1711.
- Mayden RL (1988) Vicariance biogeography, parsimony, and evolution in North American freshwater fishes. *Systematic Zoology*, **37**, 329–355.
- McDonald JH, Kreitman M (1991) Adaptive protein evolution at the *Adh* locus in *Drosophila*. *Nature*, **351**, 652–654.
- Moore WS (1995) Inferring phylogenies from mtDNA variation – mitochondrial-gene trees versus nuclear-gene trees. *Evolution*, **49**, 718–726.
- Morando M, Avila LJ, Baker J, Sites JW Jr (2004) Phylogeny and phylogeography of the *Liolaemus darwini* complex (Squamata: Liolaemidae): evidence for introgression and incomplete lineage sorting. *Evolution*, **58**, 842–861.
- Near TJ, Page LM, Mayden RL (2001) Intraspecific phylogeography of *Percina evides* (Percidae: Etheostomatinae): an additional test of the Central Highlands pre-Pleistocene vicariance hypothesis. *Molecular Ecology*, **10**, 2235–2240.
- Posada D, Crandall KA, Templeton AR (2000) GEODIS: a program for the cladistic nested analysis of the geographical distribution of genetic haplotypes. *Molecular Ecology*, **9**, 487–488.
- de Queiroz K (1998) The general lineage concept of species, species criteria, and the process of speciation: a conceptual unification and terminological recommendations. In: *Endless Forms: Species and Speciation* (eds Howard DJ, Berlocher SH), pp. 57–75. Oxford University Press, Oxford.
- Robinson-Rehavi M, Huchon D (2000) RRTree: relative-rate tests between groups of sequences on a phylogenetic tree. *Bioinformatics*, **16**, 296–297.
- Rose FL, Bush FM (1963) A new species of *Eurycea* (Amphibia: Caudata) from the southeastern United States. *Tulane Studies in Zoology*, **10**, 121–128.
- Ross RD (1969) Drainage evolution and fish distribution problems in the southern Appalachians of Virginia. In: *The Distributional History of the Biota of the Southern Appalachians, Part I, Invertebrates* (ed. Holt PC), pp. 277–292. Virginia Polytechnic Institute, Research Division Monograph I, Blacksburg, Virginia.
- Ross RD (1971) The drainage history of the Tennessee River. In: *The Distributional History of the Biota of the Southern Appalachians, Part III, Vertebrates* (ed. Holt PC), pp. 11–42. Virginia Polytechnic Institute, Research Division Monograph III, Blacksburg, Virginia.
- Routman ER, Wu R, Templeton AR (1994) Parsimony, molecular evolution, and biogeography: the case of the North American giant salamander. *Evolution*, **48**, 1799–1809.
- Rozas J, Sanchez-Delbarrio JC, Messeguer X, Rozas R (2003) DNASP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics*, **19**, 2496–2497.
- Ryan TJ (1997) Larva of *Eurycea junaluska* (Amphibia: Caudata: Plethodontidae), with comments on distribution. *Copeia*, **1997**, 210–215.
- Schneider S, Roessli DL, Excoffier L (2000) ARLEQUIN (version 2.2): A software for population genetics data analysis. Genetics and Biometry Laboratory, University of Geneva, Switzerland.
- Sever DM, Dundee HA, Sullivan CD (1976) A new *Eurycea* (Amphibia: Plethodontidae) from southwestern North Carolina. *Herpetologica*, **32**, 26–29.
- Slatkin M (1993) Isolation by distance in equilibrium and non-equilibrium populations. *Evolution*, **47**, 264–279.
- Slatkin M, Maddison WP (1989) A cladistic measure of gene flow inferred from the phylogenies of alleles. *Genetics*, **123**, 603–613.
- Smouse PE, Long JC, Sokal RR (1986) Multiple regression and correlation extensions of the Mantel tests of matrix correspondence. *Systematic Zoology*, **35**, 627–632.
- Sorenson MD (1999) *TREEROT (version 2)*. Boston University, Boston, Massachusetts.
- Starnes WC, Etnier DA (1986) Drainage evolution and fish biogeography of the Tennessee and Cumberland Rivers drainage realm. In: *The Zoogeography of North American Freshwater Fishes* (eds Hocutt CH, Wiley EO), pp. 326–359. John Wiley & Sons, New York.
- Strange RM, Burr BM (1997) Intraspecific phylogeography of North American highland fishes: a test of the Pleistocene vicariance hypothesis. *Evolution*, **51**, 885–897.
- Swofford D (2002) *PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods)*. Sinauer Associates, Sunderland, Massachusetts.
- Tajima D (1989) The effect of change in population size on DNA polymorphism. *Genetics*, **123**, 597–601.
- Tamura K, Nei M (1993) Estimation of the number of nucleotide differences in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution*, **10**, 512–526.
- Templeton AR (2001) Using phylogeographic analyses of gene trees to test species status and processes. *Molecular Ecology*, **10**, 779–791.
- Templeton AR (2004) Statistical phylogeography: methods of evaluating and minimizing inference errors. *Molecular Ecology*, **13**, 789–809.
- Templeton AR, Sing CF (1993) A cladistic analysis of phenotypic associations with haplotypes inferred from restriction endonuclease mapping. IV. Nested analyses with cladogram uncertainty and recombination. *Genetics*, **134**, 659–669.
- Templeton AR, Boerwinkle E, Sing CF (1987) A cladistic analysis of phenotypic associations inferred from restriction endonuclease mapping. *Genetics*, **117**, 343–351.
- Thornbury WD (1965) *Regional Geomorphology of the United States*. John Wiley & Sons, New York.
- Thorpe RS, Black H, Malhotra A (1996) Matrix correspondence tests on the DNA phylogeny of the Tenerife lacertid elucidate both historical causes and morphological adaptation. *Systematic Biology*, **45**, 335–343.
- Voss SR, Smith DG, Beachy CK, Heckel DG (1995) Allozyme variation in neighboring isolated populations of the plethodontid salamander *Leurognathus marmoratus*. *Journal of Herpetology*, **29**, 493–497.
- Waters JM, Craw D, Youngson JH, Wallis GP (2001) Genes meet geology: fish phylogeographic pattern reflects ancient, rather than modern, drainage connections. *Evolution*, **55**, 1844–1851.
- Watts WA (1980) The late Quaternary vegetation history of the southeastern United States. *Annual Review of Ecology and Systematics*, **11**, 387–409.
- Webb T, Bartlein PJ (1992) Global changes during the last 3 million years: climatic controls and biotic responses. *Annual Review of Ecology and Systematics*, **23**, 141–173.
- Webb SD, Hulbert RC Jr, Lambert WD (1995) Climatic implications of large-herbivore distributions in the Miocene of North America. In: *Paleoclimate and Evolution, with Emphasis on Human Origins* (eds Vrba ES, Denton GH, Partridge TC, Burkle LH), pp. 91–108. Yale University Press, New Haven, Connecticut.
- Weisrock DW, Macey JR, Ugurtas IH, Larson A, Papenfuss TJ (2001) Molecular phylogenetics and historical biogeography among salamandrids of the ‘true’ salamander clade: rapid

- branching of numerous highly divergent lineages in *Mertensiella luschani* associated with the rise of Anatolia. *Molecular Phylogenetics and Evolution*, **18**, 434–448.
- Weisrock DW, Kozak KH, Larson A (2005) Phylogeographical analysis of mitochondrial gene flow and introgression in the salamander, *Plethodon shermani*. *Molecular Ecology*, **14**, 1457–1472.
- Wiens JJ, Penkrot T (2002) Delimiting species using DNA and morphological variation and discordant species limits in spiny lizards (*Sceloporus*). *Systematic Biology*, **51**, 69–91.
- Zamudio KR, Savage WK (2003) Historical isolation, range expansion, and secondary contact of two highly divergent mitochondrial lineages in spotted salamanders (*Ambystoma maculatum*). *Evolution*, **57**, 1631–1652.
- Zhang DX, Hewitt GM (1996) Nuclear integrations: challenges for mitochondrial DNA markers. *Trends in Ecology & Evolution*, **11**, 247–251.

This study was conducted as part of Kenneth Kozak's PhD thesis focused on investigating patterns of speciation, ecological diversification, and community assembly in eastern North American plethodontid salamanders. He is currently a National Science Foundation Postdoctoral Fellow in Bioinformatics at Stony Brook University where he is investigating the relationship between landscape ecology and spatial patterns of genetic variation and species richness in woodland salamanders. Russell Blaine is a PhD candidate at Washington University. His research is focused on comparative phylogeography and landscape genetics of reptiles in the desert southwest of North America. Allan Larson is Professor of Biology at Washington University. His research specialty is molecular phylogenetics and evolution of vertebrates, particularly lizards and salamanders.
