

Gene Therapy for Cartilage Repair

Henning Madry, Patrick Orth and Magali Cucchiari
Cartilage 2011 2: 201 originally published online 11 April 2011
DOI: 10.1177/1947603510392914

The online version of this article can be found at:
<http://car.sagepub.com/content/2/3/201>

Published by:



<http://www.sagepublications.com>

On behalf of:

International Cartilage Repair Society

Additional services and information for *Cartilage* can be found at:

Email Alerts: <http://car.sagepub.com/cgi/alerts>

Subscriptions: <http://car.sagepub.com/subscriptions>

Reprints: <http://www.sagepub.com/journalsReprints.nav>

Permissions: <http://www.sagepub.com/journalsPermissions.nav>

Gene Therapy for Cartilage Repair

Cartilage
2(3) 201–225
© The Author(s) 2011
Reprints and permission:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/1947603510392914
http://cart.sagepub.com



Henning Madry¹, Patrick Orth¹, and Magali Cucchiari¹

Abstract

The concept of using gene transfer strategies for cartilage repair originates from the idea of transferring genes encoding therapeutic factors into the repair tissue, resulting in a temporarily and spatially defined delivery of therapeutic molecules to sites of cartilage damage. This review focuses on the potential benefits of using gene therapy approaches for the repair of articular cartilage and meniscal fibrocartilage, including articular cartilage defects resulting from acute trauma, osteochondritis dissecans, osteonecrosis, and osteoarthritis. Possible applications for meniscal repair comprise meniscal lesions, meniscal sutures, and meniscal transplantation. Recent studies in both small and large animal models have demonstrated the applicability of gene-based approaches for cartilage repair. Chondrogenic pathways were stimulated in the repair tissue and in osteoarthritic cartilage using genes for polypeptide growth factors and transcription factors. Although encouraging data have been generated, a successful translation of gene therapy for cartilage repair will require an ongoing combined effort of orthopedic surgeons and of basic scientists.

Keywords

gene therapy, cartilage repair, osteoarthritis, meniscal lesions, clinical trials

Introduction

Articular cartilage defects and meniscal lesions have a reduced capacity for regeneration. The concept of using gene transfer strategies for cartilage repair originates from the idea of transferring genes encoding therapeutic factors into the repair tissue, resulting in a temporarily and spatially defined delivery of the therapeutic molecule. In this review, we will focus on gene therapy approaches for the repair of articular cartilage and meniscal fibrocartilage, including articular cartilage defects resulting from acute trauma, osteochondritis dissecans, osteonecrosis, and osteoarthritis. Possible applications for meniscal repair will be described for meniscal lesions, meniscal sutures, and meniscal transplantation. As a discussion of cartilage damage resulting from rheumatoid arthritis is beyond the scope of this review, we refer to the many reviews already published on this subject.^{1–9}

Principles of Gene Therapy

Gene transfer is the introduction of foreign genes or gene sequences into different types of cells. Gene therapy is the treatment of diseases using gene transfer techniques. Gene transfer via nonviral vectors is named transfection; gene transfer using viral vectors is termed transduction. The foreign genetic material enters the cell and is next transferred towards the nucleus, where it either integrates into

the host genome or remains extrachromosomal as an episome that generally allows only for transient transgene expression. For therapeutic applications, gene transfer into a sufficiently high number of target cells is essential for the secretion of relevant concentrations of the transgene product. Current vectors available for use in gene therapy include nonviral approaches (naked DNA, physical and chemical methods) and various viral (adenoviral, HSV, retroviral, lentiviral, rAAV) vehicles (**Table 1**).

Among the nonviral systems, chemical methods of complexing DNA to various macromolecules include cationic lipids and liposomes,^{10–12} polymers,¹³ polyamines and polyethylenimines,^{14,15} and nanoparticles,¹⁶ but also calcium phosphate coprecipitates¹⁷ are mainly used. Nonviral systems avoid the risk of acquiring replication competence inherent to viral vectors, can be repeatedly administered, have the capacity to carry large therapeutic genes, are relatively easy to produce on a large scale, and do not elicit a detectable immune response. Nevertheless, their efficacy is often inferior to those of viral vectors. Moreover, the fact that they stay as episomal forms in the target cells often

¹Saarland University, Homburg, Germany

Corresponding Author:

Henning Madry, Saarland University, Kirrbergerstrasse 1, Homburg, 66424 Germany
Email: henning.madry@uks.eu

Table 1. Nonviral and Viral Gene Vectors Suitable for Gene Transfer to Cartilage Defects

	Nonviral Systems		Viral Systems			
	Liposomes	Others (Chemical, Electrical, and Mechanical Methods)	Adenovirus	Retrovirus	Herpes Simplex Virus (HSV)	Adeno-Associated Virus (AAV)
Advantages	Independent from cell cycle Noninfectious; repeatedly applicable Low toxicity Large capacity Easy to manufacture		Very high efficiency Independent from cell cycle Approved for clinical trials	High efficiency Prolonged transgene expression	High efficiency Independent from cell cycle Large capacity	Very high efficiency Prolonged transgene expression Independent from cell cycle Noninfectious; repeatedly applicable
Shortcomings	Cell-specific efficiency Short-term transgene expression		Infectious with induction of immune response; single application only Cytotoxicity Risk of replication competence Short-term transgene expression	Insertional mutagenesis Dependent from cell cycle Risk of replication competence Restricted host range	Short-term transgene expression Cytotoxicity (first- generation HSV)	Difficult to manufacture
Integration in host genome	No	No	No	Yes	No	No

Note: Properties of nonviral and viral gene vectors currently in clinical and experimental use for gene therapy approaches to cartilage defects.

results in short-term transgene expression. To avoid low gene transfer efficacy *in vivo*, nonviral gene transfer strategies are often based on the transplantation of *ex vivo*-modified cells to cartilage defects.

Viral vectors utilize natural entry pathways in human cells. Adenoviral vectors have been among the most employed gene vehicles for cartilage repair in the past.¹⁸⁻²² They allow for high transduction efficiencies and transgene expression in a variety of cells, enabling direct approaches *in vivo*. However, serious concerns about their clinical safety were raised after the death of Jesse Gelsinger, a patient included in a gene therapy trial employing adenoviral vectors. Moreover, transgene expression via adenoviral delivery is limited for about 1 to 2 weeks as the transgenes remain episomal and due to the development of host immune responses against transduction with most of the constructs derived from these viruses.

An advantage of retroviruses is their ability to integrate in the genome of the target, allowing for the replication and maintenance of the transgene over extended periods of time. Yet, this might lead to insertional mutagenesis, with the potential for activating tumor genes. Also, retroviral vectors do not transduce nondividing cells and have a restricted host range. As for nonviral systems, *ex vivo*

approaches with selection of transduced cells are usually required with retroviral vectors²³⁻²⁷ because they are produced only at relatively medium titers and do not exhibit very high efficiencies. Instead, lentiviral vectors, a subclass of retroviruses derived from the human immunodeficiency virus (HIV), can integrate in the genome of nondividing cells.²⁸ Therefore, such vectors might be good alternatives to the use of retroviruses, as they show also higher levels of transduction *in vivo* and avoid the need for cell division.^{29,30} Yet, there are common concerns associated with their application, including the potential for insertional mutagenesis and the psychological problem of introducing genetic material carrying HIV sequences.

Herpes simplex virus (HSV)-derived vectors are large vehicles that can deliver long transgenes to almost all known cell types, including nondividing cells. Although first-generation vectors induced high levels of cytotoxicity, recent work has demonstrated that second-generation HSV were less deleterious, in particular for cartilage repair.³¹ One problem remains the transient nature of transgene expression mediated by this family of vectors.

In any case, the direct application of viral vectors raises legitimate safety concerns, as potentially infectious agents or sequences (especially lentiviral vectors) might be intro-

duced *per se* in the body. This is of particular importance for the treatment of cartilage and meniscal lesions that are not life-threatening disorders. In this regard, adeno-associated viral vectors (AAV), which are based on the nonpathogenic, replication-defective human parvovirus AAV,³² might prove more adequate in direct gene therapy settings. Vectors based on AAV (rAAV) are produced by complete removal of the viral gene coding sequences, making them less immunogenic than adenoviral vectors and less toxic than HSV. Also, the latter vectors generally mediate only short-term expression of the transgenes they carry, whereas rAAV can be transcribed for months to years due to the stabilization of the episomal transgene cassettes by concatemer formation.³³⁻³⁶ Cell division and integration are not required for expression of the foreign material delivered, in marked contrast with retroviral vectors.³⁷ Redosing of vectors is practicable with rAAV, based on the manipulation of various available serotypes of the virus. For these reasons, rAAV became a preferred gene transfer method for experimental settings *in vivo* and for clinical applications.^{35,36,38,39}

The greatest obstacle to develop efficient gene transfer protocols targeting sites of articular cartilage and meniscal fibrocartilage damage so far has been the restrained accessibility of the lesions to a treatment. Therefore, the following experimental approaches are currently employed to transfer genes to sites of interest *in vivo* (Fig. 1):

1. intra-articular injection of the therapeutic formulation, and
2. administration of the therapeutic formulation to the defect via arthrotomy:
 - 2.1. direct application of a gene vector to the repair tissue,
 - 2.2. application of biomaterials carrying a gene vector, and
 - 2.3. matrix-supported application of *ex vivo* genetically modified cells.

The target cells in which genes may be transferred include the following:

1. progenitor cells (e.g., resulting from marrow-stimulating techniques or transplanted cells),
2. isolated articular chondrocytes or meniscal fibrochondrocytes that are transplanted into the defect, and
3. cells of the tissues adjacent to the defect:
 - 3.1. articular cartilage: articular chondrocytes from the adjacent cartilage, osteoblasts, and osteocytes from the subchondral bone; and
 - 3.2. meniscal tissue: meniscal fibrochondrocytes, synoviocytes from the synovial lining, and fibroblasts from the joint capsule.




Figure 1. Therapeutic genes may be transferred to sites of articular cartilage damage or to meniscal lesions *in vivo* via intra-articular injection or by direct application into the lesion. Intra-articular injection (upper panel) of the therapeutic formulation (most often a viral vector) results in a nonselective transduction of many intra-articular tissues. Direct administration of the therapeutic formulation (lower panel) to the target lesion (e.g., an articular cartilage defect) can be achieved by directly applying a gene vector to the repair tissue in the defect (left), by matrix-supported application (e.g., alginate) of target cells (e.g., articular chondrocytes, meniscal fibrochondrocytes, progenitor cells) that were previously genetically modified *ex vivo* (middle), or by application of a gene vector attached to a biomaterial (right). *In vivo*, it often includes an arthrotomy.

Articular Cartilage

Introduction

Anatomy, Function, and Pathophysiology. Adult hyaline articular cartilage is avascular and aneural and does not possess a lymphatic drainage.⁴⁰ Its major function is to allow for a smooth gliding of the articulating surfaces of a joint and to protect the subchondral bone from mechanical stress. Hyaline articular cartilage is structured in several laminar zones and formed by chondrocytes that are surrounded by an intricate network of extracellular matrix. This cartilaginous matrix is rich in proteoglycans and collagen fibrils composed of type II collagen but also contains types VI, IX, XI, and XIV collagens and a number of additional macromolecules.⁴¹ Normal hyaline articular cartilage contains about 70% to 80% water, which is mainly bound to proteoglycans. Articular chondrocytes synthesize and degrade the extracellular matrix, thereby regulating the structural and functional properties according to the applied loads.

The integrity of articular cartilage can be disrupted as a result of mainly 4 different etiologies.⁴² These include focal articular cartilage defects resulting from an acute trauma, osteoarthritis, osteonecrosis, and osteochondritis dissecans.⁴³ The resulting articular cartilage defect is characterized as being either chondral, involving only the cartilaginous zones, or osteochondral, reaching further into the subchondral bone.⁴⁴ Although a chondral defect may be in part repopulated by cells from the synovial membrane,^{45,46} it usually remains and may expand over time. An osteochondral defect is filled with a blood clot that forms if the bone marrow communicates with the defect.^{47,48} The pluripotent, undifferentiated mesenchymal cells of the blood clot differentiate into chondrocytes and osteoblasts that later form the cartilaginous repair tissue and the new subchondral bone. However, over time, this repair tissue increasingly exhibits characteristics of fibrocartilage, such as an increased type I and a decreased type II collagen content and may degenerate after several years.⁴⁸ If left untreated, secondary osteoarthritis of the joint may result.

Chondrogenic Therapeutic Factors. Strategies for enhancing chondrogenesis in an articular cartilage defect aim at improving the differentiation of mesenchymal cells into chondrocytes for cartilage repair and osteoblasts for the repair of the subchondral bone, the production and maintenance of a new cartilaginous matrix rich in type II collagen and proteoglycans, at increasing the cellularity of the repair tissue to prevent the hypertrophic differentiation of chondrocytes, and at inhibiting articular cartilage degeneration.

Growth and transcription factors are good candidates for these approaches. The therapeutic efficacy of polypeptide growth factors is, however, diminished by their short half-lives.⁴⁹⁻⁵¹ For example, the fibroblast growth factor-2

polypeptide has a plasma half-life of less than 1 hour and is cleared in some hours after intra-articular administration.⁵⁰ To overcome this problem, the idea of applying the gene encoding for a particular therapeutic protein has gained attraction.

Candidate factors to support chondrogenesis include members of the transforming growth factor beta (TGF- β) superfamily such as TGF- β 1 and TGF- β 2,^{27,52,53} bone morphogenetic protein 2 (BMP-2),^{51,53} BMP-7,^{54,55} members of the fibroblast growth factor family such as the basic fibroblast growth factor (FGF-2),⁵⁶ growth/differentiation factor 5 (GDF-5),⁵⁷ and the parathyroid hormone-related protein (PTHrP).^{58,59} Cell proliferation is promoted, among others, by FGF-2^{60,61} and the insulin-like growth factor I (IGF-I).⁶² Particularly potent candidates to stimulate matrix synthesis include IGF-I,^{63,64} BMP-2 and BMP-7, and the cartilage-derived morphogenetic proteins (CDMP).^{65,66}

Transcription factors directly modulate the expression of genes involved in chondrogenesis, such as type II collagen or aggrecan. Experimental models have demonstrated the chondrogenic properties of transcription factors, such as SOX9,⁶⁷ Cbfa-1/Runx-2,⁶⁸ Cart-1,⁶⁹ the Ets family members,⁷⁰ and various signaling molecules as well as extracellular matrix glycoproteins themselves.^{71,72} Another attractive approach is to inhibit degenerative pathways within the repair tissue. Potential targets include cytokines that mediate catabolic events, in particular the members of the interleukin-1 (IL-1),⁷³ IL-17,⁷⁴ and tumor necrosis factor (TNF)⁷⁵ families. These strategies are based on the inhibition of the production of matrix-degrading enzymes,⁷⁶ proinflammatory mediators,⁷⁵ as well as apoptotic mechanisms.⁷⁷

Traumatic Articular Cartilage Defects

Intra-articular Injection. Intra-articular injection is a convenient way to target the joint space and has been studied using naked DNA⁷⁸ or adenoviral,^{79,80} retroviral,^{81,82} HSV,⁷⁹ lentivirus,²⁹ rAAV,^{81,83,84} and nonviral vectors.^{79,85} In 1998, Ikeda *et al.*⁸⁰ injected adenoviral vectors encoding for the TGF- β 1 gene into the joints of guinea pigs and reported elevated TGF- β 1 levels in the synovial fluid for 2 weeks following gene delivery. The effectiveness of a direct intra-articular gene therapy approach in combination with a marrow stimulation technique has been shown by Morisset *et al.*⁸⁶ Full-thickness chondral defects in equine stifle and knee joints were treated by microfracturing, followed by intra-articular application of adenoviral vectors carrying the genes for interleukin-1 receptor antagonist protein (IL-1Ra) and IGF-I. Sixteen weeks postoperatively, articular cartilage defects treated with IL-1Ra and IGF-I showed increased proteoglycan content and type II collagen expression compared with defects treated using a marrow-stimulating technique alone. Yet, articular cartilage defects cannot be specifically

targeted with this approach since the transgene is expressed mainly in cells of the synovial membrane and gene transfer into articular cartilage defect is a very rare event. Therefore, many of the gene-based approaches have focused on direct gene vector delivery into a defect exposed by arthrotomy (Table 2).

Arthrotomy

Direct application of a gene vector in vivo. The direct delivery of therapeutic genes into cartilage defects in depth has long been arduous due to the reduced capability of nonviral and various viral vectors to penetrate the dense extracellular cartilaginous matrix. Following arthrotomy and gene vector application to cartilage defects, limited transgene expression was observed only in the superficial cartilage layers.⁸⁰ With the implementation of rAAV vectors, direct gene transfer to cells within defects and adjacent cartilage has met success. Reporter gene studies demonstrated efficient transgene expression in normal and osteoarthritic human articular chondrocytes within their native matrix *in situ* to depths relevant for clinical applications.⁸⁷ Moreover, transgene expression was also present in chondral and osteochondral articular cartilage defects *in vivo* for at least 4 months.⁸⁷ rAAV vectors have been manipulated recently to deliver therapeutic genes such as FGF-2 directly into osteochondral cartilage defects.³⁶ Cartilage repair was significantly enhanced 4 months after vector application.³⁶

Application of biomaterials carrying a gene vector into defects. In order to avoid a dilution of the therapeutic agents, gene vectors or modified cells can be delivered in conjunction with biomaterials such as fibrin, collagen, gelatin, carbohydrate-based polymers (polyactic acid/polyglycolic acid, hyaluronan, agarose, alginate, chitosan), and artificial polymers (dacron, teflon, carbon fibers, polyesterurethane, polybutyric acid, polyethylmethacrylate, hydroxyapatite).^{45,88} When preparations of adenoviral vectors carrying a marker gene were adsorbed onto type II collagen-glycosaminoglycan matrices and implanted into osteochondral defects, transgene expression was present until day 21.⁸⁹

Application of ex vivo genetically modified cells. The direct transplantation of cells genetically modified *ex vivo* involves their isolation, genetic modification, and reimplantation into articular cartilage defects. These modified cells can be applied without (e.g., as coagulated bone marrow aspirate) or with supportive matrices. Such components include alginate,⁹⁰⁻⁹² agarose,^{93,94} fibrin or type I collagen gels without⁹⁵⁻⁹⁷ or with a periosteal flap,^{98,99} and synthetic biodegradable scaffolds.¹⁰⁰⁻¹⁰² Kang *et al.* were the first to transplant genetically modified cells into an articular cartilage defect *in vivo*.¹⁰³ In this study, chondrocytes were transduced with a retroviral vector. Other studies used nonviral,¹⁰⁴⁻¹⁰⁶ adenoviral,^{89,96,107} retroviral,^{103,108-111} and rAAV vectors¹¹² to deliver marker genes in defects via *ex vivo*-modified cells. Although engineered chondrocytes




Figure 2. Improvement of cartilage repair in a rabbit osteochondral defect model in the trochlear groove by combined *ex vivo* gene transfer of human insulin-like growth factor I (hIGF-I) and fibroblast growth factor-2 (hFGF-2) in NIH 3T3 fibroblasts that were then embedded in alginate spheres and transplanted into the defects. Histological appearance of osteochondral defects following treatment with a *lacZ* implant (left column: **A, D**), an IGF-I implant (middle column: **B, E**), and an IGF-I/FGF-2 implant (right column: **C, F**) stained with safranin O. Images (**D-F**; 40x) are magnified views of **A** through **C** (20x), illustrating the area of integration between the repair tissue (on the left side of **D-F**) with the adjacent normal articular cartilage (on the right side of **D-F**). Implants remained in a subchondral location and are visible at the bottom of images (**A, B**). Transplantation of the cotransfected IGF-I/FGF-2 implants accelerated the formation of the subchondral bone and improved articular cartilage repair in a magnitude that was larger than with IGF-I alone or when compared to *lacZ* implants after 3 weeks *in vivo*.

are generally transplanted,^{21,96,103,104,106,111,113} fibroblasts,^{27,114} perichondrial,¹⁰⁵ periosteal,^{108,112} or muscle-derived cells¹⁰⁹ have been also applied. The data from these studies showed that transgenes can be expressed in cartilage defects via *ex vivo* strategies, remaining active for about 1 month. This is significantly longer compared with the application of recombinant proteins (Table 2). Figure 2 depicts improvements in the repair of osteochondral defects following combined gene transfer of IGF-I and FGF-2 compared with the application of a marker gene (*lacZ*) to NIH 3T3 fibroblasts.¹¹⁴

Periosteal cells transduced by a BMP-7 retroviral vector and attached to a polyglycolic acid scaffold improved cartilage repair at 8 and 12 weeks *in vivo*. Interestingly, this was the first study in which a growth factor gene was transferred into a focal defect.²⁵ Since, many reports described the use of a variety of therapeutic genes like BMP-2, BMP-7, IGF-I, FGF-2, and TGF- β .^{22,90,91,114-120} Significant improvement in articular cartilage repair was noted in these reports (Table 2). Although most of the evaluations were carried out in small animal models, Hidaka *et al.*²¹ and, more recently, Goodrich *et al.*¹²¹ performed arthroscopic implantation of chondrocytes genetically engineered by adenoviral transduction with the BMP-7²¹ or IGF-I¹²¹ gene in horses.

On the basis of such encouraging data, cartilage repair was addressed by matrix-supported implantation of genetically engineered mesenchymal stem cells (MSC). Kuroda *et al.*¹²²

Table 2. Therapeutic Gene Transfer Studies to Articular Cartilage Defects *In Vivo*

Gene	Route	Vector	Cells	Support	Defect	Size (mm)	Animal Model	Joint	Location	Period of Evaluation		
										Min (wk)	Max (wk)	Ref.
BMP-2	<i>Ex vivo</i>	Retroviral	Chondrocytes	Fibrin	Osteochondral	3.6 Ø	Rabbit	Knee	Patellar groove	4	12	120
BMP-2	<i>Ex vivo</i>	Adenoviral	Fat/muscle grafts	-/-	Osteochondral	3.0 Ø	Rabbit	Knee	Patellar groove, medial femoral condyle	6	6	124
BMP-2, IGF-1	<i>Ex vivo</i>	Adenoviral	Perichondrial cells	Fibrin	Chondral	<1.0 Ø	Rat			3	8	19
BMP-7	<i>Ex vivo</i>	Retroviral	Periosteal cells	PGA	Osteochondral	3.0 Ø	Rabbit	Knee	Patellar groove	4	12	25
BMP-7	<i>Ex vivo</i>	Adenoviral	Chondrocytes	Fibrin	Osteochondral	15.0 Ø	Horse	Knee	Lateral trochlear ridge	4	36	21
IGF-1	<i>Ex vivo</i>	FuGENE 6	Chondrocytes	Alginate	Osteochondral	3.2 Ø	Rabbit	Knee	Patellar groove	3	14	91
IGF-1	<i>Ex vivo</i>	Adenoviral	Chondrocytes	Fibrin	Chondral	15.0 Ø	Horse	Knee	Lateral trochlear ridge	4	32	121
IL-1RA + IGF-1	Intra-articularly	Adenoviral	-/-	-/-	Chondral (with microfracture)	10.0 × 10.0 □	Horse	Knee, stifle	Distal radial carpal bone, medial femoral condyle	16	16	86
IGF-1 + FGF-2	<i>Ex vivo</i>	FuGENE 6	NIH 3T3	Alginate	Osteochondral	3.2 Ø	Rabbit	Knee	Patellar groove	3	3	114
FGF-2	<i>Ex vivo</i>	FuGENE 6	Chondrocytes	Alginate	Osteochondral	3.2 Ø	Rabbit	Knee	Patellar groove	3	14	90
FGF-2	<i>In vivo</i>	rAAV	-/-	-/-	Osteochondral	3.2 Ø	Rabbit	Knee	Patellar groove	1	18	36
FGF-2	<i>Ex vivo</i>	rAAV	Chondrocytes	Type I collagen, periosteal flap	Osteochondral	5.0 Ø	Rabbit	Knee	Patellar groove	4	12	99
FGF-2	<i>In vivo</i>	rAAV	-/-	-/-	Osteochondral	5.0 Ø	Rabbit	Knee	Patellar groove	4	12	163
TGF-β	<i>Ex vivo</i>	Retroviral	NIH3T3	-/-	Chondral	3.0 × 6.0 □	Rabbit	Knee		1	6	27
TGF-β	<i>Ex vivo</i>	rAAV	hMSC	-/-	Osteochondral	1.5 Ø	Rat (athymic)	Knee	Patellar groove	4	12	38
TGF-β1	<i>Ex vivo</i>	Adenoviral	Bone marrow aspirate	-/-	Chondral	6.2 Ø	Sheep	Knee	Medial femoral condyle	26	26	125
CDMP1 (GDF-5)	<i>Ex vivo</i>	FuGENE 6	MSC	Type I collagen	Osteochondral	4.0 Ø	Rabbit	Knee	Patellar groove	2	8	66

Note: PGA = polyglycolic acid; MSC = mesenchymal stem cells; Ø = cylindrical defect; □ = rectangular defect.




Figure 3. Direct rAAV-mediated gene transfer to rabbit meniscus explants *in vitro* using an rAAV-*lacZ* (left panel) or rAAV-hFGF-2 vector (right panel) (50 μ L each vector). Persistent transgene expression after 10 days *in vitro* in meniscal explants following immunohistochemical detection of *lacZ* (A), while no signal is present in the control (B). Direct transduction of a rabbit meniscal explant with rAAV-hFGF-2 results in an increased cell density (D) compared with the control (C), indicative of the mitogenic effect of FGF-2 on meniscal fibrochondrocytes. (C, D) hematoxylin and eosin/fast green. All magnifications, 20 \times .

implanted BMP-4-transduced MSCs using fibrin glue in full-thickness cartilage defects in the trochlear groove of rabbit femurs. After 24 weeks, histological scoring of the defects revealed significantly better cartilage repair in the BMP-4 treatment group compared with defects receiving *lacZ*-transduced MSCs. Guo *et al.*¹²³ seeded TGF- β 1-engineered MSCs onto poly-L-lysine-coated polylactide scaffolds *in vitro* and allografted them into full-thickness defects in rabbits. This resulted in improved joint repair with regard to extracellular matrix formation, reconstitution of the subchondral bone, and inhibition of inflammatory immune responses. Repair of osteochondral defects was also enhanced by transplantation of MSCs transfected with the CDMP1 gene, applying a lipofection method.⁶⁶

A novel method of gene therapy for the repair of osteochondral defects has recently been published by Evans *et al.*¹²⁴ Rather than genetically modifying isolated cells, this technique describes gene transfer to biopsies of muscle and fat. An adenovirus vector carrying cDNA encoding human BMP-2 was used for genetic engineering of tissues.

These gene-activated muscle or fat pads were transplanted into osteochondral defects in rabbits. Histological analysis after 6 weeks revealed the formation of a proteoglycan-rich articular surface with subchondral bone beneath and good union with the adjacent cartilage.

Ivkovic *et al.*¹²⁵ used autologous bone marrow, transduced *ex vivo*, with adenoviral vectors containing the cDNA for TGF- β 1. Implantation of the marrow clot improved the histological, biochemical, and biomechanical parameters of partial-thickness chondral defects in sheep at 6 months.

Osteoarthritis

Osteoarthritis (OA) is the leading, most disabling human condition and prevalent form of arthritis (80%), impairing the quality of life of millions of people worldwide. OA is a chronic disorder of diarthrodial joints, mainly characterized by a slow, gradual deterioration of the articular cartilage that remains without effective treatment to date. OA not only affects the cartilage but also the subchondral bone and, to a minor degree, the synovial lining, ligaments, tendons, and muscles. Current options to manage OA, such as pharmacological therapy and reconstructive surgical interventions, do not allow for the restoration of a native cartilage. OA is a complex disorder characterized by an activation of inflammatory cascades at the molecular level, leading ultimately to cartilage breakdown, associated with alterations of the phenotype of chondrocytes and a loss of the major components of the cartilage matrix. Under mechanical or biochemical stress (presence of IL-1 and TNF- α , NO, prostaglandins, matrix degradation products), the chondrocytes undergo pathological changes in their gene expression patterns that lead to an impairment of the overall homeostasis, with diminished production of normal cartilage matrix molecules (proteoglycans, type II collagen), enhanced production of matrix-degrading enzymes (MMPs and adamalysins, including ADAMs and ADAMTs), and decreased responsiveness to reparative stimuli, ultimately leading to the degradation of the matrix and cell senescence and apoptosis (NO, Fas/FasL signaling) by alteration of cell viability.

Gene Transfer *In Vitro*. Target cells in the joint include cells of the synovial lining, chondrocytes, chondroprogenitor cells, and surrounding tissues (bone, muscle, tendons, ligaments, meniscus). Application of nonviral,^{12,79,90,91,126-135} adenoviral,^{79,126,136-154} or retroviral vectors^{25,79,126,136,140,155-159} has been achieved in these cell types with more or less success. Instead, RAAV vectors are potent alternatives as they can efficiently and durably transduce synoviocytes,¹⁶⁰⁻¹⁶⁴ chondrocytes,^{36,87,99,165-168} MSCs,^{36,38,169,170} and cells of surrounding tissues.^{126,171-177}

Regeneration of a normal structural and functional cartilage might be achieved by the following:

1. inhibiting inflammatory and catabolic pathways,
2. stimulating anabolic pathways to rebuild the matrix,
3. impeding cell senescence,
4. avoiding the pathological formation of osteophytes,
5. prevention of apoptosis, and
6. influencing several of these processes.

Inhibition of catabolic pathways has been achieved *in vitro* by expressing inhibitors of matrix-degrading enzymes (tissue inhibitor of metalloproteinases, i.e., TIMP),^{178,179} inhibitors of proinflammatory cytokines (IL-1Ra, the soluble receptors sIL-1R or Soluble Tumor Necrosis Factor Receptor),^{107,137,150,155,162} and chondroprotective cytokines (IL-4, IL-10).^{160,180,181} Activation of anabolic processes *in vitro* has been noted by single or combined administration of components of the cartilage matrix or of the enzymes that synthesize them,^{182,183} of growth factors and receptors (IGF-I, FGF-2, BMPs, TGF- β),^{36,127,131,138,143-145,147,150,151,181,184,185} and of transcription factors (SOX family of DNA-binding proteins, i.e., SOX5, SOX6, SOX9).^{130,136,149,159,167,168} Restoration of cell vitality and activation of proliferation *in vitro* have been achieved by application of IGF-I and FGF-2,^{36,90,127,131,168,185} telomerase (hTERT),¹⁸⁶ of inhibitors of apoptosis (bcl-2),¹⁸⁷ or of HSP70.¹³² Interestingly, approaches that influence several of these processes have been also successfully attempted, like combining the transfer of inhibitors of catabolism pathways and of activators of anabolic events (IGF-I/IL-1Ra or IGF-I/IL-4),^{150,151,181} as well as that of activators of anabolic and proliferative processes (FGF-2/SOX9 or FGF-2/IGF-I).¹⁶⁸

In Vivo Direct Gene Transfer. The key issue in establishing an efficient therapy against OA is the accessibility of the targets to the treatment when they reside in the joint cavity. The following approaches have been developed to deliver a molecular composition:

1. systemic delivery, and
2. intra-articular administration (via injection or arthroscopy).

Systemic approaches are better suited to target diseases that are systemic in nature like rheumatoid arthritis (RA).^{164,188-190} Local administration of components might be preferable in the case of OA that affects only a limited number of joints without major extra-articular or systemic manifestations. The foreign material may be delivered directly (gene vector preparation) or indirectly (genetically modified cells).

Several lines of evidence have demonstrated that intra-articular injection of most vector types leads to a preferential transduction of the synovium,^{29,79,81,191} being more suited for strategies aiming at inhibiting inflammatory and catabolic pathways and a common approach employed against experimental RA. Successful attempts towards

these goals have been reported by direct application of vectors coding for IL-4,^{35,192} IL-10,^{193,194} sTNFR alone¹⁶² or combined with IL-10,¹⁹⁵ IL-1Ra alone^{18,20,34,155,196,197} or combined with sTNFR,²⁰ antagonists and inhibitors of TGF- β and of the BMPs,¹⁹⁸ HSP70,¹³² gene expression silencers,¹⁹⁹ and kallistatin or thrombospondin-1.^{200,201}

Yet, even if cartilage breakdown can be contained, this will not be sufficient to fully compensate for the loss of matrix elements and cells noted during the disease progression. In this regard, increased synthesis of cartilage matrix components has been documented following injection of vectors carrying genes for anabolic factors (IGF-I).²⁰²

Ex Vivo Indirect Gene Transfer. Although more complex, *ex vivo* gene therapy is considered safer because no free vector particles are introduced in the body. Modified cells can be extensively controlled, tested, and selected while maintained in culture. Administration of cells is also a means to increase the cellularity like needed for severe OA.

Synoviocytes have been predominantly employed to deliver inhibitors of inflammatory and catabolic processes.²⁰³⁻²⁰⁸ Such pathways could be regulated by injecting synoviocytes transduced to overexpress an IL-1Ra alone²⁰³⁻²⁰⁸ or combined with IL-10.²⁰⁸ Also, dermal fibroblasts have been modified for this purpose to overexpress an IL-1Ra, sTNFR, or a combination of both.²⁰⁹

Reduced severity of the induced arthritis was associated with a decrease in cartilage breakdown, but complete resurfacing was not achieved. Successful attempts to promote the formation of new cartilage have been made by administrating dermal fibroblasts modified to express BMP-2.¹⁴⁶

Still, preparation of terminally differentiated cells from unaffected sites remains invasive, with a limited supply, and represents an additional burden for the patient. Also, committed cells generally undergo major phenotypic changes upon passaging in culture, especially chondrocytes. Multipotent cells might be more suited for transplantation purposes, possibly leading to the production of a cartilage surface of enhanced quality compared with committed cells that lead to the formation of a poorly differentiated fibrous cartilage. Progenitor cells can be easily isolated from multiple tissues (bone marrow, periosteum, perichondrium, muscle, fat, subdermis, cartilage, bone, synovial membrane, ligaments), even in OA patients, maintaining a multilineage potential with a reliability for differentiation and a capacity for expansion.^{210,211} Indeed, injection of muscle-derived stem cells modified by combined gene transfer of BMP-4 with sFlt1 (a vascular endothelial growth factor (VEGF) antagonist) allowed for cartilage repair in a rat model of OA.^{212,213}

Osteonecrosis

Osteonecrosis (ON) is primarily a disease of the subchondral bone that secondarily affects the articular cartilage.

Initially, a vascular insult is thought to cause an interference of the microcirculation of the subchondral bone, resulting in an edema that leads to an increased intraosseous pressure. This leads to ON of the affected segment of the subchondral bone, which may result in a subchondral insufficiency fracture, destabilizing the overlying articular cartilage and eventually resulting in its collapse and the creation of an osteochondral defect. Treatment options consist of conservative therapy in early stages. Precollapse lesions can be treated with retrograde core decompression, while later-stage lesions presenting with osteochondral defects require osteochondral transplants and/or osteotomies, or ultimately, partial or total knee arthroplasty.²¹⁴

Possible experimental gene therapy approaches need to be stage dependent, focusing on early stages (when the articular cartilage is not compromised) at the revascularization of the necrotic bone, while at the stage of osteochondral lesion, only gene-enhanced osteochondral transplants might be useful. Katsube *et al.*²¹⁴ applied gene transfer of VEGF, to accelerate revascularization of the necrotic bone. Using an adenoviral vector encoding for VEGF, endothelial cells of the rabbit saphenous arteries were transduced. These gene-modified arteries were then placed with its venae comitantes into necrotic iliac crest bone *in vivo*. Angiogenesis in the necrotic bone was quantified by bone blood flow measurement and assessment of vessel density following microangiography. The extent of neoangiogenesis was significantly greater in the VEGF group than the control group, reflected in an increased capillary density, length of newly formed capillaries, and increased bone blood flow at 1 week postoperatively. While this study was restricted to the bone of the iliac crest, it might serve as a paradigm for the treatment of ON in a subchondral location. Such a therapy may allow the healing of avascular necrosis before fracture and subchondral collapse occur, preventing the articular cartilage from damage. More studies with time points longer than the 1-week evaluation are needed, preferentially performed in animal models of subchondral ON, such as the femoral condyles of the knee joint, its second most common location.

Osteochondritis Dissecans

Osteochondritis dissecans (OCD) usually affects children and young adults and occurs mainly in the knee joint, characteristically in the lateral aspect of the medial femoral condyle. Possible etiological factors beside a genetic predisposition include ischemia and epiphyseal abnormalities with subsequent necrosis. For example, disruption of epiphyseal plate vessels may lead to localized avascular necrosis. Its revascularization usually occurs with the formation of a scar tissue, absorption of necrotic fragments, intertrabecular osteoid deposition, and remodeling with new

bone formation. When revascularization is delayed, an OCD lesion can occur. Clinical treatment principles focus on stimulation of revascularization or removal of necrotic subchondral bone together with its restoration (e.g., using autologous bone transplants), beside the surgical fixation of an unstable osteochondral fragment.²¹⁴

So far, no experimental gene-based treatment has been proposed for the treatment of OCD. In theory, the same principles apply for the revascularization of necrotic subchondral bone, as already outlined for ON with subsequent articular cartilage defects. It may be also possible to enhance the surgical fixation of an osteochondral fragment by applying osteoinductive genes such as the BMPs to the subchondral bone–osteochondral fragment interface to improve integration of the osteochondral fragment. It is unclear whether the integration of a chondral fragment may be achieved, a rare indication currently favored only for surgical refixation of large fragments in juvenile patients.²¹⁵ Likewise, gene-modified osteochondral transplants may be applied at later stages of deep osteochondral defects.

Meniscal Fibrocartilage

Anatomy, Function, and Pathophysiology

The menisci are semilunar fibrocartilage structures that transmit weightbearing forces and increase stability, facilitate nutrition and provide lubrication for the articular cartilage, and promote knee proprioception.^{216,217} As the medial meniscus is less mobile during joint motion,^{218,219} injuries are much more common compared to the lateral meniscus.²²⁰ Type I collagen is the predominant collagen of the meniscal tissue.²²¹ It is arranged with a circumferential orientation with interspersed radially oriented fibers.²²² The central parts of the menisci are mainly constituted of fibrochondrocytes, whereas fibroblasts are the predominant cell type in the peripheral regions.²²³ Meniscal blood supply is restricted to the peripheral 10% to 25% of the meniscal tissue.^{224,225} Nourishment in the central area is provided only by diffusion of the synovial fluid,²²⁶ perhaps playing a role in the poor healing capacity of central lesions.²²⁵⁻²²⁷ Gene transfer strategies may be applied for the following:

1. meniscal repair, and
2. meniscal reconstruction, using
 - 2.1. meniscal substitutes, and
 - 2.2. meniscal allografts.

Meniscal Repair

Meniscal tears are common^{228,229} and predispose the affected joint to develop secondary OA.²³⁰ Tears of the meniscus in the vascularized peripheral parts can be

repaired by sutures, while tears of the central avascular parts are treated by arthroscopic partial meniscectomy.

Gene Transfer Strategies: In Vitro Studies. Gene transfer strategies for the repair of meniscal tears focus on the delivery of therapeutic agents, for example, growth factors, to the site of the meniscal lesion. This can be performed either via direct application of gene vectors or by transplantation of genetically modified cells overexpressing therapeutic genes. Treatment of meniscal fibrochondrocytes with recombinant growth factor proteins such as the platelet-derived growth factor AB (PDGF-AB),²³¹⁻²³⁴ FGF-2,^{177,235-237} IGF-I,^{235,238-240} TGF- β 1,^{152,233,239,241,242} BMP-7,²³³ or TGF- β 3²³⁵ has been shown to improve the phenotypical and biochemical properties of the cells *in vitro*. Fibrochondrogenesis of stem cells is enhanced by incubation with growth factors such as TGF- β 1²⁴³ or TGF- β 3 in combination with BMP-4.²⁴⁴ The possible application of gene transfer strategies in meniscal repair has first been investigated by Goto *et al.*¹³⁹ The *lacZ* marker gene was transferred to meniscal cell cultures using retroviral and adenoviral vectors. In a next step, the marker gene was applied to human meniscal fragments and whole lapine menisci using direct adenoviral gene transfer and transplantation of meniscal fibrochondrocytes transduced with a retroviral vector. Transgene expression was detected in meniscal explants following *ex vivo* gene transfer for at least 20 weeks. Successful transfer of the *lacZ* marker gene was also achieved by rAAV-mediated transfer into human and lapine fibrochondrocytes *in vitro*.¹⁷³ Encouraged by these findings, in 2000, the group of Chris Evans transferred the gene encoding for TGF- β 1¹⁵⁸ to meniscal cells *in vitro*, resulting in enhanced synthesis of proteoglycans and collagen. Zhang *et al.* used a lipid-based gene transfer system to deliver the gene encoding for human IGF-I to meniscal fibrochondrocytes, yielding accelerated proliferation and differentiation of the modified cells.¹³⁴ Recently, we tested the hypothesis that overexpression of FGF-2 through rAAV vectors leads to detectable metabolic changes in human meniscal fibrochondrocytes and inside defects of human meniscal explants.¹⁷⁷ Application of the rAAV-hFGF-2 vector allowed for enhanced cell proliferation and survival *in vitro* (Figure 3). The idea of applying gene therapy protocols to deliver fibrochondrogenic agents to meniscal tears was supported by a significant reduction of the amplitude of meniscal tears after FGF-2 treatment in this study.¹⁷⁷

Gene Therapy: In Vivo Studies. Only few reports have evaluated the feasibility of gene therapy strategies to enhance the repair of meniscal tears *in vivo*. Experimental studies have shown that repair in the central part of the meniscus can be promoted by various chemotactic and mitogenic stimuli delivered by an autologous fibrin clot^{245,246} or a free graft of synovium^{247,248} *in vivo*. In a sheep model, longitudinal tears of the anterior horn of the medial meniscus were

sutured using VEGF-coated sutures. Interestingly, meniscal repair was not enhanced in the VEGF treatment group.^{249,250}

In 1999, methods of direct and indirect gene transfer to meniscal lesions were compared.¹³⁹ In a lapine model, a suspension of adenoviral vectors carrying the *lacZ* marker gene was mixed with whole blood, and the clot was inserted into 2-mm-long incisions in the medial meniscus. In the same study, using a canine model, retrovirally transduced allogenic meniscal fibrochondrocytes carrying the *lacZ* gene were embedded in collagen gels and transferred to partial-thickness circular defects (depth, 3 mm; diameter, 2 mm) in the medial meniscus. Gene expression persisted for at least 3 weeks in the lapine model but for 6 weeks within the transplanted meniscal fibrochondrocytes in the canine model. In another animal study,¹⁷³ longitudinal incisions were created in the avascular zone of the medial meniscus of rabbits. When rAAV-*lacZ* constructs were injected intralesionally, X-Gal staining was present by day 20 postoperatively, the longest time point evaluated.

Meniscal Reconstruction

Meniscal Substitutes. Meniscal substitutes have been proposed as a means to overcome problems associated with meniscal allografts and to promote meniscal repair of segmental defects, for example, resulting from a partial meniscectomy.^{251,252} Meniscal substitutes already in clinical use are based on porous matrices of type I collagen/glycosaminoglycan (Menaflex, ReGen Biologics, Hackensack, NJ)^{253,254} or polyurethane (Actifit, Orteq, London, UK).^{255,256}

The feasibility of genetic engineering of meniscal fibrochondrocytes has already been described above. However, in the treatment of circumscribed meniscal defects, direct gene vector administration into injured knee joints may be difficult to achieve because a loss of the bradytrophic meniscal tissue may hardly be restored by local cells, even after administration of mitogenic and anabolic genes. Therefore, gene therapy in the treatment of meniscal defects may need to be used in combination with the transplantation of modified cells or tissues.

Tissue engineering involves the combination of cells, engineered extracellular matrices, and biologically active molecules for tissue regeneration.^{257,258} Over the last 2 decades, numerous tissue engineering strategies have emerged for the replacement of meniscal tissue.²⁵⁹⁻²⁶¹ In general, 2 basic approaches for meniscal replacement can be distinguished:

1. application of acellular matrices^{262,263} versus
2. application of cell-seeded matrices.^{264,265}

Several concepts for treating circumscribed meniscal defects concentrate on meniscal replacement by acellular matrices,^{259,266-269} avoiding possible risks associated with transplantation of human allografts (e.g., failure rate,

immunoreaction,²⁷⁰ disease transmission²⁷¹). Different types of meniscal substitutes, such as decellularized allogenic and xenogenic grafts,^{262,263,272,273} collagen grafts,^{253,274} permanent synthetic scaffolds,²⁵¹ and biodegradable scaffolds based on small intestine submucosa,²⁷⁵⁻²⁷⁸ poly-lactic acid (PLA), or poly-glycolic acid (PGA),²⁷⁹⁻²⁸² have been used in experimental and clinical studies. However, after transplantation of acellular meniscal constructs into defects, the transplants are populated by synovial fibroblasts, resulting in a scar tissue with poor biomechanical properties.^{245,283} Therefore, some tissue engineering approaches focus on additional cell-seeding techniques prior to transplantation.^{251,284} Meniscal cells,^{282,285} articular chondrocytes,^{286,287} synovial fibroblasts,²⁸⁸ and MSC²⁸⁹ have been proposed as potential cell sources and have been cultivated *in vivo* and *in vitro* on various matrices.²⁶⁷ In addition, different environmental factors such as growth factors have been used to optimize cell proliferation *in vitro*.²⁹⁰

Gene therapy may aid to further enhance the fibrochondrogenic potential of tissue-engineered transplants. In 2002, Hidaka *et al.*²⁹¹ applied a gene transfer protocol to enhance the vascularization and blood supply of cell-seeded bioengineered meniscus transplants. Bovine meniscal cells overexpressing hepatocyte growth factor (HGF) were seeded onto PGA scaffolds and transplanted subcutaneously in athymic nude mice for 8 weeks. Ink injection studies showed that HGF-treated meniscal cells formed a tissue that contained significantly more blood vessels than the controls. In another preliminary *ex vivo* study, Steinert *et al.*¹⁵² transduced primary meniscus cells and bone marrow-derived MSCs with adenoviral vectors encoding for marker genes or TGF- β 1. Modified cells were seeded in type I collagen-glycosaminoglycan (GAG) matrices and transplanted into defects of bovine menisci explants. *In vitro*, the vectors efficiently transduced meniscal cells and MSCs, and transgene expression remained elevated after incorporation of the cells into matrices. Transfer of TGF- β 1 increased the fibrochondrogenic potential of modified cells, and transplantation of the TGF- β 1-transduced constructs resulted in satisfactory filling of the lesions *ex vivo* (Table 3).

A recent *in vivo* work on the use of gene transfer to enhance meniscal repair has been published by Zhang *et al.*²⁹² Following an indirect gene therapy approach without tissue engineering features, the authors created full-thickness meniscal defects in the avascular area of the anterior horn of the medial meniscus in a goat model. Bone marrow stromal cells were transfected with the gene encoding for human IGF-I using a nonviral transfection system (FuGENE 6) and suspended in calcium alginate prior to injection into the meniscal defects. After 16 weeks, the resulting repair tissue was improved according to MRI and histological and biochemical evaluation and compared with the controls (Table 3). **Meniscal Allografts.** Meniscal reconstitution with allografts²⁹³⁻³⁰⁵ is a therapeutic option especially for young and

symptomatic patients with a history of lateral meniscectomy in a normally aligned, stable joint without severe degenerative changes of the articular cartilage. A recent review³⁰⁶ suggests that meniscal allograft transplantation improves pain and function in the short and intermediate term.

Application of gene-based strategies has been suggested to improve remodeling of meniscal allografts.³⁰⁷ Martinek *et al.*³⁰⁸ studied the feasibility of gene transfer in lapine meniscal allografts *ex vivo* using a retroviral vector encoding the marker gene *lacZ*. Subsequently, unilateral meniscal replacements were performed with these engineered allografts. Transduced fibrochondrocytes migrated into the depth of the graft, while transgene expression persisted for up to 8 weeks. This investigation suggests potential promise for growth factor delivery in autografts and allografts prior to implantation.

Clinical Gene Therapy Trials

Preclinical data, as those described above, have encouraged the initiation of human clinical trials originally for arthritis. The first studies were based on the *ex vivo* retroviral gene transfer of a human IL-1Ra sequence in synoviocytes from patients with end-stage RA followed by reinjection of the modified cells in the metacarpophalangeal joint.^{23,82,309} The aim of these studies was to evaluate the possibility of transferring genes to human joints and expressing them intra-articularly in a safe fashion acceptable to the patients. The use of these protocols has permitted extensive testing of the cells prior to reimplantation, demonstrating successful expression of the transgene locally vis-à-vis control joints, without adverse events related to the treatment but with clinical improvements in some of the patients, encouraging the implementation of phase II studies (pending).^{1,5,30,310-315} Another protocol has been initiated for intra-articular plasmid³¹⁶ delivery of the HSV thymidine kinase gene to the synovial lining of RA patients followed by administration of ganciclovir to achieve synovial ablation,^{1,5,309,311-314} but this protocol has been closed because of a failure to recruit. A new phase I trial for RA involved the direct *in vivo* intra-articular injection of an rAAV vector carrying the sequence for a fusion protein as sTNFR on an immunoglobulin molecule (tgAAC94 protocol).³¹⁷ As the study revealed that the treatment was safe and well tolerated in subjects without use of concurrent systemic TNF- α antagonist,^{1,311,312,314,317} a phase I/II trial was subsequently started³¹⁸ with the possibility to include patients who were already taking systemic TNF blockers and the administration of a second injection of tgAAC94. As one of the participants who was simultaneously being treated with systemic TNF antagonist and other immunosuppressive medications died after receiving the second injection, the trial was placed on hold by the U.S. Food and Drug Administration (FDA) to investigate, in parallel with the Recombinant DNA Advisory Committee

Table 3. Therapeutic Gene Transfer Studies to Meniscal Cells *In Vitro* or Meniscal Tissue *Ex Vivo* and *In Vivo*

Gene	Strategy	Vector	Cells	Support	Experimental Model	Period of Evaluation		Major Findings	Ref.
						Min	Max		
IGF-1	<i>In vitro</i>	Liposome (FuGENE 6)	Meniscal fibrochondrocytes (human)	-/-	-/-	0 h	10 d	Transfection efficiency 16% ± 1.2% No cytotoxicity Decrease in PDT from 52.6 to 40.2 h	134
	<i>In vitro</i>	Liposome (FuGENE 6)	Bone marrow stromal cells (goat)	Calcium alginate gel	Goat	4 h	10 d	<i>In vitro</i> Transfection efficiency 22.0% ± 2.4% Elevated IGF-1 secretion (3.28 v. 1.67 ng/mL by untransfected fibrochondrocytes)	292
	<i>In vivo</i>								
HGF	<i>In vitro</i>	Adenoviral	Meniscal cells (calf)	PGA	Athymic nude mice (subcutaneous pouch)	4 wk	16 wk	<i>In vivo</i> Macroscopically and histologically improved repair tissue Elevated GAG content (14.2 v. 13.7 mg/g by untransfected BMSCs) Improved aspect of repair tissue on MRI	291
	<i>In vivo</i>					3 d	8 wk	<i>In vitro</i> Transduction efficiency N.D. No enhanced proliferation of HGF-transduced cells <i>In vivo</i> HGF expression detectable for ≥2 weeks Fibrocartilage with structural limitations (presence and organization of collagen fibrils) Enhanced vascularization of engineered constructs No improved biomechanical properties (compression testing)	
FGF-2	<i>In vitro</i>	rAAV	Meniscal fibrochondrocytes (human)	-/- (direct vector injection)	Human explants	0 d	21 d	<i>In vitro</i> Transduction efficiency 53%-59% Efficient FGF-2 transgene expression Enhanced cell proliferation and survival	177
	<i>Ex vivo</i>					5 d	15 d	<i>Ex vivo</i> Enhanced contractile markers (α-SMA) Reduction of meniscal tear amplitude in depth and width (up to 2.4-fold) No stimulation of extracellular matrix components (type I/II collagen, PG)	
TGF-β1	<i>In vitro</i>	Adenoviral	Primary meniscal cells (calf); bone marrow-derived MSCs (calf)	Type I collagen-GAG matrix	Bovine explants	3 d	3 wk	<i>In vitro</i> Transduction efficiency >75% Increased cellularity and GAG/DNA synthesis Enhanced proteoglycan and type II collagen staining Enhanced meniscal gene expression (COL I, COL II, DCN, BCN)	152
	<i>Ex vivo</i>					3 wk		<i>Ex vivo</i> Formation of highly cellular repair tissue; no differences between treatment and control group Transduction efficiency N.D. Increased transgene expression Enhanced collagen and proteoglycan synthesis (up to 15-fold)	
TGF-β1	<i>In vitro</i>	Retroviral	Meniscal cells (human and canine)	-/-	-/-	48 h			158

Note: PDT = population doubling time; GAG = glycosaminoglycan; BMSC = bone marrow stromal cell; HGF = hepatocyte growth factor; PGA = polyglycolic acid; α-SMA = alpha-smooth muscle actin; PG = proteoglycan; COL I = type I A2 collagen (537 bp); COL II = type II A1 collagen (580 bp); DCN = decorin (400 bp); BCN = biglycan (165 bp); MSC = mesenchymal stem cells; N.D. = not determined. FGF = fibroblast growth factor; TGF = transforming growth factor

Recombinant DNA Advisory Committee (RAC), the circumstances of the demise of the patient. The death was apparently due to a disseminated infection with *Histoplasma capsulatum*, a fungus endemic in the region of origin of the volunteer, and to an immunosuppression.^{312,319-321} Indeed, known serious complications of the particular TNF antagonist are susceptibility to *H. capsulatum*. The most probable explanation is that the subject was already infected with the fungus when receiving the second injection of tgAAC94. As the committee felt that the gene therapy protocol was very unlikely to have played any significant role in the event based on a large body of data from the independent investigations and since rAAV has been used safely in 47 previous human gene therapy clinical trials, the evaluation has been reopened with some modifications (exclusion of patients with elevated temperature, localized symptoms, fatigue, or with history of opportunistic infection), requiring additional monitoring (repeated blood counts, serum chemistry, vector DNA and transgene product titration, analysis of T-cell responses to AAV), as a possible role of the gene transfer in this course has not been definitely excluded (presence of neutralizing antibodies to the AAV capsid, occasional detection of vector genomes in the blood at the highest vector dose). Regarding OA, a phase I protocol is currently ongoing, based on an *ex vivo* approach using the retroviral transfer of TGF- β .³¹²

Gene Doping

Although the previously discussed gene-based approaches may have potential value for the treatment of articular cartilage defects and meniscal lesions, some of the therapeutic genes used in these studies have been also implicated for gene doping,³²² a term referring to the potential misuse of gene therapy for the purposes of enhancing athletic performance.³²³⁻³²⁵ Possible genes with such potential include, but are not limited to, growth hormone and IGF-I,³²⁶ erythropoietin (Epo),³²⁷ VEGF,³²⁸ FGF-2, and endorphins.³²⁹

IGF-I, the prime target of growth hormone action, is a potential candidate gene. A number of studies have shown that upregulation of IGF-I stimulates muscle growth and improves muscle function.³²⁶ Interestingly, this increase in muscle volume is not reflected by detectable increases in circulating IGF-I. While favorable responses have been obtained in animal studies, the transfer of such techniques to humans with the goal of a higher performance still presents many technical challenges.

The hormone Epo is produced by the peritubular capillary endothelial cells in the kidney. Under hypoxic conditions, Epo is produced and secreted, increasing the production of red blood cells. Eero Mäntyranta, a Finnish cross-country skier who won 2 gold medals in the 1964 Olympics, was born with a mutation in the Epo receptor gene that allowed his blood to carry significantly more oxygen than an average person.³³⁰ Recombinant Epo has

been used already as a performance-enhancing drug. Because of differences in its peptide sequence compared with the endogenous protein, it may be detected in blood. Recently, a viral vector for the release of Epo in response to low oxygen concentrations has been developed under the trade name Repoxygen (Oxford BioMedica, Oxford, UK). The viral vector of undisclosed origin carries the human Epo gene under the control of a hypoxia control element (HRE). At low oxygen concentrations, HRE switches on the expression of the transgene. The vector is designed to be delivered by a simple intramuscular injection, resulting in the synthesis of recombinant Epo by muscle cells, rather than by cells of the liver or kidneys. Initially developed to treat anemia, there have been speculations in the media that it has been already applied for doping purposes.³³¹

Recently, genetically engineered mice have been created with an alteration in energy metabolism based on overexpression of the gene for phosphoenolpyruvate carboxylases (PEPCK-C). PEPCK-C is an enzyme of the lyase family that plays a role in the metabolic pathway of gluconeogenesis, converting oxaloacetate into phosphoenolpyruvate and carbon dioxide. These transgenic PEPCK-C mice carry a chimeric gene in which a copy of the cDNA for PEPCK-C is placed under control of the skeletal actin gene promoter, directing overexpression of PEPCK-C exclusively to skeletal muscle. PEPCK-C mice were more active, could run longer and faster, and used fatty acids more efficiently and produced far less lactate than control animals.³³² Whether these data can be corroborated by studies in large animals remains to be determined.

Taken together, there is an emerging body of results from a number of transgenic and somatic gene transfer studies that suggest the principle of gene transfer may find application to enhance athletic performance. Many of the genes are already cloned in functional vectors, and some of them are being evaluated in clinical trials for the treatment of diseases. However, therapeutic gene transfer to humans is still technically challenging, and no clear evidence has been given that athletes have been using gene technology to enhance their performance. For antidoping authorities, the challenge will be to detect these endogenously produced gene products because of the homology between the transferred cDNA, the homology of the endogenously produced protein, and the limited specificity of indirect detection procedures.³³³ Further studies in this field are needed since a possible uncontrolled use of these gene vectors imposes potential high risks for both the athlete and the general public.

Outlook

Despite these encouraging data, application of gene transfer approaches in the treatment of articular cartilage and meniscal lesion tears is still in its infancy. Although the use

of gene therapy holds great promise, issues that need to be addressed include the duration of transgene expression, further studies in clinically relevant animal models of articular cartilage and meniscal lesions, the benefit of using *ex vivo* genetically modified cells versus direct gene transfer approaches, and the identification of (an) optimal therapeutic factor(s) for each particular clinical problem. Future studies will also have to shed light on the safety of these approaches regarding the nonlethal nature of these diseases. A successful application of gene therapy for cartilage repair requires the combined effort of orthopedic surgeons continuing to ask clinically relevant questions and of basic scientists further improving the currently available gene transfer systems.

Acknowledgments and Funding

The authors received no financial support for the research and/or authorship of this article.

Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the authorship and/or publication of this article.

References

- Evans CH, Ghivizzani SC, Herndon JH, Robbins PD. Gene therapy for the treatment of musculoskeletal diseases. *J Am Acad Orthop Surg*. 2005;13(4):230-42.
- Nakajima A. Application of cellular gene therapy for rheumatoid arthritis. *Mod Rheumatol*. 2006;16(5):269-75.
- Madry H, Kohn D, Cucchiariini M. [Gene therapy in orthopaedic surgery]. *Orthopade*. 2006;35(11):1193-202.
- Ivkovic A, Pascher A, Hudetz D, Jelic M, Haspl M, Windhager R, *et al*. Current concepts in gene therapy of the musculoskeletal system. *Acta Chir Orthop Traumatol Cech*. 2006;73(2):115-22.
- Robbins PD, Evans CH, Chernajovsky Y. Gene therapy for arthritis. *Gene Ther*. 2003;10(10):902-11.
- Abramson SB, Amin A. Blocking the effects of IL-1 in rheumatoid arthritis protects bone and cartilage. *Rheumatology (Oxford)*. 2002;41(9):972-80.
- Evans CH, Ghivizzani SC, Kang R, Muzzonigro T, Wasko MC, Herndon JH, *et al*. Gene therapy for rheumatic diseases. *Arthritis Rheum*. 1999;42(1):1-16.
- Vervoordeldonk MJ, Tak PP. Gene therapy in rheumatic diseases. *Best Pract Res Clin Rheumatol*. 2001;15(5):771-88.
- Burstein H. Gene therapy for rheumatoid arthritis. *Curr Opin Mol Ther*. 2001;3(4):362-74.
- Felgner PL, Gadek TR, Holm M, Roman R, Chan HW, Wenz M, *et al*. Lipofection: a highly efficient, lipid-mediated DNA-transfection procedure. *Proc Natl Acad Sci U S A*. 1987;84(21):7413-7.
- Schwendener RA. Liposomes in biology and medicine. *Adv Exp Med Biol*. 2007;620:117-28.
- Orth P, Weimer A, Kaul G, Kohn D, Cucchiariini M, Madry H. Analysis of novel nonviral gene transfer systems for gene delivery to cells of the musculoskeletal system. *Mol Biotechnol*. 2008;38(2):137-44.
- Hudde T, Rayner SA, Comer RM, Weber M, Isaacs JD, Waldmann H, *et al*. Activated polyamidoamine dendrimers, a non-viral vector for gene transfer to the corneal endothelium. *Gene Ther*. 1999;6(5):939-43.
- Godbey WT, Wu KK, Hirasaki GJ, Mikos AG. Improved packing of poly(ethylenimine)/DNA complexes increases transfection efficiency. *Gene Ther*. 1999;6(8):1380-8.
- Chemin I, Moradpour D, Wieland S, Offensperger WB, Walter E, Behr JP, *et al*. Liver-directed gene transfer: a linear polyethylenimine derivative mediates highly efficient DNA delivery to primary hepatocytes *in vitro* and *in vivo*. *J Viral Hepat*. 1998;5(6):369-75.
- Ravi Kumar M, Hellermann G, Lockey RF, Mohapatra SS. Nanoparticle-mediated gene delivery: state of the art. *Expert Opin Biol Ther*. 2004;4(8):1213-24.
- Graham FL, van der Eb AJ. A new technique for the assay of infectivity of human adenovirus 5 DNA. *Virology*. 1973;52(2):456-67.
- Frisbie DD, Ghivizzani SC, Robbins PD, Evans CH, McIlwraith CW. Treatment of experimental equine osteoarthritis by *in vivo* delivery of the equine interleukin-1 receptor antagonist gene. *Gene Ther*. 2002;9(1):12-20.
- Gelse K, von der Mark K, Aigner T, Park J, Schneider H. Articular cartilage repair by gene therapy using growth factor-producing mesenchymal cells. *Arthritis Rheum*. 2003;48(2):430-41.
- Ghivizzani SC, Lechman ER, Kang R, Tio C, Kolls J, Evans CH, *et al*. Direct adenovirus-mediated gene transfer of interleukin 1 and tumor necrosis factor alpha soluble receptors to rabbit knees with experimental arthritis has local and distal anti-arthritic effects. *Proc Natl Acad Sci U S A*. 1998;95(8):4613-8.
- Hidaka C, Goodrich LR, Chen CT, Warren RF, Crystal RG, Nixon AJ. Acceleration of cartilage repair by genetically modified chondrocytes over expressing bone morphogenetic protein-7. *J Orthop Res*. 2003;21(4):573-83.
- Park J, Gelse K, Frank S, von der Mark K, Aigner T, Schneider H. Transgene-activated mesenchymal cells for articular cartilage repair: a comparison of primary bone marrow-, perichondrium/periosteum- and fat-derived cells. *J Gene Med*. 2006;8(1):112-25.
- Evans CH, Robbins PD, Ghivizzani SC, Herndon JH, Kang R, Bahnson AB, *et al*. Clinical trial to assess the safety, feasibility, and efficacy of transferring a potentially anti-arthritic cytokine gene to human joints with rheumatoid arthritis. *Hum Gene Ther*. 1996;7(10):1261-80.
- Grande DA, Mason J, Light E, Dines D. Stem cells as platforms for delivery of genes to enhance cartilage repair. *J Bone Joint Surg Am*. 2003;85-A(Suppl 2):111-6.
- Mason JM, Breitbart AS, Barcia M, Porti D, Pergolizzi RG, Grande DA. Cartilage and bone regeneration using gene-enhanced tissue engineering. *Clin Orthop Relat Res*. 2000;(379 Suppl):S171-8.

26. Murphy JM, Fink DJ, Hunziker EB, Barry FP. Stem cell therapy in a caprine model of osteoarthritis. *Arthritis Rheum*. 2003;48(12):3464-74.
27. Lee KH, Song SU, Hwang TS, Yi Y, Oh IS, Lee JY, et al. Regeneration of hyaline cartilage by cell-mediated gene therapy using transforming growth factor beta 1-producing fibroblasts. *Hum Gene Ther*. 2001;12(14):1805-13.
28. Gouze E, Pawliuk R, Gouze JN, Pilapil C, Fleet C, Palmer GD, et al. Lentiviral-mediated gene delivery to synovium: potent intra-articular expression with amplification by inflammation. *Mol Ther*. 2003;7(4):460-6.
29. Gouze E, Pawliuk R, Pilapil C, Gouze JN, Fleet C, Palmer GD, et al. In vivo gene delivery to synovium by lentiviral vectors. *Mol Ther*. 2002;5(4):397-404.
30. Evans CH, Ghivizzani SC, Robbins PD. The 2003 Nicolas Andry Award: orthopaedic gene therapy. *Clin Orthop Relat Res*. 2004;(429):316-29.
31. Oligino T, Ghivizzani S, Wolfe D, Lechman E, Krisky D, Mi Z, et al. Intra-articular delivery of a herpes simplex virus IL-1Ra gene vector reduces inflammation in a rabbit model of arthritis. *Gene Ther*. 1999;6(10):1713-20.
32. Berns KI, Linden RM. The cryptic life style of adeno-associated virus. *Bioessays*. 1995;17(3):237-45.
33. Flotte TR, Afione SA, Conrad C, McGrath SA, Solow R, Oka H, et al. Stable in vivo expression of the cystic fibrosis transmembrane conductance regulator with an adeno-associated virus vector. *Proc Natl Acad Sci U S A*. 1993;90(22):10613-7.
34. Pan RY, Chen SL, Xiao X, Liu DW, Peng HJ, Tsao YP. Therapy and prevention of arthritis by recombinant adeno-associated virus vector with delivery of interleukin-1 receptor antagonist. *Arthritis Rheum*. 2000;43(2):289-97.
35. Watanabe S, Imagawa T, Boivin GP, Gao G, Wilson JM, Hirsch R. Adeno-associated virus mediates long-term gene transfer and delivery of chondroprotective IL-4 to murine synovium. *Mol Ther*. 2000;2(2):147-52.
36. Cucchiari M, Madry H, Ma C, Thurn T, Zurakowski D, Menger MD, et al. Improved tissue repair in articular cartilage defects in vivo by rAAV-mediated overexpression of human fibroblast growth factor 2. *Mol Ther*. 2005;12(2):229-38.
37. Evans CH, Ghivizzani SC, Smith P, Shuler FD, Mi Z, Robbins PD. Using gene therapy to protect and restore cartilage. *Clin Orthop Relat Res*. 2000;(379 Suppl):S214-9.
38. Pagnotto MR, Wang Z, Karpie JC, Ferretti M, Xiao X, Chu CR. Adeno-associated viral gene transfer of transforming growth factor-beta1 to human mesenchymal stem cells improves cartilage repair. *Gene Ther*. 2007;14(10):804-13.
39. Mease PJ, Wei N, Fudman EJ, Kivitz AJ, Schechtman J, Trapp RG, et al. Safety, tolerability, and clinical outcomes after intraarticular injection of a recombinant adeno-associated vector containing a tumor necrosis factor antagonist gene: results of a phase 1/2 study. *J Rheumatol*. 2010;37(4):692-703.
40. O'Driscoll SW. The healing and regeneration of articular cartilage. *J Bone Joint Surg Am*. 1998;80(12):1795-812.
41. Hunziker EB, Michel M, Studer D. Ultrastructure of adult human articular cartilage matrix after cryotechnical processing. *Microsc Res Tech*. 1997;37(4):271-84.
42. Madry H, van Dijk CN, Mueller-Gerbl M. The basic science of the subchondral bone. *Knee Surg Sports Traumatol Arthrosc*. 2010;18(4):419-33.
43. Pape D, Filardo G, Kon E, van Dijk CN, Madry H. Disease-specific clinical problems associated with the subchondral bone. *Knee Surg Sports Traumatol Arthrosc*. 2010;18(4):448-62.
44. Noyes FR, Stabler CL. A system for grading articular cartilage lesions at arthroscopy. *Am J Sports Med*. 1989;17(4):505-13.
45. Hunziker EB. Articular cartilage repair: basic science and clinical progress. A review of the current status and prospects. *Osteoarthritis Cartilage*. 2002;10(6):432-63.
46. Furukawa T, Eyre DR, Koide S, Glimcher MJ. Biochemical studies on repair cartilage resurfacing experimental defects in the rabbit knee. *J Bone Joint Surg Am*. 1980;62(1):79-89.
47. Shapiro F, Koide S, Glimcher MJ. Cell origin and differentiation in the repair of full-thickness defects of articular cartilage. *J Bone Joint Surg Am*. 1993;75(4):532-53.
48. Jackson DW, Lalor PA, Aberman HM, Simon TM. Spontaneous repair of full-thickness defects of articular cartilage in a goat model: a preliminary study. *J Bone Joint Surg Am*. 2001;83-A(1):53-64.
49. Rogachefsky RA, Dean DD, Howell DS, Altman RD. Treatment of canine osteoarthritis with insulin-like growth factor-1 (IGF-1) and sodium pentosan polysulfate. *Osteoarthritis Cartilage*. 1993;1(2):105-14.
50. Shida J, Jingushi S, Izumi T, Iwaki A, Sugioka Y. Basic fibroblast growth factor stimulates articular cartilage enlargement in young rats in vivo. *J Orthop Res*. 1996;14(2):265-72.
51. Sellers RS, Peluso D, Morris EA. The effect of recombinant human bone morphogenetic protein-2 (rhBMP-2) on the healing of full-thickness defects of articular cartilage. *J Bone Joint Surg Am*. 1997;79(10):1452-63.
52. Joyce ME, Roberts AB, Sporn MB, Bolander ME. Transforming growth factor-beta and the initiation of chondrogenesis and osteogenesis in the rat femur. *J Cell Biol*. 1990;110(6):2195-207.
53. Hanada K, Solchaga LA, Caplan AI, Hering TM, Goldberg VM, Yoo JU, et al. BMP-2 induction and TGF-beta 1 modulation of rat periosteal cell chondrogenesis. *J Cell Biochem*. 2001;81(2):284-94.
54. Asahina I, Sampath TK, Hauschka PV. Human osteogenic protein-1 induces chondroblastic, osteoblastic, and/or adipocytic differentiation of clonal murine target cells. *Exp Cell Res*. 1996;222(1):38-47.
55. Klein-Nulend J, Louwse RT, Heyligers IC, Wuisman PI, Semeins CM, Goei SW, et al. Osteogenic protein (OP-1, BMP-7) stimulates cartilage differentiation of human and goat perichondrium tissue in vitro. *J Biomed Mater Res*. 1998;40(4):614-20.
56. Jentsch KD, Wellmitz G, Heder G, Petzold E, Buntrock P, Oehme P. A bovine brain fraction with fibroblast growth factor activity inducing articular cartilage regeneration in vivo. *Acta Biol Med Ger*. 1980;39(8-9):967-71.

57. Hotten GC, Matsumoto T, Kimura M, Bechtold RF, Kron R, Ohara T, *et al.* Recombinant human growth/differentiation factor 5 stimulates mesenchyme aggregation and chondrogenesis responsible for the skeletal development of limbs. *Growth Factors*. 1996;13(1-2):65-74.
58. Vortkamp A, Lee K, Lanske B, Segre GV, Kronenberg HM, Tabin CJ. Regulation of rate of cartilage differentiation by Indian hedgehog and PTH-related protein. *Science*. 1996;273(5275):613-22.
59. Amizuka N, Warshawsky H, Henderson JE, Goltzman D, Karaplis AC. Parathyroid hormone-related peptide-depleted mice show abnormal epiphyseal cartilage development and altered endochondral bone formation. *J Cell Biol*. 1994;126(6):1611-23.
60. Trippel SB, Wroblewski J, Makower AM, Whelan MC, Schoenfeld D, Doctrow SR. Regulation of growth-plate chondrocytes by insulin-like growth-factor I and basic fibroblast growth factor. *J Bone Joint Surg Am*. 1993;75(2):177-89.
61. Fujimoto E, Ochi M, Kato Y, Mochizuki Y, Sumen Y, Ikuta Y. Beneficial effect of basic fibroblast growth factor on the repair of full-thickness defects in rabbit articular cartilage. *Arch Orthop Trauma Surg*. 1999;119(3-4):139-45.
62. Trippel SB. Growth factor actions on articular cartilage. *J Rheumatol Suppl*. 1995;43:129-32.
63. Nixon AJ, Fortier LA, Williams J, Mohammed H. Enhanced repair of extensive articular defects by insulin-like growth factor-I-laden fibrin composites. *J Orthop Res*. 1999;17(4):475-87.
64. Trippel SB, Van Wyk JJ, Foster MB, Svoboda ME. Characterization of a specific somatomedin-c receptor on isolated bovine growth plate chondrocytes. *Endocrinology*. 1983;112(6):2128-36.
65. Erlacher L, Ng CK, Ullrich R, Krieger S, Luyten FP. Presence of cartilage-derived morphogenetic proteins in articular cartilage and enhancement of matrix replacement in vitro. *Arthritis Rheum*. 1998;41(2):263-73.
66. Katayama R, Wakitani S, Tsumaki N, Morita Y, Matsushita I, Gejo R, *et al.* Repair of articular cartilage defects in rabbits using CDMP1 gene-transfected autologous mesenchymal cells derived from bone marrow. *Rheumatology (Oxford)*. 2004;43(8):980-5.
67. Bi W, Deng JM, Zhang Z, Behringer RR, de Crombrughe B. Sox9 is required for cartilage formation. *Nat Genet*. 1999;22(1):85-9.
68. Inada M, Yasui T, Nomura S, Miyake S, Deguchi K, Himeno M, *et al.* Maturational disturbance of chondrocytes in Cbfa1-deficient mice. *Dev Dyn*. 1999;214(4):279-90.
69. Zhao GQ, Eberspaecher H, Seldin MF, de Crombrughe B. The gene for the homeodomain-containing protein Cart-1 is expressed in cells that have a chondrogenic potential during embryonic development. *Mech Dev*. 1994;48(3):245-54.
70. Sumarsono SH, Wilson TJ, Tymms MJ, Venter DJ, Corrick CM, Kola R, *et al.* Down's syndrome-like skeletal abnormalities in Ets2 transgenic mice. *Nature*. 1996;379(6565):534-7.
71. Mackie EJ. Tenascin in connective tissue development and pathogenesis. *Perspect Dev Neurobiol*. 1994;2(1):125-32.
72. Lanske B, Karaplis AC, Lee K, Luz A, Vortkamp A, Pirro A, *et al.* PTH/PTHrP receptor in early development and Indian hedgehog-regulated bone growth. *Science*. 1996;273(5275):663-6.
73. Goldring MB, Fukuo K, Birkhead JR, Dudek E, Sandell LJ. Transcriptional suppression by interleukin-1 and interferon-gamma of type II collagen gene expression in human chondrocytes. *J Cell Biochem*. 1994;54(1):85-99.
74. Shalom-Barak T, Quach J, Lotz M. Interleukin-17-induced gene expression in articular chondrocytes is associated with activation of mitogen-activated protein kinases and NF-kappaB. *J Biol Chem*. 1998;273(42):27467-73.
75. Stadler J, Stefanovic-Racic M, Billiar TR, Curran RD, McIntyre LA, Georgescu HI, *et al.* Articular chondrocytes synthesize nitric oxide in response to cytokines and lipopolysaccharide. *J Immunol*. 1991;147(11):3915-20.
76. Pelletier JP, Martel-Pelletier J. [Role of synovial inflammation cytokines and IGF-1 in the physiopathology of osteoarthritis]. *Rev Rhum Ed Fr*. 1994;61(9 Pt 2):103S-8S.
77. Blanco FJ, Ochs RL, Schwarz H, Lotz M. Chondrocyte apoptosis induced by nitric oxide. *Am J Pathol*. 1995;146(1):75-85.
78. Di Cesare PE, Frenkel SR, Carlson CS, Fang C, Liu C. Regional gene therapy for full-thickness articular cartilage lesions using naked DNA with a collagen matrix. *J Orthop Res*. 2006;24(5):1118-27.
79. Nita I, Ghivizzani SC, Galea-Lauri J, Bandara G, Georgescu HI, Robbins PD, *et al.* Direct gene delivery to synovium: an evaluation of potential vectors in vitro and in vivo. *Arthritis Rheum*. 1996;39(5):820-8.
80. Ikeda T, Kubo T, Arai Y, Nakanishi T, Kobayashi K, Takahashi K, *et al.* Adenovirus mediated gene delivery to the joints of guinea pigs. *J Rheumatol*. 1998;25(9):1666-73.
81. Ghivizzani SC, Lechman ER, Tio C, Mule KM, Chada S, McCormack JE, *et al.* Direct retrovirus-mediated gene transfer to the synovium of the rabbit knee: implications for arthritis gene therapy. *Gene Ther*. 1997;4(9):977-82.
82. Wehling P, Reinecke J, Baltzer AW, Granrath M, Schulitz KP, Schultz C, *et al.* Clinical responses to gene therapy in joints of two subjects with rheumatoid arthritis. *Hum Gene Ther*. 2009;20(2):97-101.
83. Pan RY, Xiao X, Chen SL, Li J, Lin LC, Wang HJ, *et al.* Disease-inducible transgene expression from a recombinant adeno-associated virus vector in a rat arthritis model. *J Virol*. 1999;73(4):3410-7.
84. Watanabe S, Kim KN, Imagawa T, Thornton S, Grom A, Hirsch R. On the mechanism of protection of distal joints after local gene transfer in collagen-induced arthritis. *Hum Gene Ther*. 2000;11(5):751-8.
85. Tomita T, Hashimoto H, Tomita N, Morishita R, Lee SB, Hayashida K, *et al.* In vivo direct gene transfer into articular

- cartilage by intraarticular injection mediated by HVJ (Sendai virus) and liposomes. *Arthritis Rheum.* 1997;40(5):901-6.
86. Morisset S, Frisbie DD, Robbins PD, Nixon AJ, McIlwraith CW. IL-1ra/IGF-1 gene therapy modulates repair of microfractured chondral defects. *Clin Orthop Relat Res.* 2007;462:221-8.
 87. Madry H, Cucchiariini M, Terwilliger EF, Trippel SB. Recombinant adeno-associated virus vectors efficiently and persistently transduce chondrocytes in normal and osteoarthritic human articular cartilage. *Hum Gene Ther.* 2003;14(4):393-402.
 88. Coutts RD, Healey RM, Ostrander R, Sah RL, Goomer R, Amiel D. Matrices for cartilage repair. *Clin Orthop Relat Res.* 2001;(391 Suppl):S271-9.
 89. Pascher A, Palmer GD, Steinert A, Oligino T, Gouze E, Gouze JN, et al. Gene delivery to cartilage defects using coagulated bone marrow aspirate. *Gene Ther.* 2004;11(2):133-41.
 90. Kaul G, Cucchiariini M, Arntzen D, Zurakowski D, Menger MD, Kohn D, et al. Local stimulation of articular cartilage repair by transplantation of encapsulated chondrocytes overexpressing human fibroblast growth factor 2 (FGF-2) in vivo. *J Gene Med.* 2006;8(1):100-11.
 91. Madry H, Kaul G, Cucchiariini M, Stein U, Zurakowski D, Remberger K, et al. Enhanced repair of articular cartilage defects in vivo by transplanted chondrocytes overexpressing insulin-like growth factor I (IGF-I). *Gene Ther.* 2005;12(15):1171-9.
 92. Fragonas E, Valente M, Pozzi-Mucelli M, Toffanin R, Rizzo R, Silvestri F, et al. Articular cartilage repair in rabbits by using suspensions of allogenic chondrocytes in alginate. *Biomaterials.* 2000;21(8):795-801.
 93. Rahfoth B, Weisser J, Sternkopf F, Aigner T, von der Mark K, Brauer R. Transplantation of allograft chondrocytes embedded in agarose gel into cartilage defects of rabbits. *Osteoarthritis Cartilage.* 1998;6(1):50-65.
 94. Hunter CJ, Levenston ME. Maturation and integration of tissue-engineered cartilages within an in vitro defect repair model. *Tissue Eng.* 2004;10(5-6):736-46.
 95. Kawamura S, Wakitani S, Kimura T, Maeda A, Caplan AI, Shino K, et al. Articular cartilage repair: rabbit experiments with a collagen gel-biomatrix and chondrocytes cultured in it. *Acta Orthop Scand.* 1998;69(1):56-62.
 96. Ikeda T, Kubo T, Nakanishi T, Arai Y, Kobayashi K, Mazda O, et al. Ex vivo gene delivery using an adenovirus vector in treatment for cartilage defects. *J Rheumatol.* 2000;27(4):990-6.
 97. Perka C, Schultz O, Spitzer RS, Lindenhayn K. The influence of transforming growth factor beta1 on mesenchymal cell repair of full-thickness cartilage defects. *J Biomed Mater Res.* 2000;52(3):543-52.
 98. Driesang IM, Hunziker EB. Delamination rates of tissue flaps used in articular cartilage repair. *J Orthop Res.* 2000;18(6):909-11.
 99. Yokoo N, Saito T, Uesugi M, Kobayashi N, Xin KQ, Okuda K, et al. Repair of articular cartilage defect by autologous transplantation of basic fibroblast growth factor gene-transduced chondrocytes with adeno-associated virus vector. *Arthritis Rheum.* 2005;52(1):164-70.
 100. Freed LE, Grande DA, Lingbin Z, Emmanuel J, Marquis JC, Langer R. Joint resurfacing using allograft chondrocytes and synthetic biodegradable polymer scaffolds. *J Biomed Mater Res.* 1994;28(8):891-9.
 101. Perka C, Sittlinger M, Schultz O, Spitzer RS, Schlenzka D, Burmester GR. Tissue engineered cartilage repair using cryopreserved and noncryopreserved chondrocytes. *Clin Orthop Relat Res.* 2000;(378):245-54.
 102. Schaefer D, Martin I, Jundt G, Seidel J, Heberer M, Grodzinsky A, et al. Tissue-engineered composites for the repair of large osteochondral defects. *Arthritis Rheum.* 2002;46(9):2524-34.
 103. Kang R, Marui T, Ghivizzani SC, Nita IM, Georgescu HI, Suh JK, et al. Ex vivo gene transfer to chondrocytes in full-thickness articular cartilage defects: a feasibility study. *Osteoarthritis Cartilage.* 1997;5(2):139-43.
 104. Madry H, Cucchiariini M, Stein U, Remberger K, Menger MD, Kohn D, et al. Sustained transgene expression in cartilage defects in vivo after transplantation of articular chondrocytes modified by lipid-mediated gene transfer in a gel suspension delivery system. *J Gene Med.* 2003;5(6):502-9.
 105. Goomer RS, Deftos LJ, Terkeltaub R, Maris T, Lee MC, Harwood FL, et al. High-efficiency non-viral transfection of primary chondrocytes and perichondrial cells for ex-vivo gene therapy to repair articular cartilage defects. *Osteoarthritis Cartilage.* 2001;9(3):248-56.
 106. Uebliacker P, Wagner B, Kruger A, Vogt S, DeSantis G, Kennerknecht E, et al. Inducible nonviral gene expression in the treatment of osteochondral defects. *Osteoarthritis Cartilage.* 2004;12(9):711-9.
 107. Baragi VM, Renkiewicz RR, Jordan H, Bonadio J, Hartman JW, Roessler BJ. Transplantation of transduced chondrocytes protects articular cartilage from interleukin 1-induced extracellular matrix degradation. *J Clin Invest.* 1995;96(5):2454-60.
 108. Mason JM, Grande DA, Barcia M, Grant R, Pergolizzi RG, Breitbart AS. Expression of human bone morphogenic protein 7 in primary rabbit periosteal cells: potential utility in gene therapy for osteochondral repair. *Gene Ther.* 1998;5(8):1098-104.
 109. Adachi N, Sato K, Usas A, Fu FH, Ochi M, Han CW, et al. Muscle derived, cell based ex vivo gene therapy for treatment of full thickness articular cartilage defects. *J Rheumatol.* 2002;29(9):1920-30.
 110. Hirschmann F, Verhoeyen E, Wirth D, Bauwens S, Hauser H, Rudert M. Vital marking of articular chondrocytes by retroviral infection using green fluorescence protein. *Osteoarthritis Cartilage.* 2002;10(2):109-18.
 111. Mierisch CM, Wilson HA, Turner MA, Milbrandt TA, Berthou L, Hammarskjold ML, et al. Chondrocyte transplantation into articular cartilage defects with use of calcium alginate: the fate of the cells. *J Bone Joint Surg Am.* 2003;85-A(9):1757-67.
 112. Kobayashi N, Koshino T, Uesugi M, Yokoo N, Xin KQ, Okuda K, et al. Gene marking in adeno-associated virus vector infected periosteum derived cells for cartilage repair. *J Rheumatol.* 2002;29(10):2176-80.

113. Baragi VM, Renkiewicz RR, Qiu L, Brammer D, Riley JM, Sigler RE, *et al.* Transplantation of adenovirally transduced allogeneic chondrocytes into articular cartilage defects in vivo. *Osteoarthritis Cartilage*. 1997;5(4):275-82.
114. Madry H, Orth P, Kaul G, Zurakowski D, Menger MD, Kohn D, *et al.* Acceleration of articular cartilage repair by combined gene transfer of human insulin-like growth factor I and fibroblast growth factor-2 in vivo. *Arch Orthop Trauma Surg*. 2010;130(10):1311-22.
115. Turgeman G, Pittman DD, Muller R, Kurkalli BG, Zhou S, Pelled G, *et al.* Engineered human mesenchymal stem cells: a novel platform for skeletal cell mediated gene therapy. *J Gene Med*. 2001;3(3):240-51.
116. Cucchiari M, Madry H. Gene therapy for cartilage defects. *J Gene Med*. 2005;7(12):1495-509.
117. Che JH, Zhang ZR, Li GZ, Tan WH, Bai XD, Qu FJ. Application of tissue-engineered cartilage with BMP-7 gene to repair knee joint cartilage injury in rabbits. *Knee Surg Sports Traumatol Arthrosc*. 2010;18(4):496-503.
118. Gelse K, Muhle C, Franke O, Park J, Jehle M, Durst K, *et al.* Cell-based resurfacing of large cartilage defects: long-term evaluation of grafts from autologous transgene-activated periosteal cells in a porcine model of osteoarthritis. *Arthritis Rheum*. 2008;58(2):475-88.
119. Gysin R, Wergedal JE, Sheng MH, Kasukawa Y, Miyakoshi N, Chen ST, *et al.* Ex vivo gene therapy with stromal cells transduced with a retroviral vector containing the BMP4 gene completely heals critical size calvarial defect in rats. *Gene Ther*. 2002;9(15):991-9.
120. Vogt S, Wexel G, Tischer T, Schillinger U, Ueblacker P, Wagner B, *et al.* The influence of the stable expression of BMP2 in fibrin clots on the remodelling and repair of osteochondral defects. *Biomaterials*. 2009;30(12):2385-92.
121. Goodrich LR, Hidaka C, Robbins PD, Evans CH, Nixon AJ. Genetic modification of chondrocytes with insulin-like growth factor-1 enhances cartilage healing in an equine model. *J Bone Joint Surg Br*. 2007;89(5):672-85.
122. Kuroda R, Usas A, Kubo S, Corsi K, Peng H, Rose T, *et al.* Cartilage repair using bone morphogenetic protein 4 and muscle-derived stem cells. *Arthritis Rheum*. 2006;54(2):433-42.
123. Guo X, Zheng Q, Yang S, Shao Z, Yuan Q, Pan Z, *et al.* Repair of full-thickness articular cartilage defects by cultured mesenchymal stem cells transfected with the transforming growth factor beta1 gene. *Biomed Mater*. 2006;1(4):206-15.
124. Evans CH, Liu FJ, Glatt V, Hoyland JA, Kirker-Head C, Walsh A, *et al.* Use of genetically modified muscle and fat grafts to repair defects in bone and cartilage. *Eur Cell Mater*. 2009;18:96-111.
125. Ivkovic A, Pascher A, Hudetz D, Maticic D, Jelic M, Dickinson S, *et al.* Articular cartilage repair by genetically modified bone marrow aspirate in sheep. *Gene Ther*. 2010;17(6):779-89.
126. Gerich TG, Lobenhoffer HP, Fu FH, Robbins PD, Evans CH. [Virally mediated gene transfer in the patellar tendon: an experimental study in rabbits]. *Unfallchirurg*. 1997;100(5):354-62.
127. Madry H, Zurakowski D, Trippel SB. Overexpression of human insulin-like growth factor-I promotes new tissue formation in an ex vivo model of articular chondrocyte transplantation. *Gene Ther*. 2001;8(19):1443-9.
128. Goater JJ, O'Keefe RJ, Rosier RN, Puzas JE, Schwarz EM. Efficacy of ex vivo OPG gene therapy in preventing wear debris induced osteolysis. *J Orthop Res*. 2002;20(2):169-73.
129. Taniyama Y, Tachibana K, Hiraoka K, Aoki M, Yamamoto S, Matsumoto K, *et al.* Development of safe and efficient novel nonviral gene transfer using ultrasound: enhancement of transfection efficiency of naked plasmid DNA in skeletal muscle. *Gene Ther*. 2002;9(6):372-80.
130. Tsuchiya H, Kitoh H, Sugiura F, Ishiguro N. Chondrogenesis enhanced by overexpression of sox9 gene in mouse bone marrow-derived mesenchymal stem cells. *Biochem Biophys Res Commun*. 2003;301(2):338-43.
131. Madry H, Emkey G, Zurakowski D, Trippel SB. Overexpression of human fibroblast growth factor 2 stimulates cell proliferation in an ex vivo model of articular chondrocyte transplantation. *J Gene Med*. 2004;6(2):238-45.
132. Grossin L, Cournil-Henrionnet C, Pinzano A, Gaborit N, Dumas D, Etienne S, *et al.* Gene transfer with HSP 70 in rat chondrocytes confers cytoprotection in vitro and during experimental osteoarthritis. *FASEB J*. 2006;20(1):65-75.
133. Yeh LC, Lee JC. Co-transfection with the osteogenic protein (OP)-1 gene and the insulin-like growth factor (IGF)-I gene enhanced osteoblastic cell differentiation. *Biochim Biophys Acta*. 2006;1763(1):57-63.
134. Zhang HN, Leng P, Wang YZ, Zhang J. Treating human meniscal fibrochondrocytes with hIGF-1 gene by liposome. *Clin Orthop Relat Res*. 2009;467(12):3175-82.
135. Manning K, Rachakonda PS, Rai MF, Schmidt MF. Co-expression of insulin-like growth factor-1 and interleukin-4 in an in vitro inflammatory model. *Cytokine*. 2010;50(3):297-305.
136. Li Y, Tew SR, Russell AM, Gonzalez KR, Hardingham TE, Hawkins RE. Transduction of passaged human articular chondrocytes with adenoviral, retroviral, and lentiviral vectors and the effects of enhanced expression of SOX9. *Tissue Eng*. 2004;10(3-4):575-84.
137. Attur MG, Dave MN, Leung MY, Cipolletta C, Meseck M, Woo SL, *et al.* Functional genomic analysis of type II IL-1beta decoy receptor: potential for gene therapy in human arthritis and inflammation. *J Immunol*. 2002;168(4):2001-10.
138. Smith P, Shuler FD, Georgescu HI, Ghivizzani SC, Johnstone B, Niyibizi C, *et al.* Genetic enhancement of matrix synthesis by articular chondrocytes: comparison of different growth factor genes in the presence and absence of interleukin-1. *Arthritis Rheum*. 2000;43(5):1156-64.
139. Goto H, Shuler FD, Lamsam C, Moller HD, Niyibizi C, Fu FH, *et al.* Transfer of lacZ marker gene to the meniscus. *J Bone Joint Surg Am*. 1999;81(7):918-25.

140. Gerich TG, Kang R, Fu FH, Robbins PD, Evans CH. Gene transfer to the rabbit patellar tendon: potential for genetic enhancement of tendon and ligament healing. *Gene Ther.* 1996;3(12):1089-93.
141. Lou J, Kubota H, Hotokezaka S, Ludwig FJ, Manske PR. In vivo gene transfer and overexpression of focal adhesion kinase (pp125 FAK) mediated by recombinant adenovirus-induced tendon adhesion formation and epitenon cell change. *J Orthop Res.* 1997;15(6):911-8.
142. Mehrara BJ, Saadeh PB, Steinbrech DS, Dudziak M, Specator JA, Greenwald JA, et al. Adenovirus-mediated gene therapy of osteoblasts in vitro and in vivo. *J Bone Miner Res.* 1999;14(8):1290-301.
143. Nixon AJ, Brower-Toland BD, Bent SJ, Saxer RA, Wilke MJ, Robbins PD, et al. Insulinlike growth factor-I gene therapy applications for cartilage repair. *Clin Orthop Relat Res.* 2000;(379 Suppl):S201-13.
144. Shuler FD, Georgescu HI, Niyibizi C, Studer RK, Mi Z, Johnstone B, et al. Increased matrix synthesis following adenoviral transfer of a transforming growth factor beta1 gene into articular chondrocytes. *J Orthop Res.* 2000;18(4):585-92.
145. Brower-Toland BD, Saxer RA, Goodrich LR, Mi Z, Robbins PD, Evans CH, et al. Direct adenovirus-mediated insulin-like growth factor I gene transfer enhances transplant chondrocyte function. *Hum Gene Ther.* 2001;12(2):117-29.
146. Gelse K, Jiang QJ, Aigner T, Ritter T, Wagner K, Poschl E, et al. Fibroblast-mediated delivery of growth factor complementary DNA into mouse joints induces chondrogenesis but avoids the disadvantages of direct viral gene transfer. *Arthritis Rheum.* 2001;44(8):1943-53.
147. Saxer RA, Bent SJ, Brower-Toland BD, Mi Z, Robbins PD, Evans CH, et al. Gene mediated insulin-like growth factor-I delivery to the synovium. *J Orthop Res.* 2001;19(5):759-67.
148. Musgrave DS, Pruchnic R, Bosch P, Ziran BH, Whalen J, Huard J. Human skeletal muscle cells in ex vivo gene therapy to deliver bone morphogenetic protein-2. *J Bone Joint Surg Br.* 2002;84(1):120-7.
149. Ikeda T, Kamekura S, Mabuchi A, Kou I, Seki S, Takato T, et al. The combination of SOX5, SOX6, and SOX9 (the SOX trio) provides signals sufficient for induction of permanent cartilage. *Arthritis Rheum.* 2004;50(11):3561-73.
150. Haupt JL, Frisbie DD, McIlwraith CW, Robbins PD, Ghivizzani S, Evans CH, et al. Dual transduction of insulin-like growth factor-I and interleukin-1 receptor antagonist protein controls cartilage degradation in an osteoarthritic culture model. *J Orthop Res.* 2005;23(1):118-26.
151. Nixon AJ, Haupt JL, Frisbie DD, Morisset SS, McIlwraith CW, Robbins PD, et al. Gene-mediated restoration of cartilage matrix by combination insulin-like growth factor-I/interleukin-1 receptor antagonist therapy. *Gene Ther.* 2005;12(2):177-86.
152. Steinert AF, Palmer GD, Capito R, Hofstaetter JG, Pilapil C, Ghivizzani SC, et al. Genetically enhanced engineering of meniscus tissue using ex vivo delivery of transforming growth factor-beta 1 complementary deoxyribonucleic acid. *Tissue Eng.* 2007;13(9):2227-37.
153. Steinert AF, Weber M, Kunz M, Palmer GD, Noth U, Evans CH, et al. In situ IGF-1 gene delivery to cells emerging from the injured anterior cruciate ligament. *Biomaterials.* 2008;29(7):904-16.
154. Steinert AF, Proffen B, Kunz M, Hendrich C, Ghivizzani SC, Noth U, et al. Hypertrophy is induced during the in vitro chondrogenic differentiation of human mesenchymal stem cells by bone morphogenetic protein-2 and bone morphogenetic protein-4 gene transfer. *Arthritis Res Ther.* 2009;11(5):R148.
155. Roessler BJ, Hartman JW, Vallance DK, Latta JM, Janich SL, Davidson BL. Inhibition of interleukin-1-induced effects in synoviocytes transduced with the human IL-1 receptor antagonist cDNA using an adenoviral vector. *Hum Gene Ther.* 1995;6(3):307-16.
156. Baltzer AW, Whalen JD, Muzzonegro T, Georgescu HI, Robbins PD, Evans CH. [In vitro transduction of human osteoblast cell populations with retroviral vectors]. *Z Rheumatol.* 1999;58(2):88-94.
157. Hildebrand KA, Deie M, Allen CR, Smith DW, Georgescu HI, Evans CH, et al. Early expression of marker genes in the rabbit medial collateral and anterior cruciate ligaments: the use of different viral vectors and the effects of injury. *J Orthop Res.* 1999;17(1):37-42.
158. Goto H, Shuler FD, Niyibizi C, Fu FH, Robbins PD, Evans CH. Gene therapy for meniscal injury: enhanced synthesis of proteoglycan and collagen by meniscal cells transduced with a TGFbeta(1) gene. *Osteoarthritis Cartilage.* 2000;8(4):266-71.
159. Tew SR, Li Y, Pothacharoen P, Tweats LM, Hawkins RE, Hardingham TE. Retroviral transduction with SOX9 enhances re-expression of the chondrocyte phenotype in passaged osteoarthritic human articular chondrocytes. *Osteoarthritis Cartilage.* 2005;13(1):80-9.
160. Jennings K, Miyamae T, Traister R, Marinov A, Katakura S, Sowders D, et al. Proteasome inhibition enhances AAV-mediated transgene expression in human synoviocytes in vitro and in vivo. *Mol Ther.* 2005;11(4):600-7.
161. Goater J, Muller R, Kollias G, Firestein GS, Sanz I, O'Keefe RJ, et al. Empirical advantages of adeno associated viral vectors in vivo gene therapy for arthritis. *J Rheumatol.* 2000;27(4):983-9.
162. Zhang HG, Xie J, Yang P, Wang Y, Xu L, Liu D, et al. Adeno-associated virus production of soluble tumor necrosis factor receptor neutralizes tumor necrosis factor alpha and reduces arthritis. *Hum Gene Ther.* 2000;11(17):2431-42.
163. Hiraide A, Yokoo N, Xin KQ, Okuda K, Mizukami H, Ozawa K, et al. Repair of articular cartilage defect by intraarticular administration of basic fibroblast growth factor gene, using adeno-associated virus vector. *Hum Gene Ther.* 2005;16(12):1413-21.
164. Apparailly F, Millet V, Noel D, Jacquet C, Sany J, Jorgensen C. Tetracycline-inducible interleukin-10 gene transfer mediated

- by an adeno-associated virus: application to experimental arthritis. *Hum Gene Ther.* 2002;13(10):1179-88.
165. Arai Y, Kubo T, Fushiki S, Mazda O, Nakai H, Iwaki Y, *et al.* Gene delivery to human chondrocytes by an adeno associated virus vector. *J Rheumatol.* 2000;27(4):979-82.
 166. Ulrich-Vinther M, Maloney MD, Goater JJ, Soballe K, Goldring MB, O'Keefe RJ, *et al.* Light-activated gene transduction enhances adeno-associated virus vector-mediated gene expression in human articular chondrocytes. *Arthritis Rheum.* 2002;46(8):2095-104.
 167. Cucchiari M, Thurn T, Weimer A, Kohn D, Terwilliger EF, Madry H. Restoration of the extracellular matrix in human osteoarthritic articular cartilage by overexpression of the transcription factor SOX9. *Arthritis Rheum.* 2007;56(1):158-67.
 168. Cucchiari M, Terwilliger EF, Kohn D, Madry H. Remodelling of human osteoarthritic cartilage by FGF-2, alone or combined with Sox9 via rAAV gene transfer. *J Cell Mol Med.* 2009;13(8B):2476-88.
 169. Chamberlain JR, Schwarze U, Wang PR, Hirata RK, Hankenson KD, Pace JM, *et al.* Gene targeting in stem cells from individuals with osteogenesis imperfecta. *Science.* 2004;303(5661):1198-201.
 170. Ito H, Goater JJ, Tiyyapatanaputi P, Rubery PT, O'Keefe RJ, Schwarz EM. Light-activated gene transduction of recombinant adeno-associated virus in human mesenchymal stem cells. *Gene Ther.* 2004;11(1):34-41.
 171. Basile P, Dadali T, Jacobson J, Hasslund S, Ulrich-Vinther M, Soballe K, *et al.* Freeze-dried tendon allografts as tissue-engineering scaffolds for Gdf5 gene delivery. *Mol Ther.* 2008;16(3):466-73.
 172. Arsic N, Zacchigna S, Zentilin L, Ramirez-Correa G, Patarini L, Salvi A, *et al.* Vascular endothelial growth factor stimulates skeletal muscle regeneration in vivo. *Mol Ther.* 2004;10(5):844-54.
 173. Madry H, Cucchiari M, Kaul G, Kohn D, Terwilliger EF, Trippel SB. Menisci are efficiently transduced by recombinant adeno-associated virus vectors in vitro and in vivo. *Am J Sports Med.* 2004;32(8):1860-5.
 174. Ito H, Koefoed M, Tiyyapatanaputi P, Gromov K, Goater JJ, Carmouche J, *et al.* Remodeling of cortical bone allografts mediated by adherent rAAV-RANKL and VEGF gene therapy. *Nat Med.* 2005;11(3):291-7.
 175. Wang XT, Liu PY, Xin KQ, Tang JB. Tendon healing in vitro: bFGF gene transfer to tenocytes by adeno-associated viral vectors promotes expression of collagen genes. *J Hand Surg Am.* 2005;30(6):1255-61.
 176. Tang JB, Cao Y, Zhu B, Xin KQ, Wang XT, Liu PY. Adeno-associated virus-2-mediated bFGF gene transfer to digital flexor tendons significantly increases healing strength: an in vivo study. *J Bone Joint Surg Am.* 2008;90(5):1078-89.
 177. Cucchiari M, Schetting S, Terwilliger EF, Kohn D, Madry H. rAAV-mediated overexpression of FGF-2 promotes cell proliferation, survival, and alpha-SMA expression in human meniscal lesions. *Gene Ther.* 2009;16(11):1363-72.
 178. Kafienah W, Al-Fayez F, Hollander AP, Barker MD. Inhibition of cartilage degradation: a combined tissue engineering and gene therapy approach. *Arthritis Rheum.* 2003;48(3):709-18.
 179. Bondeson J, Lauder S, Wainwright S, Amos N, Evans A, Hughes C, *et al.* Adenoviral gene transfer of the endogenous inhibitor IkappaBalpha into human osteoarthritis synovial fibroblasts demonstrates that several matrix metalloproteinases and aggrecanases are nuclear factor-kappaB-dependent. *J Rheumatol.* 2007;34(3):523-33.
 180. Kim SH, Kim S, Evans CH, Ghivizzani SC, Oligino T, Robbins PD. Effective treatment of established murine collagen-induced arthritis by systemic administration of dendritic cells genetically modified to express IL-4. *J Immunol.* 2001;166(5):3499-505.
 181. Manning K, Rachakonda PS, Rai MF, Schmidt MF. Co-expression of insulin-like growth factor-1 and interleukin-4 in an in vitro inflammatory model. *Cytokine.* 2010;50(3):297-305.
 182. Dharmavaram RM, Liu G, Tuan RS, Stokes DG, Jimenez SA. Stable transfection of human fetal chondrocytes with a type II procollagen minigene: expression of the mutant protein and alterations in the structure of the extracellular matrix in vitro. *Arthritis Rheum.* 1999;42(7):1433-42.
 183. Venkatesan N, Barre L, Benani A, Netter P, Magdalou J, Fournel-Gigleux S, *et al.* Stimulation of proteoglycan synthesis by glucuronosyltransferase-I gene delivery: a strategy to promote cartilage repair. *Proc Natl Acad Sci U S A.* 2004;101(52):18087-92.
 184. Lee DK, Choi KB, Oh IS, Song SU, Hwang S, Lim CL, *et al.* Continuous transforming growth factor beta1 secretion by cell-mediated gene therapy maintains chondrocyte redifferentiation. *Tissue Eng.* 2005;11(1-2):310-8.
 185. Schmal H, Mehlhorn AT, Zwingmann J, Muller CA, Stark GB, Sudkamp NP. Stimulation of chondrocytes in vitro by gene transfer with plasmids coding for epidermal growth factor (hEGF) and basic fibroblast growth factor (bFGF). *Cytherapy.* 2005;7(3):292-300.
 186. Piera-Velazquez S, Jimenez SA, Stokes D. Increased life span of human osteoarthritic chondrocytes by exogenous expression of telomerase. *Arthritis Rheum.* 2002;46(3):683-93.
 187. Surendran S, Kim SH, Jee BK, Ahn SH, Gopinathan P, Han CW. Anti-apoptotic Bcl-2 gene transfection of human articular chondrocytes protects against nitric oxide-induced apoptosis. *J Bone Joint Surg Br.* 2006;88(12):1660-5.
 188. Chan JM, Villarreal G, Jin WW, Stepan T, Burstein H, Wahl SM. Intraarticular gene transfer of TNFR:Fc suppresses experimental arthritis with reduced systemic distribution of the gene product. *Mol Ther.* 2002;6(6):727-36.
 189. Apparailly F, Verwaerde C, Jacquet C, Aurialt C, Sany J, Jorgensen C. Adenovirus-mediated transfer of viral IL-10 gene inhibits murine collagen-induced arthritis. *J Immunol.* 1998;160(11):5213-20.
 190. Song XY, Gu M, Jin WW, Klinman DM, Wahl SM. Plasmid DNA encoding transforming growth factor-beta1 suppresses

- chronic disease in a streptococcal cell wall-induced arthritis model. *J Clin Invest*. 1998;101(12):2615-21.
191. Roessler BJ, Allen ED, Wilson JM, Hartman JW, Davidson BL. Adenoviral-mediated gene transfer to rabbit synovium in vivo. *J Clin Invest*. 1993;92(2):1085-92.
192. Kim SH, Evans CH, Kim S, Oligino T, Ghivizzani SC, Robbins PD. Gene therapy for established murine collagen-induced arthritis by local and systemic adenovirus-mediated delivery of interleukin-4. *Arthritis Res*. 2000;2(4):293-302.
193. Lechman ER, Jaffurs D, Ghivizzani SC, Gambotto A, Kovsdi I, Mi Z, et al. Direct adenoviral gene transfer of viral IL-10 to rabbit knees with experimental arthritis ameliorates disease in both injected and contralateral control knees. *J Immunol*. 1999;163(4):2202-8.
194. Keravala A, Lechman ER, Nash J, Mi Z, Robbins PD. Human, viral or mutant human IL-10 expressed after local adenovirus-mediated gene transfer are equally effective in ameliorating disease pathology in a rabbit knee model of antigen-induced arthritis. *Arthritis Res Ther*. 2006;8(4):R91.
195. Kim KN, Watanabe S, Ma Y, Thornton S, Giannini EH, Hirsch R. Viral IL-10 and soluble TNF receptor act synergistically to inhibit collagen-induced arthritis following adenovirus-mediated gene transfer. *J Immunol*. 2000;164(3):1576-81.
196. Fernandes J, Tardif G, Martel-Pelletier J, Lascau-Coman V, Dupuis M, Moldovan F, et al. In vivo transfer of interleukin-1 receptor antagonist gene in osteoarthritic rabbit knee joints: prevention of osteoarthritis progression. *Am J Pathol*. 1999;154(4):1159-69.
197. Frisbie DD, McIlwraith CW. Evaluation of gene therapy as a treatment for equine traumatic arthritis and osteoarthritis. *Clin Orthop Relat Res*. 2000;(379 Suppl):S273-87.
198. Scharstuhl A, Vitters EL, van der Kraan PM, van den Berg WB. Reduction of osteophyte formation and synovial thickening by adenoviral overexpression of transforming growth factor beta/bone morphogenetic protein inhibitors during experimental osteoarthritis. *Arthritis Rheum*. 2003;48(12):3442-51.
199. Chen LX, Lin L, Wang HJ, Wei XL, Fu X, Zhang JY, et al. Suppression of early experimental osteoarthritis by in vivo delivery of the adenoviral vector-mediated NF-kappaBp65-specific siRNA. *Osteoarthritis Cartilage*. 2008;16(2):174-84.
200. Hsieh JL, Shen PC, Shiau AL, Jou IM, Lee CH, Teo ML, et al. Adenovirus-mediated kallistatin gene transfer ameliorates disease progression in a rat model of osteoarthritis induced by anterior cruciate ligament transection. *Hum Gene Ther*. 2009;20(2):147-58.
201. Hsieh JL, Shen PC, Shiau AL, Jou IM, Lee CH, Wang CR, et al. Intraarticular gene transfer of thrombospondin-1 suppresses the disease progression of experimental osteoarthritis. *J Orthop Res*. 2010;28(10):1300-6.
202. Mi Z, Ghivizzani SC, Lechman ER, Jaffurs D, Glorioso JC, Evans CH, et al. Adenovirus-mediated gene transfer of insulin-like growth factor 1 stimulates proteoglycan synthesis in rabbit joints. *Arthritis Rheum*. 2000;43(11):2563-70.
203. Bandara G, Mueller GM, Galea-Lauri J, Tindal MH, Georgescu HI, Suchanek MK, et al. Intraarticular expression of biologically active interleukin 1-receptor-antagonist protein by ex vivo gene transfer. *Proc Natl Acad Sci U S A*. 1993;90(22):10764-8.
204. Hung GL, Galea-Lauri J, Mueller GM, Georgescu HI, Larkin LA, Suchanek MK, et al. Suppression of intra-articular responses to interleukin-1 by transfer of the interleukin-1 receptor antagonist gene to synovium. *Gene Ther*. 1994;1(1):64-9.
205. Makarov SS, Olsen JC, Johnston WN, Anderle SK, Brown RR, Baldwin AS Jr, et al. Suppression of experimental arthritis by gene transfer of interleukin 1 receptor antagonist cDNA. *Proc Natl Acad Sci U S A*. 1996;93(1):402-6.
206. Otani K, Nita I, Macaulay W, Georgescu HI, Robbins PD, Evans CH. Suppression of antigen-induced arthritis in rabbits by ex vivo gene therapy. *J Immunol*. 1996;156(9):3558-62.
207. Pelletier JP, Caron JP, Evans C, Robbins PD, Georgescu HI, Jovanovic D, et al. In vivo suppression of early experimental osteoarthritis by interleukin-1 receptor antagonist using gene therapy. *Arthritis Rheum*. 1997;40(6):1012-9.
208. Zhang X, Mao Z, Yu C. Suppression of early experimental osteoarthritis by gene transfer of interleukin-1 receptor antagonist and interleukin-10. *J Orthop Res*. 2004;22(4):742-50.
209. Kim SH, Lechman ER, Kim S, Nash J, Oligino TJ, Robbins PD. Ex vivo gene delivery of IL-1Ra and soluble TNF receptor confers a distal synergistic therapeutic effect in antigen-induced arthritis. *Mol Ther*. 2002;6(5):591-600.
210. Yoo JU, Mandell I, Angele P, Johnstone B. Chondrogenitor cells and gene therapy. *Clin Orthop Relat Res*. 2000;(379 Suppl):S164-70.
211. Barry FP, Murphy JM. Mesenchymal stem cells: clinical applications and biological characterization. *Int J Biochem Cell Biol*. 2004;36(4):568-84.
212. Matsumoto T, Cooper GM, Gharaibeh B, Meszaros LB, Li G, Usas A, et al. Blocking VEGF as a potential approach to improve cartilage healing after osteoarthritis. *J Musculoskelet Neuronal Interact*. 2008;8(4):316-7.
213. Matsumoto T, Cooper GM, Gharaibeh B, Meszaros LB, Li G, Usas A, et al. Cartilage repair in a rat model of osteoarthritis through intraarticular transplantation of muscle-derived stem cells expressing bone morphogenetic protein 4 and soluble Flt-1. *Arthritis Rheum*. 2009;60(5):1390-405.
214. Katsube K, Bishop AT, Simari RD, Yla-Herttuala S, Friedrich PF. Vascular endothelial growth factor (VEGF) gene transfer enhances surgical revascularization of necrotic bone. *J Orthop Res*. 2005;23(2):469-74.
215. Nakamura N, Horibe S, Iwahashi T, Kawano K, Shino K, Yoshikawa H. Healing of a chondral fragment of the knee in an adolescent after internal fixation: a case report. *J Bone Joint Surg Am*. 2004;86-A(12):2741-6.
216. Rath E, Richmond JC. The menisci: basic science and advances in treatment. *Br J Sports Med*. 2000;34(4):252-7.

217. Gray JC. Neural and vascular anatomy of the menisci of the human knee. *J Orthop Sports Phys Ther.* 1999;29(1):23-30.
218. Messner K, Gao J. The menisci of the knee joint: anatomical and functional characteristics, and a rationale for clinical treatment. *J Anat.* 1998;193(Pt 2):161-78.
219. Chivers MD, Howitt SD. Anatomy and physical examination of the knee menisci: a narrative review of the orthopedic literature. *J Can Chiropr Assoc.* 2009;53(4):319-33.
220. Lengsfeld M, Rudig L, von Issendorff WD, Koebke J. [Significance of shape differences between medial and lateral knee joint menisci for functional change of position]. *Unfallchirurgie.* 1991;17(6):309-15.
221. Bullough PG, Munuera L, Murphy J, Weinstein AM. The strength of the menisci of the knee as it relates to their fine structure. *J Bone Joint Surg Br.* 1970;52(3):564-7.
222. McDevitt CA, Webber RJ. The ultrastructure and biochemistry of meniscal cartilage. *Clin Orthop Relat Res.* 1990;(252):8-18.
223. Verdonk PC, Forsyth RG, Wang J, Almqvist KF, Verdonk R, Veys EM, *et al.* Characterisation of human knee meniscus cell phenotype. *Osteoarthritis Cartilage.* 2005;13(7):548-60.
224. Arnoczky SP, Warren RF. Microvasculature of the human meniscus. *Am J Sports Med.* 1982;10(2):90-5.
225. Cooper DE, Arnoczky SP, Warren RF. Meniscal repair. *Clin Sports Med.* 1991;10(3):529-48.
226. Arnoczky SP, Warren RF. The microvasculature of the meniscus and its response to injury: an experimental study in the dog. *Am J Sports Med.* 1983;11(3):131-41.
227. Englund M, Guermazi A, Roemer FW, Aliabadi P, Yang M, Lewis CE, *et al.* Meniscal tear in knees without surgery and the development of radiographic osteoarthritis among middle-aged and elderly persons: the Multicenter Osteoarthritis Study. *Arthritis Rheum.* 2009;60(3):831-9.
228. Englund M, Lohmander LS. Patellofemoral osteoarthritis coexistent with tibiofemoral osteoarthritis in a meniscectomy population. *Ann Rheum Dis.* 2005;64(12):1721-6.
229. Beaufils P, Hulet C, Dhenain M, Nizard R, Nourissat G, Pujol N. Clinical practice guidelines for the management of meniscal lesions and isolated lesions of the anterior cruciate ligament of the knee in adults. *Rev Chir Orthop Traumatol.* 2009;95(6):437-42.
230. Ding C, Martel-Pelletier J, Pelletier JP, Abram F, Raynauld JP, Cicuttini F, *et al.* Meniscal tear as an osteoarthritis risk factor in a largely non-osteoarthritic cohort: a cross-sectional study. *J Rheumatol.* 2007;34(4):776-84.
231. Tumia NS, Johnstone AJ. Platelet derived growth factor-AB enhances knee meniscal cell activity in vitro. *Knee.* 2009;16(1):73-6.
232. Bhargava MM, Hidaka C, Hannafin JA, Doty S, Warren RF. Effects of hepatocyte growth factor and platelet-derived growth factor on the repair of meniscal defects in vitro. *In Vitro Cell Dev Biol Anim.* 2005;41(8-9):305-10.
233. Lietman SA, Hobbs W, Inoue N, Reddi AH. Effects of selected growth factors on porcine meniscus in chemically defined medium. *Orthopedics.* 2003;26(8):799-803.
234. Spindler KP, Mayes CE, Miller RR, Imro AK, Davidson JM. Regional mitogenic response of the meniscus to platelet-derived growth factor (PDGF-AB). *J Orthop Res.* 1995;13(2):201-7.
235. Fox DB, Warnock JJ, Stoker AM, Luther JK, Cockrell M. Effects of growth factors on equine synovial fibroblasts seeded on synthetic scaffolds for avascular meniscal tissue engineering. *Res Vet Sci.* 2010;88(2):326-32.
236. Tumia NS, Johnstone AJ. Promoting the proliferative and synthetic activity of knee meniscal fibrochondrocytes using basic fibroblast growth factor in vitro. *Am J Sports Med.* 2004;32(4):915-20.
237. Webber RJ, Harris MG, Hough AJ Jr. Cell culture of rabbit meniscal fibrochondrocytes: proliferative and synthetic response to growth factors and ascorbate. *J Orthop Res.* 1985;3(1):36-42.
238. Imler SM, Doshi AN, Levenston ME. Combined effects of growth factors and static mechanical compression on meniscus explant biosynthesis. *Osteoarthritis Cartilage.* 2004;12(9):736-44.
239. Pangborn CA, Athanasiou KA. Growth factors and fibrochondrocytes in scaffolds. *J Orthop Res.* 2005;23(5):1184-90.
240. Tumia NS, Johnstone AJ. Regional regenerative potential of meniscal cartilage exposed to recombinant insulin-like growth factor-I in vitro. *J Bone Joint Surg Br.* 2004;86(7):1077-81.
241. Pangborn CA, Athanasiou KA. Effects of growth factors on meniscal fibrochondrocytes. *Tissue Eng.* 2005;11(7-8):1141-8.
242. Collier S, Ghosh P. Effects of transforming growth factor beta on proteoglycan synthesis by cell and explant cultures derived from the knee joint meniscus. *Osteoarthritis Cartilage.* 1995;3(2):127-38.
243. Koay EJ, Athanasiou KA. Development of serum-free, chemically defined conditions for human embryonic stem cell-derived fibrochondrogenesis. *Tissue Eng Part A.* 2009;15(8):2249-57.
244. Hoben GM, Willard VP, Athanasiou KA. Fibrochondrogenesis of hESCs: growth factor combinations and cocultures. *Stem Cells Dev.* 2009;18(2):283-92.
245. Arnoczky SP, Warren RF, Spivak JM. Meniscal repair using an exogenous fibrin clot: an experimental study in dogs. *J Bone Joint Surg Am.* 1988;70(8):1209-17.
246. Port J, Jackson DW, Lee TQ, Simon TM. Meniscal repair supplemented with exogenous fibrin clot and autogenous cultured marrow cells in the goat model. *Am J Sports Med.* 1996;24(4):547-55.
247. Shirakura K, Nijima M, Kobuna Y, Kizuki S. Free synovium promotes meniscal healing: synovium, muscle and synthetic mesh compared in dogs. *Acta Orthop Scand.* 1997;68(1):51-4.

248. Jitsuike J, Ochi M, Ikuta Y. Meniscal repair enhanced by an interpositional free synovial autograft: an experimental study in rabbits. *Arthroscopy*. 1994;10(6):659-66.
249. Petersen W, Pufe T, Starke C, Fuchs T, Kopf S, Neumann W, *et al.* The effect of locally applied vascular endothelial growth factor on meniscus healing: gross and histological findings. *Arch Orthop Trauma Surg*. 2007;127(4):235-40.
250. Petersen W, Pufe T, Starke C, Fuchs T, Kopf S, Raschke M, *et al.* Locally applied angiogenic factors: a new therapeutic tool for meniscal repair. *Ann Anat*. 2005;187(5-6):509-19.
251. van Tienen TG, Hannink G, Buma P. Meniscus replacement using synthetic materials. *Clin Sports Med*. 2009;28(1):143-56.
252. Martinek V, Imhoff A. Das künstliche meniskusimplantat. *Arthroskopie*. 2008;21:266-70.
253. Stone KR, Steadman JR, Rodkey WG, Li ST. Regeneration of meniscal cartilage with use of a collagen scaffold: analysis of preliminary data. *J Bone Joint Surg Am*. 1997;79(12):1770-7.
254. Rodkey WG, DeHaven KE, Montgomery WH 3rd, Baker CL Jr, Beck CL Jr, Hormel SE, *et al.* Comparison of the collagen meniscus implant with partial meniscectomy: a prospective randomized trial. *J Bone Joint Surg Am*. 2008;90(7):1413-26.
255. de Groot JH, de Vrijer R, Pennings AJ, Klompmaker J, Veth RP, Jansen HW. Use of porous polyurethanes for meniscal reconstruction and meniscal prostheses. *Biomaterials*. 1996;17(2):163-73.
256. Welsing RT, van Tienen TG, Ramrattan N, Heijkants R, Schouten AJ, Veth RP, *et al.* Effect on tissue differentiation and articular cartilage degradation of a polymer meniscus implant: a 2-year follow-up study in dogs. *Am J Sports Med*. 2008;36(10):1978-89.
257. Langer R, Vacanti JP. Tissue engineering. *Science*. 1993;260(5110):920-6.
258. Nerem RM. Cellular engineering. *Ann Biomed Eng*. 1991;19(5):529-45.
259. Buma P, Ramrattan NN, van Tienen TG, Veth RP. Tissue engineering of the meniscus. *Biomaterials*. 2004;25(9):1523-32.
260. Sweigart MA, Athanasiou KA. Toward tissue engineering of the knee meniscus. *Tissue Eng*. 2001;7(2):111-29.
261. Setton LA, Guilak F, Hsu EW, Vail TP. Biomechanical factors in tissue engineered meniscal repair. *Clin Orthop Relat Res*. 1999;(367 Suppl):S254-72.
262. Sandmann GH, Eichhorn S, Vogt S, Adamczyk C, Aryee S, Hoberg M, *et al.* Generation and characterization of a human acellular meniscus scaffold for tissue engineering. *J Biomed Mater Res A*. 2009;91(2):567-74.
263. Stabile KJ, Odom D, Smith TL, Northam C, Whitlock PW, Smith BP, *et al.* An acellular, allograft-derived meniscus scaffold in an ovine model. *Arthroscopy*. 2010;26(7):936-48.
264. Ibarra C, Koski JA, Warren RF. Tissue engineering meniscus: cells and matrix. *Orthop Clin North Am*. 2000;31(3):411-8.
265. Muller-Rath R, Mumme T, Miltner O, Andereya S, Schneider U. [Meniscus replacement: current aspects in the field of tissue engineering]. *Z Orthop Ihre Grenzgeb*. 2004;142(5):540-5.
266. Adams SB Jr, Randolph MA, Gill TJ. Tissue engineering for meniscus repair. *J Knee Surg*. 2005;18(1):25-30.
267. Arnoczky SP. Building a meniscus: biologic considerations. *Clin Orthop Relat Res*. 1999;(367 Suppl):S244-53.
268. Baker BM, Gee AO, Sheth NP, Huffman GR, Sennett BJ, Schaer TP, *et al.* Meniscus tissue engineering on the nanoscale: from basic principles to clinical application. *J Knee Surg*. 2009;22(1):45-59.
269. Steadman JR, Rodkey WG. Tissue-engineered collagen meniscus implants: 5- to 6-year feasibility study results. *Arthroscopy*. 2005;21(5):515-25.
270. Rodeo SA, Seneviratne A, Suzuki K, Felker K, Wickiewicz TL, Warren RF. Histological analysis of human meniscal allografts: a preliminary report. *J Bone Joint Surg Am*. 2000;82-A(8):1071-82.
271. Sohn DH, Toth AP. Meniscus transplantation: current concepts. *J Knee Surg*. 2008;21(2):163-72.
272. Elder BD, Eleswarapu SV, Athanasiou KA. Extraction techniques for the decellularization of tissue engineered articular cartilage constructs. *Biomaterials*. 2009;30(22):3749-56.
273. Stapleton TW, Ingram J, Katta J, Knight R, Korossis S, Fisher J, *et al.* Development and characterization of an acellular porcine medial meniscus for use in tissue engineering. *Tissue Eng Part A*. 2008;14(4):505-18.
274. Stone KR, Rodkey WG, Webber R, McKinney L, Steadman JR. Meniscal regeneration with copolymeric collagenscaffolds: in vitro and in vivo studies evaluated clinically, histologically, and biochemically. *Am J Sports Med*. 1992;20(2):104-11.
275. Gastel JA, Muirhead WR, Lifrak JT, Fadale PD, Hulstyn MJ, Labrador DP. Meniscal tissue regeneration using a collagenous biomaterial derived from porcine small intestine submucosa. *Arthroscopy*. 2001;17(2):151-9.
276. Welch JA, Montgomery RD, Lenz SD, Plouhar P, Shelton WR. Evaluation of small-intestinal submucosa implants for repair of meniscal defects in dogs. *Am J Vet Res*. 2002;63(3):427-31.
277. Cook JL, Fox DB, Malaviya P, Tomlinson JL, Kuroki K, Cook CR, *et al.* Long-term outcome for large meniscal defects treated with small intestinal submucosa in a dog model. *Am J Sports Med*. 2006;34(1):32-42.
278. Cook JL, Tomlinson JL, Kreeger JM, Cook CR. Induction of meniscal regeneration in dogs using a novel biomaterial. *Am J Sports Med*. 1999;27(5):658-65.
279. Klompmaker J, Jansen HW, Veth RP, de Groot JH, Nijenhuis AJ, Pennings AJ. Porous polymer implant for repair of meniscal lesions: a preliminary study in dogs. *Biomaterials*. 1991;12(9):810-6.
280. Tienen TG, Heijkants RG, de Groot JH, Pennings AJ, Schouten AJ, Veth RP, *et al.* Replacement of the knee meniscus by a porous polymer implant: a study in dogs. *Am J Sports Med*. 2006;34(1):64-71.
281. Kobayashi M, Toguchida J, Oka M. Development of an artificial meniscus using polyvinyl alcohol-hydrogel for early return

- to, and continuance of, athletic life in sportspersons with severe meniscus injury. II: animal experiments. *Knee*. 2003;10(1):53.
282. Kang SW, Son SM, Lee JS, Lee ES, Lee KY, Park SG, *et al*. Regeneration of whole meniscus using meniscal cells and polymer scaffolds in a rabbit total meniscectomy model. *J Biomed Mater Res A*. 2006;78(3):659-71.
283. Arnoczky SP, DiCarlo EF, O'Brien SJ, Warren RF. Cellular repopulation of deep-frozen meniscal autografts: an experimental study in the dog. *Arthroscopy*. 1992;8(4):428-36.
284. Marsano A, Vunjak-Novakovic G, Martin I. Towards tissue engineering of meniscus substitutes: selection of cell source and culture environment. *Conf Proc IEEE Eng Med Biol Soc*. 2006;1:3656-8.
285. Chiari C, Koller U, Kapeller B, Dorotka R, Bindreiter U, Nehrer S. Different behavior of meniscal cells in collagen II/I,III and Hyaff-11 scaffolds in vitro. *Tissue Eng Part A*. 2008;14(8):1295-304.
286. Gunja NJ, Athanasiou KA. Effects of co-cultures of meniscus cells and articular chondrocytes on PLLA scaffolds. *Biotechnol Bioeng*. 2009;103(4):808-16.
287. Kon E, Chiari C, Marcacci M, Delcogliano M, Salter DM, Martin I, *et al*. Tissue engineering for total meniscal substitution: animal study in sheep model. *Tissue Eng Part A*. 2008;14(6):1067-80.
288. Tan Y, Zhang Y, Pei M. Meniscus reconstruction through coculturing meniscus cells with synovium-derived stem cells on small intestine submucosa: a pilot study to engineer meniscus tissue constructs. *Tissue Eng Part A*. 2010;16(1):67-79.
289. Yamasaki T, Deie M, Shinomiya R, Yasunaga Y, Yanada S, Ochi M. Transplantation of meniscus regenerated by tissue engineering with a scaffold derived from a rat meniscus and mesenchymal stromal cells derived from rat bone marrow. *Artif Organs*. 2008;32(7):519-24.
290. Johns DE, Athanasiou KA. Growth factor effects on costal chondrocytes for tissue engineering fibrocartilage. *Cell Tissue Res*. 2008;333(3):439-47.
291. Hidaka C, Ibarra C, Hannafin JA, Torzilli PA, Quitoriano M, Jen SS, *et al*. Formation of vascularized meniscal tissue by combining gene therapy with tissue engineering. *Tissue Eng*. 2002;8(1):93-105.
292. Zhang H, Leng P, Zhang J. Enhanced meniscal repair by overexpression of hIGF-1 in a full-thickness model. *Clin Orthop Relat Res*. 2009;467(12):3165-74.
293. Arnoczky SP, Warren RF, McDevitt CA. Meniscal replacement using a cryopreserved allograft: an experimental study in the dog. *Clin Orthop Relat Res*. 1990;(252):121-8.
294. Cole BJ, Carter TR, Rodeo SA. Allograft meniscal transplantation: background, techniques, and results. *Instr Course Lect*. 2003;52:383-96.
295. Dienst M, Greis PE, Ellis BJ, Bachus KN, Burks RT. Effect of lateral meniscal allograft sizing on contact mechanics of the lateral tibial plateau: an experimental study in human cadaveric knee joints. *Am J Sports Med*. 2007;35(1):34-42.
296. Garrett JC, Steensen RN. Meniscal transplantation in the human knee: a preliminary report. *Arthroscopy*. 1991;7(1):57-62.
297. Greis PE, Holmstrom MC, Bardana DD, Burks RT. Meniscal injury. II: management. *J Am Acad Orthop Surg*. 2002;10(3):177-87.
298. Kohn D, Verdonk R, Aagaard H, Seil R, Dienst M. Meniscal substitutes: animal experience. *Scand J Med Sci Sports*. 1999;9(3):141-5.
299. Messner K. Meniscal regeneration or meniscal transplantation? *Scand J Med Sci Sports*. 1999;9(3):162-7.
300. Packer JD, Rodeo SA. Meniscal allograft transplantation. *Clin Sports Med*. 2009;28(2):259-83, viii.
301. Rodeo SA. Meniscal allografts: where do we stand? *Am J Sports Med*. 2001;29(2):246-61.
302. Siegel MG, Roberts CS. Meniscal allografts. *Clin Sports Med*. 1993;12(1):59-80.
303. Verdonk PC, Verstraete KL, Almqvist KF, De Cuyper K, Veys EM, Verbruggen G, *et al*. Meniscal allograft transplantation: long-term clinical results with radiological and magnetic resonance imaging correlations. *Knee Surg Sports Traumatol Arthrosc*. 2006;14(8):694-706.
304. Verdonk R. Meniscal transplantation. *Acta Orthop Belg*. 2002;68(2):118-27.
305. Verdonk R, Almqvist KF, Huysse W, Verdonk PC. Meniscal allografts: indications and outcomes. *Sports Med Arthrosc*. 2007;15(3):121-5.
306. Crook TB, Ardolino A, Williams LA, Barlow IW. Meniscal allograft transplantation: a review of the current literature. *Ann R Coll Surg Engl*. 2009;91(5):361-5.
307. Huard J, Li Y, Peng H, Fu FH. Gene therapy and tissue engineering for sports medicine. *J Gene Med*. 2003;5(2):93-108.
308. Martinek V, Usas A, Pelinkovic D, Robbins P, Fu FH, Huard J. Genetic engineering of meniscal allografts. *Tissue Eng*. 2002;8(1):107-17.
309. Evans CH, Ghivizzani SC, Herndon JH, Wasko MC, Reinecke J, Wehling P, *et al*. Clinical trials in the gene therapy of arthritis. *Clin Orthop Relat Res*. 2000;(379 Suppl):S300-7.
310. Evans CH, Gouze JN, Gouze E, Robbins PD, Ghivizzani SC. Osteoarthritis gene therapy. *Gene Ther*. 2004;11(4):379-89.
311. Evans CH, Ghivizzani SC, Robbins PD. Gene therapy for arthritis: what next? *Arthritis Rheum*. 2006;54(6):1714-29.
312. Evans CH, Ghivizzani SC, Robbins PD. Arthritis gene therapy's first death. *Arthritis Res Ther*. 2008;10(3):110.
313. Evans CH, Ghivizzani SC, Oligino TJ, Robbins PD. Gene therapy for autoimmune disorders. *J Clin Immunol*. 2000;20(5):334-46.
314. Evans CH. Gene therapy: what have we accomplished and where do we go from here? *J Rheumatol Suppl*. 2005;72:17-20.
315. Evans CH, Robbins PD, Ghivizzani SC, Wasko MC, Tomaino MM, Kang R, *et al*. Gene transfer to human joints: progress toward a gene therapy of arthritis. *Proc Natl Acad Sci U S A*. 2005;102(24):8698-703.

316. Sant SM, Suarez TM, Moalli MR, Wu BY, Blaiwas M, Laing TJ, *et al.* Molecular lysis of synovial lining cells by in vivo herpes simplex virus-thymidine kinase gene transfer. *Hum Gene Ther.* 1998;9(18):2735-43.
317. Mease PJ, Hobbs K, Chalmers A, El-Gabalawy H, Bookman A, Keystone E, *et al.* Local delivery of a recombinant adenoassociated vector containing a tumour necrosis factor alpha antagonist gene in inflammatory arthritis: a phase 1 dose-escalation safety and tolerability study. *Ann Rheum Dis.* 2009;68(8):1247-54.
318. Gladman DD, Mease PJ, Ritchlin CT, Choy EH, Sharp JT, Ory PA, *et al.* Adalimumab for long-term treatment of psoriatic arthritis: forty-eight week data from the adalimumab effectiveness in psoriatic arthritis trial. *Arthritis Rheum.* 2007;56(2):476-88.
319. Kaiser J. Clinical research: death prompts a review of gene therapy vector. *Science.* 2007;317(5838):580.
320. Williams DA. RAC reviews serious adverse event associated with AAV therapy trial. *Mol Ther.* 2007;15(12):2053-4.
321. Williams DA. NIH Recombinant DNA Advisory Committee continues to ponder adverse event associated with AAV gene therapy trial. *Mol Ther.* 2008;16(3):427-8.
322. Sweeney HL. Gene doping. *Sci Am.* 2004;291(1):62-9.
323. Adam D. Gene therapy may be up to speed for cheats at 2008 Olympics. *Nature.* 2001;414(6864):569-70.
324. Friedmann T, Koss JO. Gene transfer and athletics: an impending problem. *Mol Ther.* 2001;3(6):819-20.
325. Friedmann T, Rabin O, Frankel MS. Ethics: gene doping and sport. *Science.* 2010;327(5966):647-8.
326. Barton-Davis ER, Shoturma DI, Musaro A, Rosenthal N, Sweeney HL. Viral mediated expression of insulin-like growth factor I blocks the aging-related loss of skeletal muscle function. *Proc Natl Acad Sci U S A.* 1998;95(26):15603-7.
327. Zhou S, Murphy JE, Escobedo JA, Dwarki VJ. Adeno-associated virus-mediated delivery of erythropoietin leads to sustained elevation of hematocrit in nonhuman primates. *Gene Ther.* 1998;5(5):665-70.
328. Baumgartner I, Pieczek A, Manor O, Blair R, Kearney M, Walsh K, *et al.* Constitutive expression of phVEGF165 after intramuscular gene transfer promotes collateral vessel development in patients with critical limb ischemia. *Circulation.* 1998;97(12):1114-23.
329. Baoutina A, Alexander IE, Rasko JE, Emslie KR. Potential use of gene transfer in athletic performance enhancement. *Mol Ther.* 2007;15(10):1751-66.
330. de la Chapelle A, Traskelin AL, Juvonen E. Truncated erythropoietin receptor causes dominantly inherited benign human erythrocytosis. *Proc Natl Acad Sci U S A.* 1993;90(10):4495-9.
331. Slot O. Apocalypse now: fears of gene doping are realised. *The Times.* 2006 Feb 2.
332. Hanson RW, Hakimi P. Born to run: the story of the PEPCK-Cmus mouse. *Biochimie.* 2008;90(6):838-42.
333. Baoutina A, Alexander IE, Rasko JE, Emslie KR. Developing strategies for detection of gene doping. *J Gene Med.* 2008;10(1):3-20.