



Zootaxa 3938 (1): 001–147  
www.mapress.com/zootaxa/

Copyright © 2015 Magnolia Press

# Monograph

ISSN 1175-5326 (print edition)

**ZOOTAXA**

ISSN 1175-5334 (online edition)

<http://dx.doi.org/10.11646/zootaxa.3938.1.1>

<http://zoobank.org/urn:lsid:zoobank.org:pub:334452F1-C808-43C1-B2B4-C3A9ABA28A9C>

# ZOOTAXA

3938

## Genera of euophryine jumping spiders (Araneae: Salticidae), with a combined molecular-morphological phylogeny

JUNXIA ZHANG<sup>1</sup>\* & WAYNE P. MADDISON<sup>2</sup>

<sup>1</sup> Department of Zoology, University of British Columbia, 6270 University Boulevard, Vancouver, British Columbia, V6T 1Z4, Canada.

\*Current address: Department of Entomology, University of California, Riverside, Riverside, CA 92521, USA.

E-mail: [jxzhang1976@gmail.com](mailto:jxzhang1976@gmail.com)

<sup>2</sup> Departments of Zoology and Botany and Beaty Biodiversity Museum, University of British Columbia, 6270 University Boulevard, Vancouver, British Columbia, V6T 1Z4, Canada. E-mail: [wayne.maddison@ubc.ca](mailto:wayne.maddison@ubc.ca)



Magnolia Press  
Auckland, New Zealand

Accepted by T. Szűts: 21 Jan. 2015; published: 27 Mar. 2015

JUNXIA ZHANG & WAYNE P. MADDISON

**Genera of euophryine jumping spiders (Araneae: Salticidae), with a combined molecular-morphological phylogeny**

(*Zootaxa* 3938)

147 pp.; 30 cm.

27 Mar. 2015

ISBN 978-1-77557-667-9 (paperback)

ISBN 978-1-77557-668-6 (Online edition)

FIRST PUBLISHED IN 2015 BY

Magnolia Press

P.O. Box 41-383

Auckland 1346

New Zealand

e-mail: [zootaxa@mapress.com](mailto:zootaxa@mapress.com)

<http://www.mapress.com/zootaxa/>

© 2015 Magnolia Press

All rights reserved.

No part of this publication may be reproduced, stored, transmitted or disseminated, in any form, or by any means, without prior written permission from the publisher, to whom all requests to reproduce copyright material should be directed in writing.

This authorization does not extend to any other kind of copying, by any means, in any form, and for any purpose other than private research use.

ISSN 1175-5326 (Print edition)

ISSN 1175-5334 (Online edition)

## Table of contents

Abstract	4
Introduction	4
Material and methods	5
Techniques for morphological study	5
Phylogenetic study	5
Taxonomic decisions	7
Results	10
Phylogenetic study	10
Discussion	13
DNA vs. morphology in phylogenetic reconstruction	13
Role of morphology in phylogeny of Euophryinae	13
The major clades and genera of euophryines	14
Major euophryine clades from the New World	14
<i>Anasaitis-Corythalia</i> Clade	14
<i>Anasaitis</i> Bryant, 1950	15
<i>Corythalia</i> C. L. Koch, 1850	15
<i>Antillattus</i> Clade: <i>Antillattus</i> , <i>Petemathis</i> , <i>Truncattus</i> , and possibly <i>Allodecta</i> , <i>Caribattus</i>	16
<i>Antillattus</i> Bryant, 1943	16
<i>Agobardus</i> Clade: <i>Agobardus</i> , <i>Bythocrotus</i> , <i>Compsodecta</i> , <i>Parasaitis</i>	17
<i>Agobardus</i> Keyserling, 1885	17
<i>Compsodecta</i> Simon, 1903	18
<i>Naphrys-Corticattus</i> Clade	18
<i>Sidusa</i> Clade	19
<i>Sidusa</i> Peckham & Peckham, 1895	19
<i>Mopiopia-Saphrys</i> Clade	20
<i>Mopiopia</i> Simon, 1902	20
<i>Saphrys</i> gen. nov.	20
<i>Chapoda-Maeota</i> Clade: <i>Chapoda</i> , <i>Maeota</i> , <i>Commoris</i>	21
<i>Chapoda</i> Peckham & Peckham, 1896	21
<i>Maeota</i> Simon, 1901	22
<i>Amphidraus-Marma</i> Clade	22
<i>Amphidraus</i> Simon, 1900	22
<i>Coryphasia</i> Clade: <i>Coryphasia</i> and possibly <i>Semmolius</i>	23
<i>Coryphasia</i> Simon, 1902	23
<i>Semmolius</i> Simon, 1902	24
<i>Pensacola-Mexigonus</i> Clade	24
<i>Neonella</i> Clade: <i>Neonella</i> and <i>Darwinneon</i>	24
<i>Bellienna</i> Clade: <i>Bellienna</i> , <i>Saitidops</i> and possibly <i>Stoidis</i>	24
<i>Soesilarishius</i> Clade: <i>Soesilarishius</i> and possibly <i>Rhyphelia</i>	25
Other genera from the New World: <i>Ecuadattus</i> , <i>Ilargus</i> , <i>Popcornella</i> , <i>Tylogonus</i>	25
Major euophryine clades from the Old World	25
<i>Bathippus-Canama</i> Clade: <i>Bathippus</i> , <i>Canama</i> and possibly <i>Spilargis</i>	25
<i>Omoedus</i> Clade	26
<i>Omoedus</i> Thorell, 1881	26
<i>Bulolia-Coccorchestes</i> Clade: <i>Bulolia</i> , <i>Coccorchestes</i> , <i>Leptathamas</i> , <i>Variratina</i> and possibly <i>Athamas</i>	27
<i>Diolenius</i> Clade: <i>Chalcolecta</i> , <i>Chalcolemia</i> , <i>Diolenius</i> , <i>Efate</i> , <i>Furculattus</i> , <i>Ohilimia</i> , <i>Paraharmochirus</i> , <i>Sobasina</i> , <i>Tarodes</i> and <i>Udvardya</i>	27
<i>Pristobaeus</i> Clade: <i>Pristobaeus</i> , <i>Opisthoncana</i> , <i>Ergane</i> and <i>Asclytus</i>	27
<i>Pristobaeus</i> Simon, 1902	27
<i>Phasmolia</i> Clade: <i>Phasmolia</i> , <i>Bindax</i> , <i>Lakarobius</i> and <i>Araneotanna</i>	28
<i>Parabathippus-Parvattus</i> Clade: <i>Parabathippus</i> , <i>Parvattus</i> and possibly <i>Saitissus</i>	28
<i>Euophrys</i> Clade: <i>Chalcoscirtus</i> , <i>Euophrys</i> , <i>Pseudeuophrys</i> , <i>Talavera</i> and <i>Featheroides</i>	29
<i>Saitis</i> Clade: <i>Saitis</i> , <i>Maratus</i> , <i>Jotus</i> , <i>Prostheclina</i> , <i>Hypoblemum</i> , “ <i>Lycidas</i> ”, <i>Maileus</i> , <i>Barraina</i> and possibly <i>Margaromma</i>	29
<i>Laufeia</i> Clade: <i>Laufeia</i> and possibly <i>Magyarus</i>	30
<i>Laufeia</i> Simon, 1889	30
<i>Colyttus</i> Clade	31
<i>Colyttus</i> Thorell, 1891	31
<i>Cytaea-Euryattus</i> Clade: <i>Cytaea</i> , <i>Euryattus</i> , possibly <i>Aruattus</i> and <i>Charippus</i>	31
<i>Thiania</i> Clade	31
<i>Thiania</i> C. L. Koch, 1846	31
<i>Emathis-Lepidemathis</i> Clade	32
<i>Thyenula</i> Clade: <i>Thyenula</i> and <i>Rumburak</i> ; possibly <i>Tanzania</i> , <i>Lophostica</i> , <i>Pseudemathis</i> and <i>Yimbulunga</i>	32
Other genera from the Old World: <i>Chalcotropis</i> , <i>Chinophrys</i> , <i>Foliabitus</i> , <i>Frewena</i> , <i>Lagnus</i> , <i>Servaea</i> , <i>Sigytes</i> , <i>Thorelliola</i> , <i>Viribestus</i> , <i>Viroqua</i> , <i>Xenocytus</i> and <i>Zabkattus</i>	33
Acknowledgements	33
References	34
FIGURES	38
APPENDIX 1	135
APPENDIX 2	141

## Abstract

Morphological traits of euophryine jumping spiders were studied to clarify generic limits in the Euophryinae and to permit phylogenetic classification of genera lacking molecular data. One hundred and eight genera are recognized within the subfamily. Euophryine generic groups and the delimitation of some genera are reviewed in detail. In order to explore the effect of adding formal morphological data to previous molecular phylogenetic studies, and to find morphological synapomorphies, eighty-two morphological characters were scored for 203 euophryine species and seven outgroup species. The morphological dataset does not perform as well as the molecular dataset (genes 28S, Actin 5C; 16S-ND1, COI) in resolving the phylogeny of Euophryinae, probably because of frequent convergence and reversal. The formal morphological data were mapped on the phylogeny in order to seek synapomorphies, in hopes of extending the phylogeny to include taxa for which molecular data are not available. Because of homoplasy, few globally-applicable morphological synapomorphies for euophryine clades were found. However, synapomorphies that are unique locally in subclades still help to delimit euophryine generic groups and genera. The following synonyms of euophryine genera are proposed: *Maeotella* with *Anasaitis*; *Dinattus* with *Corythalia*; *Paradecta* with *Compsodecta*; *Cobanus*, *Chloridusa* and *Wallaba* with *Sidusa*; *Tariona* with *Mopiopia*; *Nebridia* with *Amphidraus*; *Asaphobelis* and *Siloca* with *Coryphasia*; *Ocnotelus* with *Semnolius*; *Palpeilius* with *Pristobaeus*; *Junxattus* with *Laufeia*; *Donoessus* with *Colyttus*; *Nicylla*, *Pselcis* and *Thianitara* with *Thiania*. The new genus *Saphrys* is erected for misplaced species from southern South America.

**Key words:** Salticidae, Euophryinae, generic review, phylogeny, morphology, molecules, new synonym, new combination, new genus

## Introduction

The Euophryinae is one of the largest groups in jumping spiders (Salticidae), with about 1000 species (Prószyński 2012; Zhang & Maddison 2012a, b, c, d; Platnick 2014) reported from all over the world. Most euophryine species are found in the tropics of both the Old World and New World. Spiders of this group have evolved a great variety of body forms: some are relatively large and slender with long legs (e.g. *Bathippus* spp.), some are very robust with short legs and a high carapace (e.g. *Omoedus piceus* Simon), some are flattened (e.g. *Thiania spectrum* (Simon)), some are weevil-like (*Coccorchestes* spp.), and some are ant-like (*Sobasina* spp. and *Paraharmochirus* spp.).

The Euophryinae was initially erected by Eugène Simon (1901) using the name “Evophrydeae”. In Simon’s classification (Simon 1901; 1903), it was considered as a group in Unidentati and comprised three genera: *Akela*, *Euophrys* and *Rhyphelia*, of which *Akela* is not closely related with the other two judging from its morphology (see Galiano 1989). Most of the genera currently considered as euophryines were scattered among more than 20 groups of Pluridentati, Unidentati and Fissidentati in Simon’s classification (see Table 1). More than seventy years after the Euophryinae was erected, Jerzy Prószyński (1976) clarified the group for the first time by a delimiting feature: the presence of a coiled embolus at the distal end of palpal tegulum. Based on this delineation he included 13 genera (*Admestina*, *Agobardus*, *Chalcoscirtus*, *Corythalia*, *Euophrys*, *Habrocestum*, *Laufeia*, *Marchena*, *Neonella*, *Saitis*, *Stoidis*, *Talavera*, *Thiania*) as euophryines in his partial classification of salticids. Among these, *Admestina* is now considered a marpissoid (Maddison & Hedin 2003) and *Marchena* a heliophanine (Maddison 1987). Maddison and Hedin (2003) extended the content of the subfamily and explicitly indicated 34 genera as euophryines (see Table 1). They also revised the delimitation to further specify the particular form of the embolus and tegulum in euophryines: the plane of the spiral of the embolus is more or less parallel to the longitudinal axis of the palp and a loop in the sperm duct projects towards the centre of the tegulum. A recent molecular phylogenetic study on the Euophryinae dramatically extended its content to 85 genera (Zhang & Maddison 2013). A sample of diverse euophryines was shown to be monophyletic in a recent molecular study with eight gene regions (Maddison et al. 2014).

Although the morphological and molecular delimitation for the subfamily Euophryinae is relatively clear, the taxonomy within the subfamily is a mess. There are indications from the molecular phylogeny that similar body forms in different euophryine lineages have arisen by convergence, and therefore have resulted in polyphyletic genera (e.g., the old concept of *Bathippus*). In addition, genitalic organs within this subfamily are relatively simple and usually show little interspecific variation, hindering attempts to classify genera. Most work on this subfamily has concentrated on describing species and genera, or on a regional revision of some genera (e.g. Bryant 1940, 1943, 1950; Balogh 1980a–c; Berry et al. 1996, 1997, 1998; Edwards 2002; Logunov & Azarkina 2008), and there is little systematic work on the group as a whole. Although a few genera have been reviewed in the past

Kuduk, Muse Opiang, Banak Gamui, Jim Robins, Ingi Agnarsson, Jeremy Woon, W. G. Lian and H. Q. Ma. We also would like to express our gratitude to Dr. Gitanjali S. S. Bodner for her kindness in sharing her knowledge and expertise in euophryine jumping spiders; to Dr. Tamás Szűts and Gustavo Ruiz for extensive discussions on morphological characters and classification of jumping spiders, and help in obtaining some early literature; to G. B. Edwards and Dr. Charles Griswold for constructive feedback on this work. Thanks are also due to our Chinese colleagues (Dr. Feng Zhang, Dr. Baoshi Zhang and Dr. Zizhong Yang) for their help in obtaining literature on Chinese salticid fauna. This project was funded by an NSERC Canada Discovery Grant to WPM.

## References

- Balogh, P. (1980a) Studies on the Papuan Attidae (Araneae): *Furculattus maxillosus* gen. et sp. n. *Folia Entomologica Hungarica*, 41, 25–27.
- Balogh, P. (1980b) Studies on the Papuan Attidae (Araneae): *Leptathamas paradoxus* gen. et sp. n. *Folia Entomologica Hungarica*, 41, 29–31.
- Balogh, P. (1980c) The genus *Coccorchestes* Thorell, 1881 (Araneae: Attidae). *Acta Zoologica Hungarica*, 26, 61–88.
- Benjamin, S.P. (2004) Taxonomic revision and phylogenetic hypothesis for the jumping spider subfamily Ballinae (Araneae, Salticidae). *Zoological Journal of the Linnean Society*, 142, 1–82.  
<http://dx.doi.org/10.1111/j.1096-3642.2004.00123.x>
- Berry, J.W., Beatty, J.A. & Prószyński, J. (1996) Salticidae of the Pacific Islands. I. Distributions of twelve genera, with descriptions of eighteen new species. *Journal of Arachnology*, 24, 214–253.
- Berry, J.W., Beatty, J.A. & Prószyński, J. (1997) Salticidae of the Pacific Islands. II. Distribution of nine genera, with descriptions of eleven new species. *Journal of Arachnology*, 25, 109–136.
- Berry, J.W., Beatty, J.A. & Prószyński, J. (1998) Salticidae of the Pacific Islands. III. Distribution of seven genera with descriptions of nineteen new species and two new genera. *Journal of Arachnology*, 26, 149–189.
- Bodner, G.S.S. (2002) *Biodiversity assessment and systematics of Neotropical jumping spiders (Araneae: Salticidae)*. PhD Thesis, 450 pp.
- Bodner, M.R. & Maddison, W.P. (2012) The biogeography and age of salticid spider radiations (Araneae: Salticidae). *Molecular Phylogenetics and Evolution*, 65, 213–240.  
<http://dx.doi.org/10.1016/j.ympev.2012.06.005>
- Bryant, E.B. (1940) Cuban spiders in the Museum of Comparative Zoology. *Bulletin of the Museum of Comparative Zoology*, 86, 247–554.
- Bryant, E.B. (1943) The salticid spiders of Hispaniola. *Bulletin of the Museum of Comparative Zoology*, 92, 445–529.
- Bryant, E.B. (1950) The salticid spiders of Jamaica. *Bulletin of the Museum of Comparative Zoology*, 103, 163–209.
- Chickering, A.M. (1946) The Salticidae of Panama. *Bulletin of the Museum of Comparative Zoology*, 97, 1–474.
- Coddington, J.A. (1990) Ontogeny and homology in the male palpus of orb-weaving spiders and their relatives, with comments on phylogeny (Araneocladia: Araneoidea, Deinopoidea). *Smithsonian Contributions to Zoology*, 496, 1–52.  
<http://dx.doi.org/10.5479/si.00810282.496>
- Cutler, B. (1971) *Darwinneon crypticus*, a new genus and species of jumping spider from the Galápagos Islands (Araneae: Salticidae). *Proceedings of the California Academy of Sciences*, 37, 509–513.
- Davies, V.T. & Żabka, M. (1989) Illustrated keys to the genera of jumping spiders (Araneae: Salticidae) in Australia. *Memoirs of the Queensland Museum*, 27, 189–266.
- Edmunds, M. & Prószyński, J. (2001) New species of Malaysian *Agorius* and *Sobasina* (Araneae: Salticidae). *Bulletin of the British Arachnological Society*, 12, 139–143.
- Edwards, G.B. (2002) A review of the Nearctic jumping spiders (Araneae: Salticidae) of the subfamily Euophryinae north of Mexico. *Insecta Mundi*, 16, 65–75.
- Edwards, G.B., Rinaldi, I.M.P. & Ruiz, G.R.S. (2005) A review of some South American species of jumping spiders (Araneae: Salticidae) described by Mello-Leitão from Brasil, with resolution of the genus *Asaphobelis*. *Biota Neotropica*, 5 (2), 1–31.  
<http://dx.doi.org/10.1590/S1676-06032005000300019>
- Felsenstein, J. (1985) Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, 39, 783–791.  
<http://dx.doi.org/10.2307/2408678>
- Galiano, M.E. (1960) Revisión del género *Tylogonus* Simon, 1902 (Araneae, Salticidae). *Revista de la Sociedad Entomológica Argentina*, 22, 93–103.
- Galiano, M.E. (1962) Los géneros *Amphidraus* Simon y *Marma* Simon (Araneae, Salticidae). *Acta Zoologica Lilloana*, 18, 31–44.
- Galiano, M.E. (1963) Las especies americanas de arañas de la familia Salticidae descritas por Eugène Simon: Redescripciones basadas en los ejemplares típicos. *Physis Buenos Aires (C)*, 23, 273–470.
- Galiano, M.E. (1967) Dos nuevas especies del género *Amphidraus* Simon, 1900 (Araneae, Salticidae). *Physis Buenos Aires (C)*,

- Galiano, M.E. (1968a) Nuevas especies chilenas del género *Evophrys* Koch, C. L., 1834 (Araneae-Salticidae). *Physis Buenos Aires (C)*, 27, 233–243.
- Galiano, M.E. (1968b) Revisión de los géneros *Acragas*, *Amycus*, *Encolpius*, *Hypaeus*, *Mago* y *Noegus* (Salticidae, Araneae). *Revista del Museo Argentino de Ciencias Naturales "Bernardino Rivadavia" (Entomology)*, 2, 267–360.
- Galiano, M.E. (1985) Tres nuevas especies de *Tylogonus* Simon, 1902 (Araneae, Salticidae). *Historia Natural, Corrientes*, 5 (19), 153–159.
- Galiano, M.E. (1988) New species of *Neonella* Gertsch, 1936 (Araneae, Salticidae). *Revue Suisse de Zoologie*, 95, 439–448.
- Galiano, M.E. (1989) Note on the genera *Admestina* and *Akela* (Araneae, Salticidae). *Bulletin of British Arachnological Society*, 8, 49–50.
- Gardzińska, J. (2006) A revision of the spider genus *Ohilimia* Strand, 1911 (Araneae: Salticidae). *Annales Zoologici Warszawa*, 56, 375–385.
- Gardzińska, J. & Patoleta, B. (1997) Notes on the genus *Thorelliola* Strand, 1942 (Araneae: Salticidae). *Memoirs of the Queensland Museum*, 42, 213–222.
- Gardzińska, J. & Żabka, M. (2005) A revision of the spider genus *Chalcolecta* Simon, 1884 (Araneae: Salticidae). *Annales Zoologici Warszawa*, 55, 437–448.
- Gardzińska, J. & Żabka, M. (2006) A revision of the spider genus *Diolenius* Thorell, 1870 (Araneae: Salticidae). *Annales Zoologici Warszawa*, 56, 387–433.
- Goloboff, P.A., Farris, J.S. & Nixon, K.C. (2008) TNT, a free program for phylogenetic analysis. *Cladistics*, 24 (5), 774–786. <http://dx.doi.org/10.1111/j.1096-0031.2008.00217.x>
- Hill, D.E. (2009) Euophryine jumping spiders that extend their third legs during courtship (Araneae: Salticidae: Euophryinae: *Maratus*, *Saitis*). *Peckhamia*, 74.1, 1–27.
- Hill, D.E. & Otto, J.C. (2011) Visual display by male *Maratus pavonis* (Dunn 1947) and *Maratus splendens* (Rainbow 1896) (Araneae: Salticidae: Euophryinae). *Peckhamia*, 89.1, 1–41.
- Jendrzejewska, B. (1995) Genus *Athamas* Pickard-Cambridge, 1877, an unusual salticid from the Pacific area (Araneae: Salticidae). *Genus*, 6, 181–190.
- Lewis, P.O. (2001) A likelihood approach to estimating phylogeny from discrete morphological character data. *Systematic Biology*, 50 (6), 913–925. <http://dx.doi.org/10.1080/106351501753462876>
- Logunov, D.V. (1997) Salticidae of Middle Asia. 4. A review of the genus *Euophrys* (s. str.) C. L. Koch (Araneae, Salticidae). *Bulletin of the British Arachnological Society*, 10, 344–352.
- Logunov, D.V. (1998) *Pseudeuophrys* is a valid genus of the jumping spiders (Araneae, Salticidae). *Revue Arachnologique*, 12 (11), 109–128.
- Logunov, D.V. & Azarkina, G.N. (2008) Two new genera and species of Euophryinae (Aranei: Salticidae) from SE Asia. *Arthropoda Selecta*, 17 (1–2), 111–115.
- Logunov, D.V. & Kronstedt, T. (2003) A review of the genus *Talavera* Peckham and Peckham, 1909 (Araneae, Salticidae). *Journal of Natural History*, 37, 1091–1154. <http://dx.doi.org/10.1080/00222930110098391>
- Maddison, W.P. (1987) *Marchena* and other jumping spiders with an apparent leg-carapace stridulatory mechanism (Araneae: Salticidae: Heliophaninae and Thiodininae). *Bulletin of the British Arachnology Society*, 7, 101–106.
- Maddison, W.P. (1996) Molecular approaches and the growth of phylogenetic biology. In: Ferraris, J.D. & Palumbi, S.R. (Eds.), *Molecular zoology: advances, strategies, and protocols*. Wiley-Liss, New York, pp. 47–63.
- Maddison, W.P., Bodner, M.R. & Needham, K.M. (2008) Salticid spider phylogeny revisited, with the discovery of a large Australasian clade (Araneae: Salticidae). *Zootaxa*, 1893, 46–64.
- Maddison, W.P. & Hedin, M.C. (2003) Jumping spider phylogeny (Araneae: Salticidae). *Invertebrate Systematics*, 17, 529–549. <http://dx.doi.org/10.1071/IS02044>
- Maddison, W.P., Li, D., Bodner, M., Zhang, J., Xu, X., Liu, Q. & Liu, F. (2014) The deep phylogeny of jumping spiders (Araneae, Salticidae). *Zookeys*, 440, 57–87. <http://dx.doi.org/10.3897/zookeys.440.7891>
- Maddison, W.P. & Maddison, D.R. (1992) *MacClade 3: analysis of phylogeny and character evolution*. Sunderland, Sinauer, MA.
- Maddison, W.P. & McMahon, M.M. (2000) Divergence and reticulation among montane populations of the jumping spider *Habronattus pugillis* Griswold. *Systematic Biology*, 49, 400–421. <http://dx.doi.org/10.1080/10635159950127312>
- Maddison, D.R. & Maddison, W.P. (2000) *MacClade 4: analysis of phylogeny and character evolution. Version 4.0*. Sinauer Associates, Sunderland, Massachusetts, 398 pp.
- Maddison, W.P. & Maddison, D.R. (2010) Mesquite: a modular system for evolutionary analysis. Version 2.73. Available from: <http://mesquiteproject.org> (accessed 28 January 2015)
- Maddison, W.P. & Zhang, J. (2011) Chapter 14: Salticid spiders of Papua New Guinea. In: Richards, S.J. & Gamui, B.G. (Eds), *Rapid biological assessments of the Nakanai Mountains and the upper Strickland Basin: surveying the biodiversity of*

- Papua New Guinea's sublime karst environments. RAP Bulletin of Biological Assessment* 60. Conservation International, Arlington, VA, pp. 186–189.
- Makhan, D. (2007) *Soesiladeepakius aschnae* gen. et sp. nov. and *Soesilarishius amrishi* gen. et sp. nov. from Suriname (Araneae: Salticidae). *Calodema Supplementary Papers*, 60, 1–8.
- Masta, S. & Maddison, W.P. (2002) Sexual selection driving diversification in jumping spiders. *Proceedings of the National Academy of Sciences of the United States of America*, 99 (7), 4442–4447.  
<http://dx.doi.org/10.1073/pnas.072493099>
- Otto, J.C. & Hill, D.E. (2012) Notes on *Maratus* Karsch 1878 and related jumping spiders from Australia, with five new species (Araneae: Salticidae: Euophryinae). *Peckhamia*, 103.1, 1–81.
- Oxford, G.S. & Gillespie, R.G. (1998) Evolution and ecology of spider coloration. *Annual Review of Entomology*, 43, 619–643.  
<http://dx.doi.org/10.1146/annurev.ento.43.1.619>
- Patoleta, B. (2008) Description of a new species of *Lagnus* L. Koch, 1879 from Fiji archipelago (Araneae: Salticidae). *Genus*, 19, 717–720.
- Peckham, G.W. & Peckham, E.G. (1901) On spiders of the family Attidae found in Jamaica. *Proceedings of the Zoological Society of London*, 1901 (2), 6–16.
- Peckham, G. & Peckham, E.G. (1907) The Attidae of Borneo. *Transactions of the Wisconsin Academy of Sciences, Arts and Letters*, 15, 603–653.
- Peng, X.J., Yin, C.M., Xie, L.P. & Kim, J.P. (1994) A new genus and two new species of the family Salticidae (Arachnida; Araneae) from China. *Korean Arachnology*, 10, 1–5.
- Petrunkévitch, A. (1930) The spiders of Porto Rico. Part three. *Transactions of the Connecticut Academy of Arts and Sciences*, 31, 1–191.
- Platnick, N.I. (2014) The world spider catalog, version 15. American Museum of Natural History. Available from: <http://research.amnh.org/entomology/spiders/catalog/index.html> (accessed 28 January 2015)
- Posada, D. & Buckley, T.R. (2004) Model selection and model averaging in phylogenetics: advantages of the AIC and Bayesian approaches over likelihood ratio tests. *Systematic Biology*, 53, 793–808.  
<http://dx.doi.org/10.1080/10635150490522304>
- Posada, D. & Crandall, K.A. (1998) Modeltest: testing the model of DNA substitution. *Bioinformatics*, 14 (9), 817–818.  
<http://dx.doi.org/10.1093/bioinformatics/14.9.817>
- Prószyński, J. (1971) Redescriptions of type-species of genera of Salticidae (Aranei). VIII-X. Revision of the subfamily Coccothecinae. *Annales Zoologici, Warszawa*, 28, 153–182.
- Prószyński, J. (1976) Studium systematyczno-zoogeograficzne nad rodziną Salticidae (Aranei) Regionów Palearktycznego i Nearktycznego. *Rozprawy Wyższej Szkoły Pedagogicznej*, 6, 1–260.
- Prószyński, J. (1987) *Atlas rysunków diagnostycznych mniej znanych Salticidae 2*. Zeszyty Naukowe Wyższej Szkoły Rolniczo-Pedagogicznej, Siedlce, 172 pp.
- Prószyński, J. (2012) Salticidae (Araneae) of the World. Available from: <http://www.peckhamia.com/salticidae/> (accessed 28 January 2015)
- Prószyński, J. & Deeleman-Reinhold, C.L. (2012) Description of some Salticidae (Aranei) from the Malay Archipelago. II. Salticidae of Java and Sumatra, with comments on related species. *Arthropoda Selecta*, 21 (1), 29–60.
- Richardson, B.J. (2010) A review of the jumping spider fauna (Araneae: Salticidae) of Chile. *Zootaxa*, 2418, 1–49.
- Richardson, B.J. (2013) New unidentate jumping spider genera (Araneae: Salticidae) from Australia. *Zootaxa* 3716 (3), 460–474.  
<http://dx.doi.org/10.11646/zootaxa.3716.3.8>
- Richardson, B.J. & Żabka, M. (2007) A revision of the Australian jumping spider genus *Prostheclina* Keyserling, 1892 (Araneae: Salticidae). *Records of the Australian Museum*, 59, 79–96.  
<http://dx.doi.org/10.3853/j.0067-1975.59.2007.1471>
- Roewer, C.F. (1938) Araneae. In: *Résultats scientifiques du Voyage aux indes orientales néerlandaises de la SS. AA. RR. le Prince et la Princesse*
- Ruiz, G.R.S. (2011) Systematics of the genus *Soesilarishius* (Araneae: Salticidae: Euophryinae). *Zootaxa*, 3022, 22–38.
- Ruiz, G.R.S. & Brescovit, A.D. (2005) Notes on the Venezuelan jumping spiders described by Caporiacco (Araneae, Salticidae). *Revista Brasileira de Zoologia*, 22, 753–760.  
<http://dx.doi.org/10.1590/S0101-81752005000300036>
- Simon, E. (1901) *Histoire naturelle des araignées. Vol. 2*. Paris, 288 pp. [pp. 381–668]
- Simon, E. (1903) *Histoire naturelle des araignées. Vol. 2*. Paris, 412 pp. [pp. 669–1080]
- Swofford, D.L. (2002) *PAUP.\* Phylogenetic analysis using parsimony (\*and other methods). Version 4.0b10*. Sinauer Associates: Sunderland, Massachusetts, USA.
- Szűts, T. (2002) Description of a new horned jumping spider species *Thorelliola mahunkai* (Araneae: Salticidae) from New Guinea. *Folia Entomologica Hungarica*, 63, 17–22.
- Szűts, T. (2003) On remarkable jumping spiders (Araneae: Salticidae) from Papua New Guinea. *Folia Entomologica Hungarica*, 64, 41–57.
- Szűts, T. & De Bakker, D. (2004) Taxonomical relevance of the clypeal setae in *Thorelliola* (Araneae: Salticidae). *Folia Entomologica Hungarica*, 65, 13–24.

- Thorell, T. (1881) Studi sui Ragni Malesi e Papuani. III. Ragni dell'Austro Malesia e del Capo York, conservati nel Museo civico di storia naturale di Genova. *Annali del Museo Civico di Storia Naturale di Genova*, 17, 1–727.
- Wanless, F.R. (1978) A revision of the spider genus *Sobasina* (Araneae: Salticidae). *Bulletin of the British Museum (Natural History). Zoology*, 33, 245–257.
- Wanless, F.R. (1984) A revision of the spider genus *Cyrba* (Araneae: Salticidae) with the description of a new presumptive pheromone dispersing organ. *Bulletin of the British Museum (Natural History) Zoology*, 47, 445–481.
- Wanless, F.R. (1988) A revision of the spider group Astieae (Araneae: Salticidae) in the Australian region. *New Zealand Journal of Zoology*, 15, 81–172.  
<http://dx.doi.org/10.1080/03014223.1988.10422611>
- Wanless, F.R. & Lubin, Y.D. (1986) *Diolenius minotaurus* sp. nov., a remarkable horned jumping spider from Papua New Guinea (Araneae: Salticidae). *Journal of Natural History*, 20, 1211–1220.  
<http://dx.doi.org/10.1080/00222938600770801>
- Wesołowska, W. (2001) Two new species of *Thyemula* Simon, 1902 from South Africa (Araneae: Salticidae). *Annales Zoologici Warszawa*, 51, 261–264.
- Wesołowska, W., Azarkina, G.N. & Russell-Smith, A. (2014) Euophryine jumping spiders of the Afrotropical Region—new taxa and a checklist (Araneae: Salticidae: Euophryinae). *Zootaxa*, 3785 (1), 1–72.  
<http://dx.doi.org/10.11646/zootaxa.3785.1.1>
- Wheeler, T.J. & Kececioglu, J.D. (2007) Multiple alignments by aligning alignments. *Bioinformatics*, 23, 559–568.  
<http://dx.doi.org/10.1093/bioinformatics/btm226>
- Żabka, M. (1985) Systematic and zoogeographic study on the family Salticidae (Araneae) from Viet-Nam. *Annales Zoologici Warszawa*, 39, 197–485.
- Żabka, M. (1987) Salticidae (Araneae) of Oriental, Australian and Pacific Regions, II. Genera *Lycidas* and *Maratus*. *Annales Zoologici, Warszawa*, 40 (1), 451–482.
- Żabka, M. (1996) *Bulolia*, a new genus of Salticidae (Arachnida: Araneae) from Papua New Guinea. *Revue suisse de Zoologie*, Vol. Hors Série 2, 701–707.
- Żabka, M. & Pollard, S. D. (2002) Salticidae (Arachnida: Araneae) of New Zealand: genus *Hypoblemum* Peckham and Peckham, 1886. *Records of Canterbury Museum*, 16, 64–72.
- Żabka, M. & Prószyński, J. (1998) Middle European *Euophrys* C. L. Koch, 1834 (Araneae: Salticidae)—one, two or three genera? In: Seldon, P.A. (Ed.), *Proceedings of the 17th European Colloquium of Arachnology, Edinburgh 1997*. Edinburgh, pp. 115–120.
- Zhang, J. & Maddison, W.P. (2012a) New euophryine jumping spiders from the Dominican Republic and Puerto Rico (Araneae: Salticidae: Euophryinae). *Zootaxa*, 3476, 1–54.
- Zhang, J. & Maddison, W.P. (2012b) New euophryine jumping spiders from Papua New Guinea (Araneae: Salticidae: Euophryinae). *Zootaxa*, 3491, 1–74.
- Zhang, J. & Maddison, W.P. (2012c) New euophryine jumping spiders from Southeast Asia and Africa (Araneae: Salticidae: Euophryinae). *Zootaxa*, 3581, 53–80.
- Zhang, J. & Maddison, W.P. (2012d) New euophryine jumping spiders from Central and South America (Araneae: Salticidae: Euophryinae). *Zootaxa*, 3578, 1–35.
- Zhang, J. & Maddison, W.P. (2013) Molecular phylogeny, divergence times and biogeography of the subfamily Euophryinae (Araneae: Salticidae). *Molecular Phylogenetics and Evolution*, 68 (1), 81–92.  
<http://dx.doi.org/10.1016/j.ympev.2013.03.017>
- Zhang, J. & Maddison, W.P. (2014) *Tisaniba*, a new genus of marpissoid jumping spiders from Borneo (Araneae: Salticidae). *Zootaxa*, 3852 (2), 252–272.  
<http://dx.doi.org/10.11646/zootaxa.3852.2.5>
- Zwickl, D.J. (2006) *Genetic algorithm approaches for the phylogenetic analysis of large biological sequence datasets under the maximum likelihood criterion*. PhD Dissertation. The University of Texas, Austin, 115 pp.