

# Generalisations of the Laplace–Runge–Lenz Vector

*P G L LEACH*<sup>†1†2†3</sup> and *G P FLESSAS*<sup>†1†4</sup>

<sup>†1</sup> *GEODYSYC, School of Sciences, University of the Aegean, Karlovassi 83 200, Greece*

<sup>†2</sup> *Department of Mathematics, University of the Aegean, Karlovassi 83 200, Greece*

<sup>†3</sup> *Permanent address: School of Mathematical and Statistical Sciences,  
University of Natal, Durban 4041, Republic of South Africa*

<sup>†4</sup> *Department of Information and Communication Systems Engineering,  
University of the Aegean, Karlovassi 83 200, Greece*

*Received October 17, 2002; Revised January 22, 2003; Accepted January 27, 2003*

## Abstract

The characteristic feature of the Kepler Problem is the existence of the so-called Laplace–Runge–Lenz vector which enables a very simple discussion of the properties of the orbit for the problem. It is found that there are many classes of problems, some closely related to the Kepler Problem and others somewhat remote, which share the possession of a conserved vector which plays a significant rôle in the analysis of these problems.

## Contents

<b>1</b>	<b>Introduction</b>	<b>341</b>
<b>2</b>	<b>Motions with conservation of the angular momentum vector</b>	<b>346</b>
2.1	The central force problem $\ddot{\mathbf{r}} + f\mathbf{r} = 0$ . . . . .	346
2.1.1	The conservation of the angular momentum vector, the Kepler problem and Kepler’s three laws . . . . .	346
2.1.2	The model equation $\ddot{\mathbf{r}} + f\mathbf{r} = 0$ . . . . .	350
2.2	Algebraic properties of the first integrals of the Kepler problem . . . . .	355
2.3	Extension of the model equation to nonautonomous systems . . . . .	356

<b>3</b>	<b>Conservation of the direction of angular momentum only</b>	<b>360</b>
3.1	Vector conservation laws for the equation of motion $\ddot{\mathbf{r}} + g\hat{\mathbf{r}} + h\hat{\boldsymbol{\theta}} = 0$ . . . . .	360
3.2	The orbit equation and motion in time . . . . .	362
3.3	The related Hamiltonian system . . . . .	364
3.4	The Lie algebra of the Poisson Brackets of the first integrals . . . . .	365
3.5	Invariance under time translation and the first integrals . . . . .	369
3.6	The Kepler problem with ‘drag’ . . . . .	372
3.7	Force laws admitting Keplerian orbits . . . . .	377
3.8	The geometry of the generalised Kepler problem . . . . .	380
<b>4</b>	<b>Motion with conserved <math>\mathbf{L}</math></b>	<b>385</b>
4.1	A conserved Laplace–Runge–Lenz vector for $\ddot{\mathbf{r}} + f\mathbf{L} + g\hat{\mathbf{r}} = 0$ . . . . .	385
4.2	The classical MICZ problem . . . . .	390
<b>5</b>	<b>Symmetries and conserved vectors</b>	<b>394</b>
5.1	Introduction . . . . .	394
5.2	The Lie point symmetries of $\ddot{\mathbf{r}} + f(r)\mathbf{L} + g(r)\mathbf{r} = 0$ . . . . .	396
5.2.1	The case $f = 0$ . . . . .	397
5.2.2	The case $f \neq 0$ . . . . .	398
5.2.3	Comment . . . . .	399
5.3	The equation $\ddot{\mathbf{r}} + f\dot{\mathbf{r}} + \mu g\mathbf{r} = 0$ . . . . .	400
5.4	Complete symmetry groups . . . . .	403
5.5	The method of reduction of order . . . . .	405
5.6	The symmetries of the reduced Kepler problem . . . . .	406
5.7	The reduced system for some problems possessing a conserved vector of Laplace–Runge–Lenz type . . . . .	409
5.7.1	The Kepler problem with drag . . . . .	409
5.7.2	The equation $\ddot{\mathbf{r}} - \frac{1}{2} \left( \frac{\dot{g}}{g} + 3\frac{\dot{r}}{r} \right) \dot{\mathbf{r}} + \mu g\mathbf{r} = 0$ . . . . .	411
5.7.3	An example with an angle-dependent force . . . . .	412
5.7.4	A truly three-dimensional motion . . . . .	414
	<b>References</b>	<b>416</b>

## 1 Introduction

Cosmology and its precursor, Cosmogony, have a very long history dating back to the earliest records of ancient civilisations. The practical tasks of Astronomy and its more imaginative relation, Astrology, seemed to have prompted speculation as to the nature of the Universe almost as an automatic concomitant. One aspect of these speculations related to the size and distribution of the sun and its planets. In classical times there were two schools of thought about the structure of the solar system. One placed the Earth at the centre of the Universe with the Sun, Moon and the then known planets revolving around it, the geocentric hypothesis, and the other had the Earth and the planets revolving about the Sun, the heliocentric hypothesis. The latter was proposed about 250 BC by Aristarchos of Samos. The former was the majority opinion of classical philosophers, endorsed by one of the more influential opinion-formers of that period, Aristotle, and adopted by the computer

of orbits, Claudius Ptolemaos. Aristarchos found no support until the present era. The dominance of Scholasticism in the Mediæval Ages and its dominance by Aristotlean Physics in the profane realm meant that the hypothesis of Aristarchos was known to very few by the time of the Renaissance. The heliocentric hypothesis was revived by the Pole, Nicolas Copernicus, in his book *Revolutionibus Orbium Coelestium* published in 1543. Evidently Copernicus knew of the work of Aristarchos of Samos, but the reference was deleted from the first edition, apparently by its theologically cautious publisher, the Lutheran Andrias Osiander [29, p. 319]. So effectively was the contribution of Aristarchos neglected that we find the chronicler of his work, Sir Thomas Heath, moved to entitle the book *Aristarchus of Samos: the ancient Copernicus* [58] and we find the historian of Mechanics, Dugas [30, p. 83], writing ‘From Antiquity there had been writers whose opinions were similar to those of Copernicus’! In an attempt to counteract such neglect the municipality of the town of Karlovassi on the island of Samos erected a bust of the island’s second most famous son in the main square some 200 m from where this is written with a legend (in English; one assumes that the Greeks would not be so ignorant) indicating that Copernicus was an 1800 year Johnny-come-lately.

The timing of Copernicus was not perfect. The religious turmoil which was to shake Europe for a century and a half had commenced about a quarter of a century before the book was published. Not only did physicists, entirely entangled in Aristotlean ideas, take exception to the impossible strains placed upon the fabric of the Earth and its environment, but the theologians – particularly on the reformed side – categorically rejected the theory of the motion of the Earth because they considered it contrary to Holy Writ [19, p. 22].

The eventual triumph of the heliocentric hypothesis can reasonably be attributed to the work of Kepler who, painstakingly analysing his observations and those of his predecessor as Imperial Mathematician at Prague, Tycho Brahe, endowed the hypothesis with three empirical laws. In the books *Astronomia Nova* of 1609 [73] and *Harmonices Mundi* of 1619 [74] Kepler proposed

*Itaque plane hoc est: orbita planetæ non est circulus, sed ingrediens ad latera utraque paulatim, iterumque ad circuli amplitudinem in perigæo exiens: cujusmodi figuram itineris ovalem appellitant. . . . orbitam planetæ non esse circulum, sed figura ovalis.* [73, pp. 336–337]<sup>1.1</sup>

*Cumque scirem, infinita esse puncta eccentrici, et distantias earum infinitas, subiit, in plano eccentrici has distantias omnes inesse. Nam memineram, sic olim et Archimedes, cum circumferentiæ proportionem ad diametrum quæreret, circulum in infinita triangula dissecuisse . . .* [73, p. 321]<sup>1.2</sup>

*Sed res est certissima, quod proportio, quæ est inter binorum quorumcunque planetarum tempora periodica, sit præcise sesquialtera proportionis mediarum distantiarum, id*

---

<sup>1.1</sup>Kepler’s First Law: ‘Therefore this is obvious: the path of the planet is not a circle, but gradually curves inwards on both sides and again departs to the width of the orbit at perigee: such a path is called an oval figure. . . . the orbit of the planet is not a circle but an oval figure.’

<sup>1.2</sup>Kepler’s Second Law: ‘Since I was aware that there exists an infinite number of points on the orbit and accordingly an infinite number of distances [from the Sun] it occurred to me that the sum of these distances is contained in the area of the orbit. For I remembered that in the same manner in times past, Archimedes too divided [the area] of a circle, for which he found the circumference proportional to the diameter, into an infinite number of triangles . . .’

*est orbium ipsorum . . .* [74, p. 279]<sup>1,3</sup>

Johannes Kepler had presented simple quantitative facts about the orbits of the planets. In August, 1684, Edmund Halley visited Issac Newton at Trinity College, Cambridge, and informed him of Christopher Wren’s challenge, to which there was attached a sizeable monetary reward, to derive the shape of the orbits of the planets about the sun. Newton described his solution in the tract *De Motu Corporum* and set about improving and expanding his ideas. During this period he proposed his law of universal gravitation and his three laws of motion which provided the first quantitative explanation of the motion of visible bodies. In 1686 these results were published in Newton’s masterpiece, *Philosophiæ Naturalis Principia Mathematica*, which has been claimed to mark the birth of modern science [41, p. 3], if one can isolate one moment in a period of evolution.

In the *Principia* Newton attempted to verify Kepler’s First Law by proving that, if a particle moves on an elliptical orbit under the influence of a centripetal force directed towards one focus, that force must be inversely proportional to the square of the radius [113, Prop. XI, p. 56]. He obtained the same results for focus-centred parabolic [113, Prop. XIII, p. 60] and hyperbolic motion [113, Prop. XII, p. 57]. Newton proved that equal areas were swept out by the radius vector in equal times [113, Propp. I–II, p. 40ff], which was the content of Kepler’s Second Law, and also that the square of the periodic time of an inverse square law ellipse was proportional to the cube of the semimajor axis length, thereby verifying Kepler’s Third Law [113, Prop. XV, p. 62].

Newton is also generally credited with having proved the converse of Propositions XI–XIII, *ie* that all possible orbits of a particle moving under an inverse square law force are conic sections. This is of greater significance than the proof of the Propositions above since this result verifies Newton’s law of universal gravitation by showing that elliptical planetary orbits are a natural consequence of the gravitational interaction between the Sun and the planets. The validity of this credit has been questioned by Weinstock [138, 139] who argues that proof that motion along a conic section orbit under the influence of a force directed towards one focus implies an inverse square law force does not demonstrate that a particle moving under the influence of an inverse square law force will move along a conic section orbit. The conic section orbit, which was assumed initially, is only a geometrical construction and not necessarily the actual path described by the particle.

The initial approach was to determine the gravitational force law assuming that the planets move along conic sections. In 1710 Ermanno<sup>1,4</sup>, a student of Johann Bernoulli, used the new techniques of the Leibnizian calculus to find directly the orbits given an inverse square law force. Ermanno obtained a constant of integration related to the eccentricity of the conic section [33] and summarised his results in a letter [59] to Johann Bernoulli who proceeded to generalise the results to allow for an arbitrary orientation of the orbit in the plane [12]<sup>1,5</sup>. The Ermanno–Bernoulli constants are the equivalents of the conserved

---

<sup>1,3</sup>Kepler’s Third Law: ‘But it is absolutely certain and exact that the ratio which exists between the periodic times of any two planets is precisely the ratio of the 3/2th power of the mean distances, *ie* of the spheres themselves . . .’

<sup>1,4</sup>Ermanno made a considerable impact on Italian Mathematics during his brief sojourn at Padova (1707–1713) and his place in history is based more on that. For an account of his work and extracts of many original letters and documents see the work of Mazzone and Poero [105].

<sup>1,5</sup>Some aspects of the Kepler Problem in the period from Newton to Johann Bernoulli are to be found in Speiser [132].

vector generally known as the Laplace–Runge–Lenz vector [78, 127, 93]. Almost 90 years later in his *Traité de Mécanique Céleste* [78] Laplace obtained seven first integrals for the Kepler problem. These were the three cartesian components of the angular momentum, the three cartesian components of the Laplace–Runge–Lenz vector and what was essentially the total energy. Since only five of these autonomous functions could be independent, Laplace determined the two relationships between the seven integrals. He showed that the conservation of the angular momentum implied that the orbit was planar and that the orbit equation could be expressed as a conic section [38].

In July 1845 Hamilton presented to the Royal Irish Academy a new constant of the motion for the Kepler problem which he subsequently named the ‘eccentricity’ vector [55] and which is commonly called Hamilton’s vector. In another of the ironies of science the deviser of vector notation, Josiah Gibbs, is believed to have been the first to express the classical Laplace–Runge–Lenz vector in terms of modern vector notation, thereby antedating Runge’s derivation by roughly twenty years [39].

In 1785 Charles Coulomb discovered the inverse square law of the electrostatic force. In 1913 Neils Bohr used the Kepler–Coulomb potential energy for a Rutherford model of the hydrogen atom to obtain a theoretical formula for the wavelengths of the atomic spectrum which agreed with experimental observations. The Laplace–Runge–Lenz vector<sup>1,6</sup> came to prominence with the mathematical development of Quantum Mechanics. In 1926 Pauli [119] used the vector to derive an expression for the energy levels of the hydrogen atom using the new techniques of matrix mechanics. In 1935 Fock [34] showed that the hydrogen atom possessed the four-dimensional rotation group for the bound states and the Lorentz group for positive energy states. These results were confirmed by Bargmann [10] in 1936 when he calculated the commutation relations between the components of the angular momentum and the Laplace–Runge–Lenz vectors.

The characteristic feature of the Kepler problem is the existence of the conserved vectors which enable one to determine the orbit easily. In fact what is potentially a very difficult nonlinear system can be trivially integrated [127, 14, 22, 23] to obtain the orbit and the only complexity is determining the time evolution of system. These conserved vectors play the same role for the Kepler problem as do the tensorial first integrals – the angular momentum tensor and the Jauch–Hill–Fradkin tensor [65, 35] – for that other great paradigm of Mechanics, the isotropic harmonic oscillator. What is missing for the Kepler problem are the explicitly time-dependent tensor invariants which enable one easily to specify the trajectory in the configuration space of the oscillator [92]. The Kepler problem is not the only problem in Mechanics for which there exists conserved vectors. Already in 1896 Poincaré had identified a conserved vector with properties reminiscent of angular momentum for the classical monopole problem [123].

The advent of the space age brought about some generalisations of the Kepler problem to deal with the motion of low altitude satellites, a model was proposed by Danby in 1962 [27]. Danby was able to provide an approximate solution. In the early Eighties Jezewski and Mittleman [107, 66] were able to demonstrate the existence of first integrals

---

<sup>1,6</sup>Known amongst physicists as the Lenz vector following the 1924 paper of Lenz [93] or the Runge–Lenz vector since Lenz had referred to the textbook on vector analysis by Runge [127], the contribution by Laplace has been noticed in the title of the vector only in recent decades. The lack of recognition of Ermanno and Bernoulli parallels that of Aristarchos and makes one wonder if there is some sinister influence for those who are initiators in the field of the motions of the planets.

for this problem. Subsequently many problems which could loosely be regarded as relaxations of the Kepler problem were shown to possess vector first integrals which were sufficient to provide the orbit just as the angular momentum and Laplace–Runge–Lenz vectors provided for the Kepler problem [42, 43, 44, 45, 46, 47, 48, 49, 81, 83, 87, 88, 89].

In 1994 Krause [75] used the Kepler problem as the vehicle for the introduction of his concept of the complete symmetry group of a differential equation or system of differential equations. Essentially the complete symmetry group of a differential equation is the group of the algebra of the Lie symmetries required to specify completely the differential equation. In the case of the Kepler problem the Lie symmetries required included nonlocal symmetries although it is not the case for every differential equation [5]. In the subsequent investigations of the complete symmetry groups of the Kepler and related problems by the technique of reduction of order [117] by Nucci *et al* [118] the underlying symmetry was found to be the same for a number of these problems and the critical role of the Ermanno–Bernoulli constants clearly delineated.

It is the purpose of this review to bring the details of these divers problems together and to highlight the essential role played by certain requirements on the equations of motion. Our work may be divided into three parts with the nature of the division being dependent upon the angular momentum. In the case of the Kepler problem the angular momentum is conserved. In Danby’s model for the motion of a low altitude satellite the magnitude of the angular momentum is no longer conserved. Poincaré’s comments upon the experiment of Birkland led to the introduction of a generalisation of the angular momentum as the conserved vector. The magnitude of the angular momentum was constant but not its direction. Consequently we discuss the three classes of problem, that of the conservation of angular momentum, that of the conservation of the direction of angular momentum and that of the conservation of the magnitude of angular momentum. Finally we look to the symmetry properties of these problems to reconcile the results. In this we must go beyond the point symmetries initially introduced by Lie [98, 99] to the nonlocal symmetries introduced by Krause [75] to construct the complete symmetry group of the Kepler problem and extended in both number and scope by Nucci *et al* [118].

In the various generalisations of the Kepler Problem which we treat in this paper we generally start with an equation of motion which has a certain structure consistent with the angular momentum model under consideration. These equations of motion contain arbitrary functions of, usually, unspecified variable dependence. Part of the task of the analysis is to identify the nature of the variable dependence in these functions which is possible for a conserved vector having the nature of the Laplace–Runge–Lenz vector to exist. Consequently the reader should understand that the arguments of these arbitrary functions become evident during the process of analysis. Usually the function itself remains of arbitrary form. Specific functions are used by way of example.

The position taken here in respect to the identification of these vectors as being generalisations of the Laplace–Runge–Lenz vector is based on the notion of function rather than form. The Laplace–Runge–Lenz vector of the Kepler Problem has a specific functional structure. In particular it is quadratic in the components of the velocity. Some of the conserved vectors found here are very evidently not of that structure. However, they serve the same function for their equations of motion as the Laplace–Runge–Lenz vector does for the equation of motion of the Kepler Problem. All of these conserved vectors provide a simple route to the determination of the equation of the orbit, *ie* they perform the same

function. To argue that for a conserved vector to be associated with the notion of the Laplace–Runge–Lenz vector means that the vector should be quadratic in the components of the velocity is the equivalent of using a sample of one to specify the properties of a population, a practice which is clearly statistically incorrect.

This preferential option for function over form has been given renewed strength by the results of Nucci *et al* [118] discussed in Chapter 5. There the various models treated in earlier chapters are seen to be reducible to the same linear system so that the different types of conserved vectors can in fact be regarded as different manifestations of the same object.

## 2 Motions with conservation of the angular momentum vector

### 2.1 The central force problem $\ddot{\mathbf{r}} + f\mathbf{r} = 0$

#### 2.1.1 The conservation of the angular momentum vector, the Kepler problem and Kepler’s three laws

The techniques used for this and other problems to obtain conserved vectors and scalars and equations describing the motion are natural extensions of the use of scalar and vector products in the analysis of the reduced equation for the Kepler problem (*cf* the works of Runge [127], Bleuler and Kustaanheimo [14], Collas [21], Collinson [22, 23] and Pollard [124]), *videlicet*

$$\ddot{\mathbf{r}} + \frac{\mu\mathbf{r}}{r^3} = 0, \quad (2.1.1)$$

in which the motion of the centre of mass of the system of two particles has already been removed. Such other equations as we shall study are to be regarded as generalisations of this reduced equation, (2.1.1). The vector product of  $\mathbf{r}$  with (2.1.1) immediately gives the conservation of the angular momentum vector

$$\mathbf{L} := \mathbf{r} \times \dot{\mathbf{r}}, \quad (2.1.2)$$

from which we factor the mass constant. Since  $\mathbf{L}$  is perpendicular to both  $\mathbf{r}$  and  $\dot{\mathbf{r}}$  for all time, the constancy of  $\mathbf{L}$  implies that the orbit is confined to a plane.

In the spirit of using vectorial methods to obtain the Laplace–Runge–Lenz and Hamilton’s vectors there are two possible routes. For one of them we take the vector product of (2.1.1) with  $\mathbf{L}$  and the relationship

$$\mathbf{r} \times \mathbf{L} = -r^3 \dot{\hat{\mathbf{r}}}, \quad (2.1.3)$$

where  $\hat{\mathbf{r}}$  denotes the unit vector in the direction of  $\mathbf{r}$ , to obtain

$$\begin{aligned} 0 &= \ddot{\mathbf{r}} \times \mathbf{L} - \mu \dot{\hat{\mathbf{r}}} \\ \Rightarrow \mathbf{J} &= \dot{\mathbf{r}} \times \mathbf{L} - \mu \hat{\mathbf{r}}, \end{aligned} \quad (2.1.4)$$

where  $\mathbf{J}$  is the Laplace–Runge–Lenz vector. Hamilton's vector is obtained as

$$\begin{aligned}\mathbf{K} &= \frac{1}{L^2} (\mathbf{L} \times \mathbf{J}) \\ &= \dot{\mathbf{r}} + \frac{\mu r^2 \dot{\hat{\mathbf{r}}}}{L^2},\end{aligned}\tag{2.1.5}$$

where  $\hat{\mathbf{r}} \times \mathbf{L}$  is  $r^2 \dot{\hat{\mathbf{r}}}$  and  $L$  is the constant magnitude of  $\mathbf{L}$ . For the second route we make use of the fact that the orbit is in a plane to specify plane polar coordinates,  $(r, \theta)$ , in that plane so that the three unit vectors,  $\hat{\mathbf{r}}$ ,  $\hat{\boldsymbol{\theta}}$  and  $\hat{\mathbf{L}}$  have the properties that

$$\begin{aligned}\hat{\mathbf{r}} \times \hat{\boldsymbol{\theta}} &= \hat{\mathbf{L}}, & \hat{\boldsymbol{\theta}} \times \hat{\mathbf{L}} &= \hat{\mathbf{r}}, & \hat{\mathbf{L}} \times \hat{\mathbf{r}} &= \hat{\boldsymbol{\theta}}, \\ \frac{d\hat{\mathbf{r}}}{dt} &= \dot{\theta} \hat{\boldsymbol{\theta}}, & \frac{d\hat{\boldsymbol{\theta}}}{dt} &= -\dot{\theta} \hat{\mathbf{r}}, & L &= r^2 \dot{\theta}.\end{aligned}\tag{2.1.6}$$

Then (2.1.1) can be written as

$$\begin{aligned}0 &= \ddot{\mathbf{r}} + \frac{\mu}{L} \dot{\theta} \hat{\mathbf{r}} \\ \Rightarrow \mathbf{K} &= \dot{\mathbf{r}} - \frac{\mu}{L} \hat{\boldsymbol{\theta}}.\end{aligned}\tag{2.1.7}$$

The two routes are reconciled when we express (2.1.5) in terms of the plane polar coordinates, which the constancy of the angular momentum vector enables us to do.

From the integration of the scalar product of (2.1.1) with  $\dot{\mathbf{r}}$  we obtain the conserved scalar

$$E := \frac{1}{2} \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - \frac{\mu}{r}\tag{2.1.8}$$

which is the mechanical energy of the system scaled by the mass. From the scalar products of (2.1.7) and (2.1.4) with themselves we obtain the equivalent relationships

$$K^2 = 2E + \frac{\mu^2}{L^2}\tag{2.1.9}$$

$$J^2 = 2L^2 E + \mu^2,\tag{2.1.10}$$

which give two of the constraints necessary for the ten first integrals,  $\mathbf{J}$ ,  $\mathbf{K}$ ,  $\mathbf{L}$  and  $E$ . The other constraint is the mutual orthogonality of the three conserved vectors.

Finally, if we choose the conserved vector,  $\mathbf{J}$ , as the reference line for the measurement of the polar angle,  $\theta$ , the scalar product of  $\mathbf{r}$  and  $\mathbf{J}$  gives

$$\begin{aligned}\mathbf{r} \cdot \mathbf{J} &= rJ \cos \theta = \mathbf{r} \cdot (\dot{\mathbf{r}} \times \mathbf{L}) - \mu \mathbf{r} \cdot \hat{\mathbf{r}} \\ \Rightarrow r &= \frac{L^2}{\mu + J \cos \theta}\end{aligned}\tag{2.1.11}$$

which is the polar equation for a conic section with the origin at one focus. Thus we have the orbit for the motion governed by the equation (2.1.1). The time evolution of the system may be obtained, for example, by means of the constancy of the magnitude of the



angular momentum for then we have that the relationship  $L = r^2\dot{\theta}$  can be combined with (2.1.11) to give

$$t - t_0 = \int_{\theta_0}^{\theta} \frac{L^3 d\theta}{(\mu + J \cos \theta)^2} \quad (2.1.12)$$

which is moderately simple to evaluate. In the constancy of the vector of angular momentum and the specific equation for the orbit we have two of Kepler's three laws. For a closed orbit the third law follows from the evaluation of (2.1.12) over an orbit to give the period,  $T$ , as ([51, p. 148, 2.553.3 and 2.554.3])

$$T = \frac{2\pi\mu L^3}{(\mu^2 - J^2)^{3/2}} = \frac{2\pi\mu}{(-2E)^{3/2}} \quad (2.1.13)$$

and the use of (2.1.10) and (2.1.11) to relate the semimajor axis length,  $R$ , to the energy,  $E$ , according to

$$2ER + \mu = 0 \quad \Leftrightarrow \quad R = \frac{\mu}{-2E}. \quad (2.1.14)$$

Consequently we obtain

$$\frac{T^2}{R^3} = \frac{2\pi}{\mu} \quad (2.1.15)$$

in a way somewhat different compared with the manner in which Kepler obtain the result almost 400 years ago.

It is unfortunate that not all can be calculated for the Kepler problem by these elementary methods. The evolution of the motion in time is not so simple to calculate<sup>2.1</sup>. In (2.1.12) we have the equation to be integrated to obtain the evolution of the polar angle,  $\theta$ , in time. If we differentiate the orbit equation, (2.1.11), with respect to time, we obtain

$$dt = \frac{Ldr}{J \sin \theta}. \quad (2.1.16)$$

We may eliminate  $\sin \theta$  by using the orbit equation, (2.1.12), to write

$$\sin \theta = \frac{1}{Jr} ((J^2 - \mu^2)r^2 + 2L^2\mu r - L^4)^{\frac{1}{2}} \quad (2.1.17)$$

so that (2.1.16) becomes

$$dt = \frac{rdr}{((J^2 - \mu^2)r^2/L^2 + 2\mu r - L^2)^{\frac{1}{2}}}. \quad (2.1.18)$$

Naturally, if we use (2.1.10), (2.1.18) becomes the same equation as we would obtain from the energy expression after the substitution  $\dot{\theta} = L/r^2$ . When we make this substitution and perform the integration ([51, 2.264.2 and 2.261.3]), we obtain

$$t - t_0 = \frac{(2Er^2 + 2\mu r - L^2)^{\frac{1}{2}}}{2E} + \frac{\mu}{2E(-2E)^{\frac{1}{2}}} \arcsin \left( \frac{2Er + \mu}{(2EL^2 + \mu^2)^{\frac{1}{2}}} \right) \Bigg|_{r_0}^r. \quad (2.1.19)$$

---

<sup>2.1</sup>A difficulty recognised very early in the study of the Kepler Problem through the solution of Kepler's equation, cf [31] and [24], for which see below.

Although it is a simple matter to recover Kepler's Third Law from this, the inversion to obtain an explicit function,  $r(t)$ , is not globally possible. The integration of (2.1.12) gives an equally unpleasant expression [51, 2.554.3 and 2.553.3] for the relationship between  $\theta$  and  $t$ , *videlicet*

$$t - t_0 = \frac{L}{2E} \left( \frac{J \sin \theta}{\mu + J \cos \theta} - \frac{2\mu}{L(-2E)^{\frac{1}{2}}} \arctan \frac{L(-2E)^{\frac{1}{2}} \tan \frac{1}{2}\theta}{\mu + J} \right). \quad (2.1.20)$$

The same comment applies.

There is another approach which can be taken and which dates back for several centuries. The standard equation for a conic section in polar coordinates is

$$r = \frac{1}{c(1 + e \cos \theta)}, \quad (2.1.21)$$

where  $c$  and the eccentricity,  $e$ , are constants. It is evident that

$$e = \frac{J}{\mu} \quad (2.1.22)$$

and so we may write

$$L^2 = \mu R(1 - e^2). \quad (2.1.23)$$

consequently (2.1.18) can be rewritten as

$$dt = \frac{r dr}{(-\mu r^2/R + 2\mu r - \mu R(1 - e^2))^{\frac{1}{2}}}. \quad (2.1.24)$$

The introduction of another angular variable,  $\psi$ , which is known as the eccentric anomaly (in contrast to  $\theta$  which has been known as the true anomaly since mediæval times [40, p. 99]), through the substitution

$$r = R(1 - e \cos \psi) \quad (2.1.25)$$

simplifies (2.1.24) to

$$dt = \left( \frac{R^3}{\mu} \right)^{\frac{1}{2}} (1 - e \cos \psi) d\psi \quad (2.1.26)$$

which is trivially integrated to give

$$t = \left( \frac{R^3}{\mu} \right)^{\frac{1}{2}} (\psi - e \sin \psi), \quad (2.1.27)$$

where the lower limits of integration have been taken to be zero. When (2.1.27) is evaluated at  $2\pi$ , Kepler's Third Law is recovered. Using this we may write (2.1.27) as

$$\omega t = \frac{2\pi}{T} t = \psi - e \sin \psi, \quad (2.1.28)$$

which is known as Kepler's equation. The quantity  $\omega t$  varies between 0 and  $2\pi$  as do  $\psi$  and  $\theta$  and is known as the mean anomaly. This equation is useful in the process of calculating  $r$  and  $\theta$  at a given time. It is solved to give the value of  $\psi$  at that time. From (2.1.25) one obtains  $r$ . By a manipulation of the orbit equation and (2.1.25) one obtains

$$\tan \frac{1}{2}\theta = \left( \frac{1+e}{1-e} \right)^{\frac{1}{2}} \tan \frac{1}{2}\psi. \quad (2.1.29)$$

Before the days of computers the solution of Kepler's equation attracted much attention for accurate astronomical calculations.

Many treatments dealing with the Laplace–Runge–Lenz vector have been criticised as being too *ad hoc* [67] in their derivations. Apart from the elegant simplicity of the above discussion, the same could be said of the method used here. However, in the sequel we see that the same techniques *mutatis mutandis* may be applied to a wide variety of problems and so, by this generality of application, are free from the taint of being too *ad hoc*. Certainly more sophisticated methods can be used to obtain the same results. Here we prefer to establish the classes of systems by elementary methods and then use the more advanced techniques to analyse the natures of these different classes of problems.

### 2.1.2 The model equation $\ddot{\mathbf{r}} + f\mathbf{r} = \mathbf{0}$

The central force equation

$$\ddot{\mathbf{r}} + f\mathbf{r} = \mathbf{0}, \quad (2.1.30)$$

where the variables in  $f$  are specified below, can formally be integrated to give a conserved vector of Hamilton's type, *videlicet*

$$\mathbf{K} = \dot{\mathbf{r}} + \int f\mathbf{r}dt. \quad (2.1.31)$$

The formality is removed if the integration in (2.1.31) can be rendered meaningful without the solution of the original equation (2.1.30) as an explicit function of time. The conservation of the angular momentum leads to a conserved vector of Laplace–Runge–Lenz type and immediately we can make sense of (2.1.31). More usefully for the moment we can make use of the planar nature of the motion to express everything in terms of plane polar coordinates. For the moment we measure the polar angle,  $\theta$ , from the cartesian direction,  $\hat{\mathbf{i}}$ . Then we can use the relations

$$\hat{\mathbf{r}} = \hat{\mathbf{i}} \cos \theta + \hat{\mathbf{j}} \sin \theta, \quad \hat{\boldsymbol{\theta}} = -\hat{\mathbf{i}} \sin \theta + \hat{\mathbf{j}} \cos \theta \quad (2.1.32)$$

between the variable unit vectors in the polar coordinates and the constant unit vectors in cartesian coordinates.

In central force problems the conservation of angular momentum, hence the confinement of the motion to a plane, enables one to use the relationship  $\dot{\theta} = L/r^2$  to pass from an integration with respect to time to an integration with respect to polar angle. This technique is well-established [140, p. 80], [40, Chap. 3] and [134, Chap. 3]. To reduce the formal integral in (2.1.31) to a quadrature we make the *Ansatz*

$$f\mathbf{r} = v(\theta)\dot{\boldsymbol{\theta}}, \quad (2.1.33)$$

where  $v(\theta)$  is an arbitrary function of its argument. (The only mathematical requirement on  $v$  is that it be Riemann integrable.) With the substitution (2.1.33) we have

$$\int f \mathbf{r} dt = \int v(\theta) \hat{\mathbf{r}} d\theta. \quad (2.1.34)$$

Using the cartesian decomposition of the unit vectors in plane polar coordinates in terms of the cartesian unit vectors and their inverses, *videlicet*

$$\begin{aligned} \hat{\mathbf{r}} &= \hat{\mathbf{i}} \cos \theta + \hat{\mathbf{j}} \sin \theta, & \hat{\boldsymbol{\theta}} &= -\hat{\mathbf{i}} \sin \theta + \hat{\mathbf{j}} \cos \theta, \\ \hat{\mathbf{i}} &= \hat{\mathbf{r}} \cos \theta - \hat{\boldsymbol{\theta}} \sin \theta, & \hat{\mathbf{j}} &= \hat{\mathbf{r}} \sin \theta + \hat{\boldsymbol{\theta}} \cos \theta \end{aligned} \quad (2.1.35)$$

we may write (2.1.34) as

$$\begin{aligned} \int f \mathbf{r} dt &= \hat{\mathbf{i}} \int \cos \theta v(\theta) d\theta + \hat{\mathbf{j}} \int \sin \theta v(\theta) d\theta \\ &= \hat{\mathbf{r}} \left[ \cos \theta \int_{\theta_0}^{\theta} \cos \eta v(\eta) d\eta + \sin \theta \int_{\theta_0}^{\theta} \sin \eta v(\eta) d\eta \right] \\ &\quad + \hat{\boldsymbol{\theta}} \left[ -\sin \theta \int_{\theta_0}^{\theta} \cos \eta v(\eta) d\eta + \cos \theta \int_{\theta_0}^{\theta} \sin \eta v(\eta) d\eta \right] \\ &= \hat{\mathbf{r}} z'(\theta) - \hat{\boldsymbol{\theta}} z(\theta), \end{aligned} \quad (2.1.36)$$

where

$$\begin{aligned} z(\theta) &= \int_{\theta_0}^{\theta} v(\eta) \sin(\theta - \eta) d\eta \\ z'(\theta) &= \int_{\theta_0}^{\theta} v(\eta) \cos(\theta - \eta) d\eta, \end{aligned} \quad (2.1.37)$$

and consequently we may write  $z(\theta)$  in terms of the differential equation

$$z''(\theta) + z(\theta) = v(\theta) \quad (2.1.38)$$

subject to the initial conditions  $z(\theta_0) = 0$  and  $z'(\theta_0) = 0$ . Hence we have the two conserved vectors

$$\mathbf{K} = \dot{\mathbf{r}} + z'(\theta) \hat{\mathbf{r}} - z(\theta) \hat{\boldsymbol{\theta}} \quad (2.1.39)$$

$$\mathbf{J} = \dot{\mathbf{r}} \times \hat{\mathbf{L}} - z(\theta) \hat{\mathbf{r}} - z'(\theta) \hat{\boldsymbol{\theta}}, \quad (2.1.40)$$

generalisations of Hamilton's vector and the Laplace–Runge–Lenz vector respectively.

Now that we have the Ermanno–Bernoulli constants in terms of the vector first integral,  $\mathbf{J}$ , the equation of the orbit in the plane is obtained by taking the scalar product of  $\mathbf{r}$  with  $\mathbf{J}$ . Unlike the case of the Kepler problem treated in the previous subsection here we have not assumed that  $\theta$  is measured from  $\mathbf{J}$  but from a standard cartesian direction denoted by  $\hat{\mathbf{i}}$ . We take the angle between  $\mathbf{J}$  and  $\hat{\mathbf{i}}$  to be  $\theta_0$  so that the equation of the orbit in the plane is given by

$$r(\theta) = \frac{L}{z(\theta) + J \cos(\theta - \theta_0)}. \quad (2.1.41)$$

The evolution of the motion in time can be obtained by using the conservation of angular momentum to give  $t$  as a function of  $\theta$  or by using the equation of the orbit, (2.1.41), to give  $t$  as a function of  $r$ . In either case the ease, or more likely lack of ease, in evaluating these functions will depend upon the particular function  $v(\theta)$ , which one assumes is dictated by physical considerations. Supposing that the quadratures can be performed we then have the problem of inversion to find the dependent variables  $r$  and/or  $\theta$  as explicit functions of time. Differentiating (2.1.41) with respect to time we obtain

$$\dot{r} = -z' + J \sin(\theta - \theta_0). \quad (2.1.42)$$

We can evaluate  $z'(\theta)$  from (2.1.37) and, provided all terms involving  $\theta$  can be expressed in terms of  $r$ , the relationship between  $t$  and  $r$  is given by

$$t = \int \frac{dr}{(-z'(\theta(r)) + J \sin(\theta(r) - \theta_0))}. \quad (2.1.43)$$

The angular momentum route, the more familiar one, leads to

$$t = \int \frac{Ld\theta}{(z(\theta) + J \cos(\theta - \theta_0))^2}. \quad (2.1.44)$$

Even though it may not be possible to evaluate these integrals in closed form or, if this be possible, to invert the results to give the dependent variables in terms of the independent variable  $t$ , one does know that the results of numerical integration can be trusted since the reduction of the problem to explicit quadratures removes the possibility of chaos.

Since the angular momentum is conserved, the areal velocity is a constant. In fact the area swept out by the radial vector  $\mathbf{r}$  in a time  $dt$  is given by the element of area

$$dA = \frac{1}{2}r(rd\theta). \quad (2.1.45)$$

Consequently the areal velocity is given by

$$\frac{dA}{dt} = \frac{1}{2}L \quad (2.1.46)$$

and, since  $L$  is a constant for the class of problems being considered here, (2.1.46) is integrated to give

$$A = \frac{1}{2}Lt. \quad (2.1.47)$$

In the case that the orbit for a given choice of  $v(\theta)$  has nicely defined geometric properties there may be some interest in relating (2.1.47) to these geometric properties, for example the semimajor axis length in the case of an elliptical orbit and the period which would arise from the area,  $A$ , in (2.1.47) assuming a multiple of the area enclosed by the orbit. In general one would not expect to be able to make a satisfactory relationship between the geometrical features of the orbit and the time.

We consider some examples.

**Example 2.1.** The equation of motion

$$\ddot{\mathbf{r}} + \frac{a(\theta - \theta_0) + b}{r^3} \mathbf{r} = 0, \quad (2.1.48)$$

where  $a$ ,  $b$  and  $\theta_0$  are constants, represents a force of the Kepler type with an additional inverse square law term which depends upon the polar angle  $\theta$ . The quadrature in (2.1.37) may be performed to give

$$z(\theta) = \frac{a}{L} (\theta - \theta_0 - \sin(\theta - \theta_0)) + \frac{b}{L} (1 - \cos(\theta - \theta_0)), \quad (2.1.49)$$

$$z'(\theta) = \frac{a}{L} (1 - \cos(\theta - \theta_0)) + \frac{b}{L} \sin(\theta - \theta_0). \quad (2.1.50)$$

The expressions for the conserved vectors and the orbit equation followed immediately. They are rather complicated and are not reproduced here. For increasing  $\theta$  the attraction to the centre increases and the orbit spirals inwards in a counterclockwise direction towards the origin.

**Example 2.2.** By way of contrast the equation of motion

$$\ddot{\mathbf{r}} + \frac{ae^{-(\theta-\theta_0)} + b}{r^3} \mathbf{r} = 0, \quad (2.1.51)$$

for which, again,  $a$ ,  $b$  and  $\theta_0$  are constants, contains a force of Kepler type with an additional inverse square law term which decays exponentially with increasing polar angle  $\theta$ . In this case we have

$$z(\theta) = \frac{a}{2L} \left( e^{-(\theta-\theta_0)} + \sin(\theta - \theta_0) - \cos(\theta - \theta_0) \right) + \frac{b}{L} (1 - \cos(\theta - \theta_0)), \quad (2.1.52)$$

$$z'(\theta) = \frac{a}{2L} \left( -e^{-(\theta-\theta_0)} + \sin(\theta - \theta_0) + \cos(\theta - \theta_0) \right) + \frac{b}{L} \sin(\theta - \theta_0). \quad (2.1.53)$$

The orbit contracts quickly to what appears to be a limit cycle as  $\theta$  increases.

**Example 2.3.** For the equation of motion

$$\ddot{\mathbf{r}} + \frac{a \sin 3(\theta - \theta_0) + b}{r^3} \mathbf{r} = 0, \quad (2.1.54)$$

which couples a force of Kepler type with an additional inverse square law term depending periodically on the polar angle  $\theta$ , we obtain

$$z(\theta) = \frac{a}{L} \left( -\frac{1}{8} \sin 3(\theta - \theta_0) + \frac{3}{8} \sin(\theta - \theta_0) \right) + \frac{b}{L} (1 - \cos(\theta - \theta_0)),$$

$$z'(\theta) = \frac{a}{L} \left( -\frac{3}{8} \cos 3(\theta - \theta_0) + \frac{3}{8} \cos(\theta - \theta_0) \right) + \frac{b}{L} \sin(\theta - \theta_0). \quad (2.1.55)$$

The periodicity of the central force is reflected in the orbit which has three lobes spaced at intervals of  $2\pi/3$  in  $\theta$ .

**Example 2.4.** In the present context the equation for the Kepler problem, (2.1.1), represents the choice

$$v(\theta) = \frac{\mu}{L} \tag{2.1.56}$$

for which

$$z(\theta) = \frac{\mu}{L} (1 - \cos(\theta - \theta_0)), \tag{2.1.57}$$

$$z'(\theta) = \frac{\mu}{L} \sin(\theta - \theta_0). \tag{2.1.58}$$

The conserved vectors are

$$\mathbf{K} = \dot{\mathbf{r}} - \frac{\mu}{L} (\hat{\boldsymbol{\theta}} - \hat{\boldsymbol{\theta}}_0), \tag{2.1.59}$$

$$\mathbf{J} = \dot{\mathbf{r}} \times \hat{\mathbf{L}} - \frac{\mu}{L} (\hat{\mathbf{r}} - \hat{\mathbf{r}}_0). \tag{2.1.60}$$

The expressions given here differ from the standard expressions by a factor of  $L$  in  $\mathbf{J}$  and the lack of identification of the reference line for  $\theta$  with  $\mathbf{J}$ . When he obtained his so-called ‘eccentricity vector’, Hamilton [55] immediately recognised that the velocity hodograph for the Kepler problem was a circle with its centre given by Hamilton’s vector (putting  $\hat{\boldsymbol{\theta}}_0 = 0$ ) and radius of length  $\mu/L$ . The velocity hodograph is generally attributed to Hamilton, but his biographer gives the credit to Möbius [56, p. 333]<sup>2,2</sup>.

**Example 2.5.** In the examples above  $\dot{\theta}$  does not appear in the equation of motion. In evaluating the function  $z(\theta)$  one must use the relationship  $L = r^2 \dot{\theta}$  to introduce the required  $\dot{\theta}$  for the quadrature. Here we present an elementary example in which  $f$  is explicitly dependent upon  $\dot{\theta}$  in the equation of motion. The equation of motion which we use is

$$\ddot{\mathbf{r}} + \frac{a\dot{\theta}}{r} \mathbf{r} = 0. \tag{2.1.61}$$

We find that

$$z(\theta) = a(1 - \cos(\theta - \theta_0)) \tag{2.1.62}$$

and that the equation of the orbit is

$$r = \frac{L}{a(1 - \cos(\theta - \theta_0)) + J \cos(\theta - \phi_0)}, \tag{2.1.63}$$

---

<sup>2,2</sup>In the nineteenth century the velocity hodograph attracted a considerable amount of attention for which see the works of Tait [135], Thompson (later Lord Kelvin) and Tait [71, 72], Kelland and Tait [70], Maxwell [104] and Routh [126]. The notation for the conserved vectors makes the equation of the circle obvious. The first known representation of the Laplace–Runge–Lenz vector in modern vector notation is found in the text of Gibbs and Wilson [37]. The transparency of the present representation makes for a dramatic contrast for the original representation to be found in the papers of Ermanno and Bernoulli [33, 59, 12]. Early in the twentieth century Child [20] made a complete graphical study of the velocity hodograph for the three possible orbits of the Kepler problem. There appear to be some inconsistencies in the depictions which are possibly due to roundoff errors in the numerical calculations which may well have been done on the back of an envelope. More recently the velocity hodograph has been discussed by Goldstein [39, 40] and Stickforth [133]. Possibly the most detailed discussion of the velocity hodograph as applied to the Kepler and related problems is to be found in the thesis of Gorringer [41]. Gorringer discusses also the acceleration hodograph.

where  $\phi_0$  is the angle the reference line makes with  $\mathbf{J}$ . The orbit is an ellipse, a parabola or an hyperbola depending upon the relative values of the constants.

## 2.2 Algebraic properties of the first integrals of the Kepler problem

The Hamiltonian of the Kepler problem described by equation (2.1.1) is

$$H = \frac{1}{2}\mathbf{p} \cdot \mathbf{p} - \frac{\mu}{r}, \quad (2.2.1)$$

where the canonical momentum is related to the velocity by  $\mathbf{p} = \dot{\mathbf{r}}$ , and is simply the mechanical energy which is conserved. In terms of canonical coordinates the conserved vectors are

$$\mathbf{L} = \mathbf{r} \times \mathbf{p}, \quad (2.2.2)$$

$$\mathbf{K} = \mathbf{p} - \frac{\mu}{L}\hat{\boldsymbol{\theta}}, \quad (2.2.3)$$

$$\mathbf{J} = \mathbf{p} \times \mathbf{L} - \mu\hat{\mathbf{r}} \quad (2.2.4)$$

and the components of all three vectors have zero Poisson Bracket with the Hamiltonian. Consequently, in quantum mechanical terms, the physical quantities represented by the corresponding quantum mechanical operators are simultaneously observable with the Hamiltonian. Since the three vectors are not independent, a knowledge of two means a knowledge of three. Normally the two vectors considered are the angular momentum and the Laplace–Runge–Lenz<sup>2,3</sup>. The Laplace–Runge–Lenz vector was used to explain the degeneracy of the spectrum of the hydrogen atom and is described by physicists as a dynamical symmetry in contrast to the obvious geometric symmetry of the angular momentum. The angular momentum vector may be a more frequently occurring conserved quantity than the Laplace–Runge–Lenz vector, but there is no reason to distinguish between the two. Both vectors are a consequence of the equation of motion.

If one writes each vector in terms of cartesian components, *ie* as  $\mathbf{A} = A_i\hat{\mathbf{e}}_i$ , the Poisson Bracket relationships are

$$\begin{aligned} [L_i, H] &= 0, & [J_i, H] &= 0, \\ [L_i, L_j] &= \varepsilon_{ijk}L_k, & [J_i, L_j] &= \varepsilon_{ijk}J_k, \\ [J_i, J_j] &= -2E\varepsilon_{ijk}L_k, \end{aligned} \quad (2.2.5)$$

where  $\varepsilon_{ijk}$  is Kronecker's epsilon and  $E$  is the value of the Hamiltonian. The three components of the angular momentum vector form the compact rotation algebra  $so(3)$ . The algebra of the two vectors is  $so(4)$ , the representation of the four-dimensional compact rotation group, for negative energies,  $so(3, 1)$ , the representation of the four-dimensional

---

<sup>2,3</sup>There appears to be no reason why Hamilton's vector should not be used in place of the Laplace–Runge–Lenz vector. The use of the latter is possibly an accident of history. We recall that in the early twentieth century the Ermanno–Bernoulli constants had been 'popularised' by the German writers Runge and Lenz. A large contribution to the early development of Quantum Mechanics had a Teutonic origin and it is possible that this recent work was more familiar than the older work of the quaternion-fancying Irishman, Hamilton, amongst German speakers. The Hamiltonian formulation of Quantum Mechanics by Dirac [28] was probably too late to change the situation.



noncompact rotation group, *ie* the Lorentz group, for positive energies and  $e(3)$ , the representation of the Euclidean group for zero energy. A better representation of the algebra for nonzero energy is given by defining

$$\mathcal{J}_{i\pm} = \frac{J_i}{|2E|^{1/2}} \quad (2.2.6)$$

for negative and positive energies respectively. Then (2.2.5) is replaced by

$$\begin{aligned} [L_i, H] &= 0, & [\mathcal{J}_i, H] &= 0, \\ [L_i, L_j] &= \varepsilon_{ijk} L_k, & [\mathcal{J}_i, L_j] &= \varepsilon_{ijk} \mathcal{J}_k, \\ & & [\mathcal{J}_i, \mathcal{J}_j] &= \pm \varepsilon_{ijk} L_k, \end{aligned} \quad (2.2.7)$$

for negative and positive energies respectively, which has an æsthetic appeal greater than (2.2.5).

### 2.3 Extension of the model equation to nonautonomous systems

In § 2.2 the model equation (2.1.11) was tacitly assumed to be autonomous. However, Katzin and Levine [69], using a direct method, and Leach [81], using a variant of Noether's theorem [114], showed that the time-dependent central force problem with the equation of motion

$$\ddot{\mathbf{r}} = F(r, t)\hat{\mathbf{r}} \quad (2.3.1)$$

possesses a time-dependent vector first integral of the form

$$\mathbf{J} = U(r, t)(\mathbf{L} \times \dot{\mathbf{r}}) + Z(r, t)(\mathbf{L} \times \mathbf{r}) + W(r, t)\mathbf{r}, \quad (2.3.2)$$

provided

$$\begin{aligned} U &= u(t) \neq 0, & Z &= -\dot{u}(t), \\ W &= \frac{\mu}{r}, & F &= \left(\frac{\ddot{u}r}{u}\right) - \left(\frac{\mu}{ur^2}\right). \end{aligned} \quad (2.3.3)$$

This system includes both a time-dependent Kepler problem ( $\mu \neq 0$  and  $u = (\alpha t + \beta)/\lambda$  with  $\alpha$ ,  $\beta$  and  $\lambda$  being constant – this was the subject of an earlier study by Katzin and Levine [68]) – and a particular case of the time-dependent harmonic oscillator when  $\mu = 0$ <sup>2.4</sup>. Previous techniques used to solve this problem have been algebraically rather complex. The techniques used here have been reported in Gorringer and Leach [42, 44] and Leach and Gorringer [87].

We rewrite our model equation to include the possible dependence on time explicitly as

$$\ddot{\mathbf{r}} + f(r, \theta, t)\mathbf{r} = 0 \quad (2.3.4)$$

---

<sup>2.4</sup>The time-dependent harmonic oscillator was solved in one dimension by Lewis [94, 95] using Kruskal's asymptotic method [76]. Lewis and Riesenfeld [97] used the time-dependent harmonic oscillator to construct an invariant for a charged particle moving in a time-dependent electromagnetic field. Further techniques and results have been given by Lewis and Leach [96], Moreira [110, 111] and Günther and Leach [79].

of which the system above is a special case. The conservation of the vector of angular momentum,  $\mathbf{L} := \mathbf{r} \times \dot{\mathbf{r}}$ , follows from the vector product of  $\mathbf{r}$  with (2.3.4). We seek a Hamilton-like vector for (2.3.4) through the use of an integrating factor,  $g$ . Writing

$$g\ddot{\mathbf{r}} = (g\dot{\mathbf{r}} - \dot{g}\mathbf{r})\dot{\phantom{r}} + \ddot{g}\mathbf{r} \quad (2.3.5)$$

and multiplying (2.3.4) by  $g$  we obtain

$$(g\dot{\mathbf{r}} - \dot{g}\mathbf{r})\dot{\phantom{r}} + (\ddot{g} + gf)\mathbf{r} = 0. \quad (2.3.6)$$

A second conserved vector exists if the second term of (2.3.6) can be written as a total time derivative. The presence of  $\hat{\mathbf{r}}$  which is a function of  $\theta$  signals the requirement that

$$(\ddot{g} + gf)r = v(\theta)\dot{\theta}, \quad (2.3.7)$$

for some function  $v(\theta)$ . The explicit dependence upon  $\dot{\theta}$  may be removed by using  $L = r^2\dot{\theta}$  and we then solve (2.3.7) to obtain

$$f = \frac{Lv(\theta)}{gr^3} - \frac{\ddot{g}}{g}. \quad (2.3.8)$$

Since  $f$  was assumed a function of  $r$ ,  $\theta$  and  $t$ ,  $g = g(t)$ . The equation (2.3.4) is now

$$\ddot{\mathbf{r}} + \frac{Lv(\theta)}{g(t)r^3}\mathbf{r} - \frac{\ddot{g}(t)}{g(t)}\mathbf{r} = 0, \quad (2.3.9)$$

where  $L$  is treated as a constant in the equation of motion, although it is expressed as  $r^2\dot{\theta}$  during calculations.

Direct integration of the equation of motion (2.3.9) yields the Hamilton-like vector

$$\mathbf{K} = g\dot{\mathbf{r}} - \dot{g}\mathbf{r} + z'(\theta)\hat{\mathbf{r}} - z(\theta)\hat{\boldsymbol{\theta}}, \quad (2.3.10)$$

where  $z(\theta)$  and  $z'(\theta)$  are given by (2.1.36) and (2.1.37). The analogue of the Laplace–Runge–Lenz vector is obtained by taking the vector product of  $\mathbf{K}$  with  $\hat{\mathbf{L}}$  and is

$$\mathbf{J} = (g\dot{\mathbf{r}} - \dot{g}\mathbf{r}) \times \hat{\mathbf{L}} - z(\theta)\hat{\mathbf{r}} - z'(\theta)\hat{\boldsymbol{\theta}}. \quad (2.3.11)$$

The presence of  $\mathbf{r}$  in the Hamilton-like vector  $\mathbf{K}$  precludes the possibility to express the velocity hodograph in a simple form as was the case for the Kepler problem. However, this can be partially rectified by a time-dependent linear transformation of Kummer–Liouville type [77, 100] (in Hamiltonian Mechanics it is a time-dependent linear generalised canonical transformation [79, 120]).

The scalar product of  $\mathbf{J}$  with  $\mathbf{r}$  gives

$$r(\theta) = \frac{Lg(t)}{z(\theta) + J \cos(\theta - \theta_0)}. \quad (2.3.12)$$

Since (2.3.12) contains an explicit function of time, it is not the required orbit equation. We eliminate the function of time to obtain the orbit equation. From the expression for the constant magnitude of the angular momentum we have

$$L = r^2 \frac{d\theta}{dt} = \frac{L^2 g^2(t)}{(z(\theta) + J \cos(\theta - \theta_0))^2} \frac{d\theta}{dt} \quad (2.3.13)$$

which can be rearranged to give

$$\int_0^t \frac{dt}{g^2(t)} = \int_{\theta_0}^{\theta} \frac{Ld\theta}{(z(\theta) + J \cos(\theta - \theta_0))^2}. \quad (2.3.14)$$

Provided the integrands are continuous and nonzero we can, according to the Implicit Function Theorem [16, p. 165], write the result of the integration as

$$t = N(\theta) \quad (2.3.15)$$

to obtain the orbit equation

$$r(\theta) = \frac{Lg \circ N(\theta)}{z(\theta) + J \cos(\theta - \theta_0)}, \quad (2.3.16)$$

where  $g \circ N$  denotes the composition of the functions  $g$  and  $N$ .

We consider some examples.

**Example 2.6.**  $v(\theta) = 0$ . The equation of motion, (2.3.9), is

$$\ddot{\mathbf{r}} = \frac{\ddot{g}}{g} \mathbf{r} \quad (2.3.17)$$

with the conserved vectors

$$\begin{aligned} \mathbf{K} &= g\dot{\mathbf{r}} - \dot{g}\mathbf{r}, \\ \mathbf{J} &= (g\dot{\mathbf{r}} - \dot{g}\mathbf{r}) \times \hat{\mathbf{L}} \end{aligned} \quad (2.3.18)$$

and an energylike invariant

$$\begin{aligned} I &= \frac{1}{2} \mathbf{K} \cdot \mathbf{K} \\ &= \frac{1}{2} (g\dot{\mathbf{r}} - \dot{g}\mathbf{r})^2. \end{aligned} \quad (2.3.19)$$

The Hamiltonian for (2.3.17) is

$$H = \frac{1}{2} \left( \mathbf{p} \cdot \mathbf{p} - \frac{\ddot{g}}{g} r^2 \right). \quad (2.3.20)$$

If the integral on the left side of (2.3.14) is denoted by  $M(t)$ , the orbit equation is

$$r(\theta) = \frac{Lg \circ M^{-1} \circ (L \tan(\theta - \theta_0)/J^2)}{J \cos(\theta - \theta_0)}. \quad (2.3.21)$$

**Example 2.7.**  $v(\theta) = \mu/L$ . The equation of motion is

$$\ddot{\mathbf{r}} = \frac{\ddot{g}}{g} \mathbf{r} - \frac{\mu}{gr^3} \mathbf{r} \quad (2.3.22)$$

and the conserved vectors are

$$\begin{aligned} \mathbf{K} &= g\dot{\mathbf{r}} - \dot{g}\mathbf{r} - \frac{\mu}{L}(\hat{\boldsymbol{\theta}} - \hat{\boldsymbol{\theta}}_0), \\ \mathbf{J} &= (g\dot{\mathbf{r}} - \dot{g}\mathbf{r}) \times \hat{\mathbf{L}} - \frac{\mu}{L}(\hat{\mathbf{r}} - \hat{\mathbf{r}}_0). \end{aligned} \quad (2.3.23)$$



**Example 2.8.** Another system for which an energylike integral can be found is described by the equation of motion

$$\ddot{\mathbf{r}} = \frac{\ddot{g}}{g}\mathbf{r} - \frac{\mu}{g^3r}\mathbf{r}. \quad (2.3.30)$$

An integrating factor for (2.3.30) is  $g(g\dot{\mathbf{r}} - \dot{g}\mathbf{r})$  as a scalar product. The invariant obtained is

$$I = \frac{1}{2}(g\dot{\mathbf{r}} - \dot{g}\mathbf{r})^2 + \mu r g \quad (2.3.31)$$

and the Hamiltonian is

$$H = \frac{1}{2}\mathbf{p} \cdot \mathbf{p} - \frac{1}{2}\frac{\ddot{g}}{g}r^2 + \frac{\mu r}{g^3}. \quad (2.3.32)$$

There does not seem to be a simple way to construct either a Hamilton's or Laplace–Runge–Lenz vector for (2.3.30) by the methods which have been considered here. If one follows the principles of Fradkin and others, this should be possible. However, the conserved vectors would be piecewise continuous at best and would not possess group properties but only algebraic properties. Although it is convenient to work with algebras, one must agree with Bacry [9, 8] that the important object, as far as Physics is concerned, is the group.

### 3 Conservation of the direction of angular momentum only

#### 3.1 Vector conservation laws for the equation of motion $\ddot{\mathbf{r}} + g\hat{\mathbf{r}} + h\hat{\boldsymbol{\theta}} = 0$

In the previous chapter we considered equations of motion for which the vector of angular momentum is conserved in both magnitude and direction. We saw that it was possible to obtain generalisations of Hamilton's vector and the Laplace–Runge–Lenz vector. In this chapter we examine the extent to which we may still make reasonable progress with a relaxation of one of the requirements previously imposed upon the angular momentum. The relaxation is that we require only the constancy of the direction of angular momentum and not of its magnitude. The motion is still in a plane and, by selecting the origin to be in that plane, we may still use plane polar coordinates,  $(r, \theta)$ , to describe the orbit. Consequently, for the content of this chapter at least, we do not have to move out of two dimensions. A particular advantage is that we preserve the relationship

$$\hat{\mathbf{L}} = \hat{\mathbf{r}} \times \hat{\boldsymbol{\theta}} \quad (3.1.1)$$

between the direction of the angular momentum and the two unit vectors of the coordinate system. Since  $\dot{\mathbf{L}} = \dot{L}\hat{\mathbf{L}}$  and so  $\dot{\mathbf{L}} \times \mathbf{L} = 0$ ,  $\dot{\mathbf{L}}$  and  $\mathbf{L}$  are collinear and we may write

$$\dot{\mathbf{L}} + h_1\mathbf{L} = 0, \quad (3.1.2)$$

where  $h_1$  is some arbitrary function<sup>3.1</sup>. Making use of (3.1.1) we may rewrite (3.1.2) as

$$\mathbf{r} \times \ddot{\mathbf{r}} + g_1\mathbf{r} \times \mathbf{r} + \frac{h_1L}{r}\mathbf{r} \times \hat{\boldsymbol{\theta}} = 0, \quad (3.1.3)$$

---

<sup>3.1</sup>That (3.1.2) in turn implies the constancy of  $\hat{\mathbf{L}}$  follows from writing  $\dot{\mathbf{L}} = \dot{L}\hat{\mathbf{L}} + L\dot{\hat{\mathbf{L}}}$  in (3.1.2) and taking the scalar product with  $\hat{\mathbf{L}}$  and the vector product with  $\mathbf{L}$  in turn.

where  $g_1$  is also an arbitrary function. Consequently the equation of motion

$$\ddot{\mathbf{r}} + g\hat{\mathbf{r}} + h\hat{\boldsymbol{\theta}} = 0, \quad (3.1.4)$$

where  $g = g_1 r$  and  $h = h_1 L/r$ , describes motion in a plane with  $\hat{\mathbf{L}}$  conserved. Equation (3.1.4) is not the most general equation of motion in the plane, but belongs to a class of problems for which the techniques to be discussed below can be applied. For example we could introduce additional unrestricted functions by adding terms such as  $e\mathbf{r} \times \dot{\mathbf{r}}$  and in a later section of this chapter (3.1.2) is rewritten in terms of a different set of vector products and shown to describe a model for the motion of low altitude artificial satellites in the atmosphere.

We employ a direct method, attributed to J Bertrand<sup>3,2</sup>, to find a vector of the class of Laplace–Runge–Lenz for the equation of motion (3.1.4). The general principle of a direct method is to impose some *Ansatz* on the structure of the first integral or the equation of motion so that the partial differential equation resulting from the imposition that the total time derivative of the first integral be zero can be separated into subequations. In this instance we assume that the functions  $g$  and  $h$  depend only on  $r$  and  $\theta$ . From the vector product of  $\mathbf{r}$  with (3.1.4) we obtain

$$\dot{\mathbf{L}} = -hr. \quad (3.1.5)$$

The vector product of (3.1.4) with  $\mathbf{L}$  gives

$$\dot{\mathbf{r}} \times \mathbf{L} - gL\hat{\boldsymbol{\theta}} + hL\hat{\mathbf{r}} = 0 \quad (3.1.6)$$

which can be rewritten as

$$\frac{d}{dt}(\dot{\mathbf{r}} \times \mathbf{L}) - \dot{\mathbf{r}} \times \dot{\mathbf{L}} - gL\hat{\boldsymbol{\theta}} + hL\hat{\mathbf{r}} = 0. \quad (3.1.7)$$

A Laplace–Runge–Lenz-type vector exists of the form

$$\mathbf{J} = \dot{\mathbf{r}} \times \mathbf{L} - u(r, \theta)\hat{\mathbf{r}} - v(r, \theta)\hat{\boldsymbol{\theta}} \quad (3.1.8)$$

provided

$$\frac{d}{dt}(-u\hat{\mathbf{r}} - v\hat{\boldsymbol{\theta}}) = hr\dot{\mathbf{r}} \times \hat{\mathbf{L}} - gL\hat{\boldsymbol{\theta}} + hL\hat{\mathbf{r}}. \quad (3.1.9)$$

Equating the coefficients of  $\hat{\mathbf{r}}$  and  $\hat{\boldsymbol{\theta}}$  separately to zero and using  $L = r^2\dot{\theta}$  we have

$$\begin{aligned} \hat{\mathbf{r}} : \quad & \dot{u} - v\dot{\theta} = -2r^2h\dot{\theta}, \\ \hat{\boldsymbol{\theta}} : \quad & u\dot{\theta} + \dot{v} = r\dot{r}h + gr^2\dot{\theta}. \end{aligned} \quad (3.1.10)$$

We now use the assumed functional dependence (3.1.8) to obtain the set of four partial differential equations as a sufficient condition

$$\begin{aligned} u_r &= 0, & v_r &= hr, \\ u_\theta - v &= -2hr^2, & v_\theta + u &= gr^2, \end{aligned} \quad (3.1.11)$$

---

<sup>3,2</sup>See Whittaker [140, p. 332, § 152].

where the subscripts  $r$  and  $\theta$  represent partial differentiation with respect to  $r$  and  $\theta$  respectively. We solve (3.1.11) easily to obtain

$$\begin{aligned} u &= U(\theta), & v &= U'(\theta) + 2r^{\frac{1}{2}}V(\theta), \\ h &= r^{-\frac{3}{2}}V(\theta), & g &= r^{-2} \left[ U''(\theta) + U(\theta) + 2r^{\frac{1}{2}}V'(\theta) \right], \end{aligned} \quad (3.1.12)$$

where  $U(\theta)$  and  $V(\theta)$  are arbitrary functions of their argument. The system with equation of motion

$$\dot{\mathbf{r}} + \left[ \frac{U''(\theta) + U(\theta)}{r^2} + \frac{2V'(\theta)}{r^{\frac{3}{2}}} \right] \hat{\mathbf{r}} + \frac{V(\theta)}{r^{\frac{3}{2}}} \hat{\boldsymbol{\theta}} = 0 \quad (3.1.13)$$

has the conserved vector

$$\mathbf{J} = \dot{\mathbf{r}} \times \mathbf{L} - U(\theta)\hat{\mathbf{r}} - \left[ U'(\theta) + 2r^{\frac{1}{2}}V(\theta) \right] \hat{\boldsymbol{\theta}}. \quad (3.1.14)$$

We can construct an analogue of Hamilton's vector from this vector by taking the vector product of  $\hat{\mathbf{L}}$  with  $\mathbf{J}$ . It is

$$\begin{aligned} \mathbf{K} &= \hat{\mathbf{L}} \times \mathbf{J} \\ &= L\dot{\mathbf{r}} - U(\theta)\hat{\boldsymbol{\theta}} + \left[ U'(\theta) + 2r^{\frac{1}{2}}V(\theta) \right] \hat{\mathbf{r}}. \end{aligned} \quad (3.1.15)$$

Since  $L$  is no longer conserved, we can no longer rescale either (3.1.14) or (3.1.15) by  $L$ .

### 3.2 The orbit equation and motion in time

We construct the orbit equation from the conserved vector,  $\mathbf{J}$ . If we let  $\theta_0$  be the fixed angle between  $\mathbf{J}$  and the cartesian unit vector  $\hat{\mathbf{i}}$  and take the scalar product of  $\mathbf{J}$  with  $\mathbf{r}$ , we find after the usual rearrangement that

$$r = \frac{L^2}{U(\theta) + J \cos(\theta - \theta_0)}. \quad (3.2.1)$$

This is not the orbit equation since  $L$  is not a constant. To express  $L$  in terms of  $\theta$  we write

$$\dot{L} = \frac{dL}{d\theta} \dot{\theta} = \frac{dL}{d\theta} \frac{L}{r^2}, \quad (3.2.2)$$

use (3.1.5) and the third of (3.1.12) to obtain

$$L \frac{dL}{d\theta} = -r^{\frac{3}{2}}V(\theta). \quad (3.2.3)$$

We eliminate  $r$  from (3.2.1) and (3.2.3) to express  $L$  in terms of the quadrature

$$\frac{1}{L} = \frac{1}{L_0} + \int_{\theta_0}^{\theta} \frac{V(\eta)d\eta}{[U(\eta) + J \cos(\eta - \theta_0)]^{\frac{3}{2}}}. \quad (3.2.4)$$

The orbit equation is

$$r(\theta) = \frac{1}{U(\theta) + J \cos(\theta - \theta_0)} \times \left[ \frac{1}{L_0} + \int_{\theta_0}^{\theta} \frac{V(\eta) d\eta}{[U(\eta) + J \cos(\eta - \theta_0)]^{\frac{3}{2}}} \right]^{-2}. \quad (3.2.5)$$

The orbit equation for the standard Kepler problem is recovered when we set  $U(\theta) = U_0$  and  $V(\theta) = 0$ .

The explicit determination of  $r(t)$  is not possible unless one can express  $\theta$  in terms of  $r$ . If we differentiate (3.2.1), use (3.1.5) to replace  $\dot{L}$  and again (3.2.1), we obtain

$$\dot{r} = -\frac{2hr^2}{L} - \frac{1}{L} (U'(\theta) - J \sin(\theta - \theta_0)). \quad (3.2.6)$$

The evident impossibility of expressing the right-hand side of (3.2.6) in terms of  $r$  for a general function of  $\theta$  means that the radial motion can only be evaluated numerically.

The function  $\theta(t)$  can be calculated from (3.2.4) and the scalar equation for the angular momentum

$$\dot{\theta} = \frac{(U(\theta) + J \cos(\theta - \theta_0))^2}{L^3}. \quad (3.2.7)$$

Naturally it may not be possible to invert the results of this calculation even if the quadrature can be performed to give  $\theta$  in terms of  $t$ .

The calculation of the areal velocity also poses serious practical problems. If the angular motion in time can be solved using (3.2.7) so that

$$t = f_1(\theta) \quad (3.2.8)$$

and this can be inverted to give

$$\theta = f_2(t), \quad (3.2.9)$$

we can use

$$\frac{dA}{dt} = \frac{1}{2} L (f_2(t)), \quad (3.2.10)$$

where  $L$  is calculated from (3.2.4), and the areal velocity can in principle be determined. One would expect to take the numerical route normally.

We note that the existence of the conserved vectors to enable us to perform the formal calculation of the orbit in the same, or a very similar, manner as for the Kepler problem by no means guarantees that we can perform in closed form the necessary quadratures. However, we should bear in mind that motions in the plane can be chaotic. That we are able to reduce these problems to quadratures indicates that they do not belong to the chaotic class.

**Example 3.1.** The orbit equation (3.2.5) can yield a wide variety of orbits for different choices of the functions  $U(\theta)$  and  $V(\theta)$ , not to mention for values of the parameters contained in particular choices of the functions. If  $V \equiv 0$  (indicating the conservation of the angular momentum vector and so not strictly belonging to this Chapter, but still



of some interest for purposes of comparison) and  $U$  is an increasing function of  $\theta$  and  $U(0) > J > 0$ , the orbit spirals inwards. If still  $V \equiv 0$ , but now  $U$  is a decreasing function of  $\theta$  with  $U(0) > J > 0$ , the orbit spirals outwards. For  $V \equiv 0$  and  $U$  periodic with a period incommensurate with  $2\pi$  the orbit does not close. For suitable initial conditions the orbit closes with period  $2\pi$  if  $U$  has the period  $2\pi/n$ ,  $n \in \mathbb{N}$ . If  $U$  has a period  $2n\pi$ ,  $n \in \mathbb{N}$ , the orbit circles the origin  $n$  times before closing, again for suitable values of the initial conditions. A simple model to illustrate these possibilities is

$$U(\theta) = a \sin k(\theta - \theta_0) + b \quad (3.2.11)$$

for which the corresponding equation of motion, (3.1.13), is

$$\ddot{\mathbf{r}} + \left( \frac{a(1 - k^2) \sin k(\theta - \theta_0) + b}{r^2} \right) \hat{\mathbf{r}} = 0 \quad (3.2.12)$$

and for which the conserved vectors are

$$\mathbf{K} = L\dot{\mathbf{r}} - (a \sin k(\theta - \theta_0) + b) \hat{\boldsymbol{\theta}} + ak \cos k(\theta - \theta_0) \hat{\mathbf{r}}, \quad (3.2.13)$$

$$\mathbf{J} = \dot{\mathbf{r}} \times \mathbf{L} - (a \sin k(\theta - \theta_0) + b) \hat{\mathbf{r}} - ak \cos k(\theta - \theta_0) \hat{\boldsymbol{\theta}}. \quad (3.2.14)$$

**Example 3.2.** In the case that  $U$  is constant and  $V$  is periodic with period  $2n\pi$ ,  $n \in \mathbb{N}/1$ , the orbit circles the origin  $n$  times before closing in the case that the energy permits bound orbits. For

$$U = a, \quad V = b \sin k(\theta - \theta_0) \quad (3.2.15)$$

the equation of motion is

$$\ddot{\mathbf{r}} + \left( \frac{a}{r^2} + \frac{2bk \cos k(\theta - \theta_0)}{r^{\frac{3}{2}}} \right) \hat{\mathbf{r}} + \frac{b \sin k(\theta - \theta_0)}{r^{\frac{3}{2}}} \hat{\boldsymbol{\theta}} = 0 \quad (3.2.16)$$

and the conserved vectors are

$$\mathbf{K} = L\dot{\mathbf{r}} + \left( 2r^{\frac{1}{2}} b \sin k(\theta - \theta_0) \right) \hat{\mathbf{r}} - a \hat{\boldsymbol{\theta}}, \quad (3.2.17)$$

$$\mathbf{J} = \dot{\mathbf{r}} \times \mathbf{L} - \left( 2r^{\frac{1}{2}} b \sin k(\theta - \theta_0) \right) \hat{\boldsymbol{\theta}} - a \hat{\mathbf{r}}. \quad (3.2.18)$$

This example is true to the genre in that the angular momentum is constant only in direction and not in magnitude.

Some illustrative orbits are given in Gorrington and Leach [43].

### 3.3 The related Hamiltonian system

We now consider the most general autonomous Hamiltonian which possesses a Laplace–Runge–Lenz vector and belongs to the class of problems specified by the equation of motion, (3.1.13). We assume an Hamiltonian of the form

$$H = \frac{1}{2} \mathbf{p} \cdot \mathbf{p} + W(r, \theta) \quad (3.3.1)$$

for which the Newtonian equation of motion can be calculated from

$$\ddot{\mathbf{r}} = -\nabla W(r, \theta) = \mathbf{F}(r, \theta). \quad (3.3.2)$$

In plane polar coordinates

$$\nabla W(r, \theta) = \frac{\partial W}{\partial r} \hat{\mathbf{r}} + \frac{1}{r} \frac{\partial W}{\partial \theta} \hat{\boldsymbol{\theta}} \quad (3.3.3)$$

and by comparison with (3.1.12) we have

$$\frac{\partial W}{\partial r} = r^{-2} [U''(\theta) + U(\theta)] + 2r^{-\frac{3}{2}} V'(\theta), \quad (3.3.4)$$

$$\frac{\partial W}{\partial \theta} = r^{-\frac{1}{2}} V(\theta). \quad (3.3.5)$$

A consistent solution of (3.3.4) and (3.3.5) places considerable constraints on the functions  $U(\theta)$  and  $V(\theta)$  so that we obtain the Hamiltonian

$$H = \frac{1}{2} \left( p_r^2 + \frac{p_\theta^2}{r^2} \right) - \frac{\mu}{r} - \alpha r^{-\frac{1}{2}} \cos \left( \frac{1}{2}(\theta - \beta) \right) \quad (3.3.6)$$

with the Newtonian equation of motion and conserved vectors given by

$$\ddot{\mathbf{r}} + \left( \frac{\mu}{r^2} + \frac{\alpha \cos \frac{1}{2}(\theta - \beta)}{2r^{\frac{3}{2}}} \right) \hat{\mathbf{r}} + \frac{\alpha \sin \frac{1}{2}(\theta - \beta)}{2r^{\frac{3}{2}}} \hat{\boldsymbol{\theta}} = 0, \quad (3.3.7)$$

$$\mathbf{K} = L\dot{\mathbf{r}} + \alpha r^{\frac{1}{2}} \sin \frac{1}{2}(\theta - \beta) \hat{\mathbf{r}} - \mu \hat{\boldsymbol{\theta}}, \quad (3.3.8)$$

$$\mathbf{J} = \dot{\mathbf{r}} \times \mathbf{L} - \mu \hat{\mathbf{r}} - \alpha r^{\frac{1}{2}} \sin \frac{1}{2}(\theta - \beta) \hat{\boldsymbol{\theta}}. \quad (3.3.9)$$

We note that only the constant solution for  $U(\theta)$  persists. The  $\theta$ -dependent part of  $U$  plays no role in the Hamiltonian or the equation of motion. Its sole contribution is to add a constant vector to  $\mathbf{J}$  and so it may be ignored.

### 3.4 The Lie algebra of the Poisson Brackets of the first integrals

We note a considerable diminution in the generality of the results from the Newtonian case to the Hamiltonian case just simply because of the requirement that the force be the gradient of a potential. Nevertheless it is interesting that even that requirement does not impose the condition that the force be single-valued. The two nonzero cartesian components of (3.3.9) are

$$J_1 = p_r p_\theta \sin \theta + \frac{p_\theta^2}{r} \cos \theta - \mu \cos \theta + \alpha r^{\frac{1}{2}} \sin \frac{1}{2}(\theta - \beta) \sin \theta, \quad (3.4.1)$$

$$J_2 = -p_r p_\theta \cos \theta + \frac{p_\theta^2}{r} \sin \theta - \mu \sin \theta - \alpha r^{\frac{1}{2}} \sin \frac{1}{2}(\theta - \beta) \cos \theta, \quad (3.4.2)$$

where for convenience we have written the cartesian representation in terms of the usual plane polar coordinates. An alternative derivation of (3.4.1) and (3.4.2) would be to use the structure for  $H$  (3.3.1) and  $\mathbf{J}$  (3.1.8) and impose the requirement that the Poisson

Bracket of  $H$  with each of  $J_1$  and  $J_2$  be zero. The result is the same although the machinery used is quite different. A detailed treatment may be found in Gorringe and Leach [49]. An alternate representation of the two constants is to write them as

$$\begin{aligned} J_{\pm} &= J_1 \pm iJ_2 \\ &= e^{\pm i\theta} \left[ \frac{p_{\theta}^2}{r} - \mu \mp i \left( p_r p_{\theta} + \alpha r^{\frac{1}{2}} \sin \frac{1}{2}(\theta - \beta) \right) \right] \end{aligned} \quad (3.4.3)$$

which is the type of combination of some use, as we see below, in the study of the complete symmetry groups for these problems.

Whichever form of the Ermanno–Bernoulli constants we use, the two first integrals can be used to generate a third integral using Poisson’s theorem (*cf* [140, p. 320, § 145] and from the Poisson Bracket we obtain

$$I = 2p_{\theta}H + p_r\alpha r^{\frac{1}{2}} \sin \frac{1}{2}(\theta - \beta) + p_{\theta}\alpha r^{-\frac{1}{2}} \cos \frac{1}{2}(\theta - \beta). \quad (3.4.4)$$

Since they are first integrals, all have zero Poisson Bracket with the Hamiltonian and so constitute the elements of an invariance algebra of the Hamiltonian. The Poisson Bracket relations between the three functions are

$$\begin{aligned} [J_1, J_2]_{PB} &= -I, \\ [J_1, I]_{PB} &= -2HJ_2 - \frac{1}{2}\alpha^2 \sin \beta, \\ [J_2, I]_{PB} &= 2HJ_1 + \frac{1}{2}\alpha^2 \cos \beta. \end{aligned} \quad (3.4.5)$$

The Poisson Bracket relations, (3.4.5), are neither very tidy nor very attractive. We make some adjustment to the definitions of the first integrals by writing

$$\begin{aligned} A_{\pm} &= |2H|^{-\frac{1}{2}} \left( J_1 + \frac{\alpha^2 \cos \beta}{4H} \right), \\ B_{\pm} &= |2H|^{-\frac{1}{2}} \left( J_2 + \frac{\alpha^2 \sin \beta}{4H} \right), \\ C_{\pm} &= \frac{I}{2H} \end{aligned} \quad (3.4.6)$$

in which the positive sign of the  $\pm$  refers to positive energies and the negative sign to negative energies. Now the Poisson Bracket relations are

$$\begin{aligned} [A_{\pm}, B_{\pm}]_{PB} &= \mp C_{\pm}, \\ [B_{\pm}, C_{\pm}]_{PB} &= A_{\pm}, \\ [C_{\pm}, A_{\pm}]_{PB} &= B_{\pm} \end{aligned} \quad (3.4.7)$$

which in the case of negative energy is immediately recognisable as the Lie algebra  $so(3)$  and in the case of positive energy the noncompact algebra  $so(2, 1)$ . It is remarkable how a little refinement can make things look so attractive!

In the planar version of the Kepler problem the algebra for negative and positive energies is the same as we have found for this problem. Consequently we should not be

surprised if there be a relationship between these conserved quantities as exists for the Kepler problem. After considerable manipulation one finds that the relationship is

$$\frac{1}{2H} \left\{ \left( J_1 + \frac{\alpha^2 \cos \beta}{4H} \right)^2 + \left( J_2 + \frac{\alpha^2 \sin \beta}{4H} \right)^2 - \left( \mu - \frac{\alpha^2}{4H} \right)^2 \right\} = \left( \frac{I}{2H} \right)^2 \quad (3.4.8)$$

which is also somewhat lacking in æsthetic appeal. By defining

$$\mu_{\pm} = |2H|^{-\frac{1}{2}} \left( \mu - \frac{\alpha^2}{4H} \right) \quad (3.4.9)$$

for positive and negative energies respectively we may write (3.4.8) as

$$\mu_{\pm}^2 = A_{\pm}^2 + B_{\pm}^2 \mp C_{\pm}^2 \quad (3.4.10)$$

for positive and negative energies respectively. For negative energies in the space of first integrals equation (3.4.10) represents a sphere which is naturally associated with  $so(3)$  symmetry and for positive energies an hyperboloid of one sheet, the natural geometric object associated with  $so(2, 1)$  symmetry.

When  $H$  takes the particular value of zero, the analysis commencing at (3.4.4) no longer holds true. The third integral is now

$$I = p_r \alpha r^{\frac{1}{2}} \sin \frac{1}{2}(\theta - \beta) + p_{\theta} \alpha r^{-\frac{1}{2}} \cos \frac{1}{2}(\theta - \beta) \quad (3.4.11)$$

which is no longer a true first integral but, rather, a configurational invariant as discussed by Hall [54] and Sarlet *et al* [129] since it is invariant only for the particular value,  $H = 0$ . If we introduce the unit vector  $\hat{\theta}$  defined by

$$\hat{\theta} = \hat{j}_1 \cos \beta + \hat{j}_2 \sin \beta, \quad (3.4.12)$$

where  $\hat{j}_1$  and  $\hat{j}_2$  respectively are the unit vectors along which  $J_1$  and  $J_2$  lie, we see that

$$\mathbf{J} \cdot \hat{\theta} = \frac{I^2}{\alpha^2} - \mu \quad (3.4.13)$$

or, equivalently, that

$$J_1 \cos \beta + J_2 \sin \beta = \frac{I^2}{\alpha^2} - \mu. \quad (3.4.14)$$

Equation (3.4.14) represents the natural geometric object associated with the case  $H = 0$  which describes a right parabolic cylinder with  $\hat{\theta}$  lying along the axis of symmetry.

A distinction can now be drawn between the standard Kepler problem, where, for  $H = 0$ ,  $J = \pm\mu$  from (2.1.10), and the present problem. This is the equation for a plane, unlike (3.4.14). This difference is also reflected in the form of the algebra. We define

$$\begin{aligned} A_0 &= -J_1 \sin \beta + J_2 \cos \beta, \\ B_0 &= \frac{2I}{\alpha^2}, \\ C_0 &= 1. \end{aligned} \quad (3.4.15)$$

The combination  $J_1 \cos \beta + J_2 \sin \beta$  cannot be used as this also is a function of  $I$  from (3.4.14). The Poisson Bracket relations are

$$\begin{aligned} [A_0, B_0]_{PB} &= C_0, \\ [B_0, C_0]_{PB} &= 0, \\ [C_0, A_0]_{PB} &= 0 \end{aligned} \quad (3.4.16)$$

which represents the Weyl algebra,  $W(3, 1)$ . The standard Kepler problem (in two dimensions) has the algebra  $E(2)$  when  $H = 0$ .

For  $H \neq 0$  the Hamiltonian can be expressed as a function of the three first integrals,  $J_1$ ,  $J_2$  and  $I$ , as

$$H = \frac{1}{2(J^2 - \mu^2)} [I^2 - \alpha^2(J_1 \cos \beta + J_2 \sin \beta + \mu)]. \quad (3.4.17)$$

An alternate derivation of the integrals for the Hamiltonian was proposed by Sen [130] as an extension but in the spirit of the work done on first integrals polynomial in the velocities for a variety of Hamiltonian systems by, amongst others, Gascón *et al* [36], Grammaticos *et al* [52], Thompson [136], Hietarinta [62], Leach [82] and, in the case of time-dependent systems, Lewis and Leach [96]. Sen assumes that the first integral is a polynomial in the momenta of the form

$$I_S = \sum_{i=0} \sum_{j=0} d_{ij}(x, y) p_x^i p_y^j, \quad i + j \leq n, \quad (3.4.18)$$

where  $i + j$  is either even or odd in sympathy with  $n$  since autonomous invariants for an autonomous Hamiltonian of even degree in the momenta are either even or odd in the momenta (see Thompson [136]). The extension of Sen is to make a linear canonical transformation to a complex coordinate system

$$\begin{aligned} z &= 2^{-\frac{1}{2}}(x + iy), & \bar{z} &= 2^{-\frac{1}{2}}(x - iy), \\ p_z &= 2^{-\frac{1}{2}}(p_x - ip_y), & p_{\bar{z}} &= 2^{-\frac{1}{2}}(p_x + ip_y). \end{aligned} \quad (3.4.19)$$

In the new coordinates the Hamiltonian and first integral have the forms

$$\begin{aligned} H_S &= 2p_z p_{\bar{z}} + V(z, \bar{z}), \\ I_S &= (z p_z - \bar{z} p_{\bar{z}})^{n-2} p_z p_{\bar{z}} + \sum_{i=0} \sum_{j=0} e_{ij}(z, \bar{z}) p_z^i p_{\bar{z}}^j, \end{aligned} \quad (3.4.20)$$

where now  $i + j \leq n - 2$ . The Poisson Bracket requirement that  $[I_S, H_S]_{PB} = 0$  results in a system of equations the solution of which gives the Hamiltonian

$$H_{I_S} = 2p_z p_{\bar{z}} + a z^{-\frac{1}{2}} + b \bar{z}^{-\frac{1}{2}} + c(z\bar{z})^{-\frac{1}{2}}, \quad (3.4.21)$$

the third-order first integral

$$I_3 = (z p_z - \bar{z} p_{\bar{z}}) p_z p_{\bar{z}} + \frac{1}{2} \left[ b \bar{z}^{-\frac{1}{2}} + c(z\bar{z})^{-\frac{1}{2}} \right] z p_z - \frac{1}{2} \left[ a z^{-\frac{1}{2}} + c(z\bar{z})^{-\frac{1}{2}} \right] \bar{z} p_{\bar{z}} \quad (3.4.22)$$

and the two quadratic first integrals

$$I_1 = (zp_z - \bar{z}p_{\bar{z}})p_z + \frac{1}{2} \left[ b\bar{z}^{\frac{1}{2}} - az^{-\frac{1}{2}}\bar{z} - cz^{-\frac{1}{2}}\bar{z}^{\frac{1}{2}} \right], \quad (3.4.23)$$

$$I_2 = (zp_z - \bar{z}p_{\bar{z}})p_{\bar{z}} + \frac{1}{2} \left[ bz\bar{z}^{\frac{1}{2}} - az^{\frac{1}{2}} + cz^{\frac{1}{2}}\bar{z}^{-\frac{1}{2}} \right]. \quad (3.4.24)$$

When  $b = \bar{a} = a_1 - ia_2$ , the identification of the results of Sen with those given above is

$$\begin{aligned} H_{IS} &= 2H, & -2^{-\frac{1}{2}}(I_1 - I_2) &= J_1, \\ 2iI_3 &= I, & -2^{-\frac{1}{2}}i(I_1 + I_2) &= J_2. \end{aligned} \quad (3.4.25)$$

Naturally the results of the two approaches can be mutually identified when the same matters are being discussed.

The results of Sen, derived by the direct approach of assuming a structure for the first integral, and those obtained here by means of vectorial manipulation of the equation of motion and then the imposition of the Hamiltonian structure are peculiar to two dimensions. The results of a long and tedious calculation [49] show that the only potential in three dimensions which can be permitted subject to the requirement that the Hamiltonian be autonomous and of the form  $H = T + V$  is the familiar Kepler potential. Sen [130] also considered the quantum mechanical representations of the two-dimensional Hamiltonian. The presence of the quadratic first integrals in the classical case implies that the quantum mechanical version of this potential is integrable (see Hietarinta [61]). He obtained the quantum invariants, the identical algebra to the classical algebra and the bound state energy spectrum using the Casimir operator for  $so(3)$ . Considering equation (3.4.10) above one should not be surprised.

### 3.5 Invariance under time translation and the first integrals

Sen [130] showed that the differential equation (3.3.7) possesses only the single Lie point symmetry,  $G = \partial_t$ . Sen interpreted this in the usual way as leading to the conservation of energy, which it does if one applies Noether's Theorem. Using the Lie method Leach [80] constructed the energy, the angular momentum and the components of the Laplace–Runge–Lenz vector for the Kepler problem from this single Lie point symmetry<sup>3.3</sup>.

A Lie symmetry,  $G$ , gives rise to a first integral/invariant of a differential equation for which it is a symmetry if the two conditions

$$G^{[1]}I = 0 \quad \text{and} \quad \frac{dI}{dt} \Big|_{E=0} = 0, \quad (3.5.1)$$

---

<sup>3.3</sup>As is well known (for example, see Kaplan [67]), a system of  $N$  degrees of freedom can have at most only  $2N - 1$  first integrals. The excessive number obtained by Leach followed from different routes towards obtaining the invariants of the associated Lagrange's systems which need to be solved to obtain the integrals. These integrals, energy, angular momentum and the Laplace–Runge–Lenz vector, obviously have the symmetry,  $\partial_t$ , since they are manifestly invariant under time translation. The single integral per symmetry is a feature of Noether's theorem and to obtain the integrals apart from the energy using Noether's theorem one would require additional symmetries. These symmetries would have to be generalised symmetries due to the existence of only one Lie point symmetry.

where the differential equation in question is  $E = 0$  and  $G^{[1]}$  is the first extension of  $G$ , are satisfied. The first extension is required due to the required presence of first derivatives in the first integral/invariant. The symmetry  $\partial_t$  is its own first extension and the first of (3.5.1) gives the linear partial differential equation

$$\frac{\partial I}{\partial t} + 0 \frac{\partial I}{\partial r} + 0 \frac{\partial I}{\partial \theta} + 0 \frac{\partial I}{\partial \dot{r}} + 0 \frac{\partial I}{\partial \dot{\theta}} = 0. \quad (3.5.2)$$

The characteristics obtained from the solution of the associated Lagrange's system

$$\frac{dt}{1} = \frac{dr}{0} = \frac{d\theta}{0} = \frac{d\dot{r}}{0} = \frac{d\dot{\theta}}{0} \quad (3.5.3)$$

are

$$\begin{aligned} u_1 &= r, & v_1 &= \dot{r}, \\ u_2 &= \theta, & v_2 &= \dot{\theta}, \end{aligned} \quad (3.5.4)$$

which are individually invariant under the infinitesimal transformation generated by  $G^{[1]}$ . The second condition of (3.5.1) gives the associated Lagrange's system

$$\frac{du_1}{\dot{u}_1} = \frac{du_2}{\dot{u}_2} = \frac{dv_1}{\dot{v}_1} = \frac{dv_2}{\dot{v}_2}, \quad (3.5.5)$$

where we find from the equation of motion that

$$\begin{aligned} \dot{u}_1 &= v_1, \\ \dot{v}_1 &= u_1 v_2^2 - \frac{\mu}{u_1^2} - \frac{\alpha \cos \frac{1}{2}(u_2 - \beta)}{2u_1^{\frac{3}{2}}}, \\ \dot{u}_2 &= v_2, \\ \dot{v}_2 &= -\frac{2v_1 v_2}{u_1} - \frac{\alpha \sin \frac{1}{2}(u_2 - \beta)}{2u_1^{\frac{5}{2}}}. \end{aligned} \quad (3.5.6)$$

In the case of the Kepler problem the solution of the equation corresponding to (3.5.5) was nontrivial and the righthand sides in (3.5.6) are more complex. The underlying idea of the solution of the associated Lagrange's system is to take combinations of the elements of the system to obtain equivalent elements which have the total derivative of some function in the numerator and zero in the denominator [63, p. 45]. To make the process of solution more obvious, if not precisely transparent, we indicate the combinations used. We denote the  $i$ th element of (3.5.6) by (3.5.6.i). The combination

$$\begin{aligned} &\left( u_1 v_2^2 + \frac{\mu}{u_1^2} + \frac{\alpha \cos \frac{1}{2}(u_2 - \beta)}{2u_1^{\frac{3}{2}}} \right) \quad (3.5.6.1) \\ &+ \frac{\alpha \sin \frac{1}{2}(u_2 - \beta)}{2u_1^{\frac{5}{2}}} \quad (3.5.6.2) + v_1 \quad (3.5.6.3) + u_1^2 v_2 \quad (3.5.6.4) \end{aligned} \quad (3.5.7)$$

gives one of the desired elements, *videlicet*

$$\frac{d \left[ \frac{1}{2} (v_1^2 + u_1^2 v_2^2) - \frac{\mu}{u_1} - \frac{\alpha \cos \frac{1}{2}(u_2 - \beta)}{u_1^{\frac{1}{2}}} \right]}{0}. \quad (3.5.8)$$

The term in crochets is a characteristic and hence a first integral. In the original coordinates one sees that it is the energy, which in this case is the Hamiltonian. The components of the Laplace–Runge–Lenz vector are obtained from the combinations

$$\begin{aligned} & \left( 2u_1 v_1 v_2 \sin u_2 + 3u_1^2 v_2^2 \cos u_2 + \frac{1}{2} \alpha u_1^{-\frac{1}{2}} \sin \frac{1}{2}(u_2 - \beta) \sin u_2 \right) \quad (3.5.6.1) \\ & + \left( u_1^2 v_1 v_2 \cos u_2 - u_1^3 v_2^2 \sin u_2 + \mu \sin u_2 + \frac{1}{2} \alpha u_1^{\frac{1}{2}} \cos \frac{1}{2}(u_2 - \beta) \sin u_2 \right. \\ & \left. + \alpha u_1^{\frac{1}{2}} \sin \frac{1}{2}(u_2 - \beta) \cos u_2 \right) \quad (3.5.6.2) + (u_1^2 v_2 \sin u_2) \quad (3.5.6.3) \\ & + (u_1^2 v_1 \sin u_2 + 2u_1^3 v_2 \cos u_2) \quad (3.5.6.4) \end{aligned} \quad (3.5.9)$$

for  $J_1$  and

$$\begin{aligned} & \left( -2u_1 v_1 v_2 \cos u_2 + 3u_1^2 v_2^2 \sin u_2 - \frac{1}{2} \alpha u_1^{-\frac{1}{2}} \sin \frac{1}{2}(u_2 - \beta) \cos u_2 \right) \quad (3.5.6.1) \\ & + \left( u_1^2 v_1 v_2 \sin u_2 + u_1^3 v_2^2 \cos u_2 - \mu \cos u_2 - \frac{1}{2} \alpha u_1^{\frac{1}{2}} \cos \frac{1}{2}(u_2 - \beta) \cos u_2 \right. \\ & \left. + \alpha u_1^{\frac{1}{2}} \sin \frac{1}{2}(u_2 - \beta) \sin u_2 \right) \quad (3.5.6.2) + (-u_1^2 v_2 \cos u_2) \quad (3.5.6.3) \\ & + (-u_1^2 v_1 \cos u_2 + 2u_1^3 v_2 \sin u_2) \quad (3.5.6.4) \end{aligned} \quad (3.5.10)$$

for  $J_2$ . Finally  $I$  comes from the combination

$$\begin{aligned} & \left( 4u_1 v_2 H + \frac{1}{2} u_1^{-\frac{1}{2}} v_1 \alpha \sin \frac{1}{2}(u_2 - \beta) + \frac{3}{2} u_1^{\frac{1}{2}} v_2 \alpha \cos \frac{1}{2}(u_2 - \beta) \right) \quad (3.5.6.1) \\ & + \left( \frac{1}{2} u_1^{\frac{1}{2}} v_1 \alpha \cos \frac{1}{2}(u_2 - \beta) - \frac{1}{2} u_1^{\frac{3}{2}} v_2 \alpha \sin \frac{1}{2}(u_2 - \beta) \right) \quad (3.5.6.2) \quad (3.5.11) \\ & + \left( u_1^{\frac{1}{2}} \alpha \sin \frac{1}{2}(u_2 - \beta) \right) \quad (3.5.6.3) + \left( 2u_1^2 H + u_1^{\frac{3}{2}} \alpha \cos \frac{1}{2}(u_2 - \beta) \right) \quad (3.5.6.4), \end{aligned}$$

where the invariance of  $H$ , *ie*  $dH = 0$ , has been used.

One cannot plausibly propose that the solution of (3.5.5) is transparent. However, it is one possible method where it is not essential, *a priori*, to make an *Ansatz* for the structure of the first integrals. This method is more feasible when there are several symmetries available for a complex system. Then one can impose the requirements of several symmetries simultaneously and so reduce the number of invariants before imposing the condition  $\dot{I} = 0$ , in this case the second member of (3.5.1), which is generally the major



source of difficulty<sup>3.4</sup>. An example of this procedure in practice is to be found in the paper of Cotsakis *et al* [26] devoted to the determination of the first integrals/invariants of the Bianchi models in Cosmology.

### 3.6 The Kepler problem with ‘drag’

For an equation of motion with the direction of angular momentum conserved we had, in § 3.1, the equation

$$\dot{\mathbf{L}} + h_1 \mathbf{L} = 0 \quad (3.6.1)$$

which we can rewrite as

$$\mathbf{r} \times \ddot{\mathbf{r}} + g_1 \mathbf{r} \times \mathbf{r} + h_1 \mathbf{r} \times \dot{\mathbf{r}} = 0, \quad (3.6.2)$$

where  $g_1$  and  $h_1$  are arbitrary functions for which the variable dependence is specified below when it is necessary so to do. Consequently the equation of motion

$$\ddot{\mathbf{r}} + f\dot{\mathbf{r}} + g\mathbf{r} = 0, \quad (3.6.3)$$

where we have replaced  $h_1$  and  $g_1$  with  $f$  and  $g$  respectively, describes motion in a plane with  $\hat{\mathbf{L}}$  conserved. This equation, (3.6.3), is a generalisation to a class of a problem of physical interest, the equation which models the motion of satellites in low altitude orbits and which was proposed by Brouwer and Hori [18]. They obtained a closed-form solution which included first-order corrections for drag<sup>3.5</sup> acceleration involving a quadratic velocity-dependent term to account for atmospheric effects on satellites sufficiently close to the Earth. Danby [27] modified this assumption and used a resistive term which was proportional to the velocity vector and inversely proportional to the square of the radial distance. For small values of the constant of proportionality he obtained a first-order perturbation solution. Mittleman and Jezewski [107] provided an exact solution to the same problem and in a subsequent paper [66] they demonstrated the existence of analogues of the angular momentum, energy and Laplace–Runge–Lenz vector for the Danby problem. Their approach was to manipulate the equation of motion as had been done by Collinson [22, 23], Pollard [124] and also Sarlet and Bahar [128] on various nonlinear problems. The paper of Mittleman and Jezewski [107] is quite complicated and invited attempts to provide an elegant solution to the original problem of Danby [83] and its generalisation as represented by (3.6.3) [46].

The vector product of  $\mathbf{r}$  with (3.6.3) gives

$$\dot{\mathbf{L}} = -f\mathbf{L}. \quad (3.6.4)$$

This equation may be used to eliminate  $f$  from (3.6.3) to give

$$\ddot{\mathbf{r}} - \frac{\dot{\mathbf{L}}}{L}\dot{\mathbf{r}} + g\mathbf{r} = 0. \quad (3.6.5)$$

---

<sup>3.4</sup>One need not be surprised at this since mathematicians and scientists have been looking for ways to avoid the direct solution of the equations of motion ever since they were devised several centuries ago.

<sup>3.5</sup>The use of the word ‘drag’ to describe this type of force was introduced by Jezewski and Mittleman [107, 66]. We maintain the usage, but slightly tongue in cheek as may be inferred from the ‘*’* in the title of this section.

The first two terms of (3.6.5) possess the integrating factor  $L^{-1}$  and we have

$$\frac{d}{dt} \left( \frac{\dot{\mathbf{r}}}{L} \right) + \frac{g\mathbf{r}}{L} = 0. \quad (3.6.6)$$

There exists a Hamilton-like vector if we can write

$$\mathbf{K} = \frac{\dot{\mathbf{r}}}{L} + \mathbf{u}, \quad (3.6.7)$$

where  $\mathbf{u} := \int (g\mathbf{r}/L)dt$ , explicitly without a knowledge of  $r(t)$  and  $\theta(t)$ . The analogue of the Laplace–Runge–Lenz vector is

$$\mathbf{J} = \mathbf{K} \times \hat{\mathbf{L}} = \frac{\dot{\mathbf{r}} \times \hat{\mathbf{L}}}{L} + \mathbf{u} \times \hat{\mathbf{L}}. \quad (3.6.8)$$

The orbit equation is obtained by taking the scalar product of (3.6.8) with  $\mathbf{r}$  and is

$$r = \frac{1}{-u_\theta + J \cos(\theta - \theta_0)}, \quad (3.6.9)$$

where  $\theta_0$  is the angle between  $\mathbf{J}$  and the cartesian unit vector,  $\hat{\mathbf{i}}$ , and  $u_\theta$  is the component of  $\mathbf{u}$  in the direction  $\hat{\boldsymbol{\theta}}$  (*cf* (3.6.12)). To determine  $\mathbf{u}$  we use the cartesian representation of  $\hat{\mathbf{r}}$  to write

$$\begin{aligned} \mathbf{u} &= \int \frac{g\mathbf{r}}{L} dt \\ &= \hat{\mathbf{i}} \int \frac{gr}{L} \cos \theta dt + \hat{\mathbf{j}} \int \frac{gr}{L} \sin \theta dt \end{aligned} \quad (3.6.10)$$

and the two integrals can be evaluated provided that

$$\frac{gr}{L} = v(\theta)\dot{\theta} \quad \Leftrightarrow \quad g = rv(\theta)\dot{\theta}^2. \quad (3.6.11)$$

Then we may write  $\mathbf{u}$  as (*cf* § 2.1)

$$\mathbf{u} = z'(\theta)\hat{\mathbf{r}} - z(\theta)\hat{\boldsymbol{\theta}}, \quad (3.6.12)$$

where

$$z(\theta) = \int_{\theta_0}^{\theta} v(\eta) \sin(\theta - \eta) d\eta \quad (3.6.13)$$

or, alternatively,

$$z''(\theta) + z(\theta) = v(\theta), \quad z(\theta_0) = 0 \quad \text{and} \quad z'(\theta_0) = 0. \quad (3.6.14)$$

The radial and angular components of  $\mathbf{u}$  are

$$u_r = z'(\theta), \quad u_\theta = -z(\theta) \quad (3.6.15)$$

and the two vectors are

$$\mathbf{K} = \frac{\dot{\mathbf{r}}}{L} + z'(\theta)\hat{\mathbf{r}} - z(\theta)\hat{\boldsymbol{\theta}}, \quad (3.6.16)$$

$$\mathbf{J} = \frac{\dot{\mathbf{r}} \times \hat{\mathbf{L}}}{L} - z(\theta)\hat{\mathbf{r}} - z'(\theta)\hat{\boldsymbol{\theta}}. \quad (3.6.17)$$

The orbit equation, (3.6.9), is now

$$r = \frac{1}{z(\theta) + J \cos(\theta - \theta_0)}. \quad (3.6.18)$$

We now return to (3.6.4). This can be integrated to give

$$M(L, r, \theta) = h, \quad (3.6.19)$$

where  $h$  is the constant of integration, provided

$$f = \frac{\frac{\partial M(L, r, \theta)}{\partial r} \dot{r} + \frac{\partial M(L, r, \theta)}{\partial \theta} \dot{\theta}}{L \frac{\partial M(L, r, \theta)}{\partial L}}. \quad (3.6.20)$$

We may invert (3.6.19), at least locally, to obtain

$$L = N(h, r, \theta). \quad (3.6.21)$$

With this and (3.6.11) rewritten as

$$\frac{gr^3}{L^2} = v(\theta) \quad (3.6.22)$$

by the replacement of  $\dot{\theta}$  by  $L/r^2$ , we obtain an alternative expression for  $g$ , *videlicet*

$$g = \frac{1}{r^3} N^2(h, r, \theta) v(\theta). \quad (3.6.23)$$

Equations (3.6.20) and (3.6.23) establish one possible form of (3.6.3) for which the conserved vectors  $\mathbf{K}$ , (3.6.16), and  $\mathbf{J}$ , (3.6.17), and the orbit equation, (3.6.18), exist.

For the model of the motion of a low altitude satellite Jezewski and Mittleman [66] obtained an energylike first integral by a complicated calculation. For the more general problem discussed here it is also possible to obtain a scalar energylike first integral as

$$\begin{aligned} I &= \frac{1}{2} \mathbf{K} \cdot \mathbf{K} = \frac{1}{2} \mathbf{J} \cdot \mathbf{J} \\ &= \frac{1}{2} \frac{\dot{\mathbf{r}} \cdot \dot{\mathbf{r}}}{L^2} + \frac{1}{L} \left[ \dot{r} z'(\theta) - r \dot{\theta} z(\theta) \right] + \frac{1}{2} [z^2(\theta) + z'^2(\theta)]. \end{aligned} \quad (3.6.24)$$

The radial motion in time may be found by differentiation of (3.6.18) and use of the orbit equation to remove the  $\sin(\theta - \theta_0)$  term. We obtain

$$\dot{r} = -L \left( z'(\theta) - \frac{1}{r} ((J^2 - z^2(\theta))r^2 + 2z(\theta)r - 1)^{\frac{1}{2}} \right), \quad (3.6.25)$$

where  $L = r^2\dot{\theta}$  has been used to eliminate  $\dot{\theta}$ . The only possibility that (3.6.25) is solvable in closed form is that  $z(\theta)$ ,  $z'(\theta)$  and  $L$  can be expressed as functions of  $r$ , apart from any constants of the motion. If this be possible and the subsequent quadrature performed, we obtain

$$t = f_1(r) \quad (3.6.26)$$

and are still left with the problem of the inversion to obtain  $r(t)$ . We treat one class of such problems below.

The angular motion in time is obtained from the two-dimensional angular momentum equation  $L = r^2\dot{\theta}$  and the elimination of  $r$  with (3.6.18). We obtain

$$\dot{\theta} = L(z(\theta) + J \cos(\theta - \theta_0))^2 \quad (3.6.27)$$

and, provided  $L$  can be expressed in terms of  $\theta$ , it may be possible to perform the quadrature to obtain

$$t = f_2(\theta). \quad (3.6.28)$$

The areal velocity is obtained from

$$\dot{A} = \frac{1}{2}L(t), \quad (3.6.29)$$

where  $L(t)$  is obtained as an explicit function of time through the inversion of (3.6.26) and (3.6.28) to replace  $r$  and  $\theta$  by functions of time.

We see that there is much potential for not to be able to obtain explicit expressions. However, the establishment of the existence of the conserved quantities enables us to carry out numerical computations with confidence.

These examples are described in Leach [83] and Gorrington and Leach [44]. Mavraganis [103] considered equations of the form (3.6.3) in which the resistive term varied slowly with time. With the use of a Taylor series expanded about the initial velocity he obtained expressions for the orbit equation and the Hamilton and Laplace–Runge–Lenz vectors which are similar in structure to those described by Jezewski and Mittleman [66] and Leach [83] for the Danby [27] problem.

**Example 3.3.** The equation of motion for the Danby problem is

$$\ddot{\mathbf{r}} + \frac{\alpha \dot{\mathbf{r}}}{r^2} + \frac{\mu \mathbf{r}}{r^3} = 0. \quad (3.6.30)$$

From a comparison of (3.6.30) with (3.6.3) we have

$$f = \frac{\alpha}{r^2}, \quad g = \frac{\mu}{r^3}. \quad (3.6.31)$$

From (3.6.4) the magnitude of the angular momentum satisfies the equation

$$-\frac{\dot{L}}{L} = \frac{\alpha}{r^2} = \frac{\alpha \dot{\theta}}{L} \quad (3.6.32)$$

which has the solution

$$L = k - \alpha\theta, \quad (3.6.33)$$

where  $k$  is some arbitrary constant, in agreement with the result of Jezewski and Mittleman [66]. We also find that

$$v(\theta) = \frac{\mu}{(k - \alpha\theta)^2}, \quad (3.6.34)$$

$$z(\theta) = \mu \int_{\theta_0}^{\theta} \frac{\sin(\theta - \eta) d\eta}{(k - \alpha\eta)^2}. \quad (3.6.35)$$

In terms of

$$u = \left( \frac{k}{\alpha} - \theta \right), \quad u_0 = \left( \frac{k}{\alpha} - \theta_0 \right) \quad (3.6.36)$$

and some standard integrals ([51, 2.641.1–4]) we obtain

$$z(\theta) = \frac{\mu}{\alpha^2} \left[ \frac{\sin(u - u_0)}{u_0} - \sin u (\text{si}(u) - \text{si}(u_0)) - \cos u (\text{Ci}(u) - \text{Ci}(u_0)) \right], \quad (3.6.37)$$

$$z'(\theta) = -\frac{\mu}{\alpha^2} \left[ \frac{\cos(u - u_0)}{u_0} - \frac{1}{u} + \sin u (\text{Ci}(u) - \text{Ci}(u_0)) - \cos u (\text{si}(u) - \text{si}(u_0)) \right],$$

where the sine and cosine integrals are given by

$$\text{si}(x) = -\frac{1}{2}\pi + \int_0^x \frac{\sin t}{t} dt, \quad (3.6.38)$$

$$\text{Ci}(x) = \gamma + \log x + \int_0^x \frac{\cos t - 1}{t} dt \quad (3.6.39)$$

and  $\gamma = 0.57721566490\dots$  is Euler's constant.

The explicit expressions for the conserved vectors are

$$\begin{aligned} \mathbf{K} &= \frac{\dot{\mathbf{r}}}{L} + \frac{\mu}{\alpha^2} \left( \frac{1}{u} - \sin u \text{Ci}(u) + \cos u \text{si}(u) \right) \hat{\mathbf{r}} \\ &\quad + \frac{\mu}{\alpha^2} (\sin u \text{si}(u) + \cos u \text{Ci}(u)) \hat{\boldsymbol{\theta}} \\ &\quad - \frac{\mu}{\alpha^2} \left( \frac{\hat{\mathbf{r}}_{\theta_0}}{u_0} + \text{si}(u_0) \hat{\mathbf{r}}_{k/\alpha} + \text{Ci}(u_0) \hat{\boldsymbol{\theta}}_{k/\alpha} \right), \end{aligned} \quad (3.6.40)$$

$$\begin{aligned} \mathbf{J} &= \frac{\dot{\mathbf{r}} \times \hat{\mathbf{L}}}{L} + \frac{\mu}{\alpha^2} (\sin u \text{si}(u) + \cos u \text{Ci}(u)) \hat{\mathbf{r}} \\ &\quad - \frac{\mu}{\alpha^2} \left( \frac{1}{u} - \sin u \text{Ci}(u) + \cos u \text{si}(u) \right) \hat{\boldsymbol{\theta}} \\ &\quad + \frac{\mu}{\alpha^2} \left( \frac{\hat{\boldsymbol{\theta}}_{\theta_0}}{u_0} + \text{si}(u_0) \hat{\boldsymbol{\theta}}_{k/\alpha} - \text{Ci}(u_0) \hat{\mathbf{r}}_{k/\alpha} \right). \end{aligned} \quad (3.6.41)$$

Since the final term in both (3.6.40) and (3.6.41) is a constant vector, it can be ignored in order to simplify matters. The simplified version coincides with that obtained by Jezewski and Mittleman [66] who obtained their result by means of an integrating factor. For either version the equation for the orbit comes from the scalar product of  $\mathbf{r}$  and  $\mathbf{J}$ .

A typical orbit, for which see Gorringer and Leach [44], spirals inwards which is not surprising considering what the equation of motion was intended to model.

**Example 3.4.** The density of the atmosphere is not constant for all  $\theta$ . A more realistic resistive force for the motion of a low altitude satellite which incorporates a variation in the density of the atmosphere is given by

$$f = \frac{a \cos \theta + b}{r^2}, \quad g = \frac{\mu}{r^3}. \quad (3.6.42)$$

In this case

$$z(\theta) = \mu \int_{\theta_0}^{\theta} \frac{\sin(\theta - \eta) d\eta}{(k - a \sin \eta - b\eta)^2}. \quad (3.6.43)$$

The nature of the orbit depends upon the relative values of the constants  $a$  and  $b$ . For values which give a resistive force for all values of  $\theta$ , the orbit spirals inwards, as expected, but the rate at which it spirals inwards varies with  $\theta$  to give almost the impression of a precession of the orbit.

**Example 3.5.** There is no necessity to maintain an air of reality in the model if one wants to explore the variety of orbits which can be obtained. If we make the choice

$$f = \frac{a \cos \theta + b}{Lr^2}, \quad g = \frac{\mu}{r^3}, \quad (3.6.44)$$

we find that

$$z(\theta) = \frac{1}{2}\mu \int_{\theta_0}^{\theta} \frac{\sin(\theta - \eta) d\eta}{k - a \sin \eta - b\eta}. \quad (3.6.45)$$

The “precession” of the orbit is even more pronounced. In the case that the force is not always resistive the angular momentum increases and eventually the orbit appears to depart asymptotically.

**Example 3.6.** In the case that

$$f = \frac{-ae^{-(\theta-\theta_0)}L^3}{2r^2}, \quad g = \frac{\mu}{r^3} \quad (3.6.46)$$

for which

$$z(\theta) = \frac{1}{2}\mu a \left( e^{-(\theta-\theta_0)} + \sin(\theta - \theta_0) - \cos(\theta - \theta_0) \right) + \mu b (1 - \cos(\theta - \theta_0)) \quad (3.6.47)$$

the “resistive” force decreases to zero as  $\theta$  increases. Even for small values of  $\theta$  the value of  $f$  is small. The orbit approaches a normal Keplerian orbit asymptotically. For illustrations of this and other orbits see Gorrington and Leach [44].

### 3.7 Force laws admitting Keplerian orbits

Commencing with (3.6.3) we construct the most general Keplerian orbits, *ie* conic sections centred on a focus. This is motivated by the results of Bertrand’s theorem [13] which applies specifically to central force orbits. We are not particularly concerned with the inverse problem of finding the most general equation of motion for a given orbit. This

has a long history for which see Whittaker [140] and more recently Broucke [17] and the references cited therein.

The Keplerian orbit has its origin at one of the foci and from (3.6.18) we see that, since  $z(\theta_0) = 0$ ,

$$z(\theta) = A(1 - \cos(\theta - \theta_0)), \quad (3.7.1)$$

where  $A$  is some constant. With this and (3.6.14) and (3.6.23) we obtain

$$v(\theta) = A \quad \text{and} \quad g = \frac{AL^2}{r^3}. \quad (3.7.2)$$

We solve this for  $L^2$  and differentiate with respect to time to obtain

$$2L\dot{L} = \frac{\dot{g}r^3}{A} + \frac{3r^2\dot{r}g}{A}. \quad (3.7.3)$$

Hence we find that

$$f = -\frac{1}{2} \left( \frac{\dot{g}}{g} + 3\frac{\dot{r}}{r} \right). \quad (3.7.4)$$

Thus the most general equation of motion possessing the conserved vectors  $\mathbf{K}$  and  $\mathbf{J}$  and, in addition, an orbit equation which is a conic section is given by

$$\ddot{\mathbf{r}} - \frac{1}{2} \left( \frac{\dot{g}}{g} + 3\frac{\dot{r}}{r} \right) \dot{\mathbf{r}} + g\mathbf{r} = 0. \quad (3.7.5)$$

Under the change of timescale  $t \rightarrow \rho(t)$  (3.7.5) becomes

$$\dot{\rho}^2 \frac{d^2 \mathbf{r}}{d\rho^2} + \left( \ddot{\rho} - \frac{1}{2} \left( \frac{\dot{g}}{g} + 3\frac{\dot{r}}{r} \right) \dot{\rho} \right) \frac{d\mathbf{r}}{d\rho} + g\mathbf{r} = 0. \quad (3.7.6)$$

We define the transformation by setting the middle term to zero to obtain the equation of motion

$$\frac{d^2 \mathbf{r}}{d\rho^2} + \frac{g}{\dot{\rho}^2} \mathbf{r} = 0, \quad (3.7.7)$$

with the transformation defined by

$$\ddot{\rho} - \frac{1}{2} \left( \frac{\dot{g}}{g} + 3\frac{\dot{r}}{r} \right) \dot{\rho} = 0 \quad (3.7.8)$$

$$\implies \dot{\rho}^2 = \frac{gr^3}{\mu} = \frac{AL^2}{\mu} = \frac{4A}{\mu} \dot{S}^2 = \frac{4}{\bar{L}^2} \dot{S}^2, \quad (3.7.9)$$

where  $\mu$  is the arbitrary constant in the Kepler problem

$$\frac{d^2 \mathbf{r}}{d\rho^2} + \frac{\mu}{r^3} \mathbf{r} = 0, \quad (3.7.10)$$

$\bar{L}$  is the angular momentum for (3.7.10) and  $\dot{S} = \frac{1}{2}L$  is the areal velocity of (3.7.5). Equation (3.7.5) has been transformed into an equivalent Kepler problem (3.7.10) using

the area swept out in the orbit described by (3.7.5). Equation (3.7.9) is consistent with that obtained by taking the area  $S(t)$  swept out in the orbit described by (3.7.5) in time  $t$  with the same area swept out in the Kepler problem at time  $\rho(t)$  as given by Kepler's Second Law, *ie*  $\rho(t) = 2S(t)/\bar{L}$ . The conserved vectors,  $\mathbf{K}$  and  $\mathbf{J}$ , of (3.7.5) are form invariant under the transformation  $t \rightarrow \rho(t)$  since the term  $\dot{\mathbf{r}} \times \mathbf{L}/L^2 = \mathbf{r}' \times \mathbf{L}_t/L_t^2$ , where the prime denotes differentiation with respect to  $\rho$  and  $L_t = r^2\theta'$ , and so

$$\begin{aligned}\mathbf{K} &= \frac{\mathbf{r}'}{L_t} - A(\hat{\boldsymbol{\theta}} - \hat{\boldsymbol{\theta}}_{\theta_0}), \\ \mathbf{J} &= \frac{\mathbf{r}' \times \mathbf{L}_t}{L_t} - A(\mathbf{r} - \hat{\mathbf{r}}_{\theta_0}),\end{aligned}\tag{3.7.11}$$

where  $\hat{\mathbf{r}}_{\theta_0} = \hat{\mathbf{i}} \cos \theta_0 + \hat{\mathbf{j}} \sin \theta_0$  and  $\hat{\boldsymbol{\theta}}_{\theta_0} = -\hat{\mathbf{i}} \sin \theta_0 + \hat{\mathbf{j}} \cos \theta_0$ .

The angular momentum for the equation of motion (3.7.5) is obtained by taking the vector product with  $\mathbf{r}$  to obtain

$$\dot{\mathbf{L}} - \frac{1}{2} \left( \frac{\dot{g}}{g} + 3\frac{\dot{r}}{r} \right) \mathbf{L} = 0\tag{3.7.12}$$

and, since  $\hat{\mathbf{L}}$  is constant, the scalar part of (3.7.12) gives on integration

$$L = kg^{\frac{1}{2}}r^{\frac{3}{2}},\tag{3.7.13}$$

where  $k$  is the constant of integration. With the explicit form  $L = r^2\dot{\theta}$

$$k = \left( \frac{r}{g} \right)^{\frac{1}{2}} \dot{\theta}\tag{3.7.14}$$

is also a constant of the motion and  $\mathbf{k} = k\hat{\mathbf{L}}$  can be regarded as a conserved generalised angular momentum.

We obtain the scalar energylike first integral for (3.7.5) by simplifying (3.6.24) using (3.7.1) to substitute for  $z(\theta)$  and its derivatives. The integral is

$$I = \frac{1}{2} \mathbf{J} \cdot \mathbf{J} = \frac{1}{2} A \frac{\dot{\mathbf{r}} \cdot \dot{\mathbf{r}}}{gr^3} - \frac{A}{r} + AJ.\tag{3.7.15}$$

An alternate route to obtain an energylike first integral for (3.7.5) is to take the scalar product with  $(gr^3)^{-1}\dot{\mathbf{r}}$ , which is an integrating factor, and to obtain the conserved quantity

$$E = \frac{1}{2} \frac{\dot{\mathbf{r}} \cdot \dot{\mathbf{r}}}{gr^3} - \frac{1}{r} = \frac{1}{2} J^2 A - J.\tag{3.7.16}$$

The differential equation for the radial motion in time comes from (3.7.16) and (3.7.13) and is

$$\dot{r} = [gr^2(2Er - k^2r - 2)]^{\frac{1}{2}}.\tag{3.7.17}$$

The integration of (3.7.17) can only be possible if  $g$  be expressible solely as a function of  $r$  and, even then, in general the quadrature would be formal. The angular motion in time is found from the integration of

$$\dot{\theta} = \left[ \frac{g}{A} (A + (J - A) \cos(\theta - \theta_0)) \right]^{\frac{1}{2}}.\tag{3.7.18}$$



Now the possibility of quadrature requires that  $g$  be expressible in terms of  $\theta$  only. The aerial velocity is

$$\dot{A} = \frac{1}{2}L = \frac{1}{2}k (gr^3)^{\frac{1}{2}}. \quad (3.7.19)$$

The quadrature to obtain  $A(t)$  requires that the right hand side of (3.7.19) be expressible as a function of time. In general this would require not only the ability to perform the quadrature in either (3.7.17) or (3.7.18) but also the ability to invert the result.

### 3.8 The geometry of the generalised Kepler problem

In general an expression for the area swept out by the radius vector in time cannot be found explicitly due to problems with the inversion of functions at more than a local level. However, in the case that  $g = \mu r^\alpha$ , expressions for the periodic time can be found rather elegantly and also informatively<sup>3,6</sup> [48]. The equation of motion, (3.7.5), is now

$$\ddot{\mathbf{r}} - \frac{1}{2}(\alpha + 3)\frac{\dot{r}}{r}\dot{\mathbf{r}} + \mu r^\alpha \mathbf{r} = 0. \quad (3.8.1)$$

To be consistent with the conserved vectors given for the standard Kepler problem in Chapter 2 we delete the additional constant terms in the conserved vectors given by (3.6.41). By means of the usual operations and techniques of this Chapter we obtain

$$\dot{\mathbf{L}} - \frac{1}{2}(\alpha + 3)\frac{\dot{r}}{r}\mathbf{L} = 0 \quad \implies \quad L = kr^{\frac{1}{2}(\alpha+3)}. \quad (3.8.2)$$

Since the motion is planar,  $L = r^2\dot{\theta}$  and so we have

$$k = r^{-\frac{1}{2}(\alpha-1)}\dot{\theta} \quad (3.8.3)$$

is a constant of the motion. The Hamilton's vector,

$$\mathbf{K} = \frac{\dot{\mathbf{r}}}{L} - \frac{\mu}{k^2}\hat{\boldsymbol{\theta}}, \quad (3.8.4)$$

follows from the integration of the combination of (3.8.1) divided by  $L$  and (3.8.3). The Laplace–Runge–Lenz vector is

$$\mathbf{J} = \mathbf{K} \times \hat{\mathbf{L}} = \frac{\dot{\mathbf{r}} \times \mathbf{L}}{L^2} - \frac{\mu}{k^2}\hat{\mathbf{r}} \quad (3.8.5)$$

and from this we obtain the equation of the orbit

$$r = \frac{1}{\mu/k^2 + J \cos(\theta - \theta_0)}, \quad (3.8.6)$$

where  $\theta_0$  is the angle between  $\mathbf{J}$  and  $\hat{\mathbf{i}}$ .

---

<sup>3,6</sup>Interestingly there is a change in the Lie symmetry properties of the equation of motion [121], for which see § 5.8.2.

The scalar product of  $r^{-(\alpha+3)}\dot{\mathbf{r}}$  with (3.8.1) gives the energylike integral

$$E = \frac{1}{2} \frac{\dot{\mathbf{r}} \cdot \dot{\mathbf{r}}}{r^{\alpha+3}} - \frac{\mu}{r} \quad (3.8.7)$$

after integration with respect to time. The conserved quantities  $J$ ,  $k$  and  $E$  are related according to

$$J^2 = K^2 = \frac{2E}{k^2} + \frac{\mu^2}{k^4}. \quad (3.8.8)$$

The period of the motion is found from (3.8.3) and (3.8.6) to be expressible as the quadrature

$$T = \frac{2}{k} \int_{\theta_0}^{\pi+\theta_0} \left( \frac{\mu}{k^2} + J \cos(\theta - \theta_0) \right)^{\frac{1}{2}(\alpha-1)} d\theta. \quad (3.8.9)$$

Using (3.8.8) we may rewrite the term within the large parentheses in the integrand of (3.8.9) as

$$\left( \frac{-2E}{k^2} \right)^{\frac{1}{2}} \left\{ \frac{\mu}{k(-2E)^{\frac{1}{2}}} + \left[ \left( \frac{\mu}{k(-2E)^{\frac{1}{2}}} \right)^2 - 1 \right]^{\frac{1}{2}} \cos(\theta - \theta_0) \right\}. \quad (3.8.10)$$

The definition of the Legendre function of the first kind is ([51] [8.822.1])

$$P_\nu(z) = \frac{1}{\pi} \int_0^\pi \frac{d\xi}{\left[ z + (z^2 - 1)^{\frac{1}{2}} \cos \xi \right]^{\nu+1}} = \frac{1}{\pi} \int_0^\pi \left[ z + (z^2 - 1)^{\frac{1}{2}} \cos \xi \right]^\nu d\xi. \quad (3.8.11)$$

When  $\nu$  is an integer, this becomes the Legendre polynomial.

The energylike integral, (3.8.7), is negative for an elliptical orbit. With (3.8.10) and (3.8.11) the quadrature of the right-hand side of (3.8.9) can be performed to give the period as

$$T = \frac{2\pi}{k} \left( \frac{-2E}{k^2} \right)^{\frac{1}{4}(\alpha-1)} P_{\frac{1}{2}(\alpha-1)}(z), \quad (3.8.12)$$

where the argument of the Legendre function of the first kind is given by

$$z = \frac{\mu}{k(-2E)^{\frac{1}{2}}}. \quad (3.8.13)$$

The relationship between the semilatus rectum, the semimajor axis and the eccentricity of an ellipse is

$$l = R(1 - e^2), \quad (3.8.14)$$

where the symbols are in the same order. From the orbit equation (3.8.6) the semilatus rectum and the semimajor axis are given by

$$\begin{aligned} l &= r\left(\frac{1}{2}\pi\right) & R &= \frac{1}{2}[r(0) + r(\pi)] \\ &= \frac{k^2}{\mu} & &= \frac{\mu/k^2}{\mu^2/k^4 - J^2} \\ & & &= \frac{\mu}{-2E}, \end{aligned} \quad (3.8.15)$$

where we have used (3.8.8) in the third line of the second column, so that  $z$ , the argument of the generalised Kepler's Third Law, becomes

$$z = (1 - e^2)^{-\frac{1}{2}} \quad (3.8.16)$$

and the law itself can be written as

$$T^2 R^\alpha = \frac{4\pi^2}{\mu} (1 - e^2)^{-\frac{1}{2}(\alpha+1)} P_{\frac{1}{2}(\alpha-1)}^2 \left[ (1 - e^2)^{-\frac{1}{2}} \right] \quad (3.8.17)$$

when (3.8.12) is squared.

The differential equation (3.8.1) is invariant under the similarity transformation

$$(t, \mathbf{r}) \longrightarrow (\bar{t}, \bar{\mathbf{r}} : t = \gamma\bar{t}, \mathbf{r} = \gamma^{\frac{2}{\alpha}}\bar{\mathbf{r}}). \quad (3.8.18)$$

Since

$$E = \gamma^{\frac{2}{\alpha}}\bar{E} \implies R = \gamma^{-\frac{2}{\alpha}}\bar{R}, \quad (3.8.19)$$

it follows that

$$\frac{R^{-\frac{\alpha}{2}}}{t} = \frac{\left(\gamma^{-\frac{2}{\alpha}}\bar{R}\right)^{-\frac{\alpha}{2}}}{\gamma\bar{t}} = \frac{\gamma\bar{R}^{-\frac{\alpha}{2}}}{\gamma\bar{t}} = \frac{\bar{R}^{-\frac{\alpha}{2}}}{\bar{t}} \quad (3.8.20)$$

is invariant under the transformation. This implies that  $TR^{\frac{\alpha}{2}}$  is invariant. In the case of the angular momentum

$$L = \gamma^{-\frac{4}{\alpha}-1}\bar{L} \implies l = \gamma^{-\frac{2}{\alpha}}\bar{l} \quad (3.8.21)$$

so that the ratio

$$\frac{\bar{l}}{\bar{R}} = \frac{\gamma^{\frac{2}{\alpha}}l}{\gamma^{\frac{2}{\alpha}}R} = \frac{l}{R} = (1 - e^2) \quad (3.8.22)$$

is invariant and hence the eccentricity,  $z$ , is also invariant. Alternatively we can use the first extension of the infinitesimal generator of the transformation, *videlicet*

$$G = t\partial_t - \frac{2}{\alpha}r\partial_r \quad (3.8.23)$$

to show the invariance of (3.8.17). The task is simplified somewhat by rewriting  $G^{[1]}(t, r, \dot{r}, \dot{\theta})$  in terms of the relevant conserved quantities, *ie* as  $G^{[1]}(T, R, l)$ . For then we have

$$\begin{aligned} G^{[1]} &= t\dot{t} - \frac{2}{\alpha}r\dot{r} - \frac{\alpha+2}{\alpha}\dot{r}\dot{r} - \dot{\theta}\dot{\theta} \\ \implies G^{[1]} &= T\partial_T - \frac{2}{\alpha}R\partial_R - \frac{2}{\alpha}l\partial_l \end{aligned} \quad (3.8.24)$$

when acting upon functions of  $T$ ,  $R$  and  $l$ . In the case of (3.8.17) it is easy to show that

$$G^{[1]}(T^2R^\alpha) = G^{[1]}\left(p\left(\left(1-e^2\right)^{-\frac{1}{2}}\right)\right) = G^{[1]}\left(p\left(\left(\frac{R}{l}\right)^{\frac{1}{2}}\right)\right) = 0, \quad (3.8.25)$$

where  $p(\cdot)$  denotes the function on the right side of (3.8.17), which illustrates the invariance.

In summary the equation of motion

$$\ddot{\mathbf{r}} - \frac{1}{2}(\alpha+3)\frac{\dot{r}}{r}\dot{\mathbf{r}} + \mu r^\alpha \mathbf{r} = 0 \quad (3.8.26)$$

describes motion in the plane which has the following properties:

(i) the orbit is a conic section with the origin at a focus, *ie* of the type of the Kepler problem,

(ii) the areal velocity is  $\frac{1}{2}kr^{\frac{1}{2}(\alpha+3)}$  and

(iii) in the case of an elliptical orbit the period and semimajor axis are related by

$$T^2R^\alpha = \frac{4\pi^2}{\mu}(1-e^2)^{-\frac{1}{2}(\alpha+1)}P_{\frac{1}{2}(\alpha-1)}^2\left[(1-e^2)^{-\frac{1}{2}}\right]. \quad (3.8.27)$$

For  $\alpha = -3$  in (3.8.26) we obtain the usual Kepler's laws of planetary motion and, in particular, we see that  $T^2 \propto R^3$  irrespective of the value of the eccentricity of the ellipse. In general the areal velocity is not constant. The equation (3.8.17) is the generalised Kepler's Third Law for a power law central force plus a resistive term. We note that the constant of proportionality depends explicitly upon the eccentricity of the orbit. When  $\alpha = -3$ , the selfsimilar transformation (3.8.23) maps solutions into solutions of the same eccentricity (see Prince and Eliezer [125]). This is not reflected in the relationship between the period and the semimajor axis. However, for general  $\alpha$  not only does the symmetry maps solutions into solutions of the same eccentricity but in general the relationship between the period and the semimajor axis now depends explicitly upon the eccentricity. We note that several cases arise in which the constant of proportionality in (3.8.17) does not depend upon the eccentricity. This occurs for the Kepler problem for which  $\alpha = -3$  and when  $\alpha = -1$ . In the cases  $\alpha = 1$  and  $\alpha = 3$  the constant of proportionality can be made independent of the eccentricity if the semilatus rectum,  $l = R(1-e^2)$ , is used instead of the semimajor axis length. We obtain

$$T^2l = \frac{4\pi^2}{\mu} \quad \text{and} \quad T^2l^3 = \frac{4\pi^2}{\mu} \quad (3.8.28)$$

respectively. (Note that the units of  $\mu$  depend upon the value of  $\alpha$ .) In these two instances the implication is that all orbits with the same length of semilatus rectum have the same period. In the case that  $\alpha = 0$  in (3.8.26) we have an oscillator with an additional velocity-dependent force which has the effect of changing the usual geometric-centred oscillator ellipse into a focus-centred Kepler ellipse. The isochronism of the oscillator is now only preserved for all orbits of the same eccentricity.

From the energy equation, (3.8.7), we obtain

$$\dot{r} = r^{\frac{1}{2}(\alpha+1)} (2Er^2 + 2\mu r - k^2)^{\frac{1}{2}}. \quad (3.8.29)$$

This equation can be integrated in closed form for many values of  $\alpha$ . In particular for  $\alpha = \pm 1$  we can not only perform the quadrature in closed form but also invert it ([51, formulæ 2.26ff]) to obtain  $r(t)$ . Consequently it is possible to obtain expressions for both  $\theta(t)$  and  $S(t)$ . We have

1.  $\alpha = -1$

$$\begin{aligned} r(t) &= \frac{\mu}{(-2E)} - \frac{k^2 J}{(-2E)} \cos \left( (-2E)^{\frac{1}{2}} (t - t_0) \right), \\ \theta(t) &= \theta_0 + 2 \arctan \left( \frac{k(-2E)^{\frac{1}{2}}}{(\mu - k^2 J)} \tan \left( \frac{1}{2} (-2E)^{\frac{1}{2}} (t - t_0) \right) \right), \\ S(t) &= \frac{1}{2} \left[ \frac{\mu k (t - t_0)}{(-2E)} - \frac{k^3 J}{(-2E)^{\frac{3}{2}}} \sin \left( (-2E)^{\frac{1}{2}} (t - t_0) \right) \right]. \end{aligned} \quad (3.8.30)$$

2.  $\alpha = 1$

$$\begin{aligned} r(t) &= \frac{1}{\mu/k^2 + J \cos(k(t - t_0))}, \\ \theta(t) &= \theta_0 + k(t - t_0), \\ S(t) &= \frac{1}{2} \left[ \frac{2\mu k}{(-2E)^{\frac{3}{2}}} \arctan \left( \frac{k(-2E)^{\frac{1}{2}} \tan(\frac{1}{2}k(t - t_0))}{\mu + k^2 J} \right) \right. \\ &\quad \left. - \frac{k^2 J \sin(k(t - t_0))}{(-2E)(\mu/k^2 + J \cos(k(t - t_0)))} \right]. \end{aligned} \quad (3.8.31)$$

In this case for  $S(t)$  the formulæ of [51, 2.554.3 and 2.2553.3] were used.

In the second case, that of  $\alpha = 1$ , it is rather interesting that  $\theta$  increases linearly with time. We note that the function  $\rho(t)$  introduced in (3.7.6) can be obtained explicitly in these two cases from the relationship  $\rho(t) = 2S(t)/\bar{L}$ . Since the orbit is elliptical and the area swept out in the corresponding Kepler problem is simply a constant multiple of time, one would expect intuitively that the time transformation required to transform (3.8.26) to the equivalent Kepler equation of motion should involve the area swept out in the original elliptical orbit.

## 4 Motion with conserved $L$

### 4.1 A conserved Laplace–Runge–Lenz vector for $\ddot{\mathbf{r}} + f\mathbf{L} + g\hat{\mathbf{r}} = 0$

In the previous chapter we considered equations of motion for which the direction of the angular momentum vector was constant. In this chapter we require that the magnitude of the angular momentum vector,  $L$ , be constant. We write

$$L^2 = \mathbf{L} \cdot \mathbf{L} \quad (4.1.1)$$

and upon differentiation of this obtain

$$\dot{\mathbf{L}} \cdot \mathbf{L} = 0. \quad (4.1.2)$$

Equation (4.1.2) implies that  $\dot{\mathbf{L}}$  and  $\mathbf{L}$  are orthogonal. Equations of motion arising from  $\dot{\mathbf{L}}$  and combinations of other vectors orthogonal to  $\mathbf{L}$  describe motion for which  $L$  is conserved. One such orthogonal vector is  $\mathbf{r} \times \mathbf{L}$  and so the equation

$$\dot{\mathbf{L}} + f\mathbf{r} \times \mathbf{L} = 0 \quad (4.1.3)$$

describes one possible, not to be regarded as the most general, class of equations meeting the requirement that the magnitude of the angular momentum be conserved. Since  $\dot{\mathbf{L}} = \mathbf{r} \times \ddot{\mathbf{r}}$  and  $\mathbf{r} \times \dot{\mathbf{r}} = 0$ , we may write (4.1.3) as

$$\mathbf{r} \times \ddot{\mathbf{r}} + f\mathbf{r} \times \mathbf{L} + g\mathbf{r} \times \hat{\mathbf{r}} = 0 \quad (4.1.4)$$

from which it follows that the equation of motion

$$\ddot{\mathbf{r}} + f\mathbf{L} + g\hat{\mathbf{r}} = 0, \quad (4.1.5)$$

where for the present  $f$  and  $g$  are arbitrary functions with their variables to be specified below when it is necessary, describes motion subject to the constraint that  $L$  be constant. We note that it is possible to introduce additional terms of the form  $e\dot{\mathbf{r}} \times \mathbf{L}$  into (4.1.3), but we do not consider them here. With the techniques described earlier in this work we now construct Laplace–Runge–Lenz vectors for (4.1.5) by the means of the imposition of restrictions upon the so far arbitrary functions  $f$  and  $g$ .

The vector product of (4.1.5) with  $\mathbf{L}$  gives

$$\ddot{\mathbf{r}} \times \mathbf{L} + g\hat{\mathbf{r}} \times \mathbf{L} = 0. \quad (4.1.6)$$

With the aid of (4.1.3) and making the vector triple expansion of  $\dot{\mathbf{r}} \times (\mathbf{L} \times \mathbf{r})$  we rewrite (4.1.6) in the equivalent form

$$(\dot{\mathbf{r}} \times \mathbf{L}) \cdot -f r \dot{\mathbf{r}} \mathbf{L} + \frac{g}{r} \mathbf{r} \times \mathbf{L} = 0. \quad (4.1.7)$$

The ultimate term in the left hand side of (4.1.7) may be written in the two alternate forms

$$\frac{g}{r} \mathbf{r} \times \mathbf{L} = -\frac{g}{rf} \dot{\mathbf{L}} \quad \text{and} \quad \frac{g}{r} \mathbf{r} \times \mathbf{L} = -gr^2 \dot{\hat{\mathbf{r}}} \quad (4.1.8)$$

thereby prompting the suggestion that we write

$$g = g_1 + g_2 \quad (4.1.9)$$

so that (4.1.7) becomes

$$(\dot{\mathbf{r}} \times \mathbf{L}) \cdot -f r \dot{\mathbf{r}} \mathbf{L} - \frac{g_1}{r f} \dot{\mathbf{L}} - g_2 r^2 \dot{\hat{\mathbf{r}}} = 0. \quad (4.1.10)$$

We now impose constraints on the as yet arbitrary functions  $f$ ,  $g_1$  and  $g_2$  to make (4.1.10) trivially integrable. We set

$$f = \frac{h'(r)}{r}, \quad g_1 = h(r)h'(r) \quad \text{and} \quad g_2 = \frac{k}{r^2}. \quad (4.1.11)$$

Hence the system described by the equation of motion [88]

$$\ddot{\mathbf{r}} + \frac{h'(r)}{r} \mathbf{L} + \left( h(r)h'(r) + \frac{k}{r^2} \right) \hat{\mathbf{r}} = 0 \quad (4.1.12)$$

possesses the conserved Laplace–Runge–Lenz vector

$$\mathbf{J} = \dot{\mathbf{r}} \times \mathbf{L} - h(r)\mathbf{L} - k\hat{\mathbf{r}}. \quad (4.1.13)$$

It does not seem possible to construct a Hamilton's vector for (4.1.13) for general  $h(r)$ . The scalar product of  $\dot{\mathbf{r}}$  with (4.1.12) yields the energylike integral

$$I = \frac{1}{2} \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} + \frac{1}{2} h^2(r) - \frac{k}{r}. \quad (4.1.14)$$

The conserved scalars are not independent since

$$J^2 = 2L^2 I + k^2. \quad (4.1.15)$$

If we measure the polar angle  $\theta$  from  $\mathbf{J}$ , the scalar product of (4.1.13) with  $\mathbf{r}$  gives

$$r = \frac{L^2}{k + J \cos \theta} \quad (4.1.16)$$

after some rearrangement. However, as the direction of the angular momentum vector is not constant, the motion is truly three-dimensional and we need not one but two equations to describe the orbit. We observe that (4.1.16) is independent of the azimuthal angle,  $\phi$ , *ie* we have the radial distance,  $r$ , as a function of the polar angle only. Consequently any surface upon which the position vector moves will be a surface of revolution about the polar axis. For the second equation we need to involve the azimuthal angle,  $\phi$ . We achieve this by taking the scalar product of  $\mathbf{J}$  with  $r^{-2} \hat{\phi}$ . This gives

$$\dot{r} \dot{\phi} \sin \theta + h \dot{\theta} = 0 \quad \implies \quad \dot{\phi} = -\frac{h \dot{\theta}}{\dot{r} \sin \theta}. \quad (4.1.17)$$

By differentiating (4.1.16) with respect to time we obtain

$$\dot{r} = \frac{J r^2 \sin \theta \dot{\theta}}{L^2} \quad \implies \quad \dot{\phi} = -\frac{h L^2}{J r^2 \sin 2\theta}. \quad (4.1.18)$$

In terms of spherical polar coordinates

$$L^2 = r^4 \left( \dot{\theta}^2 + \dot{\phi}^2 \sin^2 \theta \right) \quad (4.1.19)$$

and this in conjunction with (4.1.18) enables us to write

$$\dot{\theta}^2 = \frac{L^2 (J^2 \sin^2 \theta - h^2 L^2)}{J^2 r^4 \sin^2 \theta}. \quad (4.1.20)$$

We can then express the azimuthal angle as

$$\begin{aligned} \phi &= \int_{\theta_0}^{\theta} \frac{\dot{\phi}}{\dot{\theta}} d\theta \\ &= - \int_{\theta_0}^{\theta} \frac{hL}{\sin \theta} (J^2 \sin^2 \theta - h^2 L^2)^{-\frac{1}{2}} d\theta. \end{aligned} \quad (4.1.21)$$

From the orbit equation (4.1.16)  $h(r)$  can be expressed in terms of  $\theta$ . Equations (4.1.16) and (4.1.21) fully describe the orbit. Generally we would not expect to be able to express the quadrature in (4.1.21) in closed form. The relationship between  $\phi$  and  $\theta$  expressed through (4.1.21) is critical for the existence of closed rather than surface filling orbits. The existence of the term,  $\sin \theta$ , in the denominator means that in general the pole is avoided. From the second term in the denominator it is also possible that a singularity will occur for an angle in  $0 < \theta_0 < \theta < \pi$  and so the motion be confined to a zone of the surface of revolution.

From (4.1.20) we can write

$$t = \int_{\theta_0}^{\theta} \frac{JL^3 \sin \theta d\theta}{(k + J \cos \theta)^2 (J^2 \sin^2 \theta - h^2(r(\theta))L^2)^{\frac{1}{2}}}. \quad (4.1.22)$$

If this can be inverted to obtain  $\theta$  as a function of  $t$ , the orbit equation (4.1.16) can be written as

$$r(t) = \frac{L^2}{k + J \cos(\theta(t))}. \quad (4.1.23)$$

Similarly  $\phi(t)$  is obtained from (4.1.21) by the replacement of  $\theta$  with the inverse of (4.1.22). In practice, however, it is unlikely that the quadratures could be achieved in closed form and, even if the unlikely were to happen, one then has the problem of the inversion. Nevertheless everything is sufficiently well-defined for numerical integrations to be used. Finally we note that, although the direction of the areal velocity is not constant, its magnitude is and so we have equal areas swept out in equal times, *ie*

$$A = \frac{1}{2}Lt. \quad (4.1.24)$$

We have already noted that the motion occurs on a surface of revolution. If this be a recognisable surface, by expressing the area in terms of well-known geometric quantities, one could possibly obtain an expression relating the period and the geometry of the orbit. One could of course always use (4.1.22) and (4.1.21) (in the form which gives  $\phi(t)$ ) to determine the period and in the case of complex geometry there would be no choice.



**Example 4.1.** The possible orbits vary considerably depending upon the choice of the function  $h(r)$  and, indeed, even on the choice of the parameters in the chosen function. In the case of the simple choice  $h(r) = \lambda r$  we have the equation

$$\ddot{\mathbf{r}} + \frac{\lambda}{r} \mathbf{L} + \left( \lambda^2 r + \frac{k}{r^2} \right) \hat{\mathbf{r}} = 0 \quad (4.1.25)$$

which is the three-dimensional isotropic harmonic oscillator with the additional forces,  $\lambda \mathbf{L}/r$  and  $k \hat{\mathbf{r}}/r^2$ . (Alternatively we could consider this as the Kepler problem with two additional forces dependent on the parameter  $\lambda$ .) The formula (4.1.21) for the azimuthal angle takes the explicit form

$$\phi = - \int_{\theta_0}^{\theta} \left[ \sin^2 \theta (a^2 \sin^2 \theta (k + J \cos \theta)^2 - 1) \right]^{-\frac{1}{2}} d\theta \quad (4.1.26)$$

where  $a^2 = J^2 \lambda^{-2} L^{-6}$ . The performance of the quadrature in closed form does not appear to be possible and so we can only evaluate  $\phi(\theta)$  numerically. Given the comment just after (4.1.21) it is not surprising that the orbits close only for certain choices of the parameter  $k$ . As one increases the value of  $k$  from zero with values selected to close the orbit, there is an increase in the amount of rotational symmetry. The motions occur on a zone of a sphere.

**Example 4.2.** The equation of motion

$$\ddot{\mathbf{r}} = \frac{\lambda \dot{\mathbf{r}} \times \mathbf{r} + F(r) \mathbf{r}}{r^3} \quad (4.1.27)$$

was studied in some detail by Thompson [137]. He established conditions on  $F(r)$  in order that (4.1.27) possess an autonomous second first integral quadratic in the momentum independent of the energy. In summary his results were that

- (i)  $\lambda = 0$  and  $F(r) = k_1$ , a constant, which is the Kepler–Coulomb problem for  $k_1 \neq 0$  and the free particle otherwise,
- (ii)  $\lambda = 0$  and  $F(r) = k_1 r^3$  which is the three-dimensional isotropic harmonic oscillator or
- (iii)  $\lambda = 1$  and  $F(r) = k_1 - 1/r$ .

For case (iii) Thompson found the conserved vector

$$\mathbf{J} = \mathbf{L} \times \dot{\mathbf{r}} + \frac{\mathbf{L}}{r} - k_1 \mathbf{r} \quad (4.1.28)$$

which is the Laplace–Runge–Lenz vector for this problem. Case (iii) of equation (4.1.27) is, of course, a special case of (4.1.12) with  $h = -1/r$  and  $k = -k_1$ . It appears that the expression which Thompson uses for  $F(r)$  is incorrect and should be  $F(r) = k_1 + 1/r$ . The error seems to stem from the incorrect solution of an equation in his paper [137, (4.12)]. As a consequence the sign of the angular momentum term in (4.1.28) is incorrect. Thompson's Case (iii) is an apparently specific example of the more general equation

$$\ddot{\mathbf{r}} + \frac{\lambda \mathbf{L}}{r^3} + \left( \frac{k}{r^2} - \frac{\lambda^2}{r^3} \right) \hat{\mathbf{r}} = 0 \quad (4.1.29)$$

which has attracted a great deal of interest in connection with monopoles. In fact it is not a special case since (4.1.29) contains only one essential parameter which can be taken to be  $k$  by a simple rescaling of distance and time. This equation is treated in greater detail below. For  $\lambda = 1$  and  $F(r) = 0$  (4.1.27) becomes the equation of motion describing an electric charge interacting with a magnetic monopole fixed at the origin for which it has been shown that analogues of the angular momentum first integrals exist as well as some additional first integrals which are quadratic in the momentum, but which explicitly depend upon time (see Moreira [112]). The monopole system has no other time-independent scalar integrals apart from the conserved energy. It does seem that the vector approach, which we have been using, seems to offer results which are more general than the approaches adopted by other writers and with the benefit of far less mathematical complexity. It is the existence of the first integrals/invariants which determines the integrability of a given equation of motion, **not** the precise assumed form of those first integrals/invariants.

The equation of motion (4.1.29) describes the motion of an electric charge interacting with a magnetic monopole fixed at the origin with the additional centripetal forces  $k\hat{r}/r^2$  and  $-\lambda^2\hat{r}/r^3$ .

When  $h(r) = -\lambda/r$  and  $k = 0$ , from (4.1.16) we see that

$$r \cos \theta = \frac{L^2}{J}. \quad (4.1.30)$$

This means that  $\mathbf{r}$  has a constant projection onto  $\mathbf{J}$  of length  $L^2/J$ . From (4.1.21) we easily determine that

$$\phi = \operatorname{arcsec} \left( \left( 1 + \frac{L^2}{\lambda^2} \right)^{\frac{1}{2}} \sin \theta \right) \Bigg|_{\theta_0}^{\theta}. \quad (4.1.31)$$

From (4.1.30) it is evident that  $\theta$  is in the first quadrant and, since the argument of the arcsec is greater than or equal to one, we have that

$$\arcsin \frac{1}{(1 + L^2/\lambda^2)^{\frac{1}{2}}} \leq \theta \leq \frac{1}{2}\pi. \quad (4.1.32)$$

The geometry of this orbit and those to be described below are intimately connected with the orbits of the Kepler problem.

In the general case that  $h(r) = -\lambda/r$  and  $k \neq 0$  we have a truly three-dimensional motion for which there exists a Laplace–Runge–Lenz vector. The orbits may or may not be closed depending upon the relative values of  $k$  and  $J$ . The integration of (4.1.21) is not as elementary as in the case that  $k = 0$  and the expressions are somewhat more complicated. We find the three distinct cases

$J \neq k$  or  $-k$ :

$$\phi = \frac{1}{2} \arctan \left[ \frac{\frac{J^2 L^2}{\lambda^2} - k^2 - Jk - J \left( \frac{JL^2}{\lambda^2} + J + k \right) \cos \theta}{(J + k) \left( \frac{J^2 L^2}{\lambda^2} \sin^2 \theta - (k + J \cos \theta)^2 \right)^{\frac{1}{2}}} \right] \Bigg|_{\theta_0}^{\theta}$$

$$+ \frac{1}{2} \arctan \left[ \frac{\frac{J^2 L^2}{\lambda^2} + k^2 - Jk + J \left( \frac{JL^2}{\lambda^2} + J + k \right) \cos \theta}{(J - k) \left( \frac{J^2 L^2}{\lambda^2} \sin^2 \theta - (k + J \cos \theta)^2 \right)^{\frac{1}{2}}} \right] \Bigg|_{\theta_0}^{\theta}, \quad (4.1.33)$$

$J = k$ :

$$\phi = \frac{1}{2} \arctan \left[ \frac{\frac{L^2}{\lambda^2} - 2 - \left( \frac{L^2}{\lambda^2} + 2 \right) \cos \theta}{2 \left( \frac{L^2}{\lambda^2} \sin^2 \theta - (1 + \cos \theta)^2 \right)^{\frac{1}{2}}} \right] \Bigg|_{\theta_0}^{\theta}, \quad (4.1.34)$$

$J = -k$ :

$$\phi = -\frac{1}{2} \arctan \left[ \frac{\frac{L^2}{\lambda^2} - 2 - \left( \frac{L^2}{\lambda^2} + 2 \right) \cos \theta}{2 \left( \frac{L^2}{\lambda^2} \sin^2 \theta - (1 + \cos \theta)^2 \right)^{\frac{1}{2}}} \right] \Bigg|_{\theta_0}^{\theta}. \quad (4.1.35)$$

In the illustrative examples we have concentrated on the monopole because of its physical interest. We could have used any differentiable function  $h(r)$ . The point is that Laplace–Runge–Lenz vectors exist for truly three-dimensional motions for which only the magnitude of the angular momentum is conserved. To obtain the orbit we used the Laplace–Runge–Lenz vector in two separate vectorial combinations to obtain the two equations required to specify the orbit in three dimensions.

## 4.2 The classical MICZ problem

The motion of a spinless test particle in the field of a Dirac monopole plus Coulomb potential with an additional centrifugal potential has been called the MIC problem by Mladenov and Tsanov [108] and the MICZ problem by Cordani [25] after the studies of this problem by McIntosh and Cisneros [106] and also by Zwanziger [141]. The MICZ system is related to the problem of the asymptotic scattering of two self-dual monopoles by a canonical transformation [25] which gives the reduced Hamiltonian of a particle in an Euclidean Taub–NUT space. This in turn is related to the scattering of slowly moving Bogomol’nyi–Prasad–Sommerfield (BPS) monopoles [6, 7, 102].

The classical MICZ problem is described by the equation of motion

$$\ddot{\mathbf{r}} + \frac{\lambda}{r^3} \mathbf{L} + \left( \frac{\mu}{r^2} - \frac{\lambda^2}{r^3} \right) \hat{\mathbf{r}} = 0, \quad (4.2.1)$$

in which the mass is taken as unity,  $\lambda$  is the strength of the monopole and  $\mu$  is the strength of the Coulombic field. The Hamiltonian for (4.2.1) is

$$H = \frac{1}{2} \left( \mathbf{p} \cdot \mathbf{p} + \frac{\lambda^2}{r^2} \right) - \frac{\mu}{r}, \quad (4.2.2)$$

where  $\mathbf{p} = \dot{\mathbf{r}}$  is not the canonical momentum since, although  $[x_i, x_j]_{PB} = 0$  and  $[x_i, p_j]_{PB} = \delta_{ij}$  ( $x_i x_i = r^2$ ), the Poisson Bracket of the components of the mechanical momentum,  $\mathbf{p}$ , is

$$[p_i, p_j]_{PB} = \lambda \varepsilon_{ijk} \frac{x_k}{r^3}, \quad (4.2.3)$$

where  $\delta_{ij}$  and  $\varepsilon_{ijk}$  are the Kronecker delta and Kronecker epsilon respectively. The MICZ system possesses two conserved vectors which can be easily derived. We take the vector product of  $\mathbf{r}$  with (4.2.1) and integrate the result to obtain

$$\mathbf{P} = \mathbf{L} - \lambda \hat{\mathbf{r}}, \quad (4.2.4)$$

which is known as Poincare’s vector [123]. The scalar product of (4.2.4) with itself gives

$$P^2 = L^2 + \lambda^2, \quad (4.2.5)$$

which indicates that not only  $P$  is a constant but also  $L$ . The vector product of (4.2.1) with  $\mathbf{L}$  is easily integrated to give the second conserved vector

$$\mathbf{J} = \dot{\mathbf{r}} \times \mathbf{L} + \frac{\lambda}{r} \mathbf{L} - \mu \hat{\mathbf{r}}. \quad (4.2.6)$$

Hamilton’s vector for the system (4.2.1) is given by

$$\mathbf{K} = \mathbf{J} \times \mathbf{P} = (\dot{\mathbf{r}} \times \mathbf{P}) \times \mathbf{P} - \frac{\mu}{r} \mathbf{r} \times \mathbf{P}. \quad (4.2.7)$$

The integrals are related according to

$$J^2 = 2L^2H + \mu^2 \quad \text{and} \quad \mathbf{J} \cdot \mathbf{P} = \lambda\mu. \quad (4.2.8)$$

The presence of two conserved vectors for this three-dimensional motion gives rise to a rather special geometry.

If the velocity  $\dot{\mathbf{r}}$  is replaced by the momentum  $\mathbf{p}$  in  $\mathbf{P}$  and  $\mathbf{J}$ , the Poisson Bracket relations are given by [109]

$$[P_i, P_j]_{PB} = \varepsilon_{ijk} P_k, \quad [P_i, J_j]_{PB} = \varepsilon_{ijk} J_k, \quad [J_i, J_j]_{PB} = -2H \varepsilon_{ijk} P_k. \quad (4.2.9)$$

The Lie algebra of the first integrals under the operation of taking the Poisson Bracket is  $so(4)$ ,  $e(3)$  or  $so(3,1)$  depending upon whether  $H$  is negative, zero or positive. This is the same result as is found for the standard Kepler–Coulomb problem. We note, however, that the Lie algebra of the Lie point symmetries of the equation of motion, (4.2.1), is  $A_1 \oplus so(3)$  whereas for the Kepler–Coulomb problem it is  $A_2 \oplus so(3)$ , where  $A_1$  and  $A_2$  are the abelian subalgebra of dimension one and the nonabelian subalgebra of dimension two respectively [89].

The motion of a particle described by the equation of motion (4.2.1) is known to be a conic section. The only detailed study of the geometry of the orbit is to be found in the work of Gorringer [41] and this is a very detailed study indeed. McIntosh and Cisneros [106] state that they did not find any simple algebraic expressions for the parameters of the orbit in terms of constants of the motion. Gorringer comments “Using the first integrals of the MICZ system it has been possible to describe, using very elementary methods, the complete geometry of the orbits. Despite the comment of McIntosh and Cisneros [106] referred to earlier, the orbital quantities can be expressed fairly simply in terms of the conserved quantities and orbit parameters.” [41]. Unfortunately the elementariness and simplicity of the derivations is not matched by brevity and it would be out of place in

terms of the balance of this work to include it. Consequently we present only a brief summary of the general features.

The orbit of the MICZ problem is described by the intersection of a plane and a right circular cone. Consequently the orbit is a conic section. The projection of the orbit onto the azimuthal plane is a focus-centred conic section which corresponds naturally with the orbit of the standard Kepler problem. In the case of negative energy, Kepler's three laws of motion for the standard Kepler problem have a natural extension. They become<sup>4.1</sup>

- (i) The orbit is an ellipse. However, the origin is not coplanar with the ellipse.
- (ii) Equal areas are swept out on the surface of the cone in equal times.
- (iii) The relationship between the period of the motion and the semimajor axis length, *videlicet*

$$\frac{T^2}{R^3} = \frac{4\pi^2}{\mu}, \quad (4.2.10)$$

persists.

The orientation of the right circular cone is determined by Poincaré's vector,  $\mathbf{P}$ . The line of the Laplace–Runge–Lenz vector passes through the geometric centre of the orbit. Consequently the plane determined by  $\mathbf{P}$  and  $\mathbf{J}$  bisects the right circular cone and the intersection of this plane with the orbit gives the ends of the major axis. A linear combination of  $\mathbf{P}$  and  $\mathbf{J}$  naturally gives rise to two vectors one of which is parallel to the normal to the orbital plane and the other of which is parallel to the orbital plane. The second vector plays the role of a Laplace–Runge–Lenz vector in that it provides the orbital equation in a natural way using a suitable scalar product.

Let  $\pi - \alpha$  and  $\beta$  be the angles between  $\mathbf{P}$  and  $\mathbf{r}$  and  $\mathbf{P}$  and  $\mathbf{J}$  respectively. The scalar product of  $\mathbf{P}$ , (4.2.4), and  $\mathbf{r}$  and of  $\mathbf{P}$  and  $\mathbf{J}$ , (4.2.6), give

$$\cos \alpha = \frac{\lambda}{P} \quad \text{and} \quad \cos \beta = \frac{\lambda\mu}{PJ} \quad (4.2.11)$$

respectively. If the angle between  $\mathbf{J}$  and  $\mathbf{r}$  is  $\pi - \psi$ , the scalar product of  $\mathbf{J}$  and  $\mathbf{r}$  can be rearranged to give

$$r = \frac{L^2}{\mu - J \cos \psi} \quad (4.2.12)$$

which is one of the equations necessary to specify the orbit. The equation (4.2.12) describes a surface of revolution with the line of  $\mathbf{J}$  being the axis of symmetry and one focus of the surface is centred at the origin. The constancy of the angle between  $\mathbf{P}$  and  $\mathbf{r}$  implies that the particle moves on the surface of a cone with vertex at the origin. The curve of intersection of the cone and the surface of revolution, known as a conicoid, is the orbit of

---

<sup>4.1</sup>Bates [11] has approached this problem from a different direction using differential geometry. In the case of the Third Law he was able to obtain only the relationship between the period and the energy, which is the same as for the Kepler problem in the plane, since, without a knowledge of the geometry of the orbit, he was unable to relate this to the semimajor axis of the orbit.

the particle and also a plane conic section. By virtue of (4.2.4)  $\mathbf{P}$ ,  $\mathbf{L}$  and  $\mathbf{r}$  are always coplanar. However, the scalar triple product,

$$[\mathbf{P}, \mathbf{J}, \mathbf{r}] = L^2 r \dot{r}, \quad (4.2.13)$$

implies that  $\mathbf{P}$ ,  $\mathbf{J}$  and  $\mathbf{r}$  are coplanar only when  $\dot{r}$  is zero, *ie* at the extremities of the motion. We define the conserved vector

$$\mathbf{N} = \mathbf{P} - \frac{\lambda}{\mu} \mathbf{J}. \quad (4.2.14)$$

The scalar product of  $\mathbf{N}$  with  $\mathbf{r}$  gives

$$\mathbf{N} \cdot \mathbf{r} = -\frac{\lambda}{\mu} L^2, \quad (4.2.15)$$

which confirms that the orbit lies on a plane. However, the origin does not lie on this plane since the scalar product produces a nonzero value. The vector  $\mathbf{N}$  is parallel to the normal to the orbital plane. A vector  $\mathbf{S}$ , which is parallel to the orbital plane, is given by

$$\mathbf{S} = \mathbf{P} + \frac{\mu}{2\lambda H} \mathbf{J} \quad (4.2.16)$$

and is obtained by using the requirement that  $\mathbf{S} \cdot \mathbf{N} = 0$ . The angle,  $\xi$ , between  $\mathbf{P}$  and  $\mathbf{N}$  is found by taking the scalar product of these two vectors. This gives

$$\cos \xi = \sin \gamma = \frac{\mu L}{P(2H\lambda^2 + \mu^2)^{\frac{1}{2}}}, \quad (4.2.17)$$

where  $\gamma = \frac{1}{2}\pi - \xi$  is the angle between the orbital plane and the vector  $\mathbf{P}$  and can be used to find the cartesian equation of the plane. Similarly the angle between  $\mathbf{J}$  and  $\mathbf{N}$  is found from the scalar product between these two vectors and is given by

$$\cos \eta = \frac{-2HL\lambda}{(2HLL^2 + \mu^2)^{\frac{1}{2}}(2H\lambda^2 + \mu^2)^{\frac{1}{2}}}. \quad (4.2.18)$$

The projection of  $\mathbf{r}$  onto  $\mathbf{N}$ , which defines the orbital plane, is given, with the use of (4.2.15), by

$$\mathbf{r} \cdot \hat{\mathbf{N}} = r \cos \zeta = \frac{\lambda L}{(2H\lambda^2 + \mu^2)^{\frac{1}{2}}}, \quad (4.2.19)$$

where  $\zeta$  is the angle between  $\mathbf{r}$  and  $\mathbf{N}$ . If we fix  $\mathbf{P}$  to be along the  $-\mathbf{k}$  direction and rotate the orbit around  $\mathbf{P}$  so that the major axis of the orbit lies in the  $xz$ -plane, the unit normal is given by  $\hat{\mathbf{N}} = (\cos \gamma, 0, \sin \gamma)$  and correspondingly the equation of the orbital plane described by (4.2.19) can be expressed in terms of cartesian components as

$$x \cos \gamma + z \sin \gamma = \frac{\lambda P}{\mu} \sin \gamma, \quad (4.2.20)$$

where  $\gamma$  is the angle between  $\mathbf{P}$  and the plane of the orbit. The cartesian representation for the trajectory in spherical polar coordinates is given by

$$x = r \sin \alpha \cos \phi, \quad y = r \sin \alpha \sin \phi, \quad z = r \cos \alpha. \quad (4.2.21)$$

The substitution of these expressions into (4.2.20) gives

$$r = \frac{\lambda P \sin \gamma / \mu}{\cos \alpha \sin \gamma + \sin \alpha \cos \gamma \cos \phi} \quad (4.2.22)$$

which can be equated with the orbit equation, (4.2.12), to give

$$\psi = \arccos \left( \frac{\lambda^2 \mu - L^2 (2HP^2 + \mu^2)^{\frac{1}{2}} \cos \phi}{P^2 J} \right). \quad (4.2.23)$$

The general features of the motion are that the particle moves on a plane section of a cone with axis of symmetry along the line of  $\mathbf{P}$  and semivertex angle determined by  $\lambda/P$ . The ratio  $J/\mu$  determines whether the orbit is an ellipse, a parabola or an hyperbola. As  $\mathbf{r}$  moves over the cone,  $\mathbf{L}$ , which is coplanar with  $\mathbf{r}$  and  $\mathbf{P}$ , describes a circle on its own cone which has semivertex angle  $\frac{1}{2}\pi - \alpha$  with  $\mathbf{P}$  its axis of symmetry. The vector  $\mathbf{N}$  is parallel to the normal to the orbital plane and the vector  $\mathbf{S}$  is parallel to the orbital plane. From these basic considerations the geometric attributes of the orbit may be described in terms of the parameters in and the conserved quantities of the equation of motion (4.2.1) [41, pp. 150–206].

## 5 Symmetries and conserved vectors

### 5.1 Introduction

A second order ordinary differential equation,

$$\mathbf{E}(t, x, \dot{x}, \ddot{x}) = 0, \quad (5.1.1)$$

possesses a Lie symmetry,

$$G = \tau \partial_t + \eta_i \partial_{x_i}, \quad (5.1.2)$$

if

$$G^{[2]} \mathbf{E}(t, x, \dot{x}, \ddot{x})|_{\mathbf{E}=0} = 0, \quad (5.1.3)$$

where  $G$  has been twice extended to be able to operate on the first and second derivatives in the differential equation. The coefficient functions,  $\tau$  and  $\eta_i$ , can be anything provided they are twice differentiable.

If the coefficient functions depend upon  $t$  and  $x$  only, the symmetry is a Lie point symmetry. If in addition they depend upon  $\dot{x}$ , the symmetry is a generalised one. If the coefficient functions contained terms which are expressed as integrals that cannot be evaluated without a knowledge of the solution of the differential equation, the symmetry is a nonlocal symmetry.

Each of the three types of symmetry has a role to play in the discussion of the equations of motion which we have been considering. Traditionally the emphasis has been placed upon Lie point symmetries, but generalised symmetries – particularly in the context of Noether’s theorem – have been found in many instances to be necessary to be able to relate the existence of a first integral/invariant and the symmetry underlying its existence. More recently nonlocal symmetries have been found necessary to explain the existence of integrating factors for certain second order equations [84], to account for one of the integrable cases of the Hénon–Heiles problem [122] and to provide the symmetries necessary to construct the complete symmetry groups of some differential equations [75, 118, 91].

The application of (5.1.3) to the standard Kepler problem gives the following Lie point symmetries [125]

$$\begin{aligned} G_1 &= \partial_t, & G_3 &= z\partial_y - y\partial_z, \\ G_2 &= t\partial_t + \frac{2}{3}(x\partial_x + y\partial_y + z\partial_z), & G_4 &= z\partial_x - x\partial_z, \\ & & G_5 &= x\partial_y - y\partial_x \end{aligned} \tag{5.1.4}$$

which have the nonzero Lie Brackets

$$\begin{aligned} [G_1, G_2] &= G_1, \\ [G_3, G_4] &= G_5, & [G_4, G_5] &= G_3, & [G_5, G_3] &= G_4 \end{aligned} \tag{5.1.5}$$

from which it is evident that the Lie algebra is the direct sum  $A_2 \oplus so(3)$ . The three symmetries,  $G_3$ ,  $G_4$  and  $G_5$ , correspond to the invariance of (2.1.1) under rotation,  $G_1$  represents invariance under time translation and  $G_2$ , the symmetry often associated with the Laplace–Runge–Lenz vector, reflects invariance under the similarity transformation  $(t, \mathbf{r}) \longrightarrow (\bar{t}, \bar{\mathbf{r}} : t = \gamma\bar{t}, \mathbf{r} = \gamma^{\frac{2}{3}}\bar{\mathbf{r}})$ . One must realise that any of the first integrals can in fact be constructed from more than one of the symmetries in (5.1.4) [80] and so the association is not strictly accurate in the mathematical sense although it is in a geometric sense. The two-dimensional algebra,  $A_2$ , of  $G_1$  and  $G_2$ , representing invariance under time translation and rescaling, is a common combination in physical problems and has attracted attention in applications to a number of systems [15, 86, 85].

Since the Kepler problem and the time-dependent Kepler problem are related by a point transformation, they have the same algebraic structure [81].

By way of comparison the Lie point symmetries of the charge-monopole described by the equation of motion

$$\ddot{\mathbf{r}} = -\frac{\mu}{r^3}\mathbf{L}, \tag{5.1.6}$$

which were studied in some detail by Moreira *et al* [112], are

$$\begin{aligned} G_1 &= \partial_t, & G_4 &= z\partial_y - y\partial_z, \\ G_2 &= t\partial_t + \frac{1}{2}(x\partial_x + y\partial_y + z\partial_z), & G_5 &= z\partial_x - x\partial_z, \\ G_3 &= t(t\partial_t + x\partial_x + y\partial_y + z\partial_z), & G_6 &= x\partial_y - y\partial_x, \end{aligned} \tag{5.1.7}$$

and have the nonzero Lie Brackets

$$\begin{aligned} [G_1, G_2] &= G_1, & [G_2, G_3] &= G_3, & [G_3, G_1] &= -2G_2, \\ [G_4, G_5] &= G_6, & [G_5, G_6] &= G_4, & [G_6, G_4] &= G_5. \end{aligned} \tag{5.1.8}$$



The algebra is the direct sum of the subalgebras  $\{G_1, G_2, G_3\}$  and  $\{G_4, G_5, G_6\}$  and is  $sl(2, R) \oplus so(3)^{5.1}$ . Since the algebraic structure of (5.1.6) is more complex than that of the Kepler problem (2.1.1), as Thompson [137] has already noted, the charge-monopole problem is not related to the Kepler problem by a point transformation.

In this chapter we look at the symmetries of the various systems which have been considered above. In the first instance we look at the point symmetries. To explain the persistence of features such as the Laplace–Runge–Lenz vector for problems which do not have the appropriate number of point symmetries, in particular some rescaling symmetry such as  $G_2$  in (5.1.4) which is traditionally associated with the vector, we find it necessary to introduce a nonlocal symmetry. Following Krause [75] we look at the different systems from the point of view of their complete symmetry groups for which, in the case of these systems, it is also necessary to introduce nonlocal symmetries<sup>5.2</sup>. In the context of the complete symmetry group we find a remarkable unity in these diverse problems with a very interesting connection between the scalar simple harmonic oscillator and the Ermanno–Bernoulli constants [118].

## 5.2 The Lie point symmetries of $\ddot{\mathbf{r}} + f(r)\mathbf{L} + g(r)\mathbf{r} = 0$

We consider the more general equation

$$\ddot{\mathbf{r}} + f(r)\mathbf{L} + g(r)\mathbf{r} = 0 \quad (5.2.1)$$

since, when  $f = 0$  and  $g = \mu/r^3$ , we regain the Kepler problem and, when  $f = \mu/r^3$  and  $g = 0$ , we obtain the charge-monopole problem. More generally (5.2.1) contains the class of equations

$$\ddot{\mathbf{r}} + \frac{h'(r)}{r}\mathbf{L} + \left( h(r)h'(r) + \frac{k}{r^2} \right) \hat{\mathbf{r}} = 0 \quad (5.2.2)$$

which was shown in the previous chapter to possess a Laplace–Runge–Lenz vector.

The system (5.2.1) must be expressed in terms of components for the purposes of calculating its symmetries. Since the angular momentum has a regular structure in a cartesian basis, it is sensible to perform the calculation in this basis to provide a computational check through certain regularities in the determining equations. We write (5.2.1) as

$$\begin{aligned} \ddot{x} + f(y\dot{z} - z\dot{y}) + gx &= 0, \\ \ddot{y} + f(z\dot{x} - x\dot{z}) + gy &= 0, \\ \ddot{z} + f(x\dot{y} - y\dot{x}) + gz &= 0. \end{aligned} \quad (5.2.3)$$

Even with the aid of a symmetry computing package such as Alan Head's Program LIE [57, 131] one cannot do much more than obtain the determining equations using symbolic

<sup>5.1</sup>Moreira *et al* [112] use  $so(2, 1)$  rather than  $sl(2, R)$ . However, we believe that the  $sl(2, R)$  is more appropriate since the three symmetries with this algebra are common to all scalar ordinary differential equations of maximal (point) symmetry of order greater than one and the algebra is characteristic of Ermakov systems [32] of which (5.1.6) is an instance [50].

<sup>5.2</sup>Nonlocal symmetries are not always necessary to specify completely an ordinary differential equation, *eg* in the paper of Andriopoulos *et al* [5] it is shown that a linear second order equation has a complete symmetry group based on three point symmetries.

manipulation due to the unknown functions  $f$  and  $g$  in (5.2.3)<sup>5.3</sup>. There are forty-eight determining equations, twelve of which are duplicates. The solution of the thirty-six independent equations is a tedious business and we simply summarise the results. We find that two cases occur,  $f = 0$  and  $f \neq 0$ .

### 5.2.1 The case $f = 0$

In all cases we obtain the three symmetries associated with the algebra  $so(3)$  given by

$$\begin{aligned} G_1 &= z\partial_y - y\partial_z, \\ G_2 &= x\partial_z - z\partial_x, \\ G_3 &= y\partial_x - x\partial_y. \end{aligned} \tag{5.2.4}$$

Additional point symmetries depend upon the specific form of  $g(r)$ .

1.  $g(r)$  is unspecified: there is the single additional symmetry

$$G_4 = \partial_t. \tag{5.2.5}$$

The algebra is  $A_1 \oplus so(3)$ .

2.  $g = \mu r^\alpha$ : there are the two additional symmetries

$$\begin{aligned} G_4 &= \partial_t, \\ G_5 &= \alpha t \partial_t - 2r \partial_r. \end{aligned} \tag{5.2.6}$$

The algebra is  $A_2 \oplus so(3)$  as was reported for the Kepler problem, (2.1.1), by Prince and Eliezer [125] and Leach [80]. The Kepler problem is recovered when  $\alpha = -3$ .

3.  $g = \mu r^{-4}$ : the additional symmetries are

$$\begin{aligned} G_4 &= \partial_t, \\ G_5 &= 2t\partial_t + r\partial_r, \\ G_6 &= t(t\partial_t + r\partial_r). \end{aligned} \tag{5.2.7}$$

The algebra is  $sl(2, R) \oplus so(3)$  which is the same as that found for the charge-monopole problem, (5.1.6), by Moreira *et al* [112].

4.  $g = \mu r^{-4} - \epsilon$ : in this case the additional symmetries are

$$\begin{aligned} G_4 &= \partial_t, \\ G_5 &= e^{2t\epsilon^{1/2}} \left( \epsilon^{-1/2} \partial_t + r \partial_r \right), \\ G_6 &= e^{-2t\epsilon^{1/2}} \left( \epsilon^{-1/2} \partial_t - r \partial_r \right). \end{aligned} \tag{5.2.8}$$

---

<sup>5.3</sup>It would be interesting to see how far the very flexible and well-known interactive package of Nucci [115, 116] would be able to take the calculation. Unfortunately we did not have the package at the time.

The algebra is again  $sl(2, R) \oplus so(3)$ . In the case that  $\epsilon$  is a negative number, say  $-\omega^2$ , one may prefer to replace  $G_5$  and  $G_6$  with

$$\begin{aligned} X_5 &= -\frac{1}{2}i(G_5 + G_6) = \frac{\cos 2\omega t}{\omega}\partial_t + r \sin 2\omega t \partial_r, \\ X_6 &= \frac{1}{2}(G_5 - G_6) = -\frac{\sin 2\omega t}{\omega}\partial_t + r \cos 2\omega t \partial_r, \end{aligned} \quad (5.2.9)$$

although the former form is probably preferable for applications in quantum mechanics.

5.  $g = \epsilon$ : the equation of motion (5.2.1) is now that of a simple three-dimensional linear system, the free particle for  $\epsilon = 0$ , the isotropic simple harmonic oscillator for  $\epsilon > 0$  and the isotropic repulsor for  $\epsilon < 0$ . In all cases there are twenty-four symmetries and the algebra is  $sl(5, R)$  [80].

Despite the existence of a Laplace–Runge–Lenz vector in the classical Kepler problem we see that this does little to influence the construction of the symmetry algebra. Apart from the three-dimensional isotropic harmonic oscillator and the Newton–Cotes potential all power law central force problems possess the same number of symmetries and the same symmetry algebra, yet global (or even local) representations for a Laplace–Runge–Lenz vector are rarely possible. Although the Kepler problem is regarded as unusual because of the existence of the Laplace–Runge–Lenz vector, in terms of the point symmetry algebra of its equation of motion the Kepler problem is just one of an infinite number of problems.

### 5.2.2 The case $f \neq 0$

The generic result is that there exist the three generators of  $so(3)$  given in (5.2.4) and invariance under time translation given in (5.2.4). The exceptional cases are

1.  $f = \lambda r^{-3}$ ,  $g = \mu r^{-4}$ : in addition to the usual generators of  $so(3)$  there are the three additional symmetries given in (5.2.7). The algebra is again  $sl(2, R) \oplus so(3)$ . The equation of motion is

$$\ddot{\mathbf{r}} + \frac{\lambda}{r^3}\mathbf{L} + \frac{\mu}{r^4}\mathbf{r} = 0 \quad (5.2.10)$$

and the vector product of  $\mathbf{r}$  with this gives Poincare's vector

$$\mathbf{P} = \mathbf{L} - \lambda \hat{\mathbf{r}} \quad (5.2.11)$$

when integrated with respect to time. The vector product of the equation of motion, (5.2.10), with Poincare's vector,  $\mathbf{P}$ , gives

$$\frac{d(\mathbf{r} \times \mathbf{P})}{dt} + \left( \frac{L^4(\lambda^2 + \mu)}{|\mathbf{r} \times \mathbf{P}|^4} \right) (\mathbf{r} \times \mathbf{P}) = 0 \quad (5.2.12)$$

which is just the equation of motion in Case 3 above in terms of the new variable  $\mathbf{r} \times \mathbf{P}$ .

2.  $f = \lambda r^{-3}$ ,  $g = \mu r^{-4} - \epsilon$ : the additional symmetries are given by (5.2.8) in Case 4 above. The equation of motion

$$\ddot{\mathbf{r}} + \frac{\lambda}{r^3} \mathbf{L} + \left( \frac{\mu}{r^4} - \epsilon \right) \mathbf{r} = 0 \quad (5.2.13)$$

also has the Poincaré's vector (5.2.11) and, as in the previous case, the equation of motion in the variable  $\mathbf{r} \times \mathbf{P}$  is just that of Case 4 above.

### 5.2.3 Comment

The equation of motion

$$\ddot{\mathbf{r}} + f(r) \mathbf{L} + g(r) \mathbf{r} = 0 \quad (5.2.14)$$

has four Lie point symmetries for general  $f$  and  $g$  with the algebra  $A_1 \oplus so(3)$  with  $A_1$  representing invariance under time translation and  $so(3)$  the usual rotational invariance. In the special cases

$$\begin{aligned} \ddot{\mathbf{r}} + \mu r^\alpha \mathbf{r} &= 0, \\ \ddot{\mathbf{r}} + \frac{\mu}{r^4} \mathbf{r} &= 0, & \ddot{\mathbf{r}} + \left( \frac{\mu}{r^4} - \epsilon \right) \mathbf{r} &= 0, \\ \ddot{\mathbf{r}} + \epsilon \mathbf{r} &= 0, \\ \ddot{\mathbf{r}} + \frac{\lambda}{r^3} \mathbf{L} + \frac{\mu}{r^4} \mathbf{r} &= 0, & \ddot{\mathbf{r}} + \frac{\lambda}{r^3} \mathbf{L} + \left( \frac{\mu}{r^4} - \epsilon \right) \mathbf{r} &= 0 \end{aligned} \quad (5.2.15)$$

there is additional symmetry. The five-dimensional algebra of (5.2.15a) was found to be  $A_2 \oplus so(3)$  in which the  $A_2$  represents invariance under time translation and under the self-similar transformation

$$t = \gamma \bar{t}, \quad \mathbf{r} = \gamma^{-2/\alpha} \bar{\mathbf{r}}. \quad (5.2.16)$$

The interesting point about (5.2.15a) is that, as far as the Lie point symmetries are concerned, there is no distinction made between the Kepler problem and any other power law central force apart from the three-dimensional isotropic harmonic oscillator.

The equations of motion (5.2.15b) both possess the algebra  $sl(2, R) \oplus so(3)$  regardless of the values of the parameters. They may be regarded as direct extensions of the results of Moreira *et al* [112] for the algebra of the charge-monopole problem. Equation (5.2.15c) has the twenty-four-dimensional algebra  $sl(5, R)$  of the three-dimensional isotropic harmonic oscillator, free particle or repulsor depending whether  $\epsilon >, =, < 0$  respectively. Equations (5.2.15d) have the algebra  $sl(2, R) \oplus so(3)$  and also possess Poincaré vectors which can be used, through the vector product, to project and rotate them onto equations of motion equivalent to (5.2.15b) respectively in the variable  $\mathbf{r} \times \mathbf{P}$ .

The term  $\mu r r^{-4}$  can be interpreted as a centripetal force (Newton–Cotes) and the term  $\epsilon \mathbf{r}$  represents a three-dimensional isotropic harmonic oscillator, free particle or repulsor depending on whether  $\epsilon >, =, < 0$  respectively. The latter term does not affect the algebra nor the integrability of the equation. In both equations (5.2.15d)  $L$  is constant. The

motion remains on the surface of a cone due to the existence of the Poincaré vector. Another scalar integral

$$I = \frac{1}{2} \left( \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} - \frac{\mu}{r^2} - \epsilon r^2 \right) \quad (5.2.17)$$

also exists. Equation (5.2.15d2) does not belong to the class of the problems, (5.2.2), treated by Leach and Goringe [88]. However, when  $\mu = -\lambda^2$ , (5.2.15d1) belongs to the class and, in addition to the three integrals above, there is also the conserved vector

$$\begin{aligned} \mathbf{J} &= \dot{\mathbf{r}} \times \mathbf{L} + \frac{\lambda}{r} \mathbf{L} \\ &= \dot{\mathbf{r}} \times \mathbf{P} \end{aligned} \quad (5.2.18)$$

from which one of the equations specifying the orbit is easily obtained by taking the scalar product with  $\mathbf{r}$ .

The MICZ problem, (4.2.1), has the four-dimensional algebra,  $A_1 \oplus so(3)$ , which is surprising as it differs from the five-dimensional algebra,  $A_2 \oplus so(3)$ , of the Kepler problem and yet the Poincaré vector of the MICZ problem can be used to give the equation of motion for the Kepler problem in terms of the vector  $\mathbf{r} \times \mathbf{P}$ . We recall that, in the case of the Kepler problem, the self-similar symmetry is generally associated with Kepler's Third Law. However, in the MICZ problem, although an analogue of this conservation law is present, there is no suggestion of this in the Lie point symmetries of the problem. A similar observation can be made in respect of (5.2.13) in the case that  $\mu = -\lambda^2$ . Then the Poincaré vector can be used to express the equation as that of a three-dimensional isotropic harmonic oscillator in the variable  $\mathbf{r} \times \mathbf{P}$ . Apart from the change in the number of symmetries from six to twenty-four, which in itself is sufficient to cause one to pause for wonder, the self-similar symmetry of the isotropic harmonic oscillator signals a Kepler-type Third Law, in this case the oscillator's isochronism. There is no suggestion of this in (5.2.13). The obvious geometric connection between these motions on the surface of a cone and the corresponding planar motions is not reflected in the Lie point symmetries.

Finally we note that the equation (5.2.2) possesses a Laplace–Runge–Lenz vector for general functions  $h(r)$ . Can it be that there is no connection between the existence of this extraordinary vector and the underlying symmetry of the equation of motion apart from its invariance under time translation? This seems to be most unlikely.

### 5.3 The equation $\ddot{\mathbf{r}} + f\dot{\mathbf{r}} + \mu g\mathbf{r} = \mathbf{0}$

We turn now to the equation of motion we considered in Chapter 2 for which we found the conserved quantities

$$A = L (gr^3)^{-\frac{1}{2}}, \quad (5.3.1)$$

$$\mathbf{K} = \frac{\dot{\mathbf{r}}}{L} - \frac{\mu}{A^2} \hat{\boldsymbol{\theta}}, \quad (5.3.2)$$

$$\mathbf{J} = \frac{\dot{\mathbf{r}} \times \hat{\mathbf{L}}}{L} + \frac{\mu}{A^2} \hat{\mathbf{r}} \quad (5.3.3)$$

in the case that

$$f = -\frac{1}{2} \left( \frac{\dot{g}}{g} + 3\frac{\dot{r}}{r} \right), \quad (5.3.4)$$

where  $\dot{g}$  represents the total time derivative of  $g$ . We recall that the direction of the angular momentum vector,  $\hat{\mathbf{L}}$ , is a constant and we can work with plane polar coordinates,  $(r, \theta)$ . The equation of the orbit is obtained from the scalar product of (5.3.3) and  $\mathbf{r}$  and is

$$r = \frac{1}{\mu A^{-2} + J \cos \theta} \quad (5.3.5)$$

which is a conic section. Gorrington and Leach [48] reported just the four Lie point symmetries, (5.2.4) and (5.2.5), in the case that  $g$  was a function of  $r$ , but not a power law, only (and by consequence  $f$  of  $r$  and  $\dot{r}$  only) and so we are in a situation which is similar to our considerations of the previous section. In the case that  $g$  is a power law potential there is also a rescaling symmetry [48].

The Laplace–Runge–Lenz vector is invariant under time translation, but there is no other Lie point symmetry as in the case of the Kepler problem. Since we do not have a Lie point symmetry apart from  $\partial_t$  for the vector  $\mathbf{J}$ , we look for a symmetry of the form [121]

$$G = \tau \partial_t + \eta \partial_r \quad (5.3.6)$$

with no restrictions made on the coefficient functions. Note that we are not looking for a general symmetry which would include a term for  $\partial_\theta$ . We are looking for some generalisation of the similarity symmetry.

We require that (5.3.3) be invariant under the action of the first extension of the symmetry, *ie*  $G^{[1]}\mathbf{J} = 0$ . The first extension of  $G$  is given by

$$G^{[1]} = \tau \partial_t + \eta \partial_r + 0 \partial_\theta + (\dot{\eta} - \dot{r} \dot{\tau}) \partial_{\dot{r}} - \dot{\theta} \dot{\tau} \partial_{\dot{\theta}}. \quad (5.3.7)$$

(As a practical note we include the  $\partial_\theta$  term, even though its coefficient is zero, as a reminder that there is a  $\partial_{\dot{\theta}}$  term as well.) Since

$$\hat{\mathbf{r}} = \cos \theta \hat{\mathbf{i}} + \sin \theta \hat{\mathbf{j}} \quad \text{and} \quad \hat{\boldsymbol{\theta}} = -\sin \theta \hat{\mathbf{i}} + \cos \theta \hat{\mathbf{j}}, \quad (5.3.8)$$

we have

$$G^{[1]}\hat{\mathbf{r}} = 0 \quad \text{and} \quad G^{[1]}\hat{\boldsymbol{\theta}} = 0 \quad (5.3.9)$$

when  $\hat{\mathbf{r}}$  and  $\hat{\boldsymbol{\theta}}$  appear as geometrical (coordinate frame) vectors. When  $\hat{\mathbf{r}}$  and  $\hat{\boldsymbol{\theta}}$  appear as a consequence of integration of the equation of motion (as occurs in [48]), the calculation is generally more delicate [60]. Applying the first extension of  $G$  to  $\mathbf{J}$  and separating by the coefficients of the independent vectors  $\hat{\mathbf{r}}$  and  $\hat{\boldsymbol{\theta}}$  we obtain the following equations for the coefficient functions

$$\dot{\tau} + \frac{1}{2} \frac{g'}{g} \eta = 0, \quad (5.3.10)$$

$$\dot{\eta} L - \eta r \dot{\theta} = 0, \quad (5.3.11)$$

where  $g' = dg/dr$ .

Since  $\hat{\mathbf{L}}$  is conserved, we can replace  $L$  by  $r^2 \dot{\theta}$  in the solution of these equations. Hence (5.3.11) gives

$$\eta = Cr, \quad (5.3.12)$$

where  $C$  is a constant of integration. We substitute this into (5.3.10) and integrate the equation to obtain

$$\tau = -\frac{1}{2}C \int \frac{g'r}{g} dt. \quad (5.3.13)$$

We have, therefore, that

$$G = -\frac{1}{2} \left[ \int \frac{g'r}{g} dt \right] \partial_t + r \partial_r \quad (5.3.14)$$

is a symmetry of the (generalised) Laplace–Runge–Lenz vector. Note that for  $g = r^\alpha$  this is a local symmetry and corresponds to the result of Gorringer and Leach [48] quoted above.

In the case of elliptical orbits the similarity symmetry of the Kepler problem is closely connected with Kepler's Third Law. In this case we have the period  $T$  given by the quadrature

$$T = 2 \int_0^\pi \frac{d\theta}{\dot{\theta}} = 2 \int_0^\pi \left( \frac{r}{g} \right)^{\frac{1}{2}} \frac{d\theta}{A}. \quad (5.3.15)$$

The integral is evaluated by substituting for  $r$  from the orbit equation, (5.3.5). In the case that  $g$  is a power law in  $r$  Gorringer and Leach [48] were able to evaluate the integral in (5.3.15) in terms of Legendre functions. One would not expect to be able to evaluate the quadrature in closed form for general  $g$  although this would be the case were  $g$  one of some particular polynomials in  $r$ .

In terms of cartesian unit vectors in the plane of the orbit we may rewrite (5.3.3), using (5.3.1), as

$$A^2 \mathbf{J} = \hat{i} \left[ \left( \frac{\dot{\theta}^2}{g} - \mu \right) \cos \theta + \frac{\dot{r}\dot{\theta}}{gr} \sin \theta \right] + \hat{j} \left[ \left( \frac{\dot{\theta}^2}{g} - \mu \right) \sin \theta - \frac{\dot{r}\dot{\theta}}{gr} \cos \theta \right]. \quad (5.3.16)$$

(Note that in equation (36) of [121] there is an error which has been corrected in the formula given above.) The two independent components of the conserved vector, the Ermanno–Bernoulli constants, can be written in the compact form

$$J_\pm = \exp[\pm i\theta] \left( \frac{\dot{\theta}^2}{g} - \mu \pm \frac{\dot{r}\dot{\theta}}{igr} \right). \quad (5.3.17)$$

The components in (5.3.17) have a symmetry of the form (5.3.6) provided

$$\dot{\eta} - \eta \left( \frac{\dot{r}g'}{g} + \frac{\dot{r}}{r} \pm \frac{i\dot{\theta}rg'}{g} \right) = 2\dot{\tau} (\dot{r} \pm ir\dot{\theta}). \quad (5.3.18)$$

If in (5.3.18) we set  $\eta = 0$  and  $\eta = r$ , we recover the  $\partial_t$  symmetry and (5.3.14) respectively. When we put  $\tau = 0$ , (5.3.18) gives

$$\eta = gr \exp \left[ \pm i \int \frac{rg'}{g} d\theta \right], \quad (5.3.19)$$

where we recall that  $r$  and hence  $g(r)$  are functions of  $\theta$  and hence must remain in the integral, so that the components have the extra symmetry

$$G_{3\pm} = gr \exp \left[ \pm i \int \frac{rg'}{g} d\theta \right] \partial_r. \quad (5.3.20)$$

For a given  $g(r)$  the integral in (5.3.19) is evaluated using the expression for  $r$  in (5.3.5). For the Kepler problem  $g = r^{-3}$  and (5.3.20) reduces to

$$G_{3\pm} = \frac{1}{r^2} \exp [\pm 3i\theta] \partial_r \quad (5.3.21)$$

which does not seem to have been reported prior to the paper of Pillay *et al* [121]. We observe that the 3,2 combination of the similarity symmetry for the Kepler problem recurs in this symmetry. We note that this symmetry is a symmetry of the first integral and not a symmetry of the differential equation.

In the case of the Kepler problem the Ermanno–Bernoulli constants have the three Lie point symmetries

$$G_1 = \partial_t, \quad G_2 = t\partial_t + \frac{2}{3}r\partial_r, \quad G_{3\pm} = \frac{1}{r^2} \exp [\pm 3i\theta] \partial_r \quad (5.3.22)$$

with the Lie Brackets

$$[G_1, G_2] = G_1, \quad [G_1, G_{3\pm}] = 0 \quad \text{and} \quad [G_2, G_{3\pm}] = -2G_{3\pm} \quad (5.3.23)$$

which is a representation of the algebra  $A_{3,5}^{1/2}$ .

It would appear that we have now come to the beginning of a resolution of the quandary which faced us at the end of the previous section. We simply cannot expect these more complicated systems to have a Lie point symmetry naturally associated with a given first integral/invariant. In the case of simple systems, to which class most problems arising in a natural physical context belong, one has been well-served in the past by point symmetries and, perhaps consequentially, led to regard point symmetries as the norm. In cases for which the analysis of the simple system can be extended to more complicated, albeit artificial, systems we have the opportunity to extract the essence of the system and not just the simplicity of the original system.

## 5.4 Complete symmetry groups

In 1994 Krause [75] introduced a new concept into the study of the symmetries of ordinary differential equations. Krause adopted the terminology ‘complete symmetry group’, which in the past had been used to describe the group corresponding to the set of all Lie point symmetries of a differential equation, defining it by the addition of two properties to the definition of a Lie symmetry group. These were that the manifold of solutions is a homogeneous space of the group and that the group is specific to the system, *ie* no other system admits it<sup>5.4</sup>. To take a simple example the free particle in one dimension described by the equation of motion

$$\ddot{q} = 0 \quad (5.4.1)$$

---

<sup>5.4</sup>There is an exception to this in that there may be an arbitrary parameter present in the equation which can be set at any permissible value by means of a transformation, such as rescaling or translation.



has eight Lie point symmetries of which three are

$$G_1 = \partial_t, \quad G_2 = \partial_q, \quad G_3 = t\partial_t + q\partial_q \quad (5.4.2)$$

with the Lie Brackets

$$[G_1, G_2] = 0, \quad [G_1, G_3] = G_1, \quad [G_2, G_3] = G_2 \quad (5.4.3)$$

which is a representation of the algebra  $A_{3,3}$ , the semidirect product of dilations and translations,  $D \oplus_s T_2$ . If we consider the general scalar second order equation

$$\ddot{q} = f(t, q, \dot{q}), \quad (5.4.4)$$

the action of  $G_1$  is to remove the dependence of  $f$  on  $t$  and that of  $G_2$  is to remove the dependence on  $q$ . The second extension of  $G_3$  is

$$G^{[2]} = t\partial_t + q\partial_q + 0\partial_{\dot{q}} - \ddot{q}\partial_{\ddot{q}} \quad (5.4.5)$$

and this requires that the  $f(\dot{q})$  remaining be zero. Thus the algebra  $A_{3,3}$  is a representation of the complete symmetry group of (5.4.1) [5].

The vehicle which Krause used to illustrate the purpose of the introduction of the complete symmetry group was the Kepler problem. If we consider the Lie point symmetries of the Kepler problem (in two dimensions only for the sake of brevity), *videlicet*

$$G_1 = \partial_t, \quad G_2 = \partial_\theta, \quad G_3 = t\partial_t + \frac{2}{3}r\partial_r, \quad (5.4.6)$$

and apply them as in the example above, we find that the two components of the equation of motion are required to have the forms

$$\begin{aligned} \ddot{r} &= \frac{1}{r^2} f(r\dot{r}^2, r^3\dot{\theta}^2), \\ \ddot{\theta} &= \frac{1}{r^3} g(r\dot{r}^2, r^3\dot{\theta}^2), \end{aligned} \quad (5.4.7)$$

where  $f$  and  $g$  are arbitrary functions of their arguments. Evidently the three Lie point symmetries are not adequate to specify completely the Kepler problem. To obtain the complete symmetry group of the Kepler problem Krause introduced a nonlocal symmetry defined by

$$Y = \left[ \int \tau(t, x_1, \dots, x_N) dt \right] \partial_t + \eta_i(t, x_i, \dots, x_N) \partial_{x_i}. \quad (5.4.8)$$

This type of nonlocal symmetry is a very special type since only the coefficient function of  $\partial_t$  contains a nonlocal term. This type of nonlocal symmetry is potentially very useful to treat autonomous systems for then only the integrand and not the integral intrudes into the calculation. In his treatment of the three-dimensional problem Krause found the standard five Lie point symmetries and three additional nonlocal symmetries which were needed to specify completely the Kepler equation, up to the value of the gravitational constant  $\mu$ . The three additional symmetries are

$$\begin{aligned} Y_1 &= 2 \left( \int x_1 dt \right) \partial_t + x_1 r \partial_r, & Y_2 &= 2 \left( \int x_2 dt \right) \partial_t + x_2 r \partial_r, \\ Y_3 &= 2 \left( \int x_3 dt \right) \partial_t + x_3 r \partial_r \end{aligned} \quad (5.4.9)$$

in which  $r^2 = x_i x_i$ .

Krause did not give the algebra of these symmetries. He also remarked that it was not possible to obtain the symmetries of the complete symmetry group of the Kepler problem by means of Lie point methods. In 1996 Nucci [117] demonstrated that it was indeed possible to obtain the symmetries by means of Lie point methods.

### 5.5 The method of reduction of order

The technique which Nucci used to show that one could obtain the symmetries of the complete symmetry group of the Kepler problem by means of Lie point methods is one which appears to be generally useful and we summarise it here. For the purposes of our discussion we consider a system of  $N$  second order ordinary differential equations,

$$\ddot{x}_i = f_i(x, \dot{x}_i), \quad i = 1, N, \quad (5.5.1)$$

in which  $t$  is the independent variable and  $x_i$ ,  $i = 1, N$  are the  $N$  dependent variables. These equations may be considered as equations from Newtonian mechanics, which was Krause's approach and which is appropriate to the context of this paper, but there is no necessity for that to be the case. There is no necessity for the dependent variables to represent cartesian coordinates, nor is there a need for the system to be of the second order. There is no requirement that the system be autonomous. In the case of a nonautonomous system we can apply the standard procedure of introducing a new variable  $x_{N+1} = t$  and an additional first-order equation of  $\dot{x}_{N+1} = 1$  so that the system becomes formally autonomous. In our discussion we confine our attention to autonomous systems.

The first step in the method of reduction of order is to write the system (5.5.1) as a  $2N$ -dimensional first order system by means of the introduction of the variables

$$\begin{aligned} w_1 &= x_1, & w_{N+1} &= \dot{x}_1, \\ w_2 &= x_2, & w_{N+2} &= \dot{x}_2, \\ &\vdots & &\vdots \\ w_{N-1} &= x_{N-1}, & w_{2N-1} &= \dot{x}_{N-1}, \\ w_N &= x_N, & w_{2N} &= \dot{x}_N \end{aligned} \quad (5.5.2)$$

so that the system (5.5.1) becomes

$$\dot{w}_i = g_i(w), \quad i = 1, 2N, \quad (5.5.3)$$

where  $g_i = w_{N+i}$  for  $i = 1, N$  and  $g_i = f_i$  for  $i = N + 1, 2N$ . In this first step of the reduction of the original system, (5.5.1), we have simply followed the conventional method used to reduce a higher order system to a first order system. Any optimisation is performed in the further selection of the final variables. This selection may be motivated by the existence of a known first integral, such as angular momentum, or some specific symmetry in the original system, (5.5.1).

We choose one of the variables  $w_i$  to be a new independent variable  $y$ . For the purpose of the present development we make the identification  $w_N = y$ . In terms of the new independent variable we obtain the system of  $2N - 1$  first order equations

$$\frac{dw_i}{dy} = \frac{g_i}{g_N} = \frac{g_i}{w_{2N}}, \quad i = 1, \dots, N - 1, N + 1, \dots, 2N. \quad (5.5.4)$$

We do not attempt to calculate the Lie point symmetries of the system, (5.5.4), because the Lie point symmetries of a first order system are generalised symmetries and one has to impose some *Ansatz* on the form of the symmetry. The choice of *Ansatz* tends to be more of what appeals to the mind of the devisor rather than of some intrinsic property of the system. Rather we select  $n \leq N - 1$  of the variables to be new dependent variables and rewrite the system (5.5.4) as a system of  $n$  second order equations plus  $2(N - n) - 1$  first order equations<sup>5.5</sup>. The selection of the new dependent variables, denoted by  $u_1, u_2, \dots$  to distinguish them from the intermediate variables  $w_1, w_2, \dots$ , is dictated by a number of considerations. The first and foremost is that we must be able to eliminate the unwanted variables from the system (5.5.4). After this condition has been satisfied, we may look to seek variables which reflect some symmetry system, for example an ignorable coordinate such as the azimuthal angle in a central force problem.

After the symmetries of the reduced system have been calculated, they can be translated back to symmetries of the original system. In the reduction of order we are hoping that nonlocal symmetries become local, indeed point, symmetries of the reduced system and that any existing point symmetries remain as point symmetries of the reduced system. This means that the nonlocal symmetries are of Type II [1, 2, 3, 4]. As the computation of point symmetries is the easiest of all computations of symmetries to perform, the existence of Type II symmetries is desirable. Unfortunately there is also the possibility of the occurrence of Type I hidden symmetries, *ie* Lie point symmetries of the original system which become nonlocal on the reduction of order. If the symmetry associated with the reduction of order is, say,  $G_1$ , and a second symmetry  $G_2$  has the Lie Bracket  $[G_1, G_2] \neq \lambda G_1$ , where the constant  $\lambda$  may be zero,  $G_2$  will become an exponential nonlocal symmetry of the reduced system. This is the risk which one must accept in undertaking the search for symmetry by means of a change of order of the system, whether that change be a reduction of order or an increase of order. Even with simple systems the comings and goings of point symmetries with an increase or decrease in the order is quite extraordinary [90].

## 5.6 The symmetries of the reduced Kepler problem

The two components of the equation of motion for the Kepler problem in two dimensions are

$$\ddot{r} - r\dot{\theta}^2 = -\frac{\mu}{r^2}, \quad (5.6.1)$$

$$r\ddot{\theta} + 2\dot{r}\dot{\theta} = 0. \quad (5.6.2)$$

We introduce the new variables and their time derivatives

$$\begin{aligned} w_1 &= r, & \dot{w}_1 &= w_3, \\ w_2 &= \theta, & \dot{w}_2 &= w_4, \\ w_3 &= \dot{r}, & \dot{w}_3 &= w_1 w_4^2 - \frac{\mu}{w_1^2}, \\ w_4 &= \dot{\theta}, & \dot{w}_4 &= -\frac{2w_3 w_4}{w_1}. \end{aligned} \quad (5.6.3)$$

---

<sup>5.5</sup>Again there is no real necessity for the final equations to be a mixture of second and first order equations. One could include third or fourth order equations. However, from a practical point of view this may not be so productive.

Since in the original system  $\theta$  is an ignorable coordinate, we select  $w_2$  to be the new independent variable  $y$ . The right side of (5.6.3) leads to the reduced system

$$\frac{dw_1}{dy} = \frac{w_3}{w_4}, \quad (5.6.4)$$

$$\frac{dw_3}{dy} = w_1w_4 - \frac{\mu}{w_1^2w_4}, \quad (5.6.5)$$

$$\frac{dw_4}{dy} = -\frac{2w_3}{w_1}. \quad (5.6.6)$$

We proceed now to change the system of three first order equations into a system of one second order and one first order. From (5.6.4) we have  $w_3 = w_4w_1'$ , where the prime denotes differentiation with respect to the new independent variable,  $y$ , and replace (5.6.6) by

$$w_4' = -\frac{2w_4w_1'}{w_1}. \quad (5.6.7)$$

In (5.6.5) we make the same replacement to obtain the second order equation

$$w_4w_1'' - \frac{w_4w_1'^2}{w_1} = w_1w_4 - \frac{\mu}{w_1^2w_4}. \quad (5.6.8)$$

In the elimination of  $w_3$  we have not precisely decided that the variables  $u_1$  and  $u_2$  are to be  $w_1$  and  $w_4$ . We observe that the integrating factor  $w_1^2$  makes (5.6.7) exact as

$$(w_1^2w_4)' = 0 \quad (5.6.9)$$

and this prompts the choice of one of the new variables to be  $u_2 = w_1^2w_4$ . We introduce this new variable into (5.6.8) to obtain

$$w_1'' = 2\frac{w_1'^2}{w_1} + w_1 - \frac{\mu w_1^2}{u_2}. \quad (5.6.10)$$

We rearrange (5.6.10) somewhat as

$$u_2^2 \left( \frac{w_1''}{w_1^2} - \frac{2w_1'^2}{w_1^3} \right) = \frac{u_2^2}{w_1} - \mu \quad (5.6.11)$$

and immediately recognise a convenient new variable  $u_1 = \mu - u_2^2/w_1$  so that the reduced system becomes

$$\begin{aligned} u_1'' + u_1 &= 0, \\ u_2' &= 0 \end{aligned} \quad (5.6.12)$$

which is an attractively simple system. When we calculate the Lie point symmetries of (5.6.12), we obtain nine point symmetries. Some of these contain sine and cosine terms.

We combine them to give the symmetries in terms of the exponential of an imaginary variable. The symmetries are

$$\begin{aligned}
\Gamma_1 &= u_2 \partial_{u_2}, \\
\Gamma_2 &= \partial_y, \\
\Gamma_3 &= u_1 \partial_{u_1}, \\
\Gamma_{4\pm} &= e^{\pm iy} \partial_{u_1}, \\
\Gamma_{6\pm} &= e^{\pm 2iy} [\partial_y \pm i u_1 \partial_{u_1}], \\
\Gamma_{8\pm} &= e^{\pm iy} [u_1 \partial_y \pm i u_1^2 \partial_{u_1}].
\end{aligned} \tag{5.6.13}$$

In terms of the original variables we obtain

$$\begin{aligned}
\Gamma_1 &= 3t \partial_t + 2r \partial_r, \\
\Gamma_2 &= \partial_\theta, \\
\Gamma_3 &= 2 \left[ \mu \int r dt - L^2 t \right] \partial_t + r (\mu r - L^2) \partial_r, \\
\Gamma_{4\pm} &= 2 \left[ \int r e^{\pm i\theta} dt \right] \partial_t + r^2 e^{\pm i\theta} \partial_r, \\
\Gamma_{6\pm} &= 2 \left[ \int (\mu r + 3L^2) e^{\pm 2i\theta} dt \right] \partial_t + r (\mu r + 3L^2) e^{\pm 2i\theta} \partial_r + L^2 e^{\pm 2i\theta} \partial_\theta, \\
\Gamma_{8\pm} &= 2 \left[ \int \left\{ 2\dot{r} L^3 \pm ir (\mu - r^3 \dot{\theta}^2) (\mu + r^3 \dot{\theta}^2) \right\} e^{\pm i\theta} dt \right] \partial_t \\
&\quad + r \left[ 2\dot{r} L^3 \pm ir (\mu - r^3 \dot{\theta}^2) (\mu + r^3 \dot{\theta}^2) \right] \partial_r + L^2 (\mu - r^3 \dot{\theta}^2) e^{\pm i\theta} \partial_\theta
\end{aligned} \tag{5.6.14}$$

in which a factor of  $L^2$  has generally been included (the exceptions are  $\Gamma_2$  and  $\Gamma_1$ ; in the latter a factor of  $L$  has been included) to make the expressions look simpler.

In (5.6.14) we recognise in  $\Gamma_1$  and  $\Gamma_2$  the two point symmetries which were not affected by the reduction of order based on the symmetry  $\partial_t$ . The two symmetries  $\Gamma_{4\pm}$  are the two nonlocal symmetries required to be introduced to obtain the complete symmetry group for the two-dimensional Kepler problem. The remaining four symmetries are additional to the number required completely to specify the differential equation. It is curious that they were not reported by Krause [75] since they are of the same structure as the nonlocal symmetry which he introduced.

If we examine the reduced system (5.6.12), we find that the symmetries  $\Gamma_1$ ,  $\Gamma_2$  and  $\Gamma_{4\pm}$  are sufficient to specify it completely from the general system

$$\begin{aligned}
u_1'' &= f(y, u_1, u_2, u_1'), \\
u_2' &= g(y, u_1, u_2)
\end{aligned} \tag{5.6.15}$$

and so we should not be surprised that in combination with the symmetry  $\partial_t$  these symmetries completely specify the two-dimensional Kepler problem. In the case of the three-dimensional Kepler problem a third equation, of the first order, based on the constancy of the  $z$ -component of the angular momentum is added to (5.6.12). Since there are at least three representations of the algebra for the complete symmetry group of a second order linear equation [5], one would be surprised if this representation be unique.

We have seen in detail for the two-dimensional Kepler problem that one of the trivial equations of the reduced system was that of the conservation of angular momentum and have indicated for the three-dimensional Kepler problem that there will be two trivial equations, one for the conservation of the total angular momentum and the other for the conservation of the  $z$ -component of the angular momentum. The nontrivial equation is (5.6.12a), the second order equation for the simple harmonic oscillator. In defining the new variable  $u_1$  we could, although in fact we did not, have taken the standard transformation of  $u = 1/r$  which is used to convert the radial equation of the Kepler problem to an inhomogeneous oscillator equation [140] and then introduced the translation to remove the inhomogeneous term. The equation (5.6.12a) has two linear first integrals. In the notation we are using here they are

$$J_{\pm} = e^{\pm iy} (u_1 \pm iu_1') \quad (5.6.16)$$

and, if one translates these back into the original coordinates of the two-dimensional Kepler problem, one finds that these two invariants for the simple harmonic oscillator are just the Ermanno–Bernoulli constants, *ie* combinations of the two components of the Laplace–Runge–Lenz vector. This immediately raises the question whether a similar phenomenon occurs for any other other problems which possess Laplace–Runge–Lenz vectors.

## 5.7 The reduced system for some problems possessing a conserved vector of Laplace–Runge–Lenz type

### 5.7.1 The Kepler problem with drag

We recall the model proposed by Danby [27] for the motion of a satellite at low altitude subject to a resistive force due to the Earth's atmosphere described by the equation of motion

$$\ddot{\mathbf{r}} + \frac{\alpha \dot{\mathbf{r}}}{r^2} + \frac{\mu \mathbf{r}}{r^3} = 0, \quad (5.7.1)$$

where  $\alpha$  and  $\mu$  are constants. Since the direction of the angular momentum is a constant, we may use plane polar coordinates,  $(r, \theta)$ . The two components of the equation of motion are

$$\ddot{r} - r\dot{\theta}^2 + \frac{\alpha \dot{r}}{r^2} + \frac{\mu}{r^2} = 0, \quad (5.7.2)$$

$$r\ddot{\theta} + 2\dot{r}\dot{\theta} + \frac{\alpha \dot{\theta}}{r} = 0. \quad (5.7.3)$$

We introduce the same new variables as for the Kepler problem, rewrite the system (5.7.2) and (5.7.3) as the equivalent system of four first order equations and again select  $w_2$  to be the new independent variable  $y$ . We obtain the three first order equations

$$w_1' = \frac{w_3}{w_4}, \quad (5.7.4)$$

$$w_3' = w_1 w_4 - \frac{\alpha w_3}{w_1^2 w_4} - \frac{\mu}{w_1^2 w_4}, \quad (5.7.5)$$

$$w_4' = -\frac{2w_3}{w_1} - \frac{\alpha w_4}{w_1^2}, \quad (5.7.6)$$

where the prime denotes differentiation with respect to the new independent variable,  $y$ . The same convention is used below.

The choice of (5.7.4) to eliminate  $w_3$  is obvious. We obtain

$$w_1'' w_4 + w_1' w_4' = w_1 w_4 - \frac{\alpha w_1'}{w_1^2} - \frac{\mu}{w_1^2 w_4}, \quad (5.7.7)$$

$$w_4' = -\frac{2w_1' w_4}{w_1} - \frac{\alpha}{w_1^2}. \quad (5.7.8)$$

We can easily manipulate (5.7.8) to obtain

$$(w_1^2 w_4)' + \alpha = 0 \quad \Leftrightarrow \quad w_1^2 w_4 = -(\alpha y + \beta), \quad (5.7.9)$$

which indicates that the angular momentum is not conserved, and the choice of one of the new variables is then

$$u_2 = w_1^2 w_4 + \alpha y + \beta. \quad (5.7.10)$$

In (5.7.7) we use both forms of (5.7.9) to eliminate  $w_4$  and  $w_4'$  to obtain

$$\left(\frac{1}{w_1}\right)'' + \frac{1}{w_1} = \frac{\mu}{(\alpha y + \beta)^2} \quad (5.7.11)$$

and with the choice of the second new variable

$$u_1 = \frac{1}{w_1} + \mu \int^y \frac{\sin(y-s) ds}{(\alpha s + \beta)^2} \quad (5.7.12)$$

we recover the system (5.6.12).

The Lie point symmetries of the reduced system are, naturally, those listed in (5.6.13). In terms of the original variables these are

$$\begin{aligned} \Gamma_1 &= \left[ \int \frac{dt}{r^2 \dot{\theta}} \right] \partial_t, \\ \Gamma_2 &= 2 \left[ \int r I' dt \right] \partial_t + r^2 I' \partial_r + \partial_\theta, \\ \Gamma_3 &= 2 \left[ t + \int r I dt \right] \partial_t + r(1+rI) \partial_r, \\ \Gamma_{4\pm} &= 2 \left[ \int r e^{\pm i\theta} dt \right] \partial_t + r^2 e^{\pm i\theta} \partial_r, \\ \Gamma_{6\pm} &= \left[ \int \left( 2r(I' \pm iI) - \frac{\alpha}{r^2 \dot{\theta}} \right) e^{\pm 2i\theta} dt \right] \partial_t + [r(\pm i + r(I' \pm iI))] e^{\pm 2i\theta} \partial_r - e^{\pm 2i\theta} \partial_\theta, \\ \Gamma_{8\pm} &= \left[ \int \left( \frac{\dot{r}}{r^2 \dot{\theta}} + I'(1+2rI) \pm \frac{i}{r}(1+rI)^2 - \frac{\alpha}{r^3 \dot{\theta}}(1+rI) \right) e^{\pm i\theta} dt \right] \partial_t \\ &\quad + \left[ (1+rI)(rI' \pm i(1+rI)') e^{\pm i\theta} \right] \partial_r - \frac{1}{r}(1+rI) e^{\pm i\theta} \partial_\theta, \end{aligned} \quad (5.7.13)$$

where  $I$  stands for the second term in (5.7.12) and  $I'$  is its derivative with respect to  $\theta$ .

In addition to the symmetries listed in (5.7.13) equation (5.7.1) has the Lie point symmetry  $\partial_t$  which was the symmetry upon which the reduction of order was based. Consequently we can conclude that algebraically the Kepler problem and the Kepler problem with drag are identical.

### 5.7.2 The equation $\ddot{\mathbf{r}} - \frac{1}{2} \left( \frac{\dot{g}}{g} + 3\frac{\dot{r}}{r} \right) \dot{\mathbf{r}} + \mu g \mathbf{r} = \mathbf{0}$

The two components of this equation of motion in plane polar coordinates are

$$\begin{aligned} \ddot{r} &= r\dot{\theta}^2 + \frac{1}{2} \left( \frac{g'}{g} + \frac{3}{r} \right) \dot{r}^2 - \mu g r, \\ \ddot{\theta} &= \frac{1}{2} \dot{r} \dot{\theta} \left( \frac{g'}{g} - \frac{1}{r} \right), \end{aligned} \quad (5.7.14)$$

where we have written  $\dot{g}(r) = g'(r)\dot{r}$ . We proceed in the same fashion as before with the same choice of new independent variable and obtain the system of three first-order equations

$$\begin{aligned} w'_1 &= \frac{w_3}{w_4} \quad \Leftrightarrow \quad w_3 = w_4 w'_1, \\ w'_3 &= w_1 w_4 + \frac{1}{2} \left( \frac{g'}{g} + \frac{3}{r} \right) w_4 w_1'^2 - \mu g \frac{w_1}{w_4}, \\ w'_4 &= \frac{w_3}{2w_1} \left( \frac{g'}{g} w_1 - 1 \right) \end{aligned} \quad (5.7.15)$$

and again the obvious variable to be eliminated is  $w_3$  from the first of (5.7.15). With this substitution the third of (5.7.15) is easily integrated to give

$$A = \left( \frac{w_1}{g} \right)^{\frac{1}{2}} w_4 \quad (5.7.16)$$

and we use this as the definition of  $u_2$ . The second of (5.7.15) is now (after a certain amount of simplification)

$$u_2 w_1'' = w_1 u_2 + 2 \frac{u_2 w_1'^2}{w_1} - \frac{\mu w_1^2}{u_2}. \quad (5.7.17)$$

This equation is just the same equation as one obtains for the Kepler problem and we know that the correct substitution is  $u_1 = \mu - u_2^2/u_1$ . Again we have recovered the system (5.6.12) and so this problem is also algebraically equivalent to the Kepler problem. The symmetries are

$$\begin{aligned} \Gamma_1 &= \left[ \int \frac{r g'}{g} dt \right] \partial_t - 2r \partial_r, \\ \Gamma_2 &= \partial_\theta, \\ \Gamma_3 &= \frac{1}{2} \left[ \int r \left( \frac{1}{r} - \frac{g'}{g} \right) \left( \frac{\mu g}{\dot{\theta}^2} - 1 \right) dt \right] \partial_t + r \left( \frac{\mu g}{\dot{\theta}^2} - 1 \right) \partial_r, \\ \Gamma_{4\pm} &= \frac{1}{2} \left[ \int r^2 \left( \frac{1}{r} - \frac{g'}{g} \right) e^{\pm i\theta} dt \right] \partial_t + r^2 e^{\pm i\theta} \partial_r, \\ \Gamma_{6\pm} &= \pm \frac{1}{2} i \left[ \int \left( 3 + \frac{r g'}{g} + \frac{\mu(g - g'r)}{\dot{\theta}^2} \right) e^{\pm 2i\theta} dt \right] \partial_t \pm i r \left( \frac{\mu g}{\dot{\theta}^2} - 1 \right) e^{\pm 2i\theta} \partial_r + e^{\pm 2i\theta} \partial_\theta, \end{aligned}$$



$$\begin{aligned} \Gamma_{8\pm} = & \frac{1}{2} \left\{ \int e^{\pm i\theta} \left[ \frac{gr}{\dot{\theta}^2} \left( \mu - \frac{\dot{\theta}^2}{g} \right)^2 \left( \frac{1}{r} - \frac{g'}{g} \right) \pm 2i \left( \mu - \frac{\dot{\theta}^2}{g} \right) + \frac{2\dot{r}\dot{\theta}}{rg} \right] dt \right\} \partial_t \\ & + \left\{ e^{\pm i\theta} \frac{gr}{\dot{\theta}^2} \left( \mu - \frac{\dot{\theta}^2}{g} \right)^2 \right\} \partial_r + \left\{ e^{\pm i\theta} \left( \mu - \frac{\dot{\theta}^2}{g} \right) \right\} \partial_\theta. \end{aligned} \quad (5.7.18)$$

### 5.7.3 An example with an angle-dependent force

So far we have seen that by the correct choice of variables the reduced system comes down to be the same one, *videlicet* (5.6.12). One of the variables is related to the Laplace–Runge–Lenz vector. We apply the idea to the equation

$$\dot{\mathbf{r}} + g\hat{\mathbf{r}} + h\hat{\boldsymbol{\theta}} = 0, \quad (5.7.19)$$

where

$$g = \frac{U''(\theta) + U(\theta)}{r^2} + 2\frac{V'(\theta)}{r^{3/2}} \quad \text{and} \quad h = \frac{V(\theta)}{r^{3/2}}. \quad (5.7.20)$$

We make the same reduction as in previous cases to arrive at the pair of equations

$$w_1 w_1'' w_4^2 - 2w_1' w_4^2 - w_1^2 w_4^2 = w_1 h - g w_1, \quad (5.7.21)$$

$$w_1 w_4 w_4' + 2w_1' w_4^2 = -h, \quad (5.7.22)$$

where  $w_1 = r$  and  $w_4 = \dot{\theta}$  as before.

The Laplace–Runge–Lenz vector for equation (5.7.19) is

$$\mathbf{J} = \dot{\mathbf{r}} \times \mathbf{L} - U\hat{\mathbf{r}} - \left[ U' + 2r^{\frac{1}{2}}V \right] \hat{\boldsymbol{\theta}}. \quad (5.7.23)$$

If we take the two cartesian components of  $\mathbf{J}$ , *videlicet*  $J_1$  and  $J_2$ , and combine them we obtain the two Ermanno–Bernoulli constants

$$\begin{aligned} J_\pm &= -J_1 \pm iJ_2, \\ &= \left[ \left( \frac{L^2}{w_1} - U \right) \pm i \left( \frac{L^2}{w_1} - U \right)' \right] e^{\pm iy}. \end{aligned} \quad (5.7.24)$$

We see that the Ermanno–Bernoulli constants have the same structure as for the standard Kepler problem. Immediately we have the clue to the identification of one of new variables and we set

$$u_1 = \frac{L^2}{w_1} - U \quad (5.7.25)$$

so that the Ermanno–Bernoulli constants are given by

$$J_\pm = (u_1 \pm iu_1') e^{\pm iy}. \quad (5.7.26)$$

The identification of the second variable is more delicate. Equation (5.7.22) can be written as

$$LL' = -w_1^{\frac{3}{2}}V(y) \quad (5.7.27)$$

and, when (5.7.25) is taken into account, this becomes

$$0 = \frac{L'}{L^2} + \frac{V(y)}{(u_1 + U(y))^{\frac{3}{2}}}. \quad (5.7.28)$$

From (5.7.26) we have

$$u_1 = \frac{1}{2} (J_+ e^{-iy} + J_- e^{iy}) = J \cos y \quad (5.7.29)$$

since  $J_- = J_+^*$  and we have written  $J = |J_+| = |J_-|$ . We use (5.7.28) to define the second variable

$$u_2 = \frac{1}{L} - \int \frac{V(y)dy}{(J \cos y + U(y))^{\frac{3}{2}}}. \quad (5.7.30)$$

The reduced system of equations is again (5.6.12).

In addition to the symmetry  $\partial_t$  which was used to reduce the original system of equations the reduced system contributes the following symmetries to the original equation

$$\begin{aligned} \Gamma_1 &= 3 \left[ \int r^2 \dot{\theta} dt \right] \partial_t + 2r^3 \dot{\theta} \partial_r, \\ \Gamma_2 &= \left[ \int \left( \frac{2U'}{r^2 \dot{\theta}^2} + \frac{3Vr^2 \dot{\theta}}{U + J \cos \theta} \right) dt \right] \partial_t + \left[ \frac{U'}{r^2 \dot{\theta}^2} + \frac{2Vr^3 \dot{\theta}}{U + J \cos \theta} \right] \partial_r - \partial_\theta, \\ \Gamma_3 &= 2 \left[ t - \int \frac{U dt}{r^3 \dot{\theta}^2} \right] \partial_t + \left[ r - \frac{U}{r^2 \dot{\theta}^2} \right] \partial_r, \\ \Gamma_{4\pm} &= 2 \left[ \int \frac{e^{\pm i\theta}}{r^3 \dot{\theta}^2} dt \right] \partial_t + \frac{e^{\pm i\theta}}{r^2 \dot{\theta}^2} \partial_r, \\ \Gamma_{6\pm} &= \left\{ \int e^{\pm 2i\theta} \left[ \frac{3Vr^2 \dot{\theta}}{U + J \cos \theta} + \frac{2(U' \mp iU)}{r^3 \dot{\theta}^2} \right] dt \right\} \partial_t, \\ &\quad - e^{\pm 2i\theta} \left[ \frac{2Vr^3 \dot{\theta}}{U + J \cos \theta} \pm ir + \frac{U' \mp iU}{r^2 \dot{\theta}^2} \right] \partial_r + e^{\pm 2i\theta} \partial_\theta \\ \Gamma_{8\pm} &= \left\{ \int e^{\pm i\theta} \left[ 2U' \left( 1 - \frac{U}{r^3 \dot{\theta}^2} \right) - 3 \left( r^2 \dot{r} \dot{\theta} \pm ir^3 \dot{\theta}^2 - 2r^{\frac{1}{2}} V \mp iU \right) \right] dt \right\} \partial_t \quad (5.7.31) \\ &\quad - e^{\pm i\theta} \left[ \frac{2Vr^2 \dot{\theta}}{U + J \cos \theta} (U' \pm i(r^3 \dot{\theta}^2 - U)) \left( 1 - \frac{U}{r^3 \dot{\theta}^2} \right) \right] \partial_r + e^{\pm i\theta} [r^3 \dot{\theta}^2 - U] \partial_\theta. \end{aligned}$$

Again we have a system possessing a Laplace–Runge–Lenz vector having the same algebraic properties as the standard Kepler problem. The standard Kepler problem has a Hamiltonian structure whereas (5.7.19) was shown by Sen [130] to be Hamiltonian only under considerable restrictions on the functions  $U(\theta)$  and  $V(\theta)$ . It is presently an open question whether one can construct a Hamiltonian through these transformations.

### 5.7.4 A truly three-dimensional motion

We mentioned above that the three-dimensional treatment of the standard Kepler problem would require simply the addition of a second first order equation to the system (5.6.12) and that the suitable variable for this equation would be the  $z$ -component of the angular momentum. In the case of (4.1.12) we did not have a conserved  $z$ -component of the angular momentum. The magnitude of the angular momentum vector was conserved, but the direction of angular momentum was not constant and the motion was truly three-dimensional. This equation gives us a good chance to test the idea put forward above that one of the useful variables in the reduction is somehow related to the Laplace–Runge–Lenz vector in the case in which we cannot reduce the problem to one of two dimensions. We recall that the equation of motion

$$\ddot{\mathbf{r}} + \frac{h'}{r} \mathbf{L} + \left( hh' + \frac{k}{r^2} \right) \hat{\mathbf{r}} = 0 \quad (5.7.32)$$

has the Laplace–Runge–Lenz vector

$$\mathbf{J} = \dot{\mathbf{r}} \times \mathbf{L} - h(r) \mathbf{L} - k \mathbf{r}. \quad (5.7.33)$$

In addition the square of the magnitude of the angular momentum, *videlicet*

$$L^2 = r^4 \left( \dot{\theta}^2 + \dot{\phi}^2 \sin^2 \theta \right), \quad (5.7.34)$$

is a conserved scalar. In spherical polar coordinates the three components of the equation of motion, (5.7.32), are

$$\ddot{r} = r \left( \dot{\theta}^2 + \dot{\phi}^2 \sin^2 \theta \right) - hh' - \frac{k}{r^2}, \quad (5.7.35)$$

$$\ddot{\theta} = -2 \frac{\dot{r} \dot{\theta}}{r} + \dot{\phi}^2 \sin \theta \cos \theta + h' \sin \theta \dot{\phi}, \quad (5.7.36)$$

$$\ddot{\phi} \sin \theta = -2 \frac{\dot{r} \dot{\phi}}{r} \sin \theta - 2 \dot{\theta} \dot{\phi} \cos \theta - h' \dot{\theta}. \quad (5.7.37)$$

The three components of the Laplace–Runge–Lenz vector are

$$\begin{aligned} J_x &= \left( \frac{L^2}{r} - k \right) \sin \theta \cos \phi - r^2 \left( \dot{r} \dot{\theta} - h \dot{\phi} \sin \theta \right) \cos \theta \cos \phi + r^2 \left( \dot{r} \dot{\phi} \sin \theta + h \dot{\theta} \right) \sin \phi, \\ J_y &= \left( \frac{L^2}{r} - k \right) \sin \theta \sin \phi - r^2 \left( \dot{r} \dot{\theta} - h \dot{\phi} \sin \theta \right) \cos \theta \sin \phi - r^2 \left( \dot{r} \dot{\phi} \sin \theta + h \dot{\theta} \right) \cos \phi, \\ J_z &= \left( \frac{L^2}{r} - k \right) \cos \theta + r^2 \left( \dot{r} \dot{\theta} - h \dot{\phi} \sin \theta \right) \sin \theta. \end{aligned} \quad (5.7.38)$$

To obtain the Ermanno–Bernoulli constants we take the combinations

$$\begin{aligned} J_{\pm} &= J_x \pm i J_y \\ &= e^{\pm i \phi} \left[ \left( \left( \frac{L^2}{r} - k \right) \sin \theta - r^2 \left( \dot{r} \dot{\theta} - h \dot{\phi} \sin \theta \right) \cos \theta \right) \pm i \left( -r^2 \left( \dot{r} \dot{\phi} \sin \theta + h \dot{\theta} \right) \right) \right]. \end{aligned} \quad (5.7.39)$$

Since  $\phi$  does not appear in the components of the equation of motion, we choose it to be the new independent variable. It is a simple matter to verify that the derivative of the real part within crochets in (5.7.39) with respect to  $\phi$  gives the imaginary part and so we define the new variable

$$u_1 = \left( \frac{L^2}{r} - k \right) \sin \theta - r^2 \left( \dot{r}\dot{\theta} - h\dot{\phi} \sin \theta \right) \cos \theta. \quad (5.7.40)$$

Thus we have the Ermanno–Bernoulli constants in the standard form

$$J_{\pm} = e^{\pm i\phi} (u_1 \pm iu_1'), \quad (5.7.41)$$

where the new variable  $u_1$  satisfies the second order equation

$$u_1'' + u_1 = 0. \quad (5.7.42)$$

The other two variables for the reduced system can be taken to be the remaining component of the Laplace–Runge–Lenz vector and the square of the magnitude of the angular momentum. We write

$$\begin{aligned} u_2 &= r^4 \left( \dot{\theta}^2 + \dot{\phi}^2 \sin^2 \theta \right), & u_2' &= 0, \\ u_3 &= \left( \frac{L^2}{r} - k \right) \cos \theta + r^2 \left( \dot{r}\dot{\theta} - h\dot{\phi} \sin \theta \right) \sin \theta, & u_3' &= 0, \end{aligned} \quad (5.7.43)$$

and so we have the complete reduction of (5.7.32) to a system of two first-order equations and the simple harmonic oscillator. The algebra of the Lie point symmetries is  $2A_1 \oplus sl(3, R)$  if one takes the two symmetries

$$G_9 = A(u_2, u_3)\partial_{u_2} \quad \text{and} \quad G_{10} = B(u_2, u_3)\partial_{u_3}, \quad (5.7.44)$$

where  $A$  and  $B$  are arbitrary functions of the two first integrals, to be equivalent to just  $\partial_{u_2}$  and  $\partial_{u_3}$ . The involved expressions for the new variables and the additional complication of an extra dimension make the computation of the symmetries in the original variables a task of somewhat greater length than the result justifies and we omit them.

Finally we note that, of the Lie point symmetries of the system of differential equations, those associated with  $J_{\pm}$  are

$$\begin{aligned} G_{1\pm} &= \pm i\partial_y + u_1\partial_{u_1}, \\ G_{2\pm} &= e^{\pm 2iy} (\partial_y \pm iu_1\partial_{u_1}) \quad \text{and} \quad G_{3\pm} = e^{\pm iy}\partial_{u_1} \end{aligned} \quad (5.7.45)$$

in addition to the trivial  $G_9$  and  $G_{10}$ . To these three symmetries in the original representation there is added  $\partial_t$ .

## Acknowledgments

PGLL thanks the Director of GEODYSYC, Dr S Cotsakis, and the Department of Mathematics, University of the Aegean, for their hospitality and provision of facilities while this work was undertaken and acknowledges the continued support of the National Research Foundation of the Republic of South Africa and the University of Natal.

## References

- [1] Abraham-Shrauner B, Hidden Symmetries and Linearisation of the Modified Painlevé–Ince Equation, *J. Math. Phys.* **34** (1993), 4809–4816.
- [2] Abraham-Shrauner B and Guo A, Hidden Symmetries Associated with the Projective Group of Nonlinear First Order Differential Equations *J. Phys. A: Math. Gen.* **25** (1992), 5597–5608.
- [3] Abraham-Shrauner B and Guo A, Hidden and Nonlocal Symmetries in Nonlinear Differential Equations, in *Modern Group Analysis: Advanced Analytical and Computational Methods in Mathematical Physics*, Edditors: Ibragimov N H, Torrisi M and Valente A, Kluwer, Dordrecht, 1993, 1–5.
- [4] Abraham-Shrauner B and Leach P G L, Hidden Symmetries of Nonlinear Ordinary Differential Equations, in *Exploiting Symmetry in Applied and Numerical Analysis*, Editors: Allgower E, George K and Miranda R, *Lecture Notes in Applied Mathematics*, Vol. 29, American Mathematical Society, Providence, 1993, 1–10.
- [5] Andriopoulos K, Leach P G L and Flessas G P, Complete Symmetry Groups of Ordinary Differential Equations and Their Integrals: Some Basic Considerations, *J. Math. Anal. Appl.* **262** (2001), 256–273.
- [6] Atiyah M F and Hitchin N J, Low Energy Scattering of Nonabelian Magnetic Monopoles, *Phil. Trans. R. Soc. Lond.* **315** (1985), 459–469.
- [7] Atiyah M F and Hitchin N J, Low Energy Scattering of Nonabelian Monopoles, *Phys. Lett.* **A107** (1985), 21–25.
- [8] Bacry H, Comment on “The Relationship Between the Symmetries of and the Existence of Conserved Vectors for the Equation  $\ddot{\mathbf{r}} + f(r)\mathbf{L} + g(r)\mathbf{r} = 0$ ”, *J. Phys. A: Math. Gen.* **24** (1991), 1157–1159.
- [9] Bacry H, Ruegg H and Souriau J M, Dynamical Groups and Spherical Potentials in Classical Mechanics, *Commun. Math. Phys.* **3** (1966), 323–333.
- [10] Bargmann V, Zur Theorie des Wasserstoffatoms. Bemerkungen zur gleichnamigen Arbeit von V Fock, *Z. Phys.* **99** (1936), 576–582.
- [11] Bates L, Symmetry Preserving Deformations of the Kepler Problem, Preprint, Department of Mathematics and Statistics, University of Calgary, 1990.
- [12] Bernoulli J, Extrait de la Réponse de M Bernoulli à M Herman, datée de Basle le 7 Octobre, 1710, *Hist. Acad. Roy. Sci. Mém. Math. Phys.*, (1710), 521–533.
- [13] Bertrand J, Théorème relatif au mouvement d’un point attiré vers un centre fixe, *Compt. Rend. Acad. Sci. Paris* **77** (1873), 849–853.
- [14] Bleuler K and Kustaanheimo P E, Vectorielle Methoden in der Quantenmechanik I, *Ann. Acad. Sci. Fenn.* **AVI** (1967), 1–14.
- [15] Bouquet S É, Feix M R and Leach P G L, Properties of Second Order Ordinary Differential Equations Invariant under Time Translation and Selfsimilar Transformation, *J. Math. Phys.* **32** (1991), 1480–1490.
- [16] Brand L, *Advanced Calculus*, John Wiley, New York, 1955.

- [17] Broucke R, On the Construction of a Dynamical System from a Preassigned Family of Solutions, *Int. J. Eng. Sci.* **17** (1979), 1151–1162.
- [18] Brouwer D and Hori G, Theoretical Evaluation of Atmospheric Drag Effects in the Motion of an Artificial Satellite, *Astron. J.* **66** (1961), 193–225.
- [19] Caspar Max, Kepler, Dover, New York, 1993.
- [20] Child J M, Hamilton’s Hodograph, *The Monist* **25** (1915), 615–624.
- [21] Collas P, Algebraic Solution of the Kepler Problem Using the Runge–Lenz Vector, *Am. J. Phys.* **38** (1970), 253–255.
- [22] Collinson C D, Investigation of Planetary Orbits Using the Lenz–Runge Vector, *Bull. Inst. Math. Appl.* **9** (1973), 377–378.
- [23] Collinson C D, The Mathematical Simplicity of the Inverse Square Force, *Mathl. Gaz.* **61** (1977), 139–140.
- [24] Colwell P, Bessel Functions and Kepler’s Equation, *Amer. Math. Mon.* **99** (1992), 45–49.
- [25] Cordani B, On the Fock Quantisation of the Hydrogen Atom, Preprint 47/1988, Dipartimento di Matematica ‘F Enriques’, Via C Saldini 50, 20133 Milano, Italia, 1988.
- [26] Cotsakis S, Pantazi H and Leach P G L, Symmetries of Homogeneous Cosmologies, *Gravitation & Cosmology* **4** (1998), 314–325.
- [27] Danby J M A, Fundamentals of Celestial Mechanics, Macmillan, New York, 1962.
- [28] Dirac P A M, Principles of Quantum Mechanics, Cambridge University Press, Cambridge, 1932.
- [29] Dreyer J L E, A History of Astronomy from Thales to Kepler, Dover, New York, 1953.
- [30] Dugas R, A History of Mechanics, Dover, New York, 1988 (trans. by J R Maddox).
- [31] Dutka J, On the Early History of Bessel Functions, *Arch. Hist. Exact Sci.* **49** (1995), 105–134.
- [32] Ermakov V, Second Order Differential Equations. Conditions for Complete Integrability, *Univ. Izvestia Kiev Ser. III* **9** (1880), 1–25 (trans. by A O Harin).
- [33] Ermanno J, Metodo d’investigare l’Orbite de’ Pianeti, nell’ipotesi che le forze centrali o pure le gravità delgi stessi Pianeti sono in ragione reciproca de’ quadrati delle distanze, che i medesimi tengono dal Centro, a cui si dirigono le forze stesse, *G. Litt. Ital.* **2** (1710), 447–467.
- [34] Fock V, Zur Theorie des Wasserstoffatoms, *Z. Phys.* **98** (1935), 145–154.
- [35] Fradkin D M, Three-Dimensional Isotropic Harmonic Oscillator and  $SU(3)$ , *Am. J. Phys.* **33** (1965), 207–211.
- [36] Gascón F G, Ramos F B and Aguirre-Daban E, On the Polynomial First Integrals of Certain Second-Order Differential Equations, *J. Math. Phys.* **23** (1982), 2281–2285.
- [37] Gibbs J W and Wilson E B, Vector Analysis, Dover, New York, 1960.
- [38] Goldstein H, Prehistory of the ‘Runge–Lenz’ Vector, *Am. J. Phys.* **43** (1975), 737–738.

- [39] Goldstein H, More on the Prehistory of the Laplace or Runge–Lenz Vector, *Am. J. Phys.* **44** (1976), 1123–1124.
- [40] Goldstein H, *Classical Mechanics*, World Student Series Edition, Addison-Wesley, Massachusetts, 1980.
- [41] Gorringe V M, Generalisations of the Laplace–Runge–Lenz Vector in Classical Mechanics, Thesis, Faculty of Science, University of the Witwatersrand, Johannesburg, Voll I, II, 1986.
- [42] Gorringe V M and Leach P G L, Dynamical Properties of the System  $\ddot{\mathbf{r}} = \ddot{u}\mathbf{r}/u - \mu\mathbf{r}/(ur^3)$ , Preprint, Department of Computational and Applied Mathematics, University of the Witwatersrand, Johannesburg, 1986.
- [43] Gorringe V M and Leach P G L, Conserved Vectors for the Autonomous System  $\ddot{\mathbf{r}} + g(r, \theta)\hat{\mathbf{r}} + h(r, \theta)\hat{\boldsymbol{\theta}} = 0$ , *Physica* **D27** (1987), 243–248.
- [44] Gorringe V M and Leach P G L, Conserved Vectors and Orbit Equations for Motions in Two and Three Dimensions, in *Finite Dimensional Integrable Nonlinear Dynamical Systems*, Editors: Leach P G L and Steeb W-H, World Scientific, Singapore, 1988, 190–204.
- [45] Gorringe V M and Leach P G L, Hamilton-Like Vectors for a Class of Kepler Problems with a Force Proportional to the Velocity, *Celest. Mech.* **41** (1988), 125–130.
- [46] Gorringe V M and Leach P G L, Conserved Vectors and Orbit Equations for Autonomous Systems Governed by the Equation of Motion  $\ddot{\mathbf{r}} + f\dot{\mathbf{r}} + g\mathbf{r} = 0$ , *Am. J. Phys.* **57** (1989), 432–435.
- [47] Gorringe V M and Leach P G L, First Integrals Associated with the Additional Symmetry of Central Force Problems with Power Law Potentials, *Quaest. Math.* **14** (1991), 277–289.
- [48] Gorringe V M and Leach P G L, Kepler’s Third Law and the Oscillator’s Isochronism, *Am. J. Phys.* **61** (1993), 991–995.
- [49] Gorringe V M and Leach P G L, The First Integrals and Their Lie Algebra of the Most General Autonomous Hamiltonians of the Form  $H = T + V$  Possessing a Laplace–Runge–Lenz Vector, *J. Austral. Math. Soc. Ser. B* **34** (1993), 511–522.
- [50] Govinder K S, Athorne C and Leach P G L, The Algebraic Structure of Generalised Ermakov Systems in Three Dimensions, *J. Phys. A: Math. Gen.* **26** (1993), 4035–4046.
- [51] Gradshteyn I S and Ryzhik I M, *Table of Integrals, Series and Products*, Academic Press, Orlando, 1980.
- [52] Grammaticos B, Dorizzi B and Ramani A, Hamiltonians with Higher-Order Integrals and the ‘Weak-Painlevé’ Concept, *J. Math. Phys.* **25** (1984), 3470–3473.
- [53] Günther N J and Leach P G L, Generalised Invariants for the Time-Dependent Harmonic Oscillator, *J. Math. Phys.* **18** (1977), 572–576.
- [54] Hall L S, A Theory of Exact and Configurational Invariants, *Physica* **D8** (1982), 90–113.
- [55] Hamilton W R, On the Application of the Method of Quaternions to Some Dynamical Questions, *Proc. Roy. Irish Acad.* **3** (1847), 344–353, also Appendix III p xxxvi ff.
- [56] Hankins T L, *Sir William Rowan Hamilton*, John Hopkins University Press, Maryland, 1980.

- [57] Head A K, LIE, a PC Program for Lie Analysis of Differential Equations, *Comp. Phys. Commun.* **77** (1993), 241–248.
- [58] Heath T, Aristarchus of Samos: the ancient Copernicus, Dover, New York, 1981.
- [59] Herman J, Extrait d’une lettre de M Herman à M Bernoulli, datée de Padoüe le 12 Juillet, 1710, *Hist. Acad. Roy. Sci. Mém. Math. Phys.* (1710), 519–521.
- [60] Hestenes D, New Foundations of Classical Mechanics, D Riedel, Dordrecht, 1986.
- [61] Hietarinta J, Classical Versus Quantum Integrability, *J. Math. Phys.* **25** (1984), 1833–1840.
- [62] Hietarinta J, A Listing of Integrable Two-Degree-of-Freedom Hamiltonians, Preprint, University of Turku, Finland, 1985.
- [63] Ince E L, Integration of Ordinary Differential Equations, Longmans, Green & Co, London, 1927.
- [64] Jackiw R, Dynamical Symmetry of the Magnetic Monopole, *Ann. Phys. (NY)* **129** (1980), 183–200.
- [65] Jauch J M and Hill E L, On the Problem of Degeneracy in Quantum Mechanics, *Phys. Rev.* **57** (1940), 641–645.
- [66] Jezewski D J and Mittleman D, Integrals of Motion for the Classical Two-Body Problem with Drag, *Int. J. Nonlin. Mech.* **18** (1983), 119–124.
- [67] Kaplan H, The Runge–Lenz Vector as an ‘Extra’ Constant of the Motion, *Am. J. Phys.* **54** (1986), 157–161.
- [68] Katzin G H and Levine J, Time-Dependent Quadratic Constant of Motion, Symmetries, and Orbit Equations for Classical Particle Dynamical Systems with Time-Dependent Kepler Potentials, *J. Math. Phys.* **23** (1982), 552–563.
- [69] Katzin G H and Levine J, Time-Dependent Vector Constant of Motion, Symmetries, and Orbit Equations for the Dynamical System  $\ddot{\mathbf{r}} = \hat{\mathbf{I}}_r\{\ddot{U}(t)/U(t)r - [\mu_0/U(t)]r^{-2}\}$ , *J. Math. Phys.* **24** (1983), 1761–1771.
- [70] Kelland P and Tait P G, Introduction to Quaternions, Macmillan, London, 1904.
- [71] Kelvin Lord and Tait P G, Elements of Natural Philosophy, Cambridge University Press, Cambridge, 1894.
- [72] Kelvin Lord and Tait P G, Treatise on Natural Philosophy, Cambridge University Press, Cambridge, 1923.
- [73] Kepler J, *Astronomia Nova in Opera Omnia*, Editor: Frisch C H, Heyder & Zimmer, Frankfurt, Vol. III, 1860.
- [74] Kepler J, *Harmonia Mundi in Opera Omnia*, Editor: Frisch C H, Heyder & Zimmer, Frankfurt, Vol. V, 1864.
- [75] Krause J, On the Complete Symmetry Group of the Classical Kepler System, *J. Math. Phys.* **35** (1994), 5734–5748.
- [76] Kruskal M, Asymptotic Theory of Hamiltonian and Other Systems with All Solutions Nearly Periodic, *J. Math. Phys.* **3** (1962), 806–828.



- [77] Kummer E E, De generali quadam æquatione differentiali tertii ordinis, *J. für Math.* **100** (1887), 1–9 (reprinted from the Programm des evangelischen Königl. und Stadtgymnasiums in Liegnitz for the year 1834).
- [78] Laplace P S, *Traité de Mécanique Céleste*, Villars, Paris, 1799, Vol. I, p. 165ff.
- [79] Leach P G L, On the Theory of Time-Dependent Linear Canonical Transformations as Applied to Hamiltonians of Harmonic Oscillator Type, *J. Math. Phys.* **18** (1977), 1608–1611.
- [80] Leach P G L, Applications of the Lie Theory of Extended Groups in Hamiltonian Mechanics: the Oscillator and the Kepler Problem, *J. Austral. Math. Soc.* **B23** (1981), 173–186.
- [81] Leach P G L, Classes of Potentials of Time-Dependent Central Force Fields which Possess First Integrals Quadratic in the Momentum, *J. Math. Phys.* **26** (1985), 1613–1620.
- [82] Leach P G L, Comment on an Aspect of a Paper by G Thompson, *J. Math. Phys.* **27** (1986), 153–156 and Erratum **27**, 2445–2446.
- [83] Leach P G L, The First Integrals and Orbit Equation for the Kepler Problem with Drag, *J. Phys. A: Math. Gen.* **20** (1987), 1997–2002.
- [84] Leach P G L and Bouquet S É, Symmetries and Integrating Factors, *J. Nonlinear Math. Phys.* **9**, Suppl. 2 (2002), 73–91.
- [85] Leach P G L, Cotsakis S and Flessas G P, Symmetry, Singularities and Integrability in Complex Dynamics: I The Reduction Problem, *J. Nonlinear Math. Phys.* **7** (2000), 445–479.
- [86] Leach P G L, Feix M R and Bouquet S É, Analysis and Solution of a Second Order Differential Equation through Rescaling and a Dynamical Point of View, *J. Math. Phys.* **29** (1988), 2563–2569.
- [87] Leach P G L and Gorringe V M, Variations on Newton’s Keplerian Theme, *S. Afr. J. Sci.* **83** (1987), 550–555.
- [88] Leach P G L and Gorringe V M, A Conserved Laplace–Runge–Lenz-Like Vector for a Class of Three-Dimensional Motions, *Phys. Lett.* **A133** (1988), 289–294.
- [89] Leach P G L and Gorringe V M, The Relationship between the Symmetries of and the Existence of Conserved Vectors for the Equation  $\ddot{\mathbf{r}} + f(r)\mathbf{L} + g(r)\mathbf{r} = 0$ , *J. Phys. A: Math. Gen.* **23** (1990), 2765–2774.
- [90] Leach P G L, Govinder K S and Abraham-Shrauner B, Symmetries of First Integrals and their Associated Differential Equations, *J. Math. Anal. Appl.* **235** (1999), 58–83.
- [91] Leach P G L, Nucci M C and Cotsakis S, Symmetry, Singularities and Integrability in Complex Dynamics: V Complete Symmetry Groups, *J. Nonlinear Math. Phys.* **8** (2001), 475–490.
- [92] Lemmer R L, *The Paradigms of Mechanics: a Symmetry Based Approach*, Dissertation, Department of Mathematics and Applied Mathematics, Faculty of Science, University of Natal, 1996.
- [93] Lenz W, Über den Bewegungsverlauf und die Quantenzustände der gestörten Keplerbewegung, *Z. Phys.* **24** (1924), 197–207.

- [94] Lewis H Ralph Jr, Classical and Quantum Systems with Time-Dependent Harmonic-Oscillator-Type Hamiltonians, *Phys. Rev. Lett.* **18** (1967), 510–512 and Erratum **18**, 636.
- [95] Lewis H Ralph Jr, Class of Exact Invariants for Classical and Quantum Time-Dependent Harmonic Oscillators, *J. Math. Phys.* **9** (1968), 1976–1986.
- [96] Lewis H Ralph Jr and Leach P G L, A Direct Approach to Finding the Exact Invariants for One-Dimensional Time-Dependent Classical Hamiltonians, *J. Math. Phys.* **23** (1982), 2371–2374.
- [97] Lewis H Ralph Jr and Riesenfeld W B, An Exact Quantum Theory of the Time-Dependent Harmonic Oscillator and of a Charged Particle in a Time-Dependent Electromagnetic Field, *J. Math. Phys.* **10** (1969), 1458–1473.
- [98] Lie S, *Differentialgleichungen*, B G Teubner, Stuttgart, 1967.
- [99] Lie S, *Continuerliche Gruppen*, B G Teubner, Stuttgart, 1971.
- [100] Liouville J, Sur le développement des fonctions ou parties des fonctions en séries dont les divers termes sont assujettis à satisfaire à une même équation différentielle du second ordre contenant un paramètre variable, *J. de Math.* **II** (1837), 16–35.
- [101] Lutzky M, Dynamical Symmetries and Conserved Quantities, *J. Phys. A: Math. Gen.* **12** (1979), 973–981.
- [102] Manton N S, Monopole Interactions at Long Range, *Phys. Lett.* **B154** (1985), 397–400 and Erratum **B157**, 475.
- [103] Mavraganis A G, The Almost-Constant Speed Two-Body Problem with Resistance, Preprint, Department of Engineering Science, National Technical University of Athens, 1991.
- [104] Maxwell J C, *Matter and Motion*, Sheldon, London, 1925.
- [105] Mazzone S and Roero C S, *Jacob Hermann and the Diffusion of the Leibnizian Calculus in Italy*, Leo S Olschki, Firenze, 1997.
- [106] McIntosh H V and Cisneros A, Degeneracy in the Presence of a Magnetic Monopole, *J. Math. Phys.* **11** (1970), 896–916.
- [107] Mittleman D and Jezewski D J, An Analytic Solution to the Classical Two-Body Problem with Drag, *Celest. Mech.* **28** (1982), 401–413.
- [108] Mladenov I M, Scattering of Charged Particles by Dyons, *J. Phys. A: Math. Gen.* **21** (1988), L1–L4.
- [109] Mladenov I M and Tsanov V V, Geometric Quantisation of the MIC-Kepler Problem, *J. Phys. A: Math. Gen.* **20** (1987), 5865–5871.
- [110] Moreira I C, Comments on a ‘Direct Approach to Finding the Exact Invariants for One-Dimensional Time-Dependent Classical Hamiltonians’, Preprint, Universidade Federal do Rio de Janeiro, Instituto de Física, 1983.
- [111] Moreira I C, A Note on the Invariants for the Time-Dependent Oscillator, *J. Phys. A: Math. Gen.* **18** (1985), 899–907.
- [112] Moreira I C, Ritter O M and Santos F C, Lie Symmetries for the Charge-Monopole Problem, *J. Phys. A: Math. Gen.* **18** (1985), L427–L430.

- [113] Newton I, *Principia*, Editor: Cajori F, University of California Press, Berkeley, Voll I & II, 1962 (translation by A Motte).
- [114] Noether E, Invariante Variationsprobleme, *Kgl. Ges. Wiss. Nachrichten Math.-Phys. Klasse Heft 2* (1918), 235–257.
- [115] Nucci M C, Interactive REDUCE Programs for Calculating Classical, Non-Classical and Lie-Bäcklund Symmetries of Differential Equations, Preprint GT Math: 062090-051, Georgia Institute of Technology, Atlanta, 1990.
- [116] Nucci M C, Interactive REDUCE Programs for Calculating Lie Point, Nonclassical, Lie-Bäcklund, and Approximate Symmetries of Differential Equations: Manual and Floppy Disk, in *CRC Handbook of Lie Group Analysis of Differential Equations*, Vol. III: New Trends, Editor: Ibragimov N H, CRC Press, Boca Raton FL, 1996, 415–481.
- [117] Nucci M C, The Complete Kepler Group Can Be Derived by Lie Group Analysis, *J. Math. Phys.* **37** (1996), 1772–1775.
- [118] Nucci M C and Leach P G L, The Harmony in the Kepler and Related Problems, *J. Math. Phys.* **42** (2001), 746–764.
- [119] Pauli W, Über das Wasserstoffspektrum vom Standpunkt der neuen Quantenmechanik, *Z. Phys.* **36** (1926), 336–363.
- [120] Pillay T and Leach P G L, The Fate of Kepler’s Laws of Planetary Motion in the Kepler-Dirac Problem, *S. Afr. J. Sci.* **94** (1998), 482–486.
- [121] Pillay T and Leach P G L, Generalised Laplace–Runge–Lenz Vectors and Nonlocal Symmetries, *S. Afr. J. Sci.* **95** (1999), 403–407.
- [122] Pillay T and Leach P G L, Chaos, Integrability and Symmetry, *S. Afr. J. Sci.* **96** (2000), 371–376.
- [123] Poincaré H, Remarques sur une expérience de M Birkeland, *Compt. Rend. Acad. Sci. Paris* **123** (1896), 530–533.
- [124] Pollard H, *Celestial Mechanics*, The Carus Mathematical Monographs, Vol. 18, The Mathematical Association of America, Providence, 1976.
- [125] Prince G E and Eliezer C J, On the Lie Symmetries of the Classical Kepler Problem, *J. Phys. A: Math. Gen.* **14** (1981), 587–596.
- [126] Routh E J, *A Treatise on the Dynamics of a Particle*, Cambridge University Press, Cambridge, 1898.
- [127] Runge C, *Vector Analysis*, Methuen, London, Vol. I, 1923 (translation by H Levy).
- [128] Sarlet W and Bahar L Y, A Direct Construction of First Integrals for Certain Nonlinear Dynamical Systems, *Int. J. Nonlin. Mech.* **15** (1980), 133–146.
- [129] Sarlet W, Leach P G L and Cantrijn F, First Integrals Versus Configurational Invariants and a Weak Form of Complete Integrability, *Physica* **D17** (1985), 87–98.
- [130] Sen T, A Class of Integrable Potentials, *J. Math. Phys.* **28** (1987), 2841–2850.
- [131] Sherring J, Head A K and Prince G E, Dimsym and LIE: Symmetry Determining Packages, *Math. Comp. Model.* **25** (1998), 153–164.

- 
- [132] Speiser D, The Kepler Problem from Newton to Johan Bernoulli, *Arch. Hist. Exact Sci.* **50** (1996), 103–116.
- [133] Stickforth J, The Classical Kepler Problem in Momentum Space, *Am. J. Phys.* **46** (1978), 74–75.
- [134] Symon K, *Mechanics*, World Student Series Edition, Addison-Wesley, Massachusetts, 1969.
- [135] Tait P G, *An Elementary Treatise on Quaternions*, Cambridge University Press, Cambridge, 1890.
- [136] Thompson G, Polynomial Constants of Motion in Flat Space, *J. Math. Phys.* **25** (1984), 3474–3478.
- [137] Thompson G, Second Order Systems with Runge–Lenz-Type Vectors, *Lett. Math. Phys.* **14** (1987), 69–75.
- [138] Weinstock R, Dismantling a Centuries-Old Myth: Newton’s Principia and Inverse-Square Orbits, *Am. J. Phys.* **50** (1982), 610–617.
- [139] Weinstock R, Long-Buried Dismantling of a Centuries-Old Myth: Newton’s Principia and Inverse-Square Orbits, *Am. J. Phys.* **57** (1989), 846–849.
- [140] Whittaker E T, *A Treatise on the Analytical Dynamics of Particles and Rigid Bodies*, Cambridge University Press, Cambridge, 1944.
- [141] Zwansiger D, Exactly Solvable Nonrelativistic Model of a Particle with Both Electric and Magnetic Charges, *Phys. Rev.* **176**, 1480–1488.