

Generalised CR -submanifolds of an (ε, δ) -trans-Sasakian manifold with certain connection

Mohd. Danish Siddiqi¹, Janardan P. Ojha² and Mobin Ahmad³

^{1,3}Department of Mathematics, College of Science
Jazan University, Jazan, K.S.A.

²Department of Mathematics, SRMGPC
Faizabad Road, Lucknow-227105, India

Abstract: In this paper, generalised CR -submanifolds of an (ε, δ) -trans-Sasakian manifolds with semi-symmetric non-metric connection are studied. Moreover, integrability conditions of the distributions on generalised CR -submanifolds of an (ε, δ) -trans-Sasakian manifolds with semi-symmetric non-metric connection and geometry of leaves with semi-symmetric non-metric connection have been discussed.

2000 Mathematical Subject Classification : 53C21, 53C25, 53C05.

Keywords: Distributions, (ε, δ) -trans-Sasakian manifolds, Generalised CR submanifolds, Semi-symmetric non-metric connection.

I. Introduction

Ion Mihai [1] introduced a new class of submanifolds called Generalised CR -submanifolds of Kaehler manifolds and also studied generalised CR -submanifolds of Sasakian manifolds [2]. In 1985, Oubina [3] introduced a new class of almost contact Riemannian manifolds known as trans-Sasakian manifolds. After M. H. Shahid studied CR -submanifolds of trans-Sasakian manifold [4] and generic submanifolds of trans-Sasakian manifold [5]. In 2001, A. Kumar and U.C. De [6] studied generalised CR -submanifolds of a trans-Sasakian manifold. In 1993, A. Bejancu and K. L. Duggal [7] introduced the concept of (ε) -Sasakian manifolds. Then U. C. De and A. Sarkar [8] introduced (ε) -Kenmotsu manifolds. The existence of a new structure on indefinite metrics has been discussed. Moreover, Bhattacharyya [9] studied the contact CR -submanifolds of indefinite trans-Sasakian manifolds. Recently, Nagaraja et. al. [10] introduced the concept of (ε, δ) -trans-Sasakian manifolds which generalised the notion of (ε) -Sasakian as well as (δ) -Kenmotsu manifolds. In 2010, Cihan Özgür [11] studied the submanifolds of Riemannian manifold with semi-symmetric non-metric connection. Moreover, Özgür and others also studied the different structures with semi-symmetric non-metric connection in ([12], [13]). On other hand, some properties of semi-invariant submanifolds, hypersurfaces and submanifolds with semi-symmetric non-metric connection were studied in ([14], [15], [16]). Thus motivated sufficiently from the above studies, in this paper we study generalised CR -submanifolds of an (ε, δ) -trans-Sasakian manifolds with semi-symmetric non-metric connection.

We know that a connection ∇ with a Riemannian metric g on a manifold M is called metric such that $\nabla g = 0$, otherwise it is non-metric. Further it is said to be a semi-symmetric linear connection [17]. A linear connection ∇ is said to be a semi-symmetric connection if its torsion tensor is of the form

$$T(X, Y) = \eta(Y)X - \eta(X)Y,$$

where η is a 1-form. A study of semi-symmetric connection on Riemannian manifold was enriched by K. Yano [18]. In 1992, Agashe and Chaffle [19] introduced the notion of semi-symmetric non-metric connection. It is well known that a linear connection is symmetric and metric if and only if it is the Levi-Civita connection.

Semi-Symmetric metric connection plays an important role in the study of Riemannian manifolds, there are various physical problems involving the semi-symmetric metric connection. For example if a man is moving on the surface of the earth always facing one definite point, say Mekka or Jaruselam or the North pole, then this displacement is semi-symmetric and metric [20].

In this paper, we study Generalised CR -submanifolds of an (ε, δ) trans-Sasakian manifolds with a semi-symmetric non-metric connection. This paper is organized as follows. In section 2, we give a brief introduction of generalised CR -submanifolds of an (ε, δ) trans-Sasakian manifold and give an example. In

section 3, we discuss some Basic Lemmas . In section 4, integrability of some distributions discuss. In section 5, Geometry of leaves of Generalised CR -submanifolds of an (ε, δ) -trans-Sasakian manifold with semi-symmetric non-metric connection have been discussed .

II. (ε, δ) -trans-Sasakian manifolds

Let \bar{M} be an almost contact metric manifold with an almost contact metric structure (ϕ, ξ, η, g) , where ϕ is a (1,1) tensor field, ξ is a vector field, η is a 1-form and g is a compatible Riemannian metric such that

$$\phi^2 = -I + \eta \otimes \xi, \quad \phi\xi = 0, \quad \eta\phi = 0, \quad \eta(\xi) = 1, \quad (2.1)$$

$$g(\phi X, \phi Y) = g(X, Y) - \varepsilon \eta(X)\eta(Y), \quad (2.2)$$

$$g(\xi, \xi) = \varepsilon \quad (2.3)$$

$$g(X, \phi Y) = -g(\phi X, Y), \quad \varepsilon g(X, \xi) = \eta(X) \quad (2.4)$$

for all vector fields X, Y on $T\bar{M}$, where $\varepsilon = g(\xi, \xi) = \pm 1$. An (ε) - almost contact metric manifold is called an (ε, δ) -trans Sasakian manifold [10] if

$$(\bar{\nabla}_X \phi)(Y) = \alpha(g(X, Y)\xi - \varepsilon \eta(Y)X) + \beta(g(\phi X, Y)\xi - \delta \eta(Y)\phi X) \quad (2.5)$$

for some smooth functions α and β on \bar{M} and $\varepsilon = \pm 1, \delta = \pm 1$. For $\beta = 0, \alpha = 1$, an (ε, δ) -trans-Sasakian manifolds reduces to (ε) -Sasakian and for $\alpha = 0, \beta = 1$ it reduces to a (δ) -Kenmotsu manifolds. From (2.5) it follows that

$$\bar{\nabla}_X \xi = -\varepsilon \alpha \phi X - \beta \delta \phi^2 X. \quad (2.6)$$

for any vector field X tangent to \bar{M} .

Example of (ε, δ) -trans Sasakian manifolds

Consider the three dimensional manifold $M = \{(x, y, z) \in R^3 \mid z \neq 0\}$, where (x, y, z) are the cartesian coordinates in R^3 and let the vector fields are

$$e_1 = \frac{e^x}{z^2} \frac{\partial}{\partial x}, \quad e_2 = \frac{e^y}{z^2} \frac{\partial}{\partial y}, \quad e_3 = \frac{-(\varepsilon + \delta)}{2} \frac{\partial}{\partial z},$$

where e_1, e_2, e_3 are linearly independent at each point of M . Let g be the Riemannian metric defined by

$$g(e_1, e_1) = g(e_2, e_2) = g(e_3, e_3) = \varepsilon, \quad g(e_1, e_3) = g(e_2, e_3) = g(e_1, e_2) = 0,$$

where $\varepsilon = \pm 1$.

Let η be the 1-form defined by $\eta(X) = \varepsilon g(X, \xi)$ for any vector field X on M , let ϕ be the (1,1) tensor field defined by $\phi(e_1) = e_2, \phi(e_2) = -e_1, \phi(e_3) = 0$.

Then by using the linearity of ϕ and g , we have $\phi^2 X = -X + \eta(X)\xi$, with $\xi = e_3$.

Further $g(\phi X, \phi Y) = g(X, Y) - \varepsilon \eta(X)\eta(Y)$ for any vector fields X and Y on M . Hence for $e_3 = \xi$, the structure defines an (ε) -almost contact structure in R^3 .

Let ∇ be the Levi-Civita connection with respect to the metric g , then we have

$$2g(\nabla_X Y, Z) = Xg(Y, Z) + Yg(Z, X) - Zg(X, Y) - g(X, [Y, Z]) - g(Y, [X, Z]) + g(Z, [X, Y])$$

which is known as Koszul's formula.

We, also have

$$\bar{\nabla}_{e_1} e_3 = -\frac{(\varepsilon + \delta)}{z} e_1, \quad \bar{\nabla}_{e_2} e_3 = -\frac{(\varepsilon + \delta)}{z} e_2, \quad \bar{\nabla}_{e_1} e_2 = 0,$$

using the above relation, for any vector X on M , we have

$\bar{\nabla}_X \xi = -\varepsilon\alpha\phi X - \beta\delta\phi^2 X$, where $\alpha = \frac{1}{z}$ and $\beta = -\frac{1}{z}$. Hence (ϕ, ξ, η, g) structure defines the (ε, δ) -trans-Sasakian structure in R^3 .

III. Semi-symmetric non-metric connection

We remark that owing to the existence of the 1-form η , we can define a semi-symmetric non-metric connection $\bar{\nabla}$ in almost contact metric manifold by

$$\bar{\nabla}_X Y = \bar{\nabla}_X Y + \eta(Y)X, \tag{3.1}$$

where $\bar{\nabla}$ is the Riemannian connection with respect to g on n -dimensional Riemannian manifold and η is a 1-form associated with the vector field ξ on M defined by

$$\eta(X) = g(X, \xi). \tag{3.2}$$

[19] BY (3.1) the torsion tensor T of the connection $\bar{\nabla}$ is given by

$$T(X, Y) = \bar{\nabla}_X Y - \bar{\nabla}_Y X - [X, Y]. \tag{3.3}$$

Also, we have

$$T(X, Y) = \eta(Y)X - \eta(X)Y. \tag{3.4}$$

A linear connection $\bar{\nabla}$, satisfying (3.4) is called a semi-symmetric connection. $\bar{\nabla}$ is called a metric connection if

$$\bar{\nabla}g = 0$$

otherwise, if $\bar{\nabla}g \neq 0$, then $\bar{\nabla}$ is said to be non-metric connection. Furthermore, from (3.1), it is easy to see that

$$\begin{aligned} \bar{\nabla}_X g(Y, Z) &= (\bar{\nabla}_X g)(Y, Z) + g(\bar{\nabla}_X Y, Z) + g(Y, \bar{\nabla}_X Z) \\ &= (\bar{\nabla}_X g)(Y, Z) + \bar{\nabla}_X g(Y, Z) + \eta(Y)g(X, Z) + \eta(Z)g(X, Y) \end{aligned}$$

which implies

$$(\bar{\nabla}_X g)(Y, Z) = \eta(Y)g(X, Z) - \eta(Z)g(X, Y) \tag{3.5}$$

for all vector fields X, Y, Z on M . Therefore in view of (3.4) and (3.5) $\bar{\nabla}$ is a semi-symmetric non-metric connection.

for all $X, Y \in TM$. Now from (3.1), (2.5) and (2.6), we have

$$\begin{aligned} (\bar{\nabla}_X \phi)Y &= \alpha\{(g(X, Y)\xi - \varepsilon\eta(Y)X)\} + \beta(g(\phi X, Y))\xi \\ &\quad + (1 - \beta\delta)\eta(Y)\phi X. \end{aligned} \tag{3.6}$$

From (3.6) it follows that

$$\bar{\nabla}_X \xi = X - \varepsilon\alpha\phi X - \beta\delta\phi^2 X \tag{3.7}$$

for any vector field X tangent to \bar{M} .

Now, let M be a submanifold isometrically immersed in an (ε, δ) -trans-Sasakian manifold \bar{M} such that the structure vector field ξ of \bar{M} is tangent to submanifolds M . We denote by ξ is the 1-dimensional distribution spanned by ξ on M and by $\{\xi\}^\perp$ the complementary orthogonal distribution to ξ in TM .

For any $X \in TM$, we have $g(\phi X, \xi) = 0$. Then we have

$$\phi X = BX + CX, \tag{3.8}$$

where $BX \in \{\xi\}^\perp$ and $CX \in T^\perp M$. Thus $X \rightarrow BX$ is an endomorphism of the tangent bundle TM and $X \rightarrow CX$ is a normal bundle valued 1-form on M .

Definition. A submanifold of M of an almost contact metric manifolds \bar{M} with an (ε, δ) -trans-Sasakian metric structure (ϕ, ξ, η, g) is said to be a generalised CR-submanifold if

$$D_x^\perp = T_x M \cap \phi T_x^\perp M; \quad \text{for } x \in M$$

defines a differentiable sub-bundle of $T_x M$. Thus for $X \in D^\perp$ one has $BX = 0$.

We denote by D the complementary orthogonal sub-bundle to $D^\perp \oplus \{\xi\}$ in TM .

For any $X \in D$, $BX \neq 0$. Also we have $BD = D$.

Thus for a generalised CR-submanifold M , we have the orthogonal decomposition

$$TM = D \oplus D^\perp \oplus \{\xi\}. \quad (3.9)$$

IV. Basic Lemmas

Let M be a generalised CR-submanifold of an (ε, δ) -trans-Sasakian manifold \bar{M} . We denote by g both Riemannian metrics on \bar{M} and M .

For each $X \in TM$, we can write

$$X = PX + QX + \eta(X)\xi, \quad (4.1)$$

where PX and QX belong to the distribution D and D^\perp respectively.

For any $N \in T_x^\perp M$, we can write

$$\phi X = tN + fN, \quad (4.2)$$

where tN is the tangential part of ϕN and fN is the normal part of ϕN .

By using (2.2) we have

$$g(\phi X, CY) = g(\phi X, BY + CY) = g(\phi X, \phi Y) = g(X, Y) = 0,$$

for $X \in D_x^\perp$ and $Y \in D_x$. Therefore

$$g(\phi D_x^\perp, CD_x) = 0. \quad (4.3)$$

We denote by ν the orthogonal complementray vector bundle to $\phi D^\perp \oplus CD$ in $T^\perp M$.

Thus, we have

$$T^\perp M = \phi D^\perp \oplus CD \oplus \nu \quad (4.4)$$

Lemma 4.1. The morphism t and f satisfy

$$t(\phi D^\perp) = D^\perp \quad (4.5)$$

$$t(CD) \subset D \quad (4.6)$$

Proof. For $X \in D^\perp$ and $Y \in D$.

$$g(t\phi, Y) = g(t\phi X + f\phi Y, Y) = g(\phi^2 X, Y) = -g(X, Y) = 0$$

$$g(t\phi X, \xi) = g(\phi^2 X, \xi) = -g(\phi X, \phi \xi) = 0.$$

Therefore, $t(\phi D^\perp) \subset D^\perp$.

For $X \in D^\perp$, we have

$$-X = \phi^2 X = t\phi X + f\phi X, \text{ which implies } -X = t\phi X.$$

Consequently, $D^\perp \subset t(\phi D^\perp)$. Hence the equation (4.5) follows. The equation (4.6) is trivial.

Let M be a submanifold of a Riemannian manifold \bar{M} with Riemannian metric g . Then Gauss and Weingarten formulae are given respectively by

$$\bar{\nabla}_X Y = \nabla_X Y + h(X, Y) \quad (X, Y \in TM), \quad (4.7)$$

$$\bar{\nabla}_X N = -A_N X + \nabla_X^\perp N + \eta(N)X \quad (N \in T^\perp M), \quad (4.8)$$

where $\bar{\nabla}$, ∇ and ∇^\perp respectively the semi-symmetric non-metric, induced connection and induced normal connections in \bar{M} , M and the normal bundle $T^\perp M$ of M respectively and h is the second fundamental form related to A by

$$g(h(X, Y), N) = g(A_N X, Y) \quad (4.9)$$

for $X, Y \in TM$ and $N \in T^\perp M$.

We denote

$$u(X, Y) = \nabla_X BPY - A_{CPY} X - A_{\phi QY} X. \quad (4.10)$$

Lemma 4.2. Let M be a generalised CR-submanifold of an (ε, δ) -trans-Sasakian manifold \bar{M} with semi-symmetric non-metric connection. Then we have

$$P(u(X, Y)) - BP\nabla_X Y - Pth(X, Y) = -\alpha\eta(Y)PX \quad (4.11)$$

$$-(1 - \beta\delta)\eta(Y)PBX - 2\eta(CPY)PX,$$

$$Q(u(X, Y)) - Qth(X, Y) = -\alpha\eta(Y)QX - (1 - \beta\delta)\eta(Y)QBX \quad (4.12)$$

$$-2\eta(CPY)QX,$$

$$\eta(u(X, Y)) = \alpha g(\phi X, \phi Y) + \beta g(\phi BX, \phi Y) - 2\eta(CPY)\eta(X)\xi, \quad (4.13)$$

$$h(X, BPY) + \nabla_X^\perp CPY + \nabla_X^\perp \phi QY - CP\nabla_X Y - \phi Q\nabla_X Y - fh(X, Y) \quad (4.14)$$

$$= (1 - \beta\delta)\eta(Y)CX,$$

for $X, Y \in TM$.

Proof. For $X, Y \in TM$ by using (3.8), (4.1), (4.2), (4.7), (4.8) in (3.6), we have

$$\nabla_X PBY + h(X, BPY) - A_{CPY} X + \nabla_X^\perp CPY + \eta(CPY)X - A_{\phi QY} X + \nabla_X^\perp \phi QY$$

$$- BP\nabla_X Y - CP\nabla_X Y - \phi Q\nabla_X Y - Pth(X, Y) - Qth(X, Y) - fh(X, Y)$$

$$= \alpha \{ (g(X, Y)\xi - \varepsilon\eta(Y)X) \} + \beta (g(\phi X, Y))\xi + (1 - \beta\delta)\eta(Y)\phi X.$$

Then (4.11), (4.12), (4.13) and (4.14) are obtaining by taking the components of each vector bundles D , D^\perp , $\{\xi\}$ and $T^\perp(M)$ respectively.

Lemma 4.3. Let M be a generalised CR-submanifold of an (ε, δ) -trans-Sasakian manifold \bar{M} with semi-symmetric non-metric connection. Then we have

$$P(t\nabla_X^\perp N + A_{fN} X - \nabla_X tN) = BPA_N X - \eta(fN)PX, \quad (4.15)$$

$$Q(t\nabla_X^\perp N + A_{fN} X - \nabla_X tN) = -\eta(fN)QX, \quad (4.16)$$

$$\eta(A_{fN} X - \nabla_X tN) = -\beta g(CX, N) + \eta(fN)\eta(X)\xi, \quad (4.17)$$

$$h(X, tN) + \phi QA_N X + \nabla_X^\perp fN + CPA_N X = f\nabla_X^\perp N \quad (4.18)$$

for $X \in TM$ and $N \in T^\perp M$.

Proof. For $X \in TM$ and $N \in T^\perp M$ by using the equations (3.8), (4.1), (4.2), (4.7) and (4.8) in (3.6), we get

$$\begin{aligned}
 & P\nabla_X tN + Q\nabla_X tN + \eta(\nabla_X tN) + h(X, tN) - PA_{fN}X - \eta(fN)PX - QA_{fN}X \\
 & - \eta(fN)QX - \eta(A_{fN}X) + \nabla_X^\perp fN + \eta(fN)\eta(X)\xi + BPA_NX + CPA_NX \\
 & + \phi QA_NX - Pt\nabla_X^\perp N - Q\nabla_X^\perp N - f\nabla_X^\perp N = \beta g(CX, N)
 \end{aligned}$$

Then (4.15), (4.16), (4.17) and (4.18) are obtaining by taking the components of each vector bundles D , D^\perp , $\{\xi\}$ and $T^\perp(M)$ respectively.

Lemma 4.4. Let M be a generalised CR-submanifold of an (ε, δ) -trans-Sasakian manifold \bar{M} with semi-symmetric non-metric connection. Then we have

$$\nabla_X \xi = PX + \beta \delta X - \alpha \alpha BX, \text{ for } X \in D \tag{4.19}$$

$$h(X, \xi) = QX - \alpha \alpha CX \text{ and } (1 - \beta \delta)\eta(X) = 0, \text{ for } X \in D \tag{4.20}$$

$$\nabla_Y \xi = PY + \beta \delta Y, \text{ for } Y \in D^\perp \tag{4.21}$$

$$h(Y, \xi) = QY - \alpha \alpha \phi Y; \eta(Y)(1 - \beta \delta) = 0, \text{ for } Y \in D^\perp \tag{4.22}$$

$$\nabla_\xi \xi = P\xi \tag{4.23}$$

$$h(\xi, \xi) = Q\xi; \beta \delta = 1. \tag{4.24}$$

Proof. The proof of above lemma from (3.7) by using (3.8), (4.1) and (4.7).

Lemma 4.5 Let M be a generalised CR-submanifold of an (ε, δ) -trans-Sasakian manifold \bar{M} with semi-symmetric non-metric connection. Then we have

$$A_{\phi X} Y = A_{\phi Y} X, \text{ for } X, Y \in D^\perp. \tag{4.25}$$

Proof. By using (2.2), (2.3), (4.7) and (4.9), we get

$$\begin{aligned}
 g(A_{\phi X} Y, Z) &= g(h(Y, Z), \phi X) = g(\bar{\nabla}_X Y, \phi X) = -g(\phi \bar{\nabla}_Z Y, X) \\
 &= -g(\bar{\nabla}_{\phi Z} Y, X) = g(\phi Y, \bar{\nabla}_Z X) = g(h(Z, X), \phi Y) = g(h(X, Z), \phi Y) \\
 &= g(A_{\phi} YX, Z),
 \end{aligned}$$

for $X, Y \in D^\perp$ and $Z \in TM$. Hence the Lemma follows.

Lemma 4.6. Let M be a generalised CR-submanifold of an (ε, δ) -trans-Sasakian manifold \bar{M} with semi-symmetric non-metric connection. Then we have

$$\nabla_\xi V \in D^\perp, \text{ for } V \in D^\perp \text{ and} \tag{4.26}$$

$$\nabla_\xi W \in D, \text{ for } W \in D. \tag{4.27}$$

Proof. Let us take $X = \xi$ and $V = \phi N$ in (4.15), where $N \in \phi D$. Taking account that $tN = \phi N$, $fN = 0$ we get

$$P\nabla_\xi V = Pt\nabla_\xi^\perp N - BPA_N \xi. \tag{4.28}$$

The first relation of (4.20) gives

$$g(PA_N \xi, W) = g(A_N \xi, W) = g(h(W, \xi, N)) = -\alpha \alpha g(CW, N) + g(QW, N) = 0$$

for $W \in D$. Hence, (4.28) becomes

$$P\nabla_{\xi} V = P t \nabla_{\xi}^{\perp} N. \quad (4.29)$$

On the other hand (4.18) implies

$$h(\xi, V) = f \nabla_{\xi}^{\perp} N - \phi Q A_N \xi. \quad (4.30)$$

For $V \in D^{\perp}$, $h(\xi, V) = h(V, \xi) = -\varepsilon \alpha \phi V \in \phi D^{\perp}$, by (3.22)

Now for $X \in D^{\perp}$ by using the lemma (4.5) and of (4.9), we have

$$\begin{aligned} g(h(\xi, V), \phi X) &= g(h(V, \xi), \phi X) = g(A_{\phi X} V, \xi) = g(A_{\phi V} X, \xi) \\ &= g(h(X, \xi), \phi V) = g(h(X, \xi), -N) = -g(A_N \xi, X) = -g(\phi A_N \xi, \phi X) \\ &= -g(\phi P A_N \xi, \phi X) - g(\phi Q A_N \xi, \phi X) = -g(\phi Q A_N \xi, \phi X) \end{aligned}$$

since $CD^{\perp} \in \phi D^{\perp}$.

Therefore, $h(\xi, V) = -\phi Q A_N \xi$, which together with (4.30) implies $f \nabla_{\xi}^{\perp} N = 0$.

Hence $\nabla_{\xi}^{\perp} N \in \phi D^{\perp}$, since f is an automorphism of $CD \oplus \nu$. Thus, $t \nabla_{\xi}^{\perp} N \in D^{\perp}$ and from (4.29) it follows that

$$P\nabla_{\xi} V = 0, \quad \text{for all } V \in D^{\perp} \quad (4.31)$$

Next from (4.17), we have

$$\eta(\nabla_{\xi} V) = 0 \quad (4.32)$$

for all $V = \phi D \in D^{\perp}$, where $N \in \phi D^{\perp}$. Hence (4.26) follows from (4.31) and (4.32).

Finally using the (4.1), (4.23) and (4.26), we have

$$g(\nabla_{\xi} W, X) = g(\nabla_{\xi} W, PX)$$

for $X \in TM$ and $W \in D$. Thus we have $\nabla_{\xi} W \in D$, for $W \in D$ and this completes the proof.

Corollary 4.1. Let M be a generalised CR-submanifold of an (ε, δ) -trans-Sasakian manifold \overline{M} with semi-symmetric non-metric connection. Then we have

$$[Y, \xi] \in D^{\perp}, \quad \text{for } Y \in D^{\perp} \quad (4.33)$$

$$[X, \xi] \in D, \quad \text{for } X \in D \quad (4.34)$$

The above corollary follows immediate consequences of the Lemma (4.4) and Lemma (4.6).

V. Integrability of Distributions

Theorem 5.1. Let M be a generalised CR-submanifold of an (ε, δ) -trans-Sasakian manifold \overline{M} with semi-symmetric non-metric connection. Then the distribution D^{\perp} is always involutive if and only if

$$g([X, Y], \xi) - 2\beta\delta g(X, Y) = 0. \quad (5.1)$$

Proof. For $X, Y \in D^{\perp}$ by using (4.21), we get

$$g([X, Y], \xi) = g(\nabla_X Y, \xi) - g(\nabla_Y X, \xi)$$

$$g([X, Y], \xi) = g(X, \nabla_Y \xi) - g(Y, \nabla_X \xi) = 2\beta\delta g(X, Y). \quad (5.2)$$

On the other hand, from (4.10), we have

$$BP\nabla_X Y = -PA_{\phi Y} X - Pth(X, Y), \quad (5.3)$$

for $X, Y \in D^{\perp}$. Then using lemma (4.5), we get from equation (5.3)

$$BP[X, Y] = 0, \text{ for } X, Y \in D^{\perp}. \quad (5.4)$$

Theorem 5.2. Let M be a generalised CR-submanifold of an (ε, δ) -trans-Sasakian manifold \bar{M} with semi-symmetric non-metric connection. Then the distribution D is never involutive.

Proof. For $X, Y \in D$ by using (4.19), we have

$$\begin{aligned} g([X, Y], \xi) &= g(\nabla_X Y, \xi) - g(\nabla_Y X, \xi) \\ &= 2\alpha\delta g(Y, BX) + 2\beta\delta g(X, Y) + g(X, PY) - g(Y, PX). \end{aligned} \quad (5.5)$$

Taking $X \neq 0$ and $Y = BX$ in (5.5), it follows that D is not involutive.

Next we have the following theorem.

Theorem 5.3. Let M be a generalised CR-submanifold of an (ε, δ) -trans-Sasakian manifold \bar{M} with semi-symmetric non-metric connection. Then the distribution $D \oplus \{\xi\}$ is involutive if and only if

$$h(BX, Y) - h(X, BY) + \nabla_Y^\perp CX - \nabla_X^\perp CY \in CD \oplus \nu \quad (5.6)$$

Proof. Applying ϕ to equation (4.14) and taking component in D^\perp , we have

$$Q\nabla_X Y = -Qt(h(X, BY) + \nabla_X^\perp CPY - fh(X, Y))$$

for $X, Y \in D$.

Thus

$$Q[X, Y] = Qt(h(X, BY) - h(X, BY) + \nabla_Y^\perp CX - \nabla_X^\perp CY) \quad (5.7)$$

for $X, Y \in D$. Hence, the theorem follows from (5.7) and (4.34).

VI. Geometry of Leaves

Theorem 6.1. Let M be a generalised CR-submanifold of an (ε, δ) -trans-Sasakian manifold \bar{M} with semi-symmetric non-metric connection. Then the leaves of distribution D^\perp are totally geodesic in M if and only if

$$h(X, BZ) + \nabla_X^\perp CZ + \eta(CZ)X \in CD \oplus \nu \quad (6.1)$$

for $X \in D^\perp$ and $Z \in D \oplus \{\xi\}$.

Proof. For $X, Y \in D^\perp$ and $Z \in D \oplus \{\xi\}$ by using (2.2), (2.3), (4.7) and (4.8), we get

$$\begin{aligned} g(\bar{\nabla}_X Y, Z) &= -g(Y, \bar{\nabla}_X Z) = -g(\bar{\nabla}_X Z, Y) = -g(\phi\bar{\nabla}_X Z, \phi Y) \\ &= g((\bar{\nabla}_X \phi)Z, \phi Y) - g(\bar{\nabla}_X \phi Z, \phi Y) = -g(\bar{\nabla}_X BZ + \bar{\nabla}_X CZ, \phi Y) \\ &= -g(\nabla_X BZ + h(X, BZ) - A_{CZ}X + \eta(CZ)X + \nabla_X^\perp CZ, \phi Y) \\ &= -g(h(X, BZ) + \nabla_X^\perp CZ + \eta(CZ)X, \phi Y). \end{aligned} \quad (6.2)$$

Hence the theorem follows from the (6.2).

Theorem 6.2. Let M be a generalised CR-submanifold of an (ε, δ) -trans-Sasakian manifold \bar{M} with semi-symmetric non-metric connection. Then the distribution $D^\perp \oplus \{\xi\}$ is involutive and its leaves are totally geodesic in M if and only if

$$h(X, BY) + \nabla_X^\perp CY + \eta(CY)X \in CD \oplus \nu \quad (6.3)$$

for $X, Y \in D^\perp \oplus \{\xi\}$.

Proof. For $X, Y \in D^\perp \oplus \{\xi\}$ and $Z \in D^\perp$ by using (2.2), (2.3), (3.8), (4.7) and (4.8), we get

$$\begin{aligned} g(\bar{\nabla}_X Y, Z) &= g(\nabla_X Y, Z) = g(\phi\bar{\nabla}_X Y, \phi Z) = g(\nabla_X \phi Y, \phi Z) \\ &= g(\nabla_X BY + h(X, BY) - A_{CY}X + \eta(CY)X + \nabla_X^\perp CY, \phi Z). \end{aligned} \quad (6.4)$$

Hence, the theorem follows from the equation (6.4).

References

- [1]. Mihai, Geometry and Topology of Submanifolds, *World Scientific, Vol. VII*, (1995), 186-188.
- [2]. Mihai, Geometry and Topology of Submanifolds, *World Scientific, Vol. VIII*, (1996), 265-268.
- [3]. J. A. Oubina, New Class of almost contact metric structures, *Publ. Math. Debrecen*, 32(1985), 187-193.
- [4]. M. H. Shahid, CR -Submanifolds of trans-Sasakian manifold, *Indian J. Pure Appl. Math.* 22(1991), 1007-1012.
- [5]. M. H. Shahid, I. Mihai, Generic CR -Submanifolds of trans-Sasakian manifold, *Publ. Math. Debrecen*, 47(3), (1995), 209-218. Kumar, A. U. C. De, Generalised CR submanifolds of a trans-Sasakian manifold, *Indian. J. Pure. Appl. Math.* 32(4), 2001, 573-580. Bejancu, K. L. Duggal, Real Hypersurfaces on indefinite Kahler manifolds. *Int. Math. Math. Sci.* 16(3), (1993), 545-556.
- [6]. U. C. De, A. Sarkar, A., On \mathcal{E} -Kenmotsu manifolds, *Hadronic journal*, 32 (2009), 231-242. Bhattacharyya and B. Das, Contact CR -Submanifolds of an Indefinite trans-Sasakian manifold, *Int. J. Contemp. Math. Sci.* 6(26), (2011), 1271-1282.
- [7]. H. G. Nagaraja, C. R. Premalath, and G. Somashekar, On an (ε, δ) -trans-Sasakian manifolds, *Proced. Estonian Acced. Sci.* 1(61), 2012, 20-28. Özgür, On submanifolds of a Riemannian manifold with a semi-symmetric non-metric connection, *Kuwait J. Sci. engg.* 37(2), (2010), 17-30.
- [8]. S. Sular, C. Özgür, Generalized Sasakian space forms with semi-symmetric non-metric connection, *Proced. Estonian Acad. Sci.* 60(4), (2011), 251-257.
- [9]. M. Ahmad, C. Özgür, Hypersurfaces of almost F -paracontact Riemannian manifold endowed with semi-symmetric non-metric connection, *Result. Math.* 55 (2009), 1-10.
- [10]. M. Ahmad and J. B. Jun, On semi-invariant submanifolds of nearly Kenmotsu manifold with a semi-symmetric non-metric connection, *J. Chungcheong Math. Soc.* 23(2), (2010), 257-266.
- [11]. L. S. Das, M. Ahmad and A. Haseeb, On Semi-invariant submanifolds of a nearly Sasakian manifold with semi-symmetric non-metric connection. *J. Appl. Analysis*, 17(1), (2010), 119-130.
- [12]. M. Ahmad, S. Rahman and M. D. Siddiqi, Semi-invariant submanifolds of a nearly Sasakian manifold endowed with a semi-symmetric metric connection, *Bull. Allahabad Math. Soc.* 25, part 1, (2010), 23-33. Friedmann and J. A. Schouten, Über die geometrie der halbsymmetrischen übertragung *Math. Zeitschr.* 21 (1924), 211-223.
- [13]. K. Yano, On semi-symmetric metric connection, *Rev. Roumaine de Math. Pures et Appl.* 15, (1970) 1579-1586.
- [14]. N. S. Agashe, and M. R. Chaffle, A semi-symmetric non-metric connection of a Riemannian manifold, *Indian. J. Pure and Appl. Math.* 23, (1992), 399-409.
- [15]. J. A. Schouten, Ricci-Calculus, An Introduction to Tensor Analysis and Geometrical Applications. *Springer-Verlag, Berlin-Göttingen-Heidelberg*, 1954,