

GENERALIZED HERMITE INTERPOLATION AND SAMPLING THEOREM INVOLVING DERIVATIVES

CHANG EON SHIN

ABSTRACT. We derive the generalized Hermite interpolation by using the contour integral and extend the generalized Hermite interpolation to obtain the sampling expansion involving derivatives for band-limited functions f , that is, f is an entire function satisfying the following growth condition

$$|f(z)| \leq A \exp(\sigma|y|) \text{ for some } A, \sigma > 0 \text{ and any } z = x + iy \in \mathbb{C}.$$

1. Introduction

If p_n is a polynomial of degree n and z_0, z_1, \dots, z_n are distinct points in \mathbb{C} , then p_n can be constructed with the values $p_n(z_i), i = 0, \dots, n$ and polynomials independent of p_n . The polynomial $p_n(z)$ is called the *Lagrange interpolation* ([1]) and is given by

$$(1.1) \quad p_n(z) = \sum_{k=0}^n p_n(z_k) \frac{G_n(z)}{(z - z_k)G'_n(z_k)},$$

where $G_n(z) = \prod_{i=0}^n (z - z_i)$. Let $\sigma > 0$ and let B_σ^p be the set of entire functions such that there exists $A > 0$ satisfying

$$(1.2) \quad |f(z)| \leq A \exp(\sigma|y|) \quad \text{for any } z = x + iy \in \mathbb{C}$$

and

$$f|_{\mathbb{R}} \in L^p(\mathbb{R}).$$

The function f in B_σ^p for some $1 \leq p \leq \infty$ is called a *band-limited function* with band-limit σ . The following inclusions (p.24, [11]) hold

$$(1.3) \quad B_1 \subseteq B_p \subseteq B_q \subseteq B_\infty, \quad 1 \leq p \leq q \leq \infty.$$

Received February 16, 2002.

2000 Mathematics Subject Classification: 94A12.

Key words and phrases: generalized Hermite interpolation, sampling theorem, contour integral.

The Lagrange interpolation can be extended so that we obtain celebrated Shannon’s sampling theorem ([2], [11]) which is the following:

THEOREM 1.1. For any $f \in B_\sigma^p (1 \leq p < \infty)$,

$$(1.4) \quad f(z) = \sum_{n=-\infty}^{\infty} f(k_n) \frac{G(z)}{(z - k_n)G'(k_n)},$$

where $k_n = \frac{n\pi}{\sigma}, n \in \mathbb{Z}$,

$$G(z) = \frac{\sin \sigma z}{\sigma} = z \prod_{n=1}^{\infty} \left(1 - \frac{z^2}{k_n^2}\right) = \lim_{m \rightarrow \infty} \tilde{G}_m(z),$$

$$\tilde{G}_m(z) = z \prod_{n=1}^m \left(1 - \frac{z^2}{k_n^2}\right)$$

and the series converges uniformly on any compact subsets of \mathbb{C} .

In the expansion (1.4), k_n 's are called the *sampling points*, the sampling frequency or the Nyquist rate ([11]) is $\frac{\sigma}{\pi}$ and distance between two nearest sampling points is $\frac{\pi}{\sigma}$.

D. A. Linden [4] and M. D. Rawn [9] generalized Shannon’s sampling theorem by expanding $f \in B_\sigma^2$ with values of f and its derivatives. M. D. Rawn [9] used the Riesz basis to obtain the sampling expansion of f . The aim of this paper is to derive a sampling expansion for $f \in B_\sigma^p (1 \leq p \leq \infty)$ in an easier way by using the contour integral and express f in terms of values of f and derivatives at sampling points. The sampling expansion of f in this paper is simpler than Linden’s and Rawn’s results. Moreover, we regard the sampling expansion of f as a generalization of generalized Hermite interpolation.

Let $p_{\bar{N}}(z)$ be a polynomial of degree \bar{N} , $z_1 \cdots, z_n$ be n distinct points, $\alpha_1, \cdots, \alpha_n$ be n positive integers and $\bar{N} = \alpha_1 + \cdots + \alpha_n - 1$. The polynomial $p_{\bar{N}}(z)$ can be constructed with the values $p_{\bar{N}}^{(i)}(z_j), i = 0, \cdots, \alpha_i - 1, j = 1, \cdots, n$, and some polynomials independent of $p_{\bar{N}}$. The polynomial $p_{\bar{N}}(z)$ is called the *generalized Hermite interpolation* ([1], [5]).

In this paper, using the contour integral we derive the generalized Hermite interpolation $p_{\bar{N}}(z)$ which is given by

$$(1.5) \quad p_{\bar{N}}(z) = \sum_{i=1}^n \sum_{j=0}^{\alpha_i-1} \sum_{k=0}^{\alpha_i-1-j} p_{\bar{N}}^{(j)}(z_i) \frac{w(z)}{j!k!} (z - z_i)^{-\alpha_i+j+k} \frac{d^k}{dz^k} \left[\frac{(z - z_i)^{\alpha_i}}{w(z)} \right]_{z=z_i},$$

where $w(z) = \prod_{i=1}^n (z - z_i)^{\alpha_i}$. Note that the expression of the generalized Hermite interpolation in page 37 of [1] is incorrect. Clearly the Lagrange interpolation is the special case of the generalized Hermite interpolation. In fact, if $\alpha_i = 1, i = 1, \dots, n$ and $\bar{N} = n - 1$, then the generalized Hermite interpolation $p_{\bar{N}}$ in (1.5) equals the Lagrange interpolation $p_n(z)$ in (1.1).

We also extend the generalized Hermite interpolation to obtain the sampling expansion involving values of a function and its derivatives at sampling points for entire functions f satisfying

$$(1.6) \quad (1 + |x|)|f(z)| \leq A \exp(\sigma|y|) \text{ for some } A > 0, \sigma > 0$$

and any $z = x + iy \in \mathbb{C}$.

For $N \in \mathbb{N}$ and any entire function f satisfying the growth condition (1.6), we obtain the following sampling expansion

$$(1.7) \quad f(z) = \sum_{n \in \mathbb{Z}} \sum_{j=0}^{N-1} \sum_{k=0}^{N-1-j} f^{(j)}(\beta_n) \frac{(\sin \sigma z/N)^N}{j!k!}$$

$$\times (z - \beta_n)^{-N+j+k} \frac{d^k}{dz^k} \left[\left(\frac{z - \beta_n}{\sin \sigma z/N} \right)^N \right]_{z=\beta_n},$$

where $\beta_n = \frac{Nn\pi}{\sigma}, n \in \mathbb{Z}$ and the series converges uniformly on any compact subsets of \mathbb{C} . In the expansion (1.7) the set of sampling points is $\{\beta_n : n \in \mathbb{Z}\}$ and the distance between two nearest sampling points is $\frac{N\pi}{\sigma}$, whereas distance between two nearest sampling points in Shannon's sampling theorem is $\frac{\pi}{\sigma}$. However, we need the values of $f, f', \dots, f^{(N-1)}$ at each sampling point. Note that when $N = 1$, the expansion (1.7) equals the expansion (1.4).

Moreover, we show that for any $f \in B_{\sigma}^p (1 \leq p \leq \infty)$ and $\sigma' > \sigma$, the following sampling expansion holds

$$(1.8) \quad f(z) = \sum_{n \in \mathbb{Z}} \sum_{j=0}^{N-1} \sum_{k=0}^{N-1-j} f^{(j)}(\tilde{\beta}_n) \frac{(\sin \sigma' z/N)^N}{j!k!}$$

$$\times (z - \tilde{\beta}_n)^{-N+j+k} \frac{d^k}{dz^k} \left[\left(\frac{z - \tilde{\beta}_n}{\sin \sigma' z/N} \right)^N \right]_{z=\tilde{\beta}_n},$$

where $\tilde{\beta}_n = \frac{Nn\pi}{\sigma'}, n \in \mathbb{Z}$ and the series converges uniformly on any compact subsets of \mathbb{C} .

2. Generalized Hermite interpolation

Let $p_{\bar{N}}$ be a polynomial. Then $p_{\bar{N}}$ can be expressed in terms of values of $p_{\bar{N}}$, its derivatives and some polynomials independent of $p_{\bar{N}}$.

THEOREM 2.1 (Generalized Hermite Interpolation). *Let $p_{\bar{N}}$ be a polynomial of degree \bar{N} and let z_1, \dots, z_n be n distinct points in \mathbb{C} , $\alpha_1, \dots, \alpha_n$ be n positive integers and $\bar{N} = \alpha_1 + \dots + \alpha_n - 1$. Then the polynomial $p_{\bar{N}}(z)$ is given by*

$$(2.1) \quad p_{\bar{N}}(z) = \sum_{i=1}^n \sum_{j=0}^{\alpha_i-1} \sum_{k=0}^{\alpha_i-1-j} p_{\bar{N}}^{(j)}(z_i) \frac{w(z)}{j!k!} (z-z_i)^{-\alpha_i+j+k} \frac{d^k}{dz^k} \left[\frac{(z-z_i)^{\alpha_i}}{w(z)} \right]_{z=z_i},$$

where $w(z) = \prod_{i=1}^n (z-z_i)^{\alpha_i}$. The polynomial $p_{\bar{N}}(z)$ is called the generalized Hermite interpolation.

PROOF. Let z be a complex number distinct from z_i , $i = 1, \dots, n$. We define a function g on \mathbb{C} by

$$g(\zeta) = \frac{p_{\bar{N}}(\zeta)}{(\zeta-z)w(\zeta)}.$$

The function g is a meromorphic function having a simple pole at z and poles of order α_i at z_i , $i = 1, \dots, n$. Let C_n be a circular path centered at the origin with radius n . We denote by $\text{Res}(g : \bar{z})$ the residue of g at \bar{z} . Since $|p_{\bar{N}}(\zeta)| = O(|\zeta|^{\bar{N}})$, $|w(\zeta)| = O(|\zeta|^{\bar{N}+1})$ and $|g(\zeta)| = O(|\zeta|^{-2})$ as $|\zeta| \rightarrow \infty$, we can easily see that

$$\lim_{n \rightarrow \infty} \int_{C_n} g(\zeta) d\zeta = 0$$

and

$$(2.2) \quad \text{Res}(g : z) + \sum_{i=1}^n \text{Res}(g : z_i) = 0.$$

The residue at each pole is the following:

$$\text{Res}(g : z) = \frac{p_{\bar{N}}(z)}{w(z)}$$

and for $i = 1, \dots, n$, by Leibniz formula

$$\begin{aligned}
 \text{Res}(g : z_i) &= \lim_{\zeta \rightarrow z_i} \frac{1}{(\alpha_i - 1)!} \frac{d^{\alpha_i - 1}}{d\zeta^{\alpha_i - 1}} [(\zeta - z_i)^{\alpha_i} g(\zeta)] \\
 &= \lim_{\zeta \rightarrow z_i} \frac{(-1)}{(\alpha_i - 1)!} \frac{d^{\alpha_i - 1}}{d\zeta^{\alpha_i - 1}} \left[\frac{(\zeta - z_i)^{\alpha_i}}{w(\zeta)} \frac{p_{\bar{N}}(\zeta)}{(z - \zeta)} \right] \\
 &= \lim_{\zeta \rightarrow z_i} \frac{(-1)}{(\alpha_i - 1)!} \sum_{j=0}^{\alpha_i - 1} \binom{\alpha_i - 1}{j} p_{\bar{N}}^{(j)}(\zeta) \\
 &\quad \times \frac{d^{\alpha_i - 1 - j}}{d\zeta^{\alpha_i - 1 - j}} \left[\frac{(\zeta - z_i)^{\alpha_i}}{w(\zeta)} \frac{1}{(z - \zeta)} \right] \\
 &= \lim_{\zeta \rightarrow z_i} \frac{(-1)}{(\alpha_i - 1)!} \sum_{j=0}^{\alpha_i - 1} \binom{\alpha_i - 1}{j} p_{\bar{N}}^{(j)}(\zeta) \sum_{k=0}^{\alpha_i - 1 - j} \binom{\alpha_i - 1 - j}{k} \\
 &\quad \times (\alpha_i - 1 - j - k)! (z - \zeta)^{-\alpha_i + j + k} \frac{d^k}{d\zeta^k} \frac{(\zeta - z_i)^{\alpha_i}}{w(\zeta)} \\
 &= \sum_{j=0}^{\alpha_i - 1} \sum_{k=0}^{\alpha_i - 1 - j} \frac{(-1)}{j!k!} p_{\bar{N}}^{(j)}(z_i) (z - z_i)^{-\alpha_i + j + k} \\
 &\quad \times \frac{d^k}{d\zeta^k} \left[\frac{(\zeta - z_i)^{\alpha_i}}{w(\zeta)} \right]_{\zeta = z_i}.
 \end{aligned}$$

In view of (2.2) we have that

$$(2.3) \quad p_{\bar{N}}(z) = \sum_{i=1}^n \sum_{j=0}^{\alpha_i - 1} \sum_{k=0}^{\alpha_i - 1 - j} p_{\bar{N}}^{(j)}(z_i) \frac{w(z)}{j!k!} (z - z_i)^{-\alpha_i + j + k} \frac{d^k}{d\zeta^k} \left[\frac{(\zeta - z_i)^{\alpha_i}}{w(\zeta)} \right]_{\zeta = z_i}$$

for $z \neq z_i, i = 1, \dots, n$. Since both sides of (2.3) are polynomials and the equality (2.3) holds for $z \neq z_i, i = 1, \dots, n$, (2.3) holds for any $z \in \mathbb{C}$. □

The general form of the generalized Hermite interpolation is usually given in the following way.

COROLLARY 2.2. *Let z_1, \dots, z_n be n distinct points in \mathbb{C} , $\alpha_1, \dots, \alpha_n$ be n positive integers and $\bar{N} = \alpha_1 + \dots + \alpha_n - 1$. The polynomial $p_{\bar{N}}$ of*

degree \bar{N} satisfying

$$(2.4) \quad \begin{matrix} p_{\bar{N}}(z_1) = r_{10}, & p'_{\bar{N}}(z_1) = r_{11}, & \cdots & p_{\bar{N}}^{(\alpha_1-1)}(z_1) = r_{1\alpha_1-1} \\ \vdots & \vdots & \vdots & \vdots \\ p_{\bar{N}}(z_n) = r_{n0}, & p'_{\bar{N}}(z_n) = r_{n1}, & \cdots & p_{\bar{N}}^{(\alpha_n-1)}(z_n) = r_{n\alpha_n-1} \end{matrix}$$

is given by

$$(2.5) \quad p_{\bar{N}}(z) = \sum_{i=1}^n \sum_{j=0}^{\alpha_i-1} \sum_{k=0}^{\alpha_i-1-j} r_{ij} \frac{w(z)}{j!k!} (z - z_i)^{-\alpha_i+j+k} \frac{d^k}{dz^k} \left[\frac{(z - z_i)^{\alpha_i}}{w(z)} \right]_{z=z_i}.$$

3. Sampling expansion of entire functions

In this section we extend the generalized Hermite interpolation derived in Section 2 to obtain the sampling expansion involving derivatives for entire functions.

THEOREM 3.1. *Let $\sigma > 0, N \in \mathbb{N}$ and let f be an entire function such that there exists $A > 0$ satisfying*

$$(3.1) \quad (1 + |x|)|f(z)| \leq A \exp(\sigma|y|) \quad \text{for any } z = x + iy \in \mathbb{C}.$$

Then f can be expanded as

$$(3.2) \quad f(z) = \sum_{n \in \mathbb{Z}} \sum_{j=0}^{N-1} \sum_{k=0}^{N-1-j} f^{(j)}(\beta_n) \frac{(\sin \sigma z/N)^N}{j!k!} \times (z - \beta_n)^{-N+j+k} \frac{d^k}{dz^k} \left[\left(\frac{z - \beta_n}{\sin \sigma z/N} \right)^N \right]_{z=\beta_n},$$

where $\beta_n = \frac{Nn\pi}{\sigma}, n \in \mathbb{Z}$, and the series converges uniformly on any compact subsets of \mathbb{C} .

PROOF. Let $M > 1$ and

$$S = \{z \in \mathbb{C} : |z| \leq M \quad \text{and} \quad z \neq \beta_n, \quad n \in \mathbb{Z}\}.$$

Choose $z \in S$ and define a function g on the complex plane by

$$g(\zeta) = \frac{f(\zeta)}{(\zeta - z)(\sin \frac{\sigma \zeta}{N})^N}.$$

Let $\tau_n = \frac{N\pi}{\sigma}(n + \frac{1}{2})$ and let C_n be a rectangular path whose vertices are $\pm\tau_n \pm i\tau_n$. Choose $N_1 \in \mathbb{N}$ so large that $\tau_{N_1} \geq 2M$. Let $n \geq N_1$ and for $\delta > 0$ let $\alpha_\delta = \frac{e^{2\delta}-1}{2e^{2\delta}}$. Observing that for $y \geq \delta > 0, \xi, \eta \in \mathbb{R}$

$$\sinh y \geq \alpha_\delta e^y,$$

$$\begin{aligned} |\sin \frac{\sigma}{N}(\xi + i\tau_n)| &= \sqrt{\sin^2 \frac{\sigma}{N}\xi + \sinh^2 \pi(n + \frac{1}{2})} \\ &\geq \alpha_{\pi(N_1 + \frac{1}{2})} \exp(\pi(n + \frac{1}{2})), \end{aligned}$$

$$|\sin \frac{\sigma}{N}(\tau_n + i\eta)| = \sqrt{\sin^2 \pi(n + \frac{1}{2}) + \sinh^2 \frac{\sigma}{N}\eta} \geq 1$$

and for $|\eta| \geq \delta$,

$$|\sin \frac{\sigma}{N}(\tau_n + i\eta)| \geq \alpha_{\frac{\alpha\delta}{N}} \exp(\frac{\sigma}{N}|\eta|),$$

by a direct computation or following the proof of Theorem 2.1 of [10] we can see that

$$\lim_{n \rightarrow \infty} \int_{C_n} g(\zeta) d\zeta = 0$$

uniformly on S . Thus the sum of residues of g equals zero. Note that the function g has a simple pole at $\zeta = z$ and poles of order N at $\zeta = \beta_n, n \in \mathbb{Z}$. Following the computation in Section 2, we have

$$\text{Res}(g : z) = \lim_{\zeta \rightarrow z} (\zeta - z)g(\zeta) = \frac{f(z)}{(\sin \frac{\sigma}{N}z)^N},$$

and for each $n \in \mathbb{Z}$

$$\begin{aligned} \text{Res}(g : \beta_n) &= \lim_{\zeta \rightarrow \beta_n} \frac{1}{(N-1)!} \frac{d^{N-1}}{d\zeta^{N-1}} [(\zeta - \beta_n)^N g(\zeta)] \\ &= (-1) \sum_{j=0}^{N-1} \sum_{k=0}^{N-1-j} \frac{1}{j!k!} f^{(j)}(\beta_n) (z - \beta_n)^{-N+j+k} \\ &\quad \times \frac{d^k}{d\zeta^k} \left[\left(\frac{\zeta - \beta_n}{\sin \frac{\sigma}{N}\zeta} \right)^N \right]_{\zeta=\beta_n}. \end{aligned}$$

Since $\text{Res}(g : z) + \sum_{n \in \mathbb{Z}} \text{Res}(g : \beta_n) = 0$, it follows that for any $z \in S$

$$(3.3) \quad f(z) = \sum_{n \in \mathbb{Z}} \sum_{j=0}^{N-1} \sum_{k=0}^{N-1-j} f^{(j)}(\beta_n) \frac{(\sin \frac{\sigma}{N} z)^N}{j!k!} (z - \beta_n)^{-N+j+k} \\ \times \frac{d^k}{d\zeta^k} \left[\left(\frac{\zeta - \beta_n}{\sin \frac{\sigma}{N} \zeta} \right)^N \right]_{\zeta=\beta_n},$$

where the series converges uniformly on S . Since $M > 1$ is an arbitrary number, the series converges uniformly on any compact subsets of $\mathbb{C} \setminus \cup_{n \in \mathbb{Z}} \{\beta_n\}$.

We next show that the equality (3.3) holds for each $z = \beta_n, n \in \mathbb{Z}$. For convenience, we write $h(z) = (\sin \frac{\sigma}{N} z)^N$. Let $F(z)$ be the right side of (3.3). Then $f(z) = F(z)$ for $z \neq \beta_n, n \in \mathbb{N}$. We show that for any $n \in \mathbb{Z}, f(\beta_n) = F(\beta_n)$. Since $h(z)$ has zeros at each β_n with multiplicity N , the function defined by

$$\tilde{h}_n(z) = \begin{cases} \frac{h(z)}{(z-\beta_n)^N} & \text{if } z \neq \beta_n, \\ \frac{h^{(N)}(\beta_n)}{N!} & \text{if } z = \beta_n, \end{cases}$$

is an entire function and $\tilde{h}_n(\beta_n) \neq 0$. Thus $\frac{1}{\tilde{h}_n(z)}$ is analytic on some neighborhood of β_n and for any $\ell \in \mathbb{N}$ and $1 \leq k \leq N - 1$,

$$\lim_{z \rightarrow \beta_n} (z - \beta_n)^\ell \frac{d^k}{dz^k} \frac{1}{\tilde{h}_n(z)} = 0.$$

Since for $m \neq n, h(\beta_n)(\beta_n - \beta_m)^{-N+j+k} = 0$, we have

$$F(\beta_n) = \lim_{z \rightarrow \beta_n} \sum_{j=0}^{N-1} \sum_{k=0}^{N-1-j} \frac{h(z)}{j!k!} f^{(j)}(\beta_n) (z - \beta_n)^{-N+j+k} \frac{d^k}{dz^k} \frac{1}{\tilde{h}_n(z)} \\ = \lim_{z \rightarrow \beta_n} \sum_{j=0}^{N-1} \sum_{k=0}^{N-1-j} \frac{\tilde{h}_n(z)}{j!k!} f^{(j)}(\beta_n) (z - \beta_n)^{j+k} \frac{d^k}{dz^k} \frac{1}{\tilde{h}_n(z)} \\ = \tilde{h}_n(\beta_n) f(\beta_n) \frac{1}{\tilde{h}_n(\beta_n)} = f(\beta_n)$$

since when $j \neq 0$ or $k \neq 0$, the limits vanish. Hence the equality (3.3) holds for any $z \in \mathbb{C}$ and the right side of (3.3) converges uniformly on any compact subsets of \mathbb{C} . □

Following proof of the above theorem, we can replace the growth condition (3.1) by (3.5) and obtain the same result.

REMARK 3.2. Let $\sigma > 0, N \in \mathbb{N}$ and let f be an entire function such that there exists $A > 0$ satisfying

$$(3.4) \quad (1 + |y|)|f(z)| \leq A \exp(\sigma|y|) \quad \text{for any } z = x + iy \in \mathbb{C}.$$

Then f can be expanded as (3.2).

Next, for any $f \in B_\sigma^p, 1 \leq p \leq \infty$, we can expand f into a sampling series of the form (3.2).

THEOREM 3.3. Let $\sigma' > \sigma$ and let f be a band-limited function, that is, there exists $A > 0$ satisfying

$$(3.5) \quad |f(z)| \leq A \exp(\sigma|y|) \quad \text{for any } z = x + iy \in \mathbb{C}.$$

The following sampling expansion holds

$$(3.6) \quad f(z) = \sum_{n \in \mathbb{Z}} \sum_{j=0}^{N-1} \sum_{k=0}^{N-1-j} f^{(j)}(\tilde{\beta}_n) \frac{(\sin \sigma' z/N)^N}{j!k!} (z - \tilde{\beta}_n)^{-N+j+k} \\ \times \frac{d^k}{dz^k} \left[\left(\frac{z - \tilde{\beta}_n}{\sin \sigma' z/N} \right)^N \right]_{z=\tilde{\beta}_n},$$

where $\tilde{\beta}_n = \frac{Nn\pi}{\sigma'}, n \in \mathbb{Z}$ and the series converges uniformly on any compact subsets of \mathbb{C} .

PROOF. Let $\sigma' > \sigma$. Since f satisfies (3.5), there exists $A' > 0$ satisfying

$$(1 + |y|)|f(z)| \leq A' \exp(\sigma'|y|) \quad \text{for any } z = x + iy \in \mathbb{C}.$$

By Remark 3.2, the sampling expansion (3.6) holds and the series converges uniformly on any compact subsets of \mathbb{C} . \square

References

- [1] P. J. Davis, *Interpolation and Approximation*, Blaisdell Publishing Company, New York, 1963
- [2] H. R. Higgins, *Sampling Theory in Fourier and Signal Analysis: Foundations*, Oxford Science Publications, 1996
- [3] D. L. Jagerman and L. J. Fogel, *Some general aspects of the sampling theorem*, I.R.E. Trans. Inform. Theory **IT-2** (1956), 139–146.
- [4] D. A. Linden and N. M. Abramson, *A generalization of the sampling theorem*, Inform. Control **3** (1960), 26–31
- [5] G. G. Lorentz, K. Jettler, S. D. Riemenschneider, *Birkhoff Interpolation*, Addison-Wesley, 1983
- [6] F. A. Marvasti, *Nonuniform Sampling: Theory and Practice*, Plenum Pub. Corp., 2001

- [7] A. Nathan, *On sampling a function and its derivatives*, Inform. Control **22** (1973), 172–182
- [8] Z. A. Piranashvili, *On the problem of interpolation of stochastic processes*, Theor. Probability Appl. **12** (1967), 647–657
- [9] M. D. Rawn, *A Stable Nonuniform Sampling Expansion Involving Derivatives*, IEEE Trans. Inform. Theory **IT-35** No. 6 (1989), 1223–1227
- [10] C. E. Shin, S. Y. Chung, D. Kim, *General Sampling Theorem Using Contour Integral*, submitted
- [11] A. I. Zayed, *Advances in Shannon's Sampling Theory*, CRC Press, Boca Raton, FL, 1993

Department of Mathematics
Sogang University
Seoul 121-742, Korea
E-mail: shine@ccs.sogang.ac.kr