Generalized para-Bose states

A. S. Pereira, 1, 2, * A. S. Lemos, 2, † and F. A. Brito^{2, 3, ‡}

¹Instituto Federal do Pará, 68600-000 Bragança, Pará, Brazil

²Departamento de Física, Universidade Federal de Campina Grande,
Caixa Postal 10071, 58429-900 Campina Grande, Paraíba, Brazil

³Departamento de Física, Universidade Federal da Paraíba,
Caixa Postal 5008, 58051-970 João Pessoa, Paraíba, Brazil

Abstract

In this paper, we construct integrals of motion in a para-Bose formulation for a general time-dependent quadratic Hamiltonian, which, in its turn, commutes with the reflection operator. In this context, we obtain generalizations for the squeezed vacuum states (SVS) and coherent states (CS) in terms of the Wigner parameter. Furthermore, we show that there is a completeness relation for the generalized SVS owing to the Wigner parameter. In the study of the probability transition, we found that the displacement parameter acts as a transition parameter by allowing access to odd states, while the Wigner parameter controls the dispersion of the distribution. We show that the Wigner parameter is quantized by imposing that the vacuum state has even parity. We apply the general results to the case of the time-independent para-Bose oscillator and find that the mean values of the coordinate and momentum have an oscillatory behavior similarly to the simple harmonic oscillator, while the standard deviation presents corrections in terms of the squeeze, displacement, and Wigner parameters.

PACS numbers: 03.65.Sq, 03.65.Fd, 03.65.Ca

Keywords: Coherent states, squeezed states, integrals of motion, Wigner-Heisenberg algebra.

^{*}Electronic address: alfaspereira@gmail.com

[†]Electronic address: adiellemos@gmail.com

[‡]Electronic address: fabrito@df.ufcg.edu.br

I. INTRODUCTION

In 1926 Schrödinger obtained quantum states to harmonic oscillator, which allowed for a semiclassical description [1]. In the 1960s, with the advent of the laser, these states were rediscovered and widespread in the works of Glauber-Sudarshan-Klauber [2–4] by showing a semiclassical description of electromagnetic radiation, and where the term "coherent states" (CS) was presented for the first time. These states for the harmonic oscillator can be obtained in three equivalent ways: I - as eigenstates of the annihilation operator; II - by the action of a unitary displacement operator acting in the vacuum state; III - as minimum uncertainty states. The minimization of the uncertainty relation coincides with the value calculated in the vacuum state of the harmonic oscillator, which presents the same value for the standard deviation in both position and momentum. The states satisfying these properties are called canonical CS. In addition, these states have properties of continuity and form an overcomplete basis, which guarantees a prominent role in modern quantum mechanics, with various applications ranging from quantum optics [5, 6], quantum computing [7], and mathematical physics [8].

In its turn, squeezed states (SS) form a class of nonclassical states, which minimize the uncertainty relation [9–11]. At the same time, these states offer the possibility of reducing the standard deviation of a physical quantity to a value less than that calculated in the vacuum state at the expense of increasing the standard deviation of other physical quantity [12, 13]. Thus, these states have great potential for application, for example, in quantum information [14] and for the detectors of gravitational waves [15–17]. Usually, these states are constructed by the action of the unitary squeeze operator on an arbitrary state. In particular, the acting of the squeeze operator on the vacuum state leads to the squeezed vacuum states (SVS) [18].

Generalizing quantum states provides a deeper understanding of corresponding systems since they sometimes bring additional degrees of freedom. In this sense, there is a wide range of publications in the literature that seeks to generalize CS to be able to describe systems in addition to the harmonic oscillator, such as the time-dependent quadratic systems [19, 20], systems with a given Lie group [21, 22], some non-trivial generalizations [23], and squeezed coherent states (SCS) [24, 25]. On the other hand, CS can be generalized by considering deformations in the canonical commutation relation and thus allow us to study a wide range

of relevant problems. For instance, the implications of the gravitational effects in quantum mechanics [26–28], q- and f-deformed oscillators study which is applied, mainly, to quantum optics [29–31], and to study systems with singularity as the Calogero-like model via Wigner-Heisenberg algebra (WHA) (or R-deformed Heisenberg algebra) [32, 33]. In particular, we are interested in generalizing CS and SVS via WHA.

The WHA originated from the quantization proposed by Wigner [34], which generalizes the canonical commutation relation. In particular, this algebra is generated by the creation-annihilation operators and the reflection operator (parity operator), which satisfies commutation and anti-commutation relations [35]. In this formulation, it is possible to obtain a self-adjoint momentum operator on the semi-axis in terms of a parameter that deforms the canonical commutation relation [36, 37]. Furthermore, the WHA obeys the trilinear commutation and anti-commutation relations, which characterize the para-Bose operators [38]. On its turn, this algebra leads to the parastatistical description of physical systems in terms of a deformation parameter, which corresponds to generalization of the Bose-Einstein and Fermi-Dirac statistics [39–41].

According to the order of the deformation parameter, also known as the order of statistics, it is possible to describe para-Bose or para-Fermi particles, see also [42]. Although parastatistics cannot be applied to the particles described by the standard model [43], their distinctive approach has allowed presenting promising proposals, for instance, in the conjecture of candidate particles to explain dark matter [44], in paraquark models description [45], in the study of thermodynamics properties of para-Bose systems [46], and in optical physics by simulation of para-Fermi oscillators [47]. By considering the para-Bose formulation, we will construct integrals of motion, i.e., operators that commute with the Schrödinger's operator [48] in the form of a Bogoliubov transformation [49].

In the para-Bose formulation [50], the CS were obtained as eigenstates of the annihilation operator that satisfy the characteristic commutation relation of the algebra [51] – see [38, 52, 53]. Recently, some works have paid attention to this topic, for instance, the non-linear CS [54], the construction of new types of para-Bose states [55–57], and the study of "Schrödinger cat states" [58]. In this context, we will obtain the time-dependent generalized CS and SVS via integrals of motion. These states can be applied to describe Calogero-like models [59]. In its turn, the Calogero model and its generalizations are able to describe several physical phenomena [60], such as the Hall effect [61, 62], anyons [63], and fluctua-

tions in mesoscopic systems [64], for instance. Furthermore, our construction has potential application to describe the recent proposed experimental realization of para-particles [65, 66].

This paper is structured as follows. In Sec. II, the integrals of motion method in the context of WHA will be formulated. In Sec. III, the time-dependent generalized SVS in terms of the time-independent para-Bose number states will be constructed. In turn, the completeness relation and probability transition plots will be obtained. In Sec. IV, a generalization of the time-dependent CS in terms of the Wigner, squeeze and displacement parameters will be constructed. Then, the probability transition graph will be shown, and the mean values and the uncertainty relations will be calculated. In Sec. V, the generalized CS in the coordinate representation will be considered. We apply the general result to the particular case of the para-Bose oscillator in Sec. VI. Then, the concluding remarks are presented in Sec. VII.

II. INTEGRALS OF MOTION VIA WHA

The motion integral method consists of building a time-dependent operator, which commutes with the Schrödinger's operator. In turn, the eigenstates of this operator are obtained and imposed to satisfy the Schrödinger's equation. This technique introduced by Lewis and Reisenfield to study the time-dependent harmonic oscillator [67] proved to be useful in several other problems, e.g., in the study of time-dependent quadratic Hamiltonians in one dimension [68–70] and for multidimensional systems [71], Dirac equation [72], and supersymmetric quantum mechanics [73]. Here, our contribution will be to reformulate the integrals of motion method according to the WHA. This algebraic formulation allows studying time-dependent systems with a singularity at the origin characterized by centrifugal potentials.

The WHA is composed by the annihilation \hat{a} , creation \hat{a}^{\dagger} and reflection \hat{R} operators, satisfying commutation $\left(\left[\hat{b},\hat{c}\right]=\hat{b}\hat{c}-\hat{c}\hat{b}\right)$ and anti-commutation $\left(\left\{\hat{b},\hat{c}\right\}=\hat{b}\hat{c}+\hat{c}\hat{b}\right)$ relations,

$$[\hat{a}, \hat{a}^{\dagger}] = 1 + \nu \hat{R}, \quad \{\hat{R}, \hat{a}\} = 0 = \{\hat{R}, \hat{a}^{\dagger}\}, \quad \hat{R}^2 = 1,$$
 (1)

where ν is the Wigner parameter (deformation parameter) related to the fundamental energy level ε of the para-Bose oscillator given by

$$\nu = 2\varepsilon - 1. \tag{2}$$

Notice that by setting $\nu = 0 \Rightarrow \varepsilon = 1/2$ we recover the canonical commutation relation.

The operators \hat{a} and \hat{a}^{\dagger} also obey the trilinear commutation and anti-commutation relation

$$\left[\left\{\hat{a}, \hat{a}^{\dagger}\right\}, \hat{a}\right] = -2\hat{a}, \quad \left[\left\{\hat{a}, \hat{a}^{\dagger}\right\}, \hat{a}^{\dagger}\right] = 2\hat{a}^{\dagger}, \tag{3}$$

which characterize the para-Bose operators [35, 38].

For the sake of simplicity, we will consider a Hamiltonian which commutes with the reflection operator, and in that way, the eigenstates of \hat{H} can be even or odd. In this sense, the most general form for a one-dimensional time-dependent quadratic Hamiltonian is given by

$$\hat{H} = \frac{1}{2}\hbar \left(\alpha^* \hat{a}^2 + \alpha \hat{a}^{\dagger 2}\right) + \frac{1}{2}\hbar\beta \left(\hat{a}^{\dagger} \hat{a} + \hat{a} \hat{a}^{\dagger}\right) + \hbar\delta,\tag{4}$$

where $\alpha = \alpha(t)$, $\beta = \beta(t)$ and $\delta = \delta(t)$ are time-dependent functions and the signs \dagger and * denote Hermitian and complex conjugation, respectively. From the hermiticity condition $\hat{H} = \hat{H}^{\dagger}$, we have that β and δ must be real functions. By assuming the condition $\beta > |\alpha|$, we get that the Hamiltonian (4) is positive definite, i.e., it can be written in the form of an oscillator-type Hamiltonian.

In its turn, the quantum states $|\Psi\rangle$ which describe the time evolution of the system should satisfy the Schrödinger's equation

$$\hat{\Lambda} |\Psi\rangle = 0,$$

$$\hat{\Lambda} = \frac{i}{\hbar} \hat{H} + \partial_t, \quad \partial_t = \frac{\partial}{\partial t},$$
(5)

where $\hat{\Lambda}$ is the Schrödinger's operator.

Let us consider a time-dependent operator $\hat{A} = \hat{A}(t)$, as a linear combination of the \hat{a} and \hat{a}^{\dagger} operators, in the form

$$\hat{A} = f\hat{a} + g\hat{a}^{\dagger} + \varphi, \tag{6}$$

where f = f(t), g = g(t) and $\varphi = \varphi(t)$ are time-dependent complex functions. The Eq. (6) is the well-known Bogoliubov's transformation; those coefficients have been studied in Ref. [49].

In order to \hat{A} to be an integral of motion, it has to commute with the Schrödinger's operator (5), that is

$$\hat{A} = \left[\hat{\Lambda}, \hat{A}\right] = 0, \quad \hat{A} = \frac{d\hat{A}}{dt}.$$
(7)

Substituting the Eqs. (4), (5) and (6) into (7), we obtain the following equations for the functions f, g and φ :

$$\dot{f} = i \left(\beta f - \alpha^* g\right), \quad \dot{g} = i \left(f \alpha - \beta g\right), \quad \dot{\varphi} = 0,$$
 (8)

where $\varphi(t) = \varphi_0$ is a constant function for any instant of time.

The commutator between \hat{A} and \hat{A}^{\dagger} reads

$$\left[\hat{A}, \hat{A}^{\dagger}\right] = \mu \left(1 + \nu \hat{R}\right), \ \mu = |f|^2 - |g|^2 = |f_0|^2 - |g_0|^2, \text{ for arbitrary } t,$$
 (9)

where $f_{0}=f\left(0\right)$ and $g_{0}=g\left(0\right)$ are initial conditions.

It follows from (6) and (9), that

$$\hat{a} = \frac{f^* \hat{A} - g \hat{A}^{\dagger} + u}{\mu}, \ u = g \varphi_0^* - f^* \varphi_0.$$
 (10)

Notice that $|f| \neq |g|$ for all time ensures that \hat{a} is well defined for all t. It is worth highlighting that the μ -parameter will be useful to study special cases in which the Hamiltonian presents linear terms. In addition, the μ -parameter will be useful to replace the functions f and g with the squeeze and displacement parameters in the constructed states.

III. TIME-DEPENDENT GENERALIZED SVS

The SVS are pure states known to be one of the most useful nonclassical states, see for instance, [10]. In addition, these states make it possible to obtain the standard deviation of a physical quantity less than its value in the vacuum state. In this section, we will obtain the time-dependent generalized SVS via para-Bose formulation. Thus, assuming the condition

$$\varphi_0 = 0 \Longrightarrow u = 0,\tag{11}$$

one can obtain the generalized SVS following the nonunitary approach, as described in Ref. [48]. In what follows we will apply this condition.

A. Para-Bose number states

In this subsection, we will recall some properties of the para-Bose number states, since the generalized SVS can be expanded in these states. It is well-known that the para-Bose number states $|n, \varepsilon\rangle$ form a complete set and are orthonormal [38, 51], i.e.,

$$\sum_{n=0}^{\infty} |n, \varepsilon\rangle \langle \varepsilon, n| = 1, \quad \langle \varepsilon, n| m, \varepsilon\rangle = \delta_{n,m}, \tag{12}$$

where $\delta_{n,m}$ is the Kronecker delta.

Taking into account that the vacuum state has even parity $(\hat{R} | 0, \varepsilon) = |0, \varepsilon\rangle$, the action of the generators of the WHA in $|n, \varepsilon\rangle$, reads

$$\hat{a} | 2n, \varepsilon \rangle = \sqrt{2n} | 2n - 1, \varepsilon \rangle, \quad \hat{a}^{\dagger} | 2n, \varepsilon \rangle = \sqrt{2(n + \varepsilon)} | 2n + 1, \varepsilon \rangle,
\hat{a} | 2n + 1, \varepsilon \rangle = \sqrt{2(n + \varepsilon)} | 2n, \varepsilon \rangle, \quad \hat{a}^{\dagger} | 2n + 1, \varepsilon \rangle = \sqrt{2(n + 1)} | 2n + 2, \varepsilon \rangle,
\hat{n} | n, \varepsilon \rangle = \left(\frac{1}{2} \left\{ \hat{a}, \hat{a}^{\dagger} \right\} - \varepsilon \right) | n, \varepsilon \rangle = n | n, \varepsilon \rangle, \quad \hat{R} | n, \varepsilon \rangle = (-1)^n | n, \varepsilon \rangle, \quad n = 0, 1, 2, 3, \dots$$
(13)

From a recurrence relation, one can write the number states $|n, \varepsilon\rangle$ in terms of the vacuum state $|0, \varepsilon\rangle$, in the form

$$|2n,\varepsilon\rangle = \sqrt{\frac{\Gamma(\varepsilon)}{2^{2n}n!\Gamma(n+\varepsilon)}} \left(\hat{a}^{\dagger}\right)^{2n} |0,\varepsilon\rangle,$$

$$|2n+1,\varepsilon\rangle = \sqrt{\frac{\Gamma(\varepsilon)}{2^{2n+1}n!\Gamma(n+\varepsilon+1)}} \left(\hat{a}^{\dagger}\right)^{2n+1} |0,\varepsilon\rangle. \tag{14}$$

B. Time-dependent generalized SVS via para-Bose number states

In what follows, we aim to apply the nonunitary approach, as in the recent publication [48], to construct the generalized SVS. From condition (11), one can write the nonunitary operator \hat{S} , in the form

$$\hat{S} = \exp\left(\frac{1}{2}\zeta\hat{a}^{\dagger 2}\right), \quad \zeta = \frac{g}{f}, \quad \dot{\zeta} = i\alpha^*\zeta^2 - 2i\beta\zeta + i\alpha, \tag{15}$$

such that the commutators from Baker-Campbell-Hausdorff relation of the second order onwards become null. Here, $\zeta = \zeta(t)$ represents the squeeze parameter for the generalized SVS. The form of \hat{S} allows us to introduce the most convenient squeeze parameter, as well as relate the motion integral \hat{A} directly with the annihilation operator \hat{a} . Furthermore, this approach guarantees that there is a direct relationship between the eigenstates of the motion integral with the para-Bose number states, as we will see next.

Applying the Baker–Campbell–Hausdorff theorem, we can express the operator \hat{a} in terms of the integrals of motion \hat{A} , as follows

$$\hat{a} = \frac{1}{f} \hat{S} \hat{A} \hat{S}^{-1}. \tag{16}$$

The application from (16) on the vacuum state $|0, \varepsilon\rangle$, which satisfies the annihilation condition $\hat{a} |0, \varepsilon\rangle = 0$, yields:

$$\hat{A}\left|\zeta\right\rangle = 0,\tag{17}$$

whose general solution is given by

$$|\zeta\rangle = \Phi \exp\left(-\frac{1}{2}\zeta\hat{a}^{\dagger 2}\right)|0,\varepsilon\rangle,$$
 (18)

where $\Phi = \Phi(t)$ is an arbitrary function, which will be determined such that the states $|\zeta\rangle$ satisfy the Schrödinger's equation (5).

Substituting the states (18) into (5), we obtain the following equation for Φ :

$$\frac{\dot{\Phi}}{\Phi} = \frac{1}{2} \frac{\left\langle \zeta \left| \hat{a}^{\dagger 2} \right| \zeta \right\rangle}{\left\langle \zeta \right| \zeta \right\rangle} \dot{\zeta} - \frac{i}{\hbar} \frac{\left\langle \zeta \left| \hat{H} \right| \zeta \right\rangle}{\left\langle \zeta \right| \zeta \right\rangle}.$$
(19)

Using the representation (10) together with the condition (17), one can easily calculate the mean values in (19), with $\dot{\zeta}$ given in (15). Thus, the general solution from (19), reads

$$\Phi = \frac{C}{f^{\varepsilon}} \exp\left(-i \int \delta dt\right),\tag{20}$$

where C is a real normalization constant. Taking into account the normalization condition, we find that:

$$\langle \zeta | \zeta \rangle = 1 \Rightarrow C = \mu^{\varepsilon/2}.$$
 (21)

Then, the normalized states $|\zeta\rangle$ that satisfy the Schrödinger's equation are given by

$$|\zeta\rangle = \frac{\sqrt{\mu^{\varepsilon}}}{f^{\varepsilon}} \exp\left(-i\int \delta dt\right) \exp\left(-\frac{1}{2}\zeta\hat{a}^{\dagger 2}\right) |0,\varepsilon\rangle$$

$$= \frac{\sqrt{\mu^{\varepsilon}}}{f^{\varepsilon}} \exp\left(-i\int \delta dt\right) \sum_{n=0}^{\infty} (-\zeta)^{n} \sqrt{\frac{\Gamma(n+\varepsilon)}{n!\Gamma(\varepsilon)}} |2n,\varepsilon\rangle. \tag{22}$$

The above sum converges as long as the condition $|\zeta| < 1$ is satisfied.

On the other hand, we can express f and μ in terms of squeeze parameter in the form:

$$f = f_0 \exp\left[-i\int \left(\alpha^* \zeta - \beta\right) dt\right], \quad \mu = |f|^2 \left(1 - |\zeta|^2\right). \tag{23}$$

From here, the states $|\zeta\rangle$ take the form

$$|\zeta\rangle = \left(1 - |\zeta|^2\right)^{\frac{\varepsilon}{2}} e^{i\vartheta} \sum_{n=0}^{\infty} \left(-\zeta\right)^n \sqrt{\frac{\Gamma(n+\varepsilon)}{n!\Gamma(\varepsilon)}} |2n, \varepsilon\rangle, \qquad (24)$$

where the phase ϑ is given by

$$\vartheta = \int \left[\varepsilon \operatorname{Re} \left(\alpha \zeta^* \right) - \varepsilon \beta - \delta \right] dt. \tag{25}$$

In what follows, we call the time-dependent states (24) generalized SVS. Notice that, if we assume $\varepsilon = 1/2$ and $\zeta = e^{i\theta} \tanh{(r)} \Longrightarrow \mu = 1$, the Eq. (24) takes the form:

$$|\zeta\rangle = \frac{e^{i\vartheta}}{\sqrt{\cosh(r)}} \sum_{n=0}^{\infty} \frac{\sqrt{(2n)!}}{2^n n!} \left[-e^{i\theta} \tanh(r) \right]^n \left| 2n, \frac{1}{2} \right\rangle, \tag{26}$$

where $\Gamma\left(n+\frac{1}{2}\right)=\frac{\sqrt{\pi}4^{-n}(2n)!}{n!}$, $\Gamma\left(1/2\right)=\sqrt{\pi}$. So, we can conclude that the states obtained in Eq. (26) reproduce the usual case [74]. In this case, the Eq. (24) corresponds to a generalization of the SVS, differing by a time-dependent phase factor due to the time evolution that was included in our analysis. We must highlight that when identifying $\varepsilon=2k$, the states (24) coincide with the time-independent CS of the $SU\left(1,1\right)$ group constructed by Peremolov [21].

The overlap of two generalized SVS with different ζ , for example $\langle \zeta_1 | \zeta_2 \rangle$ reads

$$\langle \zeta_1 | \zeta_2 \rangle = \frac{\left(1 - |\zeta_1|^2\right)^{\frac{\varepsilon}{2}} \left(1 - |\zeta_2|^2\right)^{\frac{\varepsilon}{2}}}{\left(1 - \zeta_1^* \zeta_2\right)^{\varepsilon}} \exp\left\{i\varepsilon \int \operatorname{Re}\left[\alpha \left(\zeta_2^* - \zeta_1^*\right)\right] dt\right\}. \tag{27}$$

In turn, the probability transition $P_{2n}(\zeta,\varepsilon) = |\langle \varepsilon, 2n | \zeta \rangle|^2$ is given by

$$P_{2n}\left(\zeta,\varepsilon\right) = \frac{\left(1-\left|\zeta\right|^{2}\right)^{\varepsilon}}{\Gamma\left(\varepsilon\right)} \frac{\Gamma\left(n+\varepsilon\right)\left|\zeta\right|^{2n}}{n!}.$$
(28)

The probability transition (28) has been plotted in Fig. 1. As we see, the increase in the value of the ε parameter implies a larger dispersion of the probability transition.

C. Completeness relation on squeeze parameter

Considerations on the completeness relation for the SVS and squeezed odd number states, in the context of canonical algebra, can be seen in Refs. [74, 75]. Here, our contribution is to obtain the completeness relation in the context of the WHA, and as will be seen, its existence depends on the ε -parameter.




FIG. 1: Probability transition of the generalized SVS by considering fixed value for $|\zeta| = 0.3$.

Let us consider a weight function $w(\zeta)$ such that the states $|\zeta\rangle$ lead to the following closure relation

$$\int_{\mathbb{C}} |\zeta\rangle \langle \zeta| w(\zeta) d^{2}\zeta = 1.$$
 (29)

In turn, substituting the states (24) into Eq. (29), we find

$$\sum_{n,m=0}^{\infty} |2n,\varepsilon\rangle \langle \varepsilon, 2m| \sqrt{\frac{\Gamma(n+\varepsilon)}{n!\Gamma(\varepsilon)}} \frac{\Gamma(m+\varepsilon)}{m!\Gamma(\varepsilon)} \int_{\mathbb{C}} \left(1-|\zeta|^2\right)^{\varepsilon} (-\zeta)^n (-\zeta^*)^m w(\zeta) d^2\zeta = 1. \quad (30)$$

Now, by using polar coordinates in the above relation,

$$\zeta = r_{\zeta}e^{i\theta_{\zeta}}, \quad d^{2}\zeta = r_{\zeta}dr_{\zeta}d\theta_{\zeta}, \quad 0 \le r_{\zeta} < 1, \quad 0 \le \theta_{\zeta} \le 2\pi,$$
(31)

we get

$$\sum_{n,m=0}^{\infty} |2n,\varepsilon\rangle \langle \varepsilon, 2m| \sqrt{\frac{\Gamma(n+\varepsilon)}{n!\Gamma(\varepsilon)}} \frac{\Gamma(m+\varepsilon)}{m!\Gamma(\varepsilon)} \int_{0}^{1} \int_{0}^{2\pi} \left(1 - r_{\zeta}^{2}\right)^{\varepsilon} (-r_{\zeta})^{n} (-r_{\zeta})^{m} \times w(\zeta) \exp\left[i(n-m)\theta_{\zeta}\right] r_{\zeta} dr_{\zeta} d\theta_{\zeta} = 1.$$
(32)

On the other hand, we may readily show that $w(\zeta) = w(r_{\zeta})$, which implies that the integral on θ_{ζ} becomes

$$\int_0^{2\pi} d\theta_{\zeta} \exp\left[i\left(n - m\right)\theta_{\zeta}\right] = 2\pi \delta_{n,m}.$$
(33)

From here, we can write (32) in the form

$$2\pi \sum_{n=0}^{\infty} |2n, \varepsilon\rangle \langle \varepsilon, 2n| \frac{\Gamma(n+\varepsilon)}{n!\Gamma(\varepsilon)} \int_{0}^{1} (1 - r_{\zeta}^{2})^{\varepsilon} r_{\zeta}^{2n+1} w(r_{\zeta}) dr_{\zeta} = 1.$$
 (34)

By using the following relationship among gamma functions

$$\frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} = 2\int_0^1 (1-t^2)^{y-1} t^{2x-1} dt, \quad \operatorname{Re}(x) > 0, \quad \operatorname{Re}(y) > 0, \tag{35}$$

one can see that the weight function $w(r_{\zeta})$ must have the form

$$w(r_{\zeta}) = \frac{\Gamma(\varepsilon)}{\pi\Gamma(\varepsilon - 1)\left(1 - r_{\zeta}^{2}\right)^{2}} = \frac{\varepsilon - 1}{\pi\left(1 - r_{\zeta}^{2}\right)^{2}}, \quad \varepsilon > 1,$$
(36)

in order to ensure that the completeness relation (29) is satisfied. It is important to highlight that the condition $\varepsilon > 1$ leads to a positive weight function $w(r_{\zeta})$. At the same time, this condition shows that the canonical algebra ($\varepsilon = 1/2$) does not allow to obtain a completeness relation for the generalized SVS. It follows from (27) and (29) that these states form an overcomplete set of states on the Hilbert space. The plot of weight function can be seen in the Fig. 2.

IV. TIME-DEPENDENT GENERALIZED CS

In Ref. [48], the SCS has been built through the nonunitary approach by considering the canonical algebra. In another way, by considering the WHA, it is not possible to obtain the SCS since, from Baker–Campbell–Hausdorff theorem, we cannot establish a direct relationship between the para-Bose operators and integral of motion, due the deformed commutation relation. However, instead of following the nonunitary approach, we will construct the eigenstates of the integral of motion \hat{A} , in the form

$$\hat{A}|z,t\rangle = z|z,t\rangle, \tag{37}$$




FIG. 2: Graphics of $w(r_{\zeta})$.

where z is a complex constant. The Eq. (37) allows us to assume

$$\varphi_0 = 0 \Longrightarrow u = 0, \tag{38}$$

without loss of generality.

The states $|z,t\rangle$ can be expanded in terms of the para-Bose number states, as follows

$$|z,t\rangle = \sum_{n=0}^{\infty} c_n |n,\varepsilon\rangle = \sum_{n=0}^{\infty} \left(c_{2n} |2n,\varepsilon\rangle + c_{2n+1} |2n+1,\varepsilon\rangle \right), \tag{39}$$

where $c_n = c_n(t)$ are time-dependent coefficients and will be determined such that the Eqs. (37) and (5) will be satisfied. Substituting (6), (13) and (39) into (37), we find the following equations for the coefficients

$$c_{2n} = \sqrt{\frac{n!\Gamma(\varepsilon)}{\Gamma(n+\varepsilon)}} \left(-\frac{g}{f}\right)^n L_n^{\varepsilon-1} \left(\frac{z^2}{2gf}\right) c_0,$$

$$c_{2n+1} = \frac{z}{f} \sqrt{\frac{n!\Gamma(\varepsilon)}{2\Gamma(n+\varepsilon+1)}} \left(-\frac{g}{f}\right)^n L_n^{\varepsilon} \left(\frac{z^2}{2gf}\right) c_0,$$
(40)

where $L_n^{\alpha}(x)$ are the associated Laguerre polynomials and c_0 is a time-dependent function, which will be determined such that the states $|z,t\rangle$ satisfy the Schrödinger's equation (5).

Here, it is convenient to introduce the squeeze $\zeta = g/f$ and displacement $\xi = z/f$ parameters to rewrite the states $|z,t\rangle \longrightarrow |\zeta,\xi\rangle$, in the form

$$|\zeta,\xi\rangle = \sqrt{\Gamma(\varepsilon)}c_0 \sum_{n=0}^{\infty} (-\zeta)^n \sqrt{n!} \left[\frac{L_n^{\varepsilon-1}\left(\frac{\xi^2}{2\zeta}\right)}{\sqrt{\Gamma(n+\varepsilon)}} |2n,\varepsilon\rangle + \frac{\xi L_n^{\varepsilon}\left(\frac{\xi^2}{2\zeta}\right)}{\sqrt{2\Gamma(n+\varepsilon+1)}} |2n+1,\varepsilon\rangle \right], (41)$$

with ζ and ξ satisfy the following differential equations:

$$\dot{\zeta} = i\alpha^* \zeta^2 - 2i\beta\zeta + i\alpha, \quad \dot{\xi} = i(\alpha^* \zeta - \beta)\xi. \tag{42}$$

From normalization condition

$$\langle \xi, \zeta | \zeta, \xi \rangle = 1, \tag{43}$$

we found the following form for the function c_0 ,

$$c_{0} = \left(\frac{\xi}{\sqrt{2}}\right)^{\varepsilon - 1} \sqrt{\frac{1 - |\zeta|^{2}}{\Gamma(\varepsilon)}} \frac{\exp\left\{\frac{\zeta^{*} \xi^{2}}{2(1 - |\zeta|^{2})} + i \int \left[\operatorname{Re}\left(\alpha \zeta^{*}\right) - \beta\right] dt + i\phi\right\}}{\sqrt{I_{\varepsilon - 1}\left(\frac{|\xi|^{2}}{1 - |\zeta|^{2}}\right) + I_{\varepsilon}\left(\frac{|\xi|^{2}}{1 - |\zeta|^{2}}\right)}},$$
(44)

where $I_{\kappa}(Z)$ is the modified Bessel function of the first kind, and ϕ is a time-dependent real function. Thus, the normalized states $|\zeta,\xi\rangle$ take the form

$$|\zeta,\xi\rangle = \left(\frac{\xi}{\sqrt{2}}\right)^{\varepsilon-1} \sqrt{\frac{1-|\zeta|^2}{I_{\varepsilon-1}\left(\frac{|\xi|^2}{1-|\zeta|^2}\right) + I_{\varepsilon}\left(\frac{|\xi|^2}{1-|\zeta|^2}\right)}} \exp\left\{\frac{\zeta^*\xi^2}{2\left(1-|\zeta|^2\right)} + i\int \left[\operatorname{Re}\left(\alpha\zeta^*\right) - \beta\right] dt + i\phi\right\}$$

$$\times \sum_{n=0}^{\infty} (-\zeta)^n \sqrt{n!} \left[\frac{L_n^{\varepsilon-1}\left(\frac{\xi^2}{2\zeta}\right)}{\sqrt{\Gamma(n+\varepsilon)}} |2n,\varepsilon\rangle + \frac{\xi L_n^{\varepsilon}\left(\frac{\xi^2}{2\zeta}\right)}{\sqrt{2\Gamma(n+\varepsilon+1)}} |2n+1,\varepsilon\rangle\right], \tag{45}$$

Substituting $|\zeta,\xi\rangle$ into Schrödinger's equation, we find the following expression for ϕ ,

$$\phi = -\int \delta dt. \tag{46}$$

Therefore, the normalized states $|\zeta,\xi\rangle$ that satisfy the Schrödinger's equation are given by

$$|\zeta,\xi\rangle = \left(\frac{\xi}{\sqrt{2}}\right)^{\varepsilon-1} \sqrt{\frac{1-|\zeta|^2}{I_{\varepsilon-1}\left(\frac{|\xi|^2}{1-|\zeta|^2}\right) + I_{\varepsilon}\left(\frac{|\xi|^2}{1-|\zeta|^2}\right)}} \exp\left[\frac{\zeta^*\xi^2}{2\left(1-|\zeta|^2\right)} + i\tilde{\vartheta}\right] \times \sum_{n=0}^{\infty} (-\zeta)^n \sqrt{n!} \left[\frac{L_n^{\varepsilon-1}\left(\frac{\xi^2}{2\zeta}\right)}{\sqrt{\Gamma(n+\varepsilon)}} |2n,\varepsilon\rangle + \frac{\xi L_n^{\varepsilon}\left(\frac{\xi^2}{2\zeta}\right)}{\sqrt{2\Gamma(n+\varepsilon+1)}} |2n+1,\varepsilon\rangle\right], \quad (47)$$

where

$$\tilde{\vartheta} = \int \left[\operatorname{Re} \left(\alpha \zeta^* \right) - \beta - \delta \right] dt. \tag{48}$$

In the following, we call the time-dependent states (47) generalized para-Bose CS.

It's worth mentioning that the states (47) present a set of information that, to our knowledge, has not been previously derived, see, e.g., [38, 51–58, 65, 66]. Such information consists of the explicit form of the squeeze, displacement, and deformation parameters

(Wigner-parameter). These parameters are present explicitly in the standard deviation and probability transition. Furthermore, the time-dependent states (47) have been expanded in terms of the time-independent para-Bose number states. Thus, the states (47) lead to a deeper understanding of the physical systems described by the Hamiltonian (4). As we will see from section V, the Hamiltonian can be seen as a generalization of Calogero model [59].

In particular, taking into account the condition $\zeta = 0 \Longrightarrow \alpha = 0$, the states (47) take the form

$$|\xi\rangle = \left(\frac{\xi}{\sqrt{2}}\right)^{\varepsilon-1} \frac{\exp\left[-i\int\left(\beta+\delta\right)dt\right]}{\sqrt{I_{\varepsilon-1}\left(|\xi|^2\right) + I_{\varepsilon}\left(|\xi|^2\right)}} \sum_{n=0}^{\infty} \left(\frac{\xi^2}{2}\right)^n \left[\frac{|2n,\varepsilon\rangle}{\sqrt{n!\Gamma(n+\varepsilon)}} + \frac{\xi|2n+1,\varepsilon\rangle}{\sqrt{2n!\Gamma(n+\varepsilon+1)}}\right]$$

$$= \sqrt{\Gamma(\varepsilon)} \exp\left[-i\int\left(\beta+\delta\right)dt\right] \left(\frac{\hat{a}^{\dagger}}{\sqrt{2}}\right)^{1-\varepsilon} \frac{I_{\varepsilon-1}\left(\xi\hat{a}^{\dagger}\right) + I_{\varepsilon}\left(\xi\hat{a}^{\dagger}\right)}{\sqrt{I_{\varepsilon-1}\left(|\xi|^2\right) + I_{\varepsilon}\left(|\xi|^2\right)}} |0,\varepsilon\rangle, \tag{49}$$

with $\dot{\xi} = -i\beta\xi$ and $|\xi\rangle = |0,\xi\rangle$. Except for time evolution, these states correspond to the para-Bose CS obtained in Ref. [51]. In turn, we have that the condition $\varepsilon = 1/2$ reduces the states (49) to the form:

$$|\xi\rangle = \exp\left(\int \frac{\beta + 2\delta}{2i} dt - \frac{|\xi|^2}{2}\right) \exp\left(\xi \hat{a}^{\dagger}\right) |0\rangle, \quad |0\rangle = |0, \varepsilon = 1/2\rangle,$$
 (50)

which are the time-dependent canonical CS [76, 77].

The overlap of the states $|\zeta,\xi\rangle$ for different squeeze ζ and displacement ξ parameters is given by

$$\langle \xi_{1}, \zeta_{1} | \zeta, \xi \rangle = \left(\frac{\xi \xi_{1}^{*}}{2}\right)^{\varepsilon - 1} \sqrt{\frac{\left(1 - |\zeta|^{2}\right) \left(1 - |\zeta_{1}|^{2}\right)}{\left[I_{\varepsilon - 1}\left(\frac{|\xi|^{2}}{1 - |\zeta|^{2}}\right) + I_{\varepsilon}\left(\frac{|\xi|^{2}}{1 - |\zeta|^{2}}\right)\right] \left[I_{\varepsilon - 1}\left(\frac{|\xi_{1}|^{2}}{1 - |\zeta_{1}|^{2}}\right) + I_{\varepsilon}\left(\frac{|\xi_{1}|^{2}}{1 - |\zeta_{1}|^{2}}\right)\right]}}$$

$$\times \exp\left[\frac{\zeta^{*}\xi^{2}\left(1 - |\zeta_{1}|^{2}\right) + \zeta_{1}\xi_{1}^{*2}\left(1 - |\zeta|^{2}\right)}{2\left(1 - |\zeta|^{2}\right)} + i\int \operatorname{Re}\left(\alpha\zeta^{*} - \alpha\zeta_{1}^{*}\right) dt\right]$$

$$\times \sum_{n=0}^{\infty} \left(\zeta\zeta_{1}^{*}\right)^{n} n! \left[\frac{L_{n}^{\varepsilon - 1}\left(\frac{\xi^{2}}{2\zeta}\right)L_{n}^{\varepsilon - 1}\left(\frac{\xi_{1}^{*2}}{2\zeta_{1}^{*}}\right)}{\Gamma\left(n + \varepsilon\right)} + \frac{\xi\xi_{1}^{*}}{2}\frac{L_{n}^{\varepsilon}\left(\frac{\xi^{2}}{2\zeta}\right)L_{n}^{\varepsilon}\left(\frac{\xi_{1}^{*2}}{2\zeta_{1}^{*}}\right)}{\Gamma\left(n + \varepsilon + 1\right)}\right]. \tag{51}$$

Here, we can analyze the probability transition $P_n(\zeta, \xi, \varepsilon) = |\langle \varepsilon, n | \zeta, \xi \rangle|^2$ of the number

states $|n, \varepsilon\rangle$ to the CS $|\zeta, \xi\rangle$:

$$P_{n}\left(\zeta,\xi,\varepsilon\right) = \left(\frac{\left|\xi\right|^{2}}{2}\right)^{\varepsilon-1} \frac{\left(1-\left|\zeta\right|^{2}\right) \exp\left[\frac{\operatorname{Re}\left(\zeta^{*}\xi^{2}\right)}{1-\left|\zeta\right|^{2}}\right]}{I_{\varepsilon-1}\left(\frac{\left|\xi\right|^{2}}{1-\left|\zeta\right|^{2}}\right) + I_{\varepsilon}\left(\frac{\left|\xi\right|^{2}}{1-\left|\zeta\right|^{2}}\right)} m! \left|\zeta\right|^{2m} \left[\frac{\left|L_{m}^{\varepsilon-1}\left(\frac{\xi^{2}}{2\zeta}\right)\right|^{2} \delta_{n,2m}}{\Gamma\left(m+\varepsilon\right)} + \frac{\left|\xi\right|^{2} \left|L_{m}^{\varepsilon}\left(\frac{\xi^{2}}{2\zeta}\right)\right|^{2} \delta_{n,2m+1}}{2\Gamma\left(m+\varepsilon+1\right)}\right]$$

$$(52)$$

The probability transition $P_n(\zeta, \xi, \varepsilon)$ has been shown in Fig. 3. As we can see, the ξ parameter allows access to odd states while the ε controls the dispersion of the probability
density. Furthermore, higher values of the ε -parameter allow access to states with higher
principal quantum number n, and attenuate the access to the odd states.

A. Mean values and uncertainty relations

Let us now consider the mean value of the momentum \hat{P} and position \hat{x} operators, which satisfy the commutation relation

$$\left[\hat{x}, \hat{P}\right] = i\hbar \left[1 + (2\varepsilon - 1)\,\hat{R}\right],\tag{53}$$

in the generalized CS (47). For this, we will rewrite these operators in terms of the integrals of motion, as seen below

$$\hat{a} = \frac{\hat{A}_{f} - \zeta \hat{A}_{f^{*}}^{\dagger}}{1 - |\zeta|^{2}}, \quad \hat{A}_{f} \equiv \frac{1}{f} \hat{A},$$

$$\hat{x} = \frac{\hat{a} + \hat{a}^{\dagger}}{\sqrt{2}} l = \frac{l}{\sqrt{2}} \frac{(1 - \zeta^{*}) \hat{A}_{f} + (1 - \zeta) \hat{A}_{f^{*}}^{\dagger}}{1 - |\zeta|^{2}},$$

$$\hat{P} = \hbar \frac{\hat{a} - \hat{a}^{\dagger}}{i\sqrt{2}l} = \frac{\hbar}{i\sqrt{2}l} \frac{(1 + \zeta^{*}) \hat{A}_{f} - (1 + \zeta) \hat{A}_{f^{*}}^{\dagger}}{1 - |\zeta|^{2}},$$
(54)

where l is a length-dimensional parameter which is related to the initial standard deviation [48]. The new operator \hat{A}_f acts on states $|\zeta,\xi\rangle$ as follows

$$\hat{A}_f |\zeta, \xi\rangle = \xi |\zeta, \xi\rangle. \tag{55}$$




FIG. 3: Probability transition of the generalized CS.

Using the relations (54) and (55), we can easily calculate the mean values of the operators \hat{x} and \hat{P} ,

$$\overline{x} = \overline{x}(t) = \langle \xi, \zeta | \hat{x} | \zeta, \xi \rangle = \sqrt{2} l \frac{\operatorname{Re}\left[(1 - \zeta^*) \xi \right]}{1 - |\zeta|^2},$$

$$\overline{P} = \overline{P}(t) = \left\langle \xi, \zeta | \hat{P} | \zeta, \xi \right\rangle = \frac{\sqrt{2} \hbar}{l} \frac{\operatorname{Im}\left[(1 + \zeta^*) \xi \right]}{1 - |\zeta|^2}.$$
(56)

From here, there is a correspondence between the squeeze ζ and displacement ξ parameters with the mean values of \overline{x} and \overline{P} ,

$$\xi = \frac{1+\zeta}{\sqrt{2}l}\overline{x} + \frac{il}{\hbar}\frac{1-\zeta}{\sqrt{2}}\overline{P}.$$
 (57)

Taking the square of the operators \hat{x} and \hat{P} , we have

$$\hat{x}^{2} = \frac{l^{2}}{2} \frac{(1 - \zeta^{*})^{2} \hat{A}_{f}^{2} + (1 - \zeta)^{2} \hat{A}_{f^{*}}^{\dagger 2} + 2 |1 - \zeta|^{2} \hat{A}_{f^{*}}^{\dagger} \hat{A}_{f}}{(1 - |\zeta|^{2})^{2}} + l^{2} \frac{|1 - \zeta|^{2}}{1 - |\zeta|^{2}} \frac{1 + (2\varepsilon - 1) \hat{R}}{2},$$

$$\hat{P}^{2} = -\frac{\hbar^{2}}{2l^{2}} \frac{(1 + \zeta^{*})^{2} \hat{A}_{f}^{2} + (1 + \zeta)^{2} \hat{A}_{f^{*}}^{\dagger 2} - 2 |1 + \zeta|^{2} \hat{A}_{f^{*}}^{\dagger} \hat{A}_{f}}{(1 - |\zeta|^{2})^{2}} + \frac{\hbar^{2}}{l^{2}} \frac{|1 + \zeta|^{2}}{1 - |\zeta|^{2}} \frac{1 + (2\varepsilon - 1) \hat{R}}{2}.$$
(58)

So, from these results, we can calculate the mean values of $\overline{x^2}$ and $\overline{P^2}$, as follow

$$\overline{x^{2}} = \overline{x^{2}}(t) = \langle \xi, \zeta | \hat{x}^{2} | \zeta, \xi \rangle = \frac{2l^{2} \operatorname{Re}^{2} \left[(1 - \zeta^{*}) \xi \right]}{\left(1 - |\zeta|^{2} \right)^{2}} + l^{2} \frac{|1 - \zeta|^{2}}{1 - |\zeta|^{2}} \frac{1 + (2\varepsilon - 1) \overline{R}}{2},$$

$$\overline{P^{2}} = \overline{P^{2}}(t) = \langle \xi, \zeta | \hat{P}^{2} | \zeta, \xi \rangle = \frac{2\hbar^{2} \operatorname{Im}^{2} \left[(1 + \zeta^{*}) \xi \right]}{l^{2} \left(1 - |\zeta|^{2} \right)^{2}} + \frac{\hbar^{2}}{l^{2}} \frac{|1 + \zeta|^{2}}{1 - |\zeta|^{2}} \frac{1 + (2\varepsilon - 1) \overline{R}}{2}, \quad (59)$$

where

$$\overline{R} = \left\langle \xi, \zeta \left| \hat{R} \right| \zeta, \xi \right\rangle = \frac{I_{\varepsilon - 1} \left(\frac{|\xi|^2}{1 - |\zeta|^2} \right) - I_{\varepsilon} \left(\frac{|\xi|^2}{1 - |\zeta|^2} \right)}{I_{\varepsilon - 1} \left(\frac{|\xi|^2}{1 - |\zeta|^2} \right) + I_{\varepsilon} \left(\frac{|\xi|^2}{1 - |\zeta|^2} \right)}.$$
 (60)

Note that $\xi = 0$ leads to a well-defined even parity of the states (47), once $\overline{R} = 1$.

In what follows, we calculate the standard deviation, i.e.,

$$\sigma_{x} = \sigma_{x} (t) = \sqrt{\overline{x^{2}} - \overline{x}^{2}} = l |1 - \zeta| \sqrt{\frac{1 + (2\varepsilon - 1)\overline{R}}{2(1 - |\zeta|^{2})}},$$

$$\sigma_{P} = \sigma_{P} (t) = \sqrt{\overline{P^{2}} - \overline{P}^{2}} = \frac{\hbar}{l} |1 + \zeta| \sqrt{\frac{1 + (2\varepsilon - 1)\overline{R}}{2(1 - |\zeta|^{2})}}.$$
(61)

From Eqs. (61), it is easy to see that standard deviations in position and momentum present the squeezing property.

Finally, we aim to find the Heisenberg uncertainty relation in the generalized states (47) by considering the WHA. Therefore, in the hold of previous results, we can explicitly calculate the product $\sigma_x \sigma_P$, as shown below:

$$\sigma_{x}\sigma_{P} = \hbar \frac{|1 - \zeta| |1 + \zeta|}{1 - |\zeta|^{2}} \frac{1 + (2\varepsilon - 1)\overline{R}}{2} = \hbar \sqrt{1 + \frac{4\operatorname{Im}^{2}(\zeta)}{(1 - |\zeta|^{2})^{2}} \frac{1 + (2\varepsilon - 1)\overline{R}}{2}}.$$
 (62)

In particular, if we consider that the squeeze parameter assumes real values, this implies that $\sigma_x \sigma_P = \hbar/2 \left(1 + (2\varepsilon - 1)\overline{R}\right)$. In turn, for this particular choice, we should note that the Heisenberg uncertainty assumes the minimum value predicted by the relation

$$\sigma_A \sigma_B \ge \frac{1}{2} \left| \left\langle \left[\hat{A}, \hat{B} \right] \right\rangle \right|.$$
 (63)

It must be highlighted that in Ref. [38], a detailed analysis has been applied to show how the para-Bose uncertainty relation leads to uncertainty relation computed from canonical commutation relation.

Taking into account the covariance σ_{xP} ,

$$\sigma_{xP} = \frac{\left\langle \xi, \zeta \left| \hat{P} \hat{x} \right| \zeta, \xi \right\rangle + \left\langle \xi, \zeta \left| \hat{x} \hat{P} \right| \zeta, \xi \right\rangle}{2} - \overline{x} \left(t \right) \overline{P} \left(t \right) = -\hbar \operatorname{Im} \left(\zeta \right) \frac{1 + (2\varepsilon - 1) \overline{R}}{1 - \left| \zeta \right|^{2}}, \quad (64)$$

we can calculate the Schrödinger-Robertson uncertainty relation [78],

$$\sigma_x^2 \sigma_P^2 - \sigma_{xP}^2 = \frac{\hbar^2}{4} \left[1 + (2\varepsilon - 1)\overline{R} \right]^2, \tag{65}$$

which, as can we see, is minimized.

V. COORDINATE REPRESENTATION OF THE GENERALIZED CS

It is well-known that the phase space in the context of quantum mechanics encounters difficulties due to the uncertainty principle. Considering the states $|\zeta, \xi\rangle$ in a coordinate representation allow us to introduce the quasiprobability Wigner distribution, which plays an analogous role to the classical distributions [79–81].

According to WHA, \hat{P} is a self-adjoint operator on semi-axis $(x \ge 0)$, see [36], and have the following form:

$$\hat{P} = -i\hbar\partial_x + \frac{i\hbar}{2x}(2\varepsilon - 1)\,\hat{R}.\tag{66}$$

In this case, the annihilation operator takes the form

$$\hat{a} = \frac{1}{\sqrt{2}} \left(\frac{\hat{x}}{l} + \frac{il}{\hbar} \hat{P} \right) = \frac{l}{\sqrt{2}} \left(\partial_x - \frac{2\varepsilon - 1}{2x} \hat{R} + \frac{x}{l^2} \right). \tag{67}$$

Applying the annihilation condition

$$\hat{a}\Psi_{0,\varepsilon}(x) = 0, \quad \Psi_{0,\varepsilon}(x) = \langle x|0,\varepsilon\rangle,$$
 (68)

we can obtain a differential equation for the vacuum state, as follows:

$$\left(\partial_x + \frac{x}{l^2} - \frac{2\varepsilon - 1}{2x}\right)\Psi_{0,\varepsilon}(x) = 0, \quad \hat{R}\Psi_{0,\varepsilon}(x) = \Psi_{0,\varepsilon}(x). \tag{69}$$

The general solution reads

$$\Psi_{0,\varepsilon}(x) = Cx^{\varepsilon - \frac{1}{2}} \exp\left(-\frac{x^2}{2l^2}\right),\tag{70}$$

with C being a real constant, which will be determined through the normalization condition, as seen below,

$$2C^{2} \int_{0}^{\infty} x^{2\varepsilon - 1} \exp\left(-\frac{x^{2}}{l^{2}}\right) dx = 1 \Rightarrow C = \frac{1}{l^{\varepsilon} \sqrt{\Gamma(\varepsilon)}}.$$
 (71)

It should be noted that the condition (69) leads to the following quantization condition:

$$\varepsilon = 2\ell + \frac{1}{2}, \quad \ell = 0, 1, 2, \dots,$$
 (72)

where ℓ is analogous to the angular momentum. Such quantization was obtained in [37] to ensure that the eigenfunctions of \hat{P} are differentiable at the origin.

Thus, the vacuum state with even parity $\Psi_{0,\varepsilon}\left(x\right)\equiv\Psi_{0,\ell}^{e}\left(x\right)$ takes the form

$$\Psi_{0,\ell}^{e}(x) = \frac{1}{l^{2\ell + \frac{1}{2}} \sqrt{\Gamma\left(2\ell + \frac{1}{2}\right)}} x^{2\ell} \exp\left(-\frac{x^2}{2l^2}\right). \tag{73}$$

Taking into account that $\Psi_{\zeta,\xi}^{\ell}(x,t) = \langle x|\zeta,\xi\rangle$ and replacing the relations (14) in the states (47), we obtain

$$\Psi_{\zeta,\xi}^{\ell}(x,t) = \xi^{2\ell - \frac{1}{2}} \frac{\sqrt{\left(1 - |\zeta|^{2}\right) \Gamma\left(2\ell + \frac{1}{2}\right)} \exp\left[\frac{\zeta^{*}\xi^{2}}{2\left(1 - |\zeta|^{2}\right)} + i\tilde{\vartheta}\right]}}{2^{\frac{4\ell - 1}{4}} \sqrt{I_{2\ell - \frac{1}{2}}\left(\frac{|\xi|^{2}}{1 - |\zeta|^{2}}\right) + I_{2\ell + \frac{1}{2}}\left(\frac{|\xi|^{2}}{1 - |\zeta|^{2}}\right)}} \times \sum_{n=0}^{\infty} \left(-\frac{\zeta\hat{a}^{\dagger 2}}{2}\right)^{n} \left[\frac{L_{n}^{2\ell - \frac{1}{2}}\left(\frac{\xi^{2}}{2\zeta}\right)}{\Gamma\left(n + 2\ell + \frac{1}{2}\right)} + \frac{L_{n}^{2\ell + \frac{1}{2}}\left(\frac{\xi^{2}}{2\zeta}\right)}{\Gamma\left(n + 2\ell + \frac{3}{2}\right)} \frac{\xi\hat{a}^{\dagger}}{2}\right] \Psi_{0,\ell}^{e}(x). \tag{74}$$

Using the results below

$$(\hat{a}^{\dagger})^{2n} \Psi_{0,\ell}^{e}(x) = (-1)^{n} 2^{n} n! L_{n}^{2\ell - \frac{1}{2}} \left(\frac{x^{2}}{l^{2}}\right) \Psi_{0,\ell}^{e}(x) ,$$

$$(\hat{a}^{\dagger})^{2n+1} \Psi_{0,\ell}^{e}(x) = \frac{\sqrt{2}x}{l} (-1)^{n} 2^{n} n! L_{n}^{2\ell + \frac{1}{2}} \left(\frac{x^{2}}{l^{2}}\right) \Psi_{0,\ell}^{e}(x) ,$$

$$(75)$$

we can write (74), as follows

$$\Psi_{\zeta,\xi}^{\ell}(x,t) = \langle x|\zeta,\xi\rangle = \frac{\sqrt{1-|\zeta|^2}}{1-\zeta} \frac{\sqrt{x}}{l} \frac{I_{2\ell-\frac{1}{2}}\left(\frac{\sqrt{2\xi}x}{1-\zeta}\right) + I_{2\ell+\frac{1}{2}}\left(\frac{\sqrt{2\xi}x}{1-\zeta}\right)}{\sqrt{I_{2\ell-\frac{1}{2}}\left(\frac{|\xi|^2}{1-|\zeta|^2}\right) + I_{2\ell+\frac{1}{2}}\left(\frac{|\xi|^2}{1-|\zeta|^2}\right)}} \times \exp\left[-\frac{1+\zeta}{1-\zeta} \frac{x^2}{2l^2} - \frac{(1-\zeta^*)\xi^2}{2(1-\zeta)(1-|\zeta|^2)} + i\tilde{\vartheta}\right].$$
(76)




FIG. 4: Displayed is the probability density by assuming fixed values $l=1, \zeta=0.45$ and $\xi=i$. For $\ell=0$, we have the SCS, while for $\ell>0$, we obtained the generalized para-Bose CS.

In particular, if $\ell = 0$ we obtain the following result:

$$\Psi_{\zeta,\xi}^{0}(x,t) = \frac{\left(1 - |\zeta|^{2}\right)^{1/4}}{\sqrt{\sqrt{\pi}l(1-\zeta)}} \exp\left[-\frac{1}{2l^{2}} \frac{1+\zeta}{1-\zeta} \left(x - \frac{l\sqrt{2}\xi}{1+\zeta}\right)^{2} + \frac{(1+\zeta^{*})}{(1+\zeta)\left(1 - |\zeta|^{2}\right)} \frac{\xi^{2}}{2} - \frac{1}{2} \frac{|\xi|^{2}}{1-|\zeta|^{2}} + i\varrho\right],$$

$$\varrho = \frac{1}{2} \int \left[\operatorname{Re}\left(\alpha\zeta^{*} - \beta\right) - 2\delta\right] dt. \tag{77}$$

Performing the following identifications $\zeta = g/f$, $\xi = z/f = -\varphi/f$ and $\mu = |f|^2 (1 - |\zeta|^2) = 1$ leads to the results of the recent publication [48].

The probability density that corresponds to the generalized CS is given by

$$\rho_{\zeta,\xi}^{\ell}(x,t) = \left|\Psi_{\zeta,\xi}^{\ell}(x,t)\right|^{2} = \frac{1 - |\zeta|^{2}}{|1 - \zeta|^{2}} \frac{x}{l^{2}} \frac{\left|I_{2\ell - \frac{1}{2}}\left(\frac{\sqrt{2}\xi x}{1 - \zeta l}\right) + I_{2\ell + \frac{1}{2}}\left(\frac{\sqrt{2}\xi x}{1 - \zeta l}\right)\right|^{2}}{\left|I_{2\ell - \frac{1}{2}}\left(\frac{|\xi|^{2}}{1 - |\zeta|^{2}}\right) + I_{2\ell + \frac{1}{2}}\left(\frac{|\xi|^{2}}{1 - |\zeta|^{2}}\right)\right|} \times \exp\left[-\frac{1 - |\zeta|^{2}}{|1 - \zeta|^{2}} \frac{x^{2}}{l^{2}} - \frac{1}{1 - |\zeta|^{2}} \operatorname{Re}\left(\frac{1 - \zeta^{*}}{1 - \zeta}\xi^{2}\right)\right].$$
(78)

In Fig. 4, we have obtained some plots of the probability density. As we can see, the probability density of the generalized CS has a shape of a Gaussian distribution, which moves in space as ℓ increases.

Finally, substituting (54) into (4), the Hamiltonian takes the form

$$\hat{H} = \frac{1}{2}\hbar \left(\alpha^* \hat{a}^2 + \alpha \hat{a}^{\dagger 2}\right) + \frac{1}{2}\hbar\beta \left(\hat{a}^{\dagger} \hat{a} + \hat{a} \hat{a}^{\dagger}\right) + \hbar\delta$$

$$= \frac{\hat{P}^2}{2m} + \frac{1}{2}m\omega^2 \hat{x}^2 + \frac{1}{2}\Omega \left(\hat{P}\hat{x} + \hat{x}\hat{P}\right) + \mathcal{E},$$

$$= -\frac{\hbar^2}{2m}\partial_x^2 - i\hbar\Omega x\partial_x + \frac{\hbar^2}{2mx^2}2\ell \left(2\ell - \hat{R}\right) + \frac{1}{2}m\omega^2 x^2 + \mathcal{E} - \frac{i\hbar\Omega}{2},$$
(79)

where the time-dependent quantities m, ω , Ω and \mathcal{E} reads

$$\frac{1}{m} = \frac{l^2}{\hbar} \operatorname{Re}(\beta - \alpha), \quad m\omega^2 = \frac{\hbar}{l^2} \operatorname{Re}(\beta + \alpha), \quad \Omega = \operatorname{Im}(\alpha), \quad \mathcal{E} = \hbar\delta,
\omega^2 = \beta^2 - \operatorname{Re}^2(\alpha).$$
(80)

From Hamiltonian (79), we can describe the physical systems such as harmonic oscillators, which are described by confining potential $V_{HO} \sim x^2$. We also can establish an analogy of the potential $V_C = \frac{\hbar^2}{2mx^2} 2\ell \left(2\ell - \hat{R}\right)$ with the centrifugal potential. On the other hand, one may identify the potential V_C with the conformal sector of quantum conformal mechanics, except by the \hat{R} operator [82]. Since the reflection operator acting on even parity states leads to $\hat{R}\psi_e = \psi_e$, it must be noticed that the potential V_C has a negative correction term owing to the reflection operator. Furthermore, the repulsiveness of the V_C potential is weakened by the correction arising from the reflection operator.

Finally, we can relate the action of the reflection operator on a particular state to the reduction of the dimensionality of space. Let us analyze the centrifugal term of the d-dimensional Laplacian, which can be written in the form [37]:

$$\Delta^{(d)} = \partial_r^2 + \frac{d-1}{r} \partial_r + \frac{L(L+d-2)}{r^2}.$$
 (81)

At this point, we can establish a direct relationship between the action of the reflection operator on even parity states and the reduction of the space from three to one dimension (d=1), from the perspective of the centrifugal term by considering that $L=2\ell$.

VI. TIME-INDEPENDENT PARA-BOSE OSCILLATOR

Since the results obtained in this work can be applied to both time-dependent and time-independent physical systems, for the sake of simplicity, we apply this approach to the study of the time-independent para-Bose oscillator. First, we obtain the Hamiltonian of the time-independent para-Bose oscillator applying the following conditions $\alpha = \delta = \Omega = 0$ and $\beta = \beta_0$ on the Eq. (4), which can be written in the form:

$$\hat{H} = \frac{1}{2}\hbar\beta_0 \left(\hat{a}^{\dagger} \hat{a} + \hat{a} \hat{a}^{\dagger} \right) = \frac{\hbar^2}{2m_0} \left[-\partial_x^2 + \frac{2\ell \left(2\ell - \hat{R} \right)}{x^2} + \frac{m_0^2 \omega_0^2}{\hbar^2} x^2 \right],$$

$$\omega_0 = \beta_0, \quad m_0 = \frac{\hbar}{l^2 \beta_0},$$
(82)

where the subindex labels the initial time, and from (72) we can write the para-Bose number states $|n, \varepsilon\rangle \Longrightarrow |n, \ell\rangle$, such that

$$\hat{H}|n,\ell\rangle = \hbar\omega_0 \left(n + 2\ell + \frac{1}{2}\right)|n,\ell\rangle. \tag{83}$$

Notice that, taking $\ell = 0$ lead to the standard harmonic oscillator.

In this case, the equation system (8), take the form

$$\dot{f} = i\omega_0 f, \quad \dot{g} = -i\omega_0 g, \quad \dot{\varphi} = 0, \tag{84}$$

whose solution is given by

$$f = f_0 e^{i\omega_0 t}, \quad g = g_0 e^{-i\omega_0 t}, \quad \varphi \equiv 0.$$
 (85)

From here, we can write the squeeze and displacement parameters in the form

$$\zeta = \frac{g}{f} = \zeta_0 e^{-2i\omega_0 t}, \quad \xi = \frac{z}{f} = \xi_0 e^{-i\omega_0 t},
\zeta_0 = |\zeta_0| e^{i\theta_\zeta} = \frac{g_0}{f_0}, \quad \xi_0 = |\xi_0| e^{i\theta_\xi} = \frac{z}{f_0}.$$
(86)

In the following, we can rewrite the operators (54):

$$\hat{a} = \frac{\hat{A}_{f} - \zeta_{0}e^{-2i\omega_{0}t}\hat{A}_{f^{*}}^{\dagger}}{1 - |\zeta_{0}|^{2}}, \quad \hat{A}_{f} \equiv \frac{1}{f_{0}e^{i\omega_{0}t}}\hat{A},$$

$$\hat{x} = \frac{l}{\sqrt{2}} \frac{(1 - \zeta_{0}^{*}e^{2i\omega_{0}t})\hat{A}_{f} + (1 - \zeta_{0}e^{-2i\omega_{0}t})\hat{A}_{f^{*}}^{\dagger}}{1 - |\zeta_{0}|^{2}},$$

$$\hat{P} = \frac{\hbar}{i\sqrt{2}l} \frac{(1 + \zeta_{0}^{*}e^{2i\omega_{0}t})\hat{A}_{f} - (1 + \zeta_{0}e^{-2i\omega_{0}t})\hat{A}_{f^{*}}^{\dagger}}{1 - |\zeta_{0}|^{2}},$$
(87)

and the generalized CS (47), as follows

$$|\zeta,\xi\rangle = \left(\frac{|\xi_{0}| e^{i\theta_{\xi}}}{\sqrt{2}}\right)^{2\ell-\frac{1}{2}} \sqrt{\frac{1 - |\zeta_{0}|^{2}}{I_{2\ell-\frac{1}{2}}\left(\frac{|\xi_{0}|^{2}}{1-|\zeta_{0}|^{2}}\right) + I_{2\ell+\frac{1}{2}}\left(\frac{|\xi_{0}|^{2}}{1-|\zeta_{0}|^{2}}\right)} \exp\left[\frac{|\zeta_{0}| |\xi_{0}|^{2} e^{2i\theta_{\xi}-i\theta_{\zeta}}}{2\left(1 - |\zeta_{0}|^{2}\right)}\right] \times \\ \sum_{n=0}^{\infty} (-\zeta_{0})^{n} \sqrt{n!} e^{-2i\omega_{0}(n+\ell)t} \left[\frac{L_{n}^{2\ell-\frac{1}{2}}\left(\frac{|\xi_{0}|^{2}e^{2i\theta_{\xi}}}{2|\zeta_{0}|e^{i\theta_{\zeta}}}\right) e^{-\frac{i\omega_{0}t}{2}}}{\sqrt{\Gamma\left(n+2\ell+\frac{1}{2}\right)}} |2n,\ell\rangle + \frac{|\xi_{0}| e^{i\theta_{\xi}} L_{n}^{2\ell-\frac{1}{2}}\left(\frac{|\xi_{0}|^{2}e^{2i\theta_{\xi}}}{2|\zeta_{0}|e^{i\theta_{\zeta}}}\right) e^{-\frac{3i\omega_{0}t}{2}t}}{\sqrt{2\Gamma\left(n+2\ell+\frac{3}{2}\right)}} |2n+1,\ell\rangle\right].$$

$$(88)$$

From Eq. (56), we get \overline{x} and \overline{P} in the form

$$\overline{x} = \overline{x}_0 \cos(\omega_0 t) + \frac{\overline{P}_0}{m_0 \omega_0} \sin(\omega_0 t), \quad \overline{P} = \overline{P}_0 \cos(\omega_0 t) - m_0 \omega_0 \overline{x}_0 \sin(\omega_0 t),$$

$$\overline{x}_0 = \sqrt{2} l \left| \xi_0 \right| \frac{\cos(\theta_{\xi}) - \left| \zeta_0 \right| \cos(\theta_{\xi} - \theta_{\zeta})}{1 - \left| \zeta_0 \right|^2}, \quad \overline{P}_0 = \sqrt{2} l m_0 \omega_0 \left| \xi_0 \right| \frac{\sin(\theta_{\xi}) + \left| \zeta_0 \right| \sin(\theta_{\xi} - \theta_{\zeta})}{1 - \left| \zeta_0 \right|^2}.$$
(89)

Notice that taking $\xi_0 = 0$, the mean values of \overline{x} and \overline{P} are equal to zero, which corresponds to the mean values evaluated in the generalized SVS.

The uncertainty relation (62) and (65) becomes

$$\sigma_{x}\sigma_{P} = \hbar \sqrt{1 + \frac{4|\zeta_{0}|^{2} \sin^{2}(\theta_{\zeta} - 2\omega_{0}t)}{\left(1 - |\zeta_{0}|^{2}\right)^{2}} \frac{1 + 4\ell \overline{R}}{2}},$$

$$\sigma_{x}^{2}\sigma_{P}^{2} - \sigma_{xP}^{2} = \frac{\hbar^{2}}{4} \left(1 + 4\ell \overline{R}\right)^{2},$$
(90)

with the mean value of \hat{R} is given by the Eq. (60),

$$\overline{R} = \frac{I_{2\ell - \frac{1}{2}} \left(\frac{|\xi_0|^2}{1 - |\zeta_0|^2}\right) - I_{2\ell + \frac{1}{2}} \left(\frac{|\xi_0|^2}{1 - |\zeta_0|^2}\right)}{I_{2\ell - \frac{1}{2}} \left(\frac{|\xi_0|^2}{1 - |\zeta_0|^2}\right) + I_{2\ell + \frac{1}{2}} \left(\frac{|\xi_0|^2}{1 - |\zeta_0|^2}\right)}.$$
(91)

One can see that the uncertainty relation has an oscillatory behavior, reaching minimum values at specific points given by

$$\sin\left(\theta_{\zeta} - 2\omega_0 t_k\right) = 0, \quad t_k = \frac{\theta_{\zeta} - k\pi}{2\omega_0},\tag{92}$$

where $k = 0, \pm 1, \pm 2, \ldots$ Here t_k corresponds to the values on the time for which the uncertainty relation is minimal. The condition $\theta_{\zeta} = 0 \Longrightarrow \zeta_0 = \zeta_0^*$ leads to minimum uncertainty at t = 0. From Eqs. (61), (72) and (90), we can write the l-parameter in term of the quantities $\sigma_x(0) = \sigma_{x_0}$, ζ_0 , ξ_0 and ℓ , as follows

$$l = \sqrt{\frac{1 + \zeta_0}{1 - \zeta_0} \frac{2}{1 + 4\ell \overline{R}}} \sigma_{x_0}, \quad \sigma_{x_0} = \hbar \frac{1 + 4\ell \overline{R}}{2\sigma_{P_0}}.$$
 (93)

In what the following, let us consider ζ_0 as being a real parameter.

It must be highlighted that the para-Bose number states are eigenstates of the Hamiltonian (82), and therefore the standard deviation for this operator is null when evaluated on this basis. Since \hat{H} is time-independent, we have that $|n,\ell\rangle$ are eigenstates of \hat{H} with well-defined energy. Furthermore, it is interesting to verify the probability transition from the states with well-defined energy to generalized CS (88). Therefore, it follows from (52), (72) and (86) that

$$P_{n}(\zeta_{0}, \xi_{0}, \ell) = \left(\frac{|\xi_{0}|^{2}}{2}\right)^{2\ell - \frac{1}{2}} \frac{(1 - \zeta_{0}^{2}) \exp\left[\frac{\zeta_{0}|\xi_{0}|^{2}}{1 - \zeta_{0}^{2}} \cos\left(2\theta_{\xi}\right)\right]}{I_{2\ell - \frac{1}{2}}\left(\frac{|\xi_{0}|^{2}}{1 - \zeta_{0}^{2}}\right) + I_{2\ell + \frac{1}{2}}\left(\frac{|\xi_{0}|^{2}}{1 - \zeta_{0}^{2}}\right)}{n!\zeta_{0}^{2n}} \times \left[\frac{\left|L_{n}^{2\ell - \frac{1}{2}}\left(\frac{|\xi_{0}|^{2}e^{2i\theta_{\xi}}}{2\zeta_{0}}\right)\right|^{2}}{\Gamma\left(n + 2\ell + \frac{1}{2}\right)} + \frac{|\xi_{0}|^{2}\left|L_{n}^{2\ell + \frac{1}{2}}\left(\frac{|\xi_{0}|^{2}e^{2i\theta_{\xi}}}{2\zeta_{0}}\right)\right|^{2}}{2\Gamma\left(n + 2\ell + \frac{3}{2}\right)}\right].$$
(94)




FIG. 5: The probability transition is showed by considering fixed values $\ell = 2$, $|\xi_0| = 1$, and $\theta_{\xi} = \pi/2$. These conditions imply that $\overline{x}_0 = 0$ and $\overline{P}_0 = \sqrt{2lm_0\omega_0} |\xi_0| (1+\zeta_0) / (1-\zeta_0^2)$. In Figure (a) we recover the para-Bose CS. Meanwhile, in Figures (b), (c) and (d) we have the generalized para-Bose CS.

Notice that the probability transition is time-independent. As we saw in the figures Fig. 1 and Fig. 3, the probability transition has its shape significantly altered by the displacement and Wigner parameters. In Fig. 5, we will make an analysis considering some values for the squeeze parameter ζ_0 , keeping the other parameters fixed; namely, ξ_0 and ℓ .

From Eqs. (89), we have that the ζ_0,ξ_0 -parameters are directly related to the initial conditions that lead to the temporal evolution of the mean values in the position and momentum. As we saw, the standard deviation and the uncertainty relations are expressed in terms of special functions, whose analysis is not straightforward. Therefore, it is interesting to investigate the asymptotic form of Bessel functions of the first kind present in uncertainty

relations as follows:

a) Asymptotic form for small arguments Z and fixed κ (see Eq. 9.6.7, page 375 in Ref. [83]).

 $I_{\kappa}(Z) \sim \frac{1}{\Gamma(\kappa+1)} \left(\frac{Z}{2}\right)^{\kappa}.$ (95)

This limit implies that $|\xi_0| \to 0$ and $\zeta_0 < 1$. In its turn, the mean value of the reflection operator \overline{R} can be rewritten:

$$\overline{R} \sim \frac{(4\ell+1)\left(1-\zeta_0^2\right) - |\xi_0|^2}{(4\ell+1)\left(1-\zeta_0^2\right) + |\xi_0|^2},\tag{96}$$

and the uncertainty relations (90) become

$$\sigma_{x}\sigma_{P} \sim \frac{\hbar}{2} \sqrt{1 + \frac{4\zeta_{0}^{2} \sin^{2}(2\omega_{0}t)}{(1 - \zeta_{0}^{2})^{2}} \frac{(4\ell + 1)^{2}(1 - \zeta_{0}^{2}) - (4\ell - 1)|\xi_{0}|^{2}}{(4\ell + 1)(1 - \zeta_{0}^{2}) + |\xi_{0}|^{2}}},$$

$$\sigma_{x}^{2}\sigma_{P}^{2} - \sigma_{xP}^{2} \sim \frac{\hbar^{2}}{4} \left[\frac{(4\ell + 1)^{2}(1 - \zeta_{0}^{2}) - (4\ell - 1)|\xi_{0}|^{2}}{(4\ell + 1)(1 - \zeta_{0}^{2}) + |\xi_{0}|^{2}} \right]^{2}.$$
(97)

From (96), it is easy to see that the parity of the states is even when we take $|\xi_0| = 0$. On the other hand, the value of the uncertainties, Eq. (97), increases as ℓ increases, while for $\zeta_0 = \ell = 0$ is minimized.

b) Asymptotic form for large arguments Z and fixed κ (see Eq. 9.7.1, page 377 in Ref. [83]),

$$I_{\kappa}(Z) \sim \frac{e^Z}{\sqrt{2\pi Z}} \left[1 - \frac{1}{2Z} \left(\kappa^2 - \frac{1}{4} \right) \right].$$
 (98)

This limit can be obtained in two different ways, first it is $|\xi_0| \to \infty$ with $\zeta_0 < 1$ and second $\zeta_0 \to 1$ with $|\xi_0| < \infty$. In this second case, we have a high degree of squeeze in σ_x , which becomes smaller compared to σ_P , as ζ_0 approaches 1. From Eq. (98), we can write the mean value of \overline{R} as follows

$$\overline{R} \sim \frac{\ell (1 - \zeta_0^2)}{|\xi_0|^2 - 2\ell^2 (1 - \zeta_0^2)}.$$
 (99)

Note that the range of values $|\xi_0|^2 < 2\ell^2 (1-\zeta_0^2)$ takes $\overline{R} < 0$, indicating that only odd parity states stay at this limit. The uncertainty relations can be rewritten in the form

$$\sigma_x \sigma_P \sim \frac{\hbar}{2} \sqrt{1 + \frac{4\zeta_0^2 \sin^2(2\omega_0 t)}{(1 - \zeta_0^2)^2} \frac{|\xi_0|^2 + 2\ell^2 (1 - \zeta_0^2)}{|\xi_0|^2 - 2\ell^2 (1 - \zeta_0^2)}},$$

$$\sigma_x^2 \sigma_P^2 - \sigma_{xP}^2 \sim \frac{\hbar^2}{4} \left[\frac{|\xi_0|^2 + 2\ell^2 (1 - \zeta_0^2)}{|\xi_0|^2 - 2\ell^2 (1 - \zeta_0^2)} \right]^2.$$
(100)

It is easy to see that $\ell = 0$ lead the uncertainty relations to the value obtained when evaluated in terms of the canonical commutation relation $([\hat{x}, \hat{p}] = i\hbar, \hat{p} = -i\hbar\partial_x)$. Besides, it can be seen that the squeeze parameter ζ_0 ensures that the Heisenberg uncertainty relation oscillates over time.

This example illustrates how this general procedure can be applied to a range of problems by simply determining the parameters of the Hamiltonian, then finding the functions f and g. From these functions, we obtain the squeeze and displacement parameters that, in its turn, modify the uncertainty relation and probability transition.

VII. CONCLUDING REMARKS

In this article, we study the integrals of motion method in a para-Bose formulation. This approach generalizes the usual canonical commutation relation. In turn, we obtain a generalization of the usual SVS, which admits a completeness relation in terms of the Wigner parameter. This relation depends on a range of values for the Wigner parameter, which does not include that corresponding to canonical algebra. We also obtain a generalization of the CS in terms of the even and odd time-independent para-Bose number states. These states are thoroughly determined in terms of the time-dependent squeeze and displacement parameters, as well as the Wigner parameter. In the study of the probability transition, we saw that the displacement parameter has an additional role, which is a kind of transition parameter by allowing access to the odd states of the system. Meanwhile, the Wigner parameter has the role of controlling the "dispersion," and attenuating the access to the odd states. We show that the minimization of the Heisenberg uncertainty relation is easily obtained by taking the real value of the squeeze parameter and that the squeezing properties can be seen from the standard deviation of the position and momentum. Taking the coordinate representation of the generalized CS, we found a quantization condition on the Wigner parameter, analogous to the quantization of the angular momentum, which arises by imposing that the parity of the vacuum state is even. This quantization condition also ensures that the eigenstates of the momentum operator are differentiable at the origin. Finally, the para-Bose oscillator has been discussed in this framework.

Acknowledgments

We would like to thank CNPq, CAPES and CNPq/PRONEX/FAPESQ-PB (Grant no. 165/2018), for partial financial support. ASL and FAB acknowledge support from CNPq (Grant nos. 150601/2021-2, 312104/2018-9). ASP thanks the support of the Instituto Federal do Pará.

Data Availability Statement

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

- [1] Schrödinger E 1926 Der stetige übergang von der mikro-zur makromechanik Naturwissenschaften, Bd. 14, H. 28, 664.
- [2] Glauber R J 1963 Coherent and incoherent states of the radiation field Phys. Rev. Lett. 131, 2766.
- [3] Sudarshan E C G 1963 Equivalence of semiclassical and quantum mechanical descriptions of statistical light beams Phys. Lett. 10, 277.
- [4] Klauder J R 1963 Continuous-representation Theory. II. Generalized relation between quantum and classical dynamics J. Math. Phys. 4, 1058.
- [5] Scully M O and Zubairy M S 1997 Quantum Optics (Cambridge: Cambridge University Press).
- [6] Klauder J R and Sudarshan E C 1968 Fundamentals of Quantum Optics (New York: Benjamin).
- [7] Nielsen M and Chuang I 2000 Quantum Computation and Quantum Information (Cambridge: Cambridge University Press).
- [8] Klauder J R and Skagerstam B S 1985 Coherent States, Applications in Physics and Mathematical Physics (Singapore: World Scientific).
- [9] Bachor H A and Ralph T C 2004 A Guide to Experiments in Quantum Optics (Wiley-Vch).
- [10] Schnabel R 2017 Squeezed states of light and their applications in laser interferometers Phys. Rept. 684, 1.
- [11] Dey S 2021 An introductory review on resource theories of generalized nonclassical light J. Phys.: Conf. Ser. 2038, 012008.

- [12] Stoler D 1970 Equivalence Classes of Minimum Uncertainty Packets Phys. Rev. D 1, 3217.
- [13] Stoler D 1971 Equivalence Classes of Minimum-Uncertainty Packets. II Phys. Rev. D 4, 1925.
- [14] Slusher R E and Yurke B 1990 Squeezed light for coherent communications J. Light. Tech. 8 466.
- [15] Caves C M 1981 Quantum-mechanical noise in an interferometer Phys. Rev. D 23, 1693.
- [16] Ni W-T 1987 Quantum-mechanical noise in an interferometer: Intrinsic uncertainty versus measurement uncertainty Phys. Rev. D 35, 3002.
- [17] Chua S S Y, Slagmolen B J J, Shaddock D A and McClelland D E 2014 Quantum squeezed light in gravitational-wave detectors Class. Quant. Grav. 31, 183001.
- [18] Gerry C and Knight P 1983 Introductory Quantum Optics (Cambridge: Cambridge University Press).
- [19] Malkin I A and Man'ko V I 1979 Dynamical Symmetries and Coherent States of Quantum Systems (Moscow: Nauka).
- [20] Dodonov V V and Man'ko V I 2003 Theory of Nonclassical States of Light (London: Taylor & Fran-cis Group).
- [21] Perelomov A M 1986 Generalized Coherent States and their Applications (Verlag: Springer).
- [22] Gilmore R 1972 Geometry of symmetrized states Ann. Phys. (NY) 74, 391.
- [23] Ali S T, Antoine J P and Gazeau J P 2000 Coherent States, Wavelets and Their Generalizations (New York: Springer).
- [24] Yuen H P 1976 Two-photon coherent states of the radiation field Phys. Rev. A 13, 2226.
- [25] Satyanarayana M V 1985 Generalized coherent states and generalized squeezed coherent states Phys. Rev. D 32, 400.
- [26] Dey S and Fring A 2012 Squeezed coherent states for noncommutative spaces with minimal length uncertainty relations Phys. Rev. D 86, 064038.
- [27] Pedram P 2013 Coherent states in gravitational quantum mechanics Int. J. Mod. Phys. D 22, 1350004.
- [28] Patra P, Saha J P and Biswas K 2022 Squeezed coherent states for gravitational well in non-commutative space Indian J. Phys. 96, 309.
- [29] Man'ko V I, Marmo G, Sudarshan E C G and Zaccaria F 1997 f-oscillators and nonlinear coherent states Phys. Scr. 55, 528.
- [30] Dey S 2015 q-deformed noncommutative cat states and their nonclassical properties Phys. Rev.

- D 91, 044024.
- [31] Eremin V V and Meldianov A A 2006 The q-deformed harmonic oscillator, coherent states, and the uncertainty relation Theor. Math. Phys. 147, 709.
- [32] Agarwal G S and Chaturvedi S 1995 Calogero-Sutherland oscillator: classical behaviour and coherent states J. Phys. A: Math and Gen 28, 5747.
- [33] Brink L, Hansson T H, Konstein S and Vasiliev M A 1993 The Calogero model anyonic representation, fermionic extension and supersymmetry Nucl. Phys. B 401, 591.
- [34] Wigner E P 1950 Do the equations of motion determine the quantum mechanical commutation relations? Phys. Rev. 77, 711.
- [35] Plyushchay M S 1997 Deformed Heisenberg algebra with reflection Nucl. Phys. B 491, 619.
- [36] Mukunda N, Sudarshan E C G, Sharma J K and Mehta C L 1980 Representations and properties of para-Bose oscillator operators. I. Energy position and momentum eigenstates J. Math. Phys. 21, 2386.
- [37] Lohe M A and Thilagam A 2004 Quantum mechanical models in fractional dimensions J. Phys. A: Math. Gen. 37, 6181.
- [38] Sharma J K, Mehta C L, Mukunda N and Sudarshan E C G 1981 Representation and properties of para-Bose oscillator operators. II. Coherent states and the minimum uncertainty states J. Math. Phys. 22, 78.
- [39] Hartle J B and Taylor J R 1968 Quantum mechanics of paraparticles Phys. Rev. 178, 2043.
- [40] Green H S 1953 A generalized method of field quantization Phys. Rev. 90, 270.
- [41] Ohnuki Y and Kamefuchi S 1982 Quantum Field Theory and Parastatistics (New York: Springer).
- [42] Stoilova N I and Jeugt J V 2010 Parabosons, parafermions, and explicit representations of infinite-dimensional algebras Phys. Atom. Nuclei 73, 533.
- [43] Greenberg O W and Messiah A M L 1965 Selection rules for parafields and the absence of para particles in nature Phys. Rev. 138, B1155.
- [44] Kitabayashi T and Yasuè M 2018 Parafermionic dark matter Phys. Rev. D 98, 043504.
- [45] Bracken A J and Green H S 1973 Parastatistics and the quark model J. Math. Phys. 14, 1784.
- [46] Hama M, Sawamura M and Suzuki H 1992 Thermodynamical properties of high order parabosons Prog. Theor. Phys. 88, 149.
- [47] Rodríguez-Walton S, Ávila B J and Rodríguez-Lara B M 2020 Optical non-Hermitian para-

- Fermi oscillators Phys. Rev. A 101, 043840.
- [48] Pereira A S and Lemos A S 2021 Time-dependent coherent squeezed states in a nonunitary approach Phys. Lett. A 405, 127428.
- [49] Meyer H E and Berghe G V 1980 Generalised Bogoliubov transformation coefficients for para-Bose states J. Phys. A: Math. Gen. 13, 161.
- [50] Jordan T F, Mukunda N and Pepper S V 1963 Irreducible representations of generalized oscillator operators J. Math. Phys. 4, 1089.
- [51] Sharma J K, Metha C L and Sudarshan E C G 1978 Para-Bose coherent states J. Math. Phys. 19, 2089.
- [52] Biswas S N and Santhanam T S 1980 Coherent states of para-Bose oscillators J. Austral. Math. Soc. 22, 210.
- [53] Saxena G M and Mehta C L 1986 Dynamics of para-Bose coherent states J. Math. Phys. 27, 309.
- [54] Mojaveri B, Dehghani A and Bahrbeig R J 2018 Nonlinear coherent states of the para-Bose oscillator and their non-classical features Eur. Phys. J. Plus 133, 529.
- [55] Mojaveri B, Dehghani A and Bahrbeig R J 2018 Excitation on the para-Bose states: nonclassical properties Eur. Phys. J. Plus 133, 346.
- [56] Alderete C H and Vergara L V and Rodríguez-Lara B M 2017 Nonclassical and semiclassical para-Bose states Phys. Rev. A 95, 043835.
- [57] Bagchi B and Bhaumik D 1997 Coherent states for parabosons J. Phys. A: Math. Gen. 30, L593.
- [58] Dehghani A, Mojaveri B, Shirin S and Saedi M 2015 Cat-states in the framework of Wigner-Heisenberg algebra Ann. Phys. 362, 659.
- [59] Calogero F 1971 Solution of the one-dimensional N-body problems with quadratic and/or inversely quadratic pair potentials Jour. Math. Phys. 12, 419.
- [60] Simons B D, Lee P A and Altshuler B L 1994 Matrix models, one-dimensional fermions, and quantum chaos Phys. Rev. Lett. 72, 64.
- [61] Kawakami N 1993 Novel hierarchy of the SU(N) electron models and edge states of fractional quantum Hall effect Phys. Rev. Lett. 71, 275.
- [62] Azuma H and Iso S 1994 Explicit relation of the quantum Hall effect and the Calogero-Sutherland model Phys. Lett. B 331, 107.

- [63] Ouvry S 2001 On the relation between the anyon and Calogero models Phys. Lett. B 510, 335.
- [64] Caselle M 1995 Distribution of transmission eigenvalues in disordered wires Phy. Rev. Lett. 74, 2776.
- [65] Alderete C H, Green A M, Nguyen N H, Zhu Y, Rodríguez-Lara B M and Linke N M 2021 Experimental realization of para-particle oscillators [arXiv:2108.05471 [quant-ph]].
- [66] Alderete C H and Rodríguez-Lara B M 2017 Quantum simulation of driven para-Bose oscillators Phys. Rev. A 95, 013820.
- [67] Lewis H R Jr. and Riesenfeld W B 1969 An exact quantum theory of the time-dependent harmonic oscillator and of a charged particle in a time-dependent electromagnetic field. J. Math. Phys. 10, 1458.
- [68] Dodonov V V, Malkin I A and Man'ko V I 1975 Integrals of the motion, green functions, and coherent states of dynamical systems Int. J. Theor. Phys. 14, 37.
- [69] Bagrov V G, Gitman D M and Pereira A S 2015 Coherent states of systems with quadratic Hamiltonians Braz. J. Phys. 45, 369.
- [70] Mark Andrews 1999 Invariant operators for quadratic Hamiltonians Am. J. Phys. 67, 336.
- [71] Dodonov V V and Man'ko V I 1989 Evolution of multidimensional systems. Magnetic properties of ideal gases of charged particles. In Invariants and the Evolution of Nonstationary Quantum Systems (Proceedings of Lebedev Physics Institute, Volume 183); Markov, M.A., Ed.; Nova Science: Commack, NY, USA; pp. 263.
- [72] Man'ko V I and Mendes R V 1997 Time-dependent invariants for Dirac equation and Newton–Wigner position operator Phys. Scr. 56, 417.
- [73] Mostafazadeh A 2001 Supersymmetric dynamical invariants J. Phys. A Math. Gen. 34, 4493.
- [74] Quesne C 2001 Completeness of photon-added squeezed vacuum and one-photon states and of photon-added coherent states on a circle Phys. Lett. A 288, 241.
- [75] Chaturvedi S 1996 A direct proof of the completeness of squeezed odd number states Mod. Phys. Lett. A 11, 2805.
- [76] Glauber R J 1966 Classical behavior of system of quantum oscillators Phys. Lett. 21, 650.
- [77] Mehta C L and Sudarshan E C G 1966 Time evolution of coherent states Phys. Lett. 22, 574.
- [78] Robertson H P 1930 A general formulation of the uncertainty principle and its classical interpretation Phys. Rev. 35, 667.
- [79] Hillery M, O'Connell R F, Scully M O and Wigner E P 1984 Distribution functions in physics:

- fundamentals Phys. Rep. 106, 121.
- [80] Coffey W T, Kalmykov Y P and Titov S V 2007 Solution of the master equation for Wigner's quasiprobability distribution in phase space for the Brownian motion of a particle in a double well potential J. Chem. Phys 127, 074502.
- [81] Olivares S 2012 Quantum optics in the phase space Eur. Phys. J. Special Topics 203, 3.
- [82] Andrzejewski K 2016 Quantum conformal mechanics emerging from unitary representations of $SL(2,\mathbb{R})$ Annals Phys. 367, 227.
- [83] Abramowitz M and Stegun I A 1965 Handbook of mathematical functions (New York: Dover).