

GENERALIZED SOLUTIONS FOR THE MEAN CURVATURE EQUATION

ENRICO GIUSTI*

The purpose of this paper is to discuss general boundary value problems for the mean curvature equation

$$(0.1) \quad \operatorname{div} Tu = H(x, u); \quad Tu = Du/\sqrt{1 + |Du|^2}$$

in a bounded domain $\Omega \subset \mathbf{R}^n$. More precisely, we shall consider the problem of minimizing the functional

$$(0.2) \quad \mathcal{F}(u) = \int_{\Omega} \sqrt{1 + |Du|^2} + \int_{\Omega} \lambda(x, u) dx + \int_{\partial\Omega} \kappa(x, u) dH_{n-1}$$

where

$$(0.3) \quad \lambda(x, u) = \int_0^u H(x, t) dt.$$

It is easily seen that (0.1) is the Euler equation of the functional \mathcal{F} . The third integral in (0.2) describes the boundary conditions: if $u \in C^1(\bar{\Omega})$ and κ is of class C^1 we have

$$(0.4) \quad Tu \cdot \nu = \gamma(x, u) \quad \text{on } \partial\Omega$$

where ν denotes the interior normal to $\partial\Omega$, and

$$(0.5) \quad \kappa(x, u) = \int_0^u \gamma(x, t) dt.$$

When κ is not differentiable, as it is the case for the Dirichlet problem, condition 0.4 does not hold any longer, and we have instead the weaker condition

$$(0.6) \quad \gamma^-(x, u) \leq Tu \cdot \nu \leq \gamma^+(x, u)$$

on $\partial\Omega$, where

$$\gamma^\pm(x, u) = \lim_{t \rightarrow u^\pm} \gamma(x, t).$$

For example, for the Dirichlet problem with boundary datum $f(x)$ we have

$$\kappa(x, u) = |u - f(x)| - |f(x)|$$

and

$$\gamma(x, t) = 1 - 2\varphi_F(x, t)$$

where φ_F is the characteristic function of the subgraph F of f :

$$\varphi_F(x, t) = \begin{cases} 1 & \text{if } t < f(x) \\ 0 & \text{if } t \geq f(x) \end{cases}.$$

In this case we get the boundary conditions

$$Tu \cdot \nu = \begin{cases} 1 & \text{if } u(x) > f(x) \\ -1 & \text{if } u(x) < f(x) \\ \text{arbitrary} & \text{if } u(x) = f(x) \end{cases}.$$

Two standard methods have been developed for the study of boundary value problems. The first one consists in looking for a classical solution, i.e., a smooth function $u(x)$ satisfying equation (0.1) and the boundary conditions (0.4), or more generally (0.6). Alternatively, one may try to minimize directly the functional (0.2) in $BV(\Omega)$, the space of functions with bounded variation in Ω .

Both such approaches suffer serious limitations; in particular the variational method is not adequate when dealing with problems whose solutions may have infinite area, as it is the case for the Dirichlet problem with infinite data or in unbounded domains.

Recently, M. Miranda [10] has introduced the notion of generalized solutions for the minimal surface equation, and has used it successfully in the Dirichlet problem in infinite domains [10], and in the problem of removable singularities [11], two questions in which the area of the solution is not finite, at least in principle. The same notion of generalized solution has been used by U. Massari [8] in his paper on Dirichlet's problem with infinite data, and by the author [7] in the problem of maximal domains for the mean curvature equation.

The idea of generalized solutions originates from the observation that a function $u \in BV(\Omega)$ is a variational solution of \mathcal{F} if and only if its subgraph

$$U = \{(x, t) \in \Omega \times \mathbf{R}: t < u(x)\}$$

minimizes the functional

$$F(U) = \int_{\Omega \times R} |D\varphi_U| + \int_{\Omega \times R} H\varphi_U dx dt + \int_{\partial\Omega \times R} \gamma\varphi_U dH_n.$$

The subgraph of u has the property that the intersection of any vertical straight line with U , if it is not empty, is either the whole line or a lower half-line. Conversely, every set U with the above property is the subgraph of a function $u(x)$, taking possibly the values $+\infty$ or $-\infty$. Such a function is called a generalized solution if U is a local minimum of F .

The interest in the above definition comes mainly from the fact

that under extremely mild hypotheses on H and γ , the set of generalized solutions is compact. More precisely, from every sequence u_j of generalized solutions it is possible to extract a convergent subsequence, in the sense that the subgraphs U_j converge locally in $\Omega \times \mathbf{R}$ to a subgraph U (Lemma 1.2). Of course, the same is not true for variational solutions, for which one needs at least a uniform estimate in $L^1_{\text{loc}}(\Omega)$.

This paper deals with generalized solutions for the functional (0.2). In the first place we show the existence of such solutions, under very general assumptions for H and γ . Of course, generalized solutions may take the values $\pm\infty$; in other words the sets

$$P = \{x \in \Omega : u(x) = +\infty\}$$

and

$$N = \{x \in \Omega : u(x) = -\infty\}$$

may be nonempty. In § 2 we study the properties of these singular sets, showing that they must minimize two functionals related to F .

In § 3 we discuss more closely the relations between generalized and variational solutions and we prove that under suitable assumptions the sets P and N are empty and therefore the generalized solutions are variational. In the same section we show how a number of problems treated by various authors may find their natural place in this general setting.

As an application, we discuss in § 4 Lagrange multipliers. This problem has been studied by C. Gerhardt [4] in the case of capillarity, and by G. Williams for Dirichlet's boundary conditions [12], in the framework of variational solutions. The existence of variational solutions being not guaranteed in principle by the hypotheses, both authors introduce a perturbed functional (for which existence is granted) and then let the perturbation vanish. The use of the notion of generalized solutions, avoiding this complication and dealing directly with the original functional, permits a considerable simplification of the proof, and a generalization of the results.

1. Existence of generalized solutions. Throughout this paper we shall be concerned with the functional

$$(1.1) \quad \mathcal{F}(u) = \int_{\Omega} \sqrt{1 + |Du|^2} + \int_{\Omega} \lambda(x, u) dx + \int_{\partial\Omega} \kappa(x, u) dH_{n-1}$$

where Ω is a bounded domain in \mathbf{R}^n with smooth boundary $\partial\Omega$, and λ, κ are convex functions of u . We may suppose that $\lambda(x, 0) = \kappa(x, 0) = 0$, and therefore

$$\lambda(x, u) = \int_0^u H(x, t) dt ; \quad \kappa(x, u) = \int_0^u \gamma(x, t) dt$$

for some functions H, γ , nondecreasing in t for almost every x .

Related to (1.1) we define a second functional, operating on subsets of the cylinder $Q = \Omega \times \mathbf{R}$.

For $T > 0$ let us set

$$Q_T = \Omega \times [-T, T] \\ \partial Q_T = \partial \Omega \times [-T, T],$$

and for $U \subset Q$:

$$(1.2) \quad F_T(U) = \int_{\partial U} |D\varphi_U| + \int_{Q_T} H(x, t)\varphi_U dxdt + \int_{\partial Q_T} \gamma(x, t)\varphi_U dH_n.$$

DEFINITION 1.1. A set $A \subset Q$ is a supersolution in Q_T for the functional F if for every set $S \subset Q_T$ we have

$$F_T(A) \leqq F_T(A \cup S).$$

The set A is a subsolution in Q_T if for every set $S \subset Q_T$:

$$F_T(A) \leqq F_T(A - S).$$

Finally, A is a solution in Q_T if it is both a super- and a subsolution.

DEFINITION 1.2. A set $A \subset Q$ is a local solution for F [super-solution, subsolution] in Q if it is a solution [supersolution, subsolution] in Q_T for every $T > 0$.

The connection between the functionals (1.1) and (1.2) is apparent from the following theorem.

THEOREM 1.1. (M. Miranda [9]) *A function $u \in BV(\Omega)$ is a variational solution [supersolution, subsolution] for if and only if its subgraph*

$$U = \{(x, t) \in Q : t < u(x)\}$$

is a local solution [supersolution, subsolution] for F in Q .

The above result is the starting point for the definition of generalized solutions for the functional (1.1). Suppose we have a local solution U of the functional F , and that almost every vertical line crosses the boundary of U at most once. Then U will be the subgraph of some function $u(x)$ in Ω , taking possibly the values $+\infty$ or $-\infty$. We call such function u a generalized solution for the

functional \mathcal{F} .

DEFINITION 1.3. (M. Miranda [10]) A function $u: \Omega \rightarrow [-\infty, +\infty]$ is a generalized solution [supersolution, subsolution] for the functional \mathcal{F} if its subgraph U is a local solution [supersolution, subsolution] of the related functional F .

We shall devote the rest of this section to the proof of the existence of a generalized solution of \mathcal{F} .

LEMMA 1.1. *Let $\partial\Omega$ be a C^2 manifold. For each $\varepsilon > 0$ there exists a constant $c_1(\varepsilon)$ such that for every $w \in BV(\Omega)$ we have*

$$(1.3) \quad \int_{\partial\Omega} |w| dH_{n-1} \leq \int_{\Sigma_\varepsilon} |Dw| + c_1 \int_{\Sigma_\varepsilon} |w| dx$$

where

$$\Sigma_\varepsilon = \{x \in \Omega : \text{dist}(x, \partial\Omega) \leq \varepsilon\}.$$

Proof. Suppose first that $w \geq 0$, and let η be a C^∞ -function, $0 \leq \eta \leq 1$, $\eta = 1$ on $\partial\Omega$ and $\eta = 0$ in $\Omega - \Sigma_\varepsilon$. Since $\partial\Omega$ is of class C^2 the distance function $d(x) = \text{dist}(x, \partial\Omega)$ is of class C^2 in a neighborhood of $\partial\Omega$. We may suppose of course that ε is so small that $d \in C^2(\Sigma_\varepsilon)$. We have

$$\int_{\Omega} w \operatorname{div}(\eta Dd) dx = - \int_{\Omega} \eta Dd Dw + \int_{\partial\Omega} w \eta \nu \cdot Dd dH_{n-1}$$

and since $\nu \cdot Dd = -1$ and $\eta = 1$ on $\partial\Omega$:

$$\int_{\partial\Omega} w dH_{n-1} = - \int_{\Omega} \eta Dd Dw - \int_{\Omega} w \operatorname{div}(\eta Dd) dx \leq \int_{\Sigma_\varepsilon} |Dw| + c_1(\varepsilon) \int_{\Sigma_\varepsilon} w dx$$

where $c_1(\varepsilon) = \sup_{\Omega} |\operatorname{div}(\eta Dd)|$ depends only on ε and Ω . This proves (1.3) when $w \geq 0$. The general case follows from the inequality

$$\int_{\Omega} |Dw| \leq \int_{\Omega} |Dw|.$$

We may now prove

PROPOSITION 1.1. *Let the function $\kappa(x, u)$ satisfy*

$$(1.4) \quad |\kappa(x, u) - \kappa(x, v)| \leq |u - v|$$

for H_{n-1} almost every $x \in \partial\Omega$. Then the functional

$$\mathcal{G}(u) = \int_{\Omega} \sqrt{1 + |Du|^2} + \int_{\partial\Omega} \kappa(x, u) dH_{n-1}$$

is lower semi-continuous with respect to L^1 convergence.

Proof. Let $u_j \rightarrow u$ in $L^1(\Omega)$. We have from (1.4):

$$\mathcal{G}(u) - \mathcal{G}(u_j) \leq \int_{\Omega} \sqrt{1 + |Du|^2} - \int_{\Omega} \sqrt{1 + |Du_j|^2} + \int_{\partial\Omega} |u - u_j| dH_{n-1}$$

and from Lemma 1.1 with $w = u - u_j$:

$$\begin{aligned} \mathcal{G}(u) - \mathcal{G}(u_j) &\leq \int_{\Omega} \sqrt{1 + |Du|^2} - \int_{\Omega} \sqrt{1 + |Du_j|^2} + \int_{\Sigma_{\varepsilon}} |Du| \\ &\quad + \int_{\Sigma_{\varepsilon}} |Du_j| + c_1 \int_{\Sigma_{\varepsilon}} |u - u_j| dx \leq \int_{\Sigma_{\varepsilon}} \sqrt{1 + |Du|^2} \\ &\quad - \int_{\Sigma_{\varepsilon}} \sqrt{1 + |Du_j|^2} + 2 \int_{\Sigma_{\varepsilon}} \sqrt{1 + |Du|^2} \\ &\quad + c_1 \int_{\Sigma_{\varepsilon}} |u - u_j| dx, \end{aligned}$$

where

$$\Omega_{\varepsilon} = \Omega - \Sigma_{\varepsilon} = \{x \in \Omega : \text{dist}(x, \partial\Omega) > \varepsilon\}.$$

Let now $j \rightarrow \infty$; taking into account the lower semi-continuity of the area with respect to L^1 convergence we get:

$$\mathcal{G}(u) - \liminf_{j \rightarrow \infty} \mathcal{G}(u_j) \leq 2 \int_{\Sigma_{\varepsilon}} \sqrt{1 + |Du|^2}$$

and the result follows letting $\varepsilon \rightarrow 0$.

REMARK 1.1. An assumption equivalent to (1.4) is obviously

$$(1.5) \quad |\gamma(x, t)| \leq 1 \quad H_n - \text{a.e. on } \partial Q.$$

We have proved the lower semi-continuity of part of the functional $\mathcal{G}(u)$. For what concerns the curvature term we refer to [5] where we have proved its lower semi-continuity with respect to strong convergence in L^1 and weak convergence in $L^{n/n-1}$, under the hypothesis that $H(x, t)$ is increasing in t , and belongs to $L^n(\Omega)$ for every $t \in \mathbf{R}$.

We have in conclusion the following theorem:

THEOREM 1.2. *Let Ω be a bounded domain with C^2 boundary $\partial\Omega$, and let $H(x, t)$ and $\gamma(x, t)$ be two functions defined in Ω and $\partial\Omega$ respectively, and satisfying the following assumptions:*

- (H₁) $H(x, \cdot)$ is nondecreasing for almost every $x \in \Omega$.
- (H₂) $H(\cdot, t)$ belongs to $L^n(\Omega)$ for every t .
- (γ₁) $\gamma(x, \cdot)$ is nondecreasing for H_{n-1} -almost every $x \in \Omega$.

$$(\gamma_2) \quad |\gamma(x, t)| \leq 1 \quad H_n - \text{a.e. in } \partial Q.$$

Then for every sequence u_j , bounded in $L^{n/n-1}$ and convergent in L^1 to a function u we have:

$$\mathcal{F}(u) \leq \liminf_{j \rightarrow \infty} \mathcal{F}(u_j).$$

In order to get a generalized solution for $\mathcal{F}(u)$ we begin by minimizing this functional in the class

$$V_j = \{v \in BV(Q): |v| \leq j\}.$$

It is easily seen that \mathcal{F} is bounded from below in V_j , and that every minimizing sequence is bounded in $BV(Q)$. From well-known compactness theorems we may extract a subsequence converging in $L^1(Q)$; on the other hand it is obvious that sequence will be bounded in $L^{n/n-1}$, so that we may apply Theorem 1.2 to conclude the existence of a minimum for \mathcal{F} in V_j .

Let us denote by u_j a minimizing function. The subgraph U_j is a solution for F in Q_j . We shall now let $j \rightarrow \infty$ to show the existence of a local solution to F in Q , and whence of a generalized solution for \mathcal{F} . For that we need the following lemma:

LEMMA 1.2. Let A be a subsolution for F in Q_T . Then

$$(1.6) \quad \int_{Q_T} |D\varphi_A| \leq c_2(T).$$

Proof. We have

$$\begin{aligned} F_T(A) &= \int_{Q_T} |D\varphi_A| + \int_{Q_T} H\varphi_A dxdt + \int_{\partial Q_T} \gamma\varphi_A dH_n \leq F_T(A - Q_T) \\ &= \int_{Q_T} |D\varphi_{A-Q_T}| \leq 2|\Omega| \end{aligned}$$

and therefore

$$\int_{Q_T} |D\varphi_A| \leq 2|\Omega| + \int_{Q_T} |H| dxdt + H_n(\partial Q_T) = c_2(T).$$

In a similar way, comparing with $A \cup Q_T$, we prove (1.6) for supersolutions. In particular, (1.6) holds for solutions in Q_T .

The inequality (1.6) is the only estimate we need in order to pass to the limit as $j \rightarrow \infty$. For, let $T > 0$ and let $j > T$. Since U_j is a solution in Q_T , we have

$$\int_{Q_T} |D\varphi_{U_j}| \leq c_2(T)$$

and therefore it is possible to extract a subsequence, which we shall denote again by U_j , converging to some set U in every Q_T . It is clear that U , being limit of subgraphs, is itself a subgraph of some function $u(x)$, assuming possibly the values $\pm\infty$. It follows from the next proposition that U is a local minimum for F and hence that u is a generalized solution for \mathcal{F} .

PROPOSITION 1.2. *Let H_j and γ_j be two nondecreasing sequences converging to H and γ respectively, and let U_j minimize the functional*

$$F_T^j(A) = \int_{Q_T} |D\varphi_A| + \int_{Q_T} H_j \varphi_A dx dt + \int_{\partial Q_T} \gamma_j \varphi_A dH_n .$$

Suppose that $U_j \rightarrow U$ in Q_T and that

$$(1.7) \quad \int_{\tilde{\partial}Q_T} |\varphi_{U_j} - \varphi_U| dH_n \longrightarrow 0 , \quad \tilde{\partial}Q_T = \partial Q_T - \partial Q_T .$$

Then U minimizes F_T in Q_T .

Proof. Let $V \subset Q$, $V = U$ outside Q_T , and let

$$V_j = \begin{cases} V & \text{in } Q_T \\ U_j & \text{outside } Q_T . \end{cases}$$

We have $F_T^j(U_j) \leqq F_T^j(V_j)$ and therefore

$$\begin{aligned} & \int_{Q_T} |D\varphi_{U_j}| + \int_{Q_T} H_j \varphi_{U_j} dx dt + \int_{\partial Q_T} \gamma_j \varphi_{U_j} dH_n \\ & \leqq \int_{Q_T} |D\varphi_{V_j}| + \int_{Q_T} H_j \varphi_V dx dt + \int_{\partial Q_T} \gamma_j \varphi_V dH_n . \end{aligned}$$

On the other hand

$$\int_{Q_T} |D\varphi_{V_j}| \leqq \int_{Q_T} |D\varphi_V| + \int_{\tilde{\partial}Q_T} |\varphi_U - \varphi_{V_j}| dH_n$$

and in conclusion, if $j > s$:

$$\begin{aligned} & \int_{Q_T} |D\varphi_{U_j}| + \int_{Q_T} H_s \varphi_{U_j} dx dt + \int_{\partial Q_T} \gamma \varphi_{U_j} dH_n + \int_{\partial Q_T} (\gamma_j - \gamma) \varphi_{U_j} dH_n \\ & \leqq F_T^j(V) + \int_{\tilde{\partial}Q_T} |\varphi_U - \varphi_{V_j}| dH_n . \end{aligned}$$

Passing to the limit as $j \rightarrow \infty$:

$$\int_{Q_T} |D\varphi_U| + \int_{Q_T} H_s \varphi_U dx dt + \int_{\partial Q_T} \gamma \varphi_U dH_n \leqq F_T(V)$$

and the conclusion follows letting $s \rightarrow \infty$.

We may now apply the preceding proposition with $H_j = H$ and $\gamma_j = \gamma$; the condition (1.7) being satisfied for almost every T . In conclusion, we have proved the following existence theorem:

THEOREM 1.3. *Let Ω , γ and H satisfy the hypotheses of Theorem 1.2. Then there exists a generalized solution for the functional $\mathcal{F}(u)$.*

REMARK 1.2. Though proved for free solutions, the above theorem remains valid if some additional conditions are imposed on u . In particular, the same proof works for the obstacle problem, i.e., when u is restricted by the conditions

$$\psi_1(x) \leq u(x) \leq \psi_2(x),$$

with $\psi_1(x)$ bounded from above, and $\psi_2(x)$ from below; as well as for the “soft obstacle” problem, namely when

$$a \leq \int_{\Omega} \zeta(x)u(x)dx \leq b$$

for a given positive function $\zeta(x)$. The above includes the problem with fixed volume, when $\zeta = 1$ and $a = b$.

Combinations of these and other conditions may also be imposed, as long as they are compatible with $|u| \leq T$ for large T .

2. The structure of the sets P and N . The generalized solution u may well take the values $+\infty$ and $-\infty$. We set

$$\begin{aligned} P &= \{x \in \Omega : u(x) = +\infty\} \\ N &= \{x \in \Omega : u(x) = -\infty\}. \end{aligned}$$

The purpose of this section is to study the properties of the above sets.

Since we want to treat the obstacle problem, and even other situations such as the soft obstacles, we begin by observing that Proposition 1.2 remains valid, with the same proof, if U_j are subsolutions, provided we add to the hypotheses the assumption that U_j form a monotone decreasing sequence: $U_j \supset U_{j+1}$. In this case the limit U will be a subsolution itself.

We shall concentrate on the set P ; we remark however that if u is a supersolution for \mathcal{F} and if we set $H'(x, t) = -H(x, -t)$ and $\gamma'(x, t) = -\gamma(x, -t)$, the function $-u$ is a subsolution for \mathcal{F}' , and hence every result concerning the set P can be translated at once

into a similar result concerning the set N for supersolutions.

PROPOSITION 2.1. *Let u be a subsolution for \mathcal{F} , and let*

$$H_\infty(x) = \lim_{t \rightarrow \infty} H(x, t) : \quad \gamma_\infty(x) = \lim_{t \rightarrow \infty} \gamma(x, t) .$$

Then P is a subsolution for the functional

$$G(P) = \int_Q |D\varphi_P| + \int_Q H_\infty \varphi_P dx + \int_{\partial Q} \gamma_\infty \varphi_P dH_{n-1} .$$

Proof. For $j \in N$, let

$$U_j = \{(x, t) \in Q : t < u(x) - j\} .$$

The set U_j is obviously a subsolution for

$$F^j(A) = \int_Q |D\varphi_A| + \int_Q H_j \varphi_A dxdt + \int_{\partial Q} \gamma_j \varphi_A dH_n$$

with

$$H_j(x, t) = H(x, t + j) ; \quad \gamma_j(x, t) = \gamma(x, t + j) .$$

We have $U_j \supset U_{j+1} \supset \dots \supset W = \bigcap_{j \in N} U_j$, and hence by Proposition 1.2 and the remark above the set W is a subsolution for the functional

$$\int_Q |D\varphi_W| + \int_Q H_\infty \varphi_W dxdt + \int_{\partial Q} \gamma_\infty \varphi_W dH_n .$$

Since W is a vertical cylinder, $W = P \times \mathbf{R}$, and since H_∞ and γ_∞ are independent of t , it follows easily that P is a subsolution for G .

Before proceeding further in the discussion of the set P , we recall that if E is a set and $x_0 \in \partial E$, we say that x_0 belongs to the *reduced boundary* of E if for every $R > 0$ we have:

$$(2.1) \quad 0 < |E \cap B_R(x_0)| < |B_R(x_0)|$$

where $B_R(x_0)$ is the ball of radius R centred at x_0 . It is well known that after changing E in a set of measure zero we may suppose that ∂E coincides with the essential boundary of E .

THEOREM 1.1. *Let U be a local subsolution for the functional F , and suppose that the boundary of U , ∂U , coincides with the essential boundary. Then the same is true for P .*

Proof. It is clearly sufficient to show that the boundary of $W = P \times \mathbf{R}$ is essential. Moreover we have only to show the first

of inequalities (21), since the second follows at one from the remark that $W = \bigcap_{j \in N} U_j$.

Actually, we shall prove a stronger result, namely that if $z_0 = (x_0, t_0)$ is a point in Q and if for all $r > 0$ we have $|U_r(z_0)| > 0$, then for r small enough $|U_r(z_0)| \geq c_s r^{n+1}$, where we have set

$$U_r(z_0) = U \cap C_r(z_0); \quad C_r(z_0) = \{z = (x, t) : |z - z_0| < r, |t - t_0| < r\}.$$

Let $z_0 \in Q$ and let $0 < r < R < \text{dist}(z_0, \partial Q)$. Since U is a subsolution we have, for large T , $F_T(U) \leq F_T(U - C_r)$, and therefore:

$$(2.2) \quad \int_{C_r} |D\varphi_U| + \int_{C_r} H\varphi_U dxdt \leq \int_{\partial C_r} \varphi_U dH_n.$$

On the other hand we have for almost every $r < R$:

$$(2.3) \quad \int |D\varphi_{U_r}| = \int_{C_r} |D\varphi_U| + \int_{\partial C_r} \varphi_U dH_n$$

and whence

$$(2.4) \quad \int |D\varphi_{U_r}| + \int H\varphi_{U_r} dxdt \leq 2 \int_{\partial C_r} \varphi_U dH_n.$$

Suppose now that we have $t > 0$ in C_r (or in other words that $t_0 > R$). Then

$$\int H\varphi_{U_r} dxdt \geq \int H_0^- \varphi_{U_r} dxdt$$

where $H_0^-(x) = \min(H(x, 0), 0)$. From Lemma 2.1 below we get

$$(2.5) \quad \int H\varphi_{U_r} dxdt \geq -k(n) \|H_0^-\|_{n, B_R} \int |D\varphi_{U_r}|$$

and hence

$$(2.6) \quad (1 - k(n) \|H_0^-\|_{n, B_R}) \int |D\varphi_{U_r}| \leq 2 \int_{\partial C_r} \varphi_U dH_n,$$

where we have denoted by $k(n)$ the isoperimetric constant in \mathbb{R}^n . The right-hand side of (2.6) represents the derivative of $|U_r|$, so that we have:

$$\begin{aligned} \frac{d}{dr} |U_r| &= \int_{\partial C_r} \varphi_U dH_n \geq \frac{1}{2} (1 - k(n) \|H_0^-\|_{n, B_R}) \int |D\varphi_{U_r}| \\ &\geq \frac{1}{2k(n+1)} (1 - k(n) \|H_0^-\|_{n, B_R}) |U_r|^{n/n+1}. \end{aligned}$$

Let now R be so small that $\|H_0^-\|_{n, B_R} < 1/2k(n)$; then for almost every $r < R$ we get

$$\frac{d}{dr} |U_r| \geq \frac{1}{4k(n+1)} |U_r|^{\frac{n}{n+1}}$$

and in conclusion, if $|U_r| > 0$ for every r , we obtain

$$(2.7) \quad |U_r| \geq r^{\frac{n}{n+1}} / 4(n+1)k(n+1).$$

The estimate (2.7) holds obviously for each of the sets U_j defined in Proposition 2.1, with R independent of j . We may then pass to the limit as $j \rightarrow \infty$, getting the same estimate for W .

Finally, if $x_0 \in \Omega$ is such that $|P_r(x_0)| = |P \cap B_r(x_0)| > 0$ for every $r > 0$, we conclude that

$$(2.8) \quad |P_r(x_0)| \geq r^{\frac{n}{n+1}} / 8(n+1)k(n+1).$$

We note that the above estimate (2.8) does not hold for super-solutions, and the Theorem 2.1 is in general false. For example the function $|x|^{-1}$ is a supersolution for small $|x|(H=0)$, and $P=\{0\}$.

To conclude the proof of the theorem it remains to prove the estimate (2.5). This is done in the next lemma.

LEMMA 2.1. *Let $h(x) \in L^n(\Omega)$ and let $E \subset C_R = B_R \times I_R$. Then*

$$\int_E |h| dxdt \leq k(n) \|h\|_{n,B_R} \int |D\varphi_E|.$$

Proof. We have from Hölder's inequality

$$\int_E |h| dxdt \leq \|h\|_{n,B_R} \int |E_t|^{1-1/n} dt$$

where

$$E_t = \{x \in \Omega : (x, t) \in E\}.$$

On the other hand from the isoperimetric inequality we get

$$|E_t|^{1-1/n} \leq k(n) \int |D\varphi_{E_t}|$$

and therefore

$$\int_E |h| dxdt \leq k(n) \|h\|_{n,B_R} \int dt \int |D\varphi_{E_t}| \leq k(n) \|h\|_{n,B_R} \int |D\varphi_E|.$$

3. Variational solutions. In general the sets P and N are not empty, and sometimes they may cover the whole of Ω . The purpose of this section is to investigate under what conditions we may conclude the absence of these singular sets.

We begin with two simple remarks concerning subsolutions; similar results hold for supersolutions.

(A) If A is an open set in Ω , and if the measure of $P \cap A$ is zero, then $P \cap A$ is empty. Actually we may say more, namely if $A \subset \Omega$ and if $|P \cap A|$ is small enough (depending on A and H) then $P \cap A = \emptyset$.

(B) If $P \cap A = \emptyset$, then u is locally bounded from above in A . Assertion (B) follows from estimate (2.7), whereas (A) is a consequence of (2.8).

EXAMPLE 3.1. (Emmer [3]) Let $u(x)$ be a generalized solution for \mathcal{F} with obstacles, i.e., satisfying the conditions

$$\psi_1(x) \leq u(x) \leq \psi_2(x).$$

Suppose that $\psi_1(x) \leq M_1$, and $\psi_2(x) \geq M_2$. It is clear that $u(x)$ is a generalized subsolution in $\Omega \times (M_1, +\infty)$, and a generalized super-solution in $\Omega \times (-\infty, M_2)$.

It follows from remarks (A) and (B) above that if the obstacle $\psi_1[\psi_2]$ is finite almost everywhere in Ω , then u will be locally bounded from below [from above].

When no obstacle is present we need some assumptions on H and γ . Let us begin with some necessary conditions.

Suppose $u(x)$ is a smooth solution for \mathcal{F} in Ω . If $A \subset \Omega$, and we compare $\mathcal{F}(u)$ with $\mathcal{F}(u + t\varphi_A)$, $t > 0$, we get easily

$$\int_{\Omega} Tu \cdot D\varphi_A + \int_{\Omega} H\varphi_A dx + \int_{\partial\Omega} \gamma^+ \varphi_A dH_{n-1} \geq 0$$

where $\gamma^+(x, t) = \lim_{s \rightarrow t^+} \gamma(x, s)$.

Taking into account that $|Tu| < 1$ in Ω , and arguing as in [7], § 1, we get the inequality

$$(3.1) \quad \int_{\Omega} H\varphi_A dx + \int_{\partial\Omega} \gamma^+ \varphi_A dH_{n-1} > - \int_{\Omega} |D\varphi_A|$$

and therefore

$$(3.2) \quad \int_{\Omega} H_\infty \varphi_A dx + \int_{\partial\Omega} \gamma_\infty \varphi_A dH_{n-1} > - \int_{\Omega} |D\varphi_A|$$

for every set $A \neq \emptyset$, Ω . In a similar way:

$$(3.3) \quad \int_{\Omega} H_{-\infty} \varphi_A dx + \int_{\partial\Omega} \gamma_{-\infty} \varphi_A dH_{n-1} < \int_{\Omega} |D\varphi_A|$$

for $A \neq \emptyset$, Ω .

THEOREM 3.1. *Let the strict inequality (3.2) hold for every*

nonempty set A , and let u be a generalized subsolution for \mathcal{F} . Then $P = \emptyset$.

Proof. By Proposition 2.1, P is a subsolution for the functional

$$G(P) = \int_{\Omega} |D\varphi_P| + \int_{\Omega} H_{\infty} \varphi_P dx + \int_{\partial\Omega} \gamma_{\infty} \varphi_P dH_{n-1}.$$

On the other hand we have from (3.1), $G(A) \geq 0$, the equality holding only for $A = \emptyset$. This implies immediately that P is empty.

The same argument shows that if (3.3) holds, with strict inequality for Ω , and if u is a supersolution, then $N = \emptyset$.

EXAMPLE 3.2. Discontinuous obstacles (De Acutis [2]).

Suppose that the obstacles ψ_1 and ψ_2 of Example 3.1 are finite only in some regular subsets D_1 and D_2 of Ω . Arguing as before we may conclude that the generalized solution is locally bounded in the interior of these sets, from above in D_2 and from below in D_1 . If in addition condition (3.2) is satisfied in $\Omega - D_2$, and (3.3) in $\Omega - D_1$, we may conclude that $P = N = \emptyset$ and therefore that u is locally bounded in Ω .

In general, even when $P = N = \emptyset$, the solution u can go to $\pm\infty$ when x approaches $\partial\Omega$ (see e.g., [8]).

However this possibility can be excluded if we make some additional assumptions on the boundary function γ . In the following we shall suppose that there exist constants $\theta_0 > 0$ and α , $0 \leq \alpha < 1$, such that

$$(3.4) \quad \gamma(x, t) \geq -\alpha \quad \forall x \in \partial\Omega, \quad \forall t > \theta_0$$

$$(3.5) \quad \gamma(x, t) \leq \alpha \quad \forall x \in \partial\Omega, \quad \forall t < -\theta_0.$$

We note that (3.4), (3.5) correspond to a bounded boundary datum in the case of Dirichlet's boundary conditions ($\gamma(x, t) = 1 - 2\varphi_F(x, t)$), whereas in the case of capillarity boundary conditions they are equivalent to $|\cos\theta| \leq \alpha < 1$.

With the help of (3.4) and (3.5) we can prove the following generalization of Theorem 2.1.

THEOREM 3.2. *Let U be a local subsolution for the functional F , and let $z_0 = (x_0, t_0)$, $t_0 > \theta_0 + 1$, be a point of $\bar{\Omega}$ such that for every positive r :*

$$(3.6) \quad |U_r| = |U \cap C_r(z_0)| > 0.$$

Suppose further that (3.4) is satisfied. Then there exist constants

$R_0 > 0$ and c_4 such that for every $r \leq R_0$ we have:

$$(3.7) \quad |U_r| \geq c_4 r^{n+1}.$$

Proof. As in Theorem 2.1 we compare U with $U - C_r$, getting

$$\int_{Q \cap \sigma_r} |D\varphi_U| + \int_{Q \cap \sigma_r} H\varphi_U dxdt + \int_{\partial Q \cap \sigma_r} \gamma\varphi_U dH_n \leq \int_{\partial \sigma_r} \varphi_U dH_n$$

and therefore for almost every r :

$$\int_Q |D\varphi_{U_r}| + \int_Q H\varphi_{U_r} dxdt + \int_{\partial Q} \gamma\varphi_{U_r} dH_n \leq 2 \int_{\partial \sigma_r} \varphi_{U_r} dH_n.$$

We have from (3.4):

$$\int_{\partial Q} \gamma\varphi_{U_r} dH_n \geq -\alpha \int_{\partial Q} \varphi_{U_r} dH_n$$

for every $r < 1$, and from Lemma 1.1:

$$\begin{aligned} \int_{\partial Q} \varphi_{U_r} dH_n &\leq \int_Q |D\varphi_{U_r}| + c_1 |U_r| \leq \int_Q |D\varphi_{U_r}| \\ &\quad + c_1 k(n+1) |U_r|^{1/n+1} \int_Q |D\varphi_{U_r}|. \end{aligned}$$

If $R < 1$ is so small that $c_1 k(n+1) |C_R|^{1/n+1} < 1/2$, we get for every $r < R$:

$$(3.8) \quad \int_{\partial Q} \varphi_{U_r} dH \leq \frac{1 + c_1 k(n+1) |C_R|^{1/n+1}}{1 - c_1 k(n+1) |C_R|^{1/n+1}} \int_Q |D\varphi_{U_r}|$$

and

$$(3.9) \quad \int_Q |D\varphi_{U_r}| \leq \frac{2}{1 - c_1 k(n+1) |C_R|^{1/n+1}} \int_Q |D\varphi_{U_r}|.$$

Finally, we may estimate the curvature term as in (2.5):

$$\int_Q H\varphi_{U_r} dxdt \geq -k(n) \|H_0^-\|_{n, B_R} \int_Q |D\varphi_{U_r}|,$$

and in conclusion

$$\begin{aligned} \int_Q H\varphi_{U_r} dxdt + \int_{\partial Q} \gamma\varphi_{U_r} dH_n &\geq -\left\{ \alpha \frac{1 + c_1 k(n+1) |C_R|^{1/n+1}}{1 - c_1 k(n+1) |C_R|^{1/n+1}} \right. \\ &\quad \left. + k(n) \|H_0^-\|_{n, B_R} \frac{2}{1 - c_1 k(n+1) |C_R|^{1/n+1}} \right\} \int_Q |D\varphi_{U_r}|. \end{aligned}$$

Since $\alpha < 1$, we may choose R_0 small enough that the right-hand

side is bounded from below by $\frac{1}{2} - (1 + \alpha) \int_Q |D\varphi_{U_r}|$. We remark that the constant R_0 depends only on H through $\|H_0^-\|_{n, B_R}$ and therefore it is uniform for H_0^- in compact sets of $L^n(\Omega)$.

In conclusion we get

$$\frac{d}{dr} |U_r| = \int_{\partial C_r} \varphi_U dH_n \geq \frac{1}{2} \varepsilon_0 \int_Q |D\varphi_{U_r}| \geq \frac{1}{8} \varepsilon_0 \int_Q |D\varphi_{U_r}| \quad (\varepsilon_0 = (1 - \alpha)/2)$$

and arguing as in Theorem 2.1 we conclude for $r < R_0$:

$$|U_r| \geq c_4 r^{n+1}, \quad c_4 = \varepsilon_0 / 8(n+1).$$

REMARK 3.1. It follows from (3.7) that for $r < R_0$:

$$(3.10) \quad |P_r| = |P \cap B_r(x_0)| \geq \varepsilon_0 r^n / 16(n+1)$$

uniformly for $x_0 \in \bar{\Omega}$.

In particular, there exists a constant p_0 , depending only on α and on H_0^- (p_0 is uniform for H_0^- in compact sets of $L^n(\Omega)$) such that $|P| < p_0$ implies that P is empty and u is bounded from above in the whole of Ω . This makes possible to improve the results of Examples 3.1 and 3.2.

EXAMPLE 3.3. In Theorem 3.1 and in the above examples we have always made the assumption that the strict inequality (3.2) holds for Ω itself. It is easily seen that if the equality holds for Ω , we cannot expect in general to have a bounded solution. For example, let $\Omega = \{x \in \mathbf{R}^2 : |x| < 1\}$, and let

$$H(x, u) = \frac{2}{\pi} \operatorname{arctg} u - 3$$

$$\kappa(x, u) = |u|.$$

(Dirichlet problem with zero boundary data.) We have $H_\infty = -2$ and (3.2) is satisfied but the equality holds for Ω . In this case we have $P = \Omega$, and $u = +\infty$.

REMARK 3.2. It is clear that a bounded solution is a variational solution, namely it has finite area and minimizes the functional \mathcal{F} in $BV(\Omega)$. Moreover, if H is Lipschitz-continuous the function $u(x)$ is of class $C^{2+\alpha}$ in Ω and is a classical solution of the equation

$$\operatorname{div} Tu = H(x, u).$$

REMARK 3.3. It follows from Theorem 3.2 that every generalized solution which is almost everywhere finite is bounded. In particular this is true for every L^1 generalized solution. It is easily

seen that it is possible to give an estimate of $\sup_{\Omega} u$ in terms of the L^1 norm of u , or better in terms of $\int_{\Omega} u^+ dx$, $u^+ = \max(u, \theta_0)$.

For, let θ_0 and R_0 be as in Theorem 3.2, and let $x_0 \in \Omega$ be such that

$$\sup_{\Omega} u < u(x_0) + 1.$$

For $j \in N$ let $z_j = (x_0, \theta_0 + 2jR_0)$; we have $z_j \in U$ for $j \leq k = [u(x_0) - \theta_0]/2R_0$. From Theorem 3.2 we get

$$|U_{R_0}(z_j)| \geq c_4 R_0^{n+1}$$

and therefore

$$\int_{\Omega} u^+ dx \geq \sum_{j=1}^k |U_{R_0}(z_j)| \geq k c_4 R_0^{n+1}.$$

In conclusion

$$(3.11) \quad \sup_{\Omega} u \leq 1 + u(x_0) \leq \frac{2}{c_4 R_0^n} \int_{\Omega} u^+ dx + \theta_0 + 2R_0 + 1.$$

We remark that estimate (3.11) holds for minima with obstacle $u \geq \psi$, provided $\psi \leq M \leq \theta_0$. Moreover, the bound for $\sup u$ depends only on $\int_{\Omega} u^+ dx$ and on R_0 and θ_0 ; it is therefore uniform for H_0^- in compact subset of $L^n(\Omega)$.

4. An application: Lagrange multipliers. We apply now the results of the previous sections to the discussion of the existence of Lagrange multipliers for minima with obstacle and constant volume.

Let $\psi(x)$ be a function bounded from above, and let V be a real number, with

$$V > \int_{\Omega} \psi dx \geq -\infty.$$

We have seen in § 1 that the functional

$$(4.1) \quad \mathcal{F}(u) = \int_{\Omega} \sqrt{1 + |Du|^2} + \int_{\Omega} \lambda(x, u) dx + \int_{\partial\Omega} \kappa(x, u) dH_{n-1}$$

with the constraints

$$\begin{aligned} u &\geq \psi \\ \int_{\Omega} u dx &= V \end{aligned}$$

has a generalized solution; i.e., that there exists a function $u: \Omega \rightarrow [-\infty, +\infty]$ such that its subgraph U minimizes locally in Q

the functional

$$(4.2) \quad F(U) = \int_Q |D\varphi_U| + \int_Q H\varphi_U dxdt + \int_{\partial Q} \gamma\varphi_U dH_r$$

among all subsets $E \subset Q$ such that $E \supset \Psi$ and $|E^+| - |E^-| = V$

$$(E^\pm = \{z = (x, t) \in Q : z \in E, t \geq 0\}) .$$

Our goal is to show that there exists a variational solution to the above problem, and moreover that such solution may be obtained as a minimum for the functional

$$(4.3) \quad \mathcal{F}_q(u) = \mathcal{F}(u) - q \int_Q u dx$$

for a suitable value of the constant q (the Lagrange multiplier).

Our hypotheses will be those of Theorem 3.2, namely:

- (i) $\lambda(x, u)$ and $\kappa(x, u)$ are convex functions in u , or, what is the same, $H(x, u)$ and $\gamma(x, u)$ are nondecreasing functions of u .
- (ii) $H(x, t)$ is in $L^n(\Omega)$ for every $t \in \mathbf{R}$.
- (iii) There exist constants θ_0 and α , $0 \leq \alpha < 1$, such that

$$(4.4) \quad \gamma(x, t) \geq -\alpha \quad \forall x \in \Omega , \quad \forall t > \theta_0$$

$$(4.5) \quad \gamma(x, t) \leq \alpha \quad \forall x \in \Omega , \quad \forall t < -\theta_0 .$$

Of course we may assume that $\theta_0 \geq M = \sup \psi$.

For what concerns the obstacle ψ we shall assume that it is an upper bounded measurable function, almost everywhere finite. In this way every solution u_q for the functional \mathcal{F}_q is bounded from below (see Remark 3.1).

To prove our result we have to show that there exist a value q_0 and a variational solution u_0 to \mathcal{F}_{q_0} such that $\int_Q u_0 dx = V$.

LEMMA 4.1. *Let $h(x)$ be in $L^n(\Omega)$ and let $\gamma(x) \geq -\alpha$. Then for every set $A \subset \Omega$, satisfying*

$$c_1 k(n) |A|^{1/n} < 1$$

we have

$$(4.6) \quad \begin{aligned} \int_A h dx + \int_{\partial A} \gamma \varphi_A dH_{n-1} &\geq - \left\{ \alpha \frac{1 + c_1 k(n) |A|^{1/n}}{1 - c_1 k(n) |A|^{1/n}} \right. \\ &\quad \left. + \frac{2k(n) \|h\|_{n,A}}{1 - c_1 k(n) |A|^{1/n}} \right\} \int_Q |D\varphi_A| . \end{aligned}$$

Proof. We proceed as in Theorem 3.2. We have

$$\int_{\partial\Omega} \gamma \varphi_A dH_{n-1} \geq -\alpha \int_{\partial\Omega} \varphi_A dH_{n-1}.$$

From Lemma 1.1

$$\int_{\partial\Omega} \varphi_A dH_{n-1} \leq \int_{\Omega} |D\varphi_A| + c_1 |A| \leq \int_{\Omega} |D\varphi_A| + c_1 k(n) |A|^{1/n} \int_{\Omega} |D\varphi_A|$$

and therefore, if $c_1 k(n) |A|^{1/n} < 1$,

$$\begin{aligned} \int_{\partial\Omega} \varphi_A dH_{n-1} &\leq \frac{1 + c_1 k(n) |A|^{1/n}}{1 - c_1 k(n) |A|^{1/n}} \int_{\Omega} |D\varphi_A| \\ \int_{\Omega} |D\varphi_A| &\leq \frac{2}{1 - c_1 k(n) |A|^{1/n}} \int_{\Omega} |D\varphi_A|. \end{aligned}$$

Moreover

$$\int_A h dx \geq -\|h\|_{n,A} \frac{2k(n)}{1 - c_1 k(n) |A|^{1/n}} \int_{\Omega} |D\varphi_A|$$

from which (4.6) follows at once.

A consequence of Lemma 4.1 is that, at least for q negative and big enough, the functional \mathcal{F}_q has a variational solution. For that it is sufficient to show that condition (3.2) holds (with strict inequality even for Ω) when H_∞ is replaced by $H_\infty - q$. We distinguish two cases, depending on the size of A . If A is small, we have using Lemma 4.1:

$$\begin{aligned} \int_A (H_\infty - q) dx + \int_{\partial\Omega} \gamma_\infty \varphi_A dH_{n-1} &\geq \int_A H_0^- dx + \int_{\partial\Omega} \gamma_\infty \varphi_A dH_{n-1} \\ &< - \int_{\Omega} |D\varphi_A| \end{aligned}$$

provided $|A|$ is smaller than some constant σ_0 depending only on α and H_0^- .

On the other hand, if $|A| > \sigma_0$ we have:

$$\begin{aligned} \int_A (H_\infty - q) dx + \int_{\partial\Omega} \gamma_\infty \varphi_A dH_{n-1} &\geq -\alpha \int_{\Omega} |D\varphi_A| - c_1 |A| \\ &+ \int_{\Omega} H_0^- dx - q |A| > - \int_{\Omega} |D\varphi_A| \end{aligned}$$

whenever $-(q + \alpha c_1) \sigma_0 + \int_{\Omega} H_0^- dx \geq 0$.

Let us denote by S the set of all the values of q for which the necessary condition

$$(4.7) \quad \int_A (H_\infty - q) dx + \int_{\partial\Omega} \gamma_\infty \varphi_A dH_{n-1} \geq - \int_{\Omega} |D\varphi_A|$$

is satisfied for all $A \subset \Omega$, and let $q_0 = \sup S$.

It is clear from the above that $S \neq \emptyset$ and therefore $q_0 > -\infty$. It is easily seen that (4.7) holds for q_0 itself, and that for $q < q_0$ we have the strict inequality for every nonempty set, including Ω itself. It follows that \mathcal{F}_q has bounded (variational) solutions for all $q < q_0$ (and possibly for q_0), but not for any $q > q_0$, so that $S = (-\infty, q_0]$.

For $q < q_0$ we define $V(q)$ as the set described by the integral

$$\int_{\Omega} u dx$$

when u varies among all solutions to \mathcal{F}_q . Since \mathcal{F}_q is convex (but not strictly convex), the set $V(q)$ is a closed interval, which may of course reduce to a point.

LEMMA 4.2. *Let $p < q$ and let u, v be solutions of \mathcal{F}_q , \mathcal{F}_q respectively. Then $u \leqq v$.*

Proof. It is well known (see [1] and [6], Theorem 1.17) that for every $w \in BV(\Omega)$ there exists a sequence $w_j \in C^\infty \cap BV(\Omega)$ such that $w_j \rightarrow w$ in $L^1(\Omega)$ and $\sqrt{1 + |Dw_j|^2} \rightarrow \sqrt{1 + |Dw|^2}$. Moreover $w_j = w$ on $\partial\Omega$, and if $|w| \leqq M$ we may choose the w_j satisfying $|w_j| \leqq M$.

It is now a simple matter of computation to show that for any r , $\mathcal{F}_r(w_j) \rightarrow \mathcal{F}_r(w)$. In conclusion, for every $\varepsilon > 0$ there exist C^∞ -functions u_ε and v_ε such that

$$\begin{aligned} \|u_\varepsilon - u\|_1 + \|v_\varepsilon - v\|_1 &< \varepsilon \\ \mathcal{F}_p(u_\varepsilon) &< \mathcal{F}_p(u) + \varepsilon \\ \mathcal{F}_q(v_\varepsilon) &< \mathcal{F}_q(v) + \varepsilon. \end{aligned}$$

Let $A = \{x \in \Omega : u_\varepsilon(x) > v_\varepsilon(x)\}$; we have

$$\begin{aligned} \mathcal{F}_p(\min(u_\varepsilon, v_\varepsilon)) &= \mathcal{F}_p(u_\varepsilon) + \int_A \sqrt{1 + |Dv_\varepsilon|^2} - \int_A \sqrt{1 + |Du_\varepsilon|^2} \\ &\quad + \int_A [\lambda(x, v_\varepsilon) - \lambda(x, u_\varepsilon)] dx + \int_{\partial\Omega \cap A} [\kappa(x, v_\varepsilon) - \kappa(x, u_\varepsilon)] dH_{n-1} \\ &\quad - p \int_A (v_\varepsilon - u_\varepsilon) dx \geq \mathcal{F}_p(u) \geq \mathcal{F}_p(u_\varepsilon) - \varepsilon. \end{aligned}$$

Similarly:

$$\begin{aligned} \mathcal{F}_q(\min(u_\varepsilon, v_\varepsilon)) &= \mathcal{F}_q(v_\varepsilon) - \int_A \sqrt{1 + |Dv_\varepsilon|^2} + \int_A \sqrt{1 + |Du_\varepsilon|^2} \\ &\quad - \int_A [\lambda(x, v_\varepsilon) - \lambda(x, u_\varepsilon)] dx - \int_{\partial\Omega \cap A} [\kappa(x, v_\varepsilon) - \kappa(x, u_\varepsilon)] dH_{n-1} \end{aligned}$$

$$+q \int_A (v_\varepsilon - u_\varepsilon) dx \geq \mathcal{F}_q(v) \geq \mathcal{F}_q(v_\varepsilon) - \varepsilon.$$

In conclusion, we have

$$(q-p) \int_A (u_\varepsilon - v_\varepsilon) dx = (q-p) \int_\Omega (\max(u_\varepsilon, v_\varepsilon) - v_\varepsilon) dx < 2\varepsilon$$

and passing to the limit as $\varepsilon \rightarrow 0$:

$$(q-p) \int_\Omega (\max(u, v) - v) dx \leq 0$$

which implies $\max(u, v) = v$ and therefore $u \leq v$.

A trivial consequence of the above lemma is that if $q_1 < q_2$ and u_1, u_2 are solutions of $\mathcal{F}_{q_1}, \mathcal{F}_{q_2}$ respectively, we have

$$\int_\Omega u_1 dx \leq \int_\Omega u_2 dx.$$

In other words, the mapping $q \rightarrow V(q)$ is nondecreasing.

LEMMA 4.3. *For every V , $\inf_{q < q_0} V(q) < V < \sup_{q < q_0} V(q)$, there exists a $q' < q_0$ such that $V \in V(q')$.*

Proof. Let S_V denote the set of all q such that $V(q) < V$; and let $q' = \sup S_V$. Let $q_j \nearrow q'$ and for each j let u_j be a solution for \mathcal{F}_{q_j} . It is easily seen that the functions u_j are uniformly bounded from below, and therefore from above because the integrals $\int_\Omega u_j dx$ are uniformly bounded. Since $\mathcal{F}_{q_j}(u_j) \leq \mathcal{F}_{q_j}(0) = |\Omega|$, we conclude that u_j is a bounded sequence in BV and hence, passing possibly to a subsequence, it converges (strongly in L^1 and weakly in $L^{n/n-1}$) to some function u' . From Theorem 1.2 we deduce that u' is a solution for $\mathcal{F}_{q'}$, and $\int_\Omega u' dx \leq V$.

In a similar way, starting with a sequence p_j decreasing to q' , we arrive to a solution u'' for $\mathcal{F}_{q'}$, with $\int_\Omega u'' dx \geq V$. From the convexity of $\mathcal{F}_{q'}$ it follows that $V(q')$ is an interval, and hence $V \in V(q')$.

To conclude our proof we have only to show that every value $V > \int_\Omega \psi dx$ is reached.

LEMMA 4.4. *Let $\eta(x)$ be a smooth function in Ω , such that $\eta \geq \psi$. Let $q_j \rightarrow -\infty$ and let u_j be a solution of \mathcal{F}_{q_j} . Then*

$$\lim_{j \rightarrow \infty} \max(u_j, \eta) = \eta .$$

Proof. Let $q_1 < q_0$ and let $q < q_1$. Let u be a solution of \mathcal{F}_q and let u_ε be as in Lemma 4.2. We may compare u_ε with $v_\varepsilon = \min(u_\varepsilon, \eta)$, getting

$$\begin{aligned} q \int_{\Omega} (v_\varepsilon - u_\varepsilon) dx &\leq \int_A \sqrt{1 + |D\eta|^2} - \int_A \sqrt{1 + |Du_\varepsilon|^2} . \\ &+ \int_{\Omega} [\lambda(x, v_\varepsilon) - \lambda(x, u_\varepsilon)] dx + \int_{\partial\Omega} [\kappa(x, v_\varepsilon) - \kappa(x, u_\varepsilon)] dH_{n-1} \end{aligned}$$

We have

$$\int_{\Omega} [\lambda(x, v_\varepsilon) - \lambda(x, u_\varepsilon)] dx = - \int_A dx \int_{\eta}^{u_\varepsilon} H(x, t) dt \leq - \int_{\Omega} H(x, \eta) (u_\varepsilon - v_\varepsilon) dx$$

and

$$\begin{aligned} \int_{\partial\Omega} [\kappa(x, v_\varepsilon) - \kappa(x, u_\varepsilon)] dH_{n-1} &\leq \alpha \int_{\partial\Omega} |v_\varepsilon - u_\varepsilon| dH_{n-1} \leq \int_{\Omega} |D(v_\varepsilon - u_\varepsilon)| \\ &+ c_1 \int_{\Omega} |v_\varepsilon - u_\varepsilon| dx \leq \int_A |D\eta| + \int_A |Du_\varepsilon| + c_1 \int_{\Omega} |u_\varepsilon - v_\varepsilon| dx . \end{aligned}$$

In conclusion:

$$\begin{aligned} -(c_1 + q) \int_{\Omega} |u_\varepsilon - v_\varepsilon| dx &= -(c_1 + q) \int_{\Omega} (\max(u_\varepsilon, \eta) - \eta) dx \\ &\leq 2 \int_A \sqrt{1 + |D\eta|^2} - \int_{\Omega} H(x, \eta) (u_\varepsilon - v_\varepsilon) dx \end{aligned}$$

and letting $\varepsilon \rightarrow 0$:

$$\begin{aligned} (4.8) \quad -(c_1 + q) \int_{\Omega} (\max(u, \eta) - \eta) dx &\leq 2 \int_{\Omega} \sqrt{1 + |D\eta|^2} \\ &+ \int_{\Omega} H(x, \eta) (u - v) dx . \end{aligned}$$

On the other hand, it follows from Lemma 4.2 that $u = u_q \leq u_{q_1}$, and therefore the right-hand side of (4.8) is bounded independently of q . Letting $q \rightarrow -\infty$ we get the conclusion of the theorem.

The above result shows that $\inf V(q) = \int_{\Omega} \psi dx$. The existence of the Lagrange multiplier will follow if we show that $\sup V(q) = +\infty$.

This is easy if $q_0 = +\infty$, i.e., if the functional \mathcal{F}_q has variational solutions for every real q (this happens for instance if $H_\infty = +\infty$). In this case we have only to repeat the above argument to show that the volume $V(q)$ tends to infinity as $q \rightarrow +\infty$. More complicate is the case when $q_0 < +\infty$. Let q_j be an increasing sequence, $q_j \rightarrow q_0$, and let u_j be the corresponding sequence of solutions. We distin-

guish two cases.

$$(I) \quad \|u_j\|_1 \longrightarrow +\infty .$$

In this case, since u_j is increasing, we have $V_j = \int_{\Omega} u_j dx \rightarrow +\infty$, and therefore $\sup V(q) = +\infty$.

$$(II) \quad \|u_j\|_1 \leq M .$$

By Remark 3.3 the functions u_j are uniformly bounded:

$$|u_j(x)| \leq M_1$$

and therefore it is possible to extract from u_j a subsequence which converges to a function $u(x)$, a variational solution for \mathcal{F}_{q_0} . It is then satisfied the necessary condition:

$$(4.9) \quad -q_0 |A| + \int_A H(x, u(x)) dx + \int_{\partial\Omega} \gamma^+(x, u(x)) \varphi_A dH_{n-1} \geq - \int_{\Omega} |D\varphi_A|$$

for every set $A \subset \Omega$.

On the other hand from the true definition of q_0 we have:

$$(4.10) \quad -q_0 |A| + \int_A H_\infty dx + \int_{\partial\Omega} \gamma_\infty \varphi_A dH_{n-1} \geq - \int_{\Omega} |D\varphi_A| .$$

LEMMA 4.5. *Let the strict inequality hold in (4.10) for every nonempty set A . Then there exists $q > q_0$ such that*

$$-q |A| + \int_A H_\infty dx + \int_{\partial\Omega} \gamma_\infty \varphi_A dH_{n-1} \geq - \int_{\Omega} |D\varphi_A| ,$$

for every $A \subset \Omega$.

Proof. Let $q_j = q_0 + 1/j$, and suppose that for every j there exists a set A_j such that

$$(4.11) \quad -q_j |A_j| + \int_{A_j} H_\infty dx + \int_{\partial\Omega} \gamma_\infty \varphi_{A_j} dH_{n-1} < - \int_{\Omega} |D\varphi_{A_j}| .$$

From (4.11) we get

$$\int_{\Omega} |D\varphi_{A_j}| \leq |q_j| |\Omega| + \int_{\Omega} H_\infty^- dx + H_{n-1}(\partial\Omega)$$

and therefore, passing to a subsequence we may conclude that $A_j \rightarrow A$. From (4.6) with $h = H_\infty^- - q_j$, we conclude that the measure of A_j is bounded away from zero, and whence A has positive measure. Passing to the limit as $j \rightarrow +\infty$ we get:

$$-q_0 |A| + \int_A H_\infty dx + \int_{\partial\Omega} \gamma_\infty \varphi_A dH_{n-1} \leq - \int_{\Omega} |D\varphi_A|$$

contradicting the assumption of the lemma.

It follows from the above lemma and the definition of q_0 that the equality sign must hold in (4.10) for some nonempty set A . From the monotonicity of H and γ and from (4.9) we get then:

$$-q_0|A| + \int_A H(x, u(x))dx + \int_{\Omega} \gamma^+(x, u(x))\varphi_A dH_{n-1} = - \int_{\Omega} |D\varphi_A|$$

and therefore for almost every $x \in A$ and for every $t > u(x)$:

$$\begin{aligned} H(x, t) &= H_\infty(x) \\ \gamma(x, t) &= \gamma_\infty(x). \end{aligned}$$

In particular we may conclude that $\mathcal{F}_{q_0}(u + c\varphi_A) = \mathcal{F}_{q_0}(u)$ for every positive constant c , and hence

$$V(q_0) = \left[\int_{\Omega} u dx, +\infty \right).$$

In any case we have then

$$V(S) = \left(\int_{\Omega} \psi dx, +\infty \right]$$

thus proving the existence of a Lagrange multiplier.

REFERENCES

1. G. Anzellotti and M. Giacinta, *Funzioni BV e tracce*, to appear in Rend. Sem. Mat. Padova.
2. D. De Acutis, *Superfici non parametriche a curvatura media assegnata con ostacoli*, Ann. Univ. Ferrara, **23** (1977), 225-250.
3. M. Emmer, *Superfici di curvatura media assegnata con ostacolo*, Ann. Mat. Pura ed Appl. S. IV, **109** (1976), 371-389.
4. C. Gerhardt, *On the capillarity problem with constant volume*, Ann. Sc. Norm. Sup. Pisa S. IV, **2** (1975), 303-320.
5. E. Giusti, *Boundary value problems for non-parametric surfaces of prescribed mean curvature*, Ann. Sc. Norm. Sup. Pisa S. IV, **3** (1976), 501-548.
6. ———, *Minimal surfaces and function of bounded variation*, Notes on pure mathematics, **10**. Canberra, 1977.
7. ———, *On the equation of surfaces of prescribed mean curvature. Existence and uniqueness without boundary conditions*, Inv. Math., **46** (1978), 111-137.
8. U. Massari, *Problema di Dirichlet con dati infiniti per le superfici a curvatura media assegnate*, to appear.
9. M. Miranda, *Superfici cartesiane generalizzate ed insiemi di perimetro finito sui prodotti cartesiani*, Ann. Sc. Norm. Sup. Pisa S. III, **18** (1964), 515-542.
10. ———, *Superfici minime illimitate*, Ann. Sc. Norm. Sup. Pisa S. IV, **4** (1977), 313-322.
11. ———, *Sulle singolarità eliminabili delle soluzioni dell'equazione delle superfici minime*, Ann. Sc. Norm. Sup. Pisa S. IV, **4** (1977), 129-132.

12. G. H. Williams, *Surfaces of prescribed mean curvature with a volume constraint*, to appear.

Received June 29, 1979. Partially supported by GNAFA-CNR.

UNIVERSITA DI PISA
56100 PISA, ITALY

