

GENERATORS OF W^* -ALGEBRAS

HORST BEHNCKE

(Received Dec. 4, 1969; Revised June 12, 1970)

Recently Wogen [1] has shown that any properly infinite W^* -algebra on a separable Hilbert space is singly generated. Further results on generators of W^* -algebras have been obtained by Saito [2] and Percy [3]. In this note we shall extend these results and present new proofs for them. We shall show for example that any properly infinite W^* -algebra \mathfrak{A} is generated by a single subnormal operator T . We will also see that \mathfrak{A} is generated by two unitary operators U and V with $U^2=1$ and $V^3=1$. These results have some interesting consequences. Throughout H will be a separable Hilbert space and \mathfrak{A} will be a properly infinite W^* -algebra on H . Though some of our results are also valid for certain singly generated W^* -algebras, we formulate them only for the properly infinite ones.

LEMMA 1. \mathfrak{A} is singly generated.

PROOF. Since \mathfrak{A} is properly infinite it can be written as $\mathfrak{A}=\mathfrak{B}\otimes B(K)$, where \mathfrak{B} is a properly infinite W^* -algebra on a separable Hilbert space K and where $B(K)$ denotes the algebra of all bounded operators on the infinite dimensional Hilbert space K . In this notation $H=K\otimes K$. We shall write K as the usual sequence space $l^2(N)$, N the positive integers. Then the elements A of \mathfrak{A} can be considered as matrices with entries from \mathfrak{B} , $A=(a_{i,j})$ and $a_{i,j}\in\mathfrak{B}$. \mathfrak{B} is clearly generated by a countable number of operators b_i , $i=1,2,\dots$. Without loss of generality we shall assume that all b_i are positive invertible contractions. Now let $A=(a_{i,j})$ and $B=(b_{i,j})$ with $a_{i,j}=\delta_{i,j} 1/i\cdot 1$ and $b_{i+1,i}=b_{i,i+1}=b_i$ and $b_{i,j}=0$ otherwise. Then A and B are selfadjoint. Let \mathfrak{R} be the W^* -algebra generated by A and B and let $C^*=C\in\mathfrak{R}$. Since $CA=AC$ we see immediately that C is diagonal, $C=(c_{i,j})$ with $c_{i,j}=\delta_{i,j}c_i$. We shall denote this by $C=\text{diag}(c_1, c_2, \dots)$. Then $BC=CB$ shows $b_i c_{i+1}=c_i b_i$ $i=1,2,\dots$. The adjoint of this equation is $c_{i+1} b_i=b_i c_i$ because $c_i=c_i^*$. Thus $c_i b_i^2=b_i c_{i+1} b_i=b_i^2 c_i$. However b_i is positive, therefore $c_i b_i=b_i c_i$ and $(c_{i+1}-c_i)b_i=0$. Since b_i is invertible we see $c_{i+1}=c_i$. Thus $c_1=c_2=\dots=c$ and $cb_i=b_i c$. \mathfrak{B} is generated by the b_i . Therefore $c\in\mathfrak{B}$ and $\mathfrak{R}=\mathfrak{B}'\otimes C$ or $\mathfrak{R}=\mathfrak{A}$. Thus \mathfrak{A} is generated by the two hermitean operators A and B .

The above construction actually gives us a continuous family $\{A_k, B_k\}_{k \in (0,1)}$ of pairwise unitarily inequivalent generators of \mathfrak{A} . Simply choose for B_k $b_1 = k$. These remarks will also apply to all following results.

THEOREM 1. *\mathfrak{A} is generated by a single subnormal operator T .*

PROOF. Let a and b be positive operators on the separable Hilbert space K with $0 < a^2 < b^2$. Following [4] we define on $(K \oplus K \oplus \dots) \oplus (K \oplus K) \oplus (K \oplus K \oplus \dots) = H \oplus H' \oplus H''$ the operator

$$N = \left| \begin{array}{ccc|cc|ccc} 0 & 0 & 0 & a & 0 & & & & \\ a & 0 & 0 & 0 & c & & & & \\ b & b & \ddots & 0 & 0 & & & & \\ & & \ddots & \vdots & \vdots & & & & \\ & & & & & 0 & 0 & b & \\ & & & & & c & 0 & 0 & \\ & & & & & & & & 0 & b & \\ & & & & & & & & & 0 & b & \\ & & & & & & & & & & \ddots & \\ & & & & & & & & & & & \ddots & \end{array} \right|$$

with $c = (b^2 - a^2)^{1/2}$. A simple computation shows that N is normal. The subspace H is clearly invariant under N and thus $T = N|_H$ is subnormal.

As before write $\mathfrak{A} = \mathfrak{B} \otimes B(K')$, with \mathfrak{B} a properly infinite W^* -algebra. By lemma 1 there exist two positive operators $a, b \in \mathfrak{B}$, which generate \mathfrak{B} . We choose them such that $0 < a < 1$ and $2 < b < 3$ and form the operator N and $T = N|_H$ as above. Again denote by \mathfrak{H} the W^* -algebra generated by T . One sees easily that

$$T^n T^{*n} = \text{diag}(0, \dots, 0, b^{n-1} a^2 b^{n-1}, b^{2n}, b^{2n}, \dots)$$

$$T^{*n} T^n = \text{diag}(ab^{2n-2} a, b^{2n}, b^{2n}, \dots).$$

Thus \mathfrak{H} contains any diagonal operator

$$D_n = \text{diag}(0, \dots, 0, 1, 0, \dots).$$

Let $C = C^* \in \mathfrak{H}$ then $CD_n = D_n C$ for all n shows $C = \text{diag}(c_1, c_2, \dots)$. $CT = TC$ implies $c_2 a = ac_1, c_{i+1} b = bc_i \ i > 1$. With the same trick as in lemma 1 we can now show $c = c_2 = \dots = c$ and $ac = ca, bc = cb$ or $c \in \mathfrak{B}'$. Thus $C = c \otimes \mathbf{1} \in \mathfrak{B}' \otimes \mathbf{C}$ and $\mathfrak{H} = \mathfrak{A}$.

Wogen (private communication) has shown that any properly infinite W^* -algebra is generated by a hyponormal operator. However this construction is much

simpler than his. Since a quasinormal operator is of type I [5] the subnormality condition in theorem 1 cannot be strengthened. Here we give an independent much shorter proof for the result in [5] using C^* -algebra techniques.

DEF. An operator T on a Hilbert space H is called postliminal if the C^* -algebra \mathfrak{C} generated by T is postliminal [6, 7].

Postliminal operators are clearly of type I, i.e. they generate a W^* -algebra of type I. However the converse is not true.

THEOREM 2. *A quasinormal operator T is postliminal.*

PROOF. Since T is quasinormal T and T^* commute with (T^*T) .

Let π be an irreducible representation of \mathfrak{C} . Then $\pi(T^*T)$ commutes with $\pi(T)$ and $\pi(T^*)$ and thus with every element in $\pi(\mathfrak{C})$. By irreducibility $\pi(T^*T) = c1$ with $c > 0$. Then $\pi(T) 1/\sqrt{c}$ is an irreducible isometry. Thus $\pi(T) 1/\sqrt{c}$ is unitary or a simple unilateral shift. In the first case $\dim \pi = 1$, whereas in the second case $\pi(\mathfrak{C})$ contains all compact operators. If one applies now the method of the direct integral decomposition to the identity representation of \mathfrak{C} , one obtains the result of [5].

Now let \mathfrak{A} be a factor of type III and T a hyponormal generator of \mathfrak{A} . By \mathfrak{C} denote again the C^* -algebra generated by T . Then \mathfrak{C} is antiliminal [6]. Let π be a representation of \mathfrak{C} such that $\pi(\mathfrak{C})$ generates a W^* -algebra of finite type. Then $\pi(\mathfrak{C})'$ has a complete family of normal traces. Let φ be a trace, then $0 \leq \varphi(\pi(T^*T - TT^*)) = \varphi(\pi(T)^*\pi(T)) - \varphi(\pi(T)\pi(T)^*) = 0$. Hence $\pi(T)$ is normal. This gives us many examples of C^* -algebras with representations of type III, but none of type II_1 . This also shows that no finite nonabelian W^* -algebra is generated by a hyponormal operator.

The following corollaries are corollaries of lemma 1 or theorem 1.

COROLLARY 1. *\mathfrak{A} is generated by an operator T with $p(T) = 0$, where $p(x)$ is a polynomial of degree three or higher.*

PROOF. Let $p(x) = \prod_{i=1}^n (x - a_i)$ and let $\mathfrak{A} = \mathfrak{B} \otimes M_n$.

Then consider the operator matrix $T = (t_{i,j})$ $i, j = 1, \dots, n$ with $t_{i,i} = a_i$, $t_{1,2} = a$, $t_{n-1,n} = b$, $t_{i,i+1} = 1$ $i = 2, \dots, n-2$ and $t_{i,j} = 0$ otherwise Here a and b are positive invertible generators of \mathfrak{B} . If a and b were commuting the theorem of Hamilton

and Cayley would show $p(T)=0$. It is easy to see however that the only matrix elements of T^k $k \leqq n$, where a and b appear together are $(T^n)_{,n}$ and $(T^{n-1})_{1,n}$. Thus the theorem of Hamilton and Cayley is also applicable in this case and $p(T) = 0$. Simple matrix computation shows then as in theorem 1 that T generates \mathfrak{A} .

We should remark that an operator T satisfying a polynomial identity of degree 2 is binormal and thus generates a W^* -algebra of type $I_{\leqq 2}$. In particular \mathfrak{A} is generated by an operator T with $T^n=1, n=3, 4, \dots$. Using Weyl's trick on the bounded representation $k \rightarrow T^k$ of the cyclic group of order n this shows that T is similar to a unitary operator U with $U^n=1$. Thus \mathfrak{A} is generated by an operator T which is similar to a unitary operator.

COROLLARY 2. \mathfrak{A} is generated by a partial isometry.

PROOF. We write $\mathfrak{A} = \mathfrak{B} \otimes M_2$. By lemma 1 \mathfrak{B} is generated by a positive operator $1/2 \geqq a \geqq 1/4$ and a unitary operator u . Let $b=(1-a^2)^{1/2}u$ then $T = \begin{pmatrix} b & a \\ 0 & 0 \end{pmatrix}$ is a partial isometry with $TT^* = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$. Thus any operator $C = C^*$ commuting with T must be diagonal, because it commutes with $TT^*, C = \text{diag}(c_1, c_2)$. Then $TC = CT$ shows $bc_1 = c_1b$ and $ac_2 = c_1a$. As before this gives $c_1 = c_2 = c$ because a is positive and invertible. Thus $ac = ca$ and $cu = uc$ or $c \in \mathfrak{B}'$.

COROLLARY 3. \mathfrak{A} is generated by an (infinite) projection P and a positive operator S .

PROOF. Let $\mathfrak{A} = \mathfrak{B} \otimes M_2$ and let $P = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ and $S = \begin{pmatrix} a & b \\ b & 1 \end{pmatrix}$ where a and b are generators of \mathfrak{B} with $2 \geqq a \geqq 1$ and $1/2 \geqq b > 0$. The remainder is shown as in the previous corollary.

THEOREM 3. \mathfrak{A} is generated by three projections P_1, P_2 and P_3 , two of which may be chosen to be orthogonal, $P_1 \cdot P_2 = 0$.

PROOF. Again we write $\mathfrak{A} = \mathfrak{B} \otimes M_2$. Then by corollary 3 \mathfrak{B} is generated by a projection p and a positive invertible operator a with $1/2 \geqq a > 0$. Then let $P_1 = \begin{pmatrix} p & 0 \\ 0 & 0 \end{pmatrix}, P_2 = \begin{pmatrix} 1-p & 0 \\ 0 & 0 \end{pmatrix}, P_3 = \begin{pmatrix} a & t \\ t & 1-a \end{pmatrix}$ with $t = (a(1-a))^{1/2}$. These are obviously projections and $P_1 \cdot P_2 = 0$. Let \mathfrak{R} be the W^* -algebra generated by P_1, P_2 and P_3 and let $C = C^* \in \mathfrak{R}$. Then $C(P_1 + P_2) = (P_1 + P_2)C$ shows as before that C is diagonal, $C = \text{diag}(c_1, c_2)$. Then $CP_1 = P_1C$ and $CP_3 = P_3C$ show $c_1p = pc_1$ and $c_1a = ac_1$. By assumption $c_1 \in \mathfrak{B}'$. $CP_3 = P_3C$ gives further $tc_1 = c_2t$. We have

already shown $c_1 \in \mathfrak{B}$, thus $c_1 t = c_2 t$ or $c = c_2$ since t is invertible.

We should remark that the projections P_i $i = 1, 2, 3$ are infinite with infinite complement.

COROLLARY. *\mathfrak{A} is generated by two unitary operators U and V , which can be chosen such that $U^2 = 1$ and $V^3 = 1$.*

PROOF. Let P_1, P_2 and P_3 be the projection generators of \mathfrak{A} as we have determined them above. Let $U = 1 - 2P_3$ and $V = P_1 + P_2 e^{2\pi i/3} + (1 - P_1 - P_2) e^{4\pi i/3}$. Since the P_i $i = 1, 2, 3$ generate \mathfrak{A} also U and V generate \mathfrak{A} .

Theorem 3 improves a result by Saito [2], who did not show that two of the generating projections may be chosen orthogonal.

This corollary has an interesting consequence. Let \mathfrak{A} be a properly infinite W^* -algebra on the separable Hilbert space H and let U and V be unitary generators of \mathfrak{A} with $U^2 = 1 = V^3$. Let $G = Z_2 * Z_3$ be the free product of the cyclic group Z_2 of order 2 and Z_3 , the cyclic group of order 3, with generators α and β , $\alpha^2 = e = \beta^3$. Let $\pi(\alpha) = U$ and $\pi(\beta) = V$, then π determines a unitary representation of G such that $\pi(G)$ generates \mathfrak{A} . This shows that any properly infinite W^* -algebra \mathfrak{A} arises from a representation π of G . Because of our earlier remarks there exists even a continuous family π_k $k \in (0, 1)$ of representations of G such that $\pi_k(G)$ generates \mathfrak{A} and that the π_k are pairwise unitarily inequivalent. Conversely any W^* -algebra which comes from a representation of G is generated by two unitaries U, V with $U^2 = 1 = V^3$.

In the above theorems the separability of H cannot be dropped in general, because there exist properly infinite factors, which are not even countably generated. For example let G be the group of all finite permutations of an uncountable set and let \mathfrak{B} be the left ring of G . Then $\mathfrak{A} = \mathfrak{B} \otimes B(K)$, with K a separable infinite dimensional Hilbert space, is properly infinite. But \mathfrak{A} is not countably generated. To see this assume indirectly that \mathfrak{A} is generated by the operators $\{A_i\}_{i=1}^\infty$ with $A_i = (a_{j,k}^{(i)})$. Then \mathfrak{B} is generated by the countable set $\{a_{j,k}^{(i)}\}, i, j, k = 1, 2, \dots$. Every $a_{j,k}^{(i)}$ can be written as $a_{j,k}^{(i)} = \sum_{g \in G} \alpha_g^{(i,j,k)} U_g$ with $\sum_{g \in G} |\alpha_g^{(i,j,k)}|^2 < \infty$, where U_g is the translation by g on $\ell^2(G)$. Thus \mathfrak{B} is generated by the countable set $S = \{U_g | \alpha_g^{(i,j,k)} \neq 0 \text{ for some } (i, j, k)\}$. However the set of all $\{g \in G | U_g \in S\}$ generates a countable and thus proper subgroup H of G and thus the $U_g \in S$ do not generate \mathfrak{B} . This contradiction shows that \mathfrak{A} is not countably generated. Using some cardinal, arithmetic this result can be improved slightly.

REFERENCES

- [1] W. WOGEN, On Generators of von Neumann Algebras, Bull. Amer. Math. Soc., 1969.
- [2] T. SAITO, Generators of Certain von Neumann Algebras, Tôhoku Math. J., 20(1968), 101-105.
- [3] C. PEARCY, On Certain von Neumann Algebras, which are Generated by Partial Isometries, Proc. Amer. Math. Soc., 15(1964), 393-395.
- [4] J. G. STAMPFLI, Which Weighted Shifts are Subnormal, Pacific J. Math., 17(1966), p. 367.
- [5] A. BROWN, On a Class of Operators, Proc. Amer. Math. Soc., 4(1953), 723-728.
- [6] J. DIXMIER, Les C^* -algebres et leurs Representations, Gauthier Villars, Paris 1964.
- [7] H. BEHNCKE, Structure of Certain Nonnormal Operators, J. Math. Mech., 18(1968), 103-107.

INSTITUT FÜR ANGEWANDTE MATHEMATIK
UNIVERSITÄT HEIDELBERG
HEIDELBERG, DEUTSCHLAND