



REVIEW

Genetic factors influencing risk to orofacial clefts: today's challenges and tomorrow's opportunities [version 1; peer review: 2 approved]

Terri H. Beaty ¹, Mary L. Marazita², Elizabeth J. Leslie²

¹Department of Epidemiology, Johns Hopkins University, Baltimore, MD, 21205, USA

²Department of Oral Biology and Center for Craniofacial and Dental Genetics, University of Pittsburgh, Pittsburgh, PA, 15219, USA

V1 First published: 30 Nov 2016, 5(F1000 Faculty Rev):2800
<https://doi.org/10.12688/f1000research.9503.1>

Latest published: 30 Nov 2016, 5(F1000 Faculty Rev):2800
<https://doi.org/10.12688/f1000research.9503.1>

Abstract

Orofacial clefts include cleft lip (CL), cleft palate (CP), and cleft lip and palate (CLP), which combined represent the largest group of craniofacial malformations in humans with an overall prevalence of one per 1,000 live births. Each of these birth defects shows strong familial aggregation, suggesting a major genetic component to their etiology. Genetic studies of orofacial clefts extend back centuries, but it has proven difficult to define any single etiologic mechanism because many genes appear to influence risk. Both linkage and association studies have identified several genes influencing risk, but these differ across families and across populations. Genome-wide association studies have identified almost two dozen different genes achieving genome-wide significance, and there are broad classes of 'causal genes' for orofacial clefts: a few genes strongly associated with risk and possibly directly responsible for Mendelian syndromes which include orofacial clefts as a key phenotypic feature of the syndrome, and multiple genes with modest individual effects on risk but capable of disrupting normal craniofacial development under the right circumstances (which may include exposure to environmental risk factors). Genomic sequencing studies are now underway which will no doubt reveal additional genes/regions where variants (sequence and structural) can play a role in controlling risk to orofacial clefts. The real challenge to medicine and public health is twofold: to identify specific genes and other etiologic factors in families with affected members and then to devise effective interventions for these different biological mechanisms controlling risk to complex and heterogeneous birth defects such as orofacial clefts.

Keywords

orofacial clefts , CHD1 , ADAMTS9 , whole genome sequencing

Open Peer Review

Approval Status  

	1	2
version 1 30 Nov 2016		

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1. **Charlotte Hobbs**, University of Arkansas for Medical Sciences, Little Rock, USA
2. **Jacqueline Hecht**, University of Texas Medical School at Houston, Houston, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Terri H. Beaty (tbeaty1@jhu.edu)

Competing interests: The authors declare that they have no competing interests.

Grant information: Some of the work described in this review was partially supported by NIH grants DE014581, DE020073, and DE-018993 (T.H.B.); DE016148 (M.L.M.); DE025060 (E.J.L).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2016 Beaty TH *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Beaty TH, Marazita ML and Leslie EJ. **Genetic factors influencing risk to orofacial clefts: today's challenges and tomorrow's opportunities [version 1; peer review: 2 approved]** F1000Research 2016, 5(F1000 Faculty Rev):2800 <https://doi.org/10.12688/f1000research.9503.1>

First published: 30 Nov 2016, 5(F1000 Faculty Rev):2800 <https://doi.org/10.12688/f1000research.9503.1>

Introduction

Orofacial clefts represent a group of anatomically distinct birth defects where there is a gap or break in normal features of the mouth, most commonly the roof of the mouth (the palate) or the upper lip or both. As shown in **Figure 1**, the most common anatomical forms of orofacial clefts include cleft lip (CL), cleft palate (CP), and cleft lip and palate (CLP). Other features of the face and mouth can also be affected by orofacial clefts, but the overwhelming majority of all cases are infants with one of these three common forms¹. Normally, development of the face and mouth occurs very early in pregnancy and reflects a complex process of cell growth and migration followed by fusion of symmetric structures to form the palate separating the mouth and nasal cavity, with the outer structures of the face developing before the inner structures. Collectively CL, CLP, and CP are the most common craniofacial birth defects worldwide, and affected individuals face feeding difficulties early in life and typically require multiple corrective surgeries, therapeutic dental procedures, and speech therapy throughout childhood²⁻⁴. In addition, individuals born with an orofacial cleft have increased incidence of mental health problems and higher overall mortality rates at all stages of life, even in developed countries with good medical care^{5,6}. Where access to medical care is severely limited, infants born with CP or CLP generally have high mortality rates due to difficulty in breastfeeding, while untreated CL and CLP cases can face social discrimination throughout their lives. Thus, these birth defects have been subject to substantial selective pressure for most of human history.

Although clefts develop early in pregnancy, most epidemiologic data are based on cleft frequencies at birth. The worldwide average birth prevalence of all orofacial clefts is 9.92 per 10,000

(close to one per 1,000), but there is substantial variation among populations⁷. In general, East Asian and Native American populations have substantially higher birth prevalence rates than do European and South Asian populations, while African ancestry populations have lower birth prevalence rates⁸⁻¹⁰.

Environmental risk factors for orofacial clefts

There are several recognized environmental risk factors and multiple genes involved in the etiology of orofacial clefts, which means the etiology is complex and heterogeneous. Maternal smoking during pregnancy is a recognized risk factor for orofacial clefts, and it is estimated that 6.1% (95% confidence interval [CI]=4.4–7.7%) of orofacial clefts could be avoided by eliminating maternal smoking^{11,12}. Passive or “second-hand” exposure also appears to be a risk factor¹³. Evidence for maternal alcohol consumption as a risk factor is less consistent, although consuming large amounts over a short period of time (e.g. “binge drinking”) appears to increase risk¹⁴. Broad measures, e.g. socio-economic status, have been suggested as risk factors, but it is difficult to separate out the combined effects of maternal nutrition and health¹⁵. Several prescription medications have also been reported to increase risk when taken during the first trimester, including folate antagonists and certain other drugs exhibiting anti-folate properties⁷.

Interestingly, a substantial number of studies suggest genetic and environmental risk factors may “interact” to modify risk to orofacial clefts. Several common exposures, including maternal smoking, alcohol consumption, and vitamin supplementation (a protective factor), have yielded statistical evidence of gene–environment (GxE) interaction, although the results from different studies are not always consistent. While biologically GxE




Figure 1. Examples of orofacial clefts. A. Left unilateral cleft lip. B. Bilateral cleft lip. C. Left unilateral cleft lip plus cleft palate. D. Bilateral cleft lip plus cleft palate. E. Cleft palate.

interaction is eminently rational, replicating statistical evidence across studies is difficult and proving interaction exists remains a major challenge¹⁶.

Familial aggregation and the genetic basis of orofacial clefts

Orofacial clefts show strong familial aggregation, which suggests a major genetic component to their etiology. Marazita and Leslie¹⁷

have recently reviewed the extensive history of genetic studies of orofacial clefts, which dates back to the 18th century (see their [Table 1](#)). Despite decades of genetic research, it remains unclear exactly how many genes might control risk or how they act to influence risk to orofacial clefts. In general, close relatives of cleft cases have a greatly increased risk of being affected. Nationwide vital records from Norway¹⁸ show the relative risk to a first-degree relative of a case with CL with or without CP (CL/P, i.e. CL

Table 1. Genes achieving genome-wide significance as influencing risk to orofacial clefts.

Adapted and updated from Table 2 in Leslie and Marazita¹.

Associated locus	Candidate gene in region	Associated phenotype	Analysis method	References for genome-wide significance
1p36.13	<i>PAX7</i>	CL/P	GWAS meta-analysis, GWAS replication	36,50,51
1p36	<i>GRHL3</i>	CP	GWAS, GWAS replication	52,53
1p22	<i>ARHGAP29</i>	CL/P	GWAS	36,54
1q32.2	<i>IRF6</i>	CL/P	GWAS, linkage	25,27,33,36,54
2p13	<i>TGFA</i>	CL/P	Linkage	25
2p21	<i>THADA</i>	CL/P	GWAS meta-analysis	50
2p24	<i>FAM49A</i>	CL/P	GWAS	54
3p11	<i>EPHA3</i>	CL/P	GWAS meta-analysis	50
3q12	<i>COL8A1/FILIPIL</i>	CL/P	GWAS replication	51
3q27–28	<i>TP63</i>	CL/P	Linkage	25
8q21.3	<i>DCAF4L2</i>	CL/P	GWAS meta-analysis, GWAS replication	50,51,54
8q22.3	<i>BAALC</i>	CP & multivitamins	GWAS x E	55
8q24	Gene Desert	CL/P	GWAS	27,33,36,54,56
9q22.33	<i>FOXE1</i>	CL/P and CP	Linkage	25,28
9q31.1	<i>SMC2</i>	CP & maternal alcohol	GWAS x E	55
10q25.3	<i>VAX1</i>	CL/P	GWAS	27,36,54
12q14	<i>TBK1</i>	CP & maternal smoking	GWAS x E	55
13q31.2	<i>SPRY2</i>	CLP	GWAS meta-analysis	50
14q21–24	<i>PAX9, TGFB3, BMP4</i>	CL/P	Linkage	25
15q22	<i>TPM1</i>	CL/P	GWAS meta-analysis	50
15q24	<i>ARID3B</i>	CL/P	GWAS	54
16p13	<i>CREBBP</i>	CL/P	GWAS	57
16q24	<i>CRISPLD2</i>	CL/P	Linkage	25
17p13.1	<i>NTN1</i>	CL/P	GWAS meta-analysis, GWAS replication	49–51,54
17q22	<i>NOG</i>	CL/P	GWAS	27,49,54
17q23.2	<i>TANC2</i>	CL/P	GWAS	54
18q22	<i>ZNF236</i>	CP & maternal smoking	GWAS x E	55
19q13.11	<i>RHPN2</i>	CL/P	GWAS	54
20q12	<i>MAFB</i>	CL/P	GWAS	36,54

Abbreviations: CL/P, cleft lip with or without cleft palate; CLP, cleft lip and palate; CP, cleft palate; GWAS, genome-wide association study.

and CLP cases combined) was 32 times higher than the baseline population risk, and for relatives of a CP case this relative risk was 56 times higher than baseline risk. This increased risk drops with increasing genetic distance: nationwide registry data from Denmark spanning the entire second half of the 20th century¹⁹ show the relative risk among all first-degree relatives (parents, full siblings, and offspring) of CL/P cases was highest, while risks to second-degree relatives (half-siblings, avuncular relatives, and grandparents) were much smaller, and risks to third-degree relatives (first cousins and half-avuncular relatives) were not significantly above the baseline population risk in Denmark (see their Table 2). When rules of exclusions and phenotype definition were adjusted between these population-based studies (Norway and Denmark), both gave similar patterns of strong familial aggregation, i.e. relatives of cleft cases always had a high relative risk compared to the population's baseline risk, and there is a steep drop with increasing genetic distance between relatives.

Twin studies of orofacial clefts consistently show higher concordance rates in monozygotic (MZ or identical) twins compared to dizygotic (DZ or fraternal) twins, again suggesting a major role for genes in controlling risk of being affected. Clearly, because MZ twins have identical genes over their entire genome, anything less than 100% concordance indicates some non-genetic component to the etiology of orofacial clefts, but whenever MZ twins consistently show greater concordance than DZ twins, it suggests some proportion of the variation in risk is under genetic control. Grosen *et al.*²⁰ linked population-level vital records (birth certificate and twin registry data from Denmark) spanning most of the 20th century, achieving virtually complete ascertainment of cases for the latter part of the century. They showed MZ twins had much higher concordance rates than did DZ twins (47% MZ twins with 95% CI=31–64% compared to 8% for DZ twins with 95% CI=4–14%). These concordance rates yielded estimates of heritability (the proportion of variation in risk attributable to unobserved independent genes) of 91% for CL/P and 90% for CP. However, because the concordance rate among MZ twins is more than four times that of DZ twins, these Danish findings also raise the possibility that specific combinations of different genes (termed “epistasis”) may be important in the etiology of orofacial clefts.

While clearly orofacial cleft risk shows strong familial aggregation, it does not strictly adhere to Mendel's laws of inheritance, even within multiplex families (i.e. those with two or more affected members), which has bedeviled geneticists for decades. Statistical methods for fitting genetic models to family data are collectively termed “segregation analyses”, and these trace back to the foundations of human genetics. Statistical methods were initially developed in the early 20th century by statisticians such as R.A. Fisher, W. Weinberg, and J.B.S. Haldane^{17,21}. By the 1970s, segregation analysis had evolved to allow formal tests within a framework of “mixed models” that simultaneously considered straightforward Mendelian components (i.e. a “major gene”) and a more general residual “heritability” reflecting combined effects of many independent genes (collectively termed “polygenes”). A number of segregation analyses of orofacial clefts were performed during the early 1970s through the early 1990s for various racial/ethnic

groups²¹. Collectively, however, this modeling approach rarely resulted in definitive and reproducible findings, and the most suitable model of inheritance remained the “multifactorial threshold model”, which is exquisitely vague in defining genetic mechanisms and statistically cannot discriminate between a single major gene vs. multiple “polygenes”. This multifactorial model hypothesizes a continuous “genetic liability” for risk, plus unspecified non-genetic risk factors that could also influence risk. Under this very general model, it is possible to estimate the “heritability” or the proportion of variation in risk attributable to any number of independent, autosomal genes, but it is not possible to estimate the penetrance or allele frequencies at any one of these separate risk genes.

Still, this multifactorial threshold model adequately explains the clear gender differences in risk to CL/P and CP (where the former shows a distinct excess of affected males, while the latter has a slight excess of females, suggesting gender-specific thresholds) and the higher risk to relatives of more severe cases (i.e. bilateral versus unilateral CL/P)¹⁹. The sharp decline in relative risk to relatives of cases with increasingly distant relationships is also completely compatible with the multifactorial threshold model because the probability of sharing alleles identical by descent is constant whether one gene, a few genes, or many genes control risk. However, other explanations could potentially result in these same patterns of risk to relatives. Thus, even sophisticated statistical approaches to modeling genetic inheritance have failed to define any single biological mechanism controlling risk to orofacial clefts in all families; indeed it has been suggested that multiple genes, perhaps two to six independent genes (possibly up to 14) could determine risk to orofacial clefts²². This concept of multiple different genes involved in the etiology of orofacial clefts is increasingly borne out by the results of genomic studies, as summarized below.

Mapping genes for orofacial clefts

Different types of genetic studies create the opportunity to map genes in the absence of a well-defined model of inheritance. There are two general types of approaches: “linkage analysis” and “association analysis”. Linkage analysis always requires multiplex families and tests for co-segregation of observed genetic markers and a hypothetical gene controlling the affected vs. unaffected phenotype within a family, while association analysis simply tests for differences in frequencies of markers in samples of affected and unaffected individuals from a population, which could reflect linkage at a population level (termed “linkage disequilibrium”). Linkage analysis is a powerful approach for mapping individual genes for traits following clear Mendelian patterns within multiplex families, but it is less effective in mapping genes for complex traits (such as orofacial clefts).

Both linkage and association analysis can be used on one genetic marker or on many scattered over the entire genome. Importantly, both approaches can be adapted to accumulate evidence when multiple causal variants (in different genomic locations) exist in the data being analyzed (multiplex families or groups of cases and controls), and thus both can be used effectively to study orofacial clefts at the genome-wide level.

Numerous groups used linkage studies to map genes in multiplex cleft families²³, but limits on the number and size of these families and the underlying high level of “linkage heterogeneity” (where different families yielded evidence of co-segregation to different genes) made it extremely difficult to obtain consistent results. A consortium of multiple research groups pooling genome-wide markers (~400 microsatellite markers) identified linkage to six different chromosomal regions (chromosomes 1q32, 2p, 3q27–28, 9q21, 14q21–24, and 16q24)^{24,25}. Further fine-mapping studies of these regions showed consistent evidence of linkage to chromosome 1q32 near the *IRF6* gene. Interestingly, mutations in *IRF6* account for most of the known cases of Van der Woude syndrome, a Mendelian autosomal dominant syndrome which includes orofacial clefts as a key phenotype²³. In addition, markers in and near this gene have also consistently shown evidence of association in population-based studies. Another gene, *FOXE1* on chromosome 9q21, yielded consistent evidence of linkage in multiplex families^{25,26} and recent association studies^{27,28}. Thus, linkage studies based on multiplex families successfully identified some causal genes²⁹, but clearly multiple genes control risk to orofacial clefts and these vary from family to family.

Association studies ask a much more general question than linkage analysis, and typically this involves rather simple statistical comparisons of marker allele or genotype frequencies between groups of unrelated people. While interpreting a positive result from linkage analysis represents more compelling evidence that a putative gene controls risk, these multiplex families are rare (i.e. most cleft cases do not have any affected close relative) and individual families are not representative of the full population. Interpreting statistically significant results from an association study is more ambiguous and less definitive than results from linkage analysis, and unrecognized bias in the study design can produce spurious results (especially if the cases and controls are not drawn from the same population). As with linkage studies, one, several, or millions of markers can be compared between case and control groups, but of course as the number of tests increases some accommodation must be made for false positive results obtained by chance alone (i.e. where in truth there is no real difference between groups). Conversely, false negative results can become important too, i.e. where a key risk gene is missed because the markers used for tests of association are not informative and do a poor job of covering that gene in one population or another.

Association studies of orofacial clefts

Association studies using case–control designs have been used to examine numerous selected “candidate” genes for orofacial clefts (CL/P and less commonly CP) for over 25 years³⁰, but the results were not always consistent across studies. This inconsistency could merely be because of limited sample sizes of individual studies, subtle biases in study design, or could reflect genetic differences across populations. Interestingly, sequencing the candidate genes themselves (and the regions immediately surrounding the gene) has identified some rare mutations that would impair the function of the corresponding gene product and could be directly causal^{31,32}. While such rare variants are potentially biologically relevant and certainly justify further study of that candidate gene, their very low frequency in the population makes them difficult to incorporate either into

standard statistical tests or for use in predicting risk to individuals (which is notoriously difficult for heterogeneous disorders).

Beginning early in the 21st century and building upon the Human Genome Project, genome-wide association studies (GWAS) became feasible as genotyping technology improved to allow hundreds of thousands (or millions) of single nucleotide polymorphic (SNP) markers to be typed efficiently on large samples. The first GWAS of orofacial clefts used a classic case-control design to compare markers between CL/P cases from Germany to “universal” or unphenotyped German controls³³. The use of “universal” controls (typically adult controls from an independent study) is reasonable because the birth prevalence of orofacial clefts is relatively low. However, when conducting association studies, even subtle differences in the genetic background between the case and control groups can be a potential source of bias and must be considered. The key advantage of GWAS is its ability to identify simultaneously one or several regions of the genome likely to contain genes controlling risk to orofacial clefts. In the presence of multiple causal genes, there should be multiple genomic regions giving statistically significant results.

An alternative to the traditional case–control design is the family-based case–parent trio or “triad” design, which has the advantage of being robust to the confounding caused by “population stratification” or differences in marker allele frequencies between groups of cases and controls. Now the contrast is between the marker (allele, genotype, or haplotype) seen in the case and the set(s) that are possible given the parental mating type, yielding a comparison between the observed case and a matched set of “pseudo-controls” drawn from the parental mating type. This study design actually tests a composite null hypothesis of strict Mendelian transmission of marker alleles (or their genotypes), and rejecting this composite null represents evidence for both linkage and association^{34,35}. This design was used by Beaty *et al.*³⁶ in an international collaboration supported by the Genes and Environment Association (GENEVA) consortium³⁷ where distinct racial groups (European and Asian) were well represented.

To date, there have been seven independent GWAS for CL/P and two for CP. They have identified at least two dozen different genes/regions achieving genome-wide significance for CL/P, but only one for CP. Leslie and Marazita³⁸ listed a dozen different genes/regions attaining genome-wide significance either in one GWAS or from meta-analysis across multiple GWAS, and [Table 1](#) shows an updated list. Some of these genes have accumulated support from older candidate gene studies, and a few have been identified as causal genes for Mendelian malformation syndromes including orofacial clefts as a key phenotype. Of these genome-wide significant regions, four (*IRF6* on 1q32–41, the gene desert region on 8q24, 17q22, and 10q25.3) appear to account for 20–25% of the estimated genetic variation in risk or “heritability” to CL/P, a much larger proportion of the estimated heritability attributable to markers identified by GWAS than seen for many other complex disorders¹⁷.

However, the level of statistical support is frequently limited to one or another racial/ethnic group (e.g. the support for the gene

desert region in 8q24 is consistently strong in European ancestry groups but much less so among Asian ancestry groups). This may reflect limitations in “tagging” by widely used SNP marker panels³⁹ or true differences in the frequency of causal mutations across populations. Because this strong statistical evidence comes from SNPs in a “gene desert” with no recognized coding regions nearby, it is particularly intriguing that mouse models have suggested this region may contain a regulatory region critical for normal craniofacial development⁴⁰.

The future challenges: expanding into genomic sequencing

In many ways, the progression in genetic studies of orofacial clefts has repeatedly applied standard tools of genetics to a disorder that is unexpectedly complex and heterogeneous. Clearly there is a genetic basis to the etiology of orofacial clefts; however, it must involve multiple genes (maybe dozens) and several environmental risk factors (which may interact with different genes). Our analytical tools, both statistical and technological (for genotyping and sequencing), have greatly expanded in recent years but always seem to fall short of the mark in dealing with the high level of etiologic heterogeneity for this relatively rare disorder. Technological advances now make it feasible to sequence the entire exome (i.e. all gene regions known to code for proteins) and even the entire human genome. However, we still must carefully consider the most appropriate study design and how to analyze sequence data most effectively.

Family study designs can be very effective in identifying causal genes⁴¹, but if many rare mutations can lead to the same phenotype, it is entirely possible that each family could reflect effects of a unique gene or unique combination of genes. Bureau *et al.*^{42,43} recently showed how whole exome sequencing (WES) data in distant affected relatives (second-degree and more distant relatives) drawn from multiplex families originally recruited for linkage studies could be used to identify rare variants that modify risk to orofacial clefts. From a long list of candidate genes, this approach identified one novel variant in *CHDI* shared by three affected second cousins in a single multiplex family. *CHDI* is a strong candidate gene because mutations in this gene can cause both gastric cancer and orofacial clefts, and several studies have shown evidence of association or excess of rare variants in this gene^{44–47}. A comprehensive whole exome analysis revealed a separate novel variant in *ADAMTS9* achieving statistical significance over the entire exome in a small number of families⁴². While this gene had been identified as potentially causal in mouse models, this is the first evidence from human studies, and the risk allele is extremely rare. This same method of testing for rare shared variants can be applied to structural variants, such as deletions detected from WES data⁴⁸.

Leslie *et al.*⁴⁹ demonstrated the potential of sequencing studies in a large study of case–parent trios (where 13 regions around candidate genes were sequenced). This case–parent trio design allows the detection of *de novo* mutations, where a new mutation (not present in the parents) occurs in the child. This type of mutation simply cannot be identified in case–control studies, and such

mutations are by their nature extremely rare. Also, if a *de novo* mutation lies outside any known coding or regulatory region, it is difficult to judge its true functional status. This group concluded *de novo* variants probably don’t represent a major etiologic component for orofacial clefts, although they may be relevant for other complex diseases.

This case–parent trio study of sequencing data yielded evidence consistent with previous GWAS studies, where multiple genomic regions showed strong statistical evidence of linkage and association for common polymorphic markers, but it also was able to analyze rare variants that could themselves be directly causal for orofacial clefts. Careful analysis of low-frequency and rare variants using available methods for collapsing or combining multiple variants within a gene to assess the overall impact of that gene on risk yielded mixed results: two of 13 candidate regions (*NOG* and *NTN1*) showed some statistical evidence for distinct rare variants within the gene being important in controlling risk, while four of the 13 genes yielded no such evidence (*BMP4*, *FGFR2*, *MSX1*, and *PTCH1*). Because obtaining statistical evidence for rare variants is always difficult, even in large studies, building a biological case for a putative causal mutation requires functional studies for confirmation⁴⁹.

Large-scale whole genome sequencing (WGS) studies are now underway and are being applied to a number of complex and heterogeneous diseases (including orofacial clefts). Practical challenges in using WGS data include cleaning and processing the huge volume of data. The amount of data scales up dramatically from GWAS to WES to WGS: GWAS data represent hundreds of thousands or millions of typed markers (and potentially millions more imputed markers); WES represents 1–2% of the human genome sequenced 30–50 times and aligned to yield a consensus sequence for ~50 Mb of the genome; WGS represents similarly aligned read data on all 30 billion base pairs of the genome (although even with WGS there are gaps and some regions are not well covered). The primary advantage of WGS data is that it can be used to analyze any type of sequence variant from the reference genome (single nucleotide variants or SNVs, including the common SNP markers), small deletions (termed indels), or larger structural changes (e.g. deletions and duplications, which are collectively called copy number variants or CNVs, plus more complex structural variants such as inversions and translocations where the order of genes may be changed through chromosomal breakage and exchange). The challenge comes in defining the particular assessment/comparison being made with variants identified by WGS data (which is always a reflection of the study design and hypothesis being considered), and then interpreting any “significant” differences from expected. The study design always determines the hypothesis to be tested (strict Mendelian transmission, co-segregation, differences in frequencies, etc.), but the huge number of tests must be considered, along with the potential biases inherent in the study design. Dealing with WGS data will involve tens of millions of markers or SNVs on individuals, and most of these will be quite rare. Most observed sequence variants will occur outside known coding regions, and our ability to recognize important regulatory regions remains limited.

The more important challenge facing genetic studies of orofacial clefts is to develop strategies for dealing with multiple causal genes within any data set and applying these strategies effectively to orofacial clefts as a complex and heterogeneous birth defect. The full range of study designs (selected relatives from large multiplex families, nuclear families, case–parent trios, and groups of unrelated individuals) is available, but interpreting the biological function of a novel variant detected by sequencing will not be simple. Furthermore, because any given patient could represent the effects of one or more variants in several different genes, translating new genetic findings into meaningful medical or public health applications remains a daunting task. In reality, for orofacial clefts (or any other birth defect), because the first occurrence occurs well before birth, it is difficult to define opportunities to intervene. Nonetheless, the tools to detect genes controlling risk to orofacial clefts have expanded greatly in this first part of the 21st century, and it is our challenge to use them as effectively and efficiently as possible. It is likely that there will be classes of “causal genes” and “genetic risk factors”: a few genes with “major effects” may simultaneously control truly Mendelian malformation syndromes and show association with apparent non-syndromic forms of orofacial clefts, while possibly many “genetic risk factors” have the typical modest estimated effect sizes identified through GWAS but nonetheless achieve genome-wide significance as studies are combined via meta-analysis.

Abbreviations

CI, confidence interval; CL, cleft lip; CL/P, cleft lip with or without cleft palate; CLP, cleft lip and palate; CP, cleft palate; GWAS, genome-wide association study; GxE, gene–environment; SNP, single nucleotide polymorphism; SNV, single nucleotide variant; WES, whole exome sequencing; WGS, whole genome sequencing.

Competing interests

The authors declare that they have no competing interests.

Grant information

Some of the work described in this review was partially supported by NIH grants DE014581, DE020073, and DE-018993 (T.H.B.); DE016148 (M.L.M.); DE025060 (E.J.L.).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

Thanks to our many colleagues and mentors in the field of orofacial genetics.

References



- Leslie EJ, Marazita ML: **Genetics of Orofacial Cleft Birth Defects.** *Curr Genet Med Rep.* 2015; **3**(3): 118–26.
[Publisher Full Text](#)
- Wehby GL, Cassell CH: **The impact of orofacial clefts on quality of life and healthcare use and costs.** *Oral Dis.* 2010; **16**(1): 3–10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Semb G, Brattström V, Mølsted K, et al.: **The Eurocleft study: intercenter study of treatment outcome in patients with complete cleft lip and palate. Part 1: introduction and treatment experience.** *Cleft Palate Craniofac J.* 2005; **42**(1): 64–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Berk NW, Marazita ML: **The Costs of Cleft Lip and Palate: Personal and Societal Implications.** In: Wyszynski DF, editor. *Cleft Lip and Palate: From Origin to Treatment.* New York: Oxford University Press; 2002.
- Bille C, Winther JF, Bautz A, et al.: **Cancer risk in persons with oral cleft—a population-based study of 8,093 cases.** *Am J Epidemiol.* 2005; **161**(11): 1047–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Christensen K, Juel K, Herskind AM, et al.: **Long term follow up study of survival associated with cleft lip and palate at birth.** *BMJ.* 2004; **328**(7453): 1405.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Watkins SE, Meyer RE, Strauss RP, et al.: **Classification, epidemiology, and genetics of orofacial clefts.** *Clin Plast Surg.* 2014; **41**(2): 149–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
- IPDTC Working Group: **Prevalence at birth of cleft lip with or without cleft palate: data from the International Perinatal Database of Typical Oral Clefts (IPDTC).** *Cleft Palate Craniofac J.* 2011; **48**(1): 66–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Allagh KP, Shamanna BR, Murthy GV, et al.: **Birth prevalence of neural tube defects and orofacial clefts in India: a systematic review and meta-analysis.** *PLoS One.* 2015; **10**(3): e0118961.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Butali A, Adeyemo WL, Mossey PA, et al.: **Prevalence of orofacial clefts in Nigeria.** *Cleft Palate Craniofac J.* 2014; **51**(3): 320–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Honein MA, Devine O, Grosse SD, et al.: **Prevention of orofacial clefts caused by smoking: implications of the Surgeon General's report.** *Birth Defects Res A Clin Mol Teratol.* 2014; **100**(11): 822–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Little J, Cardy A, Munger RG: **Tobacco smoking and oral clefts: a meta-analysis.** *Bull World Health Organ.* 2004; **82**(3): 213–8.
[PubMed Abstract](#) | [Free Full Text](#)
- Kummet CM, Moreno LM, Wilcox AJ, et al.: **Passive Smoke Exposure as a Risk Factor for Oral Clefts—A Large International Population-Based Study.** *Am J Epidemiol.* 2016; **183**(9): 834–41.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- DeRoo LA, Wilcox AJ, Lie RT, et al.: **Maternal alcohol binge-drinking in the first trimester and the risk of orofacial clefts in offspring: a large population-based pooling study.** *Eur J Epidemiol.* 2016; **31**(10): 1021–1034.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
- Mossey P: **Epidemiology underpinning research in the aetiology of orofacial clefts.** *Orthod Craniofac Res.* 2007; **10**(3): 114–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Aschard H: **A perspective on interaction effects in genetic association studies.** *Genet Epidemiol.* 2016; **40**(8): 678–688.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Marazita ML, Leslie EJ: **Genetics of Nonsyndromic Clefting.** In: Losee J, Kirschner R, editors. *Comprehensive Cleft Care.* second edition ed. Boca Raton FL: CRC Press, Taylor & Francis Group; 2016; 207–24.
- Sivertsen A, Wilcox AJ, Skjaerven R, et al.: **Familial risk of oral clefts by morphological type and severity: population based cohort study of first degree relatives.** *BMJ.* 2008; **336**(7641): 432–4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Grosen D, Chevrier C, Skytthe A, et al.: **A cohort study of recurrence patterns among more than 54,000 relatives of oral cleft cases in Denmark: support for the multifactorial threshold model of inheritance.** *J Med Genet.* 2010; **47**(3): 162–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Grosen D, Bille C, Petersen I, et al.: **Risk of oral clefts in twins.** *Epidemiology.* 2011; **22**(3): 313–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

21. Marazita ML: **Segregation Analysis**. In: Wyszynski DF, editor. *Cleft Lip and Palate: From Origin to Treatment*. New York: Oxford University Press; 2002; 222–33.
22. Schliekelman P, Slatkin M: **Multiplex relative risk and estimation of the number of loci underlying an inherited disease**. *Am J Hum Genet*. 2002; 71(6): 1369–85. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
23. Dixon MJ, Marazita ML, Beaty TH, *et al.*: **Cleft lip and palate: understanding genetic and environmental influences**. *Nat Rev Genet*. 2011; 12(3): 167–78. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Marazita ML, Murray JC, Lidral AC, *et al.*: **Meta-analysis of 13 genome scans reveals multiple cleft lip/palate genes with novel loci on 9q21 and 2q32-35**. *Am J Hum Genet*. 2004; 75(2): 161–73. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Marazita ML, Lidral AC, Murray JC, *et al.*: **Genome scan, fine-mapping, and candidate gene analysis of non-syndromic cleft lip with or without cleft palate reveals phenotype-specific differences in linkage and association results**. *Hum Hered*. 2009; 68(3): 151–70. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
26. Moreno LM, Mansilla MA, Bullard SA, *et al.*: **FOXE1 association with both isolated cleft lip with or without cleft palate, and isolated cleft palate**. *Hum Mol Genet*. 2009; 18(24): 4879–96. [PubMed Abstract](#) | [Free Full Text](#)
27. Mangold E, Ludwig KU, Birnbaum S, *et al.*: **Genome-wide association study identifies two susceptibility loci for nonsyndromic cleft lip with or without cleft palate**. *Nat Genet*. 2010; 42(1): 24–6. [PubMed Abstract](#) | [Publisher Full Text](#)
28. **F** Ludwig KU, Böhmer AC, Rubini M, *et al.*: **Strong association of variants around FOXE1 and orofacial clefting**. *J Dent Res*. 2014; 93(4): 376–81. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
29. Khandelwal KD, van Bokhoven H, Roscioli T, *et al.*: **Genomic approaches for studying craniofacial disorders**. *Am J Med Genet C Semin Med Genet*. 2013; 163C(4): 218–31. [PubMed Abstract](#) | [Publisher Full Text](#)
30. Ardinger HH, Buettow KH, Bell GI, *et al.*: **Association of genetic variation of the transforming growth factor-alpha gene with cleft lip and palate**. *Am J Hum Genet*. 1989; 45(3): 348–53. [PubMed Abstract](#) | [Free Full Text](#)
31. Gowans LJ, Adeyemo WL, Eshete M, *et al.*: **Association Studies and Direct DNA Sequencing Implicate Genetic Susceptibility Loci in the Etiology of Nonsyndromic Orofacial Clefts in Sub-Saharan African Populations**. *J Dent Res*. 2016; 95(11): 1245–56. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Leslie EJ, Murray JC: **Evaluating rare coding variants as contributing causes to non-syndromic cleft lip and palate**. *Clin Genet*. 2013; 84(5): 496–500. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. **F** Birnbaum S, Ludwig KU, Reutter H, *et al.*: **Key susceptibility locus for nonsyndromic cleft lip with or without cleft palate on chromosome 8q24**. *Nat Genet*. 2009; 41(4): 473–7. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
34. Laird NM, Lange C: **Family-based designs in the age of large-scale gene-association studies**. *Nat Rev Genet*. 2006; 7(5): 385–94. [PubMed Abstract](#) | [Publisher Full Text](#)
35. Laird NM, Lange C: **Family-based methods for linkage and association analysis**. *Adv Genet*. 2008; 60: 219–52. [PubMed Abstract](#) | [Publisher Full Text](#)
36. Beaty TH, Murray JC, Marazita ML, *et al.*: **A genome-wide association study of cleft lip with and without cleft palate identifies risk variants near MAFB and ABCA4**. *Nat Genet*. 2010; 42(6): 525–9. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Cornelis MC, Agrawal A, Cole JW, *et al.*: **The Gene, Environment Association Studies consortium (GENEVA): maximizing the knowledge obtained from GWAS by collaboration across studies of multiple conditions**. *Genet Epidemiol*. 2010; 34(4): 364–72. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Leslie EJ, Marazita ML: **Genetics of cleft lip and cleft palate**. *Am J Med Genet C Semin Med Genet*. 2013; 163C(4): 246–58. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Murray T, Taub MA, Ruczinski I, *et al.*: **Examining markers in 8q24 to explain differences in evidence for association with cleft lip with/without cleft palate between Asians and Europeans**. *Genet Epidemiol*. 2012; 36(4): 392–9. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. **F** Uslu VV, Petretich M, Ruf S, *et al.*: **Long-range enhancers regulating Myc expression are required for normal facial morphogenesis**. *Nat Genet*. 2014; 46(7): 753–8. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
41. Wijsman EM: **The role of large pedigrees in an era of high-throughput sequencing**. *Hum Genet*. 2012; 131(10): 1555–63. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Bureau A, Younkin SG, Parker MM, *et al.*: **Inferring rare disease risk variants based on exact probabilities of sharing by multiple affected relatives**. *Bioinformatics*. 2014; 30(15): 2189–96. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. Bureau A, Parker MM, Ruczinski I, *et al.*: **Whole exome sequencing of distant relatives in multiplex families implicates rare variants in candidate genes for oral clefts**. *Genetics*. 2014; 197(3): 1039–44. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
44. Rafighdoost H, Hashemi M, Narouei A, *et al.*: **Association between CDH1 and MSX1 gene polymorphisms and the risk of nonsyndromic cleft lip and/or cleft palate in a southeast Iranian population**. *Cleft Palate Craniofac J*. 2013; 50(5): e98–e104. [PubMed Abstract](#) | [Publisher Full Text](#)
45. **F** Hozyasz KK, Mostowska A, Wójcicki P, *et al.*: **Nucleotide variants of the cancer predisposing gene CDH1 and the risk of non-syndromic cleft lip with or without cleft palate**. *Fam Cancer*. 2014; 13(3): 415–21. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
46. **F** Ittiwut R, Ittiwut C, Siriwan P, *et al.*: **Variants of the CDH1 (E-Cadherin) Gene Associated with Oral Clefts in the Thai Population**. *Genet Test Mol Biomarkers*. 2016; 20(7): 406–9. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
47. **F** Brito LA, Yamamoto GL, Melo S, *et al.*: **Rare Variants in the Epithelial Cadherin Gene Underlying the Genetic Etiology of Nonsyndromic Cleft Lip with or without Cleft Palate**. *Hum Mutat*. 2015; 36(11): 1029–33. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
48. Fu J, Beaty TH, Scott AF, *et al.*: **PgmNr 444: Whole exome association of rare deletions in multiplex oral cleft families**. *Genet Epidemiol*. 2016; [in press]. [Publisher Full Text](#)
49. Leslie EJ, Taub MA, Liu H, *et al.*: **Identification of functional variants for cleft lip with or without cleft palate in or near PAX7, FGFR2, and NOG by targeted sequencing of GWAS loci**. *Am J Hum Genet*. 2015; 96(3): 397–411. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Ludwig KU, Mangold E, Herms S, *et al.*: **Genome-wide meta-analyses of nonsyndromic cleft lip with or without cleft palate identify six new risk loci**. *Nat Genet*. 2012; 44(9): 968–71. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Beaty TH, Taub MA, Scott AF, *et al.*: **Confirming genes influencing risk to cleft lip with/without cleft palate in a case-parent trio study**. *Hum Genet*. 2013; 132(7): 771–81. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
52. Leslie EJ, Liu H, Carlson JC, *et al.*: **A Genome-wide Association Study of Nonsyndromic Cleft Palate Identifies an Etiologic Missense Variant in GRHL3**. *Am J Hum Genet*. 2016; 98(4): 744–54. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. **F** Mangold E, Böhmer AC, Ishorst N, *et al.*: **Sequencing the GRHL3 Coding Region Reveals Rare Truncating Mutations and a Common Susceptibility Variant for Nonsyndromic Cleft Palate**. *Am J Hum Genet*. 2016; 98(4): 755–62. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
54. Leslie EJ, Carlson JC, Shaffer JR, *et al.*: **A multi-ethnic genome-wide association study identifies novel loci for non-syndromic cleft lip with or without cleft palate on 2p24.2, 17q23 and 19q13**. *Hum Mol Genet*. 2016; pii: ddw104. [PubMed Abstract](#)
55. Beaty TH, Ruczinski I, Murray JC, *et al.*: **Evidence for gene-environment interaction in a genome wide study of nonsyndromic cleft palate**. *Genet Epidemiol*. 2011; 35(6): 469–78. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Grant SF, Wang K, Zhang H, *et al.*: **A genome-wide association study identifies a locus for nonsyndromic cleft lip with or without cleft palate on 8q24**. *J Pediatr*. 2009; 155(6): 909–13. [PubMed Abstract](#) | [Publisher Full Text](#)
57. **F** Sun Y, Huang Y, Yin A, *et al.*: **Genome-wide association study identifies a new susceptibility locus for cleft lip with or without a cleft palate**. *Nat Commun*. 2015; 6: 6414. [PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)

Open Peer Review

Current Peer Review Status:  

Editorial Note on the Review Process

Faculty Reviews are review articles written by the prestigious Members of **Faculty Opinions**. The articles are commissioned and peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

1. Jacqueline Hecht

Department of Pediatrics, University of Texas Medical School at Houston, Houston, TX, USA

Competing Interests: No competing interests were disclosed.

2. Charlotte Hobbs

Department of Pediatrics, University of Arkansas for Medical Sciences, Little Rock, AR, USA

Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research