

## ORIGINAL ARTICLE

## Genetic markers associated with abstinence length in alcohol-dependent subjects treated with acamprosate

VM Karpyak<sup>1</sup>, JM Biernacka<sup>1,2</sup>, JR Geske<sup>2</sup>, GD Jenkins<sup>2</sup>, JM Cunningham<sup>3</sup>, J Rüegg<sup>4</sup>, O Kononenko<sup>5</sup>, AA Leontovich<sup>2</sup>, OA Abulseoud<sup>1</sup>, DK Hall-Flavin<sup>1</sup>, LL Loukianova<sup>1</sup>, TD Schneekloth<sup>1</sup>, MK Skime<sup>1</sup>, J Frank<sup>6</sup>, MM Nöthen<sup>7</sup>, M Rietschel<sup>6</sup>, F Kiefer<sup>8</sup>, KF Mann<sup>8</sup>, RM Weinshilboum<sup>9</sup>, MA Frye<sup>1</sup> and DS Choi<sup>1,9</sup>

Acamprosate supports abstinence in some alcohol-dependent subjects, yet predictors of response are unknown. To identify response biomarkers, we investigated associations of abstinence length with polymorphisms in candidate genes in glycine and glutamate neurotransmission pathways and genes previously implicated in acamprosate response. Association analyses were conducted in the discovery sample of 225 alcohol-dependent subjects treated with acamprosate for 3 months in community-based treatment programs in the United States. Data from 110 alcohol-dependent males treated with acamprosate in the study PREDICT were used for replication of the top association findings. Statistical models were adjusted for relevant covariates, including recruitment site and baseline clinical variables associated with response. In the discovery sample, shorter abstinence was associated with increased intensity of alcohol craving and lower number of days between the last drink and initiation of acamprosate treatment. After adjustment for covariates, length of abstinence was associated with the *GRIN2B* rs2058878 ( $P=4.6 \times 10^{-5}$ ). In the replication sample, shorter abstinence was associated with increased craving, increased depressive mood score and higher alcohol consumption. Association of abstinence length with *GRIN2B* rs2058878 was marginally significant ( $P=0.0675$ ); as in the discovery sample, the minor A allele was associated with longer abstinence. Furthermore, rs2300272, which is in strong linkage disequilibrium with rs2058878, was also associated with abstinence length ( $P=0.049$ ). This is the first report of a replicated association of genetic markers with the length of abstinence in acamprosate-treated alcoholics. Investigation of the underlying mechanisms of this association and its usefulness for individualized treatment selection should follow.

*Translational Psychiatry* (2014) 4, e462; doi:10.1038/tp.2014.103; published online 7 October 2014

## INTRODUCTION

The economic costs associated with alcohol use and misuse are staggering, and evidence-based treatment strategies are needed.<sup>1,2</sup> Meta-analyses favor acamprosate for its ability to support abstinence,<sup>3–5</sup> which is the most stable type of remission in alcohol-dependent subjects.<sup>6</sup> Only a fraction of those treated respond to acamprosate.<sup>7</sup> Clinical characteristics, including the number of sober days before initiation of acamprosate treatment, were shown to be associated with abstinence length.<sup>8–10</sup> However, no biomarkers allowing reliable identification of potential responders to acamprosate treatment are currently known. It is expected that a pharmacogenomic approach may lead to the discovery of such biomarkers, enabling individualized recommendations for treatment selection and improved treatment outcomes.<sup>11,12</sup>

Acamprosate shares structural similarities with glycine and glutamate<sup>13</sup> and is thought to work by counteracting the 'relief craving' associated with increased glutamate levels in alcoholics with a history of withdrawal.<sup>14,15</sup> Several genes, including *GATA4*, *PER2* and *SLC29A1*, were associated with response to acamprosate treatment in human and animal studies.<sup>16–18</sup> Recent findings also

indicate that glycine, which activates N-methyl-D-aspartate (NMDA) and glycine receptors,<sup>19,20</sup> may have an important role in alcohol use disorders and treatment response.<sup>21–25</sup> Therefore, in an effort to identify genetic markers associated with abstinence in alcohol-dependent human subjects receiving acamprosate treatment, we investigated sequence variation in genes involved in composition of the glycine and NMDA receptors, glycine and glutamate reuptake, synthesis and metabolism, along with sequence variation in candidate genes previously reported to be associated with acamprosate response in human and animal studies.

To make study findings practically meaningful, we considered several potentially important issues. First, we conducted our study in samples reflecting the population where results will be applied, that is, alcohol-dependent subjects treated in community-based programs. Second, we monitored non-genetic (that is, clinical and demographic) patient characteristics and accounted for relevant covariates and potential confounders in the genetic analyses. Third, to ensure that findings were relevant and readily translated into clinical practice, we utilized diagnostic tools and outcome measures used in community-based treatment programs. In

<sup>1</sup>Department of Psychiatry and Psychology, Mayo Clinic, Rochester, MN, USA; <sup>2</sup>Department of Health Sciences Research, Mayo Clinic, Rochester, MN, USA; <sup>3</sup>Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN, USA; <sup>4</sup>Department of Clinical Neuroscience, Karolinska Institute, Stockholm, Sweden; <sup>5</sup>Department of Pharmaceutical Biosciences, Uppsala University, Uppsala, Sweden; <sup>6</sup>Department of Genetic Epidemiology in Psychiatry, Central Institute of Mental Health, Medical Faculty Mannheim/Heidelberg University, Mannheim, Germany; <sup>7</sup>Department of Genomics Life and Brain Research Centre, Institute of Human Genetics, University of Bonn, Bonn, Germany; <sup>8</sup>Department of Addictive Behavior and Addiction Medicine, Central Institute of Mental Health, Medical Faculty Mannheim/Heidelberg University, Mannheim, Germany and <sup>9</sup>Department of Molecular Pharmacology and Experimental Therapeutics, Mayo Clinic, Rochester, MN, USA. Correspondence: Dr VM Karpyak, Department of Psychiatry and Psychology, Mayo Clinic, Rochester, 200 First Street South West, Rochester, MN 55905, USA.

E-mail: karpyak.victor@mayo.edu

Received 25 July 2014; accepted 24 August 2014

accordance with these considerations, we conducted an open label, naturalistic prospective study investigating genetic markers associated with abstinence in alcohol-dependent subjects treated with acamprosate in community-based programs. Replication analyses were performed using data from a previously conducted study of acamprosate (PREDICT<sup>26</sup>).

We identified genetic markers associated with the length of sobriety during acamprosate treatment. This is an important step toward the development of personalized treatment recommendations for patients with alcohol use disorders, as genetic markers may be used for selection of patients who have the highest probability of responding to acamprosate treatment. We think that with the limited number of antidiabetic medications and lack of uniformity in treatment response, the development of such recommendations is of special importance for improvement of treatment outcomes in patients with alcohol use disorders.

## MATERIALS AND METHODS

### Discovery study

**Subjects and recruitment sites.** This study was approved by the Institutional Review Board of the Mayo Clinic Rochester and Mayo Clinic Health System, and was conducted according to the Code of Ethics of the World Medical Association (Declaration of Helsinki). All participants signed informed consent approved by the Mayo Clinic Institutional Review Board.

Both men and women between the ages of 18 and 80 with a primary diagnosis of current alcohol dependence based on DSM-IV-TR criteria with the last drink 5 or more days before enrollment were included in the study. We excluded subjects unable to provide informed consent; those unable to speak English; those with psychotic disorders or unstable psychiatric or medical conditions (see Supplementary Materials); women who were pregnant, lactating or planning to become pregnant; subjects taking disulfiram; and those allergic to acamprosate.

Participants were recruited from community-based residential and outpatient treatment programs affiliated with Mayo Clinic in Rochester, Minnesota, and the Mayo Clinic Health System sites in Austin, Minnesota, Albert Lea, Minnesota, and La Crosse, Wisconsin (Supplementary Table S1). In addition, self-referred participants residing in communities adjacent to referral sites not enrolled in treatment programs but interested in taking acamprosate were recruited and included in the analyses as a separate 'study site.' A description of programs and the number of subjects recruited at each site is presented in Supplementary Materials.

**Assessment.** A detailed description of assessments is presented in Supplementary Table S2 and summarized below. The lifetime presence of alcohol dependence and comorbid disorders was assessed using the semistructured interview PRISM.<sup>27</sup> Time Line Follow-Back<sup>28</sup> was used to assess alcohol consumption before and during treatment. Craving intensity was assessed using Penn Alcohol Craving Scale (PACS).<sup>29,30</sup> Association of alcohol use with positive or negative emotional states was assessed using the Inventory of Drug Taking Situations.<sup>31,32</sup> The intensity of depressive symptoms and anxiety were assessed using a depression scale from the patient health questionnaire,<sup>33</sup> and the Generalized Anxiety Disorder Assessment.<sup>34</sup> Alcoholic Anonymous attendance monitoring<sup>35</sup> was used to estimate utilization of support networks.

Treatment outcomes including abstinence (defined as time between initiation of acamprosate treatment and first alcohol use) or alcohol use were assessed with self-report (Timeline Follow-Back) and from available medical records during follow-up visits. Gamma-glutamyl transpeptidase measurements were used to assess the accuracy of self-reported sobriety. Compliance with acamprosate was assessed by pill count. The use of other medications was also monitored and considered as potential covariates in the analyses.

**Selection of candidate genetic targets in the discovery study.** Candidate genes (Supplementary Table S3) were selected based on the following considerations. First, we used a pathway-based approach to systematically investigate genes encoding enzymes involved in glycine metabolism, glycine transporters and subunits of glycine receptors and the NMDA receptor, which is known to be involved in glycine effects.<sup>19,36</sup> Second, because acamprosate action is thought to be associated with increased brain glutamate levels,<sup>14,15</sup> we included genes associated with glutamate

reuptake, synthesis and degradation. Third, we included candidate genes reported to be associated with acamprosate treatment outcomes in human or animal studies<sup>16–18</sup> and genes associated with alcoholism treatment response.<sup>37–39</sup>

We used SNPPicker<sup>40</sup> for selection of tag single-nucleotide polymorphisms (SNPs) to attain comprehensive coverage of each candidate gene. In addition, 28 SNPs were selected from a panel of ancestry informative markers<sup>41</sup> to verify self-reported ancestry. The list of 548 candidate SNPs included in the study is presented in Supplementary Table S4.

**Genotyping and quality control.** Genotyping was conducted using Illumina Golden Gate custom panel of 576 single-nucleotide variants.<sup>42</sup> Of the 576 SNPs included in the genotyping panel (548 candidate SNPs, 28 ancestry informative SNPs), 19 failed and 11 had minor allele frequencies < 2% and were excluded from analysis. Of the 433 subjects who started acamprosate treatment and were genotyped, four were excluded from analyses because of low call rates (< 90%). As part of genotype quality control, 18 subjects were genotyped in duplicate and a CEPH trio was genotyped eight times. No discordant genotype calls were observed for these replicated samples. Hardy-Weinberg equilibrium was evaluated; however, no SNPs were removed (Supplementary Table S4). To verify self-reported race, STRUCTURE v2.3.3 (Stanford University, Stanford, CA, USA) was used to estimate ancestry of individuals based on the 28 ancestry informative markers. Four self-reported European Americans were removed as they appeared to have > 50% non-European ancestry.

**Data analyses.** Time-to-event (first alcohol use) survival analysis methods were used to examine the association of clinical and genetic markers with treatment outcomes. Demographic and baseline clinical characteristics were first evaluated for associations with length of abstinence using univariate Cox proportional hazard models. As we found an association of treatment outcome with recruitment site, likely reflecting differences in population characteristics and non-pharmacological treatment components across sites, we controlled for recruitment site as a covariate in all subsequent analyses. Backward stepwise variable selection with covariates that were univariately related to response ( $P < 0.05$ ) was performed to identify other relevant covariates and potential confounders for the pharmacogenetic analyses. On the basis of the results of this selection process, all genetic analyses were adjusted for enrollment site, days since last drink at baseline and baseline PACS.

### Replication study

**Study sample.** To replicate top association findings in the discovery sample, we conducted similar analyses in a subset of participants from the study PREDICT, a double-blind randomized controlled trial that compared treatment outcomes including length of abstinence among alcohol-dependent subjects of German descent recruited from inpatient facilities and treated with acamprosate, naltrexone or placebo for 3 months.<sup>26</sup> Clinical and genetic data from 110 males treated with acamprosate who were also included in a genome-wide association study of alcohol dependence<sup>43</sup> and, thus, had genotype data available, were included in the replication analyses. The diagnostic assessments and instruments used in the study PREDICT were described previously.<sup>26</sup>

**Genotyping, selection of candidate genetic targets and data analysis in the replication sample.** A subset of PREDICT study participants was genotyped using Illumina Human-Hap 550, Illumina Human 610 and Illumina Human 660w quad BeadChips (Illumina, San Diego, CA, USA) for a genome-wide association study of alcohol dependence.<sup>43,44</sup> We selected four top SNPs associated with treatment outcome in the discovery sample ( $P < 0.001$ ) for replication. Three of those SNPs were not genotyped in the replication sample. Therefore, we imputed those SNPs using the ShapeIT tool, v1 (Christchurch, New Zealand) for phasing and IMPUTE (v2.3, Oxford, UK) for imputation.<sup>45,46</sup> The European sample from 1000 genomes data was used as the reference panel, and the three imputed SNPs had dosage  $R^2 > 0.99$ .

Analysis of our Replication Set data followed the same approach as the Discovery Set analysis. Specifically, Cox proportional hazard models were used to examine the association of length of abstinence with clinical variables and candidate SNPs. Backward stepwise variable selection was again used to identify covariates and potential confounders for the genetic analyses. Genetic association analyses were then performed for SNP genotypes coded as the minor allele count.

**RESULTS**

Clinical and demographic variables associated with abstinence in the discovery sample

Of the 443 subjects who started acamprosate, 225 European American subjects with available 3-month outcome data (any alcohol use  $N=93$ ; abstainers  $N=132$ ) were included in the analyses. The length of abstinence was significantly associated with the recruitment site as well as several clinical variables (Table 1). Increased craving and depression scores (measured by PACS and patient health questionnaire 9, respectively) were associated with shorter abstinence. Longer abstinence was associated with increased number of days since last drink before initiation of acamprosate, increased attendance of Alcoholics Anonymous meetings, having an Alcoholics Anonymous sponsor and attendance at counseling sessions. However, after adjustment for study site, only the baseline PACS score and the number of days between the last drink and initiation of acamprosate treatment remained strongly associated with treatment outcome ( $P < 0.0001$  and  $P = 0.0002$ , respectively). Using a backward stepwise variable selection process, covariates (including study site, PACS score and the number of days between last drink and initiation of acamprosate treatment) were selected for adjustment in pharmacogenomic analyses.

The observed changes in plasma gamma-glutamyl transpeptidase levels between baseline and 3-month follow-up were consistent with self-reported abstinence and alcohol use. As shown in Supplementary Figure S1, gamma-glutamyl transpeptidase levels were elevated at baseline and decreased markedly during treatment in abstainers and non-abstainers. At the 3-month follow-up, average gamma-glutamyl transpeptidase levels were significantly lower and closer to the normative range in abstainers compared with non-abstainers ( $P < 0.001$ ).

Single SNP association with treatment outcomes in the discovery sample

A complete list of candidate SNPs and their association with the length of abstinence is presented in Supplementary Table S4, and summarized in Figure 1. The top results ( $P < 0.001$ ) are also presented in Table 2. The strongest association finding, minor *GRIN2B* rs2058878 A allele, remains significantly associated with longer abstinence after Bonferroni correction for the number of SNPs included in the analyses ( $P = 4.6 \times 10^{-5}$ , corrected  $P = 0.024$ ). As the replication sample included only alcohol-dependent males, we also explored the association of the top four SNPs with treatment outcome in a subsample of males. As shown in Table 2, the evidence for association with treatment outcome was stronger for minor *GLRB* rs17035723 A allele in the male subset (hazard ratio = 2.88,  $P = 0.98 \times 10^{-5}$ ). The hazard ratios of the three *GRIN2B* SNPs, including rs2058878, were also higher in males; however,  $P$ -values reflecting their association with treatment outcome were less significant, possibly because of the decreased sample size in the subsample of males.


Clinical and demographic variables associated with abstinence in the replication sample

As shown in Table 3, the replication sample included only alcohol-dependent males and was characterized by higher alcohol consumption compared with the discovery sample. Similar to the discovery sample, increased craving (measured by the Obsessive Compulsive Drinking Scale) was associated with shorter abstinence. Shorter abstinence was also associated with higher alcohol consumption, increased depressive symptoms (measured by the Beck Depression Inventory) and younger age at onset of alcohol dependence. A final multivariable model of potentially relevant covariates (selected using the same process as in the discovery data set) included Beck Depression Inventory score, number of drinks per drinking day and age of onset of alcoholism; all SNP association analyses were adjusted for these covariates.

**Table 1.** Demographic and clinical measures associated with the length of abstinence during first 3 months of acamprosate treatment in 225 alcoholics in the discovery sample

Parameter	Mean $\pm$ s.d. or N (%)	Association with Relapse Risk			
		Unadjusted		Adjusted for site	
		HR (95% CI)	P-value	HR (95% CI)	P-value
Study site			<b>&lt; 0.0001</b>	—	—
Age at evaluation	44.8 $\pm$ 11.7	0.99 (0.97, 1.01)	0.35	0.99 (0.98, 1.01)	0.51
Gender (male)	148 (65.8%)	0.67 (0.45, 1.02)	0.060	0.72 (0.47, 1.09)	0.12
AA meetings/month	11.8 $\pm$ 12.8	0.96 (0.93, 0.98)	<b>0.0008</b>	0.98 (0.96, 0.99)	<b>0.043</b>
Counseling sessions/month	11.7 $\pm$ 14.4	0.98 (0.96, 1.00)	<b>0.017</b>	0.99 (0.97, 1.01)	0.27
AA sponsor (yes or no)	109 (54.0%)	0.52 (0.33, 0.81)	<b>0.0043</b>	0.62 (0.39, 0.97)	<b>0.036</b>
IDTS-negative score	56.6 $\pm$ 22.4	1.00 (0.99, 1.01)	0.51	0.99 (0.99, 1.01)	0.85
IDTS-positive score	56.3 $\pm$ 25.3	1.00 (0.99, 1.01)	0.41	1.00 (0.99, 1.01)	0.57
IDTS temptation score	48.7 $\pm$ 23.4	1.01 (1.00, 1.02)	0.083	1.00 (0.99, 1.01)	0.52
Baseline PHQ-9	9.4 $\pm$ 6.3	1.04 (1.01, 1.08)	<b>0.0073</b>	1.02 (0.99, 1.06)	0.16
Baseline GAD-7	8.8 $\pm$ 5.9	1.03 (0.99, 1.07)	0.092	1.01 (0.97, 1.04)	0.65
Baseline PACS	13.8 $\pm$ 8.5	1.09 (1.06, 1.12)	<b>&lt; 0.0001</b>	1.06 (1.03, 1.10)	<b>&lt; 0.0001</b>
Maxi drinks per day	16.4 $\pm$ 11.4	0.99 (0.98, 1.01)	0.56	1.00 (0.98, 1.01)	0.70
Average drinks per drinking day	11.3 $\pm$ 7.4	0.99 (0.96, 1.02)	0.53	1.00 (0.97, 1.03)	0.82
Antipsychotic use	6 (2.7%)	2.13 (0.78, 5.80)	0.14	1.03 (0.37, 2.89)	0.96
Benzodiazepine use	35 (15.6%)	0.87 (0.48, 1.56)	0.63	0.93 (0.56, 1.55)	0.71
Mood stabilizer use	7 (3.1%)	1.71 (0.63, 4.66)	0.29	1.45 (0.70, 2.99)	0.22
Antidepressant use	87 (38.7%)	1.02 (0.70, 1.61)	0.78	0.99 (0.79, 1.22)	0.50
Number of days between last drink and acamprosate treatment	21.8 $\pm$ 14.9	0.95 (0.93, 0.97)	<b>&lt; 0.0001</b>	0.96 (0.94, 0.98)	<b>0.0002</b>

Abbreviations: AA, Alcoholics Anonymous; CI, confidence interval; GAD-7, Generalized Anxiety Disorder assessment scale; HR, hazard ratio; IDTS, Inventory of Drug Taking situations; PACS, Penn Alcohol Craving Scale; PHQ-9, nine item depression scale from the patient health questionnaire; TLFB, The Alcohol Timeline Follow-Back.  $P$ -values below 0.05 are marked in bold.



**Figure 1.** Single-nucleotide polymorphism (SNP) association with length of abstinence in the Discovery Sample. Results adjusted for recruitment site, Penn Alcohol Craving Scale and number of days between the last drink and initiation of acamprostate treatment. Dotted line represents the Bonferroni-corrected alpha level of  $0.05/518 = 9.65E - 05$ . Dashed line represents nominal significance of  $P = 0.05$ .

**Table 2.** Candidate SNPs associated with length of abstinence in the discovery sample ( $P < 0.001$ )

SNP	Gene	Minor allele	MAF	Total sample (N = 225)		Males (N = 148)	
				HR	P-value	HR	P-value
rs2058878	GRIN2B	A	0.493	0.49	0.000046	0.41	0.00021
rs17035723	GLRB	A	0.139	2.14	0.00012	2.88	0.000098
rs2160733	GRIN2B	C	0.198	1.78	0.00065	1.85	0.0051
rs2160734	GRIN2B	G	0.488	0.57	0.00079	0.53	0.0061

Abbreviations: HR, hazard ratio; MAF, minor allele frequency. Results are adjusted for recruitment site, Penn Alcohol Craving Scale and number of between last drink and initiation of acamprostate treatment. Results for the entire sample are shown along with results for the male subset

SNP associations with treatment outcomes in the replication sample

As presented in Table 4, the association of the imputed rs2058878 genotypes with abstinence length in the replication sample was marginally significant (hazard ratio=0.72,  $P=0.068$ ). As in the discovery sample, the minor A allele was associated with lower risk of relapse (longer abstinence). To gain additional insights into this association signal, we used the available genome-wide SNP data from the replication sample to search for nearby SNPs that were in high linkage disequilibrium ( $\geq 0.9$ ) with rs2058878. We identified the GRIN2B SNP rs2300272 (minor allele frequency=0.48, linkage disequilibrium = 0.9 with rs2058878), which was genotyped in the replication sample. Analysis of this SNP revealed association of the minor rs2300272 G allele with shorter abstinence in the replication sample (hazard ratio = 1.43,  $P=0.049$ ).

The association of minor alleles of GRIN2B rs2160734 and rs11612353 (proxy for rs2160733) SNPs with abstinence length was in a reverse direction compared with the discovery sample.

**Table 3.** Demographic and clinical measures associated with the length of abstinence during first 3 months of acamprostate treatment in the replication sample of 110 male alcoholics participating in study PREDICT

Parameter	Mean $\pm$ s.d.	HR (95% CI)	P-value
Age at evaluation	45.1 $\pm$ 8.4	0.98 (0.95, 1.01)	0.13
Age of onset	31.7 $\pm$ 10.1	0.95 (0.95, 0.99)	<b>0.036</b>
Average drinks per heavy drinking day	19.2 $\pm$ 10.4	1.03 (1.01, 1.06)	<b>0.0087</b>
Average drinks per drinking day	18.8 $\pm$ 10.6	1.03 (1.01, 1.05)	<b>0.0084</b>
BDI score	6.1 $\pm$ 5.0	1.06 (1.01, 1.11)	<b>0.012</b>
OCDS total score	13.8 $\pm$ 6.6	1.02 (0.98, 1.06)	0.30
Percent drinking days	62.2 $\pm$ 24.3	1.00 (0.99, 1.01)	0.48
Percent heavy drinking days	60.6 $\pm$ 24.6	1.00 (0.99, 1.01)	0.53
Days since last drink	22.0 $\pm$ 4.2	1.03 (0.96, 1.09)	0.43

Abbreviations: BDI, Beck Depression Inventory; CI, confidence interval; HR, hazard ratio; OCDS, Obsessive Compulsive Drinking Scale.

The minor GLRB rs11100096 G allele (proxy for rs17035723 A) showed association with treatment outcome in the same direction as the discovery sample; however, the association was not statistically significant.

## DISCUSSION

Our findings indicate that the minor GRIN2B rs2058878 A allele is associated with longer abstinence during the first 3 months of acamprostate treatment. This association was replicated in an independent sample with marginally significant evidence of association. Furthermore, the replication sample provided significant evidence for association of abstinence length with rs2300272, which is in high linkage disequilibrium with

**Table 4.** Replication analysis using the PREDICT sample ( $N = 110$  acamprosate-treated subjects)

SNP	Gene	Minor allele	MAF	O/I	HR	P-value
rs2058878	<i>GRIN2B</i>	A	0.48	I	0.72	0.068
rs17035723	<i>GLRB</i>	T	0.18	I	0.82	0.48
rs2160733	<i>GRIN2B</i>	C	0.23	I	0.76	0.24
rs2160734	<i>GRIN2B</i>	C	0.47	O	1.31	0.13

Abbreviations: HR, hazard ratio; MAF, minor allele frequency; O/I, indicator of observed or imputed SNP; SNP, single-nucleotide polymorphism. Results are adjusted for the Beck Depression Inventory score, age of onset of alcoholism and the number of drinks per drinking day.

rs2058878. This is the first report showing the association of these genetic markers with the length of abstinence in alcohol-dependent subjects. It provides a compelling reason to investigate these markers as potential pharmacogenetic predictors of acamprosate treatment response.

*GRIN2B* rs2058878 was selected as a tag SNP to investigate the association of treatment outcome with corresponding gene segments. It is possible that other rarer variants located in the gene segment 'tagged' by rs2058878 and rs2300272 are responsible for physiological mechanisms contributing to the reported associations and a search for such functional variants is necessary. It is also possible that rs2058878 or rs2300272 may have a functional role associated with the expression and/or function of *GRIN2B*. Indeed, rs2058878 is located in an intronic region where enrichment of histone 3 lysine 4 mono-methylation and histone 3 lysine 27 acetylation was found in several data sets of the UCSC ENCODE database.<sup>47</sup> These marks indicate the presence of transcriptional enhancers,<sup>37,48</sup> suggesting a role for the region around rs2058878 in the regulation of *GRIN2B* expression. Interestingly, rs2058878 A allele is part of a canonical E-box sequence identified as a binding site for the transcription factors NeuroD and NeuroG in mouse and *Xenopus laevis*.<sup>49</sup> Review of the Human Brain Transcriptome database (<http://hbatlas.org/pages/hbtd>) indicates that expression of the *GRIN2B* gene in human brain areas seems to inversely correlate with expression of the NeuroD and NeuroG transcription factors. These factors are known to have an active role during neural development in humans<sup>50</sup> and to act both as transcriptional activators as well as repressors.<sup>51,52</sup> However, the predicted consensus site is abolished in the rs2058878 T allele (which is associated with shorter abstinence). Thus, altered transcription factor binding to the rs2058878-containing region and consequent changes in *GRIN2B* expression could contribute to the association between rs2058878 and duration of abstinence.

The *GRIN2B* gene encodes the GluN2B subunit of NMDA receptor. Our finding of association between *GRIN2B* variants and the length of abstinence in alcohol-dependent human subjects treated with acamprosate is in line with experimental evidence associating increased GluN2B subunit expression with chronic ethanol treatment and development of physical dependence.<sup>53</sup> Moreover, recent findings indicate that the effects of acamprosate may be related to the  $Ca^{2+}$  ion.<sup>54</sup> Providing that  $Ca^{2+}$  influx in neurons is controlled by NMDA receptor, it is of special interest to investigate the effects of rs2058878 or rs2300272 on  $Ca^{2+}$  influx.

Evidence also indicates that GluN2B-containing NMDA receptors are activated by glycine<sup>19,20</sup> and are primarily responsible for its intracellular effects<sup>56,55</sup> and related behavioral phenotypes, including alcohol intake.<sup>56</sup> Moreover, GluN2B-containing NMDA receptors are functionally important for long-term depression.<sup>20</sup> Consequently, altered production of GluN2B-containing NMDA receptors may disrupt the balance between long-term depression and long-term potentiation, which are fundamental in brain

physiology. Therefore, understanding the effects of rs2058878 and rs2300272 in human alcohol use disorders and animal models of alcohol-related phenotypes is of special interest, especially in the context of experimental evidence suggesting that glycine may have an important role in alcohol dependence and acamprosate response.<sup>21,22,57</sup>

In addition to investigating genetic variant effects, clinical and demographic variations between study subjects and recruitment sites were investigated as potential predictors of treatment outcomes in our study. Several clinical variables (for example, craving intensity, depressive symptoms and the length of sobriety before initiation of acamprosate treatment) and non-pharmacological treatment components (for example, attendance at Alcoholics Anonymous meetings and having an Alcoholics Anonymous sponsor) were found to be associated with abstinence length. The predictive role of these factors should be accounted for in the pharmacogenomic studies and further investigated in prospective clinical trials. Our findings indicate that tools routinely used for assessment of patients treated in community-based programs, including PACS and Timeline Follow-Back, can be useful for these purposes.

Results of this study should be considered in the context of the following limitations. We have not used a placebo arm in the discovery sample, which limits our ability to differentiate association findings related to acamprosate effects from those associated with sobriety independent from acamprosate effects. Prospective placebo controlled studies are needed to validate the role of our top SNPs as potential pharmacogenomic markers of abstinence length associated with or independent from acamprosate effects.

Analyses presented here are limited to the first 3 months of acamprosate treatment and, therefore, may be relevant only during early stages of abstinence. It is possible that genetic and clinical markers associated with early relapse may be different from those associated with later relapse. This possibility will be investigated in the ongoing analyses focused on genetic and clinical predictors of sobriety during the subsequent 3 months of acamprosate treatment in our study cohort.

Finally, both discovery and replication samples were relatively small, which limited power to discover all meaningful associations. However, analyses resulted in potentially important association findings.

In conclusion, our findings indicate that *GRIN2B* rs2058878 and rs2300272 SNPs are associated with abstinence length during the first 3 months of acamprosate treatment. These findings support experimental evidence implicating NMDA receptors in the treatment effects of acamprosate. Future studies should prospectively investigate the potential role of these SNPs as biomarkers of abstinence length in treatment-seeking alcoholics and determine the physiological and molecular mechanisms underlying these association findings.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## ACKNOWLEDGMENTS

We dedicate this manuscript to the late Dr David Mrazek, the founding director of the Samuel C Johnson Genomics of Addiction program at Mayo and an inspiring leader who relentlessly advocated for the advancement of research in genetic, molecular and biological psychiatry to improve clinical care for alcohol use disorders. We would like to thank Drs M Brunner, M Chauhan, G Melnyk, D Onsrud, B Proctor as well as J Wittkopp, Misty Leidal, BSN, Christine Hanzel, BA, Katrina Schaefer, BA, Evan Loehle-Conger, MA, Sharon Schulz, BSN and Joni Burns-Duenes, BSN for their assistance with patient recruitment and data collection and the Mayo Clinic Psychiatry Information Technology group for their assistance with data management. This publication was supported in part by NIH grants: 1P20AA017830 PIs: Mrazek/Choi (VMK, JMB, JRG, DKH-F, LLL, TDS, MKS, MAF and DSC); R01 GM28157 PI: Weinsilbbaum (RMW); U19

GM61388 PI: Weinshilboum (RMW); UL1 TR000135 from the National Center for Advancing Translational Sciences (NCATS) PI: Sundeeep Khosla (VMK, JMB, JRG, DKH-F, LLL, TDS, MKS, MAF and DSC). This publication was supported by Mayo Clinic Center for Individualized Medicine (CIM-2012) Grant, PI: Karpyak (VMK), and SC Johnson Genomics of Addictions Program at Mayo Clinic, PI: Choi (VMK, JMB, JRG, DKH-F, LLL, TDS, MKS, MAF and DSC). Study medication (acamprosate) was provided by Forest Pharmaceuticals, which had no other involvement in project planning, conduct, data analyses or preparation of the manuscript. This work was also supported by grant FKZ 01GS08152 from the National Genome Research Network (NGFNplus – see under www.ngfn-alkohol.de) of the German Federal Ministry of Education and Research (BMBF) and by grants FKZ 01GS0117/NGFN and FKZ EB 01011300. KFM was supported by grant 01EB0410 from the Bundesministerium für Bildung und Forschung. MMN received support from the Alfred Krupp von Bohlen und Halbach-Stiftung and is a member of the DFG-funded excellence cluster ImmunoSensation.

## DISCLAIMER

The contents of this study are solely the responsibility of the authors and do not necessarily represent the official views of the NIH.

## REFERENCES

- Bouchery EE, Harwood HJ, Sacks JJ, Simon CJ, Brewer RD. Economic costs of excessive alcohol consumption in the U.S., 2006. *Am J Prev Med* 2011; **41**: 516–524.
- Collins PY, Patel V, Joestl SS, March D, Insel TR, Daar AS et al. Grand challenges in global mental health. *Nature* 2011; **475**: 27–30.
- Bouza C, Angeles M, Munoz A, Amate JM. Efficacy and safety of naltrexone and acamprosate in the treatment of alcohol dependence: a systematic review. *Addiction* 2004; **99**: 811–828.
- Jonas DE, Amick HR, Feltner C, Bobashev G, Thomas K, Wines R et al. Pharmacotherapy for adults with alcohol use disorders in outpatient settings: a systematic review and meta-analysis. *JAMA* 2014; **311**: 1889–1900.
- Rosner S, Leucht S, Leher P, Soyka M. Acamprosate supports abstinence, naltrexone prevents excessive drinking: evidence from a meta-analysis with unreported outcomes. *J Psychopharmacol* 2008; **22**: 11–23.
- Dawson DA, Goldstein RB, Grant BF. Rates and correlates of relapse among individuals in remission from DSM-IV alcohol dependence: a 3-year follow-up. *Alcohol Clin Exp Res* 2007; **31**: 2036–2045.
- Rosner S, Hackl-Herrwerth A, Leucht S, Leher P, Vecchi S, Soyka M. Acamprosate for alcohol dependence. *Cochrane Database Syst Rev* 2010; CD004332.
- Maisel NC, Blodgett JC, Wilbourne PL, Humphreys K, Finney JW. Meta-analysis of naltrexone and acamprosate for treating alcohol use disorders: when are these medications most helpful? *Addiction* 2013; **108**: 275–293.
- Mason BJ, Goodman AM, Chabac S, Leher P. Effect of oral acamprosate on abstinence in patients with alcohol dependence in a double-blind, placebo-controlled trial: the role of patient motivation. *J Psychiatr Res* 2006; **40**: 383–393.
- Morley KC, Teesson M, Sannibale C, Baillie A, Haber PS. Clinical predictors of outcome from an Australian pharmacological relapse prevention trial. *Alcohol Alcohol* 2010; **45**: 520–526.
- Kranzler HR, Edenberg HJ. Pharmacogenetics of alcohol and alcohol dependence treatment. *Curr Pharm Des* 2010; **16**: 2141–2148.
- Litten RZ, Bradley AM, Moss HB. Alcohol biomarkers in applied settings: recent advances and future research opportunities. *Alcohol Clin Exp Res* 2010; **34**: 955–967.
- Chabenat C, Chretien P, Daoust M, Moore N, Andre D, Lhuintre JP et al. Physicochemical, pharmacological and pharmacokinetic study of a new GABAergic compound, calcium acetylhomotaurinate. *Methods Find Exp Clin Pharmacol* 1988; **10**: 311–317.
- Dahchour A, De Witte P, Bolo N, Nedelec JF, Muzet M, Durbin P et al. Central effects of acamprosate: part 1. Acamprosate blocks the glutamate increase in the nucleus accumbens microdialysate in ethanol withdrawn rats. *Psychiatry Res* 1998; **82**: 107–114.
- Littleton J. Acamprosate in alcohol dependence: how does it work? *Addiction* 1995; **90**: 1179–1188.
- Kiefer F, Witt SH, Frank J, Richter A, Treutlein J, Lemenager T et al. Involvement of the atrial natriuretic peptide transcription factor GATA4 in alcohol dependence, relapse risk and treatment response to acamprosate. *Pharmacogenomics J* 2011; **11**: 368–374.
- Lee MR, Hinton DJ, Wu J, Mishra PK, Port JD, Macura SI et al. Acamprosate reduces ethanol drinking behaviors and alters the metabolite profile in mice lacking ENT1. *Neurosci Lett* 2011; **490**: 90–95.
- Spanagel R, Pendyala G, Abarca C, Zghoul T, Sanchis-Segura C, Magnone MC et al. The clock gene *Per2* influences the glutamatergic system and modulates alcohol consumption. *Nat Med* 2005; **11**: 35–42.
- Bergeron R, Meyer TM, Coyle JTG, Greene RW. Modulation of N-methyl-D-aspartate receptor function by glycine transport. *Proc Natl Acad Sci USA* 1998; **95**: 15730–15734.
- Papouin T, Ladepeche L, Ruel J, Sacchi S, Labasque M, Hanini M et al. Synaptic and extrasynaptic NMDA receptors are gated by different endogenous coagonists. *Cell* 2012; **150**: 633–646.
- Chau P, Stomberg R, Fagerberg A, Soderpalm B, Ericson M. Glycine receptors involved in acamprosate's modulation of accumbal dopamine levels: an *in vivo* microdialysis study. *Alcohol Clin Exp Res* 2010; **34**: 32–38.
- Lido HH, Marston H, Ericson M, Soderpalm B. The glycine reuptake inhibitor Org24598 and acamprosate reduce ethanol intake in the rat; tolerance development to acamprosate but not to Org24598. *Addict Biol* 2012; **17**: 897–907.
- Molander A, Soderpalm B. Accumbal strychnine-sensitive glycine receptors: an access point for ethanol to the brain reward system. *Alcohol Clin Exp Res* 2005; **29**: 27–37.
- Molander A, Soderpalm B. Glycine receptors regulate dopamine release in the rat nucleus accumbens. *Alcohol Clin Exp Res* 2005; **29**: 17–26.
- Vengeliene V, Leonardi-Essmann F, Sommer WH, Marston HM, Spanagel R. Glycine transporter-1 blockade leads to persistently reduced relapse-like alcohol drinking in rats. *Biol Psychiatry* 2010; **68**: 704–711.
- Mann K, Kiefer F, Smolka M, Gann H, Wellek S, Heinz A. Searching for responders to acamprosate and naltrexone in alcoholism treatment: rationale and design of the PREDICT study. *Alcohol Clin Exp Res* 2009; **33**: 674–683.
- Hasin DS, Trautman KD, Miele GM, Samet S, Smith M, Endicott J. Psychiatric Research Interview for Substance and Mental Disorders (PRISM): reliability for substance abusers. *Am J Psychiatry* 1996; **153**: 1195–1201.
- Sobell L, Sobell M. Timeline follow-back. In: Litten R, Allen J (eds). *Measuring Alcohol Consumption*. Humana Press: New York, NY, 1992.
- Flannery BA, Volpicelli JR, Pettinati HM. Psychometric properties of the Penn Alcohol Craving Scale. *Alcohol Clin Exp Res* 1999; **23**: 1289–1295.
- Vander Weg MW, DeBon M, Sherrill-Mittleman D, Klesges RC, Relyea GE. Binge drinking, drinking and driving, and riding with a driver who had been drinking heavily among Air National Guard and Air Force Reserve Personnel. *Mil Med* 2006; **171**: 177–183.
- Annis HM, Martin G. *Inventory of Drug-Taking Situations*. Addiction Research Foundation: Toronto, Canada, 1985.
- Annis HM, Turner NE, Sklar SM. *Inventory of Drug-Taking Situations: User's Guide*. Addiction Research Foundation, Centre for Addiction and Mental Health: Toronto, Canada, 1997.
- Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001; **16**: 606–613.
- Spitzer RL, Kroenke K, Williams JB, Lowe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006; **166**: 1092–1097.
- Alcoholics Anonymous World Services Inc. Alcoholics Anonymous. www.aa.org, 2014. Accessed 2012.
- Cao H, Ren WH, Zhu MY, Zhao ZQ, Zhang YQ. Activation of glycine site and GluN2B subunit of NMDA receptors is necessary for ERK/CREB signaling cascade in rostral anterior cingulate cortex in rats: implications for affective pain. *Neurosci Bull* 2012; **28**: 77–87.
- Heilig M, Egli M. Pharmacological treatment of alcohol dependence: target symptoms and target mechanisms. *Pharmacol Ther* 2006; **111**: 855–876.
- Kovanen L, Saarikoski ST, Haukka J, Pirkola S, Aromaa A, Lonnqvist J et al. Circadian clock gene polymorphisms in alcohol use disorders and alcohol consumption. *Alcohol Alcohol* 2010; **45**: 303–311.
- Lutz UC, Batra A, Kolb W, Machicao F, Maurer S, Kohnke MD. Methylene-tetrahydrofolate reductase C677T-polymorphism and its association with alcohol withdrawal seizure. *Alcohol Clin Exp Res* 2006; **30**: 1966–1971.
- Sicotte H, Rider DN, Poland GA, Dhiman N, Kocher JP. SNPPicker: high quality tag SNP selection across multiple populations. *BMC Bioinformatics* 2011; **12**: 129.
- Kosoy R, Nassir R, Tian C, White PA, Butler LM, Silva G et al. Ancestry informative marker sets for determining continental origin and admixture proportions in common populations in America. *Hum Mutat* 2009; **30**: 69–78.
- Oliphant A, Barker DL, Stuelpnagel JR, Chee MS. BeadArray technology: enabling an accurate, cost-effective approach to high-throughput genotyping. *Biotechniques* 2002; **Suppl**: 56–58: 60–51.
- Treutlein J, Cichon S, Ridinger M, Wodarz N, Soyka M, Zill P et al. Genome-wide association study of alcohol dependence. *Arch Gen Psychiatry* 2009; **66**: 773–784.
- Frank J, Cichon S, Treutlein J, Ridinger M, Mattheisen M, Hoffmann P et al. Genome-wide significant association between alcohol dependence and a variant in the ADH gene cluster. *Addict Biol* 2012; **17**: 171–180.
- Delaneau O, Marchini J, Zagury JF. A linear complexity phasing method for thousands of genomes. *Nat Methods* 2012; **9**: 179–181.

- 46 Howie BN, Donnelly P, Marchini J. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genet* 2009; **5**: e1000529.
- 47 Rosenbloom KR, Sloan CA, Malladi VS, Dreszer TR, Learned K, Kirkup VM *et al*. ENCODE data in the UCSC Genome Browser: year 5 update. *Nucleic Acids Res* 2013; **41**: D56–D63.
- 48 Creyghton MP, Cheng AW, Welstead GG, Kooistra T, Carey BW, Steine EJ *et al*. Histone H3K27ac separates active from poised enhancers and predicts developmental state. *Proc Natl Acad Sci USA* 2010; **107**: 21931–21936.
- 49 Seo S, Lim JW, Yellajoshyula D, Chang LW, Kroll KL. Neurogenin and NeuroD direct transcriptional targets and their regulatory enhancers. *EMBO J* 2007; **26**: 5093–5108.
- 50 Evsen L, Sugahara S, Uchikawa M, Kondoh H, Wu DK. Progression of neurogenesis in the inner ear requires inhibition of Sox2 transcription by neurogenin1 and neurod1. *J Neurosci* 2013; **33**: 3879–3890.
- 51 Itkin-Ansari P, Marcora E, Geron I, Tyrberg B, Demeterco C, Hao E *et al*. NeuroD1 in the endocrine pancreas: localization and dual function as an activator and repressor. *Dev Dyn* 2005; **233**: 946–953.
- 52 Roybon L, Mastracci TL, Ribeiro D, Sussel L, Brundin P, Li JY. GABAergic differentiation induced by Mash1 is compromised by the bHLH proteins Neurogenin2, NeuroD1, and NeuroD2. *Cereb Cortex* 2010; **20**: 1234–1244.
- 53 Narita M, Soma M, Mizoguchi H, Tseng LF, Suzuki T. Implications of the NR2B subunit-containing NMDA receptor localized in mouse limbic forebrain in ethanol dependence. *Eur J Pharmacol* 2000; **401**: 191–195.
- 54 Spanagel R, Vengeliene V, Jandeleit B, Fischer WN, Grindstaff K, Zhang X *et al*. Acamprostate produces its anti-relapse effects via calcium. *Neuropsychopharmacology* 2014; **39**: 783–791.
- 55 Honer M, Benke D, Laube B, Kuhse J, Heckendorn R, Allgeier H *et al*. Differentiation of glycine antagonist sites of N-methyl-D-aspartate receptor subtypes. Preferential interaction of CGP 61594 with NR1/2B receptors. *J Biol Chem* 1998; **273**: 11158–11163.
- 56 Darstein M, Albrecht C, Lopez-Francos L, Knorle R, Holter SM, Spanagel R *et al*. Release and accumulation of neurotransmitters in the rat brain: acute effects of ethanol in vitro and effects of long-term voluntary ethanol intake. *Alcohol Clin Exp Res* 1998; **22**: 704–709.
- 57 Adermark L, Clarke RB, Olsson T, Hansson E, Soderpalm B, Ericson M. Implications for glycine receptors and astrocytes in ethanol-induced elevation of dopamine levels in the nucleus accumbens. *Addict Biol* 2011; **16**: 43–54.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplementary Information accompanies the paper on the Translational Psychiatry website (<http://www.nature.com/tp>)