



Published in final edited form as:

Adv Genet. 2011 ; 74: 175–230. doi:10.1016/B978-0-12-387690-4.00006-4.

Genetics of Circadian Rhythms in Mammalian Model Organisms

Phillip L. Lowrey¹ and Joseph S. Takahashi²

¹Department of Biology, Rider University, Lawrenceville, New Jersey 08648-3099
plowrey@rider.edu

²Department of Neuroscience, Howard Hughes Medical Institute, University of Texas Southwestern Medical Center, Dallas, TX 75390-9111 Joseph.Takahashi@UTSouthwestern.edu

Abstract

The mammalian circadian system is a complex hierarchical temporal network which is organized around an ensemble of uniquely coupled cells comprising the principal circadian pacemaker in the suprachiasmatic nucleus of the hypothalamus. This central pacemaker is entrained each day by the environmental light/dark cycle and transmits synchronizing cues to cell-autonomous oscillators in tissues throughout the body. Within cells of the central pacemaker and the peripheral tissues, the underlying molecular mechanism by which oscillations in gene expression occur involves interconnected feedback loops of transcription and translation. Over the past 10 years we have learned much regarding the genetics of this system, including how it is particularly resilient when challenged by single-gene mutations, how accessory transcriptional loops enhance the robustness of oscillations, how epigenetic mechanisms contribute to the control of circadian gene expression, and how, from coupled neuronal networks, emergent clock properties arise. Here we will explore the genetics of the mammalian circadian system from cell-autonomous molecular oscillations, to interactions among central and peripheral oscillators and ultimately, to the daily rhythms of behavior observed in the animal.

I. INTRODUCTION

The rising and setting of the sun each day causes predictable environmental changes to which most organisms on earth have adapted by evolving endogenous biological timing systems with a period of approximately 24 hours (Young and Kay, 2001). These circadian (~24 hr) clocks anticipate environmental cycles and control daily rhythms in biochemistry, physiology and behavior. Across phyla, all circadian clocks share several fundamental properties: they are synchronized (entrained) each day to external cues, they are self-sustained and produce oscillations that persist in the absence of any external cues, they are temperature compensated such that temperature changes in the physiological range do not alter their endogenous period, and of particular relevance to this review, they are cell-autonomous and genetically-determined. In all of the major model organisms in which circadian rhythms have been studied there has emerged a central organizing principle of the molecular clockwork: within cells a set of clock genes and their protein products together participate in autoregulatory feedback loops of transcription and translation to produce an oscillation with a period of about 24 hr (Lowrey and Takahashi, 2004; Takahashi *et al.*, 2008).

Recent work, however, has prompted a reappraisal of the transcription/translation model as the sole generative mechanism of the molecular circadian oscillator in mammals. For example, it is now clear that oscillations of some mammalian core clock components are dispensable for circadian function (Fan *et al.*, 2007; Liu *et al.*, 2008), and there is some evidence, albeit preliminary, for circadian rhythms in the absence of transcription in some mammalian cells (O'Neill and Reddy, 2011). Perhaps more importantly, however,

limitations of the conventional perturbation analysis methods that helped elucidate the transcription/translation model have become apparent. No longer is it sufficient to knock out a clock gene in a mouse and then assess the consequences on behavior (locomotor activity) or gene expression (changes in RNA and protein levels in cells) alone. We now appreciate that the mammalian circadian clock is a more complex hierarchical system than originally imagined, and thus understanding it requires analysis at many levels.

New technologies and clock models have revealed higher-order genetic properties of the mammalian clock system in which the elimination of one component may be compensated for by other components in ways that are more complex than simple redundancy, and they have demonstrated the important roles of accessory feedback loops and gene networks in conferring stability and robustness on the system (Ueda *et al.*, 2005; Ukai-Tadenuma *et al.*, 2008; Baggs *et al.*, 2009). Further, novel approaches have elucidated the importance of networks of coupled cells from which emergent circadian clock properties arise and even buffer the system against the effects of mutations (Liu *et al.*, 2007b; Abraham *et al.*, 2010; Buhr *et al.*, 2010; Ko *et al.*, 2010). These, and other advances, are making clearer the fundamental properties of each level of organization of the mammalian circadian system from cell-autonomous molecular oscillations, to tissue-specific properties, to the interaction of central and peripheral oscillators, and ultimately to the overt daily rhythms of behavior observed in the animal.

Here we present some of the key findings in the field of mammalian circadian biology over the past 10 years and introduce many of the new technologies that are revolutionizing our understanding of the clock system. Our emphasis will be primarily on work from the principal model organism used to study mammalian biology—the mouse. Indeed, for no other mammalian model is there the extensive repertoire of experimental resources and techniques as for the mouse (Silver, 1995; Nagy *et al.*, 2003; Hedrich and Bullock, 2004; Fox *et al.*, 2007; Adams and van der Weyden, 2008; Blake *et al.*, 2010). We will not, however, explore in depth the intriguing link between the mammalian circadian clock and metabolism, first proposed by McKnight and colleagues a decade ago (Rutter *et al.*, 2002), and now well established, as it is beyond the scope of this review. Instead, we refer the reader to several recent comprehensive treatments of this specific topic (Green *et al.*, 2008; Bass and Takahashi, 2010; Maury *et al.*, 2010; Asher and Schibler, 2011).

II. THE BEGINNING OF MAMMALIAN CLOCK GENETICS

A. Serendipitous discovery of the Syrian hamster *tau* mutant

Before discussing the current state of mammalian clock genetics and the details of the molecular clockwork in mammals, we would first like to reflect back briefly on the period from approximately 1985 to 2000 when the study of mammalian clock genetics began. Indeed, it was in 1985 that Martin Ralph, at the time a graduate student in the laboratory of Michael Menaker (then at the University of Oregon), identified a single outbred Syrian hamster (*Mesocricetus auratus*) with an unusually early onset of locomotor activity. Following transfer from a light/dark (LD) cycle to constant darkness (DD), this animal exhibited an endogenous free-running period of 22 hr compared to 23.5 hr, the shortest previously reported circadian period for this species. Recognizing the implications of possibly discovering the first mammalian circadian mutant, Ralph had the foresight to cross this animal to wild-type hamsters and analyze the behavioral rhythms in the offspring. The free-running periods for the resulting F₁ progeny (1:1 ratio; 22 hr : 24 hr) confirmed that the aberrant phenotype was heritable. Intercrosses produced an F₂ generation with a 1:2:1 Mendelian ratio of 20 hr : 22 hr : 24 hr periods. Thus, this spontaneous mutation designated *tau* (after the circadian symbol for period length), segregated in a semidominant manner and seemed to involve a single autosomal locus (Table 1) (Ralph and Menaker, 1988).

As the first mammalian circadian mutation, *tau* figured prominently in many studies addressing behavioral and physiological aspects of mammalian circadian biology. Perhaps the most important result obtained from the *tau* model was the definitive demonstration through transplantation experiments that the suprachiasmatic nucleus (SCN) of the hypothalamus harbors the central circadian pacemaker in mammals. Adult hamsters rendered behaviorally arrhythmic by SCN lesioning exhibited restored rhythmicity following transplantation of donor SCN tissue into the third ventricle. Not only was host rhythmicity rescued, but the restored rhythms had periods reflecting the genotype of the donor animal (Ralph *et al.*, 1990). Further, when firing rate rhythms of individual SCN neurons on fixed microelectrode plates were recorded, cells from *tau* animals helped show that the circadian period of the whole tissue/animal is determined by averaging widely dispersed periods of individual SCN clock cells. This was the first demonstration that the *tau* mutation affects circadian function in a cell-autonomous manner (Liu *et al.*, 1997b). Two additional seminal findings using the *tau* model include the first report that a diffusible signal can drive circadian rhythms in a mammal (Silver *et al.*, 1996), and that SCN-independent circadian oscillators reside in the mammalian retina (Tosini and Menaker, 1996; Tosini and Menaker, 1998).

Despite the importance to the field of mammalian circadian biology of the *tau* model, from a genetic standpoint it is unfortunate that the mutation occurred in the Syrian hamster rather than in the mouse, a mammalian model for which even in 1985 more comprehensive genetic resources were available. Efforts to develop dense genetic maps and physical mapping reagents for the mouse were well underway at the inception of the Human Genome Project and its inclusion of the mouse as a model genome sequencing project (Muller and Grossniklaus, 2010). The Syrian hamster, however, became one of many “orphan genomes” not included in the publically-funded mapping effort (Jacob, 1996). And so the quest to clone and characterize the *tau* mutation would not come to fruition until twelve years after Ralph and Menaker's 1988 report of the mutant. Our laboratory, using a comparative genomics approach called positional syntenic cloning, demonstrated that the *tau* mutation results from a single nucleotide change in the gene encoding casein kinase I epsilon (*CK1ε*) (Lowrey *et al.*, 2000).

B. Forward genetics and the *Clock* mutation

With the realization that the *tau* mutant hamster, while advantageous for physiological studies, was not immediately a genetically tractable model, our laboratory, in collaboration with William F. Dove, Lawrence H. Pinto, and Fred W. Turek, initiated a dominant circadian behavioral screen of first-generation offspring of male C57BL/6J mice mutagenized with the alkylating agent *N*-ethyl-*N*-nitrosourea (ENU). We adopted this forward genetics approach in mice encouraged by the successful mutagenesis screens in *Drosophila* for circadian defects some 20 years earlier by Seymour Benzer's group (Konopka and Benzer, 1971). Of the 304 animals tested, we recovered a mouse with a single-gene, semidominant mutation, *Clock* (for circadian locomotor output cycles kaput), that significantly lengthened its free-running period (24.8 hr) compared to wild-type C57BL/6J mice (23.3–23.8 hr). Homozygous *Clock* mutant animals exhibited an extremely long initial free-running period in DD of 26–29 hr, followed by a complete loss of rhythmicity after two weeks in DD (Table 1). We mapped the mutation to mouse chromosome 5 (Vitaterna *et al.*, 1994) and subsequently identified the gene through a combination of positional cloning and transgenic rescue of the mutant phenotype (Antoch *et al.*, 1997; King *et al.*, 1997b). Sequence analysis revealed that the *Clock* mutation is caused by an A→T transversion in a splice donor site in the intron between exons 18 and 19 of the gene, resulting in a transcript missing exon 19 (*Clock*^{Δ19}). This deletion disrupts the transactivation domain of the basic helix-loop-helix (bHLH)-PAS (*Period-Arnt-Single-*

minded) transcription factor encoded by *Clock*. Additional genetic approaches revealed that *Clock* is an antimorph—a specific type of dominant negative mutation (King *et al.*, 1997a).

Identification of the *Clock* mutation was proof of principle that, as in *Drosophila*, forward genetic screens for behavioral defects in mice are feasible (Takahashi *et al.*, 1994; Bacon *et al.*, 2004; Clark *et al.*, 2004; Siepka and Takahashi, 2005). Identifying mammalian clock genes by recovering mutants was, however, not the only approach during this “birth” of mammalian clock genetics in the 1990s. Several of what proved to be mammalian core clock genes were cloned by homology to known genes in other organisms, or by identification of paralogs in the same organism. These include three mouse orthologs of the *Drosophila period* gene (*Per1*, *Per2*, and *Per3*), two mouse *Cryptochrome* orthologs (*Cry1* and *Cry2*), and brain and muscle ARNT-like protein 1 (*Bmal1* or *Mop3*), another bHLH-PAS protein, all reviewed comprehensively elsewhere (Lowrey and Takahashi, 2000; Young and Kay, 2001; Reppert and Weaver, 2002; Lowrey and Takahashi, 2004). Indeed, the 1990s witnessed the description of a molecular model of the mammalian core circadian oscillator based on a transcription/translation feedback loop with striking similarity to models proposed for other, phylogenetically divergent organisms, including *Drosophila*, *Arabidopsis*, and *Neurospora* (Dunlap, 1999). This rapid elucidation of the basic mechanism by which mammalian cells keep time was aptly characterized by one colleague as a “clockwork explosion” (Reppert, 1998).

III. OVERVIEW OF THE MAMMALIAN CLOCK SYSTEM

The mammalian circadian system is organized around three major physiological components: an input pathway by which environmental cues (most importantly light) are transmitted to the central or 'master' pacemaker, the central pacemaker itself, and finally a set of output pathways by which the central pacemaker regulates circadian rhythms throughout the body (Takahashi *et al.*, 2001; Quintero *et al.*, 2003; Lowrey and Takahashi, 2004). Light entrainment of the circadian system relies on the eye (Nelson and Zucker, 1981; Foster *et al.*, 1991) where, within the retina, the rods and cones and a recently discovered subset (~1%) of intrinsically-photosensitive retinal ganglion cells (ipRGCs) reside (Do and Yau, 2010). The ipRGCs respond to light stimulation independently of the rod-cone system (Berson *et al.*, 2002; Hattar *et al.*, 2002), and are directly photosensitive owing to their expression of the photopigment melanopsin (Provencio *et al.*, 1998; Gooley *et al.*, 2001; Dacey *et al.*, 2005; Fu *et al.*, 2005; Melyan *et al.*, 2005; Panda *et al.*, 2005; Qiu *et al.*, 2005). Mice lacking either the rod-cone system (Freedman *et al.*, 1999), or melanopsin (*Opn4^{-/-}*) (Panda *et al.*, 2002b; Ruby *et al.*, 2002; Lucas *et al.*, 2003), exhibit normal entrainment to light. Loss of both the rod-cone system and melanopsin, however, render mice unable to entrain to photic stimuli (Hattar *et al.*, 2003; Panda *et al.*, 2003). Because the ipRGCs, via which all retinal input to the SCN is transmitted, receive synaptic input containing non-visual information from the rods and cones, selective destruction of the ipRGCs in mice also prevents circadian photoentrainment (Goz *et al.*, 2008; Hatori *et al.*, 2008).

Photic information received by the retina is transmitted via the retinohypothalamic tract (RHT) which is formed from the axons of the ipRGCs, to the bilaterally-paired suprachiasmatic nuclei of the hypothalamus, the location of the central pacemaker in mammals (Moore and Eichler, 1972; Moore and Lenn, 1972; Stephan and Zucker, 1972; Moore *et al.*, 1995; Gooley *et al.*, 2001). Light stimulation of the ipRGCs causes their axon terminals to release glutamate and pituitary adenylate cyclase-activating polypeptide (PACAP) onto postsynaptic SCN neurons (Ebling, 1996; Hannibal, 2002; Hannibal *et al.*, 2004; Michel *et al.*, 2006; Morin and Allen, 2006). Glutamate-induced calcium influx activates several protein kinase pathways in SCN neurons which ultimately lead to

phosphorylation of Ca²⁺/cAMP-response element binding protein (CREB) (Golombek and Rosenstein, 2010). Within the promoters of many core clock genes, reside Ca²⁺/cAMP-response elements (CREs) to which phospho-CREB homodimers bind to activate transcription (Zhang *et al.*, 2005). Two particularly important CREB clock targets with respect to photic entrainment are the *Per1* and *Per2* genes, both of which contain CREs in their promoters (Travnickova-Bendova *et al.*, 2002) and are rapidly induced in SCN neurons by nocturnal light exposure (Albrecht *et al.*, 1997; Shearman *et al.*, 1997; Shigeyoshi *et al.*, 1997). Circadian rhythms of locomotor activity in mice are phase advanced or phase delayed depending on the time at night during which a photic stimulus occurs (Golombek and Rosenstein, 2010). The resulting increase in PER protein presumably affects the molecular clock in SCN neurons by opposing the action of the positive effectors of the core clock feedback loop discussed later (Yan and Silver, 2004).

Each SCN of the mouse contains approximately 10,000 neurons (Abrahamson and Moore, 2001). When dissociated from SCN tissue (Welsh *et al.*, 1995; Liu *et al.*, 1997b; Herzog *et al.*, 1998; Honma *et al.*, 1998) or when grown as immortalized cells (Earnest *et al.*, 1999), these neurons can independently generate self-sustained circadian rhythms. Intact SCN neurons, however, couple to form a network that expresses synchronized rhythms (Welsh *et al.*, 2010). Ongoing work seeks to clarify the nature of the coupling mechanisms that give rise to the unique SCN network, yet it is clear that neurotransmitters, neuropeptides, gap junctions, and chemical synaptic mechanisms are involved (Welsh *et al.*, 2010). For example, the presence of vasoactive intestinal polypeptide (VIP) and its G-protein coupled receptor, VPAC₂, are important for maintaining circadian rhythmicity of gene expression in SCN cells, and for normal expression of rhythmic behavior in mice. Disrupting the genes for VIP (*Vip*^{-/-}) (Colwell *et al.*, 2003) or its receptor VPAC₂ (*Vipr2*^{-/-}) (Harmar *et al.*, 2002; Cutler *et al.*, 2003), leads to severely compromised circadian rhythms in behavior, neuronal firing, and gene expression, owing to intercellular desynchronization among SCN neurons (Table 1) (Aton *et al.*, 2005; Maywood *et al.*, 2006; Brown *et al.*, 2007; Hughes *et al.*, 2008). In normal mice, circadian rhythms generated within the SCN network are much more robust than those produced by individual neurons (Yamaguchi *et al.*, 2003), even in the presence of clock gene mutations (Liu *et al.*, 2007b). Indeed, recent work has shown that when the autonomous circadian oscillation of individual SCN neurons is eliminated by core clock gene mutation, molecular noise and intercellular coupling are sufficient to elicit stochastic, quasi-circadian oscillations as an emergent property of the SCN network (Ko *et al.*, 2010).

Photic input to the SCN via the RHT is transduced into neural and humoral output signals that synchronize other rhythms in the body, including those of temperature, hormone secretion, and rest/wake (Aston-Jones *et al.*, 2001; Buijs and Kalsbeek, 2001; Brown *et al.*, 2002). Synchronizing signals reach peripheral tissues by both autonomic neural connections (Buijs and Kalsbeek, 2001; Vujovic *et al.*, 2008), and through the release of hormones such as glucocorticoids (Balsalobre *et al.*, 2000; Le Minh *et al.*, 2001). In the absence of the SCN, circadian rhythms in most peripheral tissues damp out after a few days from desynchronization among the cells in the tissue, yet at the single cell level circadian rhythms persist (Balsalobre *et al.*, 1998; Yamazaki *et al.*, 2000; Nagoshi *et al.*, 2004; Welsh *et al.*, 2004; Yoo *et al.*, 2004). The circadian rhythm of locomotor activity commonly monitored in mice and other rodents to determine endogenous circadian period relies on diffusible signals released from the SCN (Silver *et al.*, 1996), several candidate molecules for which have been identified including transforming growth factor α (TGF α) (Kramer *et al.*, 2001; Kramer *et al.*, 2005), cardiotrophin-like cytokine (Kraves and Weitz, 2006), prokineticin 2 (PK2) (Cheng *et al.*, 2002; Li *et al.*, 2006) and potentially others (Hatcher *et al.*, 2008).

It is important to note that there are oscillators in some mammalian brain regions and tissues that, in the absence of the SCN, can drive local physiological rhythms. Two such well-

characterized regions include the retina (Tosini and Menaker, 1996) and the olfactory bulb (Granados-Fuentes *et al.*, 2004a; Granados-Fuentes *et al.*, 2004b; Granados-Fuentes *et al.*, 2006). Further, two extra-SCN pacemakers, the food-entrainable oscillator (FEO) and the methamphetamine-sensitive circadian oscillator (MASCO), can drive circadian behavioral and endocrine rhythms in the absence of the SCN or functional canonical clock genes (Honma *et al.*, 2008; Honma and Honma, 2009; Mohawk *et al.*, 2009; Storch and Weitz, 2009; Pezuk *et al.*, 2010).

IV. THE MAMMALIAN CIRCADIAN MOLECULAR OSCILLATOR

The mammalian circadian molecular oscillator model proposed following the discovery of the core clock genes described earlier encompasses our current understanding of the circadian control of gene expression in cells throughout the body. Core circadian clock genes are genes whose protein products are necessary components for the generation and regulation of circadian rhythms; that is, proteins which form the primary molecular circadian oscillatory mechanism within individual cells. In this model, positive and negative core clock components form a feedback loop with a time constant of about 24 hr per cycle (Figure 1). This loop begins during the day when two bHLH-PAS transcription factors, CLOCK and BMAL1 heterodimerize, translocate to the nucleus, and initiate transcription from genes containing E-box (5'-CACGTG-3') or E'-box (5'-CACGTT-3') *cis*-regulatory elements, including the *Per* and *Cry* genes (King *et al.*, 1997b; Gekakis *et al.*, 1998; Kume *et al.*, 1999; Bunger *et al.*, 2000; Zheng *et al.*, 2001; Yoo *et al.*, 2005). PER and CRY proteins heterodimerize and, along with other proteins such as CK1 ϵ , form a complex in the cytoplasm that translocates to the nucleus where they accumulate and subsequently represses transcription of their own (and other) genes by directly inhibiting CLOCK/BMAL1 (Griffin *et al.*, 1999; Kume *et al.*, 1999; Lee *et al.*, 2001; Sato *et al.*, 2006). Thus, the CLOCK/BMAL1 heterodimer forms the positive, or transactivating component in this loop, while the PER/CRY complex acts as the negative, or transinhibiting component (Figure 1). Following several posttranscriptional and posttranslational steps discussed later, the PER/CRY complex is targeted for degradation via the proteasomal pathway, thereby relieving inhibition such that CLOCK/BMAL1 can initiate a new cycle of transcription. This relatively straightforward feedback loop forms what has become known as the mammalian “core” oscillator mechanism.

The general molecular mechanism just described governs circadian output rhythms in all cells throughout the body, although there are tissue-specific differences. For example, in the forebrain, neuronal PAS domain protein 2 (NPAS2) appears to be the more relevant BMAL1 partner (Reick *et al.*, 2001). Thus, the CLOCK(NPAS2)/BMAL1 complex initiates the rhythmic transcription of clock-controlled genes in tissues throughout the body. Microarray studies have shown that approximately 10–15% of all mammalian transcripts exhibit a circadian oscillation from one cell type/tissue to another (Akhtar *et al.*, 2002; Duffield *et al.*, 2002; Panda *et al.*, 2002a; Storch *et al.*, 2002; Oishi *et al.*, 2003; Miller *et al.*, 2007). These studies, however, may underrepresent the true number of genes under circadian control. Powerful statistical tests are required to identify cycling transcripts from noisy microarray data sets (Doherty and Kay, 2010). Development of new nonparametric statistical algorithms promises to provide more accurate measurements of period, phase and amplitude than traditional analysis methods (Hughes *et al.*, 2010). Hence, continued improvements in data analysis methods should allow the identification of rhythmic transcripts in noisy, low-amplitude data and provide a more precise estimate of the number of genes under circadian control in various mammalian tissues.

Following the discovery of several other bona fide clock genes, it soon became evident that accessory regulatory loops interconnect with the core loop just described and add not only

robustness and stability to the clock mechanism, but also provide additional layers of control and link to a myriad of other pathways in the cell. The first of these accessory loops involves members of the large nuclear receptor family. The mouse *Bmall* promoter contains two cognate RevErbA/ROR-binding elements (ROREs) via which the nuclear receptors ROR α (retinoic acid receptor-related orphan receptor α), ROR β , or ROR γ activate (Sato *et al.*, 2004; Akashi and Takumi, 2005; Guillaumond *et al.*, 2005), and REV-ERB α (reverse orientation c-erbA α) or REV-ERB β repress (Preitner *et al.*, 2002; Triqueneaux *et al.*, 2004; Guillaumond *et al.*, 2005), *Bmall* expression, although the REV-ERBs are likely more important in this process (Figure 1) (Liu *et al.*, 2008). Interestingly, the expression of the aforementioned nuclear receptors is circadian and relies on CLOCK/BMAL1-mediated activation through E-boxes in their promoters, although the *Ror* and *Rev-erb* transcripts cycle antiphase to each other (Yang *et al.*, 2006). The ROR α -mediated activation of *Bmall* transcription is enhanced by peroxisome proliferator-activated receptor- γ (PPAR γ) coactivator 1 α (PGC1 α), the expression of which cycles in liver and muscle (Liu *et al.*, 2007c). Members of the PAR bZIP transcription factor family, including the activators DBP (D-box binding protein), TEF (thyrotroph embryonic factor), HLF (hepatic leukemia factor), and the repressor E4BP4 (E4 promoter-binding protein 4), act via D-box elements in target genes to form a second accessory feedback loop (Mitsui *et al.*, 2001; Ueda *et al.*, 2005; Ohno *et al.*, 2007).

V. BEHAVIORAL, MOLECULAR, AND CELL/TISSUE EFFECTS OF CIRCADIAN CLOCK GENE MUTATIONS

Naturally-occurring, chemically-induced, or targeted mutations exist for all of the core clock genes (Table 1). These mutations have helped define the role of each component of the molecular oscillator (Lowrey and Takahashi, 2004; Ko and Takahashi, 2006; Takahashi *et al.*, 2008). At times, however, results from disruption of clock components have been unexpected.

A. Behavioral and molecular effects

One of these surprises occurred with the generation of a mouse *Clock* knockout model. Interestingly, unlike *Clock* ^{Δ 19} animals, *Clock*^{-/-} mice continue to express circadian rhythms of locomotor activity in DD, albeit with a slightly shorter period compared to wild-type animals, and experience only modest alterations in circadian gene expression in the SCN (Table 1) (DeBruyne *et al.*, 2006). Subsequent work has revealed that in the SCN, NPAS2 can compensate for CLOCK by heterodimerizing with BMAL1 to activate transcription from E-box-containing target genes (DeBruyne *et al.*, 2007a), but that molecular circadian rhythms in peripheral tissues are dependent on the presence of CLOCK (DeBruyne *et al.*, 2007b). *Npas2*^{-/-} animals, however, experience only subtle changes in circadian locomotor activity and gene expression (Dudley *et al.*, 2003; DeBruyne *et al.*, 2007a), suggesting that CLOCK has a more important role in the molecular oscillator. Hence, although there is partial functional redundancy between CLOCK and NPAS2, it is clearly tissue specific (Reick *et al.*, 2001; DeBruyne *et al.*, 2007b). As expected, CLOCK/NPAS2 double knockout animals are completely arrhythmic in DD (DeBruyne *et al.*, 2007a).

Knockout of *Bmall* in mice results in behavioral arrhythmicity in DD and disrupted molecular rhythms of gene expression even though its paralog, BMAL2 (MOP9), is also expressed in the SCN and can form a transcriptionally-competent complex with CLOCK (Table 1) (Bunger *et al.*, 2000; Hogenesch *et al.*, 2000; Dardente *et al.*, 2007). Indeed, *Bmall* is the only core clock gene for which loss of function causes an immediate loss of circadian locomotor behavior in DD. Constitutive expression of *Bmall* in *Bmall*^{-/-} mice or cells restores circadian rhythmicity (McDearmon *et al.*, 2006; Liu *et al.*, 2008), demonstrating that

cycling *Bmal1* mRNA is not necessary for circadian rhythm generation. That BMAL2 cannot rescue the *Bmal1*^{-/-} phenotype most likely results from the dependence of *Bmal2* expression on CLOCK/BMAL1-mediated activation. Hence, disrupting *Bmal1* is likely functionally equivalent to creating a double *Bmal1/Bmal2* null animal (Shi *et al.*, 2010).

Per1 null mutations have been independently generated by three groups (Table 1). The mutant progeny from these lines exhibit subtle differences in circadian behavior. Homozygous *Per1*^{ldc} mice have about a 0.5 hr shorter free-running period in DD than wild-type controls and experience a gradual loss of rhythmicity after two weeks in DD (Bae *et al.*, 2001). *Per1*^{Brdm1} mice express a free-running period approximately 1 hr shorter than wild-type animals and maintain rhythmicity (Zheng *et al.*, 2001). This result is consistent with the circadian behavior of the *Per1*^{-/-} mutant line generated by a third group which exhibits a 0.7 hr shorter free-running period with no loss of rhythmicity (Cermakian *et al.*, 2001). The phenotypic disparities observed among the three *Per1* null studies may result from differences in targeting approaches or genetic backgrounds. Two independent null mutations in *Per2* have been reported (Table 1). Both mutant lines exhibit a loss of behavioral rhythmicity in DD, yet the *Per2*^{Brdm1} line expresses a 1.5 hr shorter period for several days before experiencing arrhythmicity (Zheng *et al.*, 1999). Most animals of the *Per2*^{ldc} line exhibit immediate arrhythmicity upon exposure to DD (Bae *et al.*, 2001). As expected from the results just presented, double *Per1/Per2* knockout animals experience behavioral and molecular arrhythmicity in constant conditions (Bae *et al.*, 2001; Zheng *et al.*, 2001). Because loss of *Per3* has no effect on circadian rhythms either in *Per1/Per3* or *Per2/Per3* double mutant mice, *Per3* is not an essential component of the circadian core clock mechanism (Shearman *et al.*, 2000a; Bae *et al.*, 2001).

Targeted disruption of either of the two *Cry* genes results in opposite effects on circadian behavior—*Cry1*^{-/-} animals have a 1 hr shorter and *Cry2*^{-/-} animals have a 1 hr longer free-running period in DD compared to wild-type animals (Thresher *et al.*, 1998; van der Horst *et al.*, 1999; Vitaterna *et al.*, 1999) (Table 1). Similar to *Per1/Per2* double knockouts, *Cry1/Cry2* double knockout animals experience a complete loss of behavioral and molecular rhythmicity when transferred to DD (Thresher *et al.*, 1998; Okamura *et al.*, 1999; van der Horst *et al.*, 1999; Vitaterna *et al.*, 1999). *Per/Cry* compound knockouts also exhibit interesting behavioral phenotypes (Table 1). *Per1*^{Brdm1}/*Cry1*^{-/-} mice have normal circadian rhythms of behavior and gene expression (Oster *et al.*, 2003). Deletion of *Per1* rescues the short period phenotype observed in *Cry1*^{-/-} mutants, revealing that *Per1* is a nonallelic suppressor of *Cry1*. *Per1*^{Brdm1}/*Cry2*^{-/-} mice have more complex phenotypes. Mutants up to six months of age express behavioral rhythms 1.5 hr longer than wild-type controls and have normal rhythms of *Per2* expression. After approximately 6 months of age, *Per1*^{Brdm1}/*Cry2*^{-/-} animals exhibit disrupted entrainment to LD cycles, and subsequently experience arrhythmicity upon release into DD (Oster *et al.*, 2003). In addition, the older animals have blunted *Per2* rhythms in the SCN revealing an age-sensitive effect in this compound mutant. *Per2*^{Brdm1}/*Cry1*^{-/-} mutants experience immediate behavioral and molecular arrhythmicity in DD (Oster *et al.*, 2002). *Per2*^{Brdm1}/*Cry2*^{-/-} animals, however, maintain behavioral and molecular circadian rhythmicity in DD with a slightly shorter free-running period compared to wild-type controls. Thus, inactivation of *Cry2* in *Per2* null animals restores circadian rhythmicity. As a result, *Cry2* is a nonallelic suppressor of *Per2* (Oster *et al.*, 2002).

The double *Per1/Per2* and *Cry1/Cry2* knockout results make sense given that the PER/CRY complex is necessary to directly inhibit CLOCK/BMAL1-mediated transcription (Griffin *et al.*, 1999; Kume *et al.*, 1999; Shearman *et al.*, 2000b; Lee *et al.*, 2001; Sato *et al.*, 2006). Until recently, transient transfection assays pointed to a more prominent role for CRY in inhibiting CLOCK/BMAL1 (Dardente *et al.*, 2007), yet new evidence suggests that PER may be the more important of the negative effectors as its constitutive expression disrupts

the circadian clock in fibroblasts and hepatocytes. Furthermore, constitutive PER2 expression in the SCN of transgenic mice results in the loss of circadian rhythms of locomotor behavior in a conditional and reversible manner (Chen *et al.*, 2009). Finally, biochemical evidence demonstrates that PER2 directly binds to the CLOCK/BMAL1 complex in a rhythmic way, and that it brings CRY into contact with CLOCK/BMAL1. Rhythmic expression of PER in turn drives the rhythmic inhibition of CLOCK/BMAL1, and it is PER that is the rate-limiting component of the inhibitor complex (Chen *et al.*, 2009). This is substantiated by independent work demonstrating that PER2 is also a more potent inhibitor of CLOCK/BMAL2-mediated transactivation than is CRY (Sasaki *et al.*, 2009).

B. Cell/tissue effects

The analysis of behavioral (locomotor activity) and molecular (RNA/protein) rhythms in mice with mutations in core circadian clock genes just described is insufficient to provide a comprehensive view of molecular clock function. For example, most of the above studies do not take into account differences in central versus peripheral oscillators, potential intercellular interactions in producing the observed phenotypes, or reveal unique properties of individual cellular oscillators. New methods allowing continuous monitoring of circadian rhythms in cultured tissues and individual cells in real time for periods of 20 days or more via bioluminescent technology have revealed many clock properties not evident from behavioral and molecular analyses alone.

By crossing clock gene knockout mice to the *Per2Luciferase* (*Per2Luc*) knockin reporter mouse line in which a luciferase gene is fused to the 3'-end of the endogenous *Per2* gene (Yoo *et al.*, 2004), one group has measured the effects of clock gene perturbations at the level of tissues, populations of cultured cells, and single dissociated cells from both the SCN and peripheral tissues (Liu *et al.*, 2007b). In SCN tissue explants, disruption of *Per1*, *Per3*, *Cry1* or *Cry2* individually has no effect on the maintenance of circadian rhythmicity, and the observed period for each mutant SCN tissue reflects the free-running behavioral period of the corresponding mutant animal model (Table 1 and Table 2). In peripheral tissue explants (e.g., liver, lung, cornea), unlike SCN explants, *Cry1*^{-/-} and *Per1*^{-/-} are required for robust, sustained circadian rhythmicity (Table 2), a property of peripheral tissues not evident from the previously described behavioral and molecular studies (Table 1). *Cry2*^{-/-} and *Per3*^{-/-} mutant peripheral tissues maintain rhythmicity with slightly longer and shorter periods, respectively, compared to wild-type controls, again consistent with behavioral results (Table 1). In dissociated fibroblast cells in culture, *Per1*, *Per2* and *Cry1* are required to maintain circadian oscillations. Thus it seems that, in fibroblast cultures at least, *Per1* and *Per2* are not functionally redundant (Liu *et al.*, 2007b).

When single fibroblast cells are imaged for circadian rhythms of bioluminescence, again *Per1* and *Cry1* prove necessary to sustain circadian oscillations, confirming the results observed in fibroblast cultures (Liu *et al.*, 2007b). Single *Cry2*^{-/-} fibroblast cells are rhythmic with a slightly longer period compared to individual wild-type cells, consistent with the behavioral phenotype of *Cry2* null mice (Table 1 and Table 2). To measure rhythms of bioluminescence from single SCN neurons, they must first be uncoupled by mechanical dissociation into single cells (Welsh *et al.*, 1995; Herzog *et al.*, 1998). Similar to the result in single fibroblasts, single *Cry2*^{-/-} SCN neurons are rhythmic with a period longer than wild-type SCN neurons (Table 2). Single *Cry1*^{-/-} and *Per1*^{-/-} SCN neurons, in contrast to single *Cry2*^{-/-} SCN neurons or *Cry1*^{-/-} and *Per1*^{-/-} SCN tissue explants, exhibit arrhythmicity (Table 2). This is an important result not apparent in earlier behavioral studies of *Per1* and *Cry1* null mutants and demonstrates that the robustness of circadian oscillations observed in *Cry1*^{-/-} and *Per1*^{-/-} SCN explant tissue is not a cell-autonomous property. Instead, the ability of *Cry1*^{-/-} and *Per1*^{-/-} SCN tissue to maintain circadian rhythmicity despite mutations in core clock components that, at the single-cell level cause arrhythmicity,

depends on intercellular coupling among SCN neurons (Liu *et al.*, 2007b). This property of intercellular coupling, and not unique intracellular molecular mechanisms, is what distinguishes SCN neurons from cells of peripheral tissues.

As mentioned previously, *Bmal1* null mutant animals experience an immediate loss of circadian behavior upon transfer to constant conditions (Bunger *et al.*, 2000). Using bioluminescence monitoring methods with *Bmal1*^{-/-} SCN and peripheral tissues similar to those just described, a recent study has elucidated interesting properties of the SCN network not discovered in previous behavioral and molecular investigations (Ko *et al.*, 2010). SCN explants from *Bmal1*^{-/-} mice crossed to the *Per2*^{Luc} reporter line exhibit rhythmic but highly variable (noisy) oscillations of PER2::LUC bioluminescence for more than 35 days in culture (Table 2). This is an unexpected result given that the behavioral and molecular rhythms of gene expression in *Bmal1*^{-/-} animals are arrhythmic (Table 1). The authors of this study refer to these quasi-circadian rhythms generated by the *Bmal1*^{-/-} SCN explants as stochastic. As expected, bioluminescence monitoring demonstrates that all peripheral tissues from *Bmal1*^{-/-} animals are arrhythmic (Table 2). Analysis of single dispersed *Bmal1*^{-/-} SCN neurons, however, reveals that they are arrhythmic and do not exhibit the stochastic rhythms of bioluminescence observed in *Bmal1*^{-/-} SCN explants. This result seems not to depend from what subtype of SCN neuron recordings are made or from what region of the SCN the neurons are obtained—all dispersed *Bmal1*^{-/-} SCN neurons exhibit arrhythmicity. Further, *Bmal1*^{-/-} SCN slices treated with tetrodotoxin (TTX) experience an immediate cessation of stochastic rhythmicity, a result of loss of rhythmicity at the single-cell level. Upon removal of TTX from SCN slices, stochastic rhythmicity is restored thereby confirming the importance of intercellular coupling in generating the observed PER2::LUC rhythms in the SCN slices. Taken together, these bioluminescence results demonstrate that the quasi-circadian rhythms in *Bmal1*^{-/-} SCN explants is not a cell-autonomous property, but rather an emergent rhythmic property of the SCN intercellular network (Ko *et al.*, 2010).

VI. POSTTRANSLATIONAL MODIFICATION OF CLOCK PROTEINS

Posttranslational modifications of the core clock components play a crucial role in generating the delays necessary to establish the ~24 hr rhythm of the mammalian circadian clock. Some of these modifications are absolutely essential to clock function while others simply fine-tune the rhythm. Phosphorylation of clock proteins was the first posttranslational process observed in the mammalian molecular clock, and we understand more about this mechanism than any other. The list of identified posttranslational modifications of mammalian clock proteins has grown rapidly and now, in addition to phosphorylation, includes dephosphorylation, ubiquitination, sumoylation, and acetylation.

A. Phosphorylation

1. Casein kinase 1 (CK1)—Posttranslational modification as an important clock-related process in higher eukaryotes became apparent with the identification in *Drosophila* of *doubletime*, a gene encoding a fly casein kinase 1 ortholog that phosphorylates PER (Kloss *et al.*, 1998; Price *et al.*, 1998). Subsequently, we identified casein kinase 1 epsilon (*CK1ε*) as the gene affected by the Syrian hamster *tau* mutation, and showed that in vitro CK1ε^{tau} is hypomorphic toward various substrates, including mammalian PER1 and PER2 proteins (Lowrey *et al.*, 2000). Others demonstrated that, in vitro and in cultured cells, CK1ε and the closely-related family member, CK1δ, can phosphorylate PER (Keesler *et al.*, 2000; Vielhaber *et al.*, 2000; Camacho *et al.*, 2001; Akashi *et al.*, 2002). Further work revealed that CK1δ/ε-mediated phosphorylation regulates PER subcellular localization and its ability to repress CLOCK/BMAL1-mediated transcription, and promotes its ubiquitin-degradation via the 26S proteasome (Vielhaber *et al.*, 2000; Eide *et al.*, 2005b; Shirogane *et al.*, 2005; Vanselow *et al.*, 2006; Ohsaki *et al.*, 2008).

Although our work showed that $CK1\epsilon^{tau}$ was a hypomorph in in vitro assays, a study using mathematical modeling and in vivo analyses has reported that the $CK1\epsilon^{tau}$ anion binding site mutation causes loss of enzyme function toward canonical acidic CK1 ϵ substrates, but gain of function toward the non-canonical β -transducin repeat-containing protein (β TrCP) binding site on PER2 (Gallego *et al.*, 2006a). Indeed, biochemical evidence substantiates the model's prediction as do behavioral, neurophysiological and cellular studies from a mouse model of the hamster *tau* mutation (Gallego *et al.*, 2006a; Meng *et al.*, 2008). Another group, however, has published findings that contradict this interpretation. They report that $CK1\epsilon^{tau}$ is actually a partial loss of function mutation as the mutant kinase is unable to phosphorylate sites that promote nuclear localization of PER2, but that it can phosphorylate amino acids required for PER2 proteasomal degradation (Vanselow *et al.*, 2006). By mapping all of the CK1 ϵ phosphorylation sites on PER2, they opine that the different sites differentially target PER2 to two cellular locations—nucleus or proteasome. Both interpretations agree, however, that the *tau* allele is a particularly interesting mutation biochemically as it differentially affects CK1 ϵ activity toward specific substrates.

The *tau* mutation focused much attention on the role of CK1 ϵ in the mammalian molecular clock, yet as mentioned above, CK1 δ also phosphorylates PER proteins and targets them for degradation (Camacho *et al.*, 2001; Xu *et al.*, 2005), and both kinases associate with PER/CRY repressor complexes in vivo (Figure 1) (Lee *et al.*, 2001). Thus, to better define the role of these two CK1 enzymes, null mutants of both have been generated independently by different laboratories. The free-running period of the locomotor activity rhythm of $CK1\epsilon^{-/-}$ mice is slightly, but significantly, longer than wild-type controls (Table 1) (Meng *et al.*, 2008; Etchegaray *et al.*, 2009). Two groups have reported that $CK1\delta^{-/-}$ mice die during the perinatal period, thus the free-running behavioral period has been studied in $CK1\delta^{+/-}$ heterozygous animals. In one case, one copy of the $CK1\delta$ null allele results in no difference in free-running period compared to controls (Xu *et al.*, 2005), while another group's heterozygous null animal exhibits a slight increase in circadian period (Etchegaray *et al.*, 2009). In addition, compared to CK1 ϵ -deficient tissue, mouse embryonic fibroblasts (MEFs) and liver tissue deficient in $CK1\delta$ have about a one to two hour longer circadian period in vitro (Etchegaray *et al.*, 2009; Lee *et al.*, 2009). When monitored from neonatal SCN explants, PER2::LUC bioluminescence rhythms from $CK1\delta^{-/-}$ mice are longer compared to wild-type controls, yet there is no significant difference in PER2::LUC rhythms from $CK1\epsilon^{-/-}$ SCN compared to controls (Etchegaray *et al.*, 2010).

Pharmacological approaches have also been used to study the roles of CK1 ϵ and CK1 δ in the mammalian clock with the general CK1 δ/ϵ inhibitors CKI-7, IC261 and D4476, all of which lengthen circadian period in cultured cells (Eide *et al.*, 2005a; Vanselow *et al.*, 2006; Reischl *et al.*, 2007; Hirota *et al.*, 2008). An inhibitor specific for CK1 ϵ (PF-4800567) has only a slight effect on the period of oscillating rat-1 fibroblasts stably transfected with a *Per2::luc* reporter compared to the dual CK1 δ/ϵ inhibitor (PF-670462) which causes an increase in fibroblast circadian period (Walton *et al.*, 2009). Single injections into rats of the dual CK1 δ/ϵ inhibitor induce large phase delays in circadian locomotor rhythms under free-running and entrained conditions (Badura *et al.*, 2007; Sprouse *et al.*, 2010). Daily treatment with PF-670462 significantly lengthens locomotor behavioral rhythms in a dose-dependent manner in wild-type, $CK1\epsilon^{tau}$, and $CK1\epsilon^{-/-}$ mice (Meng *et al.*, 2010). Selective inhibition of CK1 ϵ with PF-4800567 has no significant effect on behavioral rhythms in wild-type or $CK1\epsilon^{-/-}$ mice, yet it lengthens the free-running locomotor activity rhythm of $CK1\epsilon^{tau}$ animals. How does inhibition of the CK1 enzymes affect molecular clock function? PF-670462 seems to work by stabilizing PER2 nuclear localization in SCN neurons and peripheral tissues. This prolongs PER2-mediated negative feedback, thereby lengthening circadian period (Meng *et al.*, 2010). Together with the *CK1* knockout experiments, these results suggest that CK1 δ has a more prominent role compared to CK1 ϵ in the mammalian

clockwork; that is, the two kinases seem not to be equally redundant. It is clear, however, that loss of both CK1 ϵ and CK1 δ causes arrhythmicity in MEFs (Lee *et al.*, 2009), and that knockdown of both kinases in human U2OS (osteosarcoma) cells additively lengthens circadian period to more than 30 hr (Isojima *et al.*, 2009). It remains to be determined if this partial functional redundancy derives from unequal expression levels of the two kinases in cells throughout the body. Some evidence suggests that, at least in MEFs, CK1 δ is twice as abundant as CK1 ϵ (Lee *et al.*, 2009).

One group has reported a surprising result in their work with CK1 δ/ϵ in which a novel, non-catalytic clock-related role for these kinases is revealed (Lee *et al.*, 2009). Overexpression of the CK1 δ/ϵ -binding domain of PER2 (CKBD-P2) in MEFs severely disrupts PER2::LUC rhythms of bioluminescence, presumably because the CKBD-P2 fragment interferes (competes) with the interaction between CK1 δ/ϵ and PER. In addition, PER1 and PER2::LUC levels are lower in the CKBD-P2-expressing MEFs, while CK1 δ/ϵ levels are higher than normal. This suggests that the CKBD-P2 enhances the stability of CK1 δ/ϵ via physical interaction. Furthermore, the low levels of PER observed in the CKBD-P2-expressing MEFs may result from the inhibition of CK1 δ/ϵ -specific phosphorylation of PER, and from the reduced physical interaction between CK1 δ/ϵ and PER which, under normal circumstances, may confer stability to PER (Lee *et al.*, 2009). Overexpression of the CKBD region of PER3 in MEFs does not have an effect on the circadian rhythms in these cells, mirroring previous reports showing that PER3 does not interact with CK1 δ/ϵ . That the stabilizing role of CK1 δ/ϵ toward PER is non-catalytic is supported by experiments in which dominant-negative CK1 δ/ϵ (K38R)-expressing MEFs do not experience reduced PER levels (Lee *et al.*, 2009). Interestingly, a non-catalytic circadian role has also been reported recently for the *Drosophila* CK1 family member, DBT (Yu *et al.*, 2009).

Finally, in a high-throughput screen of approximately 120,000 compounds using U2OS cells expressing luciferase under the control of the mouse *Bmal1* promoter, another CK1 family member, CK1 α , has been identified as a mammalian clock regulatory kinase (Hirota *et al.*, 2010). A purine derivative identified in this screen, longdaysin, inhibits CK1 δ , CK1 α , and ERK2 (MAPK1) and prevents them from phosphorylating PER1, causing a dramatic 13 hr lengthening of period in U2OS cells. Knockdown by siRNA of one of the aforementioned kinases alone is insufficient to recapitulate the 13 hr period lengthening effect, yet combinatorial knockdown of all three kinases additively increases period and closely mirrors the effect of longdaysin treatment (Hirota *et al.*, 2010). Results from this interesting study suggest that multiple kinases participate in a network to enhance robustness of the molecular clock mechanism.

2. Glycogen synthase kinase-3 β (GSK-3 β)—Glycogen synthase kinase-3 (GSK-3) is a serine-threonine, phosphate-directed protein kinase of which there are two isoforms in mammals: GSK-3 α and GSK-3 β (Ali *et al.*, 2001). GSK-3 was initially characterized as a kinase involved in metabolism and energy storage, yet it has since been shown to play a role in many intracellular pathways (Doble and Woodgett, 2003). Knockout models for both isoforms have been generated, but *Gsk-3 β ^{-/-}* mice experience embryonic lethality (Hoeflich *et al.*, 2000; MacAulay *et al.*, 2007). Interestingly, GSK-3 is sensitive to lithium. Presumably, Li⁺ competes directly for binding to GSK-3 with Mg²⁺, a required cofactor for GSK-3 function (Klein and Melton, 1996; Stambolic *et al.*, 1996; Ryves and Harwood, 2001). Several studies have documented the effects of lithium treatment on circadian rhythms in mammals, including a consistent effect of lengthening the free-running period of behavioral rhythms, notably those of locomotor activity and drinking (Seggie *et al.*, 1982; LeSauter and Silver, 1993; Iwahana *et al.*, 2004). This period-lengthening effect of lithium is also observed for the firing rate rhythms in isolated mouse SCN neurons (Abe *et al.*, 2000). Other work has shown that GSK-3 β is rhythmically expressed in the SCN and liver of mice,

and that it undergoes a daily cycle in phosphorylation in vivo as well as in serum-shocked NIH3T3 mouse fibroblasts (Harada *et al.*, 2005; Iitaka *et al.*, 2005). Lithium chloride treatment phase delays, while overexpression of GSK-3 β , phase advances clock gene expression in fibroblasts (Iitaka *et al.*, 2005). In addition, *Gsk-3 α* RNAi knockdown in *Gsk-3 β ^{-/-}* MEFs, induces a phase delay in the *Per2* RNA rhythm, as does treatment of MEFs with kenpaullone, a GSK-3 antagonist (Kaladchibachi *et al.*, 2007). Surprisingly, however, a recent high-throughput screen has identified small molecule inhibitors of GSK-3 β that shorten circadian period, a result confirmed by siRNA knockdown of GSK-3 β (Hirota *et al.*, 2008). These effects of inhibition of GSK-3 β on the molecular clock in cells and on clock-controlled behavior in mammals have prompted further investigation into the potential clock-related targets of this enzyme. In addition, work in *Drosophila* has shown that *shaggy*, the fly ortholog of mammalian GSK-3 β , is an important component in determining circadian period length in that organism (Martinek *et al.*, 2001).

Mammalian targets of GSK-3 β phosphorylation include the positive and negative components of the core circadian transcriptional/translational feedback loop, CLOCK/BMAL1 and PER/CRY, respectively. Phosphorylation by GSK-3 β of PER2 promotes its nuclear localization (Iitaka *et al.*, 2005). GSK-3 β -mediated phosphorylation at Ser553 of CRY2 leads to its degradation by the proteasome (Harada *et al.*, 2005), yet GSK-3 β phosphorylation promotes the stabilization of REV-ERB α (Yin *et al.*, 2006). Lithium treatment of cultured cells results in the rapid proteasomal degradation of REV-ERB α and concomitant derepression of *Bmal1* transcription owing to inhibition of GSK-3 β (Yin *et al.*, 2006). Others have identified a phospho-degron region on the CLOCK protein that is phosphorylated by GSK-3 β , and which promotes degradation of CLOCK (Spengler *et al.*, 2009). Finally, BMAL1, the major CLOCK dimerization partner, is phosphorylated by GSK-3 β on Ser17 and Thr21. This modification targets BMAL1 for ubiquitination and subsequent degradation by the proteasome (Sahar *et al.*, 2010).

3. Casein kinase 2 (CK2)—Casein kinase 2, a serine/threonine kinase, is a tetramer composed of two α catalytic and two β regulatory subunits, and, similar to CK1 prefers acidic substrates (Meggio and Pinna, 2003). A role for this kinase in the regulation of circadian rhythms was first reported in *Arabidopsis*, *Drosophila*, and *Neurospora* (Gallego and Virshup, 2007). Recently, participation of CK2 in the mammalian molecular clock mechanism has been reported by three groups. First, BMAL1 is a substrate for CK2 α at Ser90, a residue that undergoes rhythmic phosphorylation (Tamaru *et al.*, 2009). Nuclear localization of BMAL1 seems to depend on CK2 α phosphorylation, as knockdown of CK2 α results in cytoplasmic BMAL1 accumulation and a concomitant disruption of *Per2* mRNA rhythms (Tamaru *et al.*, 2009).

PER2 is also a CK2 α substrate. One study has shown that CK2 α both phosphorylates PER2 at Ser53 and enhances CK1 ϵ -mediated PER2 destabilization (Tsuchiya *et al.*, 2009). These two actions of CK2 α on PER2 are independent, as the CK2 α -mediated potentiation of CK1 ϵ -dependent degradation of PER2 does not require phosphorylation of Ser53. Although it has not been demonstrated, it may be that CK2 α acts to phosphorylate CK1 ϵ thereby upregulating its activity toward PER2, given that a catalytically inactive form of CK2 α (CK2 α -K68A) fails to enhance CK1 ϵ -dependent PER2 degradation (Tsuchiya *et al.*, 2009). Finally, in a recent RNAi screen for clock-related components, downregulation of either CK2 α or CK2 β lengthens circadian period, while knockdown of both subunits leads to arrhythmicity (Maier *et al.*, 2009). Furthermore, overexpression of CK2 α has a period-shortening effect. This same study showed that CK2 α binds PER2 and promotes phosphorylation near the N-terminus, although the proposed sites do not correspond to Ser53 mentioned before. Also, unlike the previous study, CK2 α phosphorylation is proposed to stabilize PER2 rather than enhance its degradation. Inhibition of CK2 α was

shown to delay PER2 nuclear accumulation, suggesting that CK2 α phosphorylation of PER2 may provide a signal for nuclear entry (Maier *et al.*, 2009). These, and other questions, need to be addressed to better understand the role of CK2 in the mammalian clock mechanism.

4. Other kinases—BMAL1 has been identified as a substrate for several kinases including CK1 ϵ and GSK-3 β as described above. In addition, mitogen-activated protein kinase (MAPK) phosphorylates BMAL1 on several residues, including Thr534 which negatively regulates the transactivation potential of the CLOCK/BMAL1 complex at E-boxes (Sanada *et al.*, 2002). Another study has shown recently that RACK1 (receptor for activated C kinase-1) recruits activated protein kinase C α (PKC α) to the CLOCK/BMAL1 complex during the negative feedback phase of the circadian cycle in the nucleus where it phosphorylates BMAL1 and suppresses CLOCK/BMAL1 transcriptional activity (Robles *et al.*, 2010). Knockdown of either RACK1 or PKC α in fibroblast cultures shortens circadian period. *Prkca*^{-/-} mice have a normal circadian period, yet they experience impaired photic resetting of behavioral rhythms (Jakubcakova *et al.*, 2007). This difference between the knockout phenotype and cell culture results highlights the need for further work.

CLOCK has been shown to be phosphorylated throughout the circadian cycle, yet it is hyperphosphorylated during the negative feedback phase (Yoshitane *et al.*, 2009). Specific residues have been identified as phosphorylation sites on CLOCK, including Ser38, Ser42, and Ser427. The kinase or kinases responsible for phosphorylating these sites, however, remain undetermined. CLOCK Δ 19, which lacks the binding site for CLOCK-interacting protein circadian (CIPC), a PER/CRY-independent negative regulator of CLOCK/BMAL1-mediated transcription, is less phosphorylated and more stable than wild-type CLOCK (Zhao *et al.*, 2007). Others have shown that CLOCK is a substrate for phosphorylation by protein kinase G II (PKG-II) and by two protein kinase C isoforms (PKC α and PKC γ) (Tischkau *et al.*, 2004; Shim *et al.*, 2007).

Finally, it will be important to assess the contribution of sequential phosphorylation events on clock proteins. For example, some kinases such as CK1 δ/ϵ require priming phosphorylation of their target sites (Knippschild *et al.*, 2005). Phosphorylation of human PER2 at the Ser662 familial advanced sleep phase syndrome (FASPS) site occurs by an unidentified kinase and is necessary for subsequent CK1 δ/ϵ phosphorylation (Xu *et al.*, 2007). Similarly, GSK-3 β phosphorylation of CRY2 requires priming phosphorylation by DYRK1A (dual-specificity tyrosine-phosphorylated and regulated kinase 1A) (Kurabayashi *et al.*, 2010). In many cases, the priming kinases for these so-called “phosphate directed” kinases such as CK1 δ/ϵ and GSK-3 remain to be identified for particular circadian substrates.

B. Dephosphorylation

Protein phosphatases, although fewer in number in the mammalian genome relative to kinases, also play an important role in the molecular clock (Gallego and Virshup, 2007). Reversible phosphorylation of clock-relevant substrates presumably confers on clock the flexibility to respond appropriately to various stimuli. Few studies, however, have explored the role of phosphatases on the mammalian clock mechanism. The major clock kinases, CK1 ϵ and CK1 δ , both undergo autophosphorylation which downregulates their activity. At least eight autophosphorylation sites must be dephosphorylated by phosphatases to activate CK1 δ/ϵ (Gietzen and Virshup, 1999). Additional evidence suggests that CK1 δ/ϵ participates in a futile autophosphorylation/dephosphorylation cycle in vivo which acts to regulate kinase activity (Rivers *et al.*, 1998). One group has shown that it is protein phosphatase 5 (PP5) that dephosphorylates CK1 δ/ϵ . Furthermore, the same study demonstrated that the

CRY proteins interact with and inhibit noncompetitively, PP5. As a result, the CRY proteins seem to indirectly regulate the activity of CK1 δ/ϵ by inhibiting the phosphatase activity of PP5 (Partch *et al.*, 2006). Another study has found a role for PP1 in the clock mechanism (Gallego *et al.*, 2006b). PP1 can bind to and dephosphorylate CK1 δ/ϵ -phosphorylated PER2 thereby negatively regulating its degradation by the proteasome. Overexpression of a dominant negative form of PP1, or use of PP1 inhibitors, results in accelerated degradation of PER2 (Gallego *et al.*, 2006b).

C. Ubiquitination

One of the major pathways for protein degradation in cells is the ubiquitin-dependent proteasomal mechanism. This system requires the attachment of multiple ubiquitin molecules to lysine residues on the target protein (Ciechanover *et al.*, 2000; Nandi *et al.*, 2006). Polyubiquitinated proteins are directed to the 26S proteasome, a large multicatalytic protease, where they are then degraded to small peptides (Nandi *et al.*, 2006). Attachment of ubiquitin to a protein is a three-step process. First, ubiquitin is adenylated by an activating enzyme (E1), then transferred to a conjugating enzyme (E2), and finally linked to the target protein by a ligase (E3) (Wilkinson, 1999; Ciechanover *et al.*, 2000). The specificity of this system is determined by the E3 ligases which can be categorized into at least six subtypes. With respect to the mammalian clock mechanism, the SCF E3 subtype is the most relevant. SCF complexes are multimers and are named for their constituent protein components, Skp1, Cdc53 or Cullin, and any one of a number of proteins containing an F-box motif, each of which recognizes a particular target protein (Nandi *et al.*, 2006). It is the F-box protein that confers specificity to each SCF complex (Cardozo and Pagano, 2004). The SCF complexes are constitutively active enzymes that recognize and ubiquitinate only phosphorylated substrates (Kornitzer and Ciechanover, 2000). Consequently, this system links protein phosphorylation to proteolytic degradation by the 26S proteasome (Cardozo and Pagano, 2004).

Turnover of the PER/CRY repressor complex is an important event in relieving inhibition of CLOCK/BMAL1 such that a new circadian day can begin. Insight into how this repressor complex is cleared has come from several studies. Recently, our laboratory and another independently recovered long period behavioral mutants, *Overtime* (Siepka *et al.*, 2007) and *After-hours* (Godinho *et al.*, 2007), respectively, through ENU mutagenesis screens of mice. Both mutations affect the same gene, *Fbxl3*, which encodes the F-box and leucine-rich repeat protein 3 (FBXL3). FBXL3 is involved in ubiquitination of the CRY proteins, which targets them for degradation (Busino *et al.*, 2007). The proteasomal-mediated degradation of CRY1 and CRY2, however, appears to be differentially regulated. CRY1 is phosphorylated by AMPK (AMP-activated protein kinase) which promotes its FBXL3-dependent degradation (Figure 1) (Lamia *et al.*, 2009). CRY2, however, is phosphorylated by GSK-3 β on Ser553 (Harada *et al.*, 2005), which first requires priming phosphorylation by DYRK1A on Ser557, both of which are at the C-terminal tail of CRY2 (Kurabayashi *et al.*, 2010). This Ser557/Ser553 phosphorylation mechanism promotes proteasomal degradation of CRY2 by what is likely an FBXL3-independent mechanism. Indeed, it seems that Ser557/Ser553 phosphorylation and subsequent turnover of CRY2 slows its cytosolic accumulation rate and allows its timely nuclear translocation (Kurabayashi *et al.*, 2010). The FBXL3-dependent degradation mechanism acts during the declining phase of negative feedback when CRY2 nuclear clearance occurs (Godinho *et al.*, 2007). Taken together, these two mechanisms of CRY2 degradation suggest a model by which negative feedback is controlled.

As discussed earlier, phosphorylation of the PER proteins by CK1 δ/ϵ promotes their polyubiquitination (Eide *et al.*, 2005a; Shirogane *et al.*, 2005; Reischl *et al.*, 2007; Ohsaki *et al.*, 2008). Ubiquitination of the PERs is mediated by the F-box proteins β TrCP1 and/or β TrCP2 (Figure 1) (Vielhaber *et al.*, 2000; Eide *et al.*, 2005b; Shirogane *et al.*, 2005;

Vanselow *et al.*, 2006; Ohsaki *et al.*, 2008). Following ubiquitination, the PER proteins are degraded by the 26S proteasome (Gallego and Virshup, 2007).

D. Sumoylation

The BMAL1 protein undergoes extensive posttranslational modification in cells, including phosphorylation, acetylation, ubiquitination, and sumoylation. Small ubiquitin-like modifier (SUMO) proteins are covalently attached at lysine residues of target proteins to modify their function (Wilkinson and Henley, 2010). Sumoylation has been shown to affect nuclear/cytosolic localization of proteins, progression through the cell cycle, protein stability, and transcriptional regulation (Gareau and Lima, 2010). In mice and humans, there are three *Sumo* paralogs (*Sumo1–3*). SUMO2 and SUMO3 are 95% identical and are often referred to as SUMO2/3. Two groups have independently demonstrated that BMAL1 is rhythmically polysumoylated at a conserved lysine residue (K259) in the PAS domain linker region by all three SUMO proteins, and that this process is dependent on CLOCK, the BMAL1 dimerization partner (Cardone *et al.*, 2005; Lee *et al.*, 2008). Furthermore, sumoylation localizes BMAL1 to the promyelocytic leukemia nuclear body, potentiates CLOCK/BMAL1 transactivation of clock-controlled genes, and promotes BMAL1 ubiquitin-dependent proteasomal degradation (Lee *et al.*, 2008).

E. Acetylation, deacetylation and chromatin remodeling

Rhythmic changes in chromatin architecture participate in the activation and repression of transcription via posttranslational modifications at histone N-terminal tail regions (Imhof and Becker, 2001). Acetylation of lysine residues or phosphorylation of serine residues in histone tails induces a transcription-permissive nucleosome conformation, whereas deacetylation, dephosphorylation, or methylation of histone lysine residues promotes a transcription-inhibitory nucleosome conformation. These, and other covalent modifications that occur at histones to alter the degree of chromatin condensation, have become known as the “histone code” (Strahl and Allis, 2000; Jenuwein and Allis, 2001). Chromatin remodeling as a possible circadian regulatory mechanism was first suggested by an experiment demonstrating that in mice, light pulses during the subjective night promote phosphorylation of serine 10 of histone H3 (Crosio *et al.*, 2000). Subsequent work has shown that at the *Per1* and *Per2* promoters, lysine 9 of histone H3 is rhythmically acetylated (Etchegaray *et al.*, 2003) and that CLOCK and NPAS2 may act to recruit histone acetyltransferases (HATs) to the *Per1* promoter (Etchegaray *et al.*, 2003; Curtis *et al.*, 2004).

Interestingly, CLOCK has been shown to exhibit intrinsic HAT activity toward lysine residues of histones H3 and H4 (Doi *et al.*, 2006). This suggests that CLOCK, while activating transcription with its partner, BMAL1, may rhythmically acetylate histones at clock-controlled genes and thereby participate in chromatin remodeling. CLOCK also acetylates non-histone substrates, including BMAL1 at lysine 537 (Hirayama *et al.*, 2007), as well as the glucocorticoid receptor (Nader *et al.*, 2009). Additional work is needed to clarify the relationship of CLOCK HAT activity to the circadian control of gene expression. Recently, a role for CLOCK HAT activity in facilitating herpes simplex virus gene expression in infected mammalian cells has been reported (Kalamvoki and Roizman, 2010), hence this activity of CLOCK may not be restricted solely to circadian regulation.

Histone deacetylases (HDACs) act to remove acetyl groups from histone tails to promote transcriptional repression (Finkel *et al.*, 2009). The NAD⁺-dependent HDAC Sirtuin 1 (SIRT1) has been shown recently to be involved in the mammalian molecular clock mechanism. SIRT1 acts to deacetylate lysines 9 and 14 of histone H3 as well as lysine 16 of histone H4 leading to chromatin condensation and transcriptional repression (Vaquero *et al.*, 2004; Nakahata *et al.*, 2008). Furthermore, SIRT1 seems to bind CLOCK-BMAL1 to form a

complex that is recruited in a circadian manner to promoters of clock-controlled genes. Through its interaction with CLOCK-BMAL1, SIRT1 participates in the circadian expression of *Bmal1*, *Rorc*, *Per2* and *Cry1* (Asher *et al.*, 2008). Interestingly, SIRT1 also acts directly on two core clock components—it reverses the CLOCK-mediated acetylation of lysine 537 of BMAL1 (Nakahata *et al.*, 2008), and deacetylates and promotes the degradation of PER2 (Asher *et al.*, 2008).

It is interesting to note here that because SIRT1 is controlled by the cellular NAD⁺:NADH ratio, its activity is intimately tied to a cell's redox state. This finding fits nicely with earlier work showing that the transcriptional activity of CLOCK and NPAS2 is regulated by cellular redox state (Rutter *et al.*, 2001). Expression of the gene encoding the rate-limiting enzyme in the NAD⁺ salvage pathway, nicotinamide phosphoribosyltransferase (NAMPT), is circadian owing to E-box-mediated CLOCK/BMAL1 activation. The resulting daily oscillation in NAMPT activity produces rhythmic levels of NAD⁺ in cells (Nakahata *et al.*, 2009; Ramsey *et al.*, 2009). Hence, a novel feedback loop connecting cellular metabolism and the circadian clock has been uncovered whereby CLOCK/BMAL1 positively regulates NAD⁺ levels, and thus SIRT1 activity through the circadian control of *Nampt* expression. SIRT1, in turn, negatively regulates CLOCK/BMAL1 activity by promoting transcriptional repression and participates in the oscillation of its own coenzyme, NAD⁺. As mentioned earlier, this and several other connections between the circadian system and metabolic pathways represent an emerging area of investigation as reviewed comprehensively elsewhere (Green *et al.*, 2008; Bass and Takahashi, 2010; Bellet and Sassone-Corsi, 2010; Maury *et al.*, 2010; Asher and Schibler, 2011).

VII. POSTTRANSCRIPTIONAL CLOCK MECHANISMS

Despite significant progress in elucidating the role of posttranslational regulation of the molecular clock in mammals, only recently have the contributions of posttranscriptional regulatory processes to clock function been explored (Kojima *et al.*, 2011; Staiger and Koster, 2011). Because many of the core clock genes, as well as clock-controlled genes, exhibit circadian oscillations in their transcript levels it is important to determine what processes mediate daily mRNA turnover in mammalian cells. Furthermore, in mice only between 33–50% of the genes that encode rhythmic proteins also manifest rhythmic transcript levels, indicating that the other mRNAs are regulated at the posttranscriptional level (Reddy *et al.*, 2006; Deery *et al.*, 2009).

A. MicroRNAs

Of particular relevance to the elucidation of posttranscriptional clock mechanisms, is recent work revealing circadian control of the expression of microRNAs (miRNAs) and their role in the cycling of clock gene transcripts in mammals. Transcribed from non-coding genomic regions, miRNAs are short, single-stranded RNA molecules 19–25 nucleotides in length that interact with the 3' untranslated regions (3'UTRs) of target transcripts to induce the cleavage/destabilization of, or to repress translation of, the target mRNA (Bushati and Cohen, 2007; Rana, 2007; Bartel, 2009; Guo *et al.*, 2010). If one report is correct in positing that most mammalian mRNAs are conserved targets of miRNAs (Friedman *et al.*, 2008), it will be important to understand how this regulatory mechanism affects the molecular clock.

Results from a genome-wide screen to identify CREB-regulated miRNAs (Impey *et al.*, 2004) have encouraged analysis of miRNA expression specifically in the SCN of mice (Cheng *et al.*, 2007). Two miRNAs, miR-219 and miR-132, exhibit circadian expression in the SCN and harbor cAMP-response elements (CREs) in their promoters, yet only in miR-219 has an E-box element (noncanonical) been identified. Coexpression of CLOCK and BMAL1 in PC12 cells induces expression of miR-219, but not of miR-132. Under

constant conditions, both miRNAs exhibit a circadian oscillation solely in the SCN, peaking during the mid-subjective day. This rhythm is abolished in *Cry1/Cry2* double mutant mice. Evidence that miR-132 downregulates translation of *Per2* comes from treatment of mice with an antagomir to this miRNA. Antagomirs, modified oligoribonucleotides complementary to a specific miRNA, act to block miRNA function (Krutzfeldt *et al.*, 2005). The miR-132 antagomir results in increased PER2 expression in mice. Finally, circadian period length and light-dependent clock resetting are altered by the antagomir-mediated silencing of miR-219 and miR-132, respectively (Cheng *et al.*, 2007).

To further investigate the role of miR-132 in photic entrainment of the SCN clock, one group has generated a transgenic mouse model that conditionally expresses miR-132 in the SCN and forebrain (Alvarez-Saavedra *et al.*, 2011). Following photic stimulation, the miR-132 transgenic animals experience attenuation in light-induced resetting of behavioral rhythms, and a concomitant decline in PER1 and PER2 levels. Analysis of putative miR-132 targets in the SCN has revealed several that are involved in chromatin remodeling and translational control. Through posttranscriptional regulation of these genes, it is proposed that miR-132 is able to control chromatin remodeling and translation within SCN neurons, thus mediating clock entrainment by regulating *Per* expression. Indeed, the *Per1* and *Per2* promoters are bound to, and activated by, methyl CpG island binding protein 2 (MeCP2), whereas PABP-interacting protein 2A (PAIP2A) and B-cell translocation gene 2 (BTG2) have the opposite effect—they suppress PER translation by promoting *Per* mRNA decay (Alvarez-Saavedra *et al.*, 2011).

Another group has identified a role for the miR-192/194 cluster in the regulation of the *Per* family in HeLa and NIH3T3 cells (Nagel *et al.*, 2009). The 3' UTRs of all three *Per* genes contain putative target sites for miR-192, miR-194, or both. Evidence shows that miR-192/194 downregulates the *Per* genes by acting on mRNA. This downregulation causes a shortening of the circadian period (Nagel *et al.*, 2009).

In liver, the most abundant microRNA, miR-122, is transcribed in a rhythmic manner with pri-miR-122 peaking at zeitgeber time 0 (ZT0) and reaching a nadir at ZT12, although the levels of miR-122 remain constant throughout the day (Gatfield *et al.*, 2009). The expression of miR-122 is most likely regulated by REV-ERB α/β through two conserved ROREs in this gene's promoter as *Rev-erba*^{-/-} mice experience attenuation in pri-miR-122 accumulation. Microarray analyses of gene expression in liver tissue of mice treated with a miR-122 antisense oligonucleotide have demonstrated several mRNAs that are under circadian regulation including peroxisome proliferator-activated receptor β/δ (PPAR β/δ) and the PPAR α coactivator SMARCD1/BAF60a (SWI/SNF-related, matrix-associated, actin-dependent regulator of chromatin, subfamily d, member 1), both circadian regulators of metabolism (Yang *et al.*, 2006; Li *et al.*, 2008). Indeed, the 3' UTR of *Ppar β/δ* contains target sites for miR-122, and this protein is upregulated by two-to-threefold upon inactivation of miR-122 (Gatfield *et al.*, 2009). A similar mechanism of miR-122 regulation at the 3' UTR of *Smarcd1/Baf60a* is apparent. The identification of the circadian regulation of miR-122 expression in liver and its role in the circadian control of downstream targets involved in metabolism provides further insight into tissue-specific links between the circadian system and metabolic function.

B. RNA binding proteins

In addition to the miRNA mechanisms just described, several RNA binding proteins have been identified that regulate clock-related RNA transcripts by different mechanisms. Most RNAs contain *cis*-acting elements in their 3' UTRs to which *trans*-acting protein factors can bind and regulate splicing, transport, stability and translation (Moore, 2005; Gratacós and Brewer, 2010). The first RNA binding protein with a demonstrated effect on the mouse

molecular clockwork is LARK, for which there are two forms, LARK1 (RBM4) and LARK2 (RBM4B) (Kojima *et al.*, 2007). Although transcripts for both *Lark* isoforms are found in the SCN, only the protein levels oscillate. Interestingly, the phase of the LARK protein oscillation is similar to that for PER1. Investigation of the *Per1* 3' UTR has revealed a specific *cis* element to which LARK binds and promotes translation. Knockdown of *Lark* transcript causes a shorter circadian period, while overexpression of LARK protein increases circadian period length. Thus, LARK seems to act on *Per1* posttranscriptionally by enhancing translation and conferring robustness on the PER1 protein oscillation (Kojima *et al.*, 2007).

One of the more common *cis*-acting elements in mammalian 3' UTRs is the AU-rich element (ARE) (Gratacós and Brewer, 2010). A protein which binds these AREs and is abundant in many tissues is polypyrimidine tract-binding protein 1 (PTBP1; also known as hnRNP I). Recently, circadian oscillation of the mouse *Per2* mRNA has been shown to be regulated by PTBP1 (Woo *et al.*, 2009). PTBP1 binds to the 3' UTR of the *Per2* transcript and promotes its decay. Cytoplasmic PTBP1 levels increase at the time at which there is a concomitant rapid decline in *Per2* RNA levels. Upon knockdown of PTBP1 expression with siRNA, the peak amplitude of *Per2* expression increased. Because several other RNA binding proteins have been detected at the *Per2* 3' UTR, it may be that the effect of PTBP1 on *Per2* RNA decay relies on additional proteins (Woo *et al.*, 2009).

Another RNA binding protein implicated in the regulation of a circadian gene is heterogeneous nuclear ribonucleoprotein D (HNRNP D; also known as AUF1). HNRNP D binds to specific sequences in target mRNA 3' UTRs and regulates transcript stability or the promotion of translation (Gratacós and Brewer, 2010). One group's examination of the *Cry1* transcript has revealed a 610-bp 3' UTR which contains an ARE that, when deleted, results in *Cry1* mRNA stability (Woo *et al.*, 2010). HNRNP D has been identified as the protein responsible for binding to the *Cry1* ARE and promoting mRNA turnover. Similar to the findings for PTBP1 and *Per2*, the cytoplasmic levels of HNRNP D levels exhibit a maximum as *Cry1* mRNA levels are in decline. Knockdown of HNRNP D results in stabilized *Cry1* transcript and enhanced oscillation amplitude (Woo *et al.*, 2010). Thus, cytoplasmic destabilization of both the *Per2* and *Cry1* transcripts share many similarities. It will be interesting to determine what posttranscriptional processes act on the other core clock gene transcripts.

Decay pathways for mRNAs in mammalian cells are either deadenylation-dependent or deadenylation-independent (Gratacós and Brewer, 2010). Both PTBP1 and HNRNP D are involved in the deadenylation-dependent pathway. Deadenylation involves the removal of a transcript's poly(A) tail followed by breakdown of the RNA. It is interesting to note that a circadian deadenylase gene, *Nocturnin* (*Ccrn4l*), has been characterized in mammals where it is rhythmically expressed in multiple tissues, including the SCN (Wang *et al.*, 2001). *Noc* circadian expression is damped in the liver of *Clock* mutant mice, and is constitutively elevated in *Cry* double mutant animals (Oishi *et al.*, 2003). Although *Noc*^{-/-} mice exhibit normal locomotor activity rhythms and clock gene expression, they do have several aberrant metabolic phenotypes (Green *et al.*, 2007). Thus, *Noc* is a clock-controlled gene—a clock output. Interestingly, it has recently been shown that circadian expression of *Noc* in mouse liver is controlled by miR-122 (Kojima *et al.*, 2010).

VIII. EFFECTS OF TEMPERATURE ON THE MAMMALIAN CLOCK

A. Temperature as an entraining agent

Temperature is an important environmental entraining agent for many organisms, yet in homeothermic vertebrates, including mammals, changes in ambient temperature either do

not entrain circadian rhythms of locomotor activity, or do so poorly (Hoffmann, 1969; Aschoff and Tokura, 1986; Francis and Coleman, 1997; Palkova *et al.*, 1999). Mammals do, however, experience circadian rhythms in core body temperature with a fluctuation of 1–4°C that are regulated by the SCN (Refinetti and Menaker, 1992). As mentioned previously, cells of most peripheral tissues throughout the mammalian body harbor cell-autonomous circadian oscillators (Welsh *et al.*, 1995; Balsalobre *et al.*, 1998; Nagoshi *et al.*, 2004; Welsh *et al.*, 2004; Yoo *et al.*, 2004) which are synchronized to external cues by rhythmic signals from the SCN (Silver *et al.*, 1996; Earnest *et al.*, 1999; Buijs and Kalsbeek, 2001). These observations raise the intriguing question as to what effect, if any, the normal circadian variation in core body temperature may have upon the cell-autonomous oscillators in peripheral tissues. Could body temperature entrain peripheral oscillators in mammals and, perhaps just as interesting, what properties of the SCN prevent it from being synchronized by environmental temperature cycles?

Studies have demonstrated that circadian rhythms of gene expression in cultures of rat-1 fibroblasts (Brown *et al.*, 2002) and rat astrocytes (Prolo *et al.*, 2005) can be entrained to temperature fluctuations of 4°C and 1.5°C, respectively. Although rhythmicity in both cell types damps within a few cycles upon cessation of the temperature rhythms, damping can be delayed by exposing fibroblasts to natural body temperature oscillations (Brown *et al.*, 2002) or by co-culturing astrocytes with SCN explants (Prolo *et al.*, 2005). Furthermore, when mice are exposed to inverted environmental temperature cycles of 37°C during the day and 24°C during the night, circadian rhythms of gene expression in the liver are reversed without affecting the central clock in the SCN (Brown *et al.*, 2002). It should be mentioned that this phenomenon of decoupling peripheral circadian rhythms from the SCN has also been observed under paradigms of restricted feeding in mammals, and that restricted feeding can alter body temperature rhythms (Damiola *et al.*, 2000; Stokkan *et al.*, 2001). Indeed, it has been proposed that food availability and environmental temperature are related, but independent, entraining cues (Brown *et al.*, 2002). Thus, it is clear that in cultured nonneuronal mammalian cells, and in peripheral tissues *in vivo* in mammals, temperature cycles can entrain circadian rhythms of gene expression.

In contrast to peripheral oscillators, SCN rhythms *in vivo* are unaffected by environmental temperature changes that phase shift circadian rhythms in other brain regions (Brown *et al.*, 2002). Recent work in our lab has extended this finding by demonstrating that in mice, the resistance of the SCN to entrainment by ambient temperature is not a cell-autonomous property of individual SCN neurons. Rather, this phenomenon is an emergent property of the SCN network and is therefore dependent upon intercellular coupling among neurons (Buhr *et al.*, 2010). Compelling evidence for the role of intercellular coupling comes from experiments showing that blocking either voltage-gated Na⁺ channels with TTX or L-type, but not T-type, Ca²⁺ channels with nimodipine, renders the SCN susceptible to resetting by temperature pulses. Furthermore, communication between the neurochemically-distinct ventral and dorsal regions of the SCN is necessary as cultures of either of these regions alone are shifted in response to temperature changes (Buhr *et al.*, 2010).

Work implying a role for heat shock factor 1 (HSF1) and other components of the heat shock response pathway in circadian gene expression in mammalian liver (Kornmann *et al.*, 2007; Reinke *et al.*, 2008) has encouraged investigation of the possible involvement of this pathway in temperature resetting in mammals. When SCN cultures are pulsed for 1 hr with the heat shock pathway antagonist KNK437, no phase shifts occur. In contrast, the same treatment induces strong phase shifts in pituitary and lung cultures, indicating that in non-SCN tissues, KNK437 mimics the effect of 1-hr cool (33.5°C) pulses in reducing HSF1-mediated transcription (Buhr *et al.*, 2010). Phase shifts to warm (38.6°C) pulses observed in lung and pituitary are blocked both by KNK437 and quercetin, another HSF1 inhibitor.

Continuous inhibition of HSF1-mediated transcription via chronic KNK437 administration to SCN, lung and pituitary cultures causes an increase in circadian period. This effect is consistent with period lengthening of the circadian rhythm of locomotor activity observed in *Hsf1^{-/-}* mice (Reinke *et al.*, 2008). Taken together, these results suggest a molecular mechanism that involves HSF1 in temperature resetting in mammalian peripheral tissues (Buhr *et al.*, 2010).

It is important to mention here that two reports, in contrast to the work just presented, provide evidence that rat SCN slice cultures do exhibit entrainment to temperature cycles (Ruby *et al.*, 1999; Herzog and Huckfeldt, 2003). Several differences in these studies may account for this discrepancy. First, the work of Ruby *et al.* (1999) relied on extracellular recordings from single neurons as glass electrodes advanced along tracks through SCN slices maintained in culture for up to 60 hr. Herzog and Huckfeldt (2003) measured PER1::LUC rhythms from neonatal and juvenile rat SCN slices in cultures for up to two weeks and observed that these slice preparations were more sensitive to temperature pulses than adult tissues. In contrast, our studies used organotypic cultures of SCN and peripheral tissues from *Per2^{Luc}* reporter mice and measured in real time the PER2::LUC bioluminescence rhythm continuously for several days (Yoo *et al.*, 2004; Buhr *et al.*, 2010). The *in vivo* study of Brown *et al.* (2002) examined mice exposed to environmental temperature changes over a several day period and measured *Dbp* expression in SCN sections by *in situ* hybridization. Thus, the SCN culture techniques and output rhythms measured in each study differed. A final possibility is that species-specific differences between mice and rats account for the differences observed in some of these studies. Further work is necessary to clarify these issues.

B. Temperature compensation

One of the hallmarks of circadian rhythms in all organisms, from cyanobacteria to mammals, is that they are temperature compensated—daily rhythms remain constant as temperature increases or decreases across a physiologically viable range (Pittendrigh, 1993). This clock property can be expressed quantitatively as the Q_{10} , or temperature coefficient—the change in the rate of a biological rhythm or biochemical reaction as a result of increasing the ambient temperature by 10°C. The Q_{10} for most biochemical reactions is 2 to 3, yet circadian rhythms usually have a Q_{10} between 0.8–1.2 (Sweeney and Hastings, 1960). Efforts to elucidate a molecular mechanism underlying the temperature compensation property of circadian clocks have been unsuccessful until recently. The striking demonstration that the circadian rhythm of phosphorylation of the cyanobacterial clock protein, KaiC, can be reconstituted *in vitro* with just three Kai proteins and ATP, has led to experiments revealing that this *in vitro* oscillator is also temperature compensated (Nakajima *et al.*, 2005; Tomita *et al.*, 2005). Hence, as the cyanobacterial example aptly illustrates, temperature compensation must be an inherent property of at least some of the biochemical reactions comprising the molecular mechanism of the circadian clock of any given species.

In mammals, temperature compensation has been demonstrated in cell culture for rat-1 (Izumo *et al.*, 2003) and mouse (Tsuchiya *et al.*, 2003; Dibner *et al.*, 2009) fibroblasts, and for neonatal rat SCN neurons (Herzog and Huckfeldt, 2003). As for whole neural tissues, hamster retina (Tosini and Menaker, 1998), and rat (Ruby *et al.*, 1999), mouse (Buhr *et al.*, 2010), and ground squirrel (Ruby and Heller, 1996) SCN are temperature compensated. Several mammalian peripheral tissues also exhibit this property (Reyes *et al.*, 2008), as do human red blood cells (O'Neill and Reddy, 2011). Work by our group has also shown that the temperature compensation observed in SCN tissue is a cell-autonomous property and does not rely on intercellular coupling, unlike the resistance to temperature resetting of the SCN which is an emergent property of the SCN network (Buhr *et al.*, 2010).

Genetic mutations affecting the period of the circadian clock in mammals can also affect temperature compensation as shown for isolated *tau* mutant hamster retinal cultures (Tosini and Menaker, 1998). This is particularly intriguing in light of recent work demonstrating both in cell culture and in cell-free *in vitro* reactions, that the enzymatic activity of CK1 δ/ϵ toward circadian substrates such as PER2 is temperature-insensitive, but temperature-sensitive toward non-circadian substrates (Isojima *et al.*, 2009). Both wild-type and CK1 ϵ^{tau} exhibit temperature-insensitive phosphorylation of a peptide containing the PER2 β TrCP binding region, yet when a peptide substrate derived from the FASPS region of PER2 is tested, an increase in the temperature sensitivity of the CK1 ϵ^{tau} mutant enzyme is revealed (Isojima *et al.*, 2009). An effect of autophosphorylation state on CK1 δ/ϵ activity toward circadian-relevant substrates *in vitro* is also apparent. Thus, similar to the cyanobacterial system mentioned previously, the temperature-insensitive phosphorylation of mammalian circadian substrates by CK1 δ/ϵ can be reconstituted *in vitro*, and this property of CK1 δ/ϵ is dependent both on the substrate and on the phosphorylation state of the enzyme (Isojima *et al.*, 2009). Additional work will be necessary to replicate these results and to elucidate the biochemical mechanism underlying CK1 δ/ϵ circadian-specific temperature compensation.

IX. UNRESOLVED ISSUES AND FUTURE DIRECTIONS

As mentioned at the beginning of this review, extensive work has shown that across phyla, the primary molecular mechanism underlying cell-autonomous circadian oscillators is composed of autoregulatory feedback loops of transcription and translation. Hence, the existence of transcription-independent oscillations and a potential role for such oscillations in the function of the cellular clock in mammals and other organisms seems surprising. Indeed, it was the cell-free recapitulation in a test tube of the circadian rhythm of KaiC phosphorylation by colleagues working on the cyanobacterial circadian clock (Nakajima *et al.*, 2005; Tomita *et al.*, 2005) that generated recent interest in transcription-independent clock processes. Further work in cyanobacteria has shown that both circadian transcriptional/translational mechanisms and transcription-independent posttranslational mechanisms are necessary for a competent circadian clock in this organism (Kitayama *et al.*, 2008).

Hints that transcription-independent processes in the mammalian circadian clock exist come from several studies. The surprising discovery that rhythmic transcription of the core clock genes *Bmal1*, *Cry1*, and *Cry2* (Fan *et al.*, 2007; Liu *et al.*, 2008) is not necessary for circadian clock function in mammalian cells, and that the mammalian cellular clock is particularly resilient to attenuation of transcription (Dibner *et al.*, 2009), suggests that mechanisms other than the transcriptional/translational feedback loop are involved in the generation of oscillations. Perhaps the most striking demonstration of a transcription-independent circadian rhythm in mammals is the recent report of a daily oscillation in human red blood cells of the oxidation and subsequent monomer-to-dimer transition of peroxiredoxin (PRX), a protein that inactivates reactive oxygen species (O'Neill and Reddy, 2011). In liver cells, expression of PRX is circadian (Reddy *et al.*, 2006), but this is not possible in mature erythrocytes which have no nucleus. Inhibitors of transcription and translation have no effect on the circadian rhythm of PRX oxidation in human red blood cells further suggesting that the observed PRX oxidation rhythm in erythrocytes is a transcription-independent process. Moreover, detection of another circadian rhythm in erythrocytes—the transition of hemoglobin between dimer and tetramer states—seems also to be transcription-independent. A circadian rhythm of PRX oxidation has also been demonstrated by the same group in the green algae *Ostreococcus tauri*, a primitive eukaryote, even when gene expression is halted by exposing these cells to DD (O'Neill *et al.*, 2011). Although these findings must be repeated and validated in other model organisms, they suggest an intriguing avenue for further work.

A major goal toward understanding any biological system is to successfully model that system mathematically. Accomplishing this requires that biologists and modelers work together to incorporate experimental results into models such that they may be used to make testable predictions. The ongoing development of models of the mammalian circadian clock has been particularly helpful recently as they take into account the “combinatorial complexity” of clock component interactions as well as stochastic properties (Yamada and Forger, 2010). Model-based predictions regarding the effects of mutations in core clock genes (Liu *et al.*, 2007b), the role of posttranslational processes on clock components (Gallego *et al.*, 2006a), and the electrical properties of the SCN (Belle *et al.*, 2009), have been validated empirically. The necessity for both molecular noise and intercellular coupling to induce rhythms in a population of SCN neurons has also been predicted via modeling and subsequently confirmed by experiment (Ko *et al.*, 2010). A future challenge to clock modelers will be to address the complex connections and interactions between the mammalian circadian clock and other systems, including the many emerging links between the clock and metabolism.

While there are many potentially fruitful paths of pursuit in the study of mammalian circadian clock genetics, we will conclude by mentioning a few here. Large-scale screening of small-molecule libraries may yield promising targets for therapeutic intervention of circadian-related genetic disorders in humans including FASPS and seasonal affective disorder (Liu *et al.*, 2007a). New approaches in synthetic biology in which artificial transcriptional circuits are used to define networks of oscillating genes or to interrogate the function of circadian-related *cis*-acting elements promise to further our understanding of clock transcriptional pathways (Ueda *et al.*, 2005; Kumaki *et al.*, 2008; Ukai-Tadenuma *et al.*, 2008). Understanding circadian phenotypes that occur in mammals in the absence of underlying changes in DNA sequence—via epigenetic processes—will require ongoing work (Bellet and Sassone-Corsi, 2010; Ripperger and Meroow, 2011). Elucidation of the molecular and biochemical mechanisms of temperature-insensitive phenomena in mammalian clock systems remains an important goal (Brown *et al.*, 2002; Isojima *et al.*, 2009; Buhr *et al.*, 2010). In all of these endeavors, the mouse will remain the mammalian genetic model of choice.

X. CONCLUSION

Over the past 10 years remarkable progress has been made in our understanding of the genetics of the mammalian circadian clock. The transcriptional/translational feedback loop model of the molecular oscillator within cells, for which there is evidence across phyla, has formed the foundation of our understanding of the molecular clockwork. This model, however, must be modified given the new levels of hierarchy and complexity evident from recent work. It is necessary to study the mammalian clock at all levels from single cells, to cell-cell interactions within a tissue, to tissue-level properties, and finally, at the level of behavior. Emergent clock properties arise from the interactions among cells—properties that cannot be studied at the single-cell level. Likewise, focusing on behavior or tissue-specific processes alone will overlook cell-autonomous clock properties. New advances in reporter technology, microarrays, mathematical modeling, perturbation analysis methods, and systems biology will continue to elucidate properties of the molecular clock. Hence, a lesson learned from the work presented herein is that with respect to the mammalian circadian clock, a systems-wide approach must be advocated. In the near future we should look forward to a better understanding of how the mammalian clockwork is integrated with the other physiological systems of the body, and perhaps be better able to develop therapies for human clock-related disorders.

Acknowledgments

This work was supported by NIH grant 1R15GM086825-01 to P.L.L. and NIH grants U01 MH61915, P50 MH074924, and R01 MH078024 to J.S.T. J.S.T. is an Investigator in the Howard Hughes Medical Institute.

REFERENCES

- Abe M, Herzog ED, Block GD. Lithium lengthens the circadian period of individual suprachiasmatic nucleus neurons. *NeuroReport*. 2000; 11:3261–3264. [PubMed: 11043560]
- Abraham U, Granada AE, Westermark PO, Heine M, Kramer A, Herzel H. Coupling governs entrainment range of circadian clocks. *Mol. Syst. Biol.* 2010; 6:438. [PubMed: 21119632]
- Abrahamson EE, Moore RY. Suprachiasmatic nucleus in the mouse: retinal innervation, intrinsic organization and efferent projections. *Brain Res.* 2001; 916:172–191. [PubMed: 11597605]
- Adams DJ, van der Weyden L. Contemporary approaches for modifying the mouse genome. *Physiol. Genomics*. 2008; 34:225–238. [PubMed: 18559964]
- Akashi M, Tsuchiya Y, Yoshino T, Nishida E. Control of intracellular dynamics of mammalian period proteins by casein kinase I ϵ (CKI ϵ) and CKI δ in cultured cells. *Mol. Cell. Biol.* 2002; 22:1693–1703. [PubMed: 11865049]
- Akashi M, Takumi T. The orphan nuclear receptor ROR α regulates circadian transcription of the mammalian core-clock *Bmal1*. *Nat. Struct. Mol. Biol.* 2005; 12:441–448. [PubMed: 15821743]
- Akhtar RA, Reddy AB, Maywood ES, Clayton JD, King VM, Smith AG, Gant TW, Hastings MH, Kyriacou CP. Circadian cycling of the mouse liver transcriptome, as revealed by cDNA microarray, is driven by the suprachiasmatic nucleus. *Curr. Biol.* 2002; 12:540–550. [PubMed: 11937022]
- Albrecht U, Sun ZS, Eichele G, Lee CC. A differential response of two putative mammalian circadian regulators, *mper1* and *mper2*, to light. *Cell*. 1997; 91:1055–1064. [PubMed: 9428527]
- Ali A, Hoeflich KP, Woodgett JR. Glycogen synthase kinase-3: properties, functions, and regulation. *Chem. Rev.* 2001; 101:2527–2540. [PubMed: 11749387]
- Alvarez-Saavedra M, Antoun G, Yanagiya A, Oliva-Hernandez R, Cornejo-Palma D, Perez-Iratxeta C, Sonenberg N, Cheng HY. miRNA-132 orchestrates chromatin remodeling and translational control of the circadian clock. *Hum. Mol. Genet.* 2011; 20:731–751. [PubMed: 21118894]
- Antoch MP, Song EJ, Chang AM, Vitaterna MH, Zhao Y, Wilsbacher LD, Sangoram AM, King DP, Pinto LH, Takahashi JS. Functional identification of the mouse circadian *Clock* gene by transgenic BAC rescue. *Cell*. 1997; 89:655–667. [PubMed: 9160756]
- Aschoff J, Tokura H. Circadian activity rhythms in squirrel monkeys: entrainment by temperature cycles. *J. Biol. Rhythms*. 1986; 1:91–99. [PubMed: 2979582]
- Asher G, Gatfield D, Stratmann M, Reinke H, Dibner C, Kreppel F, Mostoslavsky R, Alt FW, Schibler U. SIRT1 regulates circadian clock gene expression through PER2 deacetylation. *Cell*. 2008; 134:317–328. [PubMed: 18662546]
- Asher G, Schibler U. Crosstalk between components of circadian and metabolic cycles in mammals. *Cell Metab.* 2011; 13:125–137. [PubMed: 21284980]
- Aston-Jones G, Chen S, Zhu Y, Oshinsky ML. A neural circuit for circadian regulation of arousal. *Nat. Neurosci.* 2001; 4:732–738. [PubMed: 11426230]
- Aton SJ, Colwell CS, Harmar AJ, Waschek J, Herzog ED. Vasoactive intestinal polypeptide mediates circadian rhythmicity and synchrony in mammalian clock neurons. *Nat. Neurosci.* 2005; 8:476–483. [PubMed: 15750589]
- Bacon Y, Ooi A, Kerr S, Shaw-Andrews L, Winchester L, Breeds S, Tymoska-Lalanne Z, Clay J, Greenfield AG, Nolan PM. Screening for novel ENU-induced rhythm, entrainment and activity mutants. *Genes Brain Behav.* 2004; 3:196–205. [PubMed: 15248865]
- Badura L, Swanson T, Adamowicz W, Adams J, Cianfrogna J, Fisher K, Holland J, Kleiman R, Nelson F, Reynolds L, et al. An inhibitor of casein kinase I ϵ induces phase delays in circadian rhythms under free-running and entrained conditions. *J. Pharmacol. Exp. Ther.* 2007; 322:730–738. [PubMed: 17502429]
- Bae K, Jin X, Maywood ES, Hastings MH, Reppert SM, Weaver DR. Differential functions of *mPer1*, *mPer2*, and *mPer3* in the SCN circadian clock. *Neuron*. 2001; 30:525–536. [PubMed: 11395012]

- Baggs JE, Price TS, DiTacchio L, Panda S, Fitzgerald GA, Hogenesch JB. Network features of the mammalian circadian clock. *PLoS Biol.* 2009; 7:e52. [PubMed: 19278294]
- Balsalobre A, Damiola F, Schibler U. A serum shock induces circadian gene expression in mammalian tissue culture cells. *Cell.* 1998; 93:929–937. [PubMed: 9635423]
- Balsalobre A, Brown SA, Marcacci L, Tronche F, Kellendonk C, Reichardt HM, Schutz G, Schibler U. Resetting of circadian time in peripheral tissues by glucocorticoid signaling. *Science.* 2000; 289:2344–2347. [PubMed: 11009419]
- Bartel DP. MicroRNAs: target recognition and regulatory functions. *Cell.* 2009; 136:215–233. [PubMed: 19167326]
- Bass J, Takahashi JS. Circadian integration of metabolism and energetics. *Science.* 2010; 330:1349–1354. [PubMed: 21127246]
- Belle MD, Diekmann CO, Forger DB, Piggins HD. Daily electrical silencing in the mammalian circadian clock. *Science.* 2009; 326:281–284. [PubMed: 19815775]
- Bellet MM, Sassone-Corsi P. Mammalian circadian clock and metabolism - the epigenetic link. *J. Cell Sci.* 2010; 123:3837–3848. [PubMed: 21048160]
- Berson DM, Dunn FA, Takao M. Phototransduction by retinal ganglion cells that set the circadian clock. *Science.* 2002; 295:1070–1073. [PubMed: 11834835]
- Blake JA, Bult CJ, Kadin JA, Richardson JE, Eppig JT. The Mouse Genome Database (MGD): premier model organism resource for mammalian genomics and genetics. *Nucleic Acids Res.* 2010; 39:D842–D848. [PubMed: 21051359]
- Brown S, Zimbrunn G, Fleury-Olela F, Preitner N, Schibler U. Rhythms of mammalian body temperature can sustain peripheral circadian clocks. *Curr. Biol.* 2002; 12:1574–1583. [PubMed: 12372249]
- Brown TM, Colwell CS, Waschek JA, Piggins HD. Disrupted neuronal activity rhythms in the suprachiasmatic nuclei of vasoactive intestinal polypeptide-deficient mice. *J. Neurophysiol.* 2007; 97:2553–2558. [PubMed: 17151217]
- Buhr ED, Yoo SH, Takahashi JS. Temperature as a universal resetting cue for mammalian circadian oscillators. *Science.* 2010; 330:379–385. [PubMed: 20947768]
- Buijs RM, Kalsbeek A. Hypothalamic integration of central and peripheral clocks. *Nat. Rev. Neurosci.* 2001; 2:521–526. [PubMed: 11433377]
- Bunger MK, Wilsbacher LD, Moran SM, Clendenin C, Radcliffe LA, Hogenesch JB, Simon C, Takahashi JS, Bradfield CA. *Mop3* is an essential component of the master circadian pacemaker in mammals. *Cell.* 2000; 103:1009–1017. [PubMed: 11163178]
- Bushati N, Cohen SM. microRNA functions. *Annu. Rev. Cell. Dev. Biol.* 2007; 23:175–205. [PubMed: 17506695]
- Busino L, Bassermann F, Maiolica A, Lee C, Nolan PM, Godinho SI, Draetta GF, Pagano M. SCF^{Fbx13} controls the oscillation of the circadian clock by directing the degradation of cryptochrome proteins. *Science.* 2007; 316:900–904. [PubMed: 17463251]
- Camacho F, Cilio M, Guo Y, Virshup DM, Patel K, Khorkova O, Styren S, Morse B, Yao Z, Keesler GA. Human casein kinase Iδ phosphorylation of human circadian clock proteins *period* 1 and 2. *FEBS Lett.* 2001; 489:159–165. [PubMed: 11165242]
- Cardone L, Hirayama J, Giordano F, Tamaru T, Palvimo JJ, Sassone-Corsi P. Circadian clock control by SUMOylation of BMAL1. *Science.* 2005; 309:1390–1394. [PubMed: 16109848]
- Cardozo T, Pagano M. The SCF ubiquitin ligase: insights into a molecular machine. *Nat. Rev. Mol. Cell Biol.* 2004; 5:739–751. [PubMed: 15340381]
- Cermakian N, Monaco L, Pando MP, Dierich A, Sassone-Corsi P. Altered behavioral rhythms and clock gene expression in mice with a targeted mutation in the *Period1* gene. *EMBO J.* 2001; 20:3967–3974. [PubMed: 11483500]
- Chen R, Schirmer A, Lee Y, Lee H, Kumar V, Yoo SH, Takahashi JS, Lee C. Rhythmic PER abundance defines a critical nodal point for negative feedback within the circadian clock mechanism. *Mol. Cell.* 2009; 36:417–430. [PubMed: 19917250]
- Cheng HY, Papp JW, Varlamova O, Dziema H, Russell B, Curfman JP, Nakazawa T, Shimizu K, Okamura H, Impey S, et al. microRNA modulation of circadian-clock period and entrainment. *Neuron.* 2007; 54:813–829. [PubMed: 17553428]

- Cheng MY, Bullock CM, Li C, Lee AG, Bermak JC, Belluzzi J, Weaver DR, Leslie FM, Zhou QY. Prokineticin 2 transmits the behavioural circadian rhythm of the suprachiasmatic nucleus. *Nature*. 2002; 417:405–410. [PubMed: 12024206]
- Ciechanover A, Orian A, Schwartz AL. Ubiquitin-mediated proteolysis: biological regulation via destruction. *Bioessays*. 2000; 22:442–451. [PubMed: 10797484]
- Clark AT, Goldowitz D, Takahashi JS, Vitaterna MH, Siepka SM, Peters LL, Frankel WN, Carlson GA, Rossant J, Nadeau JH, et al. Implementing large-scale ENU mutagenesis screens in North America. *Genetica*. 2004; 122:51–64. [PubMed: 15619961]
- Colwell CS, Michel S, Itri J, Rodriguez W, Tam J, Lelievre V, Hu Z, Liu X, Waschek JA. Disrupted circadian rhythms in VIP- and PHI-deficient mice. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 2003; 285:R939–R949. [PubMed: 12855416]
- Crosio C, Cermakian N, Allis CD, Sassone-Corsi P. Light induces chromatin modification in cells of the mammalian circadian clock. *Nat. Neurosci.* 2000; 3:1241–1247. [PubMed: 11100144]
- Curtis AM, Seo SB, Westgate EJ, Rudic RD, Smyth EM, Chakravarti D, FitzGerald GA, McNamara P. Histone acetyltransferase-dependent chromatin remodeling and the vascular clock. *J. Biol. Chem.* 2004; 279:7091–7097. [PubMed: 14645221]
- Cutler DJ, Haraura M, Reed HE, Shen S, Sheward WJ, Morrison CF, Marston HM, Harmar AJ, Piggins HD. The mouse VPAC₂ receptor confers suprachiasmatic nuclei cellular rhythmicity and responsiveness to vasoactive intestinal polypeptide *in vitro*. *Eur. J. Neurosci.* 2003; 17:197–204. [PubMed: 12542655]
- Dacey DM, Liao HW, Peterson BB, Robinson FR, Smith VC, Pokorny J, Yau KW, Gamlin PD. Melanopsin-expressing ganglion cells in primate retina signal colour and irradiance and project to the LGN. *Nature*. 2005; 433:749–754. [PubMed: 15716953]
- Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U. Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus. *Genes Dev.* 2000; 14:2950–2961. [PubMed: 11114885]
- Dardente H, Fortier EE, Martineau V, Cermakian N. Cryptochromes impair phosphorylation of transcriptional activators in the clock: a general mechanism for circadian repression. *Biochem. J.* 2007; 402:525–536. [PubMed: 17115977]
- DeBruyne JP, Noton E, Lambert CM, Maywood ES, Weaver DR, Reppert SM. A clock shock: mouse CLOCK is not required for circadian oscillator function. *Neuron*. 2006; 50:465–477. [PubMed: 16675400]
- DeBruyne JP, Weaver DR, Reppert SM. CLOCK and NPAS2 have overlapping roles in the suprachiasmatic circadian clock. *Nat. Neurosci.* 2007a; 10:543–545. [PubMed: 17417633]
- DeBruyne JP, Weaver DR, Reppert SM. Peripheral circadian oscillators require CLOCK. *Curr. Biol.* 2007b; 17:R538–R539. [PubMed: 17637349]
- Deery MJ, Maywood ES, Chesham JE, Sladek M, Karp NA, Green EW, Charles PD, Reddy AB, Kyriacou CP, Lilley KS, et al. Proteomic analysis reveals the role of synaptic vesicle cycling in sustaining the suprachiasmatic circadian clock. *Curr. Biol.* 2009; 19:2031–2036. [PubMed: 19913422]
- Dibner C, Sage D, Unser M, Bauer C, d'Eysmond T, Naef F, Schibler U. Circadian gene expression is resilient to large fluctuations in overall transcription rates. *EMBO J.* 2009; 28:123–134. [PubMed: 19078963]
- Do MT, Yau KW. Intrinsically photosensitive retinal ganglion cells. *Physiol. Rev.* 2010; 90:1547–1581. [PubMed: 20959623]
- Doble BW, Woodgett JR. GSK-3: tricks of the trade for a multi-tasking kinase. *J. Cell Sci.* 2003; 116:1175–1186. [PubMed: 12615961]
- Doherty CJ, Kay SA. Circadian control of global gene expression patterns. *Annu. Rev. Genet.* 2010; 44:419–444. [PubMed: 20809800]
- Doi M, Hirayama J, Sassone-Corsi P. Circadian regulator CLOCK is a histone acetyltransferase. *Cell.* 2006; 125:497–508. [PubMed: 16678094]
- Dudley CA, Erbel-Sieler C, Estill SJ, Reick M, Franken P, Pitts S, McKnight SL. Altered patterns of sleep and behavioral adaptability in NPAS2-deficient mice. *Science*. 2003; 301:379–383. [PubMed: 12843397]

- Duffield GE, Best JD, Meurers BH, Bittner A, Loros JJ, Dunlap JC. Circadian programs of transcriptional activation, signaling, and protein turnover revealed by microarray analysis of mammalian cells. *Curr. Biol.* 2002; 12:551–557. [PubMed: 11937023]
- Dunlap JC. Molecular bases for circadian clocks. *Cell.* 1999; 96:271–290. [PubMed: 9988221]
- Earnest DJ, Liang FQ, Ratcliff M, Cassone VM. Immortal time: circadian clock properties of rat suprachiasmatic cell lines. *Science.* 1999; 283:693–695. [PubMed: 9924030]
- Ebling FJ. The role of glutamate in the photic regulation of the suprachiasmatic nucleus. *Prog. Neurobiol.* 1996; 50:109–132. [PubMed: 8971980]
- Eide EJ, Kang H, Crapo S, Gallego M, Virshup DM. Casein kinase I in the mammalian circadian clock. *Methods Enzymol.* 2005a; 393:408–418. [PubMed: 15817302]
- Eide EJ, Woolf MF, Kang H, Woolf P, Hurst W, Camacho F, Vielhaber EL, Giovanni A, Virshup DM. Control of mammalian circadian rhythm by CKI ϵ -regulated proteasome-mediated PER2 degradation. *Mol. Cell. Biol.* 2005b; 25:2795–2807. [PubMed: 15767683]
- Etchegaray JP, Lee C, Wade PA, Reppert SM. Rhythmic histone acetylation underlies transcription in the mammalian circadian clock. *Nature.* 2003; 421:177–182. [PubMed: 12483227]
- Etchegaray JP, Machida KK, Noton E, Constance CM, Dallmann R, Di Napoli MN, DeBruyne JP, Lambert CM, Yu EA, Reppert SM, et al. Casein kinase 1 delta regulates the pace of the mammalian circadian clock. *Mol. Cell. Biol.* 2009; 29:3853–3866. [PubMed: 19414593]
- Etchegaray JP, Yu EA, Indic P, Dallmann R, Weaver DR. Casein kinase 1 delta (CK1 δ) regulates period length of the mouse suprachiasmatic circadian clock *in vitro*. *PLoS One.* 2010; 5:e10303. [PubMed: 20421981]
- Fan Y, Hida A, Anderson DA, Izumo M, Johnson CH. Cycling of CRYPTOCHROME proteins is not necessary for circadian-clock function in mammalian fibroblasts. *Curr. Biol.* 2007; 17:1091–1100. [PubMed: 17583506]
- Finkel T, Deng CX, Mostoslavsky R. Recent progress in the biology and physiology of sirtuins. *Nature.* 2009; 460:587–591. [PubMed: 19641587]
- Foster RG, Provencio I, Hudson D, Fiske S, De Grip W, Menaker M. Circadian photoreception in the retinally degenerate mouse (*rd/rd*). *J. Comp. Physiol. A.* 1991; 169:39–50. [PubMed: 1941717]
- Fox, J.; Barthold, S.; Davvison, M.; Newcomer, C.; Quimby, F.; Smith, A., editors. *The mouse in biomedical research.* 2nd ed.. Vol. 4 vols. Boston: Elsevier; 2007.
- Francis AJ, Coleman GJ. Phase response curves to ambient temperature pulses in rats. *Physiol. Behav.* 1997; 62:1211–1217. [PubMed: 9383104]
- Freedman MS, Lucas RJ, Soni B, von Schantz M, Munoz M, David-Gray Z, Foster R. Regulation of mammalian circadian behavior by non-rod, non-cone, ocular photoreceptors. *Science.* 1999; 284:502–504. [PubMed: 10205061]
- Friedman RC, Farh KKH, Burge CB, Bartel DP. Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res.* 2008; 19:92–105. [PubMed: 18955434]
- Fu Y, Zhong H, Wang MH, Luo DG, Liao HW, Maeda H, Hattar S, Frishman LJ, Yau KW. Intrinsically photosensitive retinal ganglion cells detect light with a vitamin A-based photopigment, melanopsin. *Proc. Natl. Acad. Sci. USA.* 2005; 102:10339–10344. [PubMed: 16014418]
- Gallego M, Eide EJ, Woolf MF, Virshup DM, Forger DB. An opposite role for tau in circadian rhythms revealed by mathematical modeling. *Proc. Natl. Acad. Sci. USA.* 2006a; 103:10618–10623. [PubMed: 16818876]
- Gallego M, Kang H, Virshup DM. Protein phosphatase 1 regulates the stability of the circadian protein PER2. *Biochem. J.* 2006b; 399:169–175. [PubMed: 16813562]
- Gallego M, Virshup DM. Post-translational modifications regulate the ticking of the circadian clock. *Nat. Rev. Mol. Cell Biol.* 2007; 8:139–148. [PubMed: 17245414]
- Gareau JR, Lima CD. The SUMO pathway: emerging mechanisms that shape specificity, conjugation and recognition. *Nat. Rev. Mol. Cell Biol.* 2010; 11:861–871. [PubMed: 21102611]
- Gatfield D, Le Martelot G, Vejnar CE, Gerlach D, Schaad O, Fleury-Olela F, Ruskeepaa AL, Oresic M, Esau CC, Zdobnov EM, et al. Integration of microRNA miR-122 in hepatic circadian gene expression. *Genes Dev.* 2009; 23:1313–1326. [PubMed: 19487572]

- Gekakis N, Staknis D, Nguyen HB, Davis FC, Wilsbacher LD, King DP, Takahashi JS, Weitz CJ. Role of the CLOCK protein in the mammalian circadian mechanism. *Science*. 1998; 280:1564–1569. [PubMed: 9616112]
- Gietzen KF, Virshup DM. Identification of inhibitory autophosphorylation sites in casein kinase I α . *J. Biol. Chem*. 1999; 274:32063–32070. [PubMed: 10542239]
- Godinho SI, Maywood ES, Shaw L, Tucci V, Barnard AR, Busino L, Pagano M, Kendall R, Quwaylid MM, Romero MR, et al. The after-hours mutant reveals a role for Fbx13 in determining mammalian circadian period. *Science*. 2007; 316:897–900. [PubMed: 17463252]
- Golombek DA, Rosenstein RE. Physiology of circadian entrainment. *Physiol. Rev*. 2010; 90:1063–1102. [PubMed: 20664079]
- Gooley JJ, Lu J, Chou TC, Scammell TE, Saper CB. Melanopsin in cells of origin of the retinohypothalamic tract. *Nat. Neurosci*. 2001; 4:1165. [PubMed: 11713469]
- Goz D, Studholme K, Lappi DA, Rollag MD, Provencio I, Morin LP. Targeted destruction of photosensitive retinal ganglion cells with a saporin conjugate alters the effects of light on mouse circadian rhythms. *PLoS One*. 2008; 3:e3153. [PubMed: 18773079]
- Granados-Fuentes D, Prolo LM, Abraham U, Herzog ED. The suprachiasmatic nucleus entrains, but does not sustain, circadian rhythmicity in the olfactory bulb. *J. Neurosci*. 2004a; 24:615–619. [PubMed: 14736846]
- Granados-Fuentes D, Saxena MT, Prolo LM, Aton SJ, Herzog ED. Olfactory bulb neurons express functional, entrainable circadian rhythms. *Eur. J. Neurosci*. 2004b; 19:898–906. [PubMed: 15009137]
- Granados-Fuentes D, Tseng A, Herzog ED. A circadian clock in the olfactory bulb controls olfactory responsivity. *J. Neurosci*. 2006; 26:12219–12225. [PubMed: 17122046]
- Gratacós FM, Brewer G. The role of AUF1 in regulated mRNA decay. *WIREs RNA*. 2010; 1:457–473. [PubMed: 21956942]
- Green CB, Douris N, Kojima S, Strayer CA, Fogerty J, Lourim D, Keller SR, Besharse JC. Loss of Nocturnin, a circadian deadenylase, confers resistance to hepatic steatosis and diet-induced obesity. *Proc. Natl. Acad. Sci. USA*. 2007; 104:9888–9893. [PubMed: 17517647]
- Green CB, Takahashi JS, Bass J. The meter of metabolism. *Cell*. 2008; 134:728–742. [PubMed: 18775307]
- Griffin EA Jr, Staknis D, Weitz CJ. Light-independent role of CRY1 and CRY2 in the mammalian circadian clock. *Science*. 1999; 286:768–771. [PubMed: 10531061]
- Guillaumond F, Dardente H, Giguere V, Cermakian N. Differential control of Bmal1 circadian transcription by REV-ERB and ROR nuclear receptors. *J. Biol. Rhythms*. 2005; 20:391–403. [PubMed: 16267379]
- Guo H, Ingolia NT, Weissman JS, Bartel DP. Mammalian microRNAs predominantly act to decrease target mRNA levels. *Nature*. 2010; 466:835–840. [PubMed: 20703300]
- Hannibal J. Neurotransmitters of the retino-hypothalamic tract. *Cell Tissue Res*. 2002; 309:73–88. [PubMed: 12111538]
- Hannibal J, Hindersson P, Ostergaard J, Georg B, Heegaard S, Larsen PJ, Fahrenkrug J. Melanopsin is expressed in PACAP-containing retinal ganglion cells of the human retinohypothalamic tract. *Invest. Ophthalmol. Vis. Sci*. 2004; 45:4202–4209. [PubMed: 15505076]
- Harada Y, Sakai M, Kurabayashi N, Hirota T, Fukada Y. Ser-557-phosphorylated mCRY2 is degraded upon synergistic phosphorylation by glycogen synthase kinase-3 β . *J. Biol. Chem*. 2005; 280:31714–31721. [PubMed: 15980066]
- Harmar AJ, Marston HM, Shen S, Spratt C, West KM, Sheward WJ, Morrison CF, Dorin JR, Piggins HD, Reubi JC, et al. The VPAC₂ receptor is essential for circadian function in the mouse suprachiasmatic nuclei. *Cell*. 2002; 109:497–508. [PubMed: 12086606]
- Hatcher NG, Atkins N Jr, Annangudi SP, Forbes AJ, Kelleher NL, Gillette MU, Sweedler JV. Mass spectrometry-based discovery of circadian peptides. *Proc. Natl. Acad. Sci. USA*. 2008; 105:12527–12532. [PubMed: 18719122]
- Hatori M, Le H, Vollmers C, Keding SR, Tanaka N, Buch T, Waisman A, Schmedt C, Jegla T, Panda S. Inducible ablation of melanopsin-expressing retinal ganglion cells reveals their central role in non-image forming visual responses. *PLoS One*. 2008; 3:e2451. [PubMed: 18545654]

- Hattar S, Liao HW, Takao M, Berson DM, Yau KW. Melanopsin-containing retinal ganglion cells: architecture, projections, and intrinsic photosensitivity. *Science*. 2002; 295:1065–1070. [PubMed: 11834834]
- Hattar S, Lucas RJ, Mrosovsky N, Thompson S, Douglas RH, Hankins MW, Lem J, Biel M, Hofmann F, Foster RG, et al. Melanopsin and rod-cone photoreceptive systems account for all major accessory visual functions in mice. *Nature*. 2003; 424:75–81.
- Hedrich, HJ.; Bullock, G., editors. *The Laboratory Mouse*. San Diego, CA: Elsevier Academic Press; 2004.
- Herzog ED, Takahashi JS, Block GD. *Clock* controls circadian period in isolated suprachiasmatic nucleus neurons. *Nat. Neurosci*. 1998; 1:708–713. [PubMed: 10196587]
- Herzog ED, Huckfeldt RM. Circadian entrainment to temperature, but not light, in the isolated suprachiasmatic nucleus. *J. Neurophysiol*. 2003; 90:763–770. [PubMed: 12660349]
- Hirayama J, Sahar S, Grimaldi B, Tamaru T, Takamatsu K, Nakahata Y, Sassone-Corsi P. CLOCK-mediated acetylation of BMAL1 controls circadian function. *Nature*. 2007; 450:1086–1090. [PubMed: 18075593]
- Hirota T, Lewis WG, Liu AC, Lee JW, Schultz PG, Kay SA. A chemical biology approach reveals period shortening of the mammalian circadian clock by specific inhibition of GSK-3 β . *Proc. Natl. Acad. Sci. USA*. 2008; 105:20746–20751. [PubMed: 19104043]
- Hirota T, Lee JW, Lewis WG, Zhang EE, Breton G, Liu X, Garcia M, Peters EC, Etchegaray JP, Traver D, et al. High-throughput chemical screen identifies a novel potent modulator of cellular circadian rhythms and reveals CKI α as a clock regulatory kinase. *PLoS Biol*. 2010; 8:e1000559. [PubMed: 21179498]
- Hoeflich KP, Luo J, Rubie EA, Tsao MS, Jin O, Woodgett JR. Requirement for glycogen synthase kinase-3 β in cell survival and NF- κ B activation. *Nature*. 2000; 406:86–90. [PubMed: 10894547]
- Hoffmann K. Die relative Wirksamkeit von Zeitgebern. *Oecologia*. 1969; 3:184–206.
- Hogenesch JB, Gu YZ, Moran SM, Shimomura K, Radcliffe LA, Takahashi JS, Bradfield CA. The basic helix-loop-helix-PAS protein MOP9 is a brain-specific heterodimeric partner of circadian and hypoxia factors. *J. Neurosci*. 2000; 20 RC83.
- Honma K, Honma S. The SCN-independent clocks, methamphetamine and food restriction. *Eur. J. Neurosci*. 2009; 30:1707–1717. [PubMed: 19878275]
- Honma S, Shirakawa T, Katsuno Y, Namihira M, Honma K. Circadian periods of single suprachiasmatic neurons in rats. *Neurosci. Lett*. 1998; 250:157–160. [PubMed: 9708856]
- Honma S, Yasuda T, Yasui A, van der Horst GT, Honma K. Circadian behavioral rhythms in *Cry1/Cry2* double-deficient mice induced by methamphetamine. *J. Biol. Rhythms*. 2008; 23:91–94. [PubMed: 18258761]
- Hughes AT, Guilding C, Lennox L, Samuels RE, McMahon DG, Piggins HD. Live imaging of altered *period1* expression in the suprachiasmatic nuclei of *Vipr2*^{-/-} mice. *J. Neurochem*. 2008; 106:1646–1657. [PubMed: 18554318]
- Hughes ME, Hogenesch JB, Kornacker K. JTK_CYCLE: an efficient nonparametric algorithm for detecting rhythmic components in genome-scale data sets. *J. Biol. Rhythms*. 2010; 25:372–380. [PubMed: 20876817]
- Iitaka C, Miyazaki K, Akaike T, Ishida N. A role for glycogen synthase kinase-3 β in the mammalian circadian clock. *J. Biol. Chem*. 2005; 280:29397–29402. [PubMed: 15972822]
- Imhof A, Becker PB. Modifications of the histone N-terminal domains. Evidence for an “epigenetic code”? *Mol. Biotechnol*. 2001; 17:1–13. [PubMed: 11280927]
- Impey S, McCorkle SR, Cha-Molstad H, Dwyer JM, Yochum GS, Boss JM, McWeeney S, Dunn JJ, Mandel G, Goodman RH. Defining the CREB regulon: a genome-wide analysis of transcription factor regulatory regions. *Cell*. 2004; 119:1041–1054. [PubMed: 15620361]
- Isojima Y, Nakajima M, Ukai H, Fujishima H, Yamada RG, Masumoto KH, Kiuchi R, Ishida M, Ukai-Tadenuma M, Minami Y, et al. CKI ϵ/δ -dependent phosphorylation is a temperature-insensitive, period-determining process in the mammalian circadian clock. *Proc. Natl. Acad. Sci. USA*. 2009; 106:15744–15749. [PubMed: 19805222]
- Iwahana E, Akiyama M, Miyakawa K, Uchida A, Kasahara J, Fukunaga K, Hamada T, Shibata S. Effect of lithium on the circadian rhythms of locomotor activity and glycogen synthase kinase-3

protein expression in the mouse suprachiasmatic nuclei. *Eur. J. Neurosci.* 2004; 19:2281–2287. [PubMed: 15090054]

- Izumo M, Johnson CH, Yamazaki S. Circadian gene expression in mammalian fibroblasts revealed by real-time luminescence reporting: temperature compensation and damping. *Proc. Natl. Acad. Sci. USA.* 2003; 100:16089–16094. [PubMed: 14657355]
- Jacob HJ. A landmark for orphan genomes? *Njat. Genet.* 1996; 13:14–15.
- Jakubcakova V, Oster H, Tamanini F, Cadenas C, Leitges M, van der Horst GT, Eichele G. Light entrainment of the mammalian circadian clock by a PRKCA-dependent posttranslational mechanism. *Neuron.* 2007; 54:831–843. [PubMed: 17553429]
- Jenuwein T, Allis CD. Translating the histone code. *Science.* 2001; 293:1074–1080. [PubMed: 11498575]
- Jin X, von Gall C, Pieschl RL, Gribkoff VK, Stehle JH, Reppert SM, Weaver DR. Targeted disruption of the mouse Mel_{1b} melatonin receptor. *Mol. Cell. Biol.* 2003; 23:1054–1060. [PubMed: 12529409]
- Kaladchibachi SA, Doble B, Anthopoulos N, Woodgett JR, Manoukian AS. Glycogen synthase kinase 3, circadian rhythms, and bipolar disorder: a molecular link in the therapeutic action of lithium. *J. Circadian Rhythms.* 2007; 5:3. [PubMed: 17295926]
- Kalamvoki M, Roizman B. Circadian CLOCK histone acetyl transferase localizes at ND10 nuclear bodies and enables herpes simplex virus gene expression. *Proc. Natl. Acad. Sci. USA.* 2010; 107:17721–17726. [PubMed: 20876123]
- Keesler GA, Camacho F, Guo Y, Virshup D, Mondadori C, Yao Z. Phosphorylation and destabilization of human period I clock protein by human casein kinase I epsilon. *NeuroReport.* 2000; 11:951–955. [PubMed: 10790862]
- King DP, Vitaterna MH, Chang AM, Dove WF, Pinto LH, Turek FW, Takahashi JS. The mouse *Clock* mutation behaves as an antimorph and maps within the *W*^{19H} deletion, distal of *Kit*. *Genetics.* 1997a; 146:1049–1060. [PubMed: 9215907]
- King DP, Zhao Y, Sangoram AM, Wilsbacher LD, Tanaka M, Antoch MP, Steeves TD, Vitaterna MH, Kornhauser JM, Lowrey PL, et al. Positional cloning of the mouse circadian *Clock* gene. *Cell.* 1997b; 89:641–653. [PubMed: 9160755]
- Kitayama Y, Nishiwaki T, Terauchi K, Kondo T. Dual KaiC-based oscillations constitute the circadian system of cyanobacteria. *Genes Dev.* 2008; 22:1513–1521. [PubMed: 18477603]
- Klein PS, Melton DA. A molecular mechanism for the effect of lithium on development. *Proc. Natl. Acad. Sci. USA.* 1996; 93:8455–8459. [PubMed: 8710892]
- Kloss B, Price JL, Saez L, Blau J, Rothenfluh A, Wesley CS, Young MW. The *Drosophila* clock gene *double-time* encodes a protein closely related to human casein kinase I ϵ . *Cell.* 1998; 94:97–107. [PubMed: 9674431]
- Knippschild U, Gocht A, Wolff S, Huber N, Lohler J, Stoter M. The casein kinase 1 family: participation in multiple cellular processes in eukaryotes. *Cell. Signal.* 2005; 17:675–689. [PubMed: 15722192]
- Ko CH, Takahashi JS. Molecular components of the mammalian circadian clock. *Hum. Mol. Genet.* 2006; 15 Spec(No 2):R271–R277. [PubMed: 16987893]
- Ko CH, Yamada YR, Welsh DK, Buhr ED, Liu AC, Zhang EE, Ralph MR, Kay SA, Forger DB, Takahashi JS. Emergence of noise-induced oscillations in the central circadian pacemaker. *PLoS Biol.* 2010; 8:e1000513. [PubMed: 20967239]
- Kojima S, Matsumoto K, Hirose M, Shimada M, Nagano M, Shigeyoshi Y, Hoshino S, Ui-Tei K, Saigo K, Green CB, et al. LARK activates posttranscriptional expression of an essential mammalian clock protein, PERIOD1. *Proc. Natl. Acad. Sci. USA.* 2007; 104:1859–1864. [PubMed: 17264215]
- Kojima S, Gatfield D, Esau CC, Green CB. MicroRNA-122 modulates the rhythmic expression profile of the circadian deadenylase *Nocturnin* in mouse liver. *PLoS One.* 2010; 5:e11264. [PubMed: 20582318]
- Kojima S, Shingle DL, Green CB. Post-transcriptional control of circadian rhythms. *J. Cell Sci.* 2011; 124:311–320. [PubMed: 21242310]

- Konopka RJ, Benzer S. Clock mutants of *Drosophila melanogaster*. Proc. Natl. Acad. Sci. USA. 1971; 68:2112–2116. [PubMed: 5002428]
- Kornitzer D, Ciechanover A. Modes of regulation of ubiquitin-mediated protein degradation. J. Cell. Physiol. 2000; 182:1–11. [PubMed: 10567911]
- Kornmann B, Schaad O, Bujard H, Takahashi JS, Schibler U. System-driven and oscillator-dependent circadian transcription in mice with a conditionally active liver clock. PLoS Biol. 2007; 5:e34. [PubMed: 17298173]
- Kramer A, Yang FC, Snodgrass P, Li X, Scammell TE, Davis FC, Weitz CJ. Regulation of daily locomotor activity and sleep by hypothalamic EGF receptor signaling. Science. 2001; 294:2511–2515. [PubMed: 11752569]
- Kramer A, Yang FC, Kraves S, Weitz CJ. A screen for secreted factors of the suprachiasmatic nucleus. Methods Enzymol. 2005; 393:645–663. [PubMed: 15817317]
- Kraves S, Weitz CJ. A role for cardiotrophin-like cytokine in the circadian control of mammalian locomotor activity. Nat. Neurosci. 2006; 9:212–219. [PubMed: 16429135]
- Krutzfeldt J, Rajewsky N, Braich R, Rajeev KG, Tuschl T, Manoharan M, Stoffel M. Silencing of microRNAs in vivo with 'antagomirs'. Nature. 2005; 438:685–689. [PubMed: 16258535]
- Kumaki Y, Ukai-Tadenuma M, Uno KD, Nishio J, Masumoto KH, Nagano M, Komori T, Shigeyoshi Y, Hogenesch JB, Ueda HR. Analysis and synthesis of high-amplitude *Cis*-elements in the mammalian circadian clock. Proc. Natl. Acad. Sci. USA. 2008; 105:14946–14951. [PubMed: 18815372]
- Kume K, Zylka MJ, Sriram S, Shearman LP, Weaver DR, Jin X, Maywood ES, Hastings MH, Reppert SM. mCRY1 and mCRY2 are essential components of the negative limb of the circadian clock feedback loop. Cell. 1999; 98:193–205. [PubMed: 10428031]
- Kurabayashi N, Hirota T, Sakai M, Sanada K, Fukada Y. DYRK1A and glycogen synthase kinase 3 β , a dual-kinase mechanism directing proteasomal degradation of CRY2 for circadian timekeeping. Mol. Cell. Biol. 2010; 30:1757–1768. [PubMed: 20123978]
- Lamia KA, Sachdeva UM, DiTacchio L, Williams EC, Alvarez JG, Egan DF, Vasquez DS, Juguilon H, Panda S, Shaw RJ, et al. AMPK regulates the circadian clock by cryptochrome phosphorylation and degradation. Science. 2009; 326:437–440. [PubMed: 19833968]
- Le Minh N, Damiola F, Tronche F, Schutz G, Schibler U. Glucocorticoid hormones inhibit food-induced phase-shifting of peripheral circadian oscillators. EMBO J. 2001; 20:7128–7136. [PubMed: 11742989]
- Lee C, Etchegaray JP, Cagampang FR, Loudon AS, Reppert SM. Posttranslational mechanisms regulate the mammalian circadian clock. Cell. 2001; 107:855–867. [PubMed: 11779462]
- Lee H, Chen R, Lee Y, Yoo S, Lee C. Essential roles of CKI δ and CKI ϵ in the mammalian circadian clock. Proc. Natl. Acad. Sci. USA. 2009; 106:21359–21364. [PubMed: 19948962]
- Lee J, Lee Y, Lee MJ, Park E, Kang SH, Chung CH, Lee KH, Kim K. Dual modification of BMAL1 by SUMO2/3 and ubiquitin promotes circadian activation of the CLOCK/BMAL1 complex. Mol. Cell. Biol. 2008; 28:6056–6065. [PubMed: 18644859]
- LeSauter J, Silver R. Lithium lengthens the period of circadian rhythms in lesioned hamsters bearing SCN grafts. Biol. Psychiatry. 1993; 34:75–83. [PubMed: 8251024]
- Li JD, Hu WP, Boehmer L, Cheng MY, Lee AG, Jilek A, Siegel JM, Zhou QY. Attenuated circadian rhythms in mice lacking the prokineticin 2 gene. J. Neurosci. 2006; 26:11615–11623. [PubMed: 17093083]
- Li S, Liu C, Li N, Hao T, Han T, Hill DE, Vidal M, Lin JD. Genome-wide coactivation analysis of PGC-1 α identifies BAF60a as a regulator of hepatic lipid metabolism. Cell Metab. 2008; 8:105–117. [PubMed: 18680712]
- Liu AC, Lewis WG, Kay SA. Mammalian circadian signaling networks and therapeutic targets. Nat. Chem. Biol. 2007a; 3:630–639. [PubMed: 17876320]
- Liu AC, Welsh DK, Ko CH, Tran HG, Zhang EE, Priest AA, Buhr ED, Singer O, Meeker K, Verma IM, et al. Intercellular coupling confers robustness against mutations in the SCN circadian clock network. Cell. 2007b; 129:605–616. [PubMed: 17482552]

- Liu AC, Tran HG, Zhang EE, Priest AA, Welsh DK, Kay SA. Redundant function of REV-ERB α and β and non-essential role for Bmal1 cycling in transcriptional regulation of intracellular circadian rhythms. *PLoS Genet.* 2008; 4:e1000023. [PubMed: 18454201]
- Liu C, Weaver DR, Jin X, Shearman LP, Pieschl RL, Gribkoff VK, Reppert SM. Molecular dissection of two distinct actions of melatonin on the suprachiasmatic circadian clock. *Neuron.* 1997a; 19:91–102. [PubMed: 9247266]
- Liu C, Weaver DR, Strogatz SH, Reppert SM. Cellular construction of a circadian clock: period determination in the suprachiasmatic nuclei. *Cell.* 1997b; 91:855–860. [PubMed: 9413994]
- Liu C, Li S, Liu T, Borjigin J, Lin JD. Transcriptional coactivator PGC-1 α integrates the mammalian clock and energy metabolism. *Nature.* 2007c; 447:477–481. [PubMed: 17476214]
- Lopez-Molina L, Conquet F, Dubois-Dauphin M, Schibler U. The DBP gene is expressed according to a circadian rhythm in the suprachiasmatic nucleus and influences circadian behavior. *EMBO J.* 1997; 16:6762–6771. [PubMed: 9362490]
- Lowrey PL, Shimomura K, Antoch MP, Yamazaki S, Zemenides PD, Ralph MR, Menaker M, Takahashi JS. Positional syntenic cloning and functional characterization of the mammalian circadian mutation *tau*. *Science.* 2000; 288:483–492. [PubMed: 10775102]
- Lowrey PL, Takahashi JS. Genetics of the mammalian circadian system: Photic entrainment, circadian pacemaker mechanisms, and posttranslational regulation. *Annu. Rev. Genet.* 2000; 34:533–562. [PubMed: 11092838]
- Lowrey PL, Takahashi JS. Mammalian circadian biology: elucidating genome-wide levels of temporal organization. *Annu. Rev. Genom. Hum. Genet.* 2004; 5:407–441.
- Lucas RJ, Hattar S, Takao M, Berson DM, Foster RG, Yau KW. Diminished pupillary light reflex at high irradiances in melanopsin-knockout mice. *Science.* 2003; 299:245–247. [PubMed: 12522249]
- MacAulay K, Doble BW, Patel S, Hansotia T, Sinclair EM, Drucker DJ, Nagy A, Woodgett JR. Glycogen synthase kinase 3 α -specific regulation of murine hepatic glycogen metabolism. *Cell Metab.* 2007; 6:329–337. [PubMed: 17908561]
- Maier B, Wendt S, Vanselow JT, Wallach T, Reischl S, Oehmke S, Schlosser A, Kramer A. A large-scale functional RNAi screen reveals a role for CK2 in the mammalian circadian clock. *Genes Dev.* 2009; 23:708–718. [PubMed: 19299560]
- Martinek S, Inonog S, Manoukian AS, Young MW. A role for the segment polarity gene shaggy/GSK-3 in the *Drosophila* circadian clock. *Cell.* 2001; 105:769–779. [PubMed: 11440719]
- Masana MI, Sumaya IC, Becker-Andre M, Dubocovich ML. Behavioral characterization and modulation of circadian rhythms by light and melatonin in C3H/HeN mice homozygous for the ROR β knockout. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 2007; 292:R2357–R2367. [PubMed: 17303680]
- Maury E, Ramsey KM, Bass J. Circadian rhythms and metabolic syndrome: from experimental genetics to human disease. *Circ. Res.* 2010; 106:447–462. [PubMed: 20167942]
- Maywood ES, Reddy AB, Wong GK, O'Neill JS, O'Brien JA, McMahon DG, Hattar AJ, Okamura H, Hastings MH. Synchronization and maintenance of timekeeping in suprachiasmatic circadian clock cells by neuropeptidergic signaling. *Curr. Biol.* 2006; 16:599–605. [PubMed: 16546085]
- McDearmon EL, Patel KN, Ko CH, Walisser JA, Schook AC, Chong JL, Wilsbacher LD, Song EJ, Hong HK, Bradfield CA, et al. Dissecting the functions of the mammalian clock protein BMAL1 by tissue-specific rescue in mice. *Science.* 2006; 314:1304–1308. [PubMed: 17124323]
- Meggio F, Pinna LA. One-thousand-and-one substrates of protein kinase CK2? *FASEB J.* 2003; 17:349–368. [PubMed: 12631575]
- Melyan Z, Tarttelin EE, Bellingham J, Lucas RJ, Hankins MW. Addition of human melanopsin renders mammalian cells photoresponsive. *Nature.* 2005; 433:741–745. [PubMed: 15674244]
- Meng QJ, Logunova L, Maywood ES, Gallego M, Lebiecki J, Brown TM, Sladek M, Semikhodskii AS, Glossop NR, Piggins HD, et al. Setting clock speed in mammals: the CK1 ϵ *tau* mutation in mice accelerates circadian pacemakers by selectively destabilizing PERIOD proteins. *Neuron.* 2008; 58:78–88. [PubMed: 18400165]
- Meng QJ, Maywood ES, Bechtold DA, Lu WQ, Li J, Gibbs JE, Dupre SM, Chesham JE, Rajamohan F, Knafels J, et al. Entrainment of disrupted circadian behavior through inhibition of casein

- kinase 1 (CK1) enzymes. *Proc. Natl. Acad. Sci. USA.* 2010; 107:15240–15245. [PubMed: 20696890]
- Michel S, Itri J, Han JH, Gnietczynski K, Colwell CS. Regulation of glutamatergic signalling by PACAP in the mammalian suprachiasmatic nucleus. *BMC Neurosci.* 2006; 7:15. [PubMed: 16483357]
- Miller BH, McDearmon EL, Panda S, Hayes KR, Zhang J, Andrews JL, Antoch MP, Walker JR, Esser KA, Hogenesch JB, et al. Circadian and CLOCK-controlled regulation of the mouse transcriptome and cell proliferation. *Proc. Natl. Acad. Sci. USA.* 2007; 104:3342–3347. [PubMed: 17360649]
- Mitsui S, Yamaguchi S, Matsuo T, Ishida Y, Okamura H. Antagonistic role of E4BP4 and PAR proteins in the circadian oscillatory mechanism. *Genes Dev.* 2001; 15:995–1006. [PubMed: 11316793]
- Mohawk JA, Baer ML, Menaker M. The methamphetamine-sensitive circadian oscillator does not employ canonical clock genes. *Proc. Natl. Acad. Sci. USA.* 2009; 106:3519–3524. [PubMed: 19204282]
- Moore MJ. From birth to death: the complex lives of eukaryotic mRNAs. *Science.* 2005; 309:1514–1518. [PubMed: 16141059]
- Moore RY, Eichler VB. Loss of a circadian adrenal corticosterone rhythm following suprachiasmatic lesions in the rat. *Brain Res.* 1972; 42:201–206. [PubMed: 5047187]
- Moore RY, Lenn NJ. A retinohypothalamic projection in the rat. *J. Comp. Neurol.* 1972; 146:1–14. [PubMed: 4116104]
- Moore RY, Speh JC, Card JP. The retinohypothalamic tract originates from a distinct subset of retinal ganglion cells. *J. Comp. Neurol.* 1995; 352:351–366. [PubMed: 7706557]
- Morin LP, Allen CN. The circadian visual system, 2005. *Brain Res. Rev.* 2006; 51:1–60. [PubMed: 16337005]
- Muller B, Grossniklaus U. Model organisms--A historical perspective. *J. Proteomics.* 2010; 73:2054–2063. [PubMed: 20727995]
- Nader N, Chrousos GP, Kino T. Circadian rhythm transcription factor CLOCK regulates the transcriptional activity of the glucocorticoid receptor by acetylating its hinge region lysine cluster: potential physiological implications. *FASEB J.* 2009; 23:1572–1583. [PubMed: 19141540]
- Nagel R, Clijsters L, Agami R. The miRNA-192/194 cluster regulates the *Period* gene family and the circadian clock. *FEBS J.* 2009; 276:5447–5455. [PubMed: 19682069]
- Nagoshi E, Saini C, Bauer C, Laroche T, Naef F, Schibler U. Circadian gene expression in individual fibroblasts; cell-autonomous and self-sustained oscillators pass time to daughter cells. *Cell.* 2004; 119:693–705. [PubMed: 15550250]
- Nagy, A.; Gertsenstein, M.; Vintersten, K.; Behringer, R., editors. *Manipulating the Mouse Embryo: A Laboratory Manual.* Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press; 2003.
- Nakahata Y, Kaluzova M, Grimaldi B, Sahar S, Hirayama J, Chen D, Guarente LP, Sassone-Corsi P. The NAD⁺-dependent deacetylase SIRT1 modulates CLOCK-mediated chromatin remodeling and circadian control. *Cell.* 2008; 134:329–340. [PubMed: 18662547]
- Nakahata Y, Sahar S, Astarita G, Kaluzova M, Sassone-Corsi P. Circadian control of the NAD⁺ salvage pathway by CLOCK-SIRT1. *Science.* 2009; 324:654–657. [PubMed: 19286518]
- Nakajima M, Imai K, Ito H, Nishiwaki T, Murayama Y, Iwasaki H, Oyama T, Kondo T. Reconstitution of circadian oscillation of cyanobacterial KaiC phosphorylation in vitro. *Science.* 2005; 308:414–415. [PubMed: 15831759]
- Nakashima A, Kawamoto T, Honda KK, Ueshima T, Noshiro M, Iwata T, Fujimoto K, Kubo H, Honma S, Yorioka N, et al. DEC1 modulates the circadian phase of clock gene expression. *Mol. Cell. Biol.* 2008; 28:4080–4092. [PubMed: 18411297]
- Nandi D, Tahiliani P, Kumar A, Chandu D. The ubiquitin-proteasome system. *J. Biosci.* 2006; 31:137–155. [PubMed: 16595883]
- Nelson RJ, Zucker I. Absence of extraocular photoreception in diurnal and nocturnal rodents exposed to direct sunlight. *Comp. Biochem. Physiol.* 1981; 69A:145–148.

- O'Neill JS, Reddy AB. Circadian clocks in human red blood cells. *Nature*. 2011; 469:498–503. [PubMed: 21270888]
- O'Neill JS, van Ooijen G, Dixon LE, Troein C, Corellou F, Bouget FY, Reddy AB, Millar AJ. Circadian rhythms persist without transcription in a eukaryote. *Nature*. 2011; 469:554–558. [PubMed: 21270895]
- Ohno T, Onishi Y, Ishida N. A novel E4BP4 element drives circadian expression of *mPeriod2*. *Nucleic Acids Res*. 2007; 35:648–655. [PubMed: 17182630]
- Ohsaki K, Oishi K, Kozono Y, Nakayama K, Nakayama KI, Ishida N. The role of β -TrCP1 and β -TrCP2 in circadian rhythm generation by mediating degradation of clock protein PER2. *J. Biochem. Tokyo*. 2008; 144:609–618. [PubMed: 18782782]
- Oishi K, Miyazaki K, Kadota K, Kikuno R, Nagase T, Atsumi G, Ohkura N, Azama T, Mesaki M, Yukimasa S, et al. Genome-wide expression analysis of mouse liver reveals CLOCK-regulated circadian output genes. *J. Biol. Chem*. 2003; 278:41519–41527. [PubMed: 12865428]
- Okamura H, Miyake S, Sumi Y, Yamaguchi S, Yasui A, Muijtjens M, Hoeijmakers JH, van der Horst GT. Photic induction of *mPer1* and *mPer2* in *Cry*-deficient mice lacking a biological clock. *Science*. 1999; 286:2531–2534. [PubMed: 10617474]
- Oster H, Yasui A, van der Horst GT, Albrecht U. Disruption of *mCry2* restores circadian rhythmicity in *mPer2* mutant mice. *Genes Dev*. 2002; 16:2633–2638. [PubMed: 12381662]
- Oster H, Baeriswyl S, Van Der Horst GT, Albrecht U. Loss of circadian rhythmicity in aging *mPer1^{-/-} mCry2^{-/-}* mutant mice. *Genes Dev*. 2003; 17:1366–1379. [PubMed: 12782655]
- Palkova M, Sigmund L, Erkert HG. Effect of ambient temperature on the circadian activity rhythm in common marmosets, *Callithrix j. jacchus* (primates). *Chronobiol. Int*. 1999; 16:149–161. [PubMed: 10219487]
- Panda S, Antoch MP, Miller BH, Su AI, Schook AB, Straume M, Schultz PG, Kay SA, Takahashi JS, Hogenesch JB. Coordinated transcription of key pathways in the mouse by the circadian clock. *Cell*. 2002a; 109:307–320. [PubMed: 12015981]
- Panda S, Sato TK, Castrucci AM, Rollag MD, DeGrip WJ, Hogenesch JB, Provencio I, Kay SA. Melanopsin (*Opn4*) requirement for normal light-induced circadian phase shifting. *Science*. 2002b; 298:2213–2216. [PubMed: 12481141]
- Panda S, Provencio I, Tu DC, Pires SS, Rollag MD, Castrucci AM, Pletcher MT, Sato TK, Wiltshire T, Andahazy M, et al. Melanopsin is required for non-image-forming photic responses in blind mice. *Science*. 2003; 301:525–527. [PubMed: 12829787]
- Panda S, Nayak SK, Campo B, Walker JR, Hogenesch JB, Jegla T. Illumination of the melanopsin signaling pathway. *Science*. 2005; 307:600–604. [PubMed: 15681390]
- Partch CL, Shields KF, Thompson CL, Selby CP, Sancar A. Posttranslational regulation of the mammalian circadian clock by cryptochrome and protein phosphatase 5. *Proc. Natl. Acad. Sci. USA*. 2006; 103:10467–10472. [PubMed: 16790549]
- Pezuk P, Mohawk JA, Yoshikawa T, Sellix MT, Menaker M. Circadian organization is governed by extra-SCN pacemakers. *J. Biol. Rhythms*. 2010; 25:432–441. [PubMed: 21135159]
- Pittendrigh CS. Temporal organization: reflections of a Darwinian clock-watcher. *Annu. Rev. Physiol*. 1993; 55:16–54. [PubMed: 8466172]
- Preitner N, Damiola F, Lopez-Molina L, Zakany J, Duboule D, Albrecht U, Schibler U. The orphan nuclear receptor REV-ERBa controls circadian transcription within the positive limb of the mammalian circadian oscillator. *Cell*. 2002; 110:251–260. [PubMed: 12150932]
- Price JL, Blau J, Rothenfluh A, Abodeely M, Kloss B, Young MW. *double-time* is a novel *Drosophila* clock gene that regulates PERIOD protein accumulation. *Cell*. 1998; 94:83–95. [PubMed: 9674430]
- Prolo LM, Takahashi JS, Herzog ED. Circadian rhythm generation and entrainment in astrocytes. *J. Neurosci*. 2005; 25:404–408. [PubMed: 15647483]
- Provencio I, Jiang G, De Grip WJ, Hayes WP, Rollag MD. Melanopsin: An opsin in melanophores, brain, and eye. *Proc. Natl. Acad. Sci. USA*. 1998; 95:340–345. [PubMed: 9419377]
- Qiu X, Kumbalasiri T, Carlson SM, Wong KY, Krishna V, Provencio I, Berson DM. Induction of photosensitivity by heterologous expression of melanopsin. *Nature*. 2005; 433:745–749. [PubMed: 15674243]

- Quintero JE, Kuhlman SJ, McMahon DG. The biological clock nucleus: a multiphasic oscillator network regulated by light. *J. Neurosci.* 2003; 23:8070–8076. [PubMed: 12954869]
- Ralph MR, Menaker M. A mutation of the circadian system in golden hamsters. *Science.* 1988; 241:1225–1227. [PubMed: 3413487]
- Ralph MR, Foster RG, Davis FC, Menaker M. Transplanted suprachiasmatic nucleus determines circadian period. *Science.* 1990; 247:975–978. [PubMed: 2305266]
- Ramsey KM, Yoshino J, Brace CS, Abrassart D, Kobayashi Y, Marcheva B, Hong HK, Chong JL, Buhr ED, Lee C, et al. Circadian clock feedback cycle through NAMPT-mediated NAD⁺ biosynthesis. *Science.* 2009; 324:651–654. [PubMed: 19299583]
- Rana TM. Illuminating the silence: understanding the structure and function of small RNAs. *Nat. Rev. Mol. Cell Biol.* 2007; 8:23–36. [PubMed: 17183358]
- Reddy AB, Karp NA, Maywood ES, Sage EA, Deery M, O'Neill JS, Wong GK, Chesham J, Odell M, Lilley KS, et al. Circadian orchestration of the hepatic proteome. *Curr. Biol.* 2006; 16:1107–1115. [PubMed: 16753565]
- Refinetti R, Menaker M. The circadian rhythm of body temperature. *Physiol. Behav.* 1992; 51:613–637. [PubMed: 1523238]
- Reick M, Garcia JA, Dudley C, McKnight SL. NPAS2: an analog of *Clock* operative in the mammalian forebrain. *Science.* 2001; 293:506–509. [PubMed: 11441147]
- Reinke H, Saini C, Fleury-Olela F, Dibner C, Benjamin II, Schibler U. Differential display of DNA-binding proteins reveals heat-shock factor 1 as a circadian transcription factor. *Genes Dev.* 2008; 22:331–345. [PubMed: 18245447]
- Reischl S, Vanselow K, Westermark PO, Thierfelder N, Maier B, Herzog H, Kramer A. Beta-TrCP1-mediated degradation of PERIOD2 is essential for circadian dynamics. *J. Biol. Rhythms.* 2007; 22:375–386. [PubMed: 17876059]
- Reppert SM. A clockwork explosion! *Neuron.* 1998; 21:1–4. [PubMed: 9697845]
- Reppert SM, Weaver DR. Coordination of circadian timing in mammals. *Nature.* 2002; 418:935–941. [PubMed: 12198538]
- Reyes BA, Pendergast JS, Yamazaki S. Mammalian peripheral circadian oscillators are temperature compensated. *J. Biol. Rhythms.* 2008; 23:95–98. [PubMed: 18258762]
- Ripperger JA, Mrosovsky M. Perfect timing: Epigenetic regulation of the circadian clock. *FEBS Lett.* 2011; 585:1406–1411. [PubMed: 21536041]
- Rivers A, Gietzen KF, Vielhaber E, Virshup DM. Regulation of casein kinase I ϵ and casein kinase I δ by an *in vivo* futile phosphorylation cycle. *J. Biol. Chem.* 1998; 273:15980–15984. [PubMed: 9632646]
- Robles MS, Boyault C, Knutti D, Padmanabhan K, Weitz CJ. Identification of RACK1 and protein kinase C α as integral components of the mammalian circadian clock. *Science.* 2010; 327:463–466. [PubMed: 20093473]
- Rossner MJ, Oster H, Wichert SP, Reinecke L, Wehr MC, Reinecke J, Eichele G, Taneja R, Nave KA. Disturbed clockwork resetting in Sharp-1 and Sharp-2 single and double mutant mice. *PLoS One.* 2008; 3:e2762. [PubMed: 18648504]
- Ruby NF, Heller HC. Temperature sensitivity of the suprachiasmatic nucleus of ground squirrels and rats *in vitro*. *J. Biol. Rhythms.* 1996; 11:126–136. [PubMed: 8744240]
- Ruby NF, Burns DE, Heller HC. Circadian rhythms in the suprachiasmatic nucleus are temperature-compensated and phase-shifted by heat pulses *in vitro*. *J. Neurosci.* 1999; 19:8630–8636. [PubMed: 10493763]
- Ruby NF, Brennan TJ, Xie X, Cao V, Franken P, Heller HC, O'Hara BF. Role of melanopsin in circadian responses to light. *Science.* 2002; 298:2211–2213. [PubMed: 12481140]
- Rutter J, Reick M, Wu LC, McKnight SL. Regulation of *Clock* and NPAS2 DNA binding by the redox state of NAD cofactors. *Science.* 2001; 293:510–514. [PubMed: 11441146]
- Rutter J, Reick M, McKnight SL. Metabolism and the control of circadian rhythms. *Annu. Rev. Biochem.* 2002; 71:307–331. [PubMed: 12045099]
- Ryves WJ, Harwood AJ. Lithium inhibits glycogen synthase kinase-3 by competition for magnesium. *Biochem. Biophys. Res. Commun.* 2001; 280:720–725. [PubMed: 11162580]

- Sahar S, Zocchi L, Kinoshita C, Borrelli E, Sassone-Corsi P. Regulation of BMAL1 protein stability and circadian function by GSK3 β -mediated phosphorylation. *PLoS One*. 2010; 5:e8561. [PubMed: 20049328]
- Sanada K, Okano T, Fukada Y. Mitogen-activated protein kinase phosphorylates and negatively regulates basic helix-loop-helix-PAS transcription factor BMAL1. *J. Biol. Chem*. 2002; 277:267–271. [PubMed: 11687575]
- Sasaki M, Yoshitane H, Du NH, Okano T, Fukada Y. Preferential inhibition of BMAL2-CLOCK activity by PER2 reemphasizes its negative role and a positive role of BMAL2 in the circadian transcription. *J. Biol. Chem*. 2009; 284:25149–25159. [PubMed: 19605937]
- Sato TK, Panda S, Miraglia LJ, Reyes TM, Rudic RD, McNamara P, Naik KA, FitzGerald GA, Kay SA, Hogenesch JB. A functional genomics strategy reveals Rora as a component of the mammalian circadian clock. *Neuron*. 2004; 43:527–537. [PubMed: 15312651]
- Sato TK, Yamada RG, Ukai H, Baggs JE, Miraglia LJ, Kobayashi TJ, Welsh DK, Kay SA, Ueda HR, Hogenesch JB. Feedback repression is required for mammalian circadian clock function. *Nat. Genet*. 2006; 38:312–319. [PubMed: 16474406]
- Seggie J, Werstiuik ES, Grotta L. Effect of chronic lithium treatment on twenty four hour variation in plasma and red blood cell lithium and sodium concentrations, drinking behavior, body weight, kidney weight, and corticosterone levels. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry*. 1982; 6:455–458.
- Shearman LP, Zylka MJ, Weaver DR, Kolakowski LF Jr, Reppert SM. Two *period* homologs: circadian expression and photic regulation in the suprachiasmatic nuclei. *Neuron*. 1997; 19:1261–1269. [PubMed: 9427249]
- Shearman LP, Jin X, Lee C, Reppert SM, Weaver DR. Targeted disruption of the *mPer3* gene: subtle effects on circadian clock function. *Mol. Cell. Biol*. 2000a; 20:6269–6275. [PubMed: 10938103]
- Shearman LP, Sriram S, Weaver DR, Maywood ES, Chaves I, Zheng B, Kume K, Lee CC, van der Horst GT, Hastings MH, et al. Interacting molecular loops in the mammalian circadian clock. *Science*. 2000b; 288:1013–1019. [PubMed: 10807566]
- Shi S, Hida A, McGuinness OP, Wasserman DH, Yamazaki S, Johnson CH. Circadian clock gene *Bmal1* is not essential; functional replacement with its paralog, *Bmal2*. *Curr. Biol*. 2010; 20:316–321. [PubMed: 20153195]
- Shigeyoshi Y, Taguchi K, Yamamoto S, Takekida S, Yan L, Tei H, Moriya T, Shibata S, Loros JJ, Dunlap JC, et al. Light-induced resetting of a mammalian circadian clock is associated with rapid induction of the *mPer1* transcript. *Cell*. 1997; 91:1043–1053. [PubMed: 9428526]
- Shim HS, Kim H, Lee J, Son GH, Cho S, Oh TH, Kang SH, Seen DS, Lee KH, Kim K. Rapid activation of CLOCK by Ca²⁺-dependent protein kinase C mediates resetting of the mammalian circadian clock. *EMBO Rep*. 2007; 8:366–371. [PubMed: 17347670]
- Shirogane T, Jin J, Ang XL, Harper JW. SCF ^{β} -TRCP controls clock-dependent transcription via casein kinase 1-dependent degradation of the mammalian period-1 (Per1) protein. *J. Biol. Chem*. 2005; 280:26863–26872. [PubMed: 15917222]
- Siepkha SM, Takahashi JS. Forward genetic screens to identify circadian rhythm mutants in mice. *Methods Enzymol*. 2005; 393:219–229. [PubMed: 15817290]
- Siepkha SM, Yoo SH, Park J, Song W, Kumar V, Hu Y, Lee C, Takahashi JS. Circadian mutant *Overtime* reveals F-box protein FBXL3 regulation of *Cryptochrome* and *Period* gene expression. *Cell*. 2007; 129:1011–1023. [PubMed: 17462724]
- Silver, LM. "Mouse Genetics: Concepts and Applications.". New York, NY: Oxford University Press; 1995.
- Silver R, LeSauter J, Tresco PA, Lehman MN. A diffusible coupling signal from the transplanted suprachiasmatic nucleus controlling circadian locomotor rhythms. *Nature*. 1996; 382:810–813. [PubMed: 8752274]
- Spengler ML, Kuropatwinski KK, Schumer M, Antoch MP. A serine cluster mediates BMAL1-dependent CLOCK phosphorylation and degradation. *Cell Cycle*. 2009; 8:4138–4146. [PubMed: 19946213]

- Sprouse J, Reynolds L, Kleiman R, Tate B, Swanson TA, Pickard GE. Chronic treatment with a selective inhibitor of casein kinase I δ/ϵ yields cumulative phase delays in circadian rhythms. *Psychopharmacology (Berl)*. 2010; 210:569–576. [PubMed: 20407760]
- Staiger D, Koster T. Spotlight on post-transcriptional control in the circadian system. *Cell. Mol. Life Sci*. 2011; 68:71–83. [PubMed: 20803230]
- Stambolic V, Ruel L, Woodgett JR. Lithium inhibits glycogen synthase kinase-3 activity and mimics wingless signalling in intact cells. *Curr. Biol*. 1996; 6:1664–1668. [PubMed: 8994831]
- Stephan FK, Zucker I. Circadian rhythms in drinking behavior and locomotor activity of rats are eliminated by hypothalamic lesions. *Proc. Natl. Acad. Sci. USA*. 1972; 69:1583–1586. [PubMed: 4556464]
- Stokkan KA, Yamazaki S, Tei H, Sakaki Y, Menaker M. Entrainment of the circadian clock in the liver by feeding. *Science*. 2001; 291:490–493. [PubMed: 11161204]
- Storch KF, Lipan O, Leykin I, Viswanathan N, Davis FC, Wong WH, Weitz CJ. Extensive and divergent circadian gene expression in liver and heart. *Nature*. 2002; 417:78–83. [PubMed: 11967526]
- Storch KF, Weitz CJ. Daily rhythms of food-anticipatory behavioral activity do not require the known circadian clock. *Proc. Natl. Acad. Sci. USA*. 2009; 106:6808–6813. [PubMed: 19366674]
- Strahl BD, Allis CD. The language of covalent histone modifications. *Nature*. 2000; 403:41–45. [PubMed: 10638745]
- Sweeney BM, Hastings JW. Effects of temperature upon diurnal rhythms. *Cold Spring Harbor Symp. Quant. Biol*. 1960; 25:87–104. [PubMed: 13774256]
- Takahashi JS, Pinto LH, Vitaterna MH. Forward and reverse genetic approaches to behavior in the mouse. *Science*. 1994; 264:1724–1733. [PubMed: 8209253]
- Takahashi, JS.; Turek, FW.; Moore, RY., editors. *Handbook of Behavioral Neurobiology: Circadian Clocks*. Vol. Vol. 12. New York: Kluwer Acad./Plenum Publ.; 2001.
- Takahashi JS, Hong HK, Ko CH, McDearmon EL. The genetics of mammalian circadian order and disorder: implications for physiology and disease. *Nat. Rev. Genet*. 2008; 9:764–775. [PubMed: 18802415]
- Tamaru T, Hirayama J, Isojima Y, Nagai K, Norioka S, Takamatsu K, Sassone-Corsi P. CK2 α phosphorylates BMAL1 to regulate the mammalian clock. *Nat. Struct. Mol. Biol*. 2009; 16:446–448. [PubMed: 19330005]
- Thresher RJ, Vitaterna MH, Miyamoto Y, Kazantsev A, Hsu DS, Petit C, Selby CP, Dawut L, Smithies O, Takahashi JS, et al. Role of mouse cryptochrome blue-light photoreceptor in circadian photoresponses. *Science*. 1998; 282:1490–1494. [PubMed: 9822380]
- Tischkau SA, Mitchell JW, Pace LA, Barnes JW, Barnes JA, Gillette MU. Protein kinase G type II is required for night-to-day progression of the mammalian circadian clock. *Neuron*. 2004; 43:539–549. [PubMed: 15312652]
- Tomita J, Nakajima M, Kondo T, Iwasaki H. No transcription-translation feedback in circadian rhythm of KaiC phosphorylation. *Science*. 2005; 307:251–254. [PubMed: 15550625]
- Tosini G, Menaker M. Circadian rhythms in cultured mammalian retina. *Science*. 1996; 272:419–421. [PubMed: 8602533]
- Tosini G, Menaker M. The *tau* mutation affects temperature compensation of hamster retinal circadian oscillators. *NeuroReport*. 1998; 9:1001–1005. [PubMed: 9601657]
- Travnickova-Bendova Z, Cermakian N, Reppert SM, Sassone-Corsi P. Bimodal regulation of *mPeriod* promoters by CREB-dependent signaling and CLOCK/BMAL1 activity. *Proc. Natl. Acad. Sci. USA*. 2002; 99:7728–7733. [PubMed: 12032351]
- Triqueneaux G, Thenot S, Kakizawa T, Antoch MP, Safi R, Takahashi JS, Delaunay F, Laudet V. The orphan receptor Rev-erb α gene is a target of the circadian clock pacemaker. *J. Mol. Endocrinol*. 2004; 33:585–608. [PubMed: 15591021]
- Tsuchiya Y, Akashi M, Nishida E. Temperature compensation and temperature resetting of circadian rhythms in mammalian cultured fibroblasts. *Genes Cells*. 2003; 8:713–720. [PubMed: 12875656]
- Tsuchiya Y, Akashi M, Matsuda M, Goto K, Miyata Y, Node K, Nishida E. Involvement of the protein kinase CK2 in the regulation of mammalian circadian rhythms. *Sci. Signal*. 2009; 2 ra26.

- Ueda HR, Hayashi S, Chen W, Sano M, Machida M, Shigeyoshi Y, Iino M, Hashimoto S. System-level identification of transcriptional circuits underlying mammalian circadian clocks. *Nat. Genet.* 2005; 37:187–192. [PubMed: 15665827]
- Ukai-Tadenuma M, Kasukawa T, Ueda HR. Proof-by-synthesis of the transcriptional logic of mammalian circadian clocks. *Nat. Cell Biol.* 2008; 10:1154–1163. [PubMed: 18806789]
- van der Horst GT, Muijtjens M, Kobayashi K, Takano R, Kanno S, Takao M, de Wit J, Verkerk A, Eker AP, van Leenen D, et al. Mammalian *Cry1* and *Cry2* are essential for maintenance of circadian rhythms. *Nature.* 1999; 398:627–630. [PubMed: 10217146]
- Vanselow K, Vanselow JT, Westermarck PO, Reischl S, Maier B, Korte T, Herrmann A, Herzel H, Schlosser A, Kramer A. Differential effects of PER2 phosphorylation: molecular basis for the human familial advanced sleep phase syndrome (FASPS). *Genes Dev.* 2006; 20:2660–2672. [PubMed: 16983144]
- Vaquero A, Scher M, Lee D, Erdjument-Bromage H, Tempst P, Reinberg D. Human SirT1 interacts with histone H1 and promotes formation of facultative heterochromatin. *Mol. Cell.* 2004; 16:93–105. [PubMed: 15469825]
- Vielhaber E, Eide E, Rivers A, Gao ZH, Virshup DM. Nuclear entry of the circadian regulator mPER1 is controlled by mammalian casein kinase I ϵ . *Mol. Cell. Biol.* 2000; 20:4888–4899. [PubMed: 10848614]
- Vitaterna MH, King DP, Chang AM, Kornhauser JM, Lowrey PL, McDonald JD, Dove WF, Pinto LH, Turek FW, Takahashi JS. Mutagenesis and mapping of a mouse gene, *Clock*, essential for circadian behavior. *Science.* 1994; 264:719–725. [PubMed: 8171325]
- Vitaterna MH, Selby CP, Todo T, Niwa H, Thompson C, Fruechte EM, Hitomi K, Thresher RJ, Ishikawa T, Miyazaki J, et al. Differential regulation of mammalian *period* genes and circadian rhythmicity by cryptochromes 1 and 2. *Proc. Natl. Acad. Sci. USA.* 1999; 96:12114–12119. [PubMed: 10518585]
- Vujovic N, Davidson AJ, Menaker M. Sympathetic input modulates, but does not determine, phase of peripheral circadian oscillators. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 2008; 295:R355–R360. [PubMed: 18434440]
- Walton KM, Fisher K, Rubitski D, Marconi M, Meng QJ, Sladek M, Adams J, Bass M, Chandrasekaran R, Butler T, et al. Selective inhibition of casein kinase 1 ϵ minimally alters circadian clock period. *J. Pharmacol. Exp. Ther.* 2009; 330:430–439. [PubMed: 19458106]
- Wang Y, Osterbur DL, Megaw PL, Tosini G, Fukuhara C, Green CB, Besharse JC. Rhythmic expression of *Nocturnin* mRNA in multiple tissues of the mouse. *BMC Dev. Biol.* 2001; 1:9. [PubMed: 11394964]
- Welsh DK, Logothetis DE, Meister M, Reppert SM. Individual neurons dissociated from rat suprachiasmatic nucleus express independently phased circadian firing rhythms. *Neuron.* 1995; 14:697–706. [PubMed: 7718233]
- Welsh DK, Yoo SH, Liu AC, Takahashi JS, Kay SA. Bioluminescence imaging of individual fibroblasts reveals persistent, independently phased circadian rhythms of clock gene expression. *Curr. Biol.* 2004; 14:2289–2295. [PubMed: 15620658]
- Welsh DK, Takahashi JS, Kay SA. Suprachiasmatic nucleus: cell autonomy and network properties. *Annu. Rev. Physiol.* 2010; 72:551–577. [PubMed: 20148688]
- Wilkinson, Kevin A, Henley, Jeremy M. Mechanisms, regulation and consequences of protein SUMOylation. *Biochem. J.* 2010; 428:133–145. [PubMed: 20462400]
- Wilkinson KD. Ubiquitin-dependent signaling: the role of ubiquitination in the response of cells to their environment. *J. Nutr.* 1999; 129:1933–1936. [PubMed: 10539765]
- Woo KC, Kim TD, Lee KH, Kim DY, Kim W, Lee KY, Kim KT. Mouse period 2 mRNA circadian oscillation is modulated by PTB-mediated rhythmic mRNA degradation. *Nucleic Acids Res.* 2009; 37:26–37. [PubMed: 19010962]
- Woo KC, Ha DC, Lee KH, Kim DY, Kim TD, Kim KT. Circadian amplitude of cryptochrome 1 is modulated by mRNA stability regulation via cytoplasmic hnRNP D oscillation. *Mol. Cell. Biol.* 2010; 30:197–205. [PubMed: 19858287]

- Xu Y, Padiath QS, Shapiro RE, Jones CR, Wu SC, Saigoh N, Saigoh K, Ptacek LJ, Fu YH. Functional consequences of a CK1 δ mutation causing familial advanced sleep phase syndrome. *Nature*. 2005; 434:640–644. [PubMed: 15800623]
- Xu Y, Toh KL, Jones CR, Shin JY, Fu YH, Ptacek LJ. Modeling of a human circadian mutation yields insights into clock regulation by PER2. *Cell*. 2007; 128:59–70. [PubMed: 17218255]
- Yagita K, Tamanini F, van Der Horst GT, Okamura H. Molecular mechanisms of the biological clock in cultured fibroblasts. *Science*. 2001; 292:278–281. [PubMed: 11303101]
- Yamada YR, Forger DB. Multiscale complexity in the mammalian circadian clock. *Curr. Opin. Genet. Dev.* 2010; 20:626–633. [PubMed: 20934868]
- Yamaguchi S, Isejima H, Matsuo T, Okura R, Yagita K, Kobayashi M, Okamura H. Synchronization of cellular clocks in the suprachiasmatic nucleus. *Science*. 2003; 302:1408–1412. [PubMed: 14631044]
- Yamazaki S, Numano R, Abe M, Hida A, Takahashi R, Ueda M, Block GD, Sakaki Y, Menaker M, Tei H. Resetting central and peripheral circadian oscillators in transgenic rats. *Science*. 2000; 288:682–685. [PubMed: 10784453]
- Yan L, Silver R. Resetting the brain clock: time course and localization of mPER1 and mPER2 protein expression in suprachiasmatic nuclei during phase shifts. *Eur. J. Neurosci*. 2004; 19:1105–1109. [PubMed: 15009158]
- Yang X, Downes M, Yu RT, Bookout AL, He W, Straume M, Mangelsdorf DJ, Evans RM. Nuclear receptor expression links the circadian clock to metabolism. *Cell*. 2006; 126:801–810. [PubMed: 16923398]
- Yin L, Wang J, Klein PS, Lazar MA. Nuclear receptor Rev-erba is a critical lithium-sensitive component of the circadian clock. *Science*. 2006; 311:1002–1005. [PubMed: 16484495]
- Yoo SH, Yamazaki S, Lowrey PL, Shimomura K, Ko CH, Buhr ED, Siepkha SM, Hong HK, Oh WJ, Yoo OJ, et al. PERIOD2::LUCIFERASE real-time reporting of circadian dynamics reveals persistent circadian oscillations in mouse peripheral tissues. *Proc. Natl. Acad. Sci. USA*. 2004; 101:5339–5346. [PubMed: 14963227]
- Yoo SH, Ko CH, Lowrey PL, Buhr ED, Song EJ, Chang S, Yoo OJ, Yamazaki S, Lee C, Takahashi JS. A noncanonical E-box enhancer drives mouse *Period2* circadian oscillations *in vivo*. *Proc. Natl. Acad. Sci. USA*. 2005; 102:2608–2613. [PubMed: 15699353]
- Yoshitane H, Takao T, Satomi Y, Du NH, Okano T, Fukada Y. Roles of CLOCK phosphorylation in suppression of E-box-dependent transcription. *Mol. Cell. Biol.* 2009; 29:3675–3686. [PubMed: 19414601]
- Young MW, Kay SA. Time zones: a comparative genetics of circadian clocks. *Nat. Rev. Genet.* 2001; 2:702–715. [PubMed: 11533719]
- Yu W, Zheng H, Price JL, Hardin PE. DOUBLETIME plays a noncatalytic role to mediate CLOCK phosphorylation and repress CLOCK-dependent transcription within the *Drosophila* circadian clock. *Mol. Cell. Biol.* 2009; 29:1452–1458. [PubMed: 19139270]
- Zhang X, Odom DT, Koo SH, Conkright MD, Canettieri G, Best J, Chen H, Jenner R, Herbolsheimer E, Jacobsen E, et al. Genome-wide analysis of cAMP-response element binding protein occupancy, phosphorylation, and target gene activation in human tissues. *Proc. Natl. Acad. Sci. USA*. 2005; 102:4459–4464. [PubMed: 15753290]
- Zhao WN, Malinin N, Yang FC, Staknis D, Gekakis N, Maier B, Reischl S, Kramer A, Weitz CJ. CIPC is a mammalian circadian clock protein without invertebrate homologues. *Nat. Cell Biol.* 2007; 9:268–275. [PubMed: 17310242]
- Zheng B, Larkin DW, Albrecht U, Sun ZS, Sage M, Eichele G, Lee CC, Bradley A. The *mPer2* gene encodes a functional component of the mammalian circadian clock. *Nature*. 1999; 400:169–173. [PubMed: 10408444]
- Zheng B, Albrecht U, Kaasik K, Sage M, Lu W, Vaishnav S, Li Q, Sun ZS, Eichele G, Bradley A, et al. Nonredundant roles of the *mPer1* and *mPer2* genes in the mammalian circadian clock. *Cell*. 2001; 105:683–694. [PubMed: 11389837]




Figure 1. Model of the mammalian cell-autonomous oscillator as described in the text. Abbreviations: CCG, clock-controlled gene; P, phosphate; U, ubiquitin.

Table 1

Behavioral phenotypes of mutations in mouse clock and clock-related genes

Gene(s)	Protein product(s)	Mutant allele(s)	Mutant phenotype(s)	References
<i>Clock</i>	bHLH-PAS transcription factor	<i>Clock</i> ^{<i>Δ19/Δ19</i>} <i>Clock</i> ^{-/-}	4 hr longer pd/arrhythmic 0.4 hr shorter pd	(Vitaterna <i>et al.</i> , 1994) (DeBruyne <i>et al.</i> , 2006)
<i>Npas2</i> (<i>Mop4</i>)	bHLH-PAS transcription factor	<i>Npas2</i> ^{-/-}	0.2 hr shorter pd	(Dudley <i>et al.</i> , 2003)
<i>Clock/Npas2</i>	bHLH-PAS transcription factors	<i>Clock</i> ^{-/-} / <i>Npas2</i> ^{-/-}	Arrhythmic	(DeBruyne <i>et al.</i> , 2007a)
<i>Bmal1</i> (<i>Arntl</i> , <i>Mop3</i>)	bHLH-PAS transcription factor	<i>Bmal1</i> ^{-/-}	Arrhythmic	(Bunger <i>et al.</i> , 2000)
<i>Cry1</i>	flavoprotein	<i>Cry1</i> ^{-/-}	1 hr shorter pd	(van der Horst <i>et al.</i> , 1999; Vitaterna <i>et al.</i> , 1999)
<i>Cry2</i>	flavoprotein	<i>Cry2</i> ^{-/-}	1 hr longer pd	(Thresher <i>et al.</i> , 1998; van der Horst <i>et al.</i> , 1999)
<i>Cry1/Cry2</i>	flavoproteins	<i>Cry1</i> ^{-/-} / <i>Cry2</i> ^{-/-}	Arrhythmic	(van der Horst <i>et al.</i> , 1999; Vitaterna <i>et al.</i> , 1999)
<i>Per1</i>	PAS protein	<i>Per1</i> ^{-/-}	0.7 hr shorter pd	(Cermakian <i>et al.</i> , 2001)
		<i>Per1</i> ^{<i>brdm1</i>}	1 hr shorter pd	(Zheng <i>et al.</i> , 2001)
		<i>Per1</i> ^{<i>ldc</i>}	0.5 hr shorter pd/arrhythmic	(Bae <i>et al.</i> , 2001)
<i>Per2</i>	PAS protein	<i>Per2</i> ^{<i>brdm1</i>}	1.5 hr shorter pd/arrhythmic	(Zheng <i>et al.</i> , 1999)
		<i>Per2</i> ^{<i>ldc</i>}	Arrhythmic	(Bae <i>et al.</i> , 2001)
<i>Per3</i>	PAS protein	<i>Per3</i> ^{-/-}	0–0.5 hr shorter pd	(Shearman <i>et al.</i> , 2000a)
<i>Per1/Per2</i>	PAS proteins	<i>Per1</i> ^{<i>brdm1</i>} / <i>Per2</i> ^{<i>brdm1</i>}	Arrhythmic	(Zheng <i>et al.</i> , 2001)
		<i>Per1</i> ^{<i>brdm1</i>} / <i>Per2</i> ^{<i>brdm1</i>}	Arrhythmic	(Bae <i>et al.</i> , 2001)
<i>Per1/Cry1</i>	PAS protein/flavoprotein	<i>Per1</i> ^{<i>brdm1</i>} / <i>Cry1</i> ^{-/-}	Normal behavior	(Oster <i>et al.</i> , 2003)
<i>Per1/Cry2</i>	PAS protein/flavoprotein	<i>Per1</i> ^{<i>brdm1</i>} / <i>Cry2</i> ^{-/-}	<6 months, 1.5 hr longer pd; >6 months, arrhythmic	(Oster <i>et al.</i> , 2003)
<i>Per2/Cry1</i>	PAS protein/flavoprotein	<i>Per2</i> ^{<i>brdm1</i>} / <i>Cry1</i> ^{-/-}	Arrhythmic	(Oster <i>et al.</i> , 2002)
<i>Per2/Cry2</i>	PAS protein/flavoprotein	<i>Per2</i> ^{<i>brdm1</i>} / <i>Cry2</i> ^{-/-}	0–0.4 hr shorter pd	(Oster <i>et al.</i> , 2002)
<i>Rev-erba</i> (<i>Nr1d1</i>)	nuclear receptor	<i>Rev-erba</i> ^{-/-}	0.5 hr shorter pd; disrupted entrainment	(Preitner <i>et al.</i> , 2002)
<i>Rev-erbb</i> (<i>Nr1d2</i>)	nuclear receptor	—	—	—
<i>Rora</i> (<i>Rora</i>)	nuclear receptor	<i>Rora</i> ^{-/-} (<i>staggerer</i>)	0.5 hr shorter pd; disrupted entrainment	(Sato <i>et al.</i> , 2004)
<i>Rorb</i> (<i>Rorb</i>)	nuclear receptor	<i>Rorb</i> ^{-/-}	0.5 hr longer pd	(Masana <i>et al.</i> , 2007)
<i>Rory</i> (<i>Rorc</i>)	nuclear receptor	<i>Rory</i> ^{-/-}	Normal behavior	(Liu <i>et al.</i> , 2008)
<i>Dec1</i> (<i>Bhlhe40</i> , <i>Stra13</i> , <i>Sharp-2</i>)	bHLH transcription factor	<i>Stra13</i> ^{-/-}	0.15 hr longer pd	(Nakashima <i>et al.</i> , 2008)
<i>Dec2</i> (<i>Bhlhe41</i> , <i>Sharp-1</i>)	bHLH transcription factor	<i>Sharp-1</i> ^{-/-}	Delayed resetting	(Rossner <i>et al.</i> , 2008)
<i>CK1δ</i> (<i>Csnk1d</i>)	casein kinase 1	<i>CK1δ</i> ^{+/-}	0–0.5 hr longer pd	(Xu <i>et al.</i> , 2005; Etchegaray <i>et al.</i> , 2009)
<i>CK1ε</i> (<i>Csnk1e</i>)	casein kinase 1	<i>CK1ε</i> ^{-/-}	0.2–0.4 hr longer pd	(Meng <i>et al.</i> , 2008;

Gene(s)	Protein product(s)	Mutant allele(s)	Mutant phenotype(s)	References
				Etchegaray <i>et al.</i> , 2009)
<i>CK1a</i> (<i>Csnk1a1</i>)	casein kinase 1	* <i>CK1e^{tau}</i>	4 hr shorter pd	(Lowrey <i>et al.</i> , 2000; Meng <i>et al.</i> , 2008)
<i>Fbx13</i>	F-box protein	—	—	—
		<i>Fbx13^{Ovtn}</i>	2 hr longer pd	(Siepka <i>et al.</i> , 2007)
		<i>Fbx13^{Ath}</i>	3 hr longer pd	(Godinho <i>et al.</i> , 2007)
<i>Bmal2</i> (<i>Arntl2</i> , <i>Mop9</i> , <i>Clif</i>)	bHLH-PAS transcription factor	—	—	—
<i>Pgc1a</i> (<i>Ppargc1a</i>)	transcriptional coactivator	<i>Pgc1a^{-/-}</i>	0.3 hr longer pd	(Liu <i>et al.</i> , 2007c)
<i>Mtnr1a</i> (<i>Mel1a</i>)	G protein-coupled receptor	<i>Mtnr1a^{-/-}</i>	Normal behavior	(Liu <i>et al.</i> , 1997a)
<i>Mtnr1b</i> (<i>Mel1b</i>)	G protein-coupled receptor	<i>Mtnr1b^{-/-}</i>	Normal behavior	(Jin <i>et al.</i> , 2003)
<i>Opn4</i>	melanopsin; opsin 4	<i>Opn4^{-/-}</i>	Attenuated photic responses	(Panda <i>et al.</i> , 2002b; Ruby <i>et al.</i> , 2002)
<i>Dbp</i>	PAR bZIP transcription factor	<i>Dbp^{-/-}</i>	0.5 hr shorter pd	(Lopez-Molina <i>et al.</i> , 1997)
<i>Vipr2</i>	G protein-coupled receptor	<i>Vipr2^{-/-}</i>	Disrupted locomotor rhythm	(Harmar <i>et al.</i> , 2002; Cutler <i>et al.</i> , 2003)
<i>Vip</i>	peptide hormone	<i>Vip^{-/-}</i>	1 hr shorter pd/arrhythmic	(Colwell <i>et al.</i> , 2003)
<i>Prok2</i> (<i>PK2</i>)	secreted protein	<i>Prok2^{-/-}</i>	Reduced locomotor activity	(Li <i>et al.</i> , 2006)
<i>Nocturnin</i> (<i>Ccrn4</i>)	deadenylase	<i>Noc^{-/-}</i>	Normal behavior	(Green <i>et al.</i> , 2007)

Gene symbols listed here are the predominate forms used in the scientific literature; alternate forms are given in parentheses. When the predominant symbol differs from standard mouse gene nomenclature, the standard form is given in parentheses. For genes with more than two symbols in parentheses, the form adhering to standard mouse gene nomenclature is underlined.

* First identified as a mutation in Syrian hamster (*Mesocricetus auratus*).

Table 2

Cell and tissue phenotypes of mutations in mouse clock and clock-related genes

Gene(s)	Mutant allele(s)	Cellular phenotype(s)	Tissue phenotype(s)	References
<i>Clock</i>	<i>Clock</i> ^{-/-}	—	SCN: WT; lung, liver: AR	(DeBruyne <i>et al.</i> , 2007a; DeBruyne <i>et al.</i> , 2007b)
<i>Npas2</i> (<i>Mop4</i>)	<i>Npas2</i> ^{-/-}	—	SCN, lung, liver: WT	(DeBruyne <i>et al.</i> , 2007b)
<i>Bmal1</i> (<i>Arntl</i> , <i>Mop3</i>)	<i>Bmal1</i> ^{-/-}	Fibroblasts, SCN neurons: AR	SCN: variable/stochastic; pituitary, liver, lung, cornea: AR	(Liu <i>et al.</i> , 2008; Ko <i>et al.</i> , 2010)
<i>Cry1</i>	<i>Cry1</i> ^{-/-}	Fibroblasts, SCN neurons: AR	SCN: short; lung, liver, cornea: AR	(Liu <i>et al.</i> , 2007b)
<i>Cry2</i>	<i>Cry2</i> ^{-/-}	Fibroblasts, SCN neurons: long	SCN, lung, liver, cornea: long	(Liu <i>et al.</i> , 2007b)
<i>Cry1/Cry2</i>	<i>Cry1</i> ^{-/-} / <i>Cry2</i> ^{-/-}	Fibroblasts: AR	SCN, lung, liver, cornea: AR	(Yagita <i>et al.</i> , 2001; Liu <i>et al.</i> , 2007b)
<i>Per1</i>	<i>Per1</i> ^{ldc}	Fibroblasts, SCN neurons: AR	SCN: WT; lung: AR	(Liu <i>et al.</i> , 2007b)
<i>Per2</i>	<i>Per2</i> ^{ldc}	Fibroblasts: AR	—	(Liu <i>et al.</i> , 2007b)
<i>Per3</i>	<i>Per3</i> ^{-/-}	Fibroblasts: short	SCN, lung: short	(Liu <i>et al.</i> , 2007b)
<i>Rev-erba</i> (<i>Nr1d1</i>)	<i>Rev-erba</i> ^{-/-}	Fibroblasts: WT	—	(Liu <i>et al.</i> , 2008)
<i>Rora</i> (<i>Rora</i>)	<i>Rora</i> ^{-/-} (<i>staggerer</i>)	Fibroblasts: WT	—	(Liu <i>et al.</i> , 2008)
<i>Rory</i> (<i>Rorc</i>)	<i>Rory</i> ^{-/-}	Fibroblasts: WT	Lung, liver: WT	(Liu <i>et al.</i> , 2008)
<i>CK1δ</i> (<i>Csnk1d</i>)	<i>CK1δ</i> ^{-/-}	Fibroblasts: long	*SCN, liver: long	(Etchegaray <i>et al.</i> , 2009; Lee <i>et al.</i> , 2009; Etchegaray <i>et al.</i> , 2010)
<i>CK1ε</i> (<i>Csnk1e</i>)	<i>CK1ε</i> ^{-/-}	Fibroblasts: WT	*SCN, liver: WT	(Etchegaray <i>et al.</i> , 2009; Etchegaray <i>et al.</i> , 2010)
	<i>CK1ε</i> ^{neu}	Fibroblasts: short	SCN, pituitary, lung, kidney: short	(Meng <i>et al.</i> , 2008)

Gene symbols are as in Table 1. Abbreviations: AR, arrhythmic; WT, wild type;

* neonatal tissue. Table modified from (Baggs *et al.*, 2009).