

Genome comparison of two *Exiguobacterium* strains from high altitude andean lakes with different arsenic resistance: identification and 3D modeling of the Acr3 efflux pump

OPEN ACCESS

Omar F. Ordoñez¹, Esteban Lanzarotti^{2,3}, Daniel Kurth¹, Néstor Cortez⁴, María E. Farías¹ and Adrian G. Turjanski^{2,3*}

Edited by:

Partha Basu,
Duquesne University, USA

Reviewed by:

C. French,
University of Edinburgh, UK
John W. Moreau,
University of Melbourne, Australia

*Correspondence:

Adrian G. Turjanski,
INQUIMAE-UBA/CONICET, Ciudad
Universitaria Pabellón II, Int. Guiraldes
2620, C1428EHA Buenos Aires,
Argentina
adrian@qi.fcen.uba.ar

Specialty section:

This article was submitted to
Microbiological Chemistry and
Geomicrobiology,
a section of the journal
Frontiers in Environmental Science

Received: 22 December 2014

Accepted: 02 July 2015

Published: 22 July 2015

Citation:

Ordoñez OF, Lanzarotti E, Kurth D,
Cortez N, Farías ME and Turjanski AG
(2015) Genome comparison of two
Exiguobacterium strains from high
altitude andean lakes with different
arsenic resistance: identification and
3D modeling of the Acr3 efflux pump.
Front. Environ. Sci. 3:50.
doi: 10.3389/fenvs.2015.00050

¹ Laboratorio de Investigaciones Microbiológicas de Lagunas Andinas, Planta Piloto de Procesos Industriales Microbiológicos (PROIMI), CCT-CONICET, Tucumán, Argentina, ² Departamento de Química Inorgánica, Analítica y Química Física/INQUIMAE-CONICET, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Ciudad Universitaria, Buenos Aires, Argentina, ³ Departamento de Química Biológica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Ciudad Universitaria, Buenos Aires, Argentina, ⁴ Instituto de Biología Molecular y Celular de Rosario (IBR – CONICET), Facultad de Cs. Bioquímicas y Farmacéuticas (Universidad Nacional de Rosario), Rosario, Argentina

Arsenic exists in natural systems in a variety of chemical forms, including inorganic arsenite (As [III]) and arsenate (As [V]). The majority of living organisms have evolved various mechanisms to avoid occurrence of arsenic inside the cell due to its toxicity. Common core genes include a transcriptional repressor ArsR, an arsenate reductase ArsC, and arsenite efflux pumps ArsB and Acr3. To understand arsenic resistance we have performed arsenic tolerance studies, genomic and bioinformatic analysis of two *Exiguobacterium* strains, S17 and N139, from the high-altitude Andean Lakes. In these environments high concentrations of arsenic were described in the water due to a natural geochemical phenomenon, therefore, these strains represent an attractive model system for the study of environmental stress and can be readily cultivated. Our experiments show that S17 has a greater tolerance to arsenite (10 mM) than N139, but similar growth in arsenate (150 mM). We sequenced the genome of the two *Exiguobacterium* and identified an *acr3* gene in S17 as the only difference between both species regarding known arsenic resistance genes. To further understand the Acr3 we modeled the 3D structure and identified the location of relevant residues of this protein. Our model is in agreement with previous experiments and allowed us to identify a region where a relevant cysteine lies. This Acr3 membrane efflux pump, present only in S17, may explain its increased tolerance to As(III) and is the first Acr3-family protein described in *Exiguobacterium* genus.

Keywords: arsenic resistance, *Exiguobacterium*, membrane proteins, sequencing, ACR3

Introduction

Arsenic exists in natural systems in a variety of chemical forms, including inorganic arsenite (As [III]) and arsenate (As [V]) and methylated organic forms. Arsenate behaves as a phosphate analog interfering with phosphate uptake and utilization, while arsenite is able to disrupt many enzymatic activities. The majority of living organisms have evolved various mechanisms to avoid occurrence of arsenic inside the cell, although some bacteria are able to use it as chemical energy source or need the metalloid for growth (Stolz and Oremland, 1999; Macy et al., 2000; Silver and Phung, 2005). It has been suggested that a relationship exists between arsenic toxicity and cellular antioxidant systems (Hrimpeng et al., 2006; Pandey et al., 2012).

Genes related to arsenic resistance in bacteria are usually encoded in the so called *ars* operons. These operons have varied configurations, but the common core genes include a transcriptional repressor ArsR, an arsenate reductase ArsC, and an arsenite efflux pump which extrude this toxic species (Paez-Espino et al., 2009). The pumps belong to two different unrelated families, the ArsB proteins from the Major Facilitator Superfamily, and the Acr3 proteins from the BART superfamily (Mansour et al., 2007). The ArsB proteins are best characterized, and have been shown to extrude both arsenite and antimonite in a mechanism driven either by the electrochemical gradient or by ATP hydrolysis when the ArsA ATPase is present (Rosen et al., 1999). Members of the Acr3 family are found in bacteria, archaea and fungi (Rosen, 1999; Wysocki et al., 2003) and in environmental studies *acr3* genes seem to be very widespread among arsenic resistant bacteria, even more than *arsB* genes (Achour et al., 2007; Cai et al., 2009). Nevertheless, Acr3 proteins have been functionally characterized only in a few species including *Bacillus subtilis*, *Synechocystis* sp., *Corynebacterium glutamicum* and *Saccharomyces cerevisiae* (López-Maury et al., 2003; Aaltonen and Silow, 2008; Fu et al., 2009; Maciaszczyk-Dziubinska et al., 2011; Villadangos et al., 2012), the last being the best studied microorganism where Acr3 acts as a metalloid/H(+) antiporter for arsenite and antimonite. Poirel et al. suggest that *arsB* and *acr3*, arsenite transporter genes, are appropriate biomarkers of arsenic stress that may be suitable for further exploring the adaptive response of bacterial communities to arsenic in contaminated environments (Poirel et al., 2013).

The high-altitude Andean Lakes (HAAL) are ecosystems located in the South American Andes. The HAAL ecosystems are systems of shallow lakes formed during the Tertiary geological period (Seufferheld et al., 2008; Farias et al., 2011). These ecosystems are unique not only for their geographical characteristics and broad range of extreme environments but also for their abundant biodiversity (Fernandez Zenoff et al., 2006; Dib et al., 2008; Seufferheld et al., 2008; Flores et al., 2009; Ordoñez et al., 2009; Albarracín et al., 2011; Farias et al., 2013). In these environments high concentrations of arsenic were described in the water due to a natural geochemical phenomenon (Mantelli et al., 2003; Romero et al., 2003; Escudero et al., 2007; Dib et al., 2008, 2009; Escalante et al., 2009). The high concentration of arsenic present in HAAL is strongly limiting not only for human life but also for

growth of many microorganisms (Valko et al., 2005) and selects development of arsenic tolerant bacteria (Dib et al., 2008; Ordoñez et al., 2009). Different concentrations of arsenic (maximum values) were reported in these environments, 2.3 mg L⁻¹ in Tebenquiche Lake (Farias et al., 2014), 9.1 mg L⁻¹ in Salar de Llamara (Rasuk et al., 2014), 0.8–11.8 mg L⁻¹ in Andean Lakes (Ordoñez et al., 2009), 0.9–28 mg L⁻¹ in salt lakes (Lara et al., 2012). Recently, in other HAAL higher values were detected, namely arsenic values between 18.5 and 33.8 mg L⁻¹ were observed in Laguna Socompa (Farias et al., 2011, 2013); these values are greater than the one reported for Mono Lake (200 μM HAsO₄²⁻, approximately 19.8 mg L⁻¹ As) (Oremland et al., 2004). In this place, Farias and colleagues described stromatolites forming at an altitude of 3570 m, developing in an alkaline, hypersaline environment, rich in inorganic nutrients and very rich in arsenic, and with mild temperatures (20–24°C) due to a hydrothermal input (Farias et al., 2011, 2013).

The genus *Exiguobacterium* was one of the most widespread and representative genera on the HAAL, being detected by direct (pure culture isolation) and indirect (DGGE) techniques (Ordoñez et al., 2009). *Exiguobacterium* spp. have been isolated from, or molecularly detected in, a wide range of habitats including cold and hot environments with temperature ranging from –12 up to 55°C (Vishnivetskaya et al., 2006; Rodrigues and Tiedje, 2007). This fact confers substantial interest to the genus as a potential model system for the investigation of attributes that may correlate with adaptation and evolution of organisms to diverse thermal regimes (Vishnivetskaya et al., 2009). Several studies have shown that some strains of *Exiguobacterium* possess unique properties of interest for applications in biotechnology, bioremediation, industry and agriculture (Pattanapitpaisal et al., 2002; Lopez et al., 2005; Okeke et al., 2007).

Recently, we presented the draft genome sequence of *Exiguobacterium* sp. S17, isolated from a stromatolite in L. Socompa (GenBank:ASXD00000000.1) and the draft genome sequence of *Exiguobacterium* sp. N139, a UV-B resistant bacterium isolated from L. Negra (GenBank:JMEH00000000). These strains represent an attractive model system for the study of environmental stress capable of growing readily in the laboratory. There is particular interest in the S17 strain given its arsenic-rich original niche. In a previous study, it was able to grow in high arsenic concentration, it tolerated arsenic concentration such as 10 mM of As[III] and 150 mM of As[V], but a proteomic approach did not detect unique nor arsenic specific proteins (Belfiore et al., 2013).

Based on these previous results, the aim of this study was to gain further insight into mechanisms of tolerance to high arsenic concentrations by analyzing the genome sequences of S17 and N139, recently isolated from HAAL extreme environments. We initially tested the arsenite and arsenate resistance of S17, N139 and model organisms. Interestingly, our experiments show that S17 has a greater resistance to arsenite than N139, but similar growth in arsenate. We then compared the genomes and identified that the only difference between both species regarding known arsenic resistant genes is the presence of an *acr3* gene in S17. To further understand the *acr3* of S17 we analyzed

the sequence and structural characteristics, which allowed us to identify the relevant residues.

Materials and Methods

Strain Isolation and Culture Media

The strain S17 was isolated *in situ* from L. Socompa stromatolite (Ordoñez et al., 2013) by plating samples on PY agar medium, containing per liter: 10 g peptone, 0.5 g yeast extract, 2 mL MgCl₂ 1 M, 2 mL CaCl₂ 1 M, 2.4 mL FeSO₄ 0.5% and 1.5% w/v agar, after incubation at room temperature, while the strain *Exiguobacterium* sp. N139 was isolated from water in L. Negra by plating in Lake medium (LM). The LM was used to maintain the same salinity as the isolation environment and was obtained by filtering lake water (0.22 μm Biopore filters) and adding 2.5 g yeast extract and 12 g agar (Difco) per liter. Isolated N139 was then cultured in the same isolation medium at 20°C with shaking (150 rpm). The strain *Exiguobacterium aurantiacum* DSM 6208, from the German Collection of Microorganisms and Cell Cultures (DSM) was included in the assays for comparison as external control.

To unify the tests, all cultures were routinely grown with shaking (150 rpm) in Luria-Bertani broth (LB) at 30°C overnight or in plates with LB medium supplemented with 1.5% agar (Britania). When indicated, LB was diluted to 50% final concentration (LB₅₀).

Tolerance Toward Arsenic

For qualitative assessment of arsenic tolerance in solid media, bacterial cultures collected at OD_{600nm} of 0.4 were subjected to serial dilutions. 10 μl aliquots were plated onto LB₅₀ agar plates, supplemented with different concentrations of sodium arsenate (As[V]) and sodium arsenite (As[III]) as indicated in **Figure 2**.

To quantify the resistance toward arsenic of the studied strains, cultures were grown at different concentrations of this agent. Overnight cultures were diluted to an initial OD_{600nm} between 0.05 and 0.09 into fresh LB₅₀ medium supplemented with different concentrations of arsenic: 2.5 mM, 5 mM and 10 mM of sodium arsenite (As [III]); 50, 100, and 150 mM of sodium arsenate (As [V]). A non-supplemented sample was used as positive control. The cultures were incubated at 30°C with shaking for 24 h, and OD_{600nm} was measured with a

spectrophotometer Ultrospec 10 cell density meter (Amersham Biosciences).

Phylogenetic Analyses of Sequenced *Exiguobacterium* spp.: 16S rRNA, Multilocus Sequence Analysis (MLSA), Acr3 and ArsB Proteins

We used 7 genomes of *Exiguobacterium* in these phylogenetic analyses (**Table 1**). MLSA was based on the concatenated sequences of house-keeping genes (Brown et al., 2001; Thompson et al., 2009). The 16S rRNA gene sequences, the gene sequences used for MLSA (i.e., *ftsZ*, *gyrB*, *mreB*, *pyrH*, *recA*, *rpoA*, and *topA*) and protein sequences of the ArsB and Acr3 families were obtained from the National Center for Biotechnology Information (NCBI). RAST server (Aziz et al., 2008) was used to annotate the draft genomes.

Sequences were aligned using CLUSTAL X 2.0.9, a Windows interface for the multiple alignment program ClustalW (Larkin et al., 2007). Phylogenetic analyses were carried out with the program MEGA 5 (Tamura et al., 2011), using the neighbor-joining method (Saitou and Nei, 1987). All positions containing gaps in the datasets were removed, and the confidence intervals obtained by this method were positioned using the bootstrap procedure, performing 1000 replicates (Saitou and Nei, 1987).

RNA Extraction and RT-PCR

Exiguobacterium sp. S17 was grown overnight in 50 ml of LB₅₀ and LB₅₀ with 7.5 mM of As [III] or 100 mM of As[V]. Twenty-five milliliters of each culture were used for RNA extraction. Cells were harvested by centrifugation (9200 g at 4°C during 5 min) and pellets were suspended in 1 ml of Quick-zol, total RNA was extracted following the protocol proposed by the supplier (Quick-Zol- Kalium Technologies). Acr3 differential expression was assessed by RT-PCR using the specific primers designed from the genome of S17 strain (ACR3F1: 5'-CTTGGCTTCGCGATTGGACTGAG3' - ACR3R1: 5'-GACACGGCGGCTTATGGCAAAT3') and 16S primer (357F: 5'-TACTGATAGAA.

TGTGGAGC-3' -518R: 5'-CGT ATT ACC GCG GCT GCT GG-3') was used as housekeeping gene. The reverse transcription was performed with ImProm-II TM Reverse Transcriptase (Promega), following the protocol recommended by the supplier.

TABLE 1 | Genomic features and number of arsenic resistance genes observed in *Exiguobacterium* genomes.

Organism	Accession number	Genome size (mb)	G+C (mol%)	Number of CDS	Number of contigs	Acr3	ArsB	ArsA	ArsR	ArsD	ArsC
<i>Exiguobacterium</i> sp. S17	ASXD00000000.1	3.13	53.1	3218	163	1	1	1	1	1	2
<i>Exiguobacterium</i> sp. N139	JMEH00000000	2.95	52	3007	45	0	1	1	1	1	2
<i>Exiguobacterium sibiricum</i> 255-15	CP001022.1	3.04	47.7	3015	1	0	1	0	1	0	1
<i>Exiguobacterium antarcticum</i> B7	CP003063.1	2.82	47.5	2772	1	0	1	0	1	0	1
<i>Exiguobacterium</i> sp. 8-11-1	ATKK00000000.1	2.91	52.8	2886	31	0	1	1	1	1	2
<i>Exiguobacterium pavilonensis</i> RW-2	ATCL00000000.1	3.02	52	3077	23	0	1	1	1	1	1
<i>Exiguobacterium</i> sp. AT 1b	CP001615.1	3.00	48.5	3020	1	0	1	1	1	1	1

Orthologous Analysis Using Bidirectional Best Hits

An analysis of orthologous was done comparing coding sequences present in S17 and N139 contigs, in order to explain the resistance to arsenic in both organisms. Bidirectional Best Hits (BBHs) relation was applied as described in Overbeek et al. (Overbeek et al., 1999). E-value cutoff of 1E-05 was used in Blast search and a 90% coverage cutoff was added to the criteria in order to obtain more trusted relationships. Numbers of genes belonging to BBH relation in both organisms are shown in Table S1 (see Supplementary Materials).

Transmembrane Segment Prediction

In this work we use four predictors of transmembrane segments: TMHMM (Krogh et al., 2001), DAS (Cserző et al., 1997), HMMTOP (Tusnády and Simon, 2001), TOPCONS (Bernsel et al., 2009), plus two other general secondary structure predictors: PSIPRED (Jones, 1999), PORTER (Pollastri and McLysaght, 2005). A majority voting consensus was calculated, assigning transmembrane segment structure if 4 or more predictors agree.

Template Alignment and Model Building for S17 Acr3

To find a suitable template for model construction, 5 iterations of PSI-Blast (Altschul et al., 1997) were performed to build a profile for the *Exiguobacterium* S17 Acr3 protein using UniRef90 (Suzek et al., 2007) and 1E-05 as gathering threshold. Using this profile, another PSI-Blast iteration was computed against a library of templates. This library was constructed using each amino acid

sequence underlying the PDB structures and clustering over their sequences was done using CD-HIT (Li et al., 2001) to admit at least 95% sequence identity between them. MODELLER (Sali and Blundell, 1993) was used to produce several models based on this alignment and the best model was chosen using ProQM Server (Ray et al., 2010) for model quality assessment.

Results

Phylogenetic Reconstructions by 16S rRNA and MLSA

The study of HAAL allowed isolation of bacteria resistant to UVR, antibiotic, arsenic and other extreme factors (Fernandez Zenoff et al., 2006; Dib et al., 2008; Flores et al., 2009; Ordoñez et al., 2009). In this place a predominance of *Exiguobacterium* spp. was observed (Ordoñez et al., 2009) and the strains N139 and S17 were assigned to this genus by 16S rRNA sequence analysis.

In this work were selected both conserved and variable single copy genes from the *Exiguobacterium* genomes, belonging to different functional groups used in several taxonomic studies (Brown et al., 2001; Thompson et al., 2009). Neighbor-Joining and maximum likelihood trees based on 16S rRNA gene sequences (Figure 1A) and MLSA (Figure 1B) showed similar topology using both markers. Bootstrap analysis indicated that most branches were highly significant. The phylogenetic reconstruction indicated a clear separation of groups of temperate (i.e., S17, N139, RW-2, 8-11-1, and AT1b) and cold (i.e., 255-15 and B7) origin. S17 was closely related to 8-11-1 and N139 to RW-2, all forming a single clade, while the strain *Exiguobacterium* sp. AT1b was located distant from these. On




FIGURE 1 | Phylogenetic analyses of sequenced *Exiguobacterium* spp. Phylogenetic tree based on 16S rRNA gene sequences (A) and based on MLSA of eight housekeeping genes (i.e. *ftsZ*, *rpoA*, *recA*, *topA*, *gapA*, *mreB*, *gyrB*, and *pyrH*) (B). The sequences included in the analysis were obtained from the National Center for Biotechnology Information (NCBI). The

evolutionary history of *Exiguobacterium* strains was inferred using the Neighbor-Joining method (Saitou and Nei, 1987) with 1000 repetitions. Robustness of the major branch points is indicated by bootstrap values. Phylogenetic analyses were performed with the program MEGA 5.05 (Tamura et al., 2011).

the other hand, *Exiguobacterium* sp. B7 and 255-15 formed a different clade with high similarity between them.

Tolerance to Arsenic

Soluble arsenic is present in nature as arsenate (As[V]) and arsenite (As[III]), the latter being more toxic. Tolerance against arsenic was firstly evaluated by placing drops of serially diluted culture of the studied strains on LB₅₀-agar plates, supplemented with different concentrations of arsenate or arsenite as sodium salt (see Materials and Methods). *Exiguobacterium* sp. S17 and N139 were compared, since both were isolated from a HAAL, whereas DSM 6208 was used as external control in all assays.


Exiguobacterium sp. S17, N139 and control strain DSM 6208 showed tolerance to As[V], being able to grow at concentrations up to 100 mM (Figure 2A). Interestingly, only the S17 strain was able to tolerate and grow at concentrations above 5 mM arsenite As[III] in culture media, while strains N139 and DSM 6208 did not grow at concentrations higher than 2.5 mM arsenite (Figure 2B). To quantify the inhibition due to arsenic in the culture medium, assays at different concentrations were performed in liquid media and final OD_{600nm} after 12 h of growth was determined. As shown in Figure 2C, growth of the HAALs isolates (S17 and N139) was similar to that of the control strain at all tested As[V] concentrations up to 150 mM. In contrast, at 10 mM As[III], *Exiguobacterium* N139 and DSM 6208 showed a 90% decline in growth, reaching OD_{600nm} values of 5–10% of the corresponding control without arsenic (Figure 2D). The S17 strain tolerated all tested concentrations of the metalloid,


showing only a minor impairment in growth at 10 mM of As(III) (Figure 2D).

Comparative Analysis of Arsenic Detoxification Genes from *Exiguobacterium* Genomes

In this work we used the sequence data of genes related to arsenic resistance from five sequenced *Exiguobacterium* genomes obtained from the NCBI. The characteristics, the source and the accession number of these strains are shown in Table 1. These data were utilized for comparison with the genome of *Exiguobacterium* sp. S17 and N139.

Analysis of orthologous genes shows a similar relationship between the genomes to that inferred using 16S rRNA and MLSA (see Table S1 in Supplementary Materials). As pictured in Figure 3, S17 and N139 genomes share 2221 ORFs which corresponds to 68.9% of total ORFs for S17 and 73.6% for N139. Differences between S17 and N139 biological subsystems were studied using RAST annotations assigned to non-orthologous genes (Table S2, see Supplementary Materials). Most of them were not assigned by RAST as to be in a particular subcategory: 595 and 581 for N139 and S17, respectively. In the “Virulence, Disease and Defense” category some differences were found. N139 possessed several subsystems apparently absent from S17, including genes related to tolerance of Mercury (3 ORFs), Copper (4 ORFs) and Cadmium (6 ORFs). Interestingly, S17 compared to N139, in addition to some antibiotic resistance features (3 ORFs), contains an arsenical resistance efflux pump Acr3. This seems to





be the main difference between N139 and S17 in terms of arsenic tolerance as defined by RAST.

Genetic organization of the arsenic resistance genes from the different strains is shown in **Figure 4**. Most of these genes are organized in two versions of the *ars* operon, *arsRBC* being the simplest configuration. The other configuration includes the genes *arsRDAXB* (*X* is annotated as a hypothetical protein). As shown in **Figure 4**, strains B7 and 255-15 have the smaller number of arsenic resistant genes, comprising the *arsRBC* basic detoxification system. S17, N139, AT1b, 8-11-1, and RW2 have the additional *arsD* and *arsA* genes, which would provide a more effective detoxification. In turn, S17 has a second unique cytoplasmic arsenite extrusion pump, the *acr3* gene, being the first time this gene is reported in the genus *Exiguobacterium*. In all cases *arsC* is not included in the operon, and in a few cases two *arsC* copies are found in the genomes (i.e., S17, N139 and 8-11-1). The *acr3* gene from S17 is not located in the operon with the other arsenic resistance genes. The genetic organization matches the 16S rRNA and MLSA distribution, thus B7 and 255-15, which group together in those trees (**Figures 1A,B**), have a short version of the operon, while the strains S17, N139, AT1b, 8-11-1, and RW2, which form another clade in the tree, show an extended operon.

Expression of *acr3* in S17 Strain

To assess the presence of the *acr3* transcript gen in S17, this strain was grown in presence and absence of arsenic in three different conditions: As[III] 7.5 mM, As[V] 100 mM, and control without As. RT-PCR was performed and the obtained cDNA was used to amplify the *acr3* gen with specific primers (ACR3F1-ACR3R1). The obtained results indicate that the *acr3* gene is expressed and its expression is constitutive (Figure S7).

Phylogenetic Tree of *ArsB* and *Acr3*

Resistance toward As[III] has been shown to be related to the presence of membrane efflux proteins which extrude this toxic species (Paez-Espino et al., 2009), that are included in the *ArsB* or in the *Acr3* families. We performed a phylogenetic analysis of

the proteins from *Exiguobacterium*, comparing them with other known protein sequences obtained from NCBI.

For the *ArsB* proteins, the resulting phylogenetic tree showed a distribution based on their affiliation, displaying two groups consisting of Gram negative and Gram positive members (**Figure 5A**). The sequences of the *Exiguobacterium* genus grouped with the latter. The topology of this subtree was identical to the topology observed in the 16S rRNA tree (**Figure 1A**).


On the other hand, the *Acr3*-like proteins have been classified into three subgroups, which are independent of the taxonomic affiliation (**Figure 5B**). The only known sequence from the *Exiguobacterium* genus, the protein *Acr3* from *Exiguobacterium* sp. S17, grouped in the third less characterized cluster, together with other proteins from Gram positive bacteria (Firmicutes and Actinobacteria).

Structural Analysis of *Acr3* and *ArsB* Efflux Pumps

To characterize the newly identified *acr3* gene in the *Exiguobacterium* genus we performed an structural bioinformatic approach to predict its structure and identify key residues. Initially four transmembrane segment predictors (TMHMM, DAS, HMMTOP, TOPCONS) and two general secondary structure predictors (PSIPRED, PORTER) were used in order to predict BsAcr3 topology (Figure S2, Supplementary Materials). We identified 10 transmembrane helices and the residues located in the cytoplasm and in the extracellular part. We selected the structure 3zuxA (Sodium Bile symporter) which has a 16% identity and 93% coverage (see Figure S1 in Supplementary Materials) to perform the modeling as described in the Materials and Methods Section. ProQM analysis of the model results in a quality score of 0.714 (0-low, 1-high), which represents a good model despite the low identity.

Moreover, for model validation, an alignment was computed with other two *Acr3* sequences: *Bacillus subtilis* *Acr3* (BsAcr3) (Aaltonen and Silow, 2008) and *Alkaliphilus metalliredigens* *Acr3* (AmAcr3) (Fu et al., 2009). BsAcr3 and AmAcr3 were chosen because the accessibility of certain residues for their localization in intracellular and extracellular loops has been studied. In the case of BsAcr3, accessibility was analyzed with PhoA and GFP. For AmAcr3 a scanning cysteine accessibility method (SCAM) (Bogdanov et al., 2005) was used to determine the transmembrane topology. The alignment between these three protein sequences was computed using HMMER3 (Finn et al., 2011) with the *Acr3* family HMM (TIGR00832—*acr3*: *arsenical-resistance protein*) from the TIGRFAM database (Haft et al., 2001). Using this information the residues that belong to intracellular and extracellular sides were mapped in the structure (**Figure 6**) and in the alignment (Figure S3 in Supplementary Materials).

As can be seen in **Figure 6A**, residues from S17 *Acr3* that aligned to residues previously determined to be outside by SCAM fall in loops outside the transmembrane regions. There is an exception for the leucine at position 137, but, as Aaltonen et al. pointed out -referring to a corresponding methionine in that position-, although that seems to contradict the transmembrane predictors due to its position in the middle of a transmembrane



helix (Aaltonen and Silow, 2008), there is a region that opens and close to transport the ligands across the membrane that could explain its solvent accessibility. Also, there are regions where a transmembrane segment presents a breakdown of alpha helix forming small helical hairpins reported to be a common motif in ion transporters (Scrapanti and Hunte, 2007). These helical hairpins are between positions residue 106–109 and residue 263–266, and seem to be important for Acr3 function because, as can be seen in **Figure 6B**, a conserved cysteine which is essential for the protein function is at position 107 which is in the core region of the protein placed in one of these hairpins.

Based on the analysis by Hu et al. on the Sodium/Bile Acid symporter that we have used as template structure (Hu et al., 2011), Acr3 seems to present two domains, a panel domain and a core domain, depicted in green and gray in **Figure 6B**. We also highlighted positively and negatively charged residues. As can be seen, the intracellular region is extremely populated with positive residues (Arg3, Arg57, Arg63, Arg177, Arg191, Arg241, Arg242, Arg249, Lys6, Lys119, Lys182, Lys245, Lys246) compared with the negative residues (Asp121, Asp251, Glu5, Glu190). Meanwhile, in the extracellular side, there are more negative residues (Asp33, Asp91, Glu26, Glu217, Glu250, Glu280) than positive ones (Arg29, Arg30, Arg213). The arsenite pKa is 9.2 and the *Exiguobacterium* genus is known to be alkaliphile so a few percent of arsenite would be ionized in the cytosolic side of the membrane involving positively charged residues in electrostatic interactions with arsenite due to its negative charge.

Interestingly, there are three charged residues (Arg264, Asp109, Glu294) in the core of transmembrane segments but particularly near to Cys107 and the four of them are placed right in the interface between the core domain and the panel domain. These residues could be involved in the arsenite binding together with the Cys residue. Finally, a protein cavity entrance can be appreciated in **Figure 6C**, which can be contrasted with **Figure 6B** to see how this entrance is between the two panels, it is


clear that previously mentioned leucine (Leu147 in S17 Acr3), as mentioned above, presents cytoplasmic accessibility evidence due to its position in the protein.

Transmembrane topology prediction was done for S17 *arsB* gene using sequence based predictors and results can be seen in Figure S4 and S5 (see Supplementary Materials). The *arsB* gene encodes a membrane protein of 428 amino acids in S17 and N139, showing an amino acid sequence homology of 93.5% identity between them. The amino acid sequence identity of ArsB from S17 with the protein from *E. sibiricum* 255-15 was 82.6%; *E. antarcticum* B7 83.3%; *E. pavilionensis* RW-2 93.8%; *Exiguobacterium* sp. AT1b 91.7% and *Exiguobacterium* sp. 8-11-1 93.8%, while N139-ArsB showed an identity of 83.3% with 255-15; 84.4% with B7, 98.9% with RW-2, 93.5% with AT1b and 96.4% with 8-11-1. It was observed that ArsB contains 11 probable transmembrane helices predicted by TMHMM2.0 in S17 and N139 (Figure S6, Supplementary Materials). Analysis of several published genome sequences of *Exiguobacterium* strains showed that the *arsB* gene is conserved in this genus and a similar activity is expected.

Discussion


In this work, three topics were specially emphasized. First, arsenic tolerance of *Exiguobacterium* sp. S17 and N139, isolated from extreme environments with different natural concentrations of arsenic, were compared. Second, the genomes of S17 and N139 were searched for arsenic resistance genes that were compared between strains. Finally, *in-silico* modeling was performed with the extrusion pumps ArsB and Acr3, and the first 3D model for the Acr3 protein was obtained to characterize this relevant gene.

With respect to the first topic, the tolerance to arsenic was also studied in the two mentioned strains plus a third strain as a control (DSM 6208). S17 was in fact the only strain able to tolerate and grow at concentrations above 5 mM arsenite (As[III]) (**Figures 2B,D**). In turn, all strains grew well



at increasing concentrations of arsenate (As[V]) (Figures 2A,C). Interestingly, growth was better with As[V] than for the non-supplemented cultures, implying that it somehow enhances growth. Belfiore et al. showed that glycolysis proteins seem to be more abundant in the S17 strain grown in presence of As[V], suggesting a more active metabolism which would explain the increased growth in a set timeframe (Belfiore et al., 2013).

While known arsenic resistance systems are the main factors responsible for the arsenic tolerance, additional mechanisms might also influence the extreme resistance of the S17 strain. A possible resistance mechanism could be the presence of methylases, as described for *Rhodospseudomonas palustris* (Qin et al., 2006) and cyanobacteria (Yin et al., 2011), although conserved arsenic-methylase genes were not detected neither



on the S17 genome (Ordoñez et al., 2013) nor proteome (Belfiore et al., 2013). Tolerance might also be enhanced by indirect mechanisms coping with other stresses generated by arsenic, such as oxidative stress. The proteomic study did detect overexpression of superoxide dismutase (SOD) and universal stress proteins, both in presence of As[III] and As[V] (Belfiore et al., 2013). These results are consistent with previous findings reported by Sacheti et al. (2014) and Ciprandi et al. (2012).

Regarding the second topic, whole genome sequencing and comparative genome analysis by BBHs of S17 and N139 allows us to determine that the genomes of both organisms share 2515 genes considered orthologous (Figure 3). Those genes correspond to 78.1 and 83.6% of the genes predicted in S17 and N139, respectively. BBH analysis indicates that S17 and N139 share all arsenic resistance genes, except for the presence of the Acr3 efflux pump in S17. Although we cannot be completely sure whether other detoxification systems exist in the genome and other genes could may not have been identified (Bakke et al., 2009; Poptsova and Gogarten, 2010; Richardson and Watson, 2013), the whole set of arsenic resistance genes was clearly identified and manual annotation was performed.

Comparative analysis of arsenic detoxification genes from sequenced *Exiguobacterium* genomes showed that most of the genes are organized in two versions of the *ars* operon, *arsRBC* being the simplest configuration (i.e., strains B7 and 255-15) and the other configuration includes the genes *arsRDAXB* (S17, N139, AT1b, 8-11-1, and RW2). Surprisingly, the S17 strain displayed also an uncommon version of a pump that expels cytoplasmic arsenite, the *acr3* gene (Figure 4). As was mentioned before, there are two efflux pumps, ArsB and Acr3, in this group of bacteria. While ArsB pump was found in all sequenced *Exiguobacterium* genomes, Acr3 was only found in the S17 strain. Moreover, in this work we demonstrated that the *acr3* gene in S17 is constitutively expressed as is observed in all conditions tested (Figure S7).

With respect to the third topic, the Acr3 efflux pump, we modeled the structure and showed that our structure is in agreement with previous experiments that identified

non-membrane residues. As mentioned before, a Cys residue conserved in all members of Acr3 family has been previously reported as the only cysteine required to achieve As(III) detoxification (Fu et al., 2009). In the case of this protein, Cys107 is the only cysteine in the protein, located right in the middle of a clear channel in our structure, that we now propose as the catalytic residue that binds arsenite.

Predicted topology is consistent with previous proposals for *Bacillus subtilis* and *Alkaliphilus metalliredigens* by Aaltonen et al. and Fu et al., respectively (Aaltonen and Silow, 2008; Fu et al., 2009). Ten transmembrane segments were predicted by transmembrane predictors and by homology modeling for Acr3 (Figure 6). Also, general purpose predictors are able to capture interesting secondary structure elements such as helical hairpins breaking two alpha helices in the same place as the conserved cysteine (Cys107) as mentioned in results section. This prediction agrees with the model based on a Sodium/Bile Acid symporter where there are two loops in the same region.

Based on the results presented here and previous works mentioned before, we propose that the main resistance mechanism of *Exiguobacterium* sp. S17 to arsenic would be the reduction of arsenate to arsenite and subsequent extrusion through membrane pumps as ArsB and Acr3 (Figure 7). Acr3 presents the conserved residues required for functionality as observed by Aaltonen and Silow (2008), Fu et al. (2009), and Maciaszczyk-Dziubinska et al. (2014) and would confer high resistance to arsenite (As[III]). Also, other mechanisms that overcome oxidative stress caused by arsenic would also help in the resistance to the metalloid.

The increased abundance of *acr3* gene in environments with high arsenic concentration has been reported in the literature (Achour et al., 2007; Cai et al., 2009), and this seems also the case for Socompa, where a variety of microorganisms isolated from this site bear a copy of the gene, as can be seen in the genomes of *Sphingomonas* sp. S17 (AFGG01), *Salinivibrio* sp. strains S10B, S34, and S35 (AQOE01, APMS01, AQOD01). The gene is also present in other isolates of similar locations from




FIGURE 7 | Schematic representation of genes and proteins involved in the arsenic resistance in *Exiguobacterium* sp. S17. The main arsenic resistance mechanism would be the reduction of arsenate to arsenite and subsequent extrusion through membrane pumps such as ArsB and Acr3.

the Argentinean Puna, namely *Halorubrum* sp. Strain AJ67 (CBVY01) and *Acinetobacter* sp. Ver3 (JFYLO1). These data highlights the importance of *acr3* in the environment, and is also supported by the absence of *acr3* in the known *Exiguobacterium* strains isolated from non-contaminated locations. Finally, while the HAAL strains presented higher resistance to arsenic than the control strain DSM 6208, the structural analysis of Acr3 and ArsB efflux pumps showed that the *arsB* gene is conserved in this genus, so enhanced resistance to arsenite (As[III]) in S17 may be explained by the combined work of the two membrane proteins ArsB and Acr3. Therefore, according to these

results, the appearance of Acr3 may explain the improvement in arsenic tolerance in environments with high arsenic content.

Author Contributions

OO and EL performed the research and wrote the paper. OO performed physiological experiments and data analysis, EL performed annotation, comparative genome analysis and structural modeling. DK carried out phylogenetic analysis and helped to modify the manuscript; MF, NC, and AT obtained funding for the original project idea. All authors read and approved this manuscript.

Acknowledgments

This work was supported by the Argentinean National Council for Science and Technology (CONICET; PIP 0365/12), the Argentinean National Fund for Science and Technology (FONCyT; project PICT numbers 1788 and 2805). We also want to thank Secretaria de medio ambiente de la provincia de Salta for his assistance in field trip. OO is a recipient of a postdoctoral fellowship from Bunge and Born Foundation and EL is a recipient of a doctoral fellowship from CONICET Argentina.

Supplementary Material

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fenvs.2015.00050>

References

- Aaltonen, E. K., and Silow, M. (2008). Transmembrane topology of the Acr3 family arsenite transporter from *Bacillus subtilis*. *Biochim. Biophys. Acta* 1778, 963–973. doi: 10.1016/j.bbame.2007.11.011
- Achour, A. R., Bauda, P., and Billard, P. (2007). Diversity of arsenite transporter genes from arsenic-resistant soil bacteria. *Res. Microbiol.* 158, 128–137. doi: 10.1016/j.resmic.2006.11.006
- Albarracín, V. H., Dib, J. R., Ordoñez, O. F., and Farias, M. E. (2011). “A harsh life of indigenous proteobacteria at the andean mountains: microbial diversity and resistance mechanisms towards extreme conditions,” *Book Series: Microbiology Research Advances*, ed M. L. Sezenna (Hauppauge, NY: Nova Publishers), 91–131.
- Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W., et al. (1997). Gapped blast and psi-blast: a new generation of protein database search programs. *Nucleic Acids Res.* 25, 3389–3402. doi: 10.1093/nar/25.17.3389
- Aziz, R. K., Bartels, D., Best, A. A., DeJongh, M., Disz, T., Edwards, R. A., et al. (2008). The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. doi: 10.1186/1471-2164-9-75
- Bakke, P., Carney, N., Deloache, W., Gearing, M., Ingvorsen, K., Lotz, M., et al. (2009). Evaluation of three automated genome annotations for *Halorhabdus utahensis*. *PLoS ONE* 4:e6291. doi: 10.1371/journal.pone.0006291
- Belfiore, C., Ordoñez, O. F., and Farias, M. E. (2013). Proteomic approach of adaptive response to arsenic stress in *Exiguobacterium* sp. S17, an extremophile strain isolated from a high-altitude Andean Lake stromatolite. *Extremophiles* 17, 421–431. doi: 10.1007/s00792-013-0523-y
- Bernsel, A., Viklund, H., Hennerdal, A., and Elofsson, A. (2009). TOPCONS: consensus prediction of membrane protein topology. *Nucleic Acids Res.* 37, W465–W468. doi: 10.1093/nar/gkp363
- Bogdanov, M., Zhang, W., Xie, J., and Dowhan, W. (2005). Transmembrane protein topology mapping by the substituted cysteine accessibility method (SCAM(TM)): application to lipid-specific membrane protein topogenesis. *Methods* 36, 148–171. doi: 10.1016/j.ymeth.2004.11.002
- Brown, J. R., Douady, C. J., Italia, M. J., Marshall, W. E., and Stanhope, M. J. (2001). Universal trees based on large combined protein sequence data sets. *Nat. Genet.* 28, 281–285. doi: 10.1038/90129
- Cai, L., Liu, G., Rensing, C., and Wang, G. (2009). Genes involved in arsenic transformation and resistance associated with different levels of arsenic-contaminated soils. *BMC Microbiol.* 9:4. doi: 10.1186/1471-2180-9-4
- Ciprandi, A., Baraúna, R. A., Santos, A. V., Gonçalves, E. C., Carepo, M. S. P., Schneider, M. P. C., et al. (2012). Proteomic response to arsenic stress in *Chromobacterium violaceum*. *J. Integr. OMICS* 2, 69–73. doi: 10.5584/jiomics.v2i1.84hh
- Cserző, M., Wallin, E., Simon, I., Von Heijne, G., and Elofsson, A. (1997). Prediction of transmembrane alpha-helices in prokaryotic membrane proteins: the dense alignment surface method. *Protein Eng.* 10, 673–676. doi: 10.1093/protein/10.6.673
- Dib, J., Motok, J., Zenoff, V. F., Ordoñez, O., and Farias, M. E. (2008). Occurrence of resistance to antibiotics, UV-B, and arsenic in bacteria isolated from extreme environments in high-altitude (above 4400 m) andean wetlands. *Curr. Microbiol.* 56, 510–517. doi: 10.1007/s00284-008-9103-2
- Dib, J., Weiss, A., Neumann, A., Ordoñez, O., Estevez, M. C., and Farias, M. E. (2009). Isolation of bacteria from remote high altitude Andean lakes able to grow in the presence of antibiotics. *Recent Pat. Antinfect. Drug Discov.* 4, 66–76. doi: 10.2174/157489109787236300
- Escalante, G., Campos, V. L., Valenzuela, C., Yañez, J., Zaror, C., and Mondaca, M. A. (2009). Arsenic resistant bacteria isolated from arsenic contaminated river in

- the Atacama Desert (Chile). *Bull. Environ. Contam. Toxicol.* 83, 657–661. doi: 10.1007/s00128-009-9868-4
- Escudero, L., Chong, G., Demergasso, C., Farias, M. E., Cabrol, N. A., Grin, E. A., et al. (2007). "Investigating microbial diversity and UV radiation impact at the high altitude lake Aguas Calientes, Chile," in *Proceedings SPIE Conference on Instruments, Methods, and Missions for Astrobiology*, Vol. 1, (San Diego, CA), 6694–6633.
- Farias, M. E., Contreras, M., Rasuk, M. C., Kurth, D., Flores, M. R., Poiré, D. G., et al. (2014). Characterization of bacterial diversity associated with microbial mats, gypsum evaporites and carbonate microbialites in thalassic wetlands: Tebenquiche and La Brava, Salar de Atacama, Chile. *Extremophiles* 18, 311–329. doi: 10.1007/s00792-013-0617-6
- Farias, M. E., Poire, D. G., Arrouy, M. J., and Albarracín, V. H. (2011). "Modern stromatolite ecosystems at alkaline and hypersaline high-altitude lakes at the Argentinean Puna," in *STROMATOLITES: Interaction of Microbes with Sediments Cellular Origin, Life in Extreme Habitats and Astrobiology*, Vol. 18, eds J. Seckbach and V. Tewari (New York, NY: Springer), 427–441.
- Farias, M. E., Rascovan, N., Toneatti, D. M., Albarracín, V. H., Flores, M. R., Poire, D. G., et al. (2013). The discovery of Stromatolites developing at 3570 m above sea level in a high-altitude volcanic lake Socompa, Argentinean Andes. *PLoS ONE* 8:E53497. doi: 10.1371/journal.pone.0053497
- Fernandez Zenoff, V., Sineriz, F., and Farias, M. E. (2006). Diverse responses to UV-B radiation and repair mechanisms of bacteria isolated from high-altitude aquatic environments. *Appl. Environ. Microbiol.* 72, 7857–7863. doi: 10.1128/AEM.01333-06
- Finn, R. D., Clements, J., and Eddy, S. R. (2011). HMMER web server: interactive sequence similarity searching. *Nucleic Acids Res.* 39, W29–W37. doi: 10.1093/nar/gkr367
- Flores, M. R., Ordoñez, O. F., Maldonado, J., and Farias, M. E. (2009). Isolation of UV-B resistant bacteria from two high altitude Andean lakes (4,400 m) with saline and non saline conditions. *J. Gen. Appl. Microbiol.* 55, 447–458. doi: 10.2323/jgam.55.447
- Fu, H. L., Meng, Y., Ordóñez, E., Villadangos, A. F., Bhattacharjee, H., Gil, J. A., et al. (2009). Properties of arsenite efflux permeases (Acr3) from *Alkaliphilus metalliredigens* and *Corynebacterium glutamicum*. *J. Biol. Chem.* 284, 19887–19895. doi: 10.1074/jbc.M109.011882
- Haft, D. H., Loftus, B. J., Richardson, D. L., Yang, F., Eisen, J. A., Paulsen, I. T., et al. (2001). TIGRFAMs: a protein family resource for the functional identification of proteins. *Nucleic Acids Res.* 29, 41–43. doi: 10.1093/nar/29.1.41
- Hrimpeng, K., Prapagdee, B., Banjerdikij, P., Vattanaviboon, P., Dubbs, J. M., and Mongkolsuk, S. (2006). Challenging *Xanthomonas Campestris* with low levels of arsenic mediates cross-protection against oxidant killing. *FEMS Microbiol. Lett.* 262, 121–127. doi: 10.1111/j.1574-6968.2006.00383.x
- Hu, N. J., Iwata, S., Cameron, A. D., and Drew, D. (2011). Crystal structure of a bacterial homologue of the bile acid sodium symporter ASBT. *Nature* 478, 408–411. doi: 10.1038/nature10450
- Jones, D. T. (1999). Protein secondary structure prediction based on position-specific scoring matrices. *J. Mol. Biol.* 292, 195–202. doi: 10.1006/jmbi.1999.3091
- Krogh, A., Larsson, B., Von Heijne, G., and Sonnhammer, E. L. (2001). Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J. Mol. Biol.* 305, 567–580. doi: 10.1006/jmbi.2000.4315
- Lara, J., Escudero González, L., Ferrero, M., Chong Díaz, G., Pedrós-Alió, C., and Demergasso, C. (2012). Enrichment of arsenic transforming and resistant heterotrophic bacteria from sediments of two salt lakes in Northern Chile. *Extremophiles* 16, 523–538. doi: 10.1007/s00792-012-0452-1
- Larkin, M. A., Blackshields, G., Brown, N. P., Chenna, R., McGettigan, P. A., McWilliam, H., et al. (2007). Clustal W and Clustal X version 2.0. *Bioinformatics* 23, 2947–2948. doi: 10.1093/bioinformatics/btm404
- Li, W., Jaroszewski, L., and Godzik, A. (2001). Clustering of highly homologous sequences to reduce the size of large protein databases. *Bioinformatics* 17, 282–283. doi: 10.1093/bioinformatics/17.3.282
- Lopez, L., Pozo, C., Rodelas, B., Calvo, C., Juarez, B., Martinez-Toledo, M. V., et al. (2005). Identification of bacteria isolated from an oligotrophic lake with pesticide removal capacities. *Ecotoxicology* 14, 299–312. doi: 10.1007/s10646-003-6367-y
- López-Maury, L., Florencio, F. J., and Reyes, J. C. (2003). Arsenic sensing and resistance system in the *Cyanobacterium Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* 185, 5363–5371. doi: 10.1128/JB.185.18.5363-5371.2003
- Maciaszczyk-Dziubinska, E., Migocka, M., Wawrzycka, D., Markowska, K., and Wysocki, R. (2014). Multiple cysteine residues are necessary for sorting and transport activity of the arsenite permease Acr3p from *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* 1838, 747–755. doi: 10.1016/j.bbamem.2013.11.013
- Maciaszczyk-Dziubinska, E., Migocka, M., and Wysocki, R. (2011). Acr3p is a plasma membrane antiporter that catalyzes As(III)/H(+) and Sb(III)/H(+) exchange in *Saccharomyces Cerevisiae*. *Biochim. Biophys. Acta* 1808, 1855–1859. doi: 10.1016/j.bbamem.2011.03.014
- Macy, J. M., Santini, J. M., Pauling, B. V., O'neill, A. H., and Sly, L. I. (2000). Two new arsenate/sulfate-reducing bacteria: mechanisms of arsenate reduction. *Arch. Microbiol.* 173, 49–57. doi: 10.1007/s002030050007
- Mansour, N. M., Sawhney, M., Tamang, D. G., Vogl, C., and Saier, M. H. Jr. (2007). The Bile/arsenite/riboflavin transporter (Bart) superfamily. *FEBS J.* 274, 612–629. doi: 10.1111/j.1742-4658.2006.05627.x
- Mantelli, F., Scala, C., Ronchi, A., Gatti, A., and Minoia, C. (2003). Macroconstituenti ed elementi in traccia nelle acque dei laghi salini delle Ande de Catamarca e la Rioja (Argentina). *Boll. Chim. Ig.* 54, 37–44.
- Okeke, B. C., Laymon, J., Oji, C., and Crenshaw, S. (2007). "Rapid bioreduction of hexavalent chromium in water by *Exiguobacterium* sp. GS1," in *ASM General 107th Meeting* (Toronto, ON: ASM Press), Q-199.
- Ordoñez, O. F., Flores, M. R., Dib, J. R., Paz, A., and Farias, M. E. (2009). Extremophile culture collection from Andean lakes: extreme pristine environments that host a wide diversity of microorganisms with tolerance to UV radiation. *Microb. Ecol.* 58, 461–473. doi: 10.1007/s00248-009-9527-7
- Ordoñez, O. F., Lanzarotti, E., Kurth, D., Gorriti, M. F., Revale, S., Cortez, N., et al. (2013). Draft genome sequence of the polyextremophilic *Exiguobacterium* sp. strain S17, isolated from hyperarsenic lakes in the Argentinian Puna. *Genome Announc.* 1, pii: e00480–13. doi: 10.1128/genomeA.00480-13
- Oremland, R. S., Stolz, J. F., and Hollibaugh, J. T. (2004). The microbial arsenic cycle in Mono Lake, California. *FEMS Microb. Ecol.* 48, 15–27. doi: 10.1016/j.femsec.2003.12.016
- Overbeek, R., Fonstern, M., D'Souza, M., Pusch, G. D., and Maltsev, N. (1999). The use of gene clusters to infer functional coupling. *Proc. Natl. Acad. Sci. U.S.A.* 96, 2896–2901. doi: 10.1073/pnas.96.6.2896
- Paez-Espino, D., Tamames, J., De Lorenzo, V., and Canovas, D. (2009). Microbial responses to environmental arsenic. *Biomaterials* 22, 117–130. doi: 10.1007/s10534-008-9195-y
- Pandey, S., Shrivastava, A. K., Singh, V. K., Rai, R., Singh, P. K., Rai, S., et al. (2012). A New arsenate reductase involved in arsenic detoxification in *Anabaena* Sp. Pcc7120. *Funct. Integr. Genom.* 13, 43–55. doi: 10.1007/s10142-012-0296-x
- Pattanapitpaisal, P., Mabbett, A. N., Finlay, J. A., Beswick, A. J., Paterson-Beedle, M., Essa, A., et al. (2002). Reduction of Cr(VI) and bioaccumulation of chromium by Gram-positive and Gram-negative microorganisms not previously exposed to Cr-stress. *Environ. Technol.* 23, 731–745. doi: 10.1080/09593332308618367
- Poirel, J., Joulian, C., Leyval, C., and Billard, P. (2013). Arsenite-induced changes in abundance and expression of arsenite transporter and arsenite oxidase genes of a soil microbial community. *Res. Microbiol.* 164, 457–465. doi: 10.1016/j.resmic.2013.01.012
- Pollastri, G., and McLysaght, A. (2005). Porter: a new, accurate server for protein secondary structure prediction. *Bioinformatics* 21, 1719–1720. doi: 10.1093/bioinformatics/bti203
- Poptsova, M. S., and Gogarten, J. P. (2010). Using comparative genome analysis to identify problems in annotated microbial genomes. *Microbiology* 156, 1909–1917. doi: 10.1099/mic.0.033811-0
- Qin, J., Rosen, B. P., Zhang, Y., Wang, G., Franke, S., and Rensing, C. (2006). Arsenic detoxification and evolution of trimethylarsine gas by a microbial arsenite S-adenosylmethionine methyltransferase. *Proc. Natl. Acad. Sci. U.S.A.* 103, 2075–2080. doi: 10.1073/pnas.0506836103
- Rasuk, M. C., Kurth, D., Flores, M. R., Contreras, M., Novoa, F., Poiré, D. G., et al. (2014). Microbial characterization of microbial ecosystems associated to evaporites domes of gypsum in Salar de Llamara in Atacama desert. *Microb. Ecol.* 68, 483–494. doi: 10.1007/s00248-014-0431-4

- Ray, A., Lindahl, E., and Wallner, B. (2010). Model quality assessment for membrane proteins. *Bioinformatics* 26, 3067–3074. doi: 10.1093/bioinformatics/btq581
- Richardson, E. J., and Watson, M. (2013). The automatic annotation of bacterial genomes. *Brief. Bioinformatics* 14, 1–12. doi: 10.1093/bib/bbs007
- Rodrigues, D. F., and Tiedje, J. M. (2007). Multi-locus real-time pcr for quantitation of bacteria in the environment reveals *Exiguobacterium* to be prevalent in permafrost. *FEMS Microb. Ecol.* 59, 489–499. doi: 10.1111/j.1574-6941.2006.00233.x
- Romero, L., Alonso, H., Campano, P., Fanfani, L., Cidu, R., Dadea, C., et al. (2003). Arsenic enrichment in waters and sediments of the Rio Loa (Second Region, Chile). *Appl. Geochem.* 18, 1399–1416. doi: 10.1016/S0883-2927(03)00059-3
- Rosen, B. P. (1999). Families of arsenic transporters. *Trends Microbiol.* 7, 207–212. doi: 10.1016/S0966-842X(99)01494-8
- Rosen, B. P., Bhattacharjee, H., Zhou, T., and Walmsley, A. R. (1999). Mechanism of the ArsA ATPase. *Biochim. Biophys. Acta* 1461, 207–215. doi: 10.1016/S0005-2736(99)00159-5
- Sacheti, P., Patil, R., Dube, A., Bhonsle, H., Thombre, D., Marathe, S., et al. (2014). Proteomics of arsenic stress in the gram-positive organism *Exiguobacterium* sp. PS NCIM 5463. *Appl. Microbiol. Biotechnol.* 98, 6761–6773. doi: 10.1007/s00253-014-5873-6
- Saitou, N., and Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4, 406–425.
- Sali, A., and Blundell, T. L. (1993). Comparative protein modelling by satisfaction of spatial restraints. *J. Mol. Biol.* 234, 779–815. doi: 10.1006/jmbi.1993.1626
- Screpanti, E., and Hunte, C. (2007). Discontinuous membrane helices in transport proteins and their correlation with function. *J. Struct. Biol.* 159, 261–267. doi: 10.1016/j.jsb.2007.01.011
- Seufferheld, M. J., Alvarez, H., and Farias, M. E. (2008). Role of polyphosphates in microbial adaptation to extreme environments. *Appl. Environ. Microbiol.* 74, 5867–5874. doi: 10.1128/AEM.00501-08
- Silver, S., and Phung, L. T. (2005). Genes and enzymes involved in bacterial oxidation and reduction of inorganic arsenic. *Appl. Environ. Microbiol.* 71, 599–608. doi: 10.1128/AEM.71.2.599-608.2005
- Stolz, J. F., and Oremland, R. S. (1999). Bacterial respiration of arsenic and selenium. *FEMS Microbiol. Rev.* 23, 615–627. doi: 10.1111/j.1574-6976.1999.tb00416.x
- Suzek, B. E., Huang, H., McGarvey, P., Mazumder, R., and Wu, C. H. (2007). UniRef: comprehensive and non-redundant UniProt reference clusters. *Bioinformatics* 23, 1282–1288. doi: 10.1093/bioinformatics/btm098
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., and Kumar, S. (2011). Mega5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731–2739. doi: 10.1093/molbev/msr121
- Thompson, C. C., Vicente, A. C., Souza, R. C., Vasconcelos, A. T., Vesth, T., Alves, N. J., et al. (2009). Genomic taxonomy of *Vibrios*. *BMC Evol. Biol.* 27:258. doi: 10.1186/1471-2148-9-258
- Tusnády, G. E., and Simon, I. (2001). The HMMTOP transmembrane topology prediction server. *Bioinformatics* 17, 849–850. doi: 10.1093/bioinformatics/17.9.849
- Valko, M., Morris, H., and Cronin, M. T. (2005). Metals toxicity and oxidative stress. *Curr. Med. Chem.* 12, 1161–1208. doi: 10.2174/0929867053764635
- Villadangos, A. F., Fu, H. L., Gil, J. A., Messens, J., Rosen, B. P., and Mateos, L. M. (2012). Efflux permease CgAcr3-1 of *Corynebacterium glutamicum* is an arsenite-specific antiporter. *J. Biol. Chem.* 287, 723–735. doi: 10.1074/jbc.M111.263335
- Vishnivetskaya, T. A., Kathariou, S., and Tiedje, J. M. (2009). The *Exiguobacterium* genus: biodiversity and biogeography. *Extremophiles* 13, 541–555. doi: 10.1007/s00792-009-0243-5
- Vishnivetskaya, T. A., Petrova, M. A., Urbance, J., Ponder, M., Moyer, C. L., Gilichinsky, D. A., et al. (2006). Bacterial community in ancient siberian permafrost as characterized by culture and culture-independent methods. *Astrobiology* 6, 400–414. doi: 10.1089/ast.2006.6.400
- Wysocki, R., Clemens, S., Augustyniak, D., Golik, P., Maciaszczyk, E., Tamas, M. J., et al. (2003). Metalloid tolerance based on phytochelatin is not functionally equivalent to the arsenite transporter Acr3p. *Biochem. Biophys. Res. Commun.* 304, 293–300. doi: 10.1016/S0006-291X(03)00584-9
- Yin, X. X., Chen, J., Qin, J., Sun, G. X., Rosen, B. P., and Zhu, Y. G. (2011). Biotransformation and volatilization of arsenic by three photosynthetic cyanobacteria. *Plant Physiol.* 156, 1631–1638. doi: 10.1104/pp.111.178947

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Ordoñez, Lanzarotti, Kurth, Cortez, Farias and Turjanski. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.