

Lawrence Berkeley National Laboratory

Lawrence Berkeley National Laboratory

Title

Genomic Characterization of Methanomicrobiales Reveals Three Classes of Methanogens

Permalink

<https://escholarship.org/uc/item/7zt6c5pt>

Author

Anderson, Iain

Publication Date

2009-10-06

Genomic Characterization of Methanomicrobiales Reveals Three Classes of Methanogens

Iain Anderson¹, Luke E. Ulrich², Boguslaw Lupa³, Dwi Susanti^{4,7}, Iris Porat^{3,11}, Sean D. Hooper¹, Athanasios Lykidis¹, Magdalena Sieprawska-Lupa³, Lakshmi Dharmarajan^{4,7}, Eugene Goltsman¹, Alla Lapidus¹, Elizabeth Saunders⁸, Cliff Han⁸, Miriam Land⁹, Susan Lucas¹, Biswarup Mukhopadhyay^{4,5,6,7}, William B. Whitman³, Carl Woese¹⁰, James Bristow¹, and Nikos Kyrpides¹

¹Joint Genome Institute, 2800 Mitchell Drive, Walnut Creek, California, USA

²Joint Institute for Computational Sciences, University of Tennessee – Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

³Department of Microbiology, University of Georgia, Athens, Georgia, USA

⁴Virginia Bioinformatics Institute and Departments of ⁵Biochemistry and ⁶Biological Sciences, ⁷Genetics, Bioinformatics and Computational Biology Graduate Program, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, USA

⁸Joint Genome Institute, Los Alamos National Laboratory, Bioscience Division, Los Alamos, New Mexico, USA

⁹Bioscience Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

¹⁰Department of Microbiology, University of Illinois, 601 S. Goodwin Ave., Urbana, Illinois, USA

¹¹Present address: Biological and Environmental Sciences Directorate, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

Corresponding author: Iain Anderson
Joint Genome Institute
2800 Mitchell Drive
Walnut Creek, CA 94598
Phone: 925-296-5784
Fax: 925-296-5850
Email: IJAnderson@lbl.gov

Abstract

Background

Methanomicrobiales is the least studied order of methanogens. While these organisms appear to be more closely related to the Methanosarcinales in ribosomal-based phylogenetic analyses, they are metabolically more similar to Class I methanogens.

Methodology/Principal Findings

In order to improve our understanding of this lineage, we have completely sequenced the genomes of two members of this order, *Methanocorpusculum labreanum* Z and *Methanoculleus marisnigri* JR1, and compared them with the genome of a third, *Methanospirillum hungatei* JF-1. Similar to Class I methanogens, Methanomicrobiales use a partial reductive citric acid cycle for 2-oxoglutarate biosynthesis, and they have the Eha energy-converting hydrogenase. In common with Methanosarcinales, Methanomicrobiales possess the Ech hydrogenase and at least some of them may couple formylmethanofuran formation and heterodisulfide reduction to transmembrane ion gradients. Uniquely, *M. labreanum* and *M. hungatei* contain hydrogenases similar to the *Pyrococcus furiosus* Mbh hydrogenase, and all three Methanomicrobiales have anti-sigma factor and anti-anti-sigma factor regulatory proteins not found in other methanogens. Phylogenetic analysis based on seven core proteins of methanogenesis and cofactor biosynthesis places the Methanomicrobiales equidistant from Class I methanogens and Methanosarcinales.

Conclusions/Significance

Our results indicate that Methanomicrobiales, rather than being similar to Class I methanogens or Methanosarcinales, share some features of both and have some unique

properties. We find that there are three distinct classes of methanogens: the Class I methanogens, the Methanomicrobiales (Class II), and the Methanosarcinales (Class III).

Introduction

The Archaea were discovered to form a distinct domain in 1977 [1] and subsequently were found to be comprised of two major kingdoms, the Crenarchaeota and the Euryarchaeota [2]. The Crenarchaeota consist mainly of thermophiles and thermoacidophiles while the Euryarchaeota contains a wider variety of organisms including the methanogens, the extreme halophiles, thermophiles, and thermoacidophiles. Recently a third kingdom, Thaumarchaeota, has been proposed that includes mesophilic organisms previously classified as Crenarchaeota [3].

Methanogens play a major role in the global carbon cycle [4] by carrying out the final steps in the anaerobic degradation of organic material. In the process, they are estimated to produce close to 400 million metric tons of methane per year. Much of the methane is converted back to carbon dioxide by methanotrophs, but some is released to the atmosphere where it is a potent greenhouse gas. As a result of human activities, the concentration of methane in the atmosphere has almost tripled in the last 200 years [5].

Methanogens are currently classified in five orders: Methanobacteriales, Methanococcales, Methanopyrales, Methanomicrobiales, and Methanosarcinales. It has been recognized that the methanogens can be divided into two major groups based on phylogenetic analysis [6,7]. The first group contains the orders Methanobacteriales, Methanococcales, and Methanopyrales, and has been named Class I methanogens by Baptiste et al. [7]. The second group, the Class II methanogens, includes Methanomicrobiales and Methanosarcinales. However, the Methanomicrobiales are physiologically more similar to the Class I methanogens than to the Methanosarcinales, growing on H_2/CO_2 or formate, while members of the Methanosarcinales can produce

methane from acetate, methanol, methylamines, and other C-1 compounds. Recently Thauer et al. [8] have argued that methanogens can be divided into two groups based on the presence or lack of cytochromes, with Methanosarcinales alone possessing cytochromes. The Methanomicrobiales thus belong to the phylogenetic group of Class II methanogens, but to the physiological group of methanogens without cytochromes.

Members of the order Methanomicrobiales have few known unique properties. However their membrane lipid composition is distinctive, and they are unique in possessing aminopentane tetrols in their lipids (reviewed in [9]). In addition to growth on H₂/CO₂ or formate, some are capable of using secondary alcohols as electron donors [10]. Methanomicrobiales have been detected in marine environments, in landfills and wastewater reactors, and as symbionts of ciliates (reviewed in [9]).

This is the first publication to describe genomes from the order Methanomicrobiales. We report here the genome of the marine methanogen *Methanoculleus marisnigri* JR1 [11] and that of *Methanocorpusculum labreanum* Z, a methanogen isolated from tar pit sediments [12]. We include comparisons of these two with the genome sequence of *Methanospirillum hungatei* JF-1, a spiral-shaped methanogen isolated from sewage sludge [13]. We also present a comparative analysis of Methanomicrobiales genomes with those of Methanosarcinales and Class I methanogens.

Results

General features

The genomes of *M. labreanum* and *M. marisnigri* consist of one chromosome and no plasmids (Table 1), and the same is true for *M. hungatei*. The size of the *M. hungatei* genome is substantially larger than those of the other two. *M. marisnigri* has only one

ribosomal RNA (rRNA) operon, while *M. labreanum* has three and *M. hungatei* has four. In two of the *M. hungatei* rRNA operons, there are two copies of the 5S rRNA.

Methanogenesis

As expected, the three Methanomicrobiales have all of the genes required for methanogenesis from hydrogen and carbon dioxide. All three species are capable of utilizing formate, and they have formate transporters as well as cytosolic formate dehydrogenases that probably reduce coenzyme F₄₂₀. No homologs were found to C-1 compound:corrinoid methyltransferases, corrinoid proteins, and methylcobalamin:Coenzyme M methyltransferases involved in methanogenesis from methanol and methylamines (no BLAST hit to *Methanosarcina acetivorans* proteins with 10⁻⁵ cutoff value). Some methanogens, including *M. marisnigri*, can utilize secondary alcohols as electron donors for methanogenesis [10], whereas *M. hungatei* JF-1 can not [14], and *M. labreanum* has not been tested. Alcohol dehydrogenases that oxidize secondary alcohols and use the electrons to reduce coenzyme F₄₂₀ have been characterized [15] and the structure has been determined for one enzyme [16]. *M. marisnigri* has a gene (Memar_0783) that is closely related to the F₄₂₀-dependent secondary alcohol dehydrogenase from *Methanoculleus thermophilus*, but *M. labreanum* and *M. hungatei* do not (no BLAST hit with cutoff of 10⁻¹⁰).

In Class I methanogens, F₄₂₀-non-reducing hydrogenase provides electrons to heterodisulfide reductase, and its D subunit interfaces with heterodisulfide reductase [17]. In all three Methanomicrobiales the gene for the D subunit of the hydrogenase (Mlab_0242, Memar_0622, Mhun_1839) is adjacent to the genes for heterodisulfide reductase, but phylogenetic analysis of hydrogenase alpha subunits (not shown) reveals

that only *M. marisnigri* possesses the F₄₂₀-non-reducing hydrogenase (Memar_1007-1008) (Table 2). Apparently, *M. labreanum* and *M. hungatei* use a different source of electrons for their heterodisulfide reductase. Based on the lack of F₄₂₀-nonreducing hydrogenase and the fact that the Eha hydrogenase is located adjacent to formylmethanofuran dehydrogenase (Fmd) (see below), we propose that, in at least some Methanomicrobiales, Fmd and heterodisulfide reductase are linked to transmembrane proton or sodium ion transport (Figure 1) rather than flavin-based electron bifurcation as proposed by Thauer et al. [8].

M. labreanum has a hydrogen-forming methylene-tetrahydromethanopterin dehydrogenase (COG4074), an enzyme previously found only in Class I methanogens. The other two genomes lack genes assigned to COG4074 and thus are unlikely to have this enzyme. The enzyme functions under conditions of nickel limitation (reviewed in [18]), so this suggests that *M. labreanum* can tolerate lower environmental nickel concentrations. When this gene is found in a Class I methanogen genome, it is often accompanied by one or two paralogs of unknown function belonging to COG4007. However, *M. labreanum* lacks these paralogs.

Membrane-Bound Hydrogenases

Methanogens have several families of membrane-bound hydrogenases that participate in various processes including methanogenesis and biosynthesis (reviewed in [19]). These hydrogenases are encoded by a core of conserved genes that includes from six to more than 20 subunits. The three Methanomicrobiales genomes encode two to three membrane-bound hydrogenases (Table 2). All three possess the genes for a membrane-bound hydrogenase similar to that encoded by the *Methanothermobacter*

thermautotrophicus eha operon (Memar_1172-1185, Mlab_0561-0573, Mhun_2094-2106). Their genes for the enzyme subunits are in the same order as those in the *M. thermautotrophicus eha* operon. However, some of the smaller subunits have diverged so extensively that homology can not be detected, and subunits A and M are absent. Adjacent to the hydrogenase operon are genes for the subunits of formylmethanofuran dehydrogenase, suggesting that the Eha hydrogenase may reduce the ferredoxin used by this enzyme (Figure 1).

All three genomes also have a six-subunit membrane-bound hydrogenase operon similar to Ech hydrogenase (Mlab_1619-1624, Memar_0359-0364, Mhun_1741-1746), which has multiple functions in *Methanosarcina barkeri* [20]. *M. labreanum* and *M. hungatei*, but not *M. marisnigri*, also have an operon very similar to the *mbh* operon of *Pyrococcus furiosus*. Since this hydrogenase is found in the two Methanomicrobiales genomes that lack F₄₂₀-nonreducing hydrogenase, the Mbh hydrogenase may be involved in heterodisulfide reduction (Figure 1). *M. hungatei* has another operon similar to membrane-bound hydrogenases (Mhun_1817-1822). Homologous operons are absent from the other two Methanomicrobiales, but they are found in two Methanosarcinales, *Methanosarcina acetivorans* and *Methanosarcina mazei*. However, the hydrogenase large subunits of these operons appear to lack the cysteine residues necessary for binding to the nickel-iron center, so these operons may not encode hydrogenases.

Metabolism and Transport

The Embden-Meyerhof pathway is present in many methanogens, where it is thought to play a role in the metabolism of stored glycogen. Although *M. hungatei* and *M. marisnigri* have putative glycogen phosphorylases (Mhun_1203, Memar_1262,

Memar_2480), and *M. marisnigri* has a putative glycogen branching enzyme (Memar_1265), none of the three Methanomicrobiales has an identifiable glycogen synthase. *M. marisnigri* and *M. hungatei* appear to encode a complete glycolysis pathway. This suggests that they may be able to utilize glucose from the environment (although they lack identifiable sugar transporters) or that they have a novel glycogen synthase. *M. hungatei* was previously reported to lack phosphofructokinase activity [21], but the genome contains two putative phosphofructokinase genes (Mhun_0556 and Mhun_1465). The Embden-Meyerhof pathway appears to be absent from *M. labreanum* as it lacks both phosphofructokinase and pyruvate kinase. A gluconeogenesis pathway is present in all three, as it is necessary for biosynthesis of pentoses and hexoses.

The pathway for 2-oxoglutarate production differs significantly between the Methanosarcinales and the Class I methanogens. Methanosarcinales generate 2-oxoglutarate through a partial oxidative TCA cycle with isocitrate as an intermediate, while Class I methanogens use a partial reductive TCA cycle with succinate as an intermediate (Figure 2). Methanomicrobiales appear to use the partial reductive TCA cycle, similar to the Class I methanogens, as they have genes for all of the necessary enzymes and they lack genes for citrate synthase and isocitrate dehydrogenase. They possess genes encoding the two subunits of the predicted archaeal aconitase [22], but this enzymatic activity has not been verified experimentally.

Sigma Factor Regulators

Both *M. labreanum* and *M. marisnigri* contain an anti-anti-sigma factor (Memar_02467, Mlab_1451), an anti-sigma factor (Memar_2469, Mlab_1452), and a serine phosphatase (Memar_2468, Mlab_1450) that are similar to the

SpoIIAA/SpoIIAB/SpoIIIE components of the *Bacillus subtilis* sporulation pathway. Moreover, these SpoII-type proteins are also found in *M. hungatei*, but not outside the order Methanomicrobiales. This finding is intriguing given that no *bona fide* sigma factors have been identified in Archaea. Kyrpides and Ouzounis identified proteins in *M. jannaschii* with similarity to conserved region 4 of bacterial sigma factors [23], and the Methanomicrobiales have homologs of three of these proteins (MJ0173, MJ0272, and MJ1243). However, the SpoIIAB anti-sigma factor binds to three separate regions of sigma F [24] corresponding to conserved regions 2, 3, and 4, and regions 2 and 3 are not present in the archaeal proteins. Therefore the targets of these archaeal anti-sigma factors can not be determined from the genome sequence.

Phylogenetic Analysis of Enzymes for Methanogenesis and Cofactor Biosynthesis

Bapteste et al. [7] determined the relationships among the various groups of methanogens by generating phylogenetic trees for enzymes of methanogenesis and cofactor biosynthesis. Their analysis found that methanogens could be divided into two groups: Class I and Class II methanogens. We present here an updated analysis that includes additional sequenced genomes. Furthermore, the protein-coding genes that we used in the analysis (see Materials and Methods) are present in only one copy per genome. Inclusion of the additional genomes reveals that, surprisingly, Methanomicrobiales are equally distant from Class I methanogens and from the Methanosarcinales (Figure 3). Therefore there appear to be three distinct classes of methanogens: the Class I methanogens, the Methanomicrobiales (that we have termed Class II methanogens), and the Methanosarcinales (that we have termed Class III methanogens).

Comparative genomics of methanogens

Now that several sequenced genomes from the order Methanomicrobiales are available, it is possible to carry out comparative genomic analyses between this order and the other methanogens. We used a protein clustering method to identify and cluster related proteins from 15 species representing Class I, II, and III methanogens (see Materials and Methods for the list of organisms). We then searched for the signature clusters, i.e. clusters of homologous proteins that are present in all members of a phylogenetic group and absent from other groups. Of particular interest are the exclusive signature clusters, those whose member proteins are found in all sequenced genomes from only one class. We also identified shared signature clusters (present in only two classes) and common signature clusters (present in all three).

We found 413 common signature clusters (Figure 4, Supplementary Table S1). These proteins are involved primarily in core information processing and essential metabolic activities (i.e. transcription, translation, methanogenesis, etc.). We found 62 exclusive signature clusters for Methanomicrobiales, 24 for Class I methanogens, and 48 for Methanosarcinales. Given the relatively close phylogenetic relationship between Methanomicrobiales and Methanosarcinales in ribosomal RNA and ribosomal protein-based trees, it is surprising that they share only 33 clusters to the exclusion of the Class I methanogens. While this is more than either class shares with the Class I methanogens, it represents but a very small proportion of the genome. In the following sections we describe some of the signature proteins associated with each of the three classes, as well as those shared by Methanomicrobiales and Methanosarcinales.

Class I: The Class I methanogen exclusive signature clusters include two LSU ribosomal proteins (L34E and L14E) and three enzymes of coenzyme M (CoM) biosynthesis (phosphosulfolactate synthase, phosphosulfolactate phosphatase, and sulfolactate dehydrogenase). This suggests that other methanogens possess either unrelated genes for these enzymes or a different pathway for CoM biosynthesis. Also present in only Class I methanogens is 2-phosphoglycerate kinase, an enzyme used in the synthesis of cyclic 2,3-diphosphoglycerate, which is thought to be a thermoprotectant. Its presence in mesophilic Class I methanogens suggests that it carries out a different function in these organisms. The second enzyme of the pathway, cyclic 2,3-diphosphoglycerate synthetase, is found only in a subset of Class I methanogens and is not part of the signature.

All Class I methanogens also have a homolog of seryl-tRNA(Sec) selenium transferase, used for the synthesis of selenocysteine in bacteria. However, this gene is likely to have a different function in archaea because not all of the Class I methanogens use selenocysteine [25], and those that do utilize a different pathway for selenocysteine synthesis, one that is shared with eukaryotes [26,27]. Experimental testing of this protein found that it did not catalyze selenocysteine formation [28].

Methanomicrobiales (Class II): Of the 62 exclusive signature clusters for Methanomicrobiales, 26 are hypothetical proteins, reflecting the fact that this order has been less studied. A serine/threonine kinase and a serine phosphatase, both of which regulate sigma factors in bacteria (see the Sigma Factor Regulators section above), are part of the Methanomicrobiales signature. In addition to a full-length heterodisulfide reductase subunit A (HdrA), Methanomicrobiales also contain a homolog that is

truncated at both the N- and C-terminus. Similarly, the A and G subunits of their tetrahydromethanopterin S-methyltransferase are fused. A separate A subunit was found, but no other G subunit is present.

Methanosarcinales (Class III): Among the exclusive signature proteins found in Methanosarcinales are subunits A, K, and N of reduced coenzyme F₄₂₀ (F₄₂₀H₂) dehydrogenase. Since only Methanosarcinales can use methyl compounds as a substrate for methanogenesis, it is not surprising that this enzyme, used for growth on methyl compounds, is not found in the other methanogens. All methanogens have the archaeal bisphosphoglycerate-independent phosphoglycerate mutase, but Methanosarcinales also have a bacterial version. Similarly, Methanosarcinales use the bacterial adenylate kinase while other methanogens have the archaeal enzyme.

Methanosarcinales exclusive signature proteins include phage shock protein A, a protein that functions in the repair of damaged cell membranes. Likewise, they encode two proteins involved in DNA compaction: the non-histone chromosomal protein MC1 and a unique variant (~200 amino acids longer) of the ScpB subunit of the condensin complex. That these two chromosome condensation proteins are found only in Methanosarcinales may be related to the larger genome size of some members. The other two components of the condensin complex, ScpA and Smc, are present in most methanogens, including Methanosarcinales. In addition to these DNA condensation proteins, all methanogens have at least one histone gene. Most also have a copy of the gene encoding the Alba protein, but among Methanosarcinales it is present only in *Methanosaeta thermophila*.

Missing from Class I: There are 33 clusters shared by Methanomicrobiales and Methanosarcinales that are absent from Class I methanogens. Among these are the DNA mismatch repair proteins MutL and MutS. MutH, however, is not present in any methanogen. This suggests that, if Class I methanogens have methyl-directed mismatch repair, they use a different system. Class I methanogens also lack DNA gyrase subunits A and B. This is unexpected as several Class I methanogens were found to be sensitive to coumarins that target bacterial DNA gyrase [29]. Furthermore, DNA gyrase is the only protein known to introduce negative supercoils into DNA, and these are required for many cellular processes including transcription and DNA replication [30]. Another enzyme missing from Class I methanogens is 5-amino-4-imidazolecarboxamide ribonucleotide (AICAR) transformylase in the pathway for *de novo* purine synthesis. Since most Class I methanogens are autotrophs, they must have this capability provided by a protein unrelated to the known enzyme.

Another shared cluster is the one containing Ech hydrogenase subunit A. Although *M. acetivorans* lacks Ech, it does have the F₄₂₀H₂ dehydrogenase subunit L and a subunit of multisubunit sodium/proton antiporters, both of which cluster with EchA.

Lastly, Methanomicrobiales and Methanosarcinales have one form of glucose-6-phosphate isomerase (COG2140), while most Class I methanogens use another (COG0166). Since a glucose-6-phosphate isomerase could not be identified in *Methanopyrus kandleri* or in Methanobacteriales, there is probably a third form of this enzyme.

Discussion

Phylogenetics: The sequencing of the genomes of *M. labreanum* and *M. marisnigri* reported in this paper, combined with the previously sequenced genome of *M. hungatei*, has enabled further characterization of the order Methanomicrobiales and clarification of its relationship to other methanogens. Our analyses including these species reveal that the order Methanomicrobiales is clearly distinct from other methanogens. The phylogenetic tree built for seven core methanogenesis and cofactor biosynthesis enzymes reveals three discrete groups of methanogens: the Class I methanogens, the Methanomicrobiales (termed here Class II), and the Methanosarcinales (termed here Class III). This classification differs significantly from the previous study by Baptiste et al. [7] that divided the methanogens into two major groups. In that earlier study, the order Methanosarcinales was represented by only species from the genus *Methanosarcina*, whereas our study also included two genomes from other genera. Likewise, their analysis included only one representative of the Methanomicrobiales, while we included four species from this order. Because our study encompassed more species and greater diversity, our results may be a more accurate representation of the relationships among these groups. A relatively close relationship was previously seen between Methanosarcinales and Methanomicrobiales in 16S rRNA trees [3,31] and ribosomal protein trees [3,7]. In contrast, Methanomicrobiales are equally distant from Class I methanogens and Methanosarcinales in the tree built in this study from core methanogenesis proteins.

Genomic Analyses: The protein clustering results reported also suggest a significant distance between Methanomicrobiales and all other methanogens. They share only 6 signature clusters with Class I methanogens and 33 with Methanosarcinales. In addition,

the number of exclusive signature clusters for the Methanomicrobiales is of the same magnitude as the signatures for the other two groups. The complement of membrane-bound hydrogenases also shows the uniqueness of Methanomicrobiales. They all have the Eha hydrogenase similar to Class I methanogens and the Ech hydrogenase found in Methanosarcinales, while some of them have hydrogenases similar to Mbh from *P. furiosus* and a putative membrane-bound hydrogenase from Methanosarcinales.

Methanomicrobiales share some capabilities with Class I methanogens to the exclusion of Methanosarcinales. Both groups are capable of using only H₂/CO₂ or formate for methanogenesis. The genomes show that they also share the pathway for 2-oxoglutarate synthesis. Both use a partial reductive TCA cycle, while Methanosarcinales use a partial oxidative TCA cycle. This could reflect the observations that Methanomicrobiales efficiently use low concentrations of H₂, while the Methanosarcinales dominate in environments in which acetate is plentiful. The partial oxidative TCA cycle results in the loss of one carbon as CO₂, therefore the use of the reductive cycle by Methanomicrobiales and Class I methanogens would be predicted to preserve more fixed carbon. On the other hand, we propose that, similar to Methanosarcinales, Methanomicrobiales link formylmethanofuran synthesis and heterodisulfide reduction to membrane ion gradients, even though they lack cytochromes and methanophenazine that are present in Methanosarcinales.

Hydrogenases: Methanomicrobiales encode from two to four membrane-bound hydrogenases. In all three genomes (*M. labreanum*, *M. marisnigri*, and *M. hungatei*), the genes for Eha hydrogenase are found adjacent to genes for formylmethanofuran dehydrogenase (Fmd), suggesting that the Eha hydrogenase may reduce a low potential

ferredoxin that is required for the reduction of CO₂ to formylmethanofuran. In contrast, in the Class I methanogen *Methanococcus maripaludis*, the *eha* and *fmd* operons are not linked, and Eha hydrogenase presumably plays a role in carbon assimilation similar to Ehb and not methanogenesis [8,32,33].

All Methanomicrobiales also contain genes for the Ech hydrogenase that has been characterized in *M. barkeri*. Ech hydrogenase is involved in reduction of ferredoxin for the first step of methanogenesis from H₂/CO₂, in the reduction of ferredoxin for biosynthesis, and in the formation of H₂ from ferredoxin during acetoclastic methanogenesis [20]. Since the Ech hydrogenase is found in all three Methanomicrobiales, it is likely that its function is common to all three, e.g. the reduction of ferredoxin for 2-oxoglutarate synthesis. Another putative membrane-bound hydrogenase (Pmh) is found only in *M. hungatei* where it may perform a function that is unique to this organism, such as producing ferredoxin for acetyl-CoA decarbonylase/synthase, an enzyme that is absent from the other two.

Experimental evidence is needed to determine the functions of these hydrogenases. Nevertheless, their distribution within the Methanomicrobiales is clearly distinct from that in the Class I methanogens and the Methanosarcinales (Table 2), supporting the functional and evolutionary uniqueness of this group. Their distribution and other features of the operons suggest that their roles in energy conservation differ in Class I methanogens, Methanosarcinales, and Methanomicrobiales.

Materials and Methods

DNA Preparation. *M. marisnigri* strain JR1 was obtained from the ATCC (ATCC 35101). It was cultured at room temperature in modified McC medium [34] that

contained 0.1 M NaCl, 3 g/L of sodium bicarbonate, 2 g/L of Trypticase (replacing yeast extract), and 0.17 g/L of Na₂S₉H₂O. *M. labreanum* strain Z was obtained from the ATCC (ATCC 43576). It was cultured at 37°C in MS-OCM Base Medium with 2.5 g/L NaCl, 5 mM sodium acetate, 50 mM sodium formate, and 2.5% (v/v) of rumen fluid.

For DNA isolation, cells were suspended in TE buffer (10 mM Tris, 1 mM EDTA, pH 8.0). Sodium dodecyl sulfate was added to a final concentration of 0.5% and proteinase K was added to make 100 micrograms/ml, then the solution was incubated at 37°C for 1 hour. After adding NaCl to 0.5 M concentration, the solution was approximately 0.9 ml. Next, 0.5 ml chloroform:isopropyl alcohol (24:1) was added. The solution was mixed and then centrifuged at 13,000 x g for 10 minutes. The aqueous phase was transferred to a new tube, combined with 0.5 ml phenol:chloroform:isoamyl alcohol (25:24:1), mixed, and centrifuged at 13,000 x g for 10 minutes. The aqueous phase was collected, combined with 0.6 ml isopropanol, incubated at room temperature for 30 minutes, and then centrifuged at 13,000 x g for 5 minutes. The pellet was washed with 70% ethanol, resuspended in TE + RNase (100 micrograms/ml), and incubated at 37°C for 20 minutes.

Genome Sequencing and Assembly. The genome of *M. labreanum* Z was sequenced at the Joint Genome Institute (JGI) using a combination of Sanger shotgun sequencing and 454 sequencing-by-synthesis technology. All general aspects of library construction and sequencing performed at the JGI can be found at http://www.jgi.doe.gov/sequencing/protocols/prots_production.html. Draft assemblies were based on 26,432 Sanger shotgun and 390,106 pyrosequencing reads. The combined reads provided 34x coverage of the genome. The Newbler assembly software (www.454.com) and the Paracel Genome Assembler (Paracel, Pasadena, CA) were used

for fragment assembly, and the Consed finishing package (www.phrap.org) was used for quality assessment and editing. All mis-assemblies were corrected and all gaps between contigs were closed by custom primer walk using subclones or PCR products as templates. A total of 196 additional reactions were run to close gaps and to raise the quality of the finished sequence.

The genome of *M. marisnigri* JR1 was sequenced at the Joint Genome Institute (JGI) using a combination of 3 kb, 7 kb and 36 kb (fosmid) DNA libraries. Draft assemblies were based on 29,769 total reads. The three libraries combined provided 11x coverage of the genome. The Phred/Phrap/Consed software package (www.phrap.com) was used for sequence assembly and quality assessment [35-37]. All mis-assemblies were corrected and all gaps between contigs were closed by custom primer walk using subclones or PCR products as templates. A total of 702 primer walk reactions, PCR end reads and 3 mini-libraries were required to close gaps and to raise the quality of the finished sequence.

Genome Analysis. Automatic genome annotation was performed at Oak Ridge National Laboratory. Genes were identified using a combination of Critica [38] and Glimmer [39]. In addition, predicted coding regions (CDSs) were manually curated using JGI's Gene-PRIMP Quality Assurance pipeline (<http://tunis.jgi-psf.org/geneprimp>) (Pati et al., in preparation). Comparative genome analysis was performed within the Integrated Microbial Genomes (IMG) system [40]. CRISPR repeats were identified with the CRISPR Recognition Tool [41].

A phylogenetic tree was constructed using the concatenated sequences of seven core proteins found in all methanogens and involved in methanogenesis and cofactor biosynthesis. The genes included are F₄₂₀-dependent methylenetetrahydromethanopterin

dehydrogenase (*mtd*, COG1927), tetrahydromethanopterin:coenzyme M methyltransferase subunits B (*mtrB*, COG4062), C (*mtrC*, COG4061), D (*mtrD*, COG4060) and E (*mtrE*, COG4059), F_O synthase subunit 1 (*cofG*), and sulfopyruvate decarboxylase alpha subunit (*comD*). Protein sequences were downloaded from IMG [40]. The concatenated amino acid sequences were aligned with Clustal W [42], and the tree was generated with MrBayes 3.1.2 [43] with 1,000,000 generations sampled every 100 generations. The first 250,000 generations were discarded as burn-in. The tree was visualized with TreeView [44].

For protein clustering, methanogens were included from all three groups: six Class I methanogens, four Methanomicrobiales, and five Methanosarcinales. Class I included *Methanocaldococcus jannaschii*, *Methanothermobacter thermautotrophicus*, *Methanopyrus kandleri*, *Methanococcus maripaludis* S2, *Methanobrevibacter smithii*, and *Methanosphaera stadtmanae*. Methanomicrobiales included *Methanocorpusculum labreanum*, *Methanoculleus marisnigri*, *Methanospirillum hungatei*, and Candidatus *Methanoregula boonei*. Methanosarcinales included *Methanosarcina acetivorans*, *Methanosarcina mazei*, *Methanosarcina barkeri*, *Methanococcoides burtonii*, and *Methanosaeta thermophila*. Protein sequences for these organisms were downloaded from IMG [40]. We applied a spectral clustering procedure [45,46] to cluster similar proteins based on the topology of their similarity graph, rather than using a fixed threshold value for sequence similarity. The proteins are represented as nodes in a connected undirected graph with edges that carry weights based on node-to-node similarity according to the protein identity. The clustering procedure is analogous to a random walk of a particle moving on this graph from one node to another. In each node

the particle moves to another node based on the probabilities corresponding to the weights of the edges. The amount of time the particle spends in a given subgraph will determine whether this is indeed a cluster of its own or not.

The second eigenvalue of the transition matrix is a measure of how easily a graph (i.e. a cluster) can be partitioned. A cutoff value of 0.8 was applied; if the second eigenvalue exceeds 0.8, the cluster is further partitioned. This approach provides a relatively flexible partitioning that can reveal protein similarities despite sequence differences due to phylogenetic distance.

Signature protein clusters were identified as clusters for which a member protein was present in every analyzed species from one (or more) class of methanogens. Those clusters were binned into groups: exclusive signature clusters found in all members of only one class, shared signature clusters found in all members of a specified pair of classes, and common clusters found in all three classes. The resultant cluster distribution was visualized as a Venn diagram.

Accession Numbers. The genome sequences of *Methanoculleus marisnigri* JR1, *Methanocorpusculum labreanum* Z, and *Methanospirillum hungatei* JF-1 can be accessed in GenBank (CP000562, CP000559, and CP000254, respectively). The Genomes OnLine Database accession numbers are Gc00512, Gc00506, and Gc00350, respectively.

Acknowledgments

This work was performed under the auspices of the US Department of Energy's Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Berkeley National Laboratory under Contract No. DE-AC02-05CH11231, Lawrence Livermore National Laboratory under Contract No. DE-AC52-07NA27344, and Los Alamos National Laboratory under Contract No. DE-AC02-06NA25396. L. E. U. was supported by grant number GM72285 from the National Institutes of Health. B. L., I. P., M. S.-L., and W. B. W. were supported by DOE contract number DE-FG02-97ER20269. D. S., L. D., and B. M. were supported by NASA Astrobiology, Exobiology, and Evolutionary Biology grant NNG05GP24G. M. L. was supported by the Department of Energy under contract DE-AC05-000R22725.

References

1. Woese CR, Fox GE (1977) Phylogenetic structure of the prokaryotic domain: the primary kingdoms. *Proc Natl Acad Sci U S A* 74: 5088-5090.
2. Woese CR, Kandler O, Wheelis ML (1990) Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc Natl Acad Sci U S A* 87: 4576-4579.
3. Brochier-Armanet C, Boussau B, Gribaldo S, Forterre P (2008) Mesophilic crenarchaeota: proposal for a third archaeal phylum, the Thaumarchaeota. *Nat Rev Microbiol* 6: 245-252.
4. Ferry JG (1997) Methane: small molecule, big impact. *Science* 278: 1413-1414.
5. Etheridge DM, Steele LP, Francey RJ, Langenfelds RL (1998) Atmospheric methane between 1000 A.D. and present: evidence of anthropogenic emissions and climatic variability. *J Geophys Res* 103: 15979-15993.
6. Fox GE, Magrum LJ, Balch WE, Wolfe RS, Woese CR (1977) Classification of methanogenic bacteria by 16S ribosomal RNA characterization. *Proc Natl Acad Sci U S A* 74: 4537-4541.
7. Bapteste É, Brochier C, Boucher Y (2005) Higher-level classification of the Archaea: evolution of methanogenesis and methanogens. *Archaea* 1: 353-363.
8. Thauer RK, Kaster A-K, Seedorf H, Buckel W, Hedderich R (2008) Methanogenic archaea: ecologically relevant differences in energy conservation. *Nat Rev Microbiol* 6: 579-591.
9. Garcia J-L, Ollivier B, Whitman WB (2006) The order Methanomicrobiales. *Prokaryotes* 3: 208-230.
10. Zellner G, Winter J (1987) Secondary alcohols as hydrogen donors for CO₂-reduction by methanogens. *FEMS Microbiol Lett* 44: 323-328.
11. Romesser JA, Wolfe RS, Mayer F, Spiess E, Walther-Mauruschat A (1979) *Methanogenium*, a new genus of marine methanogenic bacteria, and characterization of *Methanogenium cariaci* sp. nov. and *Methanogenium marisnigri* sp. nov. *Arch Microbiol* 121: 147-153.
12. Zhao Y, Boone DR, Mah RA, Boone JE, Xun L (1989) Isolation and characterization of *Methanocorpusculum labreanum* sp. nov. from the LaBrea tar pits. *Int J Syst Bacteriol* 39: 10-13.

13. Ferry JG, Smith PH, Wolfe RS (1974) *Methanospirillum*, a new genus of methanogenic bacteria, and characterization of *Methanospirillum hungatii* sp. nov. Int. J. Syst. Bacteriol. 24: 465-469.
14. Widdel F, Rouvière PE, Wolfe RS (1988) Classification of secondary alcohol-utilizing methanogens including a new thermophilic isolate. Arch Microbiol 150: 477-481.
15. Bleicher K, Winter J (1991) Purification and properties of F₄₂₀- and NADP⁺-dependent alcohol dehydrogenases of *Methanogenium liminatans* and *Methanobacterium palustre*, specific for secondary alcohols. Eur J Biochem 200: 43-51.
16. Aufhammer SW, Warkentin E, Berk H, Shima S, Thauer RK *et al.* (2004) Coenzyme binding in F₄₂₀-dependent secondary alcohol dehydrogenase, a member of the bacterial luciferase family. Structure 12: 361-370.
17. Stojanowic A, Mander GJ, Duin EC, Hedderich R (2003) Physiological role of the F₄₂₀-non-reducing hydrogenase (Mvh) from *Methanothermobacter marburgensis*. Arch Microbiol 180: 194-203.
18. Shima S, Thauer RK (2007) A third type of hydrogenase catalyzing H₂ activation. Chem Rec 7: 37-46.
19. Hedderich R, Forzi L (2005) Energy-converting [NiFe] hydrogenases: more than just H₂ activation. J Mol Microbiol Biotechnol 10: 92-104.
20. Meuer J, Kuettner HC, Zhang JK, Hedderich R, Metcalf WW (2002) Genetic analysis of the archaeon *Methanosarcina barkeri* Fusaro reveals a central role for Ech hydrogenase and ferredoxin in methanogenesis and carbon fixation. Proc Natl Acad Sci U S A 99: 5632-5637.
21. Verhees CH, Tuininga JE, Kengen SWM, Stams AJM, van der Oost J *et al.* (2001) ADP-dependent phosphofructokinases in mesophilic and thermophilic methanogenic archaea. J Bacteriol 183: 7145-7153.
22. Makarova KS, Koonin EV (2003) Filling a gap in the central metabolism of archaea: prediction of a novel aconitase by comparative-genomic analysis. FEMS Microbiol Lett 227: 17-23.
23. Kyrpides NC, Ouzounis CA (1997) Bacterial sigma 70 transcription factor DNA-binding domains in the archaeon *Methanococcus jannaschii*. J Mol Evol 45: 706-707.
24. Decatur AL, Losick R (1996) Three sites of contact between the *Bacillus subtilis* transcription factor sigma F and its antisigma factor SpoIIAB. Genes Dev 10: 2348-2358.

25. Zhang Y, Turanov AA, Hatfield DL, Gladyshev VN (2008) *In silico* identification of genes involved in selenium metabolism: evidence for a third selenium utilization trait. *BMC Genomics* 9: 251.
26. Yuan J, Palioura S, Salazar JC, Su D, O'Donoghue P *et al.* (2006) RNA-dependent conversion of phosphoserine forms selenocysteine in eukaryotes and archaea. *Proc Natl Acad Sci U S A* 103: 18923-18927.
27. Xu X-M, Carlson BA, Mix H, Zhang Y, Saira K *et al.* (2007) Biosynthesis of selenocysteine on its tRNA in eukaryotes. *PLoS Biol* 5: e4.
28. Kaiser JT, Gromadski K, Rother M, Engelhardt H, Rodnina MV *et al.* (2005) Structural and functional investigation of a putative archaeal selenocysteine synthase. *Biochemistry* 44: 13315-13327.
29. Sioud M, Possot O, Elie C, Sibold L, Forterre P (1989) Coumarin and quinolone action in archaebacteria: evidence for the presence of a DNA gyrase-like enzyme. *J Bacteriol* 170: 946-953.
30. Nöllman M, Crisona NJ, Arimondo PB (2007) Thirty years of *Escherichia coli* DNA gyrase: From *in vivo* function to single-molecule mechanism. *Biochimie* 89: 490-499.
31. Schleper C, Jurgens G, Jonuscheit M (2005) Genomic studies of uncultivated archaea. *Nat Rev Microbiol* 3: 479-488.
32. Porat I, Kim W, Hendrickson EL, Xia Q, Zhang Y *et al.* (2006) Disruption of the operon encoding Ehb hydrogenase limits anabolic CO₂ assimilation in the archaeon *Methanococcus maripaludis*. *J Bacteriol* 188: 1373-1380.
33. Lupa B, Hendrickson EL, Leigh JA, Whitman WB (2008) Formate-dependent H₂ production by the mesophilic methanogen *Methanococcus maripaludis*. *Appl Env Microbiol* 74: 6584-6590.
34. Whitman WB, Shieh J-S, Sohn S-H, Caras DS, Premachandran U (1986) Isolation and characterization of 22 mesophilic methanococci. *Syst Appl Microbiol* 7: 235-240.
35. Ewing B, Green P (1998) Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res.* 8: 186-194.
36. Ewing B, Hillier L, Wendl MC, Green P (1998) Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res.* 8: 175-185
37. Gordon D, Abajian C, Green P (1998) Consed: a graphical tool for sequence finishing. *Genome Res.* 8: 195-202.

38. Badger JH, Olsen GJ (1999) CRITICA: coding region identification tool invoking comparative analysis. *Mol Biol Evol* 16: 512-524.
39. Delcher AL, Harmon D, Kasif S, White O, Salzberg SL (1999) Improved microbial gene identification with GLIMMER. *Nucleic Acids Res* 27: 4636-4641.
40. Markowitz VM, Szeto E, Palaniappan K, Grechkin Y, Chu K *et al.* (2008) The integrated microbial genomes (IMG) system in 2007: data content and analysis tool extensions. *Nucleic Acids Res* 36: D528-D533.
41. Bland C, Ramsey TL, Sabree F, Lowe M, Brown K *et al.* (2007) CRISPR Recognition Tool (CRT): a tool for automatic detection of clustered regularly interspaced palindromic repeats. *BMC Bioinformatics* 8: 209.
42. Thompson JD, Higgins DG, Gibson TJ (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 22: 4673-4680.
43. Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572-1574.
44. Page RDM (1996) TREEVIEW: an application to display phylogenetic trees on personal computers. *Comput Appl Biosci* 12: 357-358.
45. Paccanaro A, Casbon JA, Saqi MAS (2006) Spectral clustering of protein sequences. *Nucleic Acids Res* 34: 1571-1580.
46. Brewer ML (2007) Development of a spectral clustering method for the analysis of molecular data sets. *J Chem Inf Model* 47: 1727-1733.

Figure legends

Figure 1. Proposed pathway for methanogenesis in Methanomicrobiales.

Methanomicrobiales are predicted to couple formylmethanofuran formation and CoM-CoB heterodisulfide reduction to ion gradients. Fd: ferredoxin; MF: methanofuran; H₄MPT: tetrahydromethanopterin.

Figure 2. Alternate pathways for synthesis of 2-oxoglutarate from oxaloacetate. Class I methanogens and Methanomicrobiales use a partial reductive citric acid cycle while Methanosarcinales use a partial oxidative citric acid cycle.

Figure 3. Phylogenetic tree of methanogens based on seven core enzymes of methanogenesis and cofactor biosynthesis. See Materials and Methods for a list of the proteins and organisms included. Protein sequences were concatenated and aligned with Clustal W. The tree was generated with MrBayes 3.1.2 and viewed with TreeView.

Figure 4. Venn diagram of signature clusters. The clusters were generated using a spectral clustering procedure (see Materials and Methods section for details). Signature protein clusters were identified as clusters for which a member protein was present in every analyzed species from one or more classes of methanogens. The number of exclusive, shared, and common signature clusters associated with each methanogen class are shown. The functions of characterized proteins belonging to exclusive signature clusters and to clusters shared between the Methanomicrobiales and the Methanosarcinales are also noted.

Table 1. General genome statistics

	<i>M. labreanum</i>	<i>M. marisnigri</i>	<i>M. hungatei</i>
Genome size (bp)	1,804,962	2,478,101	3,544,738
G+C content (bp)	902,600 (50.0%)	1,537,981 (62.1%)	1,600,415 (45.1%)
Number of genes	1828	2559	3305
RNA genes	63 (3.4%)	53 (2.1%)	66 (2.0%)
Protein-coding genes	1765 (96.6%)	2506 (97.9%)	3239 (98.0%)
Pseudogenes	26 (1.4%)	17 (0.7%)	99 (3.0%)
Genes in ortholog clusters	1676 (91.7%)	2294 (89.6%)	3031 (91.7%)
Genes assigned to COGs	1358 (74.3%)	1832 (71.6%)	2314 (70.0%)
Genes assigned to Pfam domains	1335 (73.0%)	1790 (69.9%)	2326 (70.4%)
Genes with signal peptides	406 (22.2%)	620 (24.2%)	771 (23.3%)
Genes with transmembrane helices	368 (20.1%)	595 (23.3%)	762 (23.1%)
Fusion genes	73 (4.0%)	104 (4.1%)	171 (5.2%)
Transposable elements	0	3	76
CRISPR-associated genes	8	1	21
CRISPR repeat arrays	1	0	6

Table 2. Hydrogenases in methanogen genomes

	Frh	Mvh	Eha	Ehb	Ech	Mbh
Class I methanogens	All	All	all except Msp	all except Mka		
Methanosarcinales	Mac, Mba, Mmz				Mba, Mmz	
Methanomicrobiales	All	Mmar	All		All	Mlab, Mhun

Frh: F₄₂₀-reducing hydrogenase; Mvh: F₄₂₀-non-reducing hydrogenase; Eha: energy-converting hydrogenase A; Ehb: energy-converting hydrogenase B; Ech: energy-converting hydrogenase; Mbh: membrane-bound hydrogenase; Msp: *Methanosphaera stadtmanae*; Mka: *Methanopyrus kandleri*; Mac: *Methanosarcina acetivorans*; Mba: *Methanosarcina barkeri*; Mmz: *Methanosarcina mazei*; Mmar: *Methanoculleus marisnigri*; Mlab: *Methanocorpusculum labreanum*; Mhun: *Methanospirillum hungatei*.

Figure 1




Figure 2.




Figure 3.




Figure 4.

