

UC Davis

UC Davis Previously Published Works

Title

Genomic Characterization of the Evolutionary Potential of the Sea Urchin *Strongylocentrotus droebachiensis* Facing Ocean Acidification.

Permalink

<https://escholarship.org/uc/item/02d5f9g3>

Journal

Genome biology and evolution, 8(12)

ISSN

1759-6653

Authors

Runcie, Daniel E
Dorey, Narimane
Garfield, David A
et al.

Publication Date

2016-12-01

DOI

10.1093/gbe/evw272

Peer reviewed

Genomic Characterization of the Evolutionary Potential of the Sea Urchin *Strongylocentrotus droebachiensis* Facing Ocean Acidification

Daniel E. Runcie^{1,2,*}, Narimane Dorey³, David A. Garfield^{1,4}, Meike Stumpff^{3,5}, Sam Dupont³, and Gregory A. Wray^{1,6}

¹Department of Biology, Duke University, Durham, NC

²Department of Plant Sciences, University of California, Davis

³Department of Biological and Environmental Sciences, University of Gothenburg, Fiskebäckskil, Sweden

⁴Integrative Research Institute for the Life Sciences, Humboldt University, Berlin, Germany

⁵Helmholtz Centre for Ocean Sciences (GEOMAR), Kiel, Germany

⁶Center for Genomic and Computational Biology, Duke University, Durham, NC

*Corresponding author: E-mail: deruncie@ucdavis.edu.

Accepted: November 10, 2016

Data deposition: All RNAseq reads generated for the manuscript have been deposited at the NCBI Sequence Read Archive under BioProject: PRJNA301890 and biosamples: SAMN04260159 - SAMN04260200. All computational scripts will be posted in Dryad after the manuscript is accepted. All computational scripts and processed data have been deposited at <http://dx.doi.org/10.5061/dryad.1f6t8> with DOI: doi:10.5061/dryad.1f6t8

Abstract

Ocean acidification (OA) is increasing due to anthropogenic CO₂ emissions and poses a threat to marine species and communities worldwide. To better project the effects of acidification on organisms' health and persistence, an understanding is needed of the 1) mechanisms underlying developmental and physiological tolerance and 2) potential populations have for rapid evolutionary adaptation. This is especially challenging in nonmodel species where targeted assays of metabolism and stress physiology may not be available or economical for large-scale assessments of genetic constraints. We used mRNA sequencing and a quantitative genetics breeding design to study mechanisms underlying genetic variability and tolerance to decreased seawater pH (-0.4 pH units) in larvae of the sea urchin *Strongylocentrotus droebachiensis*. We used a gene ontology-based approach to integrate expression profiles into indirect measures of cellular and biochemical traits underlying variation in larval performance (i.e., growth rates). Molecular responses to OA were complex, involving changes to several functions such as growth rates, cell division, metabolism, and immune activities. Surprisingly, the magnitude of pH effects on molecular traits tended to be small relative to variation attributable to segregating functional genetic variation in this species. We discuss how the application of transcriptomics and quantitative genetics approaches across diverse species can enrich our understanding of the biological impacts of climate change.

Key words: System genetics, climate change, genetic variation, plasticity, RNAseq, gene set variation analysis.

Introduction

The rapid pace of climate change threatens the persistence of species worldwide by exposing individuals to environmental conditions outside the range of recent historical precedents (IPCC 2014). In the marine realm, the combination of elevated temperatures and decreased pH (i.e., ocean acidification [OA]) poses a particularly acute challenge (Caldeira and Wickett

2003; Doney et al. 2009). Phenotypic responses to environmental factors may be a key determinant of a species' prospects for survival, by buffering critical life processes via altered behavior, phenology, or stress response mechanisms (Williams et al. 2008). Evolutionary adaptation provides another path for species persistence and may be particularly advantageous over the long term (Davis and Shaw 2001; Davis et al. 2005). The prospects for evolutionary rescue depend on many factors,

© The Author(s) 2016. Published by Oxford University Press on behalf of the Society for Molecular Biology and Evolution.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

including demography, life history, and the amount of standing genetic variation in traits directly related to the tolerance of the environmental challenge (Lynch and Lande 1993; Bürger and Lynch 1995; Gomulkiewicz and Holt 1995). Even if a population harbors extensive genetic variation that influences traits necessary for adaptation, genetic or developmental trade-offs can place limitations on its ability to adapt (Lande and Arnold 1983). Although many studies have documented genetic variation that influences the tolerance of stressors related to climate change, evidence for recent adaptation—particularly adaptation that can keep up with the rapid pace of climate change—remains limited (Gienapp et al. 2008; Hendry et al. 2008; Kelly et al. 2013; Vander Wal et al. 2013).

Most evolutionary and ecological studies focus on accessible traits such as growth, survival, or phenology (e.g., Parmesan and Yohe 2003; Hendry et al. 2008; Gienapp et al. 2008), whereas variation in physiological, metabolic, and developmental traits is less commonly considered even though they are also key determinants of environmental responses (Wikelski and Cooke 2006; Chown and Gaston 2008; Somero 2010; Pan et al. 2015). Importantly, the link between these internal traits and external metrics of performance is often not obvious (Houle 2010). A comprehensive, unbiased approach is therefore needed to understand the genetic basis for plasticity and evolutionary responses to climate change.

A central challenge facing such efforts is that key physiological and developmental traits are often difficult, time consuming, or expensive to measure, making it impractical to survey a population for genetic variation affecting these traits. Genomic approaches, particularly ones that combine high-throughput molecular phenotyping with measures of high-level performance traits, provide a partial but highly practical solution. Transcriptome-wide gene expression assays are increasingly affordable and produce a rich source of data, both on variation in gene expression and on variation in cellular, physiological, and developmental processes that leave signatures in the covariation of functionally related genes (Alvarez et al. 2015). To list just a few examples, recent studies using transcriptomics in evolutionary ecology have identified gene networks underlying adaptive developmental plasticity in cichlid fish (Schneider et al. 2014), physiological mechanisms of cold and hypoxia tolerance in deer mice (Cheviron et al. 2014), and hormone signals regulating drought responses in *Arabidopsis* (Marais Des et al. 2012). Analytical approaches that leverage databases of gene function or genetic pathways (gene set enrichment analyses; Subramanian et al. 2005) are particularly promising: they reduce the high-dimensional data to a more accessible scale, highlight variation in informative molecular processes, and increase the power to detect subtle signals in noisy gene expression data. Here, we apply a transcriptome-based systems genetics approach (Civelek and Lusis 2013) to study the molecular response to changes in ocean pH

relevant in the context of OA in larvae of the green sea urchin, *Strongylocentrotus droebachiensis*.

In marine systems, OA is one of the most immediate results of climate change and is recognized as a major threat to many species and communities (IPCC 2014; Breitbart et al. 2015; Gaylord et al. 2015). CO₂ from the atmosphere dissolves in the ocean and reacts with water, lowering seawater pH and altering the carbonate balance of seawater (Doney et al. 2009). OA can affect a wide range of physiological processes in marine species, including acid–base balance, metabolism, development, deposition of skeleton, and growth rates (Dupont and Pörtner 2013). A number of studies in recent years have investigated the tolerance of sea urchins to decreased pH (reviewed in Dupont et al. 2010; Byrne 2011). In general, sea urchins have proven surprisingly robust to low pH (Dupont and Thorndyke 2013). However, large differences in tolerance are apparent even among closely related species, different life history stages can differ in their tolerance of low pH, and effects may increase when combined with other climate-related stressors such as increasing temperatures (Byrne 2011; Gianguzza et al. 2014). At the molecular level, larvae grown under low pH show increased expression of genes related to metabolic and ion regulation, and differential regulation of several genes involved in stress responses, apoptosis, calcification and skeletal formation (Todgham and Hofmann 2009; Martin et al. 2011; Stumpp et al. 2011a; Evans et al. 2013; Padilla-Gamiño et al. 2013; Pespeni et al. 2013). Several studies have also observed genetic variation in the response of growth rates (Sunday et al. 2011; Kelly et al. 2013) and development (Foo et al. 2012) to decreased pH in sea urchin embryos and larvae, as well as genetic changes in allele frequencies in artificial populations grown under simulated future conditions (Pespeni et al. 2013).

We extend these studies using transcriptome-wide gene expression profiling and a quantitative genetic breeding design to quantify molecular plasticity and test for segregating genetic variation affecting cellular and physiological traits in cultures of larvae spawned from *S. droebachiensis* individuals collected in northern Europe. We demonstrate that the molecular response to decreased pH in these larvae is broad and complex. Surprisingly, we also find that the gene expression plasticity to pH is dwarfed in magnitude and frequency by the influence of genetic variation among families. Finally, we show how combining signals over many functionally related genes can provide detailed information regarding 1) molecular mechanisms underlying responses to decreased pH and 2) aspects of these mechanisms that are mostly likely to contribute to near-term adaptations.

Materials and Methods

Experimental Design

Traits with ample genetic variation can efficiently respond to natural selection. A traditional tool to assess genetic variation




FIG. 1.—Experimental layout. We used a three-way factorial layout to assess the effects of pH (control vs low pH), male parent (seven individuals), and female parent (three individuals). Cultures were created representing all 42 combinations of these three factors. Larvae were raised for 14 days, and monitored daily for mortality, growth rate (days 1–8), and seawater chemistry (table 1).

in a population is to test for consistent differences among the offspring with different fathers (Lynch and Walsh 1998) because sperm is thought to contribute little to zygotes beyond DNA although the generality of this conclusion has been challenged by recent studies (Krawetz 2005; Curley et al. 2011; Crean et al. 2013). Consistent differences among offspring with different mothers may also be caused by genetic (heritable) variation, but may additionally be caused by maternal environmental variation that is passed transgenerationally through egg quality or epigenetic modifications. We used a three-way factorial breeding design to quantify effects of genetic background and seawater pH on larval gene expression, survival, and growth. The three factors were as follows: father (seven levels), mother (two levels), and pH (two levels; 8.0 and 7.6). A decrease by 0.4 units from the average pH at the study site (pH_T 8.0) is within the range of the projections from the IPCC (2014) for average pH in 2100. Additionally, taking into account local environmental variability, a pH of 7.6 is the extreme of present environmental variability (Dorey et al. 2013). Cultures of larvae representing all 42 combinations of these three sets of factors were created and monitored for up to 9 days.

Animal Collection and Rearing

Adult *S. droebachiensis* were collected during the fall of 2009 from northern Norway (three males and one female; Hammerfest: 70°39'N/23°39'E, collected by SCAN-AQUA) and northern Denmark (four males and two females; Anholt: 56°42'N/11°31'E, collected by the Marine Biological Laboratory of Helsingør). The original experimental design intended to test for differences between the populations in early-life responses to low pH; however, we do not focus on

these differences here. Adults were maintained in a deep-water flow-through aquarium at the Sven Lovén Centre for Marine Sciences - Kristineberg (Fiskebäckskil, Sweden) and were fed ad libitum using *Ulva* spp. for 6–7 months before starting the experiment (March 2010). Spawning was induced by injecting ~2 ml 0.5 mM KCl into the coelomic cavity. Sperm (four males from Denmark and three from Norway) was collected by pipette and kept dry until fertilization. Eggs (two females from Denmark and one from Norway) were collected in 250-ml beakers of filtered ambient seawater (FSW; 0.22 μm) and washed once. Total egg number was estimated by counting small-volume samples of the egg solution, and ~200K–350K eggs were divided among seven 250-ml beakers to be fertilized, each with sperm from a different male. Fertilization was followed by monitoring the elevation of the vitelline envelope under a stereomicroscope (fertilization in all cultures was >95%). This created 21 families of full-sib embryos (fig. 1).

Each family of embryos was split between two 5-l Erlenmeyer flasks of FSW for culturing at an initial density of 3–5 embryos/ml. All culture vessels were preequilibrated at the target pH (table 1. control: ca. $\text{pH}_T \approx 8.00$, $\approx 420 \mu\text{atm}$, and low pH: $\text{pH}_T \approx 7.6$, $\approx 1,150 \mu\text{atm}$). Culture vessels were aerated and gently mixed with a stream of bubbles of pressurized air and kept in the dark. Cultures were monitored for growth and survival until sampling.

Seawater pH, temperature, and dissolved inorganic carbon (DIC) were monitored throughout the experiment (table 1). Culture pH was maintained (± 0.04 units) using Aqua Medic pH controllers (NBS scale, AquaMedic, Germany) that controlled valves for injection of pure CO_2 into the culture flasks once pH_{NBS} exceeded the target value. pH-system settings were adjusted from pH measurements on the total scale

Table 1

Seawater Carbonate Chemistry in Cultures at Each pH Treatment

Treatment	Control (n = 41)	Low pH (n = 96)
pH _T	8.00 ± 0.03	7.62 ± 0.11
T (°C)	10.17 ± 0.20	10.34 ± 0.24
DIC (mmol/kg)	2.00 ± 0.08	2.13 ± 0.06
pCO ₂ (μatm)	419 ± 39	1145 ± 296
Ω _{ar}	1.76 ± 0.14	0.83 ± 0.25
Ω _{ca}	2.77 ± 0.22	1.30 ± 0.40

NOTE.—Seawater total scale pH (pH_T), temperature (T), and DIC were measured daily in four control bottles and eight to 20 randomly-chosen acidified bottles (10 days). These measurements were used to calculate CO₂ partial pressure (pCO₂) as well as aragonite and calcite saturation states (respectively Ω_{ar} and Ω_{ca}), assuming a salinity of 34.7, using the package *seacarb* for R. All the values are expressed as mean ± SD.

(pH_T) using TRIS (Tris/HCl) and AMP (2-aminopyridine/HCl) buffer solutions with a salinity of 32.0 (provided by Unit  d'Oc anographie Chimique, Universit  de Li ge, Belgium). DIC was measured with an automated carbon dioxide analyzer (CIBA Corning 965 UK). pCO₂ and calcium carbonate saturation states for calcite and aragonite (Ω_{ca} and Ω_{ar}) were calculated from DIC and pH_T using the R package *seacarb* (Gattuso et al. 2015).

Larval Mortality

On day 6, two 10-ml samples of each culture were aspirated and counted. This provided an estimate of the concentration of surviving larvae in each culture. Larval mortality associated with father or pH treatment was tested by comparing larval concentrations among the cultures with the same mother using a three-way analysis of variance (ANOVA) because the batch of eggs from each mother was distributed equally among all of her cultures. The ANOVA table is presented in [supplementary table S1, Supplementary Material](#) online.

Larval Growth Rate

Average growth rates were estimated for each culture over the first 7 days. Each day, 10 larvae were randomly sampled from each culture, fixed with 4% PFA, and imaged with a Leica microscope mounted with a DFC295 camera. From these images, the average body length (see measurement from Stumpp et al. 2011b) of each culture on each day was calculated. Growth rates were calculated in mm/day using linear regression of body length on age.

Gene Expression Measurements

Gene expression was measured using RNAseq on pools of young unfed embryos from each of the 42 cultures. Collection times were calibrated for each culture individually based on the per-culture growth rates and varied between 5.4 and 9.4 days when the average larval body length was approximately 0.36 mm. Sampling based on size rather than on

physical time accounts for potential effects of pH on developmental rates (P rtner et al. 2010; Stumpp et al. 2011a, 2011b). Developmental stages were consistent at the time of sampling, with most larvae in all cultures just starting to form buds for the posterodorsal pair of skeletal arms ([supplementary figure S1, Supplementary Material](#) online).

Approximately 3,000 larvae were collected by gently pouring 1-l of each culture into a 1.5-cm sieve consisting of a submerged nylon mesh and then pipetting the concentrated larvae into a clear 2-ml screw-top microcentrifuge tube. Once in the tube, the larvae were quickly mixed again and two 100-μl samples of highly concentrated larvae (50–200 larvae each) were collected and fixed in 4% PFA to estimate the number of collected larvae and the average larval size. Remaining larvae were pelleted by centrifugation, the seawater removed by aspiration, and 600 ml of RLT buffer (Qiagen, Hilden, Germany) added to stabilize the RNA. Tubes were vortexed rapidly for 15 s to lyse the cells and then stored at –80°C. The entire process from sieve to RLT took about 1 minute. RNA was extracted with the RNeasy kit (Qiagen) after treatment with DNase, and RNA quality was assessed with a total RNA analysis ng sensitivity (Eukaryote) assay on an Agilent 2100 Bioanalyzer. No samples showed signs of RNA degradation. The resulting mRNA was purified by poly-A selection.

RNAseq libraries were prepared by the Duke GCB Sequencing and Genomic Technologies Shared Resource using the Illumina TruSeq v1 kit with poly-A selection to purify mRNA. Six samples were pooled per lane to be run on the Illumina HiSeq 2000 system with version 3 chemistry. 1.3×10^9 50-bp single-end reads were generated, representing $1.8\text{--}5.3 \times 10^7$ reads per sample (median 3.2×10^7). Because no genome sequence of *S. droebachiensis* is available, we mapped reads to the genome sequence of the closely related species, *Strongylocentrotus purpuratus* (v3.1; Sodergren et al. 2006). We used the program bowtie2 (v2.0.0-beta5; Langmead et al. 2009; Langmead and Salzberg 2012) to map reads with the “-local” option and settings “-D 20 -R 3 -N 1 -L 20 -i S,1,0.50.” Across individuals, 74–86% of reads mapped to the *S. purpuratus* genome, and 67% of these mapped uniquely. Only uniquely mapped reads were used to estimate gene expression.

To convert mapped reads into gene expression measurements, we used the program HTseq-count (Anders et al. 2015) to count the number of reads that aligned uniquely to any exon, or either the 5' or 3' untranslated region of any isoform of the 29,016 v3.1 GLEAN gene models (GLEAN-UTR-3.1.gff3, <http://www.echinobase.org/Echinobase/SpDownloads>; last accessed July 26, 2012). Gene expression analyses were performed in R (R Development Core Team 2010). Sample normalization factors were calculated with the *calcnormfactors* function of the *edgeR* package (Robinson and Oshlack 2010; Robinson et al. 2010) using default parameters. Genes with fewer than 10 counts over the 42 samples were not analyzed further, leaving a total of 22,430 genes.

Gene Set Variation Analysis

We used gene set variation analysis (GSVA; Hänzelmann et al. 2013) implemented in the *R* package *GSVA* to explore variation in higher-order molecular traits. GSVA defines a set of synthetic traits based on a list of gene ontology terms. Each trait's value represents the extent to which genes labeled with a specific term tend to be up- or downregulated in that sample, measured using a Kolmogorov–Smirnov-like random walk statistic. Coordinated regulation of genes in the same pathway or involved in similar processes is evidence for changes in the activity of specific cellular functions (Subramanian et al. 2005).

We used two sources for gene function assignments: the PANTHER database (Mi et al. 2013) and a sea urchin-specific database of 130 ontology terms annotated by the sea urchin research community ("hand-annotated ontology": Sodergren et al. 2006; Tu et al. 2012). The PANTHER ontology is a reduced set terms from the Gene Ontology Project. For the sea urchin, the PANTHER ontology (PTHR8.1_sea_urchin) consists of 128 molecular function (MF) terms, 165 biological process (BP) terms, 22 cellular component (CC) terms, and 171 protein class (PC) terms with at least 10 and not >1,000 genes. We report results for all classes of ontology terms in the [Supplementary Material](#) online, but only discuss results from the MF and hand-annotated ontologies because we find them to be the most straight-forward to interpret. The hand-annotated ontologies are less comprehensive than the PANTHER ontologies, but include annotations for processes that have been especially studied in sea urchins, including biomineralization-related genes. Phenotypic correlations among gene set traits within and among ontologies can help guide interpretation and are presented as heatmaps in [supplementary figure S1](#), [Supplementary Material](#) online.

We used the bubble plot method described by Supek et al. (2011) to visualize relationships among gene set traits. Within each of the PANTHER ontologies, specific terms are associated in a hierarchical tree, with more general terms (containing more genes) toward the base of the tree and more specific terms (containing fewer genes) toward the tips. We calculated the *SimREL* Semantic Similarity (Schlicker et al. 2006) distance between terms in the ontology based on the tree distance and term sizes and represent this distance on the 2D plane of the bubble plot using multidimensional scaling (*R* function *sammon*).

Statistical Analyses

All statistical analyses were performed in *R* v3.2.2 (R Development Core Team 2010).

Our experimental design consisted of three sets of factors: father, mother, and pH treatment [fig. 1](#). The goal of the statistical analysis was to identify traits affected by each of these factors. For each trait, we considered a factor to be important if the main effect or any of the interactions of that factor with

other factors were significant. Therefore, our lists of pH-responsive traits include all traits affected by pH in the offspring of any father or any mother. Traits are considered to be genetically variable if father effects were detected in either pH treatment or offspring of any of the three mothers.

To quantify the importance of each factor, we used the *R* Bioconductor package *limma* (Ritchie et al. 2015) to fit the following Gaussian linear models to the variation in each trait:

$$Y_{ijk} = \mu + \text{pH}_i + \text{PopF}_j + \text{PopM}_k + \text{pH} \times \text{PopF}_j + \text{pH} \times \text{PopM}_k + \text{PopF}_j \times \text{PopM}_k + F_j + M_k + \text{pH}_i \times F_j + \text{pH}_i \times M_k + F_j \times M_k + e_{ijk} \quad (1)$$

where Y_{ijk} was the trait value of the sample in pH treatment i with father j and mother k , pH_i , F_j , and M_k are effects of the specific levels of each factor, PopF_j and PopM_k are effects of the source population (Norway or Denmark) of each parent, terms with x 's denote interactions, and μ is a global intercept. The source population terms accounted for any systematic differences (genetic or environmental) between urchins collected from each population. Although genetic differences among populations can serve as a source of variation for adaptation, trans-generational environmental effects cannot. To be conservative, we excluded differences in trait means or responses to pH between source populations from our estimates of the amount of genetic variation in each trait. Given our limited sample size (3–4 fathers and 1–2 mothers per population), we were unable to effectively test for population differences in any trait.

For the gene expression data, we converted counts to log2CPM (counts-per-million, with prior.count = 1) with the function *cpm* (Robinson et al. 2010), calculated sample-weights (accounting for variation in sample quality) with the *arrayWeights* function (Ritchie et al. 2006) and then fit model (1) and calculated smoothed Type III F-statistics for the sets of coefficients associated with each factor using the functions *lmFit* and *eBayes* (Smyth 2004) using the *limma*-trend method (Law et al. 2014). The sample-weights calculated for the gene expression data were carried over to the analysis of the gene set variation traits because these relied on the same gene expression values, but not to the analyses of survival and growth rate traits. Empirical Bayes smoothing of SEs was performed separately for the gene expression traits, each class of gene set traits, growth rate, and the survival traits.

We calculated the percentage of variance explained by a factor as the total weighted sum-of-squares for that factor (including interactions) after accounting for all other factors divided by the total weighted sum-of-squares of the observations around the global mean. We used the *qvalue* package (Storey et al. 2015) to calculate family-wise false discovery rates (FDR) and estimated numbers of true positives ($1 - \pi_0$) for each set of tests. We tested for enrichments of genes with




Fig. 2.—Larval growth rates are reduced in low pH seawater. Boxplots show median and quartiles of the distributions of growth rates for cultures grown in control or low pH seawater ($N=21$ for each). The effect of pH treatment on growth rate was significant ($P=2.78 \times 10^{-8}$). Growth rates (mm/day) were calculated using linear regression of daily measures of ~ 10 larvae/culture over the first 6–9 days of development.

functional annotations among genes with small P values using the Wilcoxon rank-sum test (*Wilcox.test*).

We used multiple linear regressions to test for associations between gene expression variation and variation in performance. For each performance trait (growth rate or survival fraction), we ran a penalized regression of the trait value on either the entire gene expression matrix (22,430 genes), or one of the classes of gene set traits using the LASSO penalty (Tibshirani 1996) implemented in the *R* function *glmnet* (Friedman et al. 2010). For each regression, we used leave-one-out cross-validation to select the tuning parameter lambda and scored models based on the improvement in the mean squared error of prediction.

Results

All trait values are presented in [supplementary table S2, Supplementary Material](#) online, and statistics for all tests of pH and parental effects are presented in [supplementary table S3, Supplementary Material](#) online.

Impact of Seawater pH

We did not observe any effects of low pH seawater on larval mortality ([supplementary table S1, Supplementary Material](#) online). Low pH seawater caused a 14% reduction in larval growth rate (low pH: 0.033 mm/day, control: 0.039 mm/day; $P < 0.0001$, [fig. 2](#)).

Gene expression responses to low pH seawater were common but generally subtle. Using the *qvalue* function

(Storey et al. 2015), we estimated that 43.4% of the 22,430 measured genes were affected by the pH treatment in any of the family groups (cultures sharing a father or a mother), but the average response to low pH across all cultures was smaller than 2-fold for all but nine genes ($\log_2FC < \pm 1$, [fig. 3A, supplementary table S1, Supplementary Material](#) online). Overall, among the 517 genes, we could declare significant at an FDR of 5% (corresponding to a P value threshold of 0.002, [fig. 3A, supplementary table S1, Supplementary Material](#) online), many more genes were upregulated than downregulated at low pH (65%).

In order to gain insight into biological responses to low pH, we used gene ontology annotations to group genes into functionally similar modules and measured how these modules responded to the pH treatment. The set of gene ontology annotations we used included annotations for 14,951 genes, which were significantly enriched for responses to the pH treatment (Wilcoxon sign-rank $W: 5.49 \times 10^7$, $P=0.016$). Using GSVA (Hänzelmann et al. 2013), we combined the expression levels of these 14,951 genes into measures of 616 synthetic gene set traits, divided into five categories (PANTHER ontologies: MF: 128 terms, BP: 165 terms, CC: 22 terms, PC: 171 terms, and hand-annotated ontology: 130 terms). We focused on MF and hand-annotated categories that are the easiest to interpret. Results for all gene set traits are presented in [supplementary table S1, Supplementary Material](#) online. pH responses in the 128 MF gene set traits were larger in magnitude (relative to the total among-sample variance, Wilcoxon sign-rank $W: 1.73 \times 10^6$, $P < 0.0001$) and were more common than the responses to pH of the individual genes ([fig. 3B](#)). We estimated that 60.9% of these MF gene set traits were affected by the pH treatment. pH treatment effects (either as main effects or as interactions with genetic backgrounds) accounted for $>25\%$ of the among-sample variation in 38 of these traits and $>50\%$ of the variation in 2. The bubble plot in [figure 4](#) summarizes the 25 MF gene set traits that we could declare to be significantly affected by low pH in any cohort at a 5% FDR. Several key groups of MF gene sets constitute the core of the response to pH. Downregulated gene sets were mostly related to DNA and RNA metabolism and cell division. Upregulated gene sets were related to cell signaling and other transmembrane functions such as ion channel activities and G-protein coupled receptor activity.

Results were largely congruent for the 130 gene set traits based on the hand-annotated ontologies. Low pH seawater affected 71.2% of these traits and explained $>25\%$ of the variance in 46. The 62 hand-annotated gene set traits we could declare to be significantly affected by low pH in any cohort primarily involved processes such as the cell cycle, cell–cell signaling and immune-related processes ([supplementary table S3, Supplementary Material](#) online). The category of




Fig. 3.—Physiological and molecular responses attributed to seawater pH are common but subtle relative to variation attributed to father or mother effects. (A) Mean (across all cultures) log₂ responses to low pH seawater for 22,430 genes plotted against mean log₂(expression). Genes with significant responses to low pH seawater (5% FDR) are highlighted in blue. (B) Boxplots show median and quartiles of the distributions of percentage of total among-culture variation in each gene expression trait ($n = 22,430$) or each MF gene set ($n = 128$) trait that could be attributed to each experimental factor (mother, father or pH).




Fig. 4.—Summary of the response to pH in Molecular Function gene set traits. Bubble plot of MF gene set variation traits with significant (5% FDR) responses to low pH seawater. Bubble plots represent: (i) The a priori relationship among the MF gene sets. Bubble centers are arranged based on a multidimensional scaling projection (*R* function *sammon*) of the *SimREL* distances (Schlicker et al. 2006) among the PANTHER MF ontology terms. This distance takes into account both the tree-relationships among terms and the number of genes in each category. (ii) The number of genes linked to each MF term (bubble area is proportional to gene number). (iii) The percentage of variation in each gene set explained by the pH treatment and the mean direction of the response to low pH (orange = increase in expression, blue = decrease in expression). This plot is based on the REVIGO scatter plot (Supek et al. 2011).

biomineralization genes was significantly downregulated by low pH (-0.204).

Genetic and Maternal Effects on Larval Traits

Across both performance and molecular traits, differences associated with larval parentage tended to be larger and more

common than differences induced by low seawater pH (fig. 3B). We detected significant mother effects on growth rate ($P = 0.001$), but no significant father effects ($P = 0.44$). Among the 22,430 genes, 77.4% had expression variation associated with inherited father effects and 48.9% had expression variation associated with mother effects (note that the experiment included seven fathers and only three mothers,

but only among parent differences within the same source population were counted). Again, genes with functional annotations were enriched for significant father or mother effects relative to unannotated genes. Gene expression profiles integrated into gene set traits also showed high levels of parent-associated variation: 93.8% of the 128 MF gene set traits showed father-associated variation and 55.7% showed mother-associated variation.

Interactions between Genetic Variation and the Response to Low pH

By splitting cultures of genetically related larvae between the two pH treatments, we were able to test for differences in responses to pH among culture with different fathers or mothers, and differences in the father effects when crossed to different mothers. We estimated that 20% of genes had a different expression response to pH across families with different fathers and 6.7% of genes had a different response across cultures with different mothers. Among the 128 MF gene set traits, 1% had a different response to pH among cultures with different mothers, but we found no evidence for a different response to pH in any of these traits among cultures with different fathers. We estimated that 6.7% of genes and 28.6% of MF gene set traits had different father effects depending on the mother. However, due to the low level of replication of these interactions in our experimental design, we were not able to declare more than a handful to be significant while maintaining a $FDR < 5\%$ (supplementary table S3, Supplementary Material online) and the true frequencies of these gene–environment, or gene–gene interactions may be higher.

Relationship between Molecular Traits and Larval Performance

To test if the molecular traits could help explain variation in larval performance, we used multiple regression with a LASSO penalty to try to predict the variation in growth rate among cultures based on the whole matrix of 22,430 genes, or each of the five classes of gene set traits. We used the LASSO penalty because it provides variable selection among the predictors and helps with model regularization when there are more predictors than samples. The BP and hand-annotated gene set traits were the best predictors of growth rate variation, as measured by cross-validation performance, and outperformed the raw gene expression data (table 2). The best model selected 25 BP gene set traits and improved the mean squared error by 79.2% relative to a model with only culture pH. These 25 BP gene set traits are displayed in figure 5. In this model, the BP traits pyrimidine base metabolic process (positive) and RNA localization (negative) were selected as the most strongly associated with growth rate.

Table 2

Percent Decrease in Mean Squared Error of Prediction (MSE) Values Relative to pH Treatment Alone for the Regression of Growth Rate on Each Set of Molecular Traits

Molecular Trait Class	Growth Rate
BP	79.2% (24)
MF	46.8% (13)
CC	18.5% (9)
PC	36.9% (29)
Hand	64.7% (41)
Gene expression	56% (40)

NOTE.—MSE was calculated using the *cv.glmnet* function with $\alpha=1$ for the LASSO penalty. Values represent $1 - \text{MSE}(\text{full})/\text{MSE}(\text{pH})$ for each model at the optimal value of the lambda tuning parameter. The number in parentheses is the number of molecular traits with non-zero regression coefficients in the best model.

Discussion

Molecular Responses to Low pH

Our results demonstrate that the molecular response to low pH in larvae of *S. droebachiensis* is wide-ranging and complex. We observed plasticity in response to low pH in the expression of nearly half of all assayed genes, and in three quarters of the higher-order molecular traits inferred by integrating the expression variation across groups of functionally related genes. Previous studies investigating the effects of low pH on gene expression in sea urchin larvae have reported changes in the expression of genes related to calcification and biomineralization (Todgham and Hofmann 2009; Martin et al. 2011; Stumpp et al. 2011a; Padilla-Gamiño et al. 2013; Evans et al. 2013), cytoskeleton and cell division (Todgham and Hofmann 2009; Padilla-Gamiño et al. 2013), acid/base regulation (Todgham and Hofmann 2009; Stumpp et al. 2011a; Stumpp et al. 2015), and metabolism (Todgham and Hofmann 2009; Stumpp et al. 2011a), with the direction and strengths of the responses varying by species and the particulars of the environment treatments (e.g., simultaneous increases in temperature). We observed changes in many of these same processes in *S. droebachiensis* larvae. This is consistent with physiological studies showing that *S. droebachiensis* pluteus larvae are unable to compensate for an extracellular acidosis (pHe) resulting from an exposure to low pH. However, the calcifying primary mesenchyme cells are able to fully compensate an induced intracellular acidosis (pHi) using a bicarbonate buffer mechanism involving secondary active Na^+ -dependent membrane transport proteins (Stumpp et al. 2012). Additional energetic costs also derived from compensatory mechanisms associated with larval gastric pH changes (Stumpp et al. 2013). These extra costs lead to a shift in energy budget, with less energy available for growth and leading to a delay in development (Dupont and Thorndyke 2013; Jager et al. 2016). Our results also highlighted a strong signal of increased expression of many genes involved in immune




Fig. 5.—Biological process gene set traits associated with variation in larval growth rates. Bubble plot show the 22 BP gene set traits selected by the LASSO penalized regression as important predictors of growth rate variation. Color hue and intensity represents the sign and magnitude of each of the non-zero regression coefficients in the best model. The lambda parameter was chosen by leave-one-out cross validation.

responses and other responses to extracellular environments, perhaps indicating an involvement of the immune system in pH tolerance or an increased sensitivity of the larvae to infections or additional environmental stresses. This is in contrast with studies of mussels (Bibby et al. 2008) and starfish (Hernroth et al. 2011) both showed a depressed immune system under low pH but consistent with an increased in cellular immune response in two adult echinoderm species, including *S. droebachiensis* (Dupont and Thorndyke 2012). Dupont and Thorndyke (2012) hypothesized that there may be a direct link between pHe and cellular immune-response. As it was demonstrated that decreased environmental pH also lead to an uncompensated pHe decrease in pluteus larvae, the observed upregulation of the immune system may result from such a link. However, other measures of cellular stress did not show evidence of a pH response: no consistent expression changes were observed in the class of heat shock proteins, or other molecular chaperones, unlike what has been seen in other environmental stressors, such as temperature (Hammond and Hofmann 2010; Runcie et al. 2012).

The increased number of significant effects in our data relative to previous studies is largely a function of our increased sample size (21 cultures per treatment), as the magnitude of pH effects on gene expression were generally small as previously reported (Todgham and Hofmann 2009; Pespeni et al. 2013). Despite this molecular plasticity, we found a general robustness of larval development to predicted near-future levels of pH, with the larvae appearing healthy despite slower growth rates and no increase in larval mortality due to pH. This replicates similar findings in previous studies (Dupont and Thorndyke 2008; Martin et al. 2011; Stumpp et al. 2011b). In the absence of data from long-term cultures, however, potential impacts of lower pH on survival to metamorphosis and lifetime fecundity remain unclear.

Natural Variation in Larval Traits

Despite clear evidence of physiological, developmental, and growth impacts of pH on the larvae, the pH treatments did not account for a particularly large portion of the total gene expression variation among cultures of larvae. pH treatment effects accounted for <25% of the among-culture variation in the majority of the gene expression traits measured (fig. 3B). Instead, our results show that genetic variants and/or effects of differences in paternal or maternal environments contributed to a much larger fraction of the trait variation (fig. 3B). Although it is not possible using our experimental design to unequivocally differentiate between paternal environmental effects and genetic variation, a high frequency and magnitude of genetic variation in gene expression has been observed before in sea urchins (Runcie et al. 2012; Garfield et al. 2013), as well as in other taxa including fish, plants, *Drosophila*, and primates (Oleksiak et al. 2002; Schadt et al. 2003; Nuzhdin et al. 2008; Ayroles et al. 2009). We extended these results by demonstrating that there is a high level of genetic variation in higher-order molecular traits, captured by the gene set traits defined by ontology terms from the PANTHER gene ontology (Mi et al. 2013), and by the sea urchin-specific gene function ontology (Sodergren et al. 2006; Tu et al. 2012). For example, the father and mother effects on the MF signatures were as frequent and of as large a magnitude (proportionally to the total amount of trait variation) as the parental effects on the expression of individual genes.

Prospects for Evolutionary Rescue

The large amount of genetic variation in molecular gene expression and survival fractions suggests that rapid adaptation to low pH seawater might be possible in this species. Large amounts of standing genetic variation can accelerate

adaptation if the variation contributes to variation in fitness, and if genetic correlations among traits with contrasting effects on fitness are not too strong (Walsh and Blows 2009). We observed considerably more genetic variation (father-associated variation) in the molecular traits than in larval growth rates (fig. 3B). It is possible that genetic variation can accumulate in low-level traits like gene expression because gene networks are structured to buffer much of this variation from affecting high-order phenotypes such as morphology (Runcie et al. 2012; Garfield et al. 2013). However, at least a portion of the variation in the gene expression traits was correlated with variation in growth rate (fig. 5 and table 2) and thus may be relevant for fitness. It is also possible that this molecular variation may be a source of "cryptic" genetic variation that could be exposed by changing environments (Gibson and Dworkin 2004), and thus accelerate adaptation in changing environments. We found evidence for gene and environment interactions being important for a small percentage of the gene expression traits, but our sample sizes were too small to clearly identify any patterns as to which traits might show a change in genetic variance in acidified seawater. The fact that we observed as strong evidence for genetic variation in the higher order gene set traits as in the individual genes suggests that much of this genetic variation may be highly pleiotropic. The possibility of extensive pleiotropy is an important caveat in the interpretation of our finding that many genes show variation in expression. This is because pleiotropy places genetic constraints on the direction of evolutionary change (Walsh and Blows 2009), and thus the capacity of a population to adapt in any particular way may be more limited than the magnitude of genetic variation in each trait initially suggests.

Another limitation of our finding is the use of the two scenarios used in this study: the average pH at present (pH_T 8.0) and the average pH projected for 2100 (pH_T 7.6). However, both pH values are within the present range of natural variability (Dorey et al. 2013). Previous physiological works on the impact of acidification across a wide range of pH on the development of *S. droebachiensis* have demonstrated a clearly pH-dependent increase in stress with decreasing pH, including an apparent tipping point around a pH of 7.5 (Jager et al. 2016). Below that threshold, different physiological processes were involved leading to increased mortality at low pH (Dorey et al. 2013). Different evolutionary processes may also be involved, as demonstrated in copepods. When exposed to decreased pH with the present day's range of natural variability, copepods responded mostly through phenotypic plasticity. When exposed to pH below present day's natural variability, transgenerational effects (including genetic adaptation) set in (Thor and Dupont 2015; De Wit et al. 2015).

Gene Expression Profiling in Climate Change Research

The use of RNAseq to address problems in ecological and evolutionary functional genomics is growing rapidly and shows

great potential to greatly improve our understanding of the key traits underlying adaptation and acclimation to changing climates (Gracey 2007; Whitehead 2012; Alvarez et al. 2015; Papetti et al. 2016; Todd et al. 2016). Transcriptome profiling by RNAseq can now be applied to nearly any organism. The principal advantage of RNAseq for climate change research is that it rapidly profiles a wide range of molecular processes, many of which are difficult to measure directly (Evans and Hofmann 2012). For example, techniques to measure respiration and immune system activity are technically challenging to apply to large numbers of individuals (Hernroth et al. 2011; Stumpp et al. 2011b). With gene expression profiling, we can quickly sort through hundreds of conceivable molecular processes to identify ones that are likely relevant and are worth further study. As sequencing costs continue to decline rapidly, it is becoming feasible to measure gene expression in many hundreds or even thousands of samples. Such sample sizes are necessary for accurately estimating genetic variances and genetic correlations among traits that are necessary for predicting evolutionary change (Roff 1995).

By themselves, transcriptome-wide gene expression data can be overwhelming in their scale. Individual gene expression measurements are often too noisy (whether due to measurement error or systematic variation such as genetic effects) to use as reliable signatures of underlying molecular trait variation. Directly incorporating information on gene function (Hänzelmann et al. 2013) and modeling the correlation among traits (Runcie and Mukherjee 2013) can dramatically increase both the power to detect molecular variation, and make the results easier to interpret. There are important limitations: characterizing traits based on functional classification of genes is inherently imprecise would benefit from explicit information about how genes in gene sets interact (Khatri et al. 2012) and requires follow-up studies to directly measure the traits identified by gene expression signatures. Nevertheless, by simultaneously measuring many aspects of physiology and development, gene expression can provide a vastly more comprehensive estimate of the potential for adaptation to climate change (De Wit et al. 2015; Rose et al. 2016).

Supplementary Material

Supplementary data are available at *Genome Biology and Evolution* online.

Acknowledgments

This work was supported by the National Science Foundation [grant numbers 1202838 to D.E.R., 0614509 to G.A.W.]; the Linnaeus Centre for Marine Evolutionary Biology at the University of Gothenburg; and the Swedish Research Councils VR and Formas (Linnaeus grant) to N.D. and S.D. We would like to thank Bengt Lundve and Aude Griffet for their assistance with rearing the urchin cultures.

Literature Cited

- Alvarez M, Schrey AW, Richards CL. 2015. Ten years of transcriptomics in wild populations: what have we learned about their ecology and evolution?. *Mol Ecol* 24:710–725.
- Anders S, Pyl PT, Huber W. 2015. *HTSeq* - a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31:166–169.
- Ayroles JF, et al. 2009. Systems genetics of complex traits in *Drosophila melanogaster*. *Nat Genet.* 41:299–307.
- Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate - a practical and powerful approach to multiple testing. *J Roy Stat Soc B.* 57:289–300.
- Bibby R, Widdicombe S, Parry H, Spicer J, Pipe R. 2008. Effects of ocean acidification on the immune response of the blue mussel *Mytilus edulis*. *Aquat Biol.* 2:67–74.
- Breitburg DL, et al. 2015. And on top of all that... coping with ocean acidification in the midst of many stressors. *Oceanog* 25:48–61.
- Bürger R, Lynch M. 1995. Evolution and extinction in a changing environment: a quantitative-genetic analysis. *Evolution* 49:151–163.
- Byrne M. 2011. Impact of ocean warming and ocean acidification on marine invertebrate life history stages: vulnerabilities and potential for persistence in a changing ocean. In: Gibson, RN, Atkinson, R, Gordon, J, Smith, IP, & Hughes, DJ, editors. *Oceanography and marine biology: an annual review*. Vol. 49. Taylor & Francis, New York: CRC Press. p. 1–42.
- Caldeira K, Wickett ME. 2003. Oceanography: anthropogenic carbon and ocean pH. *Nature* 425:365.
- Cheviron ZA, Connaty AD, McClelland GB, Storz JF. 2014. Functional genomics of adaptation to hypoxic cold-stress in high-altitude deer mice: transcriptomic plasticity and thermogenic performance. *Evolution* 68:48–62.
- Chown SL, Gaston KJ. 2008. Macrophysiology for a changing world. *Proc Biol Sci.* 275:1469–1478.
- Civelek M, Lusi AJ. 2013. Systems genetics approaches to understand complex traits. *Nat Rev Genet.* 15:34–48.
- Crean AJ, Dwyer JM, Marshall DJ. 2013. Adaptive paternal effects? Experimental evidence that the paternal environment affects offspring performance. *Ecology* 94:2575–2582.
- Curley JP, Mashoodh R, Champagne FA. 2011. Epigenetics and the origins of paternal effects. *Horm Behav* 59:306–314.
- Davis MB, Shaw RG. 2001. Range shifts and adaptive responses to Quaternary climate change. *Science* 292:673–679.
- Davis MB, Shaw RG, Etterson JR. 2005. Evolutionary responses to changing climate. *Ecology* 86:1704–1714.
- De Wit P, Dupont S, Thor P. 2015. Selection on oxidative phosphorylation and ribosomal structure as a multigenerational response to ocean acidification in the common copepod *Pseudocalanus acuspes*. *Evol Appl.* 9:1112–1123.
- Doney SC, Fabry VJ, Feely RA, Kleypas JA. 2009. Ocean acidification: the other CO₂ problem. *Ann Rev Mar Sci.* 1:169–192.
- Dorey N, Lancon P, Thorndyke MC, Dupont S. 2013. Assessing physiological tipping point of sea urchin larvae exposed to a broad range of pH. *Glob Chang Biol.* 19:3355–3367.
- Dupont S, Ortega-Martinez O, Thorndyke MC. 2010. Impact of near-future ocean acidification on echinoderms. *Ecotoxicology* 19:449–462.
- Dupont S, Pörtner HO. 2013. A snapshot of ocean acidification research. *Mar Biol.* 160:1765–1771.
- Dupont S, Thorndyke MC. 2008. Ocean acidification and its impact on the early life-history stages of marine animals. In: *Impacts of acidification on biological, chemical and physical systems in the Mediterranean and Black Seas*. CIESM Workshop Monographs n. 36 pp. 89–97.
- Dupont S, Thorndyke MC. 2012. Relationship between CO₂-driven changes in extracellular acid–base balance and cellular immune response in two polar echinoderm species. *J Exp Mar Biol Ecol* 424–425:32–37.
- Dupont S, Thorndyke MC. 2013. Direct impacts of near-future ocean acidification on sea urchins. In: Fernández-Palacios JM, de Nascimento L, Hernández JC, Clemente S, González A & Díaz-González JP, editors. *Climate change perspective from the Atlantic: past, present and future*. Servicio de Publicaciones, Universidad de La Laguna. pp. 461–485.
- Evans TG, Chan F, Menge BA, Hofmann GE. 2013. Transcriptomic responses to ocean acidification in larval sea urchins from a naturally variable pH environment. *Mol Ecol* 22:1609–1625.
- Evans TG, Hofmann GE. 2012. Defining the limits of physiological plasticity: how gene expression can assess and predict the consequences of ocean change. *Philos T Roy Soc B.* 367:1733–1745.
- Foo SA, Dworjanyn SA, Poore AGB, Byrne M. 2012. Adaptive capacity of the habitat modifying sea urchin *Centrostephanus rodgersii* to ocean warming and ocean acidification: performance of early embryos. *PLoS ONE* 7:e42497.
- Friedman J, Hastie T, Tibshirani R. 2010. Regularization paths for generalized linear models via coordinate descent. *J Stat Softw* 33:1–22.
- Garfield DA, et al. 2013. The impact of gene expression variation on the robustness and evolvability of a developmental gene regulatory network. *PLoS Biol.* 11:e1001696.
- Gattuso J-P, Epitalon J-M, Lavigne H. 2015. Seawater Carbonate Chemistry. <http://CRAN.R-project.org/package=seacarb>.
- Gaylord B, et al. 2015. Ocean acidification through the lens of ecological theory. *Ecology* 96:3–15.
- Gianguzza P, et al. 2014. Temperature modulates the response of the thermophilous sea urchin *Arbacia lixula* early life stages to CO₂-driven acidification. *Mar Environ Res.* 93:70–77.
- Gibson G, Dworkin I. 2004. Uncovering cryptic genetic variation. *Nat Rev Genet.* 5:681–690.
- Gienapp P, Teplitsky C, Alho JS, Mills JA, Merilä J. 2008. Climate change and evolution: disentangling environmental and genetic responses. *Mol Ecol.* 17:167–178.
- Gomulkiewicz R, Holt RD. 1995. When does evolution by natural selection prevent extinction? *Evolution* 49:201.
- Grace AY. 2007. Interpreting physiological responses to environmental change through gene expression profiling. *J Exp Biol.* 210:1584–1592.
- Hammond LM, Hofmann GE. 2010. Thermal tolerance of *Strongylocentrotus purpuratus* early life history stages: mortality, stress-induced gene expression and biogeographic patterns. *Mar Biol.* 157:2677–2687.
- Hänzelmann S, Castelo R, Guinney J. 2013. GSEA: gene set variation analysis for microarray and RNA-seq data. *BMC Bioinformatics* 14:7.
- Hendry AP, Farrugia TJ, Kinnison MT. 2008. Human influences on rates of phenotypic change in wild animal populations. *Mol Ecol* 17:20–29.
- Hernroth B, Baden S, Thorndyke M, Dupont S. 2011. Immune suppression of the echinoderm *Asterias rubens* (L.) following long-term ocean acidification. *Aquat Toxicol.* 103:222–224.
- Houle D. 2010. Colloquium papers: numbering the hairs on our heads: the shared challenge and promise of phenomics. *Proc. Natl. Acad. Sci. U.S.A* 107 (Suppl 1):1793–1799.
- IPCC. 2014. *Climate Change 2013: The Physical Science Basis*. Cambridge, UK: Cambridge University Press.
- Jager T, Ravagnan E, Dupont S. 2016. Near-future ocean acidification impacts maintenance costs in sea-urchin larvae: Identification of stress factors and tipping points using a DEB modelling approach. *J Exp Mar Biol Ecol.* 474:11–17.
- Kelly MW, Padilla-Gamiño JL, Hofmann GE. 2013. Natural variation and the capacity to adapt to ocean acidification in the keystone sea urchin *Strongylocentrotus purpuratus*. *Glob Chang Biol.* 19:2536–2546.
- Khatri P, Sirota M, Butte AJ. 2012. Ten years of pathway analysis: current approaches and outstanding challenges. *PLoS Comput Biol.* 8:e1002375.

- Krawetz SA. 2005. Paternal contribution: new insights and future challenges. *Nat Rev Genet.* 6:633–642.
- Lande R, Arnold S. 1983. The measurement of selection on correlated characters. *Evolution.* 37(6):1210–1226.
- Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359.
- Langmead B, Trapnell C, Pop M, Salzberg SL. 2009 Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* 10:R25.
- Law CW, Chen Y, Shi W, Smyth GK. 2014. *voom*: precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biology* 15:R29.
- Lynch M, Lande R. 1993. Evolution and extinction in response to environmental change. In: Kareiva, PM, Kingsolver, JG, and Huey, RB, editors. *Biotic interactions and global change.* Sunderland (MA): Sinauer Associates Inc. p. 234–250.
- Lynch M, Walsh B. 1998. *Genetics and analysis of quantitative traits.* 1st ed. Sunderland (MA): Sinauer Associates Inc.
- Marais Des DL, et al. 2012. Physiological genomics of response to soil drying in diverse *Arabidopsis* accessions. *Plant Cell* 24:893–914.
- Martin S, et al. 2011. Early development and molecular plasticity in the Mediterranean sea urchin *Paracentrotus lividus* exposed to CO₂-driven acidification. *J Exp Biol.* 214:1357–1368.
- Mi H, Muruganujan A, Thomas PD. 2013. PANTHER in 2013: modeling the evolution of gene function, and other gene attributes, in the context of phylogenetic trees. *Nucl. Acids Res.* 41:D377–D386.
- Nuzhdin SV, Tufts DM, Hahn MW. 2008 Abundant genetic variation in transcript level during early *Drosophila* development. *Evol Dev.* 10:683–689.
- Oleksiak MF, Churchill GA, Crawford DL. 2002. Variation in gene expression within and among natural populations. *Nat Genet.* 32:261–266.
- Padilla-Gamiño JL, Kelly MW, Evans TG, Hofmann GE. 2013. Temperature and CO₂ additively regulate physiology, morphology and genomic responses of larval sea urchins, *Strongylocentrotus purpuratus*. *Proc Roy Soc B.* 280:20130155.
- Pan TCF, Applebaum SL, Manahan DT. 2015. Experimental ocean acidification alters the allocation of metabolic energy. *Proc. Natl. Acad. Sci. U.S.A.* 112:4696–4701.
- Papetti C, Lucassen M, Pörtner H-O. 2016. Integrated studies of organismal plasticity through physiological and transcriptomic approaches: examples from marine polar regions. *Brief Funct Genomics* 15:365–372.
- Parmesan C, Yohe G. 2003. A globally coherent fingerprint of climate change impacts across natural systems. *Nature* 421:37–42.
- Pespeni MH, et al. 2013. Evolutionary change during experimental ocean acidification. *Proc Natl Acad Sci. U S A.* 110:6937–6942.
- Pörtner HO, Dupont S, Melzner F, Storch D. 2010. Studies of metabolic rate and other characters across life stages. In: Riebesell, U, Fabry, VJ, Hansson, L, and Gattuso, J-P, editors. *Guide to best practices for ocean acidification research and data reporting.* Luxembourg: Publications Office of the European Union. p. 167–180.
- R Development Core Team. 2015. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- Ritchie ME, et al. 2006. Empirical array quality weights in the analysis of microarray data. *BMC Bioinformatics* 7:261.
- Ritchie ME, et al. 2015. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucl Acids Res.* 43:e47–e47.
- Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26:139–140.
- Robinson MD, Oshlack A. 2010 A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol.* 11:R25.
- Roff D. 1995. The estimation of genetic correlations from phenotypic correlations: a test of Cheverud's conjecture. *Heredity* 74:481–490.
- Rose NH, Seneca FO, Palumbi SR. 2016. Gene networks in the wild: identifying transcriptional modules that mediate coral resistance to experimental heat stress. *Genome Biol Evol.* 8:243–252.
- Runcie D, et al. 2012. Genetics of gene expression responses to temperature stress in a sea urchin gene network. *Mol Ecol.* 21:4547–4562.
- Runcie D, Mukherjee S. 2013. Dissecting high-dimensional phenotypes with bayesian sparse factor analysis of genetic covariance matrices. *Genetics* 194:753–767.
- Schadt EE, et al. 2003. Genetics of gene expression surveyed in maize, mouse and man. *Nature* 422:297–302.
- Schlicker A, Domingues FS, Rahnenführer J, Lengauer T. 2006. A new measure for functional similarity of gene products based on Gene Ontology. *BMC Bioinformatics* 7:302.
- Schneider RF, Li Y, Meyer A, Gunter HM. 2014. Regulatory gene networks that shape the development of adaptive phenotypic plasticity in a cichlid fish. *Mol Ecol.* 23:4511–4526.
- Smyth GK. 2004. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mo B.* 3:Article3–Articl25.
- Sodergren E, et al. 2006. The genome of the sea urchin *Strongylocentrotus purpuratus*. *Science* 314:941–952.
- Somero GN. 2010. The physiology of climate change: how potentials for acclimatization and genetic adaptation will determine 'winners' and 'losers'. *J Exp Biol.* 213:912–920.
- Storey JD, Bass AJ, Dabney A, Robinson D. 2015. *qvalue: Q-value estimation for false discovery rate control.*
- Stumpp M, et al. 2012. Acidified seawater impacts sea urchin larvae pH regulatory systems relevant for calcification. *Proc Natl Acad Sci U S A.* 109:18192–18197.
- Stumpp M, et al. 2013. Digestion in sea urchin larvae impaired under ocean acidification. *Nat Clim Chang.* 3:1044–1049.
- Stumpp M, et al. 2015. Evolution of extreme stomach pH in bilateria inferred from gastric alkalization mechanisms in basal deuterostomes. *Sci Rep.* 5:10421.
- Stumpp M, Dupont S, Thorndyke MC, Melzner F. 2011a CO₂ induced seawater acidification impacts sea urchin larval development II: gene expression patterns in pluteus larvae. *Comp Biochem Physiol A Mol Integr Physiol.* 160:320–330.
- Stumpp M, Wren J, et al. 2011b. CO₂ induced seawater acidification impacts sea urchin larval development I: elevated metabolic rates decrease scope for growth and induce developmental delay. *Comp Biochem Physiol A Mol Integr Physiol.* 160:331–340.
- Subramanian A, et al. 2005. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci. U S A.* 102:15545–15550.
- Sunday JM, Crim RN, Harley CDG, Hart MW. 2011. Quantifying rates of evolutionary adaptation in response to ocean acidification. *PLoS ONE* 6:e22881.
- Supek F, Bošnjak M, Škunca N, Šmuc T. 2011. REVIGO summarizes and visualizes long lists of gene ontology terms. *PLoS ONE* 6:e21800.
- Thor P, Dupont S. 2015. Transgenerational effects alleviate severe fecundity loss during ocean acidification in a ubiquitous planktonic copepod. *Glob Chang Biol.* 21:2261–2271.
- Tibshirani R. 1996. Regression Shrinkage and Selection via the Lasso. *J Roy Stat Soc B.* 58:267–288.
- Todd EV, Black MA, Gemmell NJ. 2016. The power and promise of RNA-seq in ecology and evolution. *Mol Ecol.* 25:1224–1241.
- Todgham AE, Hofmann GE. 2009. Transcriptomic response of sea urchin larvae *Strongylocentrotus purpuratus* to CO₂-driven seawater acidification. *J Exp Biol.* 212:2579–2594.
- Tu Q, Cameron RA, Worley KC, Gibbs RA, Davidson EH. 2012. Gene structure in the sea urchin *Strongylocentrotus purpuratus* based on transcriptome analysis. *Genome Res.* 22:2079–2087.

- Vander Wal E, Garant D, Festa-Bianchet M, Pelletier F. 2013. Evolutionary rescue in vertebrates: evidence, applications and uncertainty. *Philos Trans R Soci Lond B*. 368:20120090–20120090.
- Walsh B, Blows MW. 2009. Abundant genetic variation plus strong selection = multivariate genetic constraints: a geometric view of adaptation. *Annu Rev Ecol Evol S* 40:41–59.
- Warnes GR, Bolker B, Bonebakker L, Gentleman R, Liaw WHA, Lumley T, Maechler M, Magnusson A, Moeller S, Schwartz M, Venables B. 2015. gplots: Various R Programming Tools for Plotting Data. Data. R package version 2.17.0. <http://CRAN.R-project.org/package=gplots>.
- Whitehead A. 2012. Comparative genomics in ecological physiology: toward a more nuanced understanding of acclimation and adaptation. *J Exp Biol*. 215:884–891.
- Wikelski M, Cooke SJ. 2006. Conservation physiology. *Trends Ecol Evol*. 21:38–46.
- Williams SE, et al. *PLoS Biol*. 6:e325.

Associate editor: Ross Hardison