

Published in final edited form as:

*Science*. 2013 March 1; 339(6123): 1063–1067. doi:10.1126/science.1230422.

## Genomic diversity and evolution of the head crest in the rock pigeon

Michael D. Shapiro<sup>1,†</sup>, Zev Kronenberg<sup>2</sup>, Cai Li<sup>3,4</sup>, Eric T. Domyan<sup>1</sup>, Hailin Pan<sup>3</sup>, Michael Campbell<sup>2</sup>, Hao Tan<sup>3</sup>, Chad D. Huff<sup>2,5</sup>, Haofu Hu<sup>3</sup>, Anna I. Vickrey<sup>1</sup>, Sandra C.A. Nielsen<sup>4</sup>, Sydney A. Stringham<sup>1</sup>, Hao Hu<sup>5</sup>, Eske Willerslev<sup>4</sup>, M. Thomas P. Gilbert<sup>4,6</sup>, Mark Yandell<sup>2</sup>, Guojie Zhang<sup>3</sup>, and Jun Wang<sup>3,7,8,†</sup>

<sup>1</sup>Department of Biology, University of Utah, Salt Lake City, UT 84112, USA

<sup>2</sup>Department of Human Genetics, University of Utah, Salt Lake City, UT 84112, USA

<sup>3</sup>BGI-Shenzhen, Shenzhen, 518083, China

<sup>4</sup>Centre for GeoGenetics, Natural History Museum of Denmark, University of Copenhagen, Øster Voldgade 5-7, 1350 Copenhagen, Denmark

<sup>5</sup>Department of Epidemiology, The University of Texas M.D. Anderson Cancer Center, Houston, TX 77030, USA

<sup>6</sup>Ancient DNA Laboratory, Murdoch University, WA 6150, Australia

<sup>7</sup>Department of Biology, University of Copenhagen, DK-1165 Copenhagen, Denmark

<sup>8</sup>The Novo Nordisk Foundation Center for Basic Metabolic Research, University of Copenhagen, DK-1165 Copenhagen, Denmark

### Abstract

The geographic origins of breeds and genetic basis of variation within the widely distributed and phenotypically diverse domestic rock pigeon (*Columba livia*) remain largely unknown. We generated a rock pigeon reference genome and additional genome sequences representing domestic and feral populations. We find evidence for the origins of major breed groups in the Middle East, and contributions from a racing breed to North American feral populations. We identify *EphB2* as a strong candidate for the derived head crest phenotype shared by numerous breeds, an important trait in mate selection in many avian species. We also find evidence that this trait evolved just once and spread throughout the species, and that the crest originates early in development by the localized molecular reversal of feather bud polarity.

Since the initial domestication of the rock pigeon in Neolithic times (1), breeders have selected striking differences in behavior, vocalizations, skeletal morphology, feather ornaments, colors, and color patterns to establish over 350 breeds (2). In many cases, the number and magnitude of differences among breeds are more characteristic of macroevolutionary changes than of changes within a single species (2, 3). Indeed, Charles Darwin was so fascinated by domestic pigeons that he repeatedly called attention to this

<sup>†</sup>Authors for correspondence: mike.shapiro@utah.edu, wangj@genomics.org.cn.

Contact person during manuscript review:

Michael D. Shapiro, Department of Biology, University of Utah, 257 South 1400 East, Salt Lake City, UT 84112 USA; phone: (801) 581 5690; fax: (801) 581 4668; mike.shapiro@utah.edu

This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession AKCR000000000 (first version described here is AKCR010000000; raw reads, SRA052637); RNA-seq data for annotation, GSE39333; raw reads for resequenced genomes, SRA054391.

dramatic example of diversity within a species to communicate his ideas about natural selection (3, 4).

The genetic architecture for many derived traits in pigeons is probably relatively simple (5, 6), likely more so than interspecific trait variation among many wild species, as breeders often focus on qualitative rather than quantitative variation; this increases the chance of identifying genes responsible for differences among breeds. Additionally, several morphological traits show similar patterns of variation in different breeds, making it possible to test whether the same or different genes underlie similar phenotypes. Despite these advantages, the pigeon is underused as a model for the molecular genetic basis of avian variation due to the paucity of genetic and genomic resources for this bird.

We examined genomic diversity, genetic structure, and phylogenetic relationships among domestic breeds and feral populations (free-living birds descended from escaped domestics) of the rock pigeon. The pigeon reference genome was sequenced from a male Danish tumbler with the Illumina HiSeq2000 platform, and we also resequenced 40 additional *C. livia* genomes to 8-to 26-fold coverage (38 individuals from 36 domestic breeds and two feral pigeons) (7). Genome-wide nucleotide diversity in the rock pigeon ( $\pi=3.6\times10^{-3}$ ) and the mutation rate estimate in the pigeon lineage ( $1.42\times10^{-9}$  substitutions site<sup>-1</sup> year<sup>-1</sup>  $\pm$   $2.60\times10^{-12}$  SE) are comparable to other avian species (8, 9). Observed heterozygosity indicates a large effective population size for the rock pigeon of  $N_e\approx521,000$ ; demographic inferences based on the allele frequency spectrum indicate that, aside from a very recent bottleneck,  $N_e$  has been remarkably stable over the past 1.5 million generations (7).

Patterns of linkage disequilibrium (LD) are indicative of haplotype sizes and genome-wide recombination rates, and inform decisions about genetic mapping strategies. Using genotype data from the 40 resequenced *C. livia* genomes, we found that mean “useful LD” (10) ( $r^2>0.3$ ) decays in 2.2 kb (Fig. S10J). This suggests that we should expect little LD between typical pairs of genes in an analysis across breeds; thus, the pigeon is well suited for association-mapping strategies.

We leveraged our whole-genome data to determine breed relationships using 1.48 million variable loci. A neighbor-joining tree rooted on *C. rupestris*, the sister species of *C. livia* (11), yielded several well-supported groups (Figs. 1, S16). Notably, the two feral pigeons grouped with the wattle and homer breeds (Fig. 1, pink branches), supporting the idea that escaped racing homers are probably major contributors to feral populations (12). As with many domesticated species, pigeon evolution is probably not exclusively linear or hierarchical (12). We therefore examined genetic structure among breeds by analyzing 3,950 loci with ADMIXTURE (13), and found a best model fit at  $K=1$  (a single population, where  $K$  is the number of assumed ancestral populations). However, higher values of  $K$  can also be biologically informative (Figs. S17–S20). Our analysis includes some of the oldest lineages of domestic pigeons and breeds that were not exported from the Middle East until the late nineteenth or early twentieth centuries (14), providing information about likely geographic origins of breeds and their exchange along ancient trade routes (7).

Derived traits in domesticated birds tend to evolve along a predictable temporal trajectory, with color variation appearing in the earliest stages of domestication, followed by plumage and structural (skeletal and soft tissue) variation, and finally behavioral differences (2). One of the genetically simplest derived traits of pigeons is the head crest. Head crests are common ornaments in many bird species (2) and are important display structures in mate selection (15). In pigeons, head crests consist of neck and occipital feathers with reversed growth polarity, such that the feathers grow toward the top of the head instead of down the neck. Crests can be as small and simple as a peak of feathers, or as elaborate as the hood of

the Jacobin that envelops the head (Fig. 2A). Remarkably, classical genetics experiments suggest that the head crest segregates as a simple Mendelian recessive trait (6, 14). Moreover, previous studies suggest that the *same* locus controls the presence of a crest in numerous breeds, either with alternative alleles at this locus or additional modifier loci controlling the extent of crest development (6, 14).

We resequenced 8 individuals with head crests to directly test if the same mutation controls crest development in different breeds. We sorted genomic variants from birds with and without head crests into separate bins, and calculated allele frequency differentiation ( $F_{ST}$ ) across the genome (Fig. 2B). We identified a region of high differentiation between crested and uncrested birds in the pigeon ortholog of *Ephrin receptor B2* (*EphB2*;  $F_{ST}=0.94$ , top hit genome wide; Fig. S22A) (Fig. 2D). The role of *EphB2* in feather growth is not known, but it plays important roles in tissue patterning and morphogenesis, and is a member of a receptor tyrosine kinase family that mediates development of the feather cytoskeleton (16, 17). All 8 crested birds were homozygous for a T nucleotide at scaffold 612, position 596613 (hereafter, “*cr*” allele), while uncrested birds were heterozygous ( $n=3$ ) or homozygous ( $n=30$ , including the uncrested outgroup *C. rupestris*) for the putatively ancestral C nucleotide (“+” allele). These results were consistent with the known simple recessive architecture of the trait, and implicated a common polymorphism associated with head crest development in multiple breeds with different genetic histories (Fig. 1). This trend extended well beyond our resequencing panel: we genotyped an additional 61 crested birds from 22 breeds, and 69 uncrested birds from 57 breeds, and found a perfect association between *cr/cr* genotype and the crest phenotype (Fig. 2F). By treating the genomes of crested and uncrested birds as separate populations, we also found suggestive evidence for positive selection around the *cr* allele using cross-population extended haplotype homozygosity analysis (Figs. 2D, S21, S22B).

We then used the Variant Annotation, Analysis, and Search Tool (VAAST (18)) to interrogate the pigeon genomes for additional coding changes associated with the head crest phenotype. This identified one gene with genome-wide significance: *EphB2*, and specifically the *cr* SNP ( $P_{\text{genome}} = 2.0 \times 10^{-8}$ ) (Fig. 2C,D). The *cr* allele has a predicted charge-changing arginine (basic) to cysteine (polar uncharged) transition in the catalytic loop of the intracellular tyrosine kinase domain of *EphB2* (Fig. 2E). This amino acid position is invariant among other vertebrates suggesting strong purifying selection for conserved protein function. Notably, the same DLAARN to DLAACN motif change we observe in *EphB2* is sufficient to abrogate kinase activity in human and mouse orthologs of the protein tyrosine kinase ZAP-70, and in both mammals and pigeons the mutant phenotypes are inherited recessively (19). Hence, the pigeon *cr* mutation probably abrogates kinase activity in *EphB2* and disrupts downstream signal propagation, consistent with the high VAAST score for this gene. *EphB2* is therefore a convincing candidate for the *cr* locus of classical pigeon genetics (5–7, 14).

In several wild and domesticated species, the repeated evolution of a derived trait has occurred by selection on the same gene, possibly due to the repeated selection on the same allele or haplotype (20–22). Similarly, the *cr* SNP is part of a 27.4-kb haplotype that is shared by all crested pigeons, suggesting that the mutation occurred just once and spread to multiple breeds by introgression among domestic breeds, or was selected repeatedly from a standing variant in wild rock pigeons (Figs. 2G, S23; the core haplotype containing the *cr* mutation is reduced to 11 kb with the inclusion of uncrested heterozygotes). The only gene present in the shared *cr* haplotype is *EphB2* (Fig. 2D, green bar), although at this time we cannot rule out the presence of regulatory variants that might alter the expression of another gene. Crested members of the toy, fantail, Iranian, Jacobin, and owl breed groups are not more closely related to each other than to uncrested breeds (Fig. 1). Nevertheless, members

of these groups had head crests hundreds of years ago (14), so some of these introgression events must have occurred in the distant past. Breeds with a wide variety of crest phenotypes share the same derived allele; therefore, allelic variation at the *cr* locus alone does not control all aspects of crest development (14). Other genetic and developmental factors beyond this locus must contribute to variation in crest morphology, akin to the presumed complex genetic architecture of species-level divergence in feather ornaments (2).

In crested pigeons, feather placode polarity and bud outgrowth are inverted during embryogenesis (Fig. 3). Expression of *EphB2* is not polarized in early placodes (Fig. S26), so the effects of the *cr* mutation on feather polarity are probably exerted earlier in development. Why might the crest phenotype be limited to the head and neck? In Naked neck chicken mutants, regionalized production of retinoic acid allows uniform upregulation of *Bmp7* expression to change skin phenotypes in the neck but not the body (23). Similarly, the head crests of several chicken breeds, in which feathers are elongated but do not have a reversed growth trajectory as in pigeons, are localized to the top of the head probably due to ectopic expression of *Hox* positional cues (24). Together, these examples provide evidence for regionalization of the developing head and neck skin in the chicken. We propose that analogous mechanisms might underlie skin regionalization in the pigeon and allow *cr* to change feather polarity in the occiput and neck, but not elsewhere.

Our study of domestic rock pigeons illustrates how combining comparative genomics and population-based analyses forwards our understanding of genetic relationships and the genomic basis of traits. Many of the traits that vary among pigeon breeds also vary among wild species of birds and other animals (2, 25); thus, pigeons represent a model for identifying the genetic basis of variation in traits of general interest. Moreover, variation in many traits in domestic pigeons, including the head crest phenotype described here, is constructive rather than regressive: breeds derived from the ancestral rock pigeon possess traits that the ancestor does not have. While adaptive regressive traits are important, the genetic basis of constructive traits in vertebrates remains comparatively poorly understood. The domestic pigeon is thus a promising model to explore the genetic architecture of derived, constructive phenotypes in a bird that is amenable to genetic, genomic, and developmental investigation.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank the University of Washington Burke Museum for the *C. rupestris* tissue sample (UWBM 59803); J. Oldham, K. Wright, Utah Pigeon Club, National Pigeon Association, and A. and H.O. Christiansen for domestic pigeon samples; D. Clayton for feral samples; and D.C., M. Horvath, D. Kingsley, R. Nielsen, and W. Warren for discussion and comments. Supported by a Burroughs Wellcome Fund Career Award in the Biomedical Sciences, NSF CAREER DEB-1149160, University of Utah Research Foundation (M.D.S.); NIH training grant T32GM007464 (S.A.S.); NIH training grant T32HD07491 (E.T.D.); NSF EDEN internship (A.I.V.); NIH/NHGRI R01HG004694, NIH ARRA GO RC2HG005619 (M.Y.); Danish National Research Foundation (M.T.P.G., E.W.). We acknowledge a computer time allocation from the Center for High Performance Computing at the University of Utah.

## References and Notes


1. Driscoll CA, Macdonald DW, O'Brien SJ. From wild animals to domestic pets, an evolutionary view of domestication. *Proc Natl Acad Sci U S A*. 2009 Jun 16;106(Suppl 1):9971. [PubMed: 19528637]

2. Price TD. Domesticated birds as a model for the genetics of speciation by sexual selection. *Genetica*. 2002; 116:311. [PubMed: 12555787]
3. Darwin, C. On the Origin of Species by Means of Natural Selection. London: John Murray; 1859.
4. Darwin, CR. The Variation of Animals and Plants Under Domestication. Vol. vol. 1. London: John Murray; 1868. p. 412
5. Morgan TH. Notes on two crosses between different races of pigeons. *Biological Bulletin*. 1911; 21:215.
6. Sell, A. Breeding and Inheritance in Pigeons. Germany: Schober Verlags-GmbH, Hengersberg; 1994. p. 202
7. Supplementary Online Materials
8. Balakrishnan CN, Edwards SV. Nucleotide variation, linkage disequilibrium and founder-facilitated speciation in wild populations of the zebra finch (*Taeniopygia guttata*). *Genetics*. 2009 Feb.181:645. [PubMed: 19047416]
9. Ellegren H, et al. The genomic landscape of species divergence in *Ficedula* flycatchers. *Nature*. 2012 Nov 29.491:756. [PubMed: 23103876]
10. Aerts J, et al. Extent of linkage disequilibrium in chicken. *Cytogenet Genome Res*. 2007; 117:338. [PubMed: 17675876]
11. Johnson KP, et al. A molecular phylogeny of the dove genera *Streptopelia* and *Columba*. *The Auk*. 2001; 118:874.
12. Stringham SA, et al. Divergence, convergence, and the ancestry of feral populations in the domestic rock pigeon. *Curr Biol*. 2012
13. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in unrelated individuals. *Genome Res*. 2009 Sep.19:1655. [PubMed: 19648217]
14. Levi, WM. The Pigeon. Second Revised. Sumpter, SC., editor. Levi Publishing Co., Inc.; 1986. p. 667
15. Amundsen T. Why are female birds ornamented? *Trends Ecol Evol*. 2000 Apr.15:149. [PubMed: 10717684]
16. McKinnell IW, Makarenkova H, de Curtis I, Turmaine M, Patel K. EphA4, RhoB and the molecular development of feather buds are maintained by the integrity of the actin cytoskeleton. *Dev Biol*. 2004 Jun 1.270:94. [PubMed: 15136143]
17. Kelsh RN, Harris ML, Colanesi S, Erickson CA. Stripes and belly-spots - a review of pigment cell morphogenesis in vertebrates. *Semin Cell Dev Biol*. 2009 Feb.20:90. [PubMed: 18977309]
18. Yandell M, et al. A probabilistic disease-gene finder for personal genomes. *Genome Res*. 2011 Jun 23.
19. Elder ME, et al. Distinct T cell developmental consequences in humans and mice expressing identical mutations in the DLAARN motif of ZAP-70. *J Immunol*. 2001 Jan 1.166:656. [PubMed: 11123350]
20. Colosimo PF, et al. Widespread parallel evolution in sticklebacks by repeated fixation of Ectodysplasin alleles. *Science*. 2005 Mar 25.307:1928. [PubMed: 15790847]
21. Sutter NB, et al. A single IGF1 allele is a major determinant of small size in dogs. *Science*. 2007 Apr 6.316:112. [PubMed: 17412960]
22. Van Laere AS, et al. A regulatory mutation in IGF2 causes a major QTL effect on muscle growth in the pig. *Nature*. 2003 Oct 23.425:832. [PubMed: 14574411]
23. Mou C, et al. Cryptic patterning of avian skin confers a developmental facility for loss of neck feathering. *PLoS Biol*. 2011 Mar.9:e1001028. [PubMed: 21423653]
24. Wang Y, et al. The crest phenotype in chicken is associated with ectopic expression of HOXC8 in cranial skin. *PLoS One*. 2012; 7:e34012. [PubMed: 22514613]
25. Baptista LF, Gomez Martinez JE, Horblit HM. Darwin's pigeons and the evolution of the columbiforms: recapitulation of ancient genes. *Acta Zoológica Mexicana*. 2009 Oct 29.25:719.
26. Li R, et al. The sequence and de novo assembly of the giant panda genome. *Nature*. 2010 Jan 21.463:311. [PubMed: 20010809]
27. Li R, et al. SOAP2: an improved ultrafast tool for short read alignment. *Bioinformatics*. 2009; 25:1966. [PubMed: 19497933]




28. Benson G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Res.* 1999 Jan 15;27:573. [PubMed: 9862982]
29. Dalloul RA, et al. Multi-platform next-generation sequencing of the domestic turkey (*Meleagris gallopavo*): genome assembly and analysis. *PLoS Biol.* 2010; 8
30. Hillier L, Miller W, Birney E, Warren W, Hardison R. Sequence and comparative analysis of the chicken genome provide unique perspectives on vertebrate evolution. *Nature.* 2004 Dec 9;432:695. [PubMed: 15592404]
31. Warren WC, et al. The genome of a songbird. *Nature.* 2010 Apr 1;464:757. [PubMed: 20360741]
32. Birney E, Clamp M, Durbin R. GeneWise and Genomewise. *Genome Res.* 2004 May;14:988. [PubMed: 15123596]
33. Trapnell C, Pachter L, Salzberg SL. TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics.* 2009 May 1;25:1105. [PubMed: 19289445]
34. Trapnell C, et al. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nature biotechnology.* 2010 May; 28:511.
35. Stanke M, Waack S. Gene prediction with a hidden Markov model and a new intron submodel. *Bioinformatics.* 2003 Oct;19(Suppl 2):ii215. [PubMed: 14534192]
36. Burge C, Karlin S. Prediction of complete gene structures in human genomic DNA. *Journal of molecular biology.* 1997 Apr 25;268:78. [PubMed: 9149143]
37. Elsik CG, et al. Creating a honey bee consensus gene set. *Genome Biol.* 2007; 8:R13. [PubMed: 17241472]
38. Apweiler R, et al. UniProt: the Universal Protein knowledgebase. *Nucleic Acids Res.* 2004 Jan 1;32:D115. [PubMed: 14681372]
39. Apweiler R, et al. The InterPro database, an integrated documentation resource for protein families, domains and functional sites. *Nucleic Acids Res.* 2001 Jan 1;29:37. [PubMed: 11125043]
40. Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* 2000 Jan 1;28:27. [PubMed: 10592173]
41. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 1997 Mar 1;25:955. [PubMed: 9023104]
42. Nawrocki EP, Kolbe DL, Eddy SR. Infernal 1.0: inference of RNA alignments. *Bioinformatics.* 2009 May 15;25:1335. [PubMed: 19307242]
43. Griffiths-Jones S, Bateman A, Marshall M, Khanna A, Eddy SR. Rfam: an RNA family database. *Nucleic Acids Res.* 2003 Jan 1;31:439. [PubMed: 12520045]
44. Li H, et al. TreeFam: a curated database of phylogenetic trees of animal gene families. *Nucleic Acids Res.* 2006 Jan 1;34:D572. [PubMed: 16381935]
45. Huang DW, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* 2009 Jan;37:1. [PubMed: 19033363]
46. De Bie T, Cristianini N, Demuth JP, Hahn MW. CAFE: a computational tool for the study of gene family evolution. *Bioinformatics.* 2006 May 15;22:1269. [PubMed: 16543274]
47. Guindon S, Delsuc F, Dufayard JF, Gascuel O. Estimating maximum likelihood phylogenies with PhyML. *Methods in molecular biology.* 2009; 537:113. [PubMed: 19378142]
48. Morgulis A, et al. Database indexing for production MegaBLAST searches. *Bioinformatics.* 2008 Aug 15;24:1757. [PubMed: 18567917]
49. R Development Core Team. R Foundation for Statistical Computing. Vienna, Austria: 2008.
50. Paradis E, Claude J, Strimmer K. APE: Analyses of Phylogenetics and Evolution in R language. *Bioinformatics.* 2004 Jan 22;20:289. [PubMed: 14734327]
51. Purcell S, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet.* 2007 Sep;81:559. [PubMed: 17701901]
52. Rosenberg NA. DISTRUCT: a program for the graphical display of population structure. *Molecular Ecology Notes.* 2004; 4:137.
53. Rogers AR, Huff C. Linkage disequilibrium between loci with unknown phase. *Genetics.* 2009 Jul; 182:839. [PubMed: 19433632]

54. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *Journal of molecular biology*. 1990 Oct 5.215:403. [PubMed: 2231712]
55. Posada D, Crandall KA. MODELTEST: testing the model of DNA substitution. *Bioinformatics*. 1998; 14:817. [PubMed: 9918953]
56. Pacheco MA, et al. Evolution of modern birds revealed by mitogenomics: timing the radiation and origin of major orders. *Mol Biol Evol*. 2011 Jun.28:1927. [PubMed: 21242529]
57. Yang Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol*. 2007 Aug. 24:1586. [PubMed: 17483113]
58. Nam K, et al. Molecular evolution of genes in avian genomes. *Genome Biol*. 2010; 11:R68. [PubMed: 20573239]
59. Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD. Inferring the joint demographic history of multiple populations from multidimensional SNP frequency data. *PLoS Genet*. 2009 Oct.5:e1000695. [PubMed: 19851460]
60. Hudson RR. Generating samples under a Wright-Fisher neutral model of genetic variation. *Bioinformatics*. 2002 Feb.18:337. [PubMed: 11847089]
61. Weir BS, Cockerham CC. Estimating F-Statistics for the Analysis of Population Structure. *Evolution*. 1984; 38:1358.
62. Browning SR, Browning BL. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. *Am J Hum Genet*. 2007 Nov.81:1084. [PubMed: 17924348]
63. Clement M, Posada D, Crandall KA. TCS: a computer program to estimate gene genealogies. *Mol Ecol*. 2000 Oct.9:1657. [PubMed: 11050560]
64. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res*. 1994 Nov 11.22:4673. [PubMed: 7984417]
65. Abler LL, et al. A high throughput in situ hybridization method to characterize mRNA expression patterns in the fetal mouse lower urogenital tract. *JoVE*. 2011
66. Fazl A. The art of training pigeons in the East [annotated translation from Ain-i-Akbari, 1590]. *The Zoologist* (London). 1888; 12:167.
67. Tegetmeier, WB. Pigeons: Their Structure, Varieties, Habits, and Management. London: George Routledge and Sons; 1868. p. 178
68. National Pigeon Association. 2010 National Pigeon Association Book of Standards. Goodlettsville, TN: Purebred Pigeon Publishing; 2010.
69. Pelak K, et al. The characterization of twenty sequenced human genomes. *PLoS Genet*. 2010 Sep.6
70. Kent W. BLAT - the BLAST-like alignment tool. *Genome Res*. 2002; 12:656. [PubMed: 11932250]
71. Jones DT, Taylor WR, Thornton JM. The rapid generation of mutation data matrices from protein sequences. *CABIOS*. 1992 Jun.8:275. [PubMed: 1633570]
72. Tamura K, et al. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol*. 2011 Oct.28:2731. [PubMed: 21546353]




**Fig. 1.** Relationships among rock pigeons and the hill pigeon *Columba rupestris*. Consensus neighbor-joining tree based on 1.48 million genomic SNPs and 1000 bootstrap replicates (see Fig. S16 for bootstrap support). Branches are colored by traditional breed group (12) and/or geographic affinities: orange, toy breeds; brown, pouters and utility breeds; light blue, Indian and Iranian breeds; green, tumblers and highflyers; pink, homers and wattle breeds; red, Mediterranean and owl breeds; black, voice characteristics (14). Bold, red lettering indicates breeds with the head crest phenotype. Scale bar, Euclidean distance. Photo credits: T. Hellmann (domestic breeds) and M.V. Shreeram (*C. rupestris*).



**Fig. 2.**

*EphB2* is associated with the derived head crest phenotype. **(A)** Head crests are variable among breeds (left to right: Indian fantail, Old German owl, Old Dutch capuchin, Jacobin). **(B)**  $F_{ST}$  between crested and uncrested pigeons, maximum value for individual SNPs plotted for non-overlapping 100-kb windows across the genome. Red star, window with the highest score. Dashed red line, top 1% of scores. **(C)** Genome-wide VAAST scan. Each dot represents a single gene. Red star, gene with the highest score. Dashed red line, genome-wide significance cutoff. **(D)** Magnification of scaffold 612 in shaded region of **(B–C)**. Black trace, maximum  $F_{ST}$  between crested and uncrested birds over a 300-SNP window.

Red trace, unstandardized cross-population extended haplotype homozygosity (XP-EHH); higher values are evidence of selection (see Fig. S21, genome-wide plot). Dashed vertical line, position of the lone genome-wide significant VAAST hit. Green bar, 27.4-kb haplotype shared by all crested birds, includes only the *EphB2* gene. Blue bars, gene predictions on + and – DNA strands. (E) The *cr* mutation induces a charge-changing amino acid substitution; black bar, highly conserved DLAARN motif of catalytic loop. (F) Genotypes of 159 birds from 79 breeds at the *cr* locus are perfectly associated with the crest phenotype under a recessive model. (G) Network diagram of the minimal 11-kb haplotype shared by all resequenced rock pigeons with the *cr* mutation (also see Fig. S23). Many haplotypes contain the + allele (blue), but only one contains the *cr*SNP (red). Sizes of circles are proportional to the number of chromosomes containing a haplotype. Line segments represent single nucleotide differences. Jacobin photo credit: T. Hellmann.

**Fig. 3.**

Feather bud polarity is reversed in the *cr* mutant. (A,B) Expression of the feather structural gene *Ctnnb1* reveals the direction of outgrowth of early feather buds. (A) Neck and occipital head expression of *Ctnnb1* in an embryo of the uncrested racing homer. Feather buds point downward along the contour of the head and neck (arrowheads). (B) Occipital feathers buds point upward in the equivalent region of the crested English trumpeter, indicating morphological reversal of feather orientation. (C,D) Expression of the polarity marker *EphA4* was assayed at an earlier developmental stage to test if feather placodes, the ectodermal thickenings that give rise to feather buds, are also reversed. (C) Polarity marker *EphA4* is expressed posteriorly (arrowheads) in feather placodes of the racing homer. (D)

Polarity of placodes is reversed in the English trumpeter. Expression of *EphB2* in the skin is weak and unpolarized at this stage in both morphs (Fig. S26).