

GENUS GROUP OF FINITE GALOIS EXTENSIONS

TERUO TAKEUCHI

ABSTRACT. Let K/k be a Galois extension of finite degree, and let K' denote the maximal abelian extension over k contained in the Hilbert class field of K . We give formulas about the group structure of $\text{Gal}(K'/k)$ and the genus group of K/k , which refine the ordinary genus formula.

For an algebraic number field k of finite degree, let $\text{Cl}(k)$ denote the ideal class group of k . For a modulus S of k (i.e., a finite product of primes of k), let $I_k(S)$, $P_k(S)$, and P_{kS} denote the group of ideals of k prime to S , the group of principal ideals of k prime to S , and the ray ideal group modulo S in k , respectively. Similarly, let $k(S)$ and k_S denote the group of elements of k prime to S and the ray number group of k modulo S , respectively. Let K/k be a Galois extension of finite degree. Let K' be the maximal abelian extension over k contained in the Hilbert class field \bar{K} of K . Then, by definition, the genus field K^* of K/k in the wide sense is $K \cdot K'$, and the genus group of K/k is $\text{Gal}(K^*/K)$.

The following lemma is well known and proved by a standard manner in class field theory.

LEMMA 1. *Let \mathfrak{f}' be the conductor of K'/k ; then K' is a class field over k corresponding to $I_k(\mathfrak{f}')/N_{K/k}(P_K(\mathfrak{f}'))P_{k\mathfrak{f}'}$. Moreover, if K/k is abelian, then \mathfrak{f}' coincides with the conductor of K/k .*

For a modulus \mathfrak{f} with $\mathfrak{f}'|\mathfrak{f}$, put $\mathfrak{G}(K/k) = I_k(\mathfrak{f})/N_{K/k}(P_K(\mathfrak{f}))P_{k\mathfrak{f}}$. Clearly, $\mathfrak{G}(K/k)$ does not depend on the choice of such \mathfrak{f} (up to isomorphisms). The purpose of this paper is to describe the l^i -rank of $\mathfrak{G}(K/k)$. Let l be a prime number. Throughout this paper we fix l unless otherwise stated. For an abelian group A written multiplicatively, let $\text{rank}_l(A)$ denote the l^i -rank of A , i.e., the F_l -dimension of $A^{l^{i-1}}/A^{l^i}$. For $i \geq 0$, put $F_i = \{a \in k^\times \mid (a) \in P_k^{l^i}\}$. Then $k^\times \supset F_1 \supset F_2 \supset \cdots \supset F_i \supset \cdots \supset E_k$ and $F_i \supset F_{i-1}^l E_k$, where E_k denotes the group of units in k . Put $F_i(S) = F_i \cap k(S)$.

LEMMA 2. *Let $1 \rightarrow N \rightarrow M \rightarrow L \rightarrow 1$ ($N \subset M$) be an exact sequence of finite abelian groups. Put $N_i = N \cap M^{l^i}$. Then for $i \geq 1$, we have*

$$\begin{aligned}\text{rank}_l(M) &= \text{rank}_l(L) + \text{rank}_1(N_{i-1}/N_i) \\ &= \text{rank}_l(L) + \log_l \{ \#(N_{i-1}/N_i) \}.\end{aligned}$$

Received by the editors October 11, 1985.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 11R37, 11R65.

©1986 American Mathematical Society
0002-9939/86 \$1.00 + \$.25 per page

PROOF. By the definition of l^i -rank, we see $\text{rank}_i(L) = \text{rank}_1(L^{l^{i-1}}/L^l) = \text{rank}_1(M^{l^{i-1}}N/M^lN)$. On the other hand,

$$\begin{aligned}\#(M^{l^{i-1}}N/M^lN) &= \#(M^{l^{i-1}}/M^{l^{i-1}} \cap N)/\#(M^l/M^l \cap N) \\ &= \#(M^{l^{i-1}}/M^l)/\#(N_{i-1}/N_i).\end{aligned}$$

This proves the lemma.

PROPOSITION 1. Put $\mathcal{N}_i(\mathfrak{f}) = k(\mathfrak{f})^l N_{K/k}(K(\mathfrak{f}))k_{\mathfrak{f}}$. Then

$$\begin{aligned}\text{rank}_i(\mathfrak{G}(K/k)) &= \text{rank}_i(\text{Cl}(k)) + \text{rank}_i(k(\mathfrak{f})/N_{K/k}(K(\mathfrak{f}))k_{\mathfrak{f}}) \\ &\quad + \log_l \{ \#(F_{i-1}(\mathfrak{f})/F_{i-1}(\mathfrak{f}) \cap \mathcal{N}_{i-1}(\mathfrak{f}))/\#(F_i(\mathfrak{f})/F_i(\mathfrak{f}) \cap \mathcal{N}_i(\mathfrak{f})) \}.\end{aligned}$$

PROOF. We apply Lemma 2 to an exact sequence

$$\begin{aligned}1 \rightarrow P_k(\mathfrak{f})/N_{K/k}(P_K(\mathfrak{f}))P_{k\mathfrak{f}} &\rightarrow I_k(\mathfrak{f})/N_{K/k}(P_K(\mathfrak{f}))P_{k\mathfrak{f}} \\ &\rightarrow I_k(\mathfrak{f})/P_k(\mathfrak{f}) \rightarrow 1.\end{aligned}$$

Then with the notations in Lemma 2 we have

$$\begin{aligned}N_i &= (P_k(F) \cap I_k(\mathfrak{f})^l N_{K/k}(P_K(\mathfrak{f}))P_{k\mathfrak{f}})/N_{K/k}(P_K(\mathfrak{f}))P_{k\mathfrak{f}} \\ &\cong F_i(\mathfrak{f})N_{K/k}(K(\mathfrak{f}))k_{\mathfrak{f}}/E_k N_{K/k}(K(\mathfrak{f}))k_{\mathfrak{f}}.\end{aligned}$$

Hence

$$N_{i-1}/N_i \cong F_{i-1}(\mathfrak{f})N_{K/k}(K(\mathfrak{f}))k_{\mathfrak{f}}/F_i(\mathfrak{f})N_{K/k}(K(\mathfrak{f}))k_{\mathfrak{f}}.$$

So

$$\begin{aligned}\#(N_{i-1}/N_i) &= \left\{ \frac{\#(F_{i-1}(\mathfrak{f})N_{K/k}(K(\mathfrak{f}))k_{\mathfrak{f}}/\mathcal{N}_{i-1}(\mathfrak{f}))}{\#(F_i(\mathfrak{f})N_{K/k}(K(\mathfrak{f}))k_{\mathfrak{f}}/\mathcal{N}_i(\mathfrak{f}))} \right\} \#(\mathcal{N}_{i-1}(\mathfrak{f})/\mathcal{N}_i(\mathfrak{f})) \\ &= \left\{ \frac{\#(F_{i-1}(\mathfrak{f})/F_{i-1}(\mathfrak{f}) \cap \mathcal{N}_{i-1}(\mathfrak{f}))}{\#(F_i(\mathfrak{f})/F_i(\mathfrak{f}) \cap \mathcal{N}_i(\mathfrak{f}))} \right\} \#(\mathcal{N}_{i-1}(\mathfrak{f})/\mathcal{N}_i(\mathfrak{f})).\end{aligned}$$

Thus Lemma 2 implies the assertion.

Let v be a prime of k ramified in K , and let V be a prime divisor of v in \bar{K} . We also denote the restriction of V to an intermediate field of \bar{K}/k . Let \bar{K}_V , K_V , K'_V , and k_v denote the completion of \bar{K} , K , K' , and k by V , respectively. Further, let $(K_V)''$ be the maximal abelian subextension of \bar{K}_V/k_v , then $K'_V \subset (K_V)''$. Moreover, let $(K_V)_2$ be the maximal abelian subextension of K_V/k_v , and let \mathfrak{f}_V and T_V denote the conductor of $(K_V)_2/k_v$ and the inertia group of v in $(K_V)_2/k_v$, respectively. Since K/k is Galois, the conductor \mathfrak{f}_V and the group T_V do not depend on the choice of a prime divisor of v . Therefore, we write \mathfrak{f}_v and T_v instead of \mathfrak{f}_V and T_V . On the other hand, the conductor of $(K_V)''/k_v$ coincides with \mathfrak{f}_v since $N_{\bar{K}_V/k_v}(U(\bar{K}_V)) = N_{K_V/k_v}(U(K_V))$, where $U(K_V)$ denotes the group of units of K_V . Thus if we put $\mathfrak{f}^* = \prod_v \mathfrak{f}_v$, then $\mathfrak{f}' | \mathfrak{f}^*$, so we can apply the above results to \mathfrak{f}^* .

THEOREM. *Let the notation be as above. For $i \geq 1$, we have*

$$\begin{aligned} \text{rank}_i(\mathfrak{G}(K/k)) &= \text{rank}_i(\text{Cl}(k)) + \sum_v \text{rank}_i(T_v) \\ &\quad + \log_l \left\{ \frac{\#(F_{i-1}(\mathfrak{f}^*)/F_{i-1}(\mathfrak{f}^*) \cap \mathcal{N}_{i-1}(\mathfrak{f}^*))}{\#(F_i(\mathfrak{f}^*)/F_i(\mathfrak{f}^*) \cap \mathcal{N}_i(\mathfrak{f}^*))} \right\}. \end{aligned}$$

Moreover, if K/k is abelian, then \mathfrak{f}^* is the conductor of K/k and T_v is the inertia group of v in K/k .

PROOF. We apply Proposition 1 for $\mathfrak{f} = \mathfrak{f}^*$. Then it suffices to prove the assertion about the second term in the right-hand side of the above formula. Clearly,

$$k(\mathfrak{f})/N_{K/k}(K(\mathfrak{f}))k_{\mathfrak{f}} = \prod_v (k(\mathfrak{f}_v)/N_{K/k}(K(\mathfrak{f}_v))k_{\mathfrak{f}_v}) \quad (\text{direct})$$

holds. Further, $k_{\mathfrak{f}_v} \subset N_{K_v/k_v}(U(K_v))$ since \mathfrak{f}_v is the conductor of $(K_v)''/k_v$. So we have a natural homomorphism: $k(\mathfrak{f}_v)/N_{K/k}(K(\mathfrak{f}_v))k_{\mathfrak{f}_v} \rightarrow U(k_v)/N_{K_v/k_v}(U(K_v))$; but noting K/k is Galois, we can easily check that this gives an isomorphism. On the other hand, local class field theory states $U(k_v)/N_{K_v/k_v}(U(K_v)) \cong T_v$, which proves the theorem.

Here the last term of the above is rewritten as

$$\log_l \left\{ \frac{\left[\#(F_{i-1}(\mathfrak{f})/k(\mathfrak{f})^{\mu^{-1}}) / \#(F_{i-1}(\mathfrak{f}) \cap \mathcal{N}_{i-1}(\mathfrak{f})/k(\mathfrak{f})^{\mu^{-1}}) \right]}{\left[\#(F_i(\mathfrak{f})/k(\mathfrak{f})^{\mu}) / \#(F_i(\mathfrak{f}) \cap \mathcal{N}_i(\mathfrak{f})/k(\mathfrak{f})^{\mu}) \right]} \right\};$$

hence from [3, Lemma 1] we have the following:

COROLLARY.

$$\begin{aligned} \text{rank}_i(\mathfrak{G}(K/k)) &= \sum_v \text{rank}_i(T_v) - \text{rank}_i(E_k) \\ &\quad + \log_l \left\{ \frac{\#(F_i(\mathfrak{f}^*) \cap \mathcal{N}_i(\mathfrak{f}^*)/k(\mathfrak{f}^*)^{\mu})}{\#(F_{i-1}(\mathfrak{f}^*) \cap \mathcal{N}_{i-1}(\mathfrak{f}^*)/k(\mathfrak{f}^*)^{\mu^{-1}})} \right\}. \end{aligned}$$

REMARK 1. We know for sufficiently large i and j (independent of l), $F_i(\mathfrak{f}^*)/F_i(\mathfrak{f}^*) \cap \mathcal{N}_i(\mathfrak{f}^*) \cong E_k/E_k \cap \mathcal{N}_i(\mathfrak{f}^*)$ and $E_k \cap \mathcal{N}_i(\mathfrak{f}^*) = E_k \cap \mathcal{N}_j(\mathfrak{f}^*)$. Moreover, taking a product for all l , we have $E_k/E_k \cap N_{K/k}(K(\mathfrak{f}^*))k_{\mathfrak{f}^*} \cong \prod_l (E_k/E_k \cap \mathcal{N}^{(l)}(\mathfrak{f}^*))$, where $\mathcal{N}^{(l)}$ denotes \mathcal{N}_i corresponding to l . Thus multiplying the formulas for all l and i in the above theorem, we have

$$\#(\mathfrak{G}(K/k)) = h(k) \cdot \prod_v \#(T_v) / [E_k : E_k \cap N_{K/k}(K(\mathfrak{f}^*))k_{\mathfrak{f}^*}].$$

Let K_1 denote the maximal abelian subextension of K/k . Then $[K^* : K] = [K' : K_1] = \#(\mathfrak{G}(K/k))/[K_1 : k]$, so the above is nothing but the genus formula (e.g. see [1]). Thus the above theorem refines the genus formula.

REMARK 2. Now we consider an abelian case. Let T be a finite set of primes of k . Let M_i be the maximal abelian extension at most of index l^i in which only primes in T are ramified. For the conductor \mathfrak{f} of M_i , let M be the ray class field modulo \mathfrak{f} . Then M_i is the maximal subfield of M at most of index l^i . Hence $\log_l \#(\text{Gal}(M_i/k)) = \sum_{j=1}^i \text{rank}_j \mathfrak{G}(M/k)$. Put $h_i = \#\{x \in \text{Cl}(k) \mid x^{l^i} = 1\}$ and $t_i(v) = \#\{x \in T_v \mid x^{l^i} = 1\}$. Then the above theorem implies

$$\#(\text{Gal}(M_i/k)) = h_i(k) \prod_{v \in T} t_i(v) / [F_i(\mathfrak{f}) : F_i(\mathfrak{f}) \cap \mathcal{N}_i(\mathfrak{f})].$$

Since M is the ray class field, $\mathcal{N}_i(\mathfrak{f}) = k(\mathfrak{f})^{l^i} k_{\mathfrak{f}}$. Thus the above theorem gives a generalization of Kubota-Miki's formula [3, Theorem 1] (cf. [2]). Of course, in this case if $i = 1$, then the corollary to the above theorem is [4, Theorem 1].

Finally we study the genus group of K/k . In general, the genus group is not determined only by $\mathfrak{G}(K/k)$ and $\text{Gal}(K/k)$. Indeed, as is easily seen there are abelian fields K/Q and L/Q such that $\text{Gal}(K/Q) \cong \text{Gal}(L/Q)$ and $K^* = L^*$ although $\text{Gal}(K^*/K)$ is not isomorphic to $\text{Gal}(L^*/L)$. However, in the following case the genus group is completely determined by $\mathfrak{G}(K/k)$ and $\text{Gal}(K_1/k)$ since $\text{Gal}(K^*/K) \cong \text{Gal}(K'/K_1)$. The proof of Proposition 2 is easy, so we omit it.

PROPOSITION 2. *If there exists a finite set T of primes of k such that $\text{Gal}(K_1/k) = \prod_{v \in T} T_v$ (direct product), then*

$$\mathfrak{G}(K/k) \cong \text{Gal}(K_1/k) \oplus \text{Gal}(K'/K_1).$$

REMARK 3. In this paper we deal with the wide sense, but a similar argument holds in the narrow sense with a few changes of parts about infinite primes.

REFERENCES

1. Y. Furuta, *The genus field and genus number in algebraic number fields*, Nagoya Math. J. **29** (1967), 281–285.
2. T. Kubota, *Galois group of the maximal abelian extension over an algebraic number field*, Nagoya Math. J. **12** (1957), 177–189.
3. H. Miki, *On the maximal abelian l -extension of a finite algebraic number field with given ramification*, Nagoya Math. J. **70** (1978), 183–202.
4. I. R. Šfarevič, *Extensions with given points of ramification*, Inst. Hautes Études Sci. Publ. Math. **18** (1963), 71–95; Amer. Math. Soc. Transl. **59** (1966), 128–149.

DEPARTMENT OF MATHEMATICS, FACULTY OF GENERAL EDUCATION, NIIGATA UNIVERSITY, NIIGATA 950-21, JAPAN