



Contents lists available at ScienceDirect

Proceedings of the Geologists' Association

journal homepage: www.elsevier.com/locate/pgeola

Geoarchaeology of Pleistocene open-air sites in the Vila Nova da Barquinha-Santa Cita area (Lower Tejo River basin, central Portugal)

António A. Martins^{a,*}, Pedro P. Cunha^b, Pierluigi Rosina^c, Luiz Osterbeek^c, Sara Cura^d, Stefano Grimaldi^e, José Gomes^f, Jan-Pieter Buylaert^g, Andrew S. Murray^g, João Matos^h

^a Centro de Geofísica, Dep. Geociências University Évora, Portugal

^b Dep. of Earth Sciences, Marine and Environmental Research Centre, Univ. Coimbra, Portugal

^c Instituto Politécnico de Tomar, Centro de Geociências Univ. Coimbra, Portugal

^d Museu de Arte Pré-Histórica de Mação, Centro de Geociências Univ. Coimbra, Portugal

^e Università degli Studi di Trento, Centro de Geociências Univ. Coimbra, Portugal

^f Centro de Interpretação de Arqueologia do Alto Ribatejo, Centro de Geociências Univ. Coimbra, Portugal

^g Nordic Laboratory for Luminescence Dating, Aarhus Univ., Risø DTU, Denmark

^h Bolseiro BIC (proj. POCI/CTE-GEX/58120/2004), Dep. Geociências Univ. Évora, Portugal

ARTICLE INFO

Article history:

Received 14 April 2009

Received in revised form 25 August 2009

Accepted 21 January 2010

Available online 19 February 2010

Keywords:

Palaeolithic

Pleistocene

Luminescence dating

Terraces

Tejo River

Iberia

ABSTRACT

This paper aims to provide insight into human occupation and landscape change during the Pleistocene in a central area of the Lower Tejo basin (Portugal). Detailed geomorphological mapping, coupled with lithostratigraphy, sedimentology and luminescence dating, supports the identification of a complete terrace staircase sequence. It consists of six gravely terraces located below the culminant (Pliocene) basin unit. A chronological framework for the sedimentary sequences and associated human industries is proposed and correlated with marine oxygen isotope stages (MIS): T1 terrace, not dated; T2, not dated; T3, >300 ka; T4, ~300–160 ka (MIS8, MIS7 and MIS6); T5, ~136–75 ka (MIS5); T6, ~62–30 ka (MIS3); colluvium and aeolian sands, ~30–14 ka (MIS2); valley fill deposits, ~14 ka to present (MIS1). The oldest artefacts were found at the base of the T4 terrace, with the local stratigraphic level dated to $\geq 175 \pm 6$ ka (Middle Pleistocene). The lithic assemblages collected from distinct stratigraphic levels (T4, T5 top, T6 terraces and colluvium) are characterized by the predominance of opportunistic technological choices, a feature that can be attributed partly to the preferential exploitation of the available raw material, dominated by local-sourced quartzites and quartz pebbles. The adaptation to local raw material (texture and volume), together with subsistence patterns and behaviours, could explain the rarity of Acheulian types (handaxes and cleavers) and picks in the T4 terraces of the Tejo tributaries; this is in contrast to the same terrace of the Tejo valley, in which these types are found. Interpretation of the environmental conditions (controlled by climate and glacio-eustatic sea-level changes) affecting the hunter-gatherer human groups is also presented.

© 2010 The Geologists' Association. Published by Elsevier Ltd. All rights reserved.

1. Introduction

The Iberian Peninsula is influenced by both the Atlantic and Mediterranean domains. Pleistocene fluvial terrace sequences are well developed in most of the major river systems in Iberia, with many of them driven by climate changes, although tectonic and sea-level effects can also be observed (Bridgland and Westaway, 2007; Santisteban and Schulte, 2007).

The ~1100 km long Tejo River originates in central-eastern Spain at an altitude of 1839 m and has a catchment area of

81,947 km² with a present average flow of ~17,253 Hm³ yr⁻¹ (Benito et al., 2003). In the Portuguese sector the catchment comprises ~25,000 km² and the river is 230 km long, with a mean gradient of 0.03%. Today the river is characterized by extreme seasonal and annual flow variability, with peak discharges more than 30 times the average discharge (Benito et al., 2003; Bettencourt and Ramos, 2003). The Tejo drains two Cenozoic basins (the Madrid and the Lower Tejo basins, respectively) towards the west and southwest. Its mouth, at the Atlantic Ocean, is located near Lisboa (central western Portugal). The Lower Tejo valley provides a long record (ca. 3.4 Ma) of alluvial and cultural history, from which it is possible to interpret landscape development, environmental changes and human occupation, all influenced by base-level and climate changes (Cunha et al., 2005, 2008).

In the Lower Tejo basin, the aggradational terraces consist mainly of coarse-grained siliciclastic gravels and sands (Cunha

* Corresponding author at: Departamento de Geociências da Univ. de Évora, Rua Romão Ramalho, 59, 7000-671, Évora, Portugal. Tel.: +351 266745301; fax: +351 266745397.

E-mail address: aam@uevora.pt (A.A. Martins).