

## GEOMETRY OF COMPLEX MANIFOLDS SIMILAR TO THE CALABI-ECKMANN MANIFOLDS

DAVID E. BLAIR, GERALD D. LUDDEN & KENTARO YANO

In [4] Calabi and Eckmann showed that the product of two odd-dimensional spheres  $S^{2p+1} \times S^{2q+1}$  ( $p, q \geq 1$ ) is a complex manifold. As  $S^{2p+1} \times S^{2q+1}$  is not Kaehlerian, the fundamental 2-form  $\Omega$  of the Hermitian structure is not closed. However,  $d\Omega$  does have a special form on  $S^{2p+1} \times S^{2q+1}$ ; in fact,  $S^{2p+1} \times S^{2q+1}$  admits two nonvanishing vector fields which are both Killing and analytic, and whose covariant forms determine  $\Omega$ . Our purpose here is to study complex manifolds whose complex structures are similar to the complex structure on  $S^{2p+1} \times S^{2q+1}$ .

In § 1 we review the geometry of the Calabi-Eckmann manifolds. In § 2 we give some elementary properties of vector fields on a Hermitian manifold, and introduce the notion of a holomorphic pair of automorphisms and of a bicontact manifold. § 3 continues the author's paper [2] on the differential geometry of principal toroidal bundles for the present case. In § 4 we discuss bicontact manifolds and, in particular, the integrable distributions of a bicontact structure on a Hermitian manifold. Finally in § 5 we give some results on the curvatures of a Hermitian manifold admitting a holomorphic pair of automorphisms.

### 1. The Hermitian structure on the Calabi-Eckmann manifolds

The construction of the complex structure on  $S^{2p+1} \times S^{2q+1}$  which we will give is due to Morimoto [6]. It is well known that an odd-dimensional sphere  $S^{2p+1}$  carries a contact structure, i.e., a nonvanishing 1-form  $\eta$  such that  $\eta \wedge (d\eta)^p \neq 0$ . Let  $G$  be the standard metric on  $S^{2p+1}$ . Then there exist on  $S^{2p+1}$  (see e.g. [8]) a contact form  $\eta$ , a vector field  $\xi$ , and a tensor field  $\varphi$  of type (1, 1) such that

$$\begin{aligned}\eta(\xi) &= 1, \quad \varphi\xi = 0, \quad \eta \circ \varphi = 0, \quad \varphi^2 = -I + \eta \otimes \xi, \\ G(\xi, X) &= \eta(X), \quad G(\varphi X, \varphi Y) = G(X, Y) - \eta(X)\eta(Y),\end{aligned}$$

i.e.,  $S^{2p+1}$  carries an almost contact metric structure. For a contact structure  $\eta \wedge (d\eta)^p \neq 0$ ,  $\varphi$ ,  $\xi$  and  $G$  may be chosen such that  $d\eta(X, Y) = G(\varphi X, Y)$ ,

as happens in the sphere example. Moreover, the contact metric structure on  $S^{2p+1}$  is normal, i.e.,

$$[\varphi, \varphi] + d\eta \otimes \xi = 0,$$

where  $[\varphi, \varphi]$  is the Nijenhuis torsion of  $\varphi$ . Thus  $S^{2p+1}$  carries a normal contact metric or *Sasakian* structure.

Now let  $(\varphi, \xi, \eta, G)$  and  $(\bar{\varphi}, \bar{\xi}, \bar{\eta}, \bar{G})$  be Sasakian structures on  $S^{2p+1}$  and  $S^{2q+1}$  respectively. Then define an almost complex structure  $J$  on  $S^{2p+1} \times S^{2q+1}$  by

$$J(X, \bar{X}) = (\varphi X - \bar{\eta}(\bar{X})\xi, \bar{\varphi}\bar{X} + \eta(X)\bar{\xi}),$$

and let  $g$  be the product metric. Then direct computations show [6] that  $J^2 = -I$ ,  $g(J(X, \bar{X}), J(Y, \bar{Y})) = g((X, \bar{X}), (Y, \bar{Y}))$  and, using normality, that  $[J, J] = 0$ . Thus  $S^{2p+1} \times S^{2q+1}$  is a Hermitian manifold.

Defining the fundamental 2-form  $\Omega$  of the Hermitian structure by

$$\Omega((X, \bar{X}), (Y, \bar{Y})) = g(J(X, \bar{X}), (Y, \bar{Y})),$$

we find that

$$\Omega = d\eta + d\bar{\eta} + \eta \wedge \bar{\eta},$$

where we view  $\eta$  and  $\bar{\eta}$  as 1-forms extended to the product. Thus the fundamental 2-form  $\Omega$  of the Hermitian structure on  $S^{2p+1} \times S^{2q+1}$  satisfies

$$d\Omega = d\eta \wedge \bar{\eta} - \eta \wedge d\bar{\eta}.$$

Finally we remark that from the Hopf fibration  $\pi': S^{2p+1} \rightarrow PC^p$  of an odd-dimensional sphere as a principal circle bundle over complex projective space, we obtain a natural fibration  $\pi: S^{2p+1} \times S^{2q+1} \rightarrow PC^p \times PC^q$  of a Calabi-Eckmann manifold as a principal  $T^2$  (2-dimensional torus) bundle over a product of complex projective spaces. In fact the complex coordinates of  $S^{2p+1} \times S^{2q+1}$  are essentially those of  $PC^p \times PC^q$  together with a fibre coordinate [4], [5].

## 2. Some elementary properties of vector fields on a Hermitian manifold

Let  $M^{2n}$  be a Hermitian manifold with complex structure  $J$  and Hermitian metric  $g$ . Let  $U$  be an analytic vector field<sup>1</sup> on  $M^{2n}$ , i.e.,  $\mathfrak{L}_U J = 0$  where  $\mathfrak{L}$  denotes Lie differentiation.

<sup>1</sup> More generally on an almost complex manifold a vector field  $U$  is said to be almost analytic if  $\mathfrak{L}_U J = 0$  and  $[J, J](U, X) = 0$  for all vector fields  $X$ .

**Proposition 2.1.** *If  $U$  is an analytic vector field on  $M^{2n}$ , then so is  $V = JU$ .  
*Proof.**

$$\begin{aligned} 0 &= [J, J](U, X) = -[U, X] + [V, JX] - J[V, X] - J[U, JX] \\ &= -J(\mathfrak{L}_V J)X + (\mathfrak{L}_V J)X = (\mathfrak{L}_V J)X . \end{aligned}$$

Thus, if  $U$  is an infinitesimal automorphism of  $J$ , so is  $JU$ ; but if  $U$  is Killing (an automorphism of  $g$ ),  $JU$  is not in general Killing. We therefore make the following definition.

**Definition.** By a holomorphic pair of automorphisms we mean a unit vector field  $U$  such that  $U$  and  $V = JU$  are infinitesimal automorphisms of the Hermitian structure.

Let  $u$  and  $v$  denote the covariant forms of  $U$  and  $V$  respectively. We begin with some elementary properties of a holomorphic pair of automorphisms ( $U, V = JU$ ).

**Lemma 2.2.**  $[U, V] = 0$ .

*Proof.*  $0 = (\mathfrak{L}_V J)U = [U, JU] - J[U, U] = [U, V]$ .

**Lemma 2.3.**  $du(U, X) = 0, du(V, X) = 0, dv(U, X) = 0, dv(V, X) = 0$ .

*Proof.* We give the proof for  $du$ , the proof for  $dv$  being similar. Since  $U$  is Killing and unit, we have

$$\begin{aligned} du(U, X) &= (\nabla_U u)(X) - (\nabla_X u)(U) = g(\nabla_U U, X) - g(\nabla_X U, U) \\ &= -2g(\nabla_X U, U) = 0 , \end{aligned}$$

where  $\nabla$  denotes the Riemannian connection of  $g$ . Similarly since  $[U, V] = 0$  and  $V$  is also Killing, we have

$$du(V, X) = g(\nabla_V U, X) - g(\nabla_X U, V) = g(\nabla_V V, X) + g(\nabla_X V, U) = 0 .$$

**Proposition 2.4.** *At each point of  $M^{2n}$ ,  $u$  and  $v$  have odd rank, i.e., there exist nonnegative integers  $p$  and  $q$  such that  $u \wedge (du)^p \neq 0, v \wedge (dv)^q \neq 0, (du)^{p+1} = 0, (dv)^{q+1} = 0$ .*

*Proof.* First note that  $(du)^n = 0$ ; for evaluating  $(du)^n$  on a  $J$ -basis containing  $U$  and  $V$  each term in

$$(du)^n(U, V, X_3, \dots, X_{2n})$$

vanishes by Lemma 2.3; here we have set  $X_1 = U, X_2 = JU = V$  and  $\{X_i\}$  a  $J$ -basis. Suppose now that at  $m \in M^{2n}$ ,  $(du)^p \neq 0$  and  $(du)^{p+1} = 0$ . Then evaluating  $(u \wedge (du)^p)(U, Y_1, \dots, Y_{2p})$  where  $Y_1, \dots, Y_{2p}$  are vector fields such that  $du(Y_i, Y_j) \neq 0$ , we have that  $u \wedge (du)^p \neq 0$ . Similarly  $v$  has rank  $2q + 1$ .

**Definition.** We say that a differentiable manifold  $M^{2n}$  is bicontact if it admits 1-forms  $u$  and  $v$  such that  $u \wedge v \wedge (du)^p \wedge (dv)^q \neq 0, (du)^{p+1} = 0$

and  $(dv)^{q+1} = 0$  with  $p + q + 1 = n$ .  $M^{2n}$  is called a Hermitian bicontact manifold if  $M^{2n}$  is both Hermitian and bicontact, and the 1-forms  $u$  and  $v$  are the covariant forms of a holomorphic pair of automorphisms.

**Lemma 2.5.** *If  $du$  is of bidegree  $(1, 1)$  with respect to the complex structure  $J$ , then so is  $dv$ .*

*Proof.* Recall that the Nijenhuis torsion of a vector-valued 1-form  $h$  is given by its action on a 1-form  $\theta$ . This action is

$$[h, h]\theta = -h^{(2)}d\theta + h^{(1)}d(\theta \circ h) - d(\theta \circ h^2) ,$$

where for a 2-form  $\theta$ ,

$$(h^{(1)}\theta)(X, Y) = \theta(hX, Y) + \theta(X, hY) , \quad (h^{(2)}\theta)(X, Y) = \theta(hX, hY) .$$

$h^{(1)}\theta$  is often denoted by  $\theta \frown h$ . Now since  $v = -u \circ J$  and  $du$  is of bidegree  $(1, 1)$ , we have

$$\begin{aligned} 0 &= ([J, J]u)(X, Y) \\ &= -du(JX, JY) - dv(JX, Y) - dv(X, JY) + du(X, Y) \\ &= -dv(JX, Y) - dv(X, JY) , \end{aligned}$$

and hence  $dv$  is of bidegree  $(1, 1)$ .

**Remark.** The above proof also shows that if  $du = dv$ , then  $[J, J] = 0$  implies that  $du(=dv)$  is of bidegree  $(1, 1)$ . The authors have studied certain manifolds admitting independent 1-forms  $u$  and  $v$  with  $du = dv$ , [1], [2].

**Proposition 2.6.** *If  $M^{2n}$  is Kaehlerian, then  $du = dv = 0$ .*

*Proof.* First since  $V$  is analytic, we have

$$0 = (\mathfrak{L}_V J)X = \nabla_V JX - \nabla_{JX} V - J\nabla_V X + J\nabla_X V = -\nabla_{JX} V + J\nabla_X V .$$

Now since  $V$  is Killing,

$$\begin{aligned} du(X, Y) &= g(\nabla_X U, Y) - g(\nabla_Y U, X) = g(-\nabla_X J V, Y) - g(-\nabla_Y J V, X) \\ &= g(\nabla_X V, JY) + g(J\nabla_Y V, X) = -g(\nabla_{JY} V, X) + g(J\nabla_Y V, X) = 0 . \end{aligned}$$

Similarly one can show that  $dv = 0$ .

In [9] one of the authors introduced the notion of an  $f$ -structure on a differentiable manifold, i.e., the manifold admits a tensor field  $f \neq 0$  of type  $(1, 1)$  satisfying  $f^3 + f = 0$  (see also [1], [7]).

**Proposition 2.7.** *Let  $(M^{2n}, J, g)$  be an almost Hermitian manifold admitting a nonvanishing vector field  $U$ , then  $U, V = JU, u, v$  (the covariant forms of  $U$  and  $V$ ) and  $f = J + v \otimes U - u \otimes V$  define an  $f$ -structure with complemented frames and rank  $(f) = 2n - 2$  on  $M^{2n}$ , i.e., we have*

$$f^2 = -I + u \otimes U + v \otimes V, \quad fU = fV = 0, \quad u \circ f = v \circ f = 0, \\ u(U) = v(V) = 1, \quad u(V) = v(U) = 0.$$

The proof of this proposition is a straightforward computation and will be omitted.

An  $f$ -structure with complemented frames  $(f, U, V, u, v)$  is said to be *normal* if the tensor  $S$  defined by

$$S(X, Y) = [f, f](X, Y) + du(X, Y)U + dv(X, Y)V$$

vanishes. Computing  $S$  in our case gives

$$S(X, Y) = [J, J](X, Y) - (du \wedge J)(X, Y) - (dv \wedge J)(X, Y) \\ + u(X)(\mathfrak{L}_V J)Y - u(Y)(\mathfrak{L}_V J)X + v(X)(\mathfrak{L}_U J)Y - v(Y)(\mathfrak{L}_U J)X.$$

Thus we have the following result.

**Proposition 2.8.** *On a Hermitian manifold with a nonvanishing analytic vector field  $U$ , if  $du$  is of bidegree  $(1, 1)$ , then the  $f$ -structure  $(f, U, V, u, v)$  is normal.*

It is well known (see e.g. [7]) that for a normal  $f$ -structure with complemented frames, we have

$$\mathfrak{L}_U f = 0, \quad \mathfrak{L}_U u = 0, \quad \mathfrak{L}_U v = 0, \quad \mathfrak{L}_V f = 0, \quad \mathfrak{L}_V u = 0, \quad \mathfrak{L}_V v = 0, \\ du \wedge f = 0, \quad dv \wedge f = 0, \quad [U, V] = 0.$$

Thus a straightforward computation shows that  $S = 0$  implies  $[J, J] = 0$ .

Now if  $g$  is the Hermitian metric on  $M^{2n}$ , then

$$g(fX, fY) = g(X, Y) - u(X)u(Y) - v(X)v(Y), \\ u(X) = g(U, X), \quad v(X) = g(V, X),$$

that is,  $(f, g, u, v)$  defines a metric  $f$ -structure with complemented frames.

Finally we define the fundamental 2-forms  $\Omega$  and  $F$  of the structures by

$$\Omega(X, Y) = g(JX, Y), \quad F(X, Y) = g(fX, Y).$$

Then a short computation gives

$$F = \Omega - u \wedge v.$$

### 3. Fiberings by a holomorphic pair of automorphisms

In [2] the authors proved the following result.

**Theorem.** *Let  $M^{2m+s}$  be a compact connected manifold with a regular normal  $f$ -structure of rank  $2m$ . Then  $M^{2m+s}$  is the bundle space of a principal toroidal bundle over a complex manifold  $N^{2m}$ .*

Now if a complex manifold  $M^{2n}$  admits a regular analytic vector field  $U$  (i.e., every point  $m \in M^{2n}$  has a neighborhood such that the integral curve of  $U$  through  $m$  passes through the neighborhood only once), the vector field  $V = JU$  is not necessarily regular. Thus we say that a holomorphic pair of automorphisms is regular if both  $U$  and  $V$  are regular vector fields. Then using the above theorem and Proposition 2.8 we can prove the following result.

**Theorem 3.1.** *If a compact Hermitian manifold  $(M^{2n}, J, g)$  admits a regular holomorphic pair of automorphisms  $(U, V = JU)$  with  $du$  of bidegree  $(1, 1)$ , then  $M^{2n}$  is a principal  $T^2$  bundle over a Hermitian manifold  $N^{2n-2}$ .*

*Proof.* From the above theorem and Proposition 2.8 we obtain the desired fibration. Thus we shall only exhibit the Hermitian structure on  $N^{2n-2}$ . As  $U$  and  $V$  are analytic,  $J$  is projectable and we define  $J'$  on  $N^{2n-2}$  by

$$J'X = \pi_* J \tilde{\pi} X,$$

where  $\tilde{\pi}$  denotes the horizontal lift with respect to the Riemannian connection of  $g$  (in the nonmetric case one can use the pair  $(u, v)$  as a Lie algebra valued connection form to determine  $\tilde{\pi}$  [2]). Then it is easy to check that  $J'^2 = -I$ . Moreover we have

$$\begin{aligned} [J', J'](X, Y) &= -[\pi_* \tilde{\pi} X, \pi_* \tilde{\pi} Y] + [\pi_* J \tilde{\pi} X, \pi_* J \tilde{\pi} Y] \\ &\quad - \pi_* J \tilde{\pi} [\pi_* J \tilde{\pi} X, \pi_* \tilde{\pi} Y] - \pi_* J \tilde{\pi} [\pi_* \tilde{\pi} X, \pi_* J \tilde{\pi} Y] \\ &= \pi_* [J, J](\tilde{\pi} X, \tilde{\pi} Y) = 0. \end{aligned}$$

Finally as  $U$  and  $V$  are Killing, the metric  $g$  is projectable to a metric  $g'$  on  $N^{2n-2}$  given by  $g'(X, Y) \circ \pi = g(\tilde{\pi} X, \tilde{\pi} Y)$ . Then

$$g'(J'X, J'Y) \circ \pi = g(J \tilde{\pi} X, J \tilde{\pi} Y) = g(\tilde{\pi} X, \tilde{\pi} Y) = g'(X, Y) \circ \pi,$$

and hence the structure on  $N^{2n-2}$  is Hermitian.

We now compute the fundamental 2-form  $F$  of the  $f$ -structure  $(f, U, V, u, v)$  on  $M^{2n}$ . First of all it is clear that  $F(U, X) = 0$  and  $F(V, X) = 0$ . Thus it is enough to compute  $F$  on vector fields of the form  $\tilde{\pi} X, \tilde{\pi} Y$  where  $X$  and  $Y$  are vector fields on  $N^{2n-2}$ .

$$\begin{aligned} F(\tilde{\pi} X, \tilde{\pi} Y) &= g(f \tilde{\pi} X, \tilde{\pi} Y) = g(J \tilde{\pi} X, \tilde{\pi} Y) = g(\tilde{\pi} J' X, \tilde{\pi} Y) \\ &= g'(J' X, Y) \circ \pi = \mathcal{O}'(X, Y) \circ \pi, \end{aligned}$$

where  $\mathcal{O}'$  is the fundamental 2-form on  $N^{2n-2}$ . Hence we have  $F = \pi^* \mathcal{O}'$ . Now  $dF = d\pi^* \mathcal{O}' = \pi^* d\mathcal{O}'$  and  $dF = d\mathcal{O} - du \wedge v + u \wedge dv$ , from which we get the following result.

**Theorem 3.2.** *The base manifold  $(N^{2n-2}, J', g')$  of the above fibration is Kaehlerian if and only if*

$$d\Omega = du \wedge v - u \wedge dv$$

on  $M^{2n}$ .

Note also that by Proposition 2.6,  $d\Omega = 0$  implies  $du = dv = 0$  and hence  $dF = 0$ . Thus we have the following result.

**Proposition 3.3.** *If  $M^{2n}$  is Kaehlerian, then the base manifold  $N^{2n-2}$  is also Kaehlerian.*

#### 4. Hermitian bicontact manifolds

We begin with the following elementary result on the topology of a compact bicontact manifold.

**Theorem 4.1.** *Let  $M^{2n}$  be a compact bicontact manifold, and let  $2p + 1$  and  $2q + 1$  denote the ranks of the bicontact forms  $u$  and  $v$ . Then the betti numbers  $b_{2p+1}$  and  $b_{2q+1}$  are nonzero.*

*Proof.* As  $(2p + 1) + (2q + 1) = 2n$  it suffices to show that  $b_{2p+1}$  is nonzero. We shall show that  $u \wedge (du)^p$  has nonzero harmonic part. Suppose  $u \wedge (du)^p$  has no harmonic part, then as  $(du)^{p+1} = 0$ ,  $u \wedge (du)^p$  is exact, say  $d\alpha$ . Now on a bicontact manifold  $u \wedge (du)^p \wedge v \wedge (dv)^q$  is a volume element, hence, since  $(dv)^{q+1} = 0$ , we have

$$0 \neq \int_M u \wedge (du)^p \wedge v \wedge (dv)^q = \int_M d\alpha \wedge v \wedge (dv)^q = \int_M d(\alpha \wedge v \wedge (dv)^q) = 0,$$

a contradiction.

We shall now digress briefly to introduce the notion of a semi-invariant submanifold [3]. Let  $M^{2n}$  be an almost complex manifold with a vector field  $U$  and a 1-form  $u$  with  $u(U) = 1$ , and set  $V = JU$ ,  $v = -u \circ J$ . Let  $\bar{M} \rightarrow M^{2n}$  be a submanifold of  $M^{2n}$  such that 1) the transform of a vector tangent to  $\bar{M}$  by  $J$  is in the space spanned by the tangent space of  $\bar{M}$  and the vector  $U$ , 2)  $V$  is tangent to  $\bar{M}$ , and 3)  $u \circ \iota_* = 0$ ; we then say that  $\bar{M}$  is *semi-invariant with respect to  $U$* . Note that  $U$  is never tangent to  $\bar{M}$ , for if it were, then  $U = \iota_* \bar{U}$ , and  $1 = u(U) = u(\iota_* \bar{U}) = 0$ , a contradiction.

Now define a tensor field  $\varphi$  of type  $(1, 1)$ , a vector field  $\xi$ , and a 1-form  $\eta$  on  $\bar{M}$  by

$$J\iota_*X = \iota_*\varphi X - \eta(X)U, \quad V = \iota_*\xi.$$

We then have

$$-\iota_*X = \iota_*\varphi^2X - \eta(\varphi X)U - \eta(X)\iota_*\xi,$$

from which it follows that

$$\varphi^2 = -I + \eta \otimes \xi, \quad \eta \circ \varphi = 0.$$

Also

$$-U = JV = J\iota_*\xi = \iota_*\varphi\xi - \eta(\xi)U ,$$

giving

$$\varphi\xi = 0 , \quad \eta(\xi) = 1 .$$

Thus we have the following result.

**Proposition 4.2.** *A submanifold of  $M^{2n}$ , which is semi-invariant with respect to  $U$ , admits an almost contact structure.*

Now computing  $[J, J](\iota_*X, \iota_*Y)$  we have

$$\begin{aligned} [J, J](\iota_*X, \iota_*Y) &= \iota_*[\varphi, \varphi](X, Y) + d\eta(X, Y)\iota_*\xi - \eta(X)(\mathfrak{L}_U J)\iota_*Y \\ &\quad + \eta(Y)(\mathfrak{L}_U J)\iota_*X - ((\mathfrak{L}_{\varphi X}\eta)(Y) - (\mathfrak{L}_{\varphi Y}\eta)(X))U , \end{aligned}$$

from which we obtain the following result.

**Proposition 4.3.** *If a submanifold is semi-invariant with respect to an analytic vector field  $U$  on a complex manifold  $M^{2n}$ , then its almost contact structure is normal.*

Returning to the bicontact case, we assume for the remainder of this section that  $M^{2n}$  is a Hermitian bicontact manifold as defined in § 2. We define a distribution  $\mathcal{U}$  of dimension  $2q + 1$  by

$$\mathcal{U} = \{X \mid i(X)u = 0, i(X)du = 0\} ,$$

where  $i$  denotes the interior product operator. We shall show that  $\mathcal{U}$  is integrable. Let  $X$  and  $Y$  be vector fields belonging to  $\mathcal{U}$ . Then

$$0 = du(X, Y) = Xu(Y) - Yu(X) - u([X, Y]) = -u([X, Y]) .$$

Also for any  $Z$

$$0 = du(X, Z) = Xu(Z) - u([X, Z]) = (\mathfrak{L}_X u)(Z) ,$$

and therefore

$$\begin{aligned} du([X, Y], Z) &= [X, Y]u(Z) - Zu([X, Y]) - u([[X, Y], Z]) \\ &= (\mathfrak{L}_{[X, Y]}u)(Z) = ((\mathfrak{L}_X\mathfrak{L}_Y - \mathfrak{L}_Y\mathfrak{L}_X)u)(Z) = 0 . \end{aligned}$$

Similarly the complementary distribution  $\mathcal{V} = \{X \mid i(X)v = 0, i(X)dv = 0\}$  of dimension  $2p + 1$  is integrable.

**Theorem 4.4.** *A Hermitian bicontact manifold  $M^{2n}$  with  $du$  of bidegree  $(1, 1)$  is locally the product of two normal contact manifolds  $M^{2p+1}$  and  $M^{2q+1}$ .*

*Proof.* As noted above the distributions  $\mathcal{U}$  and  $\mathcal{V}$  are complementary and integrable. Thus  $M^{2n}$  is locally the product of the respective maximal integral

submanifolds  $M^{2q+1}$  and  $M^{2p+1}$ . We shall show that the integral submanifold  $M^{2q+1}$  of  $\mathcal{U}$  is semi-invariant with respect to  $U$ . Let  $\iota: M^{2q+1} \rightarrow M^{2n}$  denote the immersion, and let  $X$  be tangent to  $M^{2q+1}$ , i.e.,  $\iota_*X \in \mathcal{U}$ . Set  $Y = J\iota_*X + v(\iota_*X)U$ . Then

$$u(Y) = u(J\iota_*X) + v(\iota_*X) = -v(\iota_*X) + v(\iota_*X) = 0 ,$$

and

$$du(Y, Z) = du(J\iota_*X + v(\iota_*X)U, Z) = du(J\iota_*X, Z) = -du(\iota_*X, JZ) = 0$$

since  $du$  is of bidegree  $(1, 1)$ . Thus  $Y \in \mathcal{U}$  so that  $M^{2q+1}$  is semi-invariant with respect to  $U$ , and hence by Proposition 4.3 its almost contact structure is normal. Finally as

$$\eta(X) = -g(J\iota_*X, U) = g(\iota_*X, V) = v(\iota_*X) ,$$

we have that  $\eta \wedge (d\eta)^q \neq 0$  on  $M^{2q+1}$ . Similarly,  $M^{2p+1}$  is semi-invariant with respect to  $V$ , and is thus a normal contact manifold completing the proof.

Now let  $P$  and  $Q$  denote the projection maps to the tangent spaces of  $M^{2p+1}$  and  $M^{2q+1}$  respectively. We note for later use that  $J(P - u \otimes U) = (P - u \otimes U)J$  as is easily verified, and hence that

$$JP = PJ + u \otimes V + v \otimes U .$$

We now compute the Lie derivative of  $P$  with respect to  $U$  and  $V$ . First note that

$$(\mathcal{L}_U P)X = [U, PX] - P[U, X] .$$

Thus, if  $X$  is  $U$  or  $V$ , we clearly have  $(\mathcal{L}_U P)X = 0$ . If  $X$  is orthogonal to  $U$  but also tangent to  $M^{2p+1}$ , then  $PX = X$  and  $[U, X]$  is again tangent to  $M^{2p+1}$  so that

$$(\mathcal{L}_U P)X = [U, X] - [U, X] = 0 .$$

Finally, if  $X$  is orthogonal to  $V$  and tangent to  $M^{2q+1}$ , then  $PX = 0$ . Let  $Y$  be arbitrary. Then as  $U$  is Killing and  $P$  symmetric, we have

$$\begin{aligned} g((\mathcal{L}_U P)X, Y) &= -g(P[U, X], Y) = -g(\nabla_U X, PY) + g(\nabla_X U, PY) \\ &= g(X, \nabla_U PY) - g(X, \nabla_{PY} U) = g(X, [U, PY]) = 0 . \end{aligned}$$

Similarly  $\mathcal{L}_V P = 0$ , and thus  $P$  and  $Q = I - P$  are projectable by the fibration of § 3.

On the base manifold  $N^{2n-2}$  of the fibration we define an almost product structure as follows.

$$P'X = \pi_* P \bar{\pi} X, \quad Q'X = \pi_* Q \bar{\pi} X.$$

Then a direct computation shows that

$$P'^2 = P', \quad Q'^2 = Q', \quad P'Q' = Q'P' = 0, \quad P' + Q' = I.$$

Moreover as both the distributions  $\mathcal{U}$  and  $\mathcal{V}$  are integrable,  $[P, P] = 0$  so that

$$\begin{aligned} [P', P'](X, Y) &= \pi_* P^2 \bar{\pi} [\pi_* \bar{\pi} X, \pi_* \bar{\pi} Y] + [\pi_* P \bar{\pi} X, \pi_* P \bar{\pi} Y] \\ &\quad - \pi_* P \bar{\pi} [\pi_* P \bar{\pi} X, \pi_* \bar{\pi} Y] - \pi_* P \bar{\pi} [\pi_* \bar{\pi} X, \pi_* P \bar{\pi} Y] \\ &= \pi_* [P, P](\bar{\pi} X, \bar{\pi} Y) = 0. \end{aligned}$$

Thus the induced almost product structure on  $N^{2n-2}$  is integrable, and so  $N^{2n-2}$  is locally the product of two manifolds  $N^{2p}$  and  $N^{2q}$ .

We have already seen that  $J$  is projectable since  $U$  and  $V$  are analytic, and that  $(J' = \pi_* J \bar{\pi}, g')$  is a Hermitian structure on  $N^{2n-2}$ . Now let  $\iota' : N^{2p} \rightarrow N^{2n-2}$  denote the immersion of  $N^{2p}$  in  $N^{2n-2}$ , and let  $X$  be a vector field on  $N^{2p}$ . Then using  $J'P = PJ + u \otimes V + v \otimes U$ , we have

$$\begin{aligned} J' \iota'_* X &= \pi_* J \bar{\pi} P' \iota'_* X = \pi_* J P \bar{\pi} \iota'_* X = \pi_* P J \bar{\pi} \iota'_* X \\ &= \pi_* P \bar{\pi} J' \iota'_* X = P' J' \iota'_* X, \end{aligned}$$

and hence  $N^{2p}$  is an invariant submanifold of  $N^{2n-2}$  and consequently is a Hermitian submanifold of  $N^{2n-2}$ . Moreover, if  $N^{2n-2}$  is Kaehlerian, so is  $N^{2p}$  and similarly  $N^{2q}$ . Also, if each of the induced structures on  $N^{2p}$  and  $N^{2q}$  are Kaehlerian, so is the structure on  $N^{2n-2}$ . Thus using Theorems 3.1 and 4.4 and Proposition 3.2 we have

**Theorem 4.5.** *Let  $M^{2n}$  be a regular Hermitian bicontact manifold with du of bidegree (1, 1). Then the base manifold  $N^{2n-2}$  of the induced fibration is locally the product of two Hermitian manifolds. Moreover,  $N^{2n-2}$  is locally the product of two Kaehler manifolds if and only if the fundamental 2-form  $\Omega$  on  $M^{2n}$  satisfies  $d\Omega = du \wedge v - u \wedge dv$ .*

### 5. Curvature

In this section we give some results on the curvature of a Hermitian manifold admitting a holomorphic pair of automorphisms.

**Proposition 5.1.** *Let  $(M^{2n}, J, g)$  be a Hermitian manifold admitting a holomorphic pair of automorphisms  $(U, V = JU)$ . Then the sectional curvature of a section spanned by  $U$  and  $V$  vanishes.*

*Proof.* Since  $U$  is Killing, from  $g(\nabla_V U, X) - g(\nabla_X U, V) = 0$  which was derived in the proof of Lemma 2.3 it follows that  $2g(\nabla_V U, X) = 0$  and hence that  $\nabla_V U = 0$ . Moreover as  $U$  is a unit vector field, we have  $0 = g(\nabla_X U, U) = -g(\nabla_U U, X)$  giving  $\nabla_U U = 0$ . Thus  $g(R_{UV} U, V) = 0$ , where  $R$  is the

curvature tensor of  $g$ , and hence the sectional curvature of a section spanned by  $U$  and  $V$  vanishes.

**Theorem 5.2.** *If the Hermitian manifold  $M^{2n}$  of Theorem 3.1 has non-negative sectional curvature, then the base manifold  $N^{2n-2}$  also has nonnegative curvature.*

*Proof.* First we note some relations.

$$[\tilde{\pi}X, \tilde{\pi}Y] = \tilde{\pi}[X, Y] + u([\tilde{\pi}X, \tilde{\pi}Y])U + v([\tilde{\pi}X, \tilde{\pi}Y])V .$$

Since  $U$  and  $V$  are Killing, we have

$$\begin{aligned} g(\nabla_{\tilde{\pi}X}\tilde{\pi}Y, U) &= -g(\tilde{\pi}Y, \nabla_{\tilde{\pi}X}U) = -\frac{1}{2}du(\tilde{\pi}X, \tilde{\pi}Y) , \\ g(\nabla_{\tilde{\pi}X}\tilde{\pi}Y, V) &= -g(\tilde{\pi}Y, \nabla_{\tilde{\pi}X}V) = -\frac{1}{2}dv(\tilde{\pi}X, \tilde{\pi}Y) , \end{aligned}$$

and hence

$$\nabla_{\tilde{\pi}X}\tilde{\pi}Y = \tilde{\pi}\nabla'_X Y - \frac{1}{2}du(\tilde{\pi}X, \tilde{\pi}Y)U - \frac{1}{2}dv(\tilde{\pi}X, \tilde{\pi}Y)V ,$$

where  $\nabla'$  is the Riemannian connection of  $g'$ . Also, since  $[U, \tilde{\pi}X]$  is vertical,  $g(\nabla_U\tilde{\pi}X, \tilde{\pi}Y) = g(\nabla_{\tilde{\pi}X}U + [U, \tilde{\pi}X], \tilde{\pi}Y) = \frac{1}{2}du(\tilde{\pi}X, \tilde{\pi}Y)$ , and similarly  $g(\nabla_V\tilde{\pi}X, \tilde{\pi}Y) = \frac{1}{2}dv(\tilde{\pi}X, \tilde{\pi}Y)$ .

We now compute the curvature.

$$\begin{aligned} g(R_{\tilde{\pi}X\tilde{\pi}Y}\tilde{\pi}X, \tilde{\pi}Y) &= g(\nabla_{\tilde{\pi}X}\nabla_{\tilde{\pi}Y}\tilde{\pi}X - \nabla_{\tilde{\pi}Y}\nabla_{\tilde{\pi}X}\tilde{\pi}X - \nabla_{[\tilde{\pi}X, \tilde{\pi}Y]}\tilde{\pi}X, \tilde{\pi}Y) \\ &= g(\nabla_{\tilde{\pi}X}(\tilde{\pi}\nabla'_Y X - \frac{1}{2}du(\tilde{\pi}Y, \tilde{\pi}X)U - \frac{1}{2}dv(\tilde{\pi}Y, \tilde{\pi}X)V) \\ &\quad - \nabla_{\tilde{\pi}Y}\tilde{\pi}\nabla'_X X - \nabla_{[\tilde{\pi}X, \tilde{\pi}Y]}\tilde{\pi}X, \tilde{\pi}Y) \\ &= g(\tilde{\pi}\nabla'_X\nabla'_Y X, \tilde{\pi}Y) - \frac{1}{2}du(\tilde{\pi}Y, \tilde{\pi}X)g(\nabla_{\tilde{\pi}X}U, \tilde{\pi}Y) \\ &\quad - \frac{1}{2}dv(\tilde{\pi}Y, \tilde{\pi}X)g(\nabla_{\tilde{\pi}X}V, \tilde{\pi}Y) - g(\tilde{\pi}\nabla'_Y\nabla'_X X, \tilde{\pi}Y) \\ &\quad - g(\tilde{\pi}\nabla'_{[X, Y]}X, \tilde{\pi}Y) - u([\tilde{\pi}X, \tilde{\pi}Y])g(\nabla_U\tilde{\pi}X, \tilde{\pi}Y) \\ &\quad - v([\tilde{\pi}X, \tilde{\pi}Y])g(\nabla_V\tilde{\pi}X, \tilde{\pi}Y) \\ &= g'(R'_{XY}X, Y) \circ \pi + \frac{3}{4}du(\tilde{\pi}X, \tilde{\pi}Y)^2 + \frac{3}{4}dv(\tilde{\pi}X, \tilde{\pi}Y)^2 \end{aligned}$$

since  $du(\tilde{\pi}X, \tilde{\pi}Y) = \tilde{\pi}Xu(\tilde{\pi}Y) - \tilde{\pi}Yu(\tilde{\pi}X) - u([\tilde{\pi}X, \tilde{\pi}Y]) = -u([\tilde{\pi}X, \tilde{\pi}Y])$ . Now for the sectional curvature  $K$  we have

$$K(\tilde{\pi}X, \tilde{\pi}Y) = \frac{-g(R_{\tilde{\pi}X\tilde{\pi}Y}\tilde{\pi}X, \tilde{\pi}Y)}{g(\tilde{\pi}X, \tilde{\pi}X)g(\tilde{\pi}Y, \tilde{\pi}Y) - g(\tilde{\pi}X, \tilde{\pi}Y)^2} .$$

Thus, if  $K \geq 0$ , then  $g(R_{\tilde{\pi}X\tilde{\pi}Y}\tilde{\pi}X, \tilde{\pi}Y) \leq 0$  and hence

$$-g'(R'_{XY}X, Y) \circ \pi \geq \frac{3}{4}(du(\tilde{\pi}X, \tilde{\pi}Y)^2 + dv(\tilde{\pi}X, \tilde{\pi}Y)^2) ,$$

from which it follows that the sectional curvature  $K'(X, Y) \geq 0$ .

## References

- [ 1 ] D. E. Blair, *Geometry of manifolds with structural group  $\mathcal{U}(n) \times \mathcal{O}(s)$* , J. Differential Geometry **4** (1970) 155–167.
- [ 2 ] D. E. Blair, G. D. Ludden & K. Yano, *Differential geometric structures on principal toroidal bundles*, Trans. Amer. Math. Soc. **181** (1973) 175–184.
- [ 3 ] ———, *Semi-invariant immersions*, to appear in Kōdai Math. Sem. Rep.
- [ 4 ] E. Calabi & B. Eckmann, *A class of compact complex manifolds which are not algebraic*, Ann. of Math. **58** (1953) 494–500.
- [ 5 ] S. S. Chern, *Complex manifolds without potential theory*, Math. Studies No. 15, Van Nostrand, Princeton, 1967.
- [ 6 ] A. Morimoto, *On normal almost contact structures*, J. Math. Soc. Japan **15** (1963) 420–436.
- [ 7 ] H. Nakagawa, *f-structures induced on submanifolds in spaces, almost Hermitian or almost Kaehlerian*, Kōdai Math. Sem. Rep. **18** (1966) 161–183.
- [ 8 ] S. Sasaki & Y. Hatakeyama, *On differentiable manifolds with contact metric structures*, J. Math. Soc. Japan **14** (1962) 249–271.
- [ 9 ] K. Yano, *On a structure defined by a tensor field  $f$  of type  $(1, 1)$  satisfying  $f^3 + f = 0$* , Tensor **15** (1963) 99–109.

MICHIGAN STATE UNIVERSITY