

## Exceptional Case

# Gitelman syndrome and pregnancy

Michael N. Moustakakis<sup>1</sup> and Margarita Bockorny<sup>2</sup>

<sup>1</sup>Section of Nephrology, Saint Francis Hospital and Medical Center, Hartford, CT, USA and <sup>2</sup>University of Connecticut Health Center, Farmington, CT, USA

Correspondence and offprint requests to: Michael N. Moustakakis; E-mail: m.moustakakis@gmail.com

### Abstract

Gitelman syndrome (GS) is an autosomal-recessive condition characterized by hypokalemia, hypomagnesemia and hypocalciuria. Very little information is available in the literature to guide the management of pregnant patients with GS. We report a case of a 27-year-old woman with GS who became pregnant and despite persistent hypokalemia and hypomagnesemia during pregnancy and labor, had a successful maternal and fetal outcome.

**Keywords:** Gitelman syndrome; hypokalemia; pregnancy

### Introduction

Gitelman syndrome (GS) is a rare autosomal-recessive inherited disorder with 99% penetrance caused by inactivating mutations in the SLC12A3 gene, localized on the 16q13 chromosome. This genetic defect impairs the function of the thiazide-sensitive sodium-chloride cotransporter in the distal convoluted renal tubule. GS is the hypocalciuric, hypomagnesemic variant of Bartter syndrome, which was first described in 1966 [1]. The impact of this condition on maternal and fetal outcome is still unclear. Here, we describe the management and pregnancy course of a 27-year-old patient with GS.

### Case presentation

We report the course of a pregnancy in a 27-year-old woman previously diagnosed with GS. Her pregnancy was uneventful and she remained asymptomatic despite persistent hypokalemia. Potassium levels were followed at a regular basis and ranged between 2.3 and 3.1 mmol/L (2.3 and 3.1 mEq/L). She was electively admitted to hospital at 39 weeks of gestational age for full-term induction of labor. A PICC line was placed to allow replacement of electrolytes. The patient received 240 mEq of intravenous potassium chloride and 180 mEq per oral, as well as 4 g of intravenous magnesium sulfate and 1200 mEq of oral magnesium oxide. She was continued on 150 mg daily of eplerenone. Despite aggressive electrolyte replacement, her potassium and magnesium levels ranged between 2.0 and 3.0 mmol/L (2.0 and 3.0 mEq/L) and 1.1 and 2.6 mmol/L (1.1 and 2.6 mEq/L), respectively. The baby was born healthy with 2863 g, and Apgar measurements were 8 and 9 at 1 and 5 min, respectively. She was discharged two days after giving birth and has remained symptom free.

### Discussion

First described in 1966 by Gitelman *et al.*, GS is a rare autosomal-recessive condition characterized by hypokalemia, metabolic alkalosis and hypomagnesemia [1]. It has since been shown to result from a mutation in the SLC12A3 gene on chromosome 16 leading to a loss of function mutation in the sodium-chloride cotransporter in the distal convoluted tubule. This leads to potassium and magnesium wasting into the cortical collecting duct as the increase in delivered sodium is reabsorbed through the epithelial sodium channel. Despite being normotensive, the patients clinically have activation of the renin-angiotensin-aldosterone axis [2]. To date, 24 pregnancies in 18 women with GS have been described in the literature [3–15]. Twenty of the pregnancies had no reported fetal complication (Table 1), six pregnancies were complicated by oligohydramnios [5, 8, 9, 12, 13] and one was complicated by intrauterine growth retardation [8]. The management of maternal GS during pregnancy continues to be a challenge. Renin, angiotensin II and aldosterone levels are known to increase during pregnancy [16]. Kaliuresis does not typically ensue presumably due to the antimineralocorticoid effects of progesterone demonstrated by Ehrlich and Lindheimer in 1972 [17]. De Haan *et al.* have suggested that this protective mechanism may be impaired in GS patients thus resulting in an exacerbation of potassium and magnesium losses [8]. Oral potassium and magnesium supplements continue to be the mainstay of therapy. The increase in the requirement for oral potassium during pregnancy has been documented by several authors including Talaulikar *et al.* in 2005 who reported a 6-fold increase in potassium and magnesium requirements in their patient [7]. Six of the pregnancies document the use of intravenous potassium [7–12]. Intravenous magnesium was required in three cases [7–9]. Two authors report achievement of

**Table 1.** Summary with the cases of Gitelman syndrome during pregnancy

Author	Age at diagnosis (years)	Diagnosis	Maternal age (years)	Number of patients	Number of pregnancies	Symptoms
Jones JM <i>et al.</i> [5]	35	1st trimester	35	1	2	Initially presented with new onset of seizures N/A
Basu A <i>et al.</i> [6]	17	Before pregnancy	1) 24 2) 25 3) 27	1	3	1st- asymptomatic except for tiredness in 3rd trimester 2nd- asymptomatic 3rd- asymptomatic
Talaulikar GS <i>et al.</i> [7]	20	Before pregnancy	1) 22	1	2	1) Increasing tiredness, muscle cramps. Required 3 hospitalizations 2) Not described
Kwan <i>et al.</i> [10] Srinivas SK <i>et al.</i> [11]	17	2nd trimester (17 weeks)	2) 30 17	1	1	Nausea, vomiting, proximal lower extremity muscle weakness. VF arrest, seizures, prolonged QTc
De Haan J <i>et al.</i> [8]	N/A	Before pregnancy	1) 24	2	1	1st- symptomatic, not specified
De Arriba G <i>et al.</i> [12]	18	Before pregnancy	2) 27 36	1	1	2nd- muscle weakness, fatigue Limb paresthesias and cramps, myalgia
Daskalakis G <i>et al.</i> [13]	20	1st trimester (10 weeks)	20	1	1	Tiredness, muscle weakness, decreased muscle strength and tenderness bilaterally in lower extremities
McCarthy FP <i>et al.</i> [9]	32	1st trimester (9 weeks)	32	1	1	Episodic severe fatigue, postural hypotension
Raffi <i>et al.</i> [14]	22	Before pregnancy	27	1	1	Not described
Morton A <i>et al.</i> [15]	21	Before pregnancy	N/A	1	1	Asymptomatic
Mascetti L <i>et al.</i> [4]	N/A	N/A	1) 27 2) 32 3) 28 4) 33 5) 33/35	5	6	1, 2, 3, 4) uneventful 5) tiredness and tetanic seizures
Calo LA <i>et al.</i> [3]	N/A	Before pregnancy	39 24	2	3	Intermittent nausea and vomiting Nausea // N/A
Gestational complications	Delivery	Neonate	Gestational age	Potassium values	Magnesium values	Treatment
Oligohydramnios	Induced vaginal delivery	Healthy male infant. 2509 g	37 weeks	~3.2 mEq/L	~1.3 mEq/L	K and Mg supplementation
Oligohydramnios	Induced vaginal delivery	Healthy female infant, 2410 g	36 weeks	N/A	N/A	K and Mg supplementation
1, 2 and 3) None	1st- spontaneous labor	1st- healthy male infant. 2620 g	1st-35 weeks	1st- K range = 2.3–2.8 mmol/L	1st- Mg range = 0.53–0.58 mmol/L	1st- K and Mg supplementation
	2nd- spontaneous labor	2nd- healthy female infant, 2910 g	2nd- 38 weeks	2nd- maximum level = 3.1 mmol/L	2nd- maximum level = 0.56 mmol/L	2nd- K and Mg supplementation
	3rd-spontaneous labor	3- healthy female infant. 3480 g	3rd- 38 weeks	3rd- N/A	3rd- N/A	3rd-N/A
1) None	Normal delivery	Healthy infant	1) N/A	K range = 2.3–3.4 mmol/L	1 and 2)N/A	1 and 2) K and Mg supplementation
2) Gestational diabetes	Elective C-section	N/A	2) 39 weeks	N/A	N/A	K and Mg supplementation
None	Uncomplicated vaginal delivery, induced for post-term pregnancy	Healthy female infant, 3250 g	41 weeks and 4 days	N/A	N/A	K and Mg supplementation
1st-Oligohydramnios, partial placenta previa, intrauterine growth restriction.	1st- primary C-section, healthy infant 2nd- secondary C-section. healthy infant	Healthy female infant, 3970 g Healthy male infant, 3845 g	1 and 2) 38 weeks	N/A	N/A	K and Mg supplementation. 1st case required i.v. supplementation
Oligohydramnios	Programmed C-section, w/o complications	Healthy infant, 3000 g	N/A	2.4–2.8 mEq/L	N/A	K and Mg supplementation, Spironolactone.
Oligohydramnios	Elective C-section (breech presentation)	Healthy female baby, 3350 g	273	Maximum level = 3.0 mmol/L	Maximum level = 0.68 mmol/L	K and Mg supplementation

(continued)

Table 1. Continued

Gestational complications	Delivery	Neonate	Gestational age	Potassium values	Magnesium values	Treatment
Oligohydramnios	Induced C-section (failure to progress in the first stage)	Healthy female baby, 2940 g	266	K range = 2.6–3.3 mmol/L	Mg range = 0.47–0.66 mmol/L	K and Mg supplementation, amiloride. Required 39 hospitalizations i.v. supplementation of K and Mg
Fetal macrosomia (patient had gestational diabetes)	Induced C-section	Healthy male baby, 3080 g	N/A	N/A	N/A	K and Mg supplementation, amiloride (1st trimester). Required 1 hospitalization for i.v. supplementation of K and Mg.
None	Vaginal delivery	Healthy female baby, 3630 g	273	2.6–2.9 mmol/L	N/A	Epleronone
1, 2, 3, 4 and 5) None	N/A	Healthy male baby, 4080 g	40 weeks	N/A	N/A	K and Mg Homeopathy K, amiloride K, Mg None (1st pregnancy) / K (2nd pregnancy)
		Healthy female baby, 3380 g	38 weeks			
		Healthy female baby, 2420 g	37 weeks			
		Healthy male baby, 2810 g	41 weeks			
1 and 2) None	N/A	Healthy male baby, 2830 g/ healthy male baby, 2640 g	38 weeks/36 weeks	N/A	N/A	1 and 2) K and Mg supplementation
		Healthy female baby 2900 g	At term			
		Healthy female baby 3125 g/ healthy female baby 3400 g	At term/ At term			

potassium levels in the 3.6–4 mmol/L (3.6–4.0 mEq/L) range [10, 14]. The remainder of the documented cases had peak potassium levels ranging from 2.8 to 3.3 mmol/L (2.8–3.3 mEq/L). It has been suggested that normalization of potassium and magnesium levels is not required for a good obstetric and neonatal outcome [6]. Worsening of the symptoms of GS such as fatigue, cramping, tetany and dizziness has been described in the GS pregnancies [7, 12, 13]. The symptoms may be exacerbated by hyperemesis and fetal demand for potassium [9, 12]. Worsening of symptoms is cited as a reason for the modification of therapy [3, 4, 6–11, 13]. Our patient did not exhibit symptoms during her pregnancy, and we felt that intravenous potassium loading would be inefficient and unnecessary in the absence of symptoms. Beyond the use of supplemental cations, the management of patients with GS has included the use of potassium-sparing diuretics. The US Food and Drug Administration has deemed spironolactone a category C drug in pregnancy. De Arriba *et al.* describe the use of spironolactone in one pregnancy [12]. The authors note that no feminization was seen in the male newborn. This is consistent with the findings of Mascetti *et al.* who describe a series of spironolactone-exposed children born to a group of mothers with Bartter syndrome, another potassium-wasting nephropathy [4]. The use of amiloride and eplerenone, both class B drugs in pregnancy, has been previously documented in GS [4, 9, 14, 15]. In contrast to Bartter syndrome, the use of non-steroidal anti-inflammatory drugs (NSAIDs) to inhibit prostaglandin synthase

is theoretically of no benefit as GS is not a hyperprostaglandinemic state [18]. There is a risk of a lack of ductus arteriosus closure with NSAID use. Angiotensin-converting enzyme inhibitors are contraindicated in pregnancy due to teratogenic effects as well as impaired fetal growth. There has been a concern over the potential for a ventricular tachyarrhythmia during pregnancy and childbirth due to hypokalemia. Fortunately, no such complication has been observed in any of the documented GS pregnancies. The extensive work of Calo *et al.* in this area suggests however that the upregulation of the nitric oxide system and vasodilation seen in GS may limit the physiologic response to increased myocardial demand [19, 20]. It is also suggested in subsequent work that increased angiotensin 1–7 levels seen in GS may be anti-arrhythmic at low levels and proarrhythmic at very high levels [21]. The small number of documented GS pregnancies precludes any rigorously studied recommendations for management in the peripartum period. The anesthesia literature describes a risk of complications in GS patients in the perioperative setting. These include electrocardiographic changes as well as vasodilation, electrolyte imbalance and alkalemia complicating the ventilatory management of the patient [22]. Uneventful spinal anesthesia for cesarean delivery is also described in the literature [23]. Given the expected physiologic demands of labor and delivery, it appears that an elective, multidisciplinary approach to the delivery of the child is most prudent. Baseline electrocardiography should be obtained as up to 50% of patients with GS

have QT interval prolongation. Cardiac telemetry and central venous access should be considered at the time of delivery. Frequent monitoring of electrolytes is advisable as is blood pressure monitoring given the propensity of GS patients have vasodilation. A drop in serum potassium during labor is not previously described in the literature. The postpartum period is associated with natriuresis; however, this would not explain a drop in potassium during active labor. It may be, however, that the drop in serum potassium observed here is related to a shift in potassium from the extracellular to intracellular space as could be seen in any patient and is not unique to GS. In our patient, we suspect that increased  $\beta$ -adrenergic tone may have played a role. The effects of epinephrine on hypokalemia were described in 1983 by Struthers *et al.* Pretreatment with a thiazide diuretic was associated with a more significant drop in serum potassium during epinephrine loading than that seen in the control group receiving epinephrine alone [24]. As GS mimics the effects of thiazide diuretics, one could assume a similar effect in these patients. Cellular shift of potassium may also occur because of rising blood glucose concentrations.

In conclusion, the experience of clinicians treating pregnant women with GS is limited but is also being described with more frequency. There is no evidence of significant risk to the fetus but maternal symptoms may worsen during the pregnancy. The management of GS should be focused on the replacement of potassium and magnesium though routine intravenous supplementation and should be reserved for cases of worsened symptoms. Cautious use of adjunctive treatment with potassium-sparing agents should be considered as the potassium requirement may increase significantly during pregnancy. In addition to GS-related potassium and magnesium loss, hypokalemia can be exacerbated by cellular shift as well as extrarenal losses of cations which can affect any patient. No consensus recommendations exist for the perinatal management of the pregnant GS patient but an elective delivery in a setting with appropriate cardiac monitoring and central venous access is prudent as potassium and magnesium aberrations during active labor may be seen.

*Conflict of interest statement.* None declared.

## References

- Gitelman HJ, Graham JB, Welt LG. A new familial disorder characterized by hypokalemia and hypomagnesemia. *Trans Assoc Am Phys* 1966; 79: 221–235
- Naesens M, Steels P, Verberckmoes R *et al.* Bartter's and Gitelman's syndromes: from gene to clinic. *Nephron Physiol* 2004; 96: 65–78
- Calo LA, Caielli P. Gitelman's syndrome and pregnancy: new potential pathophysiological influencing factors, therapeutic approach and materno-fetal outcome. *J Matern Fetal Neonatal Med* 2011; 1–3
- Mascetti L, Bettinelli A, Simonetti GD *et al.* Pregnancy in inherited hypokalemic salt-losing renal tubular disorder. *Obstet Gynecol* 2011; 117: 512–516
- Jones JM, Dorrel S. Outcome of two pregnancies in a patient with Gitelman's syndrome: a case report. *J Matern Fetal Investig* 1998; 8: 147–148
- Basu A, Dillon RD, Taylor R *et al.* Is normalisation of serum potassium and magnesium always necessary in Gitelman syndrome for a successful obstetric outcome? *BJOG* 2004; 111: 630–634
- Talaulikar GS, Falk MC. Outcome of pregnancy in a patient with Gitelman syndrome: a case report. *Nephron Physiol* 2005; 101: 35–38
- De Haan J, Geers T, Berghout A. Gitelman syndrome in pregnancy. *Int J Gynaecol Obstet* 2008; 103: 69–71
- McCarthy FP, Magee CN, Plant WD *et al.* Gitelman's syndrome in pregnancy: case report and review of literature. *Nephrol Dial Transplant* 2010; 25: 1338–1340
- Kwan TK, Falk MC. Second pregnancy outcome in a patient with Gitelman syndrome without the use of parenteral electrolyte supplementation. *Aust N Z J Obstet Gynaecol* 2011; 51: 94–95
- Srinivas SK, Sukhan S, Elovitz MA. Nausea, emesis, and muscle weakness in a pregnant adolescent. *Obstet Gynecol* 2006; 107: 481–484
- De Arriba G, Sanchez-Heras M, Basterrechea MA. Gitelman syndrome during pregnancy: a therapeutic challenge. *Arch Gynecol Obstet* 2009; 280: 807–809
- Daskalakis G, Marinopoulos S, Mousiolis A *et al.* Gitelman syndrome-associated severe hypokalemia and hypomagnesemia: case report and review of literature. *J Matern Fetal Neonatal Med* 2010; 23: 1301–1304
- Raffi F, Fairlie FM, Madhuvrata P *et al.* Pregnancy with Gitelman syndrome. *Obstet Med* 2011; 4: 39–41
- Morton A, Panitz B, Bush A. Eplerenone for Gitelman syndrome in pregnancy. *Nephrology (Carlton)* 2011; 16: 349
- Baylis C, Davison JM. The normal renal physiological changes which occur during pregnancy. In: Davison AM, Cameron JS, Grunfeld JP *et al.* (eds) *Oxford Textbook of Clinical Nephrology*, 3rd ed. Oxford: Oxford University Press, 2005, pp. 2213–2231
- Ehrlich EN, Lindheimer MD. Effect of administered mineralocorticoids or ACTH in pregnant women: attenuation of kaliuretic influence of mineralocorticoids during pregnancy. *J Clin Invest* 1972; 51: 1301–1309
- Luthy C, Bettinelli A, Iselin S *et al.* Normal prostaglandinuria E2 in Gitelman's syndrome, the hypocalciuric variant of Bartter's syndrome. *Am J Kidney Dis* 1995; 25: 824
- Calò L, Davis PA, Milani M *et al.* Increased endothelial nitric oxide synthase mRNA level in Bartter's and Gitelman's syndrome. Relationship to vascular reactivity. *Clin Nephrol* 1999; 51: 12–17
- Calò L, D'Angelo A, Cantaro S *et al.* Increased urinary NO<sub>2</sub>/NO<sub>3</sub>- and cyclic guanosine monophosphate levels in patients with Bartter's syndrome: Relationship to vascular reactivity. *Am J Kidney Dis* 1996; 27: 784–789
- Calò LA, Schiavo S, Davis PA *et al.* ACE2 and angiotensin 1–7 are increased in a human model of cardiovascular hyporeactivity: Pathophysiological implications. *J Nephrol* 2010; 23: 472–477
- Farmer JD, Vasdev GM, Martin DP. Perioperative considerations in patients with Gitelman syndrome: a case series. *J Clin Anesth* 2012; 24: 14–18
- Shanbhag S, Neil J, Howell C. Anaesthesia for cesarean section in a patient with Gitelman's syndrome. *Int J Obstet Anesth* 2010; 19: 451–453
- Struthers AD, Whitesmith R, Reid JL. Prior thiazide diuretic treatment increases adrenaline-induced hypokalaemia. *Lancet* 1983; 321: 1358–1361

Received for publication: 1.7.12; Accepted in revised form: 9.8.12