

Glandular Odontogenic Cysts (GOCs) Lack *MAML2* Rearrangements: A Finding to Discredit the Putative Nature of GOC as a Precursor to Central Mucoepidermoid Carcinoma

Justin A. Bishop · Raluca Yonescu ·
Denise Batista · Gary R. Warnock ·
William H. Westra

Received: 3 January 2014 / Accepted: 11 March 2014 / Published online: 20 March 2014
© Springer Science+Business Media New York 2014

Abstract Glandular odontogenic cyst (GOC) is a cyst of the gnathic bones that is characterized by squamous and glandular differentiation. The histopathologic features of GOC overlap considerably with central mucoepidermoid carcinoma (MEC), suggesting that GOC could be a precursor lesion to, or even a low-grade form of, central MEC. Differentiating the two lesions may be difficult or impossible on a limited biopsy. *MAML2* rearrangements have been recently found to be specific for MEC, even those arising in the jaws. An analysis of *MAML2* in GOCs could help clarify its relationship with central MEC. Tissue blocks from 21 GOCs and 5 central MECs were retrieved from the surgical pathology archives of The Johns Hopkins Hospital. Each MEC exhibited solid areas and clear-cut stromal invasion. In addition, 4 of the MECs demonstrated cystic areas that were histologically similar to GOC. Break-

apart fluorescence in situ hybridization for *MAML2* was performed. For the MECs, analysis was performed on both the solid components and the cystic areas that resembled GOC. *MAML2* rearrangements were identified in all 5 of the MECs, but in none of the 21 GOCs (100 vs. 0 %; $p < 0.0001$, Fisher's Exact). In the MECs, the rearrangement was present in both the solid and GOC-like cystic areas. While central MECs consistently harbor the *MAML2* rearrangement, even in low-grade cystic areas that resemble a pre-existing GOC, true GOCs do not. Accordingly, GOC does not appear to represent an early or low-grade form of central MEC, but rather an unrelated lesion. The high sensitivity and specificity of *MAML2* rearrangement for MECs points to its utility as a diagnostic adjunct in separating mucinous cystic lesions of the gnathic bones.

Keywords Glandular odontogenic cyst · Central mucoepidermoid carcinoma · *MAML2*

J. A. Bishop · R. Yonescu · D. Batista ·
G. R. Warnock · W. H. Westra
Department of Pathology, The Johns Hopkins Medical
Institutions, Baltimore, MD, USA

J. A. Bishop · W. H. Westra
Department of Otolaryngology/Head and Neck Surgery, The
Johns Hopkins Medical Institutions, Baltimore, MD, USA

J. A. Bishop (✉)
The Johns Hopkins University School of Medicine, 401 N.
Broadway, Weinberg 2249, Baltimore, MD 21231, USA
e-mail: jbisho16@jhmi.edu; jbishop@jhmi.edu


G. R. Warnock · W. H. Westra
Department of Dermatology, The Johns Hopkins Medical
Institutions, Baltimore, MD, USA

W. H. Westra
Department of Oncology, The Johns Hopkins Medical
Institutions, Baltimore, MD, USA

Introduction

Central (i.e., primary intraosseous) mucoepidermoid carcinoma (MEC) is a rare malignant neoplasm of the jaws that is identical in most respects to MEC of the salivary glands, but arises entirely within bone [1–5]. Glandular odontogenic cyst (GOC) is an uncommon benign cystic lesion of the gnathic bones initially described in 1987 as “sialo-odontogenic cyst” or “mucoepidermoid odontogenic cyst.” [6–16] Since its first description, many investigators have recognized that GOC shares many histologic features with central MEC [1, 2, 5, 7, 9, 10, 16–20]. Indeed, it is common to see areas indistinguishable from GOC within a central MEC [2]. Conversely, “MEC-like” areas within the walls of GOCs have been described [10].

Fig. 1 In this case of central mucoepidermoid carcinoma, the solid areas of clear cut invasive tumor (a) were positive for the *MAML2* rearrangement by break apart FISH (b). In very cystic areas of the tumor that mimicked glandular odontogenic cyst (c), the translocation was also present (d)



These morphologic similarities have prompted speculation that the GOC represents a precursor to, or even a low-grade form of, central MEC [2, 18, 19]. Although not always easy on morphologic grounds, distinguishing GOC from central MEC is important because MECs have a higher rate of recurrence, carry a potential to metastasize to regional lymph nodes, and can in some cases be lethal [1, 4, 5, 21, 22].

Rearrangements of *MAML2* have recently been detected in up to 75 % of MECs of salivary glands, and are very specific for this tumor type [23–27]. They preferentially occur in low/intermediate grade MECs with favorable prognosis where they are regarded as an early genetic alteration that drives tumorigenesis [28]. *MAML2* rearrangements have been reported in two of three central MECs [29, 30], but the *MAML2* status of GOC is not known. We sought to clarify the relationship of GOC to central MEC by performing *MAML2* molecular analysis on these lesions.

Methods

Cases

The surgical pathology archives of The Johns Hopkins Hospital were searched for cases of GOC and central (i.e., primary intraosseous) MEC. Slides and tissue blocks from 21 GOCs and 5 central MECs were retrieved. Hematoxylin and eosin-stained sections were reviewed to confirm the


diagnoses. Each MEC exhibited solid areas and clear-cut stromal invasion. In addition, four of the MECs demonstrated focal cystic areas that were histologically similar to GOC (Fig. 1).

Fluorescence In Situ Hybridization

Fluorescence in situ hybridization (FISH) was performed on formalin fixed paraffin embedded tissue sections using the commercially available *MAML2* dual color break apart probe (Z-2014-200, Zytovision, Germany). Prior to hybridization the slides were deparaffinized utilizing a VP 2000 processor (Abbott Molecular, Des Plaines, IL) with pretreatment with protease I. Following deparaffinization the slides and the *MAML2* probe were co-denatured at 80 °C for 7 min and allowed to hybridize for 22 h at 37 °C in a humidified atmosphere. The slides were then washed with agitation in 2× SSC/0.3 % NP-40 for 2 min at 72 °C and for 2 min at room temperature. Traces of detergent were removed by washing the slides in 2× SSC at room temperature. The slides were then counterstained with DAPI and a cover slip was applied using Vectashield mounting medium (H-1000, Vector Laboratories, Inc.).

A fluorescence microscope was used to evaluate the probe pattern for each case. Cells with two fusion signals of one orange and one green fluorochrome were scored as normal. Cells with rearrangements for *MAML2* gene had one normal fusion signal and one orange and one green signal at a distance from each other. A parotid gland MEC known to harbor the *MAML2* rearrangement served as a

Fig. 2 Despite their histologic similarity to cystic mucoepidermoid carcinoma, all cases of glandular odontogenic cyst (a) were negative for the *MAML2* rearrangement by break apart FISH (b)



positive control, while benign dental follicular tissue served as a negative control.

Results

MAML2 rearrangement was identified in all five cases of central MEC (Fig. 1). In contrast, all 21 GOCs were negative for the *MAML2* rearrangement (100 vs. 0 %; $p < 0.0001$, Fisher's Exact) (Fig. 2). In the 4 MECs where an attenuated cystic area that resembled GOC was present, the rearrangement was detected in both the solid and the cystic zones (Fig. 1).

Discussion

While MEC is the most common malignancy of salivary glands, it may also rarely occur as a primary neoplasm of the jaws (i.e., central MEC). The definitive explanation for the presence of a salivary gland tumor within gnathic bones has been elusive. One theory is that central MECs arise from ectopically displaced benign salivary tissue [1, 4, 5]. A second theory is that central MECs arise from pre-existing benign odontogenic cysts [1, 2, 4, 21]. In light of significant morphologic overlap, GOC in particular has been identified as a likely progenitor for central MEC [2, 18]. Like low grade MECs, GOCs are cystic with an epithelial lining comprised of mixed cell types including mucinous cells and non-keratinizing squamous cells. *MAML2* rearrangements occur in up to 75 % of MEC, but their detection is not site-specific. *MAML2* rearrangements are found not just in MECs derived from the major salivary glands [23, 24, 31], but have been reported in MEC of the lung [32], uterine cervix [33], and thymus [34]. Indeed, 2 of 3 tested central MECs of the gnathic bones have been found to harbor *MAML2* rearrangements [29, 30]. GOCs have not been previously analyzed for the chromosomal translocation, but determination of *MAML2* status could

help clarify their putative relationship to central MECs as a precursor lesion.

In this study, a *MAML2* rearrangement was detected in all 5 of the central MECs, but in none of the 21 GOCs. Moreover, in the central MECs the *MAML2* rearrangement was uniformly distributed throughout the solid, invasive and lining components. The lining component morphologically resembles a GOC and, based on this resemblance, GOC has been incriminated by some as a precursor from which central MECs arise [2, 18, 19]. The striking disparity in *MAML2* status suggests that GOC and central MEC are separate entities, and that GOC should not be regarded as an early or low grade form of MEC, or even as a precursor of MEC. Admittedly, some genetic alterations occur at later stages of tumorigenesis and may not be routinely encountered in early lesions, but this is not the case for *MAML2* in the development of MECs. *MAML2* rearrangements are driver alterations that are necessary for initiating and maintaining MEC tumorigenesis [35]. Indeed, they are consistently detected in the lowest grades of MEC where they represent the first and only detectable metaphase genetic alteration [28].

In addition to providing biologic insight into the relationship between GOC and central MEC, our findings also have a practical diagnostic application. In those cases where diagnosis of a cystic MEC or GOC cannot be easily made on morphologic grounds alone, the detection of a *MAML2* rearrangement would provide compelling evidence to support MEC. Potentially, a similar approach could be used for other tumors that might enter the differential diagnosis of central MEC (e.g., clear cell odontogenic carcinoma) provided that the specificity of *MAML2* rearrangements for MEC is confirmed across other types of odontogenic tumors just as we have confirmed *MAML2* status for GOCs.

Acknowledgments This study was funded in part by the National Institutes of Health/National Institute of Dental and Craniofacial Research (NIH/NIDCR) Head and Neck SPORE Grant P50 DE019032.

References

- Ellis G, Auclair PL. Central (primary intraosseous) mucoepidermoid carcinoma. AFIP atlas of tumor pathology: tumors of the salivary glands. Silver Spring, MD: ARP Press; 2008. p. 193–6.
- Waldron CA, Koh ML. Central mucoepidermoid carcinoma of the jaws: report of four cases with analysis of the literature and discussion of the relationship to mucoepidermoid, sialodontogenic, and glandular odontogenic cysts. *J Oral Maxillofac Surg.* 1990;48:871–7.
- Browand BC, Waldron CA. Central mucoepidermoid tumors of the jaws. Report of nine cases and review of the literature. *Oral Surg Oral Med Oral Pathol.* 1975;40:631–43.
- Brookstone MS, Huvos AG. Central salivary gland tumors of the maxilla and mandible: a clinicopathologic study of 11 cases with an analysis of the literature. *J Oral Maxillofac Surg.* 1992;50:229–36.
- Martinez-Madrigal F, Pineda-Daboin K, Casiraghi O, Luna MA. Salivary gland tumors of the mandible. *Ann Diagn Pathol.* 2000;4:347–53.
- Sadeghi EM, Weldon LL, Kwon PH, Sampson E. Mucoepidermoid odontogenic cyst. *Int J Oral Maxillofac Surg.* 1991;20:142–3.
- Kaplan I, Anavi Y, Hirshberg A. Glandular odontogenic cyst: a challenge in diagnosis and treatment. *Oral Dis.* 2008;14:575–81.
- Araujo de Moraes HH, Jose de Holanda Vasconcellos R, de Santana Santos T, Guedes Queiroz LM, Dantas da Silveira EJ. Glandular odontogenic cyst: case report and review of diagnostic criteria. *J Craniomaxillofac Surg.* 2012;40:e46–50.
- Robinson RA, Vincent SD. Glandular odontogenic cyst. AFIP atlas of tumor pathology: tumors and cysts of the jaws. Silver Spring, MD: ARP Press; 2012. p. 26–30.
- Fowler CB, Brannon RB, Kessler HP, Castle JT, Kahn MA. Glandular odontogenic cyst: analysis of 46 cases with special emphasis on microscopic criteria for diagnosis. *Head Neck Pathol.* 2011;5:364–75.
- Shen J, Fan M, Chen X, Wang S, Wang L, Li Y. Glandular odontogenic cyst in China: report of 12 cases and immunohistochemical study. *J Oral Pathol Med.* 2006;35:175–82.
- Noffke C, Raubenheimer EJ. The glandular odontogenic cyst: clinical and radiological features; review of the literature and report of nine cases. *Dentomaxillofac Radiol.* 2002;31:333–8.
- Gardner DG, Kessler HP, Morency R, Schaffner DL. The glandular odontogenic cyst: an apparent entity. *J Oral Pathol.* 1988;17:359–66.
- Patron M, Colmenero C, Larrauri J. Glandular odontogenic cyst: clinicopathologic analysis of three cases. *Oral Surg Oral Med Oral Pathol.* 1991;72:71–4.
- Semba I, Kitano M, Mimura T, Sonoda S, Miyawaki A. Glandular odontogenic cyst: analysis of cytokeratin expression and clinicopathological features. *J Oral Pathol Med.* 1994;23:377–82.
- Padayachee A, Van Wyk CW. Two cystic lesions with features of both the botryoid odontogenic cyst and the central mucoepidermoid tumour: sialo-odontogenic cyst? *J Oral Pathol.* 1987;16:499–504.
- Prabhu S, Rekha K, Kumar G. Glandular odontogenic cyst mimicking central mucoepidermoid carcinoma. *J Oral Maxillofac Pathol.* 2010;14:12–5.
- Manojlovic S, Grgurevic J, Knezevic G, Kruslin B. Glandular odontogenic cyst: a case report and clinicopathologic analysis of the relationship to central mucoepidermoid carcinoma. *Head Neck.* 1997;19:227–31.
- Pires FR, Chen SY, da Cruz Perez DE, de Almeida OP, Kowalski LP. Cytokeratin expression in central mucoepidermoid carcinoma and glandular odontogenic cyst. *Oral Oncol.* 2004;40:545–51.
- Vered M, Allon I, Buchner A, Dayan D. Is maspin immunolocalization a tool to differentiate central low-grade mucoepidermoid carcinoma from glandular odontogenic cyst? *Acta Histochem.* 2010;112:161–8.
- Eversole LR, Sabes WR, Rovin S. Aggressive growth and neoplastic potential of odontogenic cysts: with special reference to central epidermoid and mucoepidermoid carcinomas. *Cancer.* 1975;35:270–82.
- Bruner JM, Batsakis JG. Salivary neoplasms of the jaw bones with particular reference to central mucoepidermoid carcinomas. *Ann Otol Rhinol Laryngol.* 1991;100:954–5.
- Behboudi A, Enlund F, Winnes M, Andren Y, Nordkvist A, Leivo I, et al. Molecular classification of mucoepidermoid carcinomas—prognostic significance of the MECT1-MAML2 fusion oncogene. *Genes Chromosomes Cancer.* 2006;45:470–81.
- Chiose SI, Dacic S, Nikiforova MN, Seethala RR. Prospective testing of mucoepidermoid carcinoma for the MAML2 translocation: clinical implications. *Laryngoscope.* 2012;122:1690–4.
- Seethala RR, Dacic S, Cieply K, Kelly LM, Nikiforova MN. A reappraisal of the MECT1/MAML2 translocation in salivary mucoepidermoid carcinomas. *Am J Surg Pathol.* 2010;34:1106–21.
- Garcia JJ, Hunt JL, Weinreb I, McHugh JB, Barnes EL, Cieply K, et al. Fluorescence in situ hybridization for detection of MAML2 rearrangements in oncocytic mucoepidermoid carcinomas: utility as a diagnostic test. *Hum Pathol.* 2011;42:2001–9.
- Okabe M, Miyabe S, Nagatsuka H, Terada A, Hanai N, Yokoi M, et al. MECT1-MAML2 fusion transcript defines a favorable subset of mucoepidermoid carcinoma. *Clin Cancer Res.* 2006;12:3902–7.
- El-Naggar AK, Lovell M, Killary AM, Clayman GL, Batsakis JG. A mucoepidermoid carcinoma of minor salivary gland with t(11;19)(q21;p13.1) as the only karyotypic abnormality. *Cancer Genet Cytogenet.* 1996;87:29–33.
- Bell D, Holsinger CF, El-Naggar AK. CRTC1/MAML2 fusion transcript in central mucoepidermoid carcinoma of mandible—diagnostic and histogenetic implications. *Ann Diagn Pathol.* 2010;14:396–401.
- Khan HA, Loya A, Azhar R, Din NU, Bell D. Central mucoepidermoid carcinoma, a case report with molecular analysis of the TORC1/MAML2 gene fusion. *Head Neck Pathol.* 2010;4:261–4.
- Seethala RR, Dacic S, Cieply K, Kelly LM, Nikiforova MN. A reappraisal of the MECT1/MAML2 translocation in salivary mucoepidermoid carcinomas. *Am J Surg Pathol.* 2010;34:1106–21.
- Achcar Rde O, Nikiforova MN, Dacic S, Nicholson AG, Yousem SA. Mammalian mastermind like 2 11q21 gene rearrangement in bronchopulmonary mucoepidermoid carcinoma. *Hum Pathol.* 2009;40:854–60.
- Lennerz JK, Perry A, Mills JC, Huettner PC, Pfeifer JD. Mucoepidermoid carcinoma of the cervix: another tumor with the t(11;19)-associated CRTC1-MAML2 gene fusion. *Am J Surg Pathol.* 2009;33(6):835–43.
- Roden AC, Erickson-Johnson MR, Yi ES, Garcia JJ. Analysis of MAML2 rearrangement in mucoepidermoid carcinoma of the thymus. *Hum Pathol.* 2013;44:2799–805.
- Bell D, El-Naggar AK. Molecular heterogeneity in mucoepidermoid carcinoma: conceptual and practical implications. *Head Neck Pathol.* 2013;7:23–7.