

PAPER • OPEN ACCESS

## Global and national assessment of the incidence of asthma in children and adolescents from major sources of ambient NO<sub>2</sub>

To cite this article: Sourangsu Chowdhury *et al* 2021 *Environ. Res. Lett.* **16** 035020

View the [article online](#) for updates and enhancements.

You may also like

- [Oral health and halitosis among type 1 diabetic and healthy children](#)  
Tayyibe Aslihan Iscan, Cansu Ozsin-Ozler, Tulin Ileri-Keceli et al.
- [On the importance of developing a new generation of breath tests for \*Helicobacter pylori\* detection](#)  
levgeniia Kushch, Nikolai Korenev, Lyudmila Kamarchuk et al.
- [Associations of sex, age and adiposity in endothelium-independent dilation in children](#)  
Michelle M Harbin, Hanan Zavala, Justin R Ryder et al.

ENVIRONMENTAL RESEARCH  
LETTERS

## LETTER

Global and national assessment of the incidence of asthma in children and adolescents from major sources of ambient NO<sub>2</sub>

## OPEN ACCESS

## RECEIVED

28 October 2020

## REVISED

12 February 2021

## ACCEPTED FOR PUBLICATION

23 February 2021

## PUBLISHED

8 March 2021

Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Sourangsu Chowdhury<sup>1</sup>, Andy Haines<sup>2</sup>, Klaus Klingmüller<sup>1</sup>, Vinod Kumar<sup>3</sup>, Andrea Pozzer<sup>1</sup>, Chandra Venkataraman<sup>4</sup>, Christian Witt<sup>5</sup> and Jos Lelieveld<sup>1,6</sup>

<sup>1</sup> Department of Atmospheric Chemistry, Max Planck Institute for Chemistry, 55128 Mainz, Germany

<sup>2</sup> Centre on Climate Change and Planetary Health, Department of Public Health, Environments and Society and Department of Population Health, London School of Hygiene and Tropical Medicine, London WC1 9SH, United Kingdom

<sup>3</sup> Satellite Remote Sensing Division, Max Planck Institute for Chemistry, 55128 Mainz, Germany

<sup>4</sup> Interdisciplinary Programme in Climate Studies, and Department of Chemical Engineering, Indian Institute of Technology Bombay, Mumbai 400076, India

<sup>5</sup> Charité—University Medicine Berlin, 10117 Berlin, Germany

<sup>6</sup> Climate and Atmosphere Research Center, The Cyprus Institute, 1645 Nicosia, Cyprus

E-mail: [s.chowdhury@mpic.de](mailto:s.chowdhury@mpic.de)

**Keywords:** asthma, health impact assessment, children, NO<sub>2</sub>

Supplementary material for this article is available [online](#)

**Abstract**

Pediatric asthma incidence has been associated with exposure to nitrogen dioxide (NO<sub>2</sub>) in ambient air. NO<sub>2</sub> is predominantly emitted through fossil fuel use in land transportation, power generation and the burning of solid biofuels in households. We simulated NO<sub>2</sub> with a global atmospheric chemistry model, combined with a land use regression model, to estimate NO<sub>2</sub> exposure in all countries worldwide. The global asthma incidence among children and adolescents attributable to NO<sub>2</sub> was estimated by deriving an exposure-response function from a meta-analysis which included epidemiological studies from multiple countries, baseline incidence rates from the Global Burden of Disease and gridded population data. The sectoral contribution to pediatric asthma from NO<sub>2</sub> exposure (NO<sub>2</sub>-related asthma incidence: NINC) was estimated for different source categories to provide guidance to mitigation policies. We estimate 3.52 (2.1–6.0) million NINC per year globally, being about 14% of the total asthma incidence cases among children and adolescents. We find that emissions from land transportation are the leading contributor to NINC globally (~44%), followed by the domestic burning of solid fuels (~10.3%) and power generation from fossil fuels (~8.7%). Biogenic emissions which are not anthropogenically induced may contribute ~14% to the total NINC. Our results show large regional differences in source contributions, as the domestic burning of solid fuels is a main contributor to NINC in India and Nepal (~25%), while emissions from shipping are the leading source in Scandinavian countries (~40%), for example. While only 5% of all children and adolescents live in areas where NO<sub>2</sub> exceeds the WHO annual guideline of 21.25 ppb (40 μg m<sup>-3</sup>) for NO<sub>2</sub>, about 90% of the NINC is found in regions that meet the WHO guideline, related to the uneven distribution of children and adolescents in the population. This suggests the need for stricter policies to reduce NO<sub>2</sub> exposure, and revisiting the current WHO guideline to reduce the health risks of children and adolescents.

**1. Introduction**

Chronic respiratory diseases are among the leading contributors to morbidity worldwide (Soriano *et al* 2017, Stanaway *et al* 2018, Murray *et al* 2020). Asthma is a common respiratory condition leading to about 0.5 (95% confidence interval (CI) : 0.33–0.65) million

premature deaths per year, associated with the loss of 22 (18–28) million disability adjusted life years in 2017 among all age groups (Soriano *et al* 2017, Stanaway *et al* 2018). Although the premature mortality from asthma decreased by 17.5% from 1990 to 2017, asthma affected about 330 million people in 2016) and according to the Global Burden of Disease

(GBD) its prevalence increased by 30% in the same time period (Soriano *et al* 2017, Stanaway *et al* 2018). In low- and middle-income countries, where access to medical facilities and asthma medication is often limited, the increase is expected to continue in future (Foreman *et al* 2018, Global Asthma Network 2018).

Long term exposure to ambient air pollution not only exacerbates pre-existing asthma (Friedman *et al* 2001, Shima *et al* 2002, O'Connor *et al* 2008, Guarnieri and Balmes 2014, Brandt *et al* 2015), but is also a major driver of new asthma cases (Gauderman *et al* 2004, Jerrett *et al* 2008, Gehring *et al* 2013, Nishimura *et al* 2013, Guarnieri and Balmes 2014, Wijga *et al* 2014, Gehring *et al* 2020). Some studies indicate that air pollution increases the risk of the development of asthma during adulthood (Jacquemin *et al* 2009, Modig *et al* 2009), while infants (<5 years) are generally at greatest risk considering the immaturity of their lungs (USEPA 2008, 2016, Guarnieri and Balmes 2014). The human respiratory system develops fully in the first 18–20 years, and during this period there is a higher intrinsic risk for the development of asthma (USEPA 2016, Gehring *et al* 2020). The asthma incidence rate is about twice as high in young people ( $\leq 19$  years) than among adults ( $> 19$  years) (Soriano *et al* 2017, Stanaway *et al* 2018).

The GBD has not yet included the impact of air pollutants on asthma. Recent studies demonstrate associations between both long-term and short-term ambient NO<sub>2</sub> exposure and asthma incidence in children, supported by experimental studies that characterize a probable adverse effect of NO<sub>2</sub> on the respiratory system (Gauderman *et al* 2004, Gehring *et al* 2013, Kudo *et al* 2013, USEPA 2016, Global Asthma Network 2018, Stevens *et al* 2019). The biological pathway of NO<sub>2</sub> exposure leading to airway inflammation, airflow obstruction and new asthma development is also well established (Fukuto *et al* 2012, Kim *et al* 2013, USEPA 2016, Stevens *et al* 2019). After assessing the evidence from recent cohort studies on exposure to NO<sub>2</sub>, the 2016 integrated science assessment report by the United States Environmental Protection Agency (US EPA) changed the formulation of causal determination from long-term exposure to NO<sub>2</sub> on respiratory effects from ‘sufficient to infer a likely causal relationship’ in the 2008 Integrated Science Assessment report to a ‘causal relationship’ (USEPA 2008, p 201, 2016). The World Health Organization (WHO) sets the annual guideline for NO<sub>2</sub> at 40  $\mu\text{g m}^{-3}$  ( $\sim 21.25$  ppb at 25 °C temperature and 1 atmosphere pressure). However, this was proposed in the early 2000s when a robust basis for an annual average guideline for NO<sub>2</sub> through direct health effects was still lacking (WHO 2006).

Since the conversion between NO and NO<sub>2</sub> is rapid, typically they are considered as the tracer family NO<sub>x</sub> (Seinfeld and Pandis 2016). NO<sub>x</sub> is predominantly emitted as NO by anthropogenic

activities involving the burning of fossil fuels (about 10% is directly emitted as NO<sub>2</sub>), but it rapidly reacts with tropospheric ozone (O<sub>3</sub>) to form NO<sub>2</sub>. Previous health impact studies used satellite retrievals and land use regression (LUR) models to estimate the exposure to NO<sub>2</sub>, and assumed the incident cases of asthma to be linked to transportation related emissions (Anenberg *et al* 2018, Achakulwisut *et al* 2019). While studies have shown that transportation is a major source of NO<sub>x</sub>, other emission sectors like biomass burning, agricultural waste burning, agricultural soils, power generation, industries, shipping and domestic solid fuel use also emit significant amounts of NO<sub>x</sub> (Gerstle *et al* 1965, Jaegle *et al* 2005, Tian *et al* 2013, Boersma *et al* 2015, Janssens-Maenhout *et al* 2015, USEPA 2016, Ding *et al* 2017, Degraeuwe *et al* 2019). Here, we combine the outputs from an atmospheric chemistry model and an LUR model to estimate the burden of incident cases of asthma among children and adolescents from the exposure to ambient NO<sub>2</sub> and attribute it to different source categories. Our study adds to previous work (Anenberg *et al* 2018, Achakulwisut *et al* 2019, Khreis *et al* 2019) by using an atmospheric chemistry model to distinguish the contributions by the above-mentioned NO<sub>2</sub> sources, to guide policies aimed at regulating NO<sub>2</sub> exposure.

## 2. Methods

### 2.1. Model setup and emission inventory

We used the ECHAM5/MESy atmospheric chemistry (EMAC)–general circulation model at a T106 horizontal spectral resolution ( $1.12^\circ \times 1.12^\circ$  latitude  $\times$  longitude), with 31 vertical hybrid terrain-following and pressure levels up to 10 hPa in the lower stratosphere (Sander *et al* 2005, Kerkweg *et al* 2006, Jöckel *et al* 2010, De Meij *et al* 2012, Pozzer *et al* 2012, Lelieveld *et al* 2015, Joeckel *et al* 2016). The core atmospheric model is the 5th generation European Centre Hamburg (ECHAM5) general circulation model. EMAC includes sub-models that represent tropospheric and stratospheric processes and their interaction with oceans, land and human influences. We used the Modular Earth Submodel System (MESy, v.2.54, (Joeckel *et al* 2010)) to link submodels that describe emissions, atmospheric chemistry, aerosol and deposition processes. The gas phase and heterogeneous chemistry was simulated through the Module Efficiently Calculating the Chemistry of the Atmosphere (MECCA) submodel (Sander *et al* 2011, 2019), which accounts for the photochemical oxidation of natural and anthropogenic emissions, also accounting for the volatile organic compounds. The evolution of organic aerosol compounds is simulated with the ORACLE submodel (Tsimpidi *et al* 2014, 2018), which effectively accounts for primary and secondary combustion products from biofuel, fossil fuel, biomass burning

and their chemical oxidation during any atmospheric transport. Further, EMAC was nudged to simulate actual meteorological conditions, applying European Reanalysis Interim (ERA Interim) data for a period of 2 years (2014–2015), using 2014 as model spin-up.

We applied monthly varying sector-specific anthropogenic emissions to the model from the Community Emissions Data System (CEDS (Hoesly *et al* 2018)) at  $0.5^\circ \times 0.5^\circ$  resolution for the primary emitted species like sulfur dioxide (SO<sub>2</sub>), oxides of nitrogen (NO<sub>x</sub>), carbon monoxide (CO), black carbon (BC), organic carbon (OC), ammonia (NH<sub>3</sub>) and speciated non-methane volatile organic compounds. We improved the model emissions over India using region-specific and updated emission inventories (Venkataraman *et al* 2020). Biomass burning emissions were obtained from the Global Fire Assimilation System (GFAS, Kaiser *et al* 2012). The emissions data were then pre-processed by distributing them in six emission heights; please see Pozzer *et al* (2009) for further details on distribution of emission from different sources to the six emission heights. Ambient NO<sub>2</sub> exposure was first estimated for all sources and then all source categories were removed one by one to obtain the contribution of nine sectors to ambient NO<sub>2</sub> exposure (Lelieveld *et al* 2015, 2019a).

The sectors considered are (a) land transportation (TRA) which includes emissions from road transportation and railways, (b) industries (IND) with emissions from iron and steel, paper and pulp, chemical, food, solvent and other manufacturing, oil refineries and fuel production, (c) domestic energy use (DOM) by the burning of fuels in households for cooking, lighting, domestic gas heating, diesel generators and biofuel use, (d) power generation (PGN) by thermal, oil and gas based power plants, (e) agricultural soils (AGS) that cause emissions from manure and fertilizer application and cultivation practices, (f) agricultural waste and residue burning (AWB), (g) emissions from ships and other water navigation (SHP), (h) biomass burning (BMB) which includes tropical, savanna, middle and high latitude forest fires, deforestation, savanna and shrub fires and (i) biogenic emissions from soils and plants (BGN).

## 2.2. Estimating the burden of incident cases of asthma

A recent study performed an extensive review of epidemiological studies that reported impacts of long-term exposure to NO<sub>2</sub> on NINC in children and adolescents (Anenberg *et al* 2018). Among five meta-analyses (Gasana *et al* 2012, Takenoue *et al* 2012, Anderson *et al* 2013, Bowatte *et al* 2015, Khreis *et al* 2017) identified by the study, Khreis *et al* (2017) is the most recent and included the largest number of epidemiological studies. The exposure-response function (ERF) from Khreis *et al* (2017) was also used

by recent studies to estimate the total NINC in children (Achakulwisut *et al* 2019, Khreis *et al* 2019). Khreis *et al* (2017) included 41 epidemiological studies performed mostly in East Asia, Europe and North America and reported an relative risk (RR) of 1.26 (95% confidence interval (CI) 1.10–1.37) per 10 ppb increase in ambient NO<sub>2</sub>. The information from (Khreis *et al* 2017) was utilized here to build an ERF for impacts of NO<sub>2</sub> on NINC using the following relationship:

$$RR = \exp^{\beta \Delta x}, \quad \Delta x = \max[0, (NO_2 - LCC)]. \quad (1)$$

$\beta$  was first estimated by taking ln-RR of 1.12 (95% Uncertainty Interval: 1.08, 1.16) for a 10 ppb increase in NO<sub>2</sub>. The  $\beta$  obtained was then used to obtain the RR for exposure to NO<sub>2</sub> using a low-concentration cut-off (LCC: 0.8 ppb). The LCC was selected based on the lowest concentration at which health impact was recorded (Khreis *et al* 2017, 2019). We performed sensitivity studies with lower (0 ppb) and higher LCCs' (1 and 2 ppb). We evaluated the ERFs obtained from Khreis *et al* (2017) against those from the other four meta-analyses (Gasana *et al* 2012, Takenoue *et al* 2012, Anderson *et al* 2013, Bowatte *et al* 2015) and found that the estimates of NINCs from the ERFs drawn from all the five studies overlap within their uncertainty bounds (see: figures S1 and S12(b) (available online at [stacks.iop.org/ERL/16/035020/mmedia](https://stacks.iop.org/ERL/16/035020/mmedia))).

The burden of NINC among children and adolescents for 2015 was estimated using equation (2). A similar relation was applied in our earlier studies (Lelieveld *et al* 2019b, Chowdhury *et al* 2020) to estimate excess mortality

$$NINC = \left( \frac{RR - 1}{RR} \right) \times p \times BINC, \quad (2)$$

where  $p$  is the exposed population with age  $\leq 19$  years for 2015. The age profile of the population for 2015 was obtained from the GBD and was merged with population data at  $\sim 1$  km spatial resolution obtained from Global Human Settlement Layers (Florczyk *et al* 2019), resampled at  $5 \times 5$  km spatial resolution. Age-specific baseline incidence rates (BINC) for the age classes <5, 5–9, 10–14 and 15–19 for asthma were obtained for each country from the GBD (<https://vizhub.healthdata.org/gbd-compare/>). All estimates of NINC in this study are accompanied by 95% CIs. The 95% CI ranges were estimated by combining the uncertainty of the BINC obtained from GBD and the uncertainty in the RR curves estimated by 10 000 random samples over the log-normal distribution of baseline incidence rates and RR. 95% CI was obtained from the resulting 10 000 estimates of NINC for each grid. Subsequently, the results were integrated and are presented at country level.

### 3. Results and discussions

#### 3.1. Modeled NO<sub>2</sub> exposure and bias-correction

The EMAC model simulated global, gridded NO<sub>2</sub> at 15 min time intervals has been averaged, and the output stored at monthly time periods. The spatial distribution of NO<sub>2</sub><sup>EMAC</sup> is depicted in figure S2(a). The NO<sub>2</sub><sup>EMAC</sup> was then evaluated against annual averaged *in-situ* observation data for 2015 (NO<sub>2</sub><sup>OBS</sup>), which were collected from ~5000 sites around the world. The density of measurement locations is highest over Europe, East Asia and North America. NO<sub>2</sub><sup>EMAC</sup> was found to have a low-bias compared to ground-based observations NO<sub>2</sub><sup>OBS</sup> across the globe (figure S2(b)). The global population weighted NO<sub>2</sub><sup>EMAC</sup> was estimated at 4.3 ppb. The low-bias in NO<sub>2</sub><sup>EMAC</sup> may be due to: (a) its simulation at coarse horizontal resolution (1.12° × 1.12°) which averages out local gradients, this is especially relevant for urban areas and other locations with dense traffic, where mixing ratios are expected to be relatively high (below we will discuss the importance of traffic); (b), The lifetime of NO<sub>x</sub> (NO + NO<sub>2</sub>) is short, typically of the order of a few hours to a day, hence spatial-temporal gradients between sources and locations downwind can be steep, which are not captured well at the applied horizontal and time resolution. However, the EMAC tropospheric column densities of NO<sub>2</sub> compared well with those from the ozone monitoring instrument (Boersma *et al* 2018), indicating that sources and sinks are overall well represented. Please see supplementary information sections S1, figure S3 for more details on the comparison.


The bias towards low values was most notable over densely populated regions, whereas in background locations EMAC performs rather well (figures S4(a) and (b), detailed description on identification of background sites across the globe is provided in supplementary information sections S2). To amend the low-bias in densely populated areas, we utilized a publicly available LUR model (Larkin *et al* 2017) with which a high-resolution NO<sub>2</sub> dataset was derived, henceforth referred to as NO<sub>2</sub><sup>LUR</sup>. NO<sub>2</sub><sup>LUR</sup> was established for 2011 at 100 m resolution globally, by using information from 5220 monitoring sites located in 58 countries globally and predictor variables like active fires, population density, proximity to major and minor roads, tree cover, impervious surface area, normalized difference vegetation index, water body and satellite retrieved NO<sub>2</sub>. NO<sub>2</sub><sup>LUR</sup> was able to predict 54% of global variability, with a mean absolute error of 3.7 ppb (see Larkin *et al* (2017) for further details on formulation of NO<sub>2</sub><sup>LUR</sup>). The LUR model was shown to perform better compared to NO<sub>2</sub><sup>EMAC</sup> over the densely populated and urban areas, and was able to capture the spatial variability over such locations. However, it was found to overestimate NO<sub>2</sub> compared to observations over the background locations (figure S5), with a mean bias (MB) of -44%

for NO<sub>2</sub><sup>OBS</sup> compared to NO<sub>2</sub><sup>LUR</sup>. Also note the high values of NO<sub>2</sub><sup>LUR</sup> over the regions with sparse availability of predictor variables and low monitoring station density (e.g. Saudi Arabia, Saharan Africa, Midwest USA, central Australia; see figure S5(a)). Sparse availability of predictor variables also impacted the efficiency of NO<sub>2</sub><sup>LUR</sup> in biomass burning dominated regions of Central Africa (figure S5a).

To optimize the contributions of both NO<sub>2</sub><sup>EMAC</sup> and NO<sub>2</sub><sup>LUR</sup>, we combined the two data sets with information from the global human settlement layers settlement model (GHS-SMOD). The GHS-SMOD classifies each 1 km grid by population clusters into eight broad settlement classes: urban center, dense urban cluster, semi-dense urban cluster, suburban/peri-urban, rural cluster, low density rural, very low-density rural and water grid cells (see Florczyk *et al* (2019), Melchiorri *et al* (2018) for more details on GHSL-SMOD classifications). Merging of the data was performed by first resampling the NO<sub>2</sub><sup>LUR</sup> at 1 × 1 km resolution and then implementing NO<sub>2</sub><sup>EMAC</sup> over the background grids while using NO<sub>2</sub><sup>LUR</sup> over the other grids. The grid cells with the GHSL-SMOD classifications low-density rural and very low-density rural were considered as background (more details on background grid classification in supplementary material section S2). A sensitivity study was performed by also including rural cluster grid cells into the background grid. Subsequently, the merged NO<sub>2</sub> exposure data were weighted by population (of children and adolescents ≤ 19 years) to obtain NO<sub>2</sub><sup>F</sup> (NO<sub>2</sub><sup>EMAC</sup> fused ('F') with NO<sub>2</sub><sup>LUR</sup>) at 5 × 5 km resolution. Figures S6 and S7 depicts the algorithm for obtaining NO<sub>2</sub><sup>F</sup> by combining NO<sub>2</sub><sup>LUR</sup> and NO<sub>2</sub><sup>EMAC</sup> by summarizing globally and also zooming in over a city.

The global population weighted NO<sub>2</sub><sup>LUR</sup> was estimated at 9.8 ppb. Since 26% of the global child and adolescent population were estimated to reside in background grid locations (who were assigned NO<sub>2</sub><sup>EMAC</sup> for exposure), the global population weighted NO<sub>2</sub><sup>F</sup> was estimated to be lower at 8.1 ppb. The percentages of the population in all countries residing in background grid locations are depicted in figure S8(a). A large proportion of the child and adolescent population in Africa (31%) resides in background locations. In India, China, USA and Western Europe\* 26%, 27%, 30% and 27% of the total child and adolescent population reside in background locations. We estimated the global population weighted NO<sub>2</sub><sup>F</sup> exposure over background grid cells to be 2.9 ppb, compared to 6.3 ppb estimated with NO<sub>2</sub><sup>LUR</sup>. The population weighted NO<sub>2</sub><sup>F</sup> exposure at country-level is depicted in figure 1(a). We find that South Korea, Kuwait and Qatar have the

\* Countries considered in Western Europe: The Netherlands, Belgium, Germany, France, Austria, Portugal, Spain, Switzerland, Luxembourg, Italy, Republic of Ireland and the United Kingdom.



highest population weighted  $\text{NO}_2^{\text{F}}$  exposure at 18.6, 17.4 and 17.2 ppb, respectively. Population weighted  $\text{NO}_2^{\text{F}}$  exposure for China, the USA and India were estimated to be 12.5, 10.5 and 7.8 ppb, respectively. Figure 1(b) depicts the comparison between  $\text{NO}_2^{\text{OBS}}$  and  $\text{NO}_2^{\text{F}}$  across the world. The correlation coefficient for comparison between  $\text{NO}_2^{\text{OBS}}$  and  $\text{NO}_2^{\text{F}}$  ( $R = 0.66$ ) is higher compared to comparison with  $\text{NO}_2^{\text{LUR}}$  ( $R = 0.63$ ) and  $\text{NO}_2^{\text{EMAC}}$  ( $R = 0.43$ ).

A histogram of the pediatric population- $\text{NO}_2$  exposure curve was generated using the global gridded  $\text{NO}_2^{\text{F}}$  (figure 1(c)). The distribution was built by summing up the child and adolescent population in 500 logarithmically spaced bins from <1 ppb to 47 ppb and was found to have a distinct peak at 8 ppb. This feature is dominated by high exposure to  $\text{NO}_2^{\text{F}}$  among children and adolescents in India and China. The child and adolescent population- $\text{NO}_2$  exposure distribution of the five major countries are depicted in figure S9(a). We estimate that globally about 5% of all children and adolescents reside in areas where  $\text{NO}_2^{\text{F}}$  exceeds the WHO annual guideline level. These percentages are higher in South Korea (37%), Qatar (22%), Japan (16%), Egypt (12%) and China (10%) (figure S9(b)), while only ~2% of the children and adolescents in India and Western Europe live in areas where exposure exceeds the WHO guideline. We performed a sensitivity analysis by considering the GHSL-SMOD threshold to classify background grid cells by combining the rural cluster with the low density rural and very low-density rural grid cells as background locations and then merging  $\text{NO}_2^{\text{EMAC}}$

and  $\text{NO}_2^{\text{LUR}}$  to obtain  $\text{NO}_2^{\text{FR}}$ . About 36% of the global child and adolescent population lives in these (rural cluster, low density rural, very low-density rural) settlement grids (figure S8(b)). Global population weighted  $\text{NO}_2^{\text{FR}}$  was estimated to be 7.8 ppb. Population weighted  $\text{NO}_2^{\text{F}}$  was higher than  $\text{NO}_2^{\text{FR}}$  for all countries, and figure S8(c) depicts the percentage difference between population weighted  $\text{NO}_2^{\text{F}}$  and  $\text{NO}_2^{\text{FR}}$ .

### 3.2. Burden of asthma incidence

Globally, we attribute 3.52 (2.16–6.05) million  $\text{yr}^{-1}$  NINC among children and adolescents to the long-term exposure to  $\text{NO}_2$ , among which 56.5%, 18.1%, 16.4% and 10.8% occur among population of ages <5 years, 5–9 years, 10–14 years and 15–19 years, respectively. This amounts to ~13% of the total incidence. The NINC is highest in China, followed by India, Brazil, the USA and Indonesia, in that order, (figure 2, table 1). The numbers for the top nine countries (ranked according to NINC) plus Western Europe are listed in table 1. The supplementary data present this information for all countries worldwide considered in this study. The total burden of NINC depends on the size of the exposed child and adolescent population, BINC and  $\text{NO}_2^{\text{F}}$ . For example, India has the highest number of exposed children and adolescents (520 million), being 1.75 times that in China; however, the lower BINC in India (425 per 100 000 children and adolescent) compared to China (917 per 100 000) and significantly lower population weighted  $\text{NO}_2^{\text{F}}$  (38% lower) in India result in 47%




Figure 2. Incident cases of asthma among children and adolescents (NINC) attributed to exposure to ambient  $\text{NO}_2^{\text{F}}$ .

Table 1. Incident asthma cases attributed to ambient  $\text{NO}_2$  exposure in children, and the percentage contribution of major source categories in the countries with highest incidence plus Western Europe. Data of incident cases of asthma from exposure to  $\text{NO}_2^{\text{F}}$  by country are listed in SI DATA1.

Countries	NINC (95% CI) in million	Contribution from the major sectors (%)								
		TRA	IND	DOM	PGN	AGS	AWB	SHP	BMB	BGN
China	0.57 (0.33–0.96)	41.5	22.9	7.1	20.7	1.3	0.1	0.4	0.1	5.8
India	0.3 (0.17–0.52)	26.1	2.2	25.6	2.6	24.6	0.8	0.5	0.5	17
Brazil	0.2 (0.12–0.33)	53.4	7.5	2.3	4.6	3.3	0.2	6.8	5.3	16.6
United States of America	0.2 (0.12–0.3)	58.8	6.1	15.8	6.1	0	0	4.6	0.9	7.6
Indonesia	0.15 (0.09–0.24)	65.7	3.3	3	10.8	2.9	0	2	6.2	6.2
Mexico	0.13 (0.08–0.22)	73.4	3.3	3.6	2.7	1.8	0.1	1.9	2.1	11
Egypt	0.13 (0.08–0.22)	58.7	2.1	6.4	5.3	11.3	0.1	5.7	0	10.4
Pakistan	0.1 (0.06–0.17)	36.5	0.9	18.4	0.7	27.5	1.4	0.1	0	14.5
Nigeria	0.09 (0.06–0.18)	33	2.8	23.8	1.1	3.5	0.8	1.7	10	23.4
Western Europe	0.11 (0.06–0.17)	56.8	4	9	5.2	2	0	15.5	0.1	7.4


lower NINC. To adjust for the exposed population, the rate of NINC (per 100 000 children and adolescents) was calculated. The annual rate of NINC is highest in the Middle East, e.g. in the United Arab Emirates (UAE), Kuwait and Qatar [612 (412–1101), 564(367–992) and 446 (304–859) per 100 000 children and adolescent, respectively]. The spatial distribution is depicted in figure S10(a). In Brazil, the USA, China and India, the annual rate of NINC is estimated at 313 (187–513), 235 (144–359), 193 (113–324), and 57 (33–100) per 100 000, respectively. We estimate that in Kuwait 32% of all pediatric NINC may be attributed to  $\text{NO}_2$  exposure. The estimates for China, Brazil, India, United States of America and Western Europe are 22%, 14%, 13.8%, 19% and 18%, respectively (figure S10(b)). Using  $\text{NO}_2^{\text{FR}}$ , we attribute 3.33(2.06–5.78) million NINC to long-term  $\text{NO}_2$  exposure. Using  $\text{NO}_2^{\text{LUR}}$ , 4.07 (2.5–7.01) million NINC (15.5% higher compared to  $\text{NO}_2^{\text{F}}$ ) was estimated, which is close to the result (4 million) obtained by a recent study that used the LUR model (Achakulwisut *et al* 2019). Figure S11 depicts the percentage differences between NINC estimated using  $\text{NO}_2^{\text{LUR}}$  and  $\text{NO}_2^{\text{F}}$  and  $\text{NO}_2^{\text{FR}}$ . Large differences (up to 55% in the sub-Saharan countries) were found between NINC calculated with  $\text{NO}_2^{\text{LUR}}$  and  $\text{NO}_2^{\text{F}}$  in

the countries with a large fraction of background grids (see figure S8). For China, India, Brazil and the USA the difference was estimated at 3.8%, 12%, 12% and 13.5% respectively.

### 3.3. Source sector attribution

The impacts of the nine aforementioned source categories was determined by omitting them one by one from the total  $\text{NO}_x$  emissions in the global model results, following an earlier study (Lelieveld *et al* 2015). The relative difference between incident cases of asthma estimated by the simulation with total  $\text{NO}_x$  and the simulation without emissions from a sector determines the contribution by the sector. The source contribution in the nine countries with highest NINC burden and Western Europe are depicted in figure 3 and listed in table 1. Land transportation (TRA) is the single largest contributor to NINC globally, accounting for 44%, followed by BGN, DOM, PGN, IND, AGS, SHP, BMB and AWB with 14.1%, 10.3%, 8.7%, 7%, 6.6%, 4%, 5% and 0.4%, respectively. The source contributions for all countries are presented in the supplementary data.

TRA is the largest contributor to NINC in most countries in East Asia, Central Asia, Europe, North and South America (figure 4), associated with



1.56(0.95–3) million  $\text{yr}^{-1}$  NINCs. In Mexico and the USA, TRA contributes up to 73% and 59%, respectively. In Iran, Iraq, Saudi Arabia, Qatar and Kuwait, TRA is estimated to contribute to 60%–70% of the total NINC. We found it to be associated with 42%, 57% and 54% of NINC in China, Western Europe and Brazil, respectively. Its contribution is found to be lower in India (26%) and Nepal (29%) where the domestic use of solid fuels (DOM) is also a prevalent source of ambient  $\text{NO}_2$  at the surface.

DOM is associated with 0.37 (0.22–0.63) million  $\text{yr}^{-1}$  NINCs globally, of which almost 21% occur in India (0.08 (0.04–1.33) million  $\text{yr}^{-1}$ ). In Nepal, Pakistan and China, its contribution is 33.4%, 18.4% and 8%, respectively. DOM is also a major contributor to NINCs in Central African countries such as Rwanda (32%), Burundi (27%), Ethiopia (25%), Uganda (25%) and Nigeria (24%). In Argentina, its contribution to NINC is 42%. In high-income countries such as the USA (15%), Japan (11.4%) and Australia (3%) the DOM contribution is comparatively lower (figure 4). DOM contributes to 9% of NINC in Western Europe.


Power generation from fossil fuels (PGN) is a major source of  $\text{NO}_2$  as well (Tian *et al* 2013). It is associated with 0.3 (0.18–0.51) million  $\text{yr}^{-1}$  NINCs globally. In Russia, the Philippines and China it is a major contributor to NINC with 25%, 23% and 21%, respectively. Its role is considerably less in the USA (6.3%), Western Europe (5%), Brazil (4.6%) and India (2.6%). Emissions from fuel combustion in industries and industrial processes (IND) contribute comparatively less. With a global fraction of  $\sim 7\%$ , it is associated with 0.24 (0.15–0.41) million  $\text{yr}^{-1}$  NINCs. It is relatively important in Japan (12%), South Korea (16%) and China (17.8%) while in most western high-income countries its contribution is comparatively low (4–8%).

The application of manure and fertilizers to soils for agriculture (AGS) is the next largest contributor at 6.6% accounting for 0.23 (0.14–0.4) NINCs. Use of nitrogen-based fertilizers has increased over time and is largely driven by intensive farming practices. In South Asia (e.g. India, Pakistan, Nepal, Bhutan and Bangladesh) where intensive farming is predominant, AGS contributes  $\sim 20\%$  to the total NINC. In Europe and North America, the contribution is small (up to 5%). Agricultural waste and residue burning is a minor contributor to NINC with 0.01 (0.006–0.02) million  $\text{yr}^{-1}$ .

$\text{NO}_x$  from ship emissions (SHP) is expected to result in NINCs in coastal locations, major harbors and countries with coastlines adjacent to the major shipping routes. Globally, SHP is associated with 0.14 (0.09–0.23) million  $\text{yr}^{-1}$  NINCs. In Scandinavian countries it is a relatively large contributor to NINC, e.g. in Denmark (69%), Sweden (48%), Norway (44%) and Finland (20%). It also contributes significantly in island states, e.g. Iceland (65%), the UK (17%), Ireland (37%), Australia (16%) and New Zealand (14%).

Biomass burning, forest and grassland fires (BMB) account for 5% or 0.18 (0.11–0.31) million  $\text{yr}^{-1}$  NINCs globally. It plays a major role in sub-Saharan and particularly central African countries (figure 4) where it can contribute more than 60%. It contributes  $\sim 15\%$  in Southeast Asia (e.g. Cambodia, Laos, Indonesia). BMB is also important (6%–10%) in South America and Australia. BGN from natural vegetation and soil is associated with 0.49 (0.31–0.87) million NINCs. It was found to be a major contributor to NINCs in sub-Saharan, central and south African countries (12%–20%). In China, USA, Brazil and India it contributes about 6%, 7.5%, 16.7% and 16.9% respectively.





### 3.4. Uncertainties and assumptions

All estimates of NINC in this study are accompanied by 95% CIs. The 95% CI ranges were estimated by combining the uncertainty of the BINC and the uncertainty in the RR curves by distributing the RR and BINC lognormally across the 95% CIs to obtain 10 000 estimates of NINC for each grid cell, from which the overall 95% CIs were estimated (see Chowdhury *et al* (2020) for details). Several other limitations may affect the results presented in this study. Firstly, the model was simulated to estimate  $\text{NO}_2$  exposure at  $1.12^\circ \times 1.12^\circ$  horizontal resolution, which may be responsible for averaging out the local scale gradients in  $\text{NO}_2$  exposure (USEPA 2016, Larkin *et al* 2017, Achakulwisut *et al* 2019). Nevertheless, this was alleviated by merging the model simulations (of 2015) with fine resolution LUR model calculations of  $\text{NO}_2$  (of 2011). However, the source attribution was performed at the coarse model resolution and applied to the merged  $\text{NO}_2^{\text{F}}$  data, assuming only the steep  $\text{NO}_2$  concentration gradients in the vicinity of sources, e.g. in urban centers, and not the emission strengths of the sources, which are not well represented.

Secondly, the main results of this study were obtained by using an LCC of 0.8 ppb, following the lowest concentration at which causality was noted in a recent analysis (Khreis *et al* 2017). We performed sensitivity calculations by considering no threshold (i.e. LCC = 0 ppb), and threshold levels of 1 ppb and 2 ppb (based on Achakulwisut *et al* (2019)) (figure S12(a)). The estimates of NINC were 12% higher and 3% and 16% lower when LCC was changed to 0, 1 and 2 ppb respectively (figure S12(a)). We estimated 3.93 (2.41–6.75) million, 3.42 (2.09–5.87) million and 2.95 (1.8–5.06) million incident cases of asthma

from exposure to  $\text{NO}_2$  per year among children and adolescents using 0, 1 and 2 ppb as LCC, respectively, which corresponds to 14.5%, 12% and 10.9% of the total global incidence of asthma, respectively.

Thirdly, the Khreis *et al* (2017) meta-analysis included epidemiological studies from East Asia, Europe and North America, therefore missing out on epidemiological evidence from the rest of the world. Also, an ERF derived from a single-pollutant epidemiological model was used to assume a causal relationship between  $\text{NO}_2$  exposure and asthma incidence (because multipollutant models that adjust for other co-pollutants are currently not available). Nonetheless, the evidence is suggestive of more robust associations asthma incidence with  $\text{NO}_2$  exposure, than for other air pollutants like  $\text{PM}_{2.5}$  (Anderson *et al* 2013, USEPA 2016). However, it may be possible that the reported associations are sensitive to control for co-pollutants which may potentially change the estimates presented in this study. A sensitivity study was performed by estimating the NINC based on ERF curves from other available meta-analyses (figure S1). It was found that all the estimates of NINC overlap within the uncertainty bounds (figure S12(b)).

A recent study by Holst *et al* (2020) that inspected association of multiple pollutants on onset of asthma found that only  $\text{PM}_{2.5}$  remained significantly associated with asthma after controlling for other pollutants ( $\text{NO}_2$ ,  $\text{PM}_{10}$ ) among children of parents with asthma and mothers who smoked during pregnancy. However, after adjusting for parental asthma, parental income and parental education, a significant association was obtained for  $\text{NO}_2$  and new incidence of asthma among children. Another recent study (Gehring *et al* 2020) found stronger association of long-term exposure to  $\text{NO}_2$  and incidence of asthma

among children and adolescents compared to exposure to PM<sub>2.5</sub>. Nonetheless, we estimated the incident cases of asthma from long-term exposure to PM<sub>2.5</sub> (PINC) simulated by the EMAC model as a sensitivity study to our estimates of NINC (supplementary information section S3). Globally, we estimated PINC to be ~2 times higher than NINC, which is consistent with a previous study (Anenberg *et al* 2018). However we emphasize that, unlike for long-term exposure to NO<sub>2</sub>, there is no consensus among studies that associate exposure to long-term exposure to PM<sub>2.5</sub> with incidence of asthma (Gasana *et al* 2012, Anderson *et al* 2013, Guarneri and Balmes 2014, USEPA 2016). We reiterate that our estimates of NINC and PINC are not additive considering significant overlap between them due to the use of single-pollutant epidemiological models in this study in absence of models that control for co-pollutants. We refer to supplementary information section S3 for further discussion.

#### 4. Summary and conclusions

We estimate that 3.52 (1.2–7.2) million yr<sup>-1</sup> new incident cases of asthma in children and adolescents are attributable to exposure to ambient NO<sub>2</sub>. Our result is slightly lower than that of a recent study (Achakulwisut *et al* 2019), which used exposure obtained from an LUR model only, but is three times that from a prior study (Anenberg *et al* 2018). We combined results from a global LUR model for NO<sub>2</sub> with atmospheric chemistry model simulations to obtain the NO<sub>2</sub> exposure metric. We estimate that globally 5% of all children and adolescents live in areas where NO<sub>2</sub> exposure exceeds the WHO annual guideline of 21.25 ppb (the WHO guideline of 40 µg m<sup>-3</sup> was converted to ppb at standard temperature and pressure). However, ~90% of the total NINC burden occurs in locations where the WHO annual guideline is met. This calls for revisiting the WHO annual guideline (WHO 2006) for NO<sub>2</sub>. It was formulated in 2006 when a robust basis for the guideline from epidemiological studies was not yet available (WHO 2006, USEPA 2008). If the areas where the WHO guideline for NO<sub>2</sub> is currently exceeded would actually meet the guideline, ~2% [0.07 (0.04–0.11) million yr<sup>-1</sup>] of the NINC could be averted. Note that we do not account for the effect of NO<sub>2</sub> in causing exacerbations of symptoms in children and adolescents with pre-existing asthma, and hence we provide a lower limit of the full effect.

To help direct policy decision making, we provide the first estimates of source sector specific contributions of NO<sub>2</sub>-induced NINC, whereas previous work attributed the exposure generally to traffic related sources (Anenberg *et al* 2018, Achakulwisut *et al* 2019). Nevertheless, our results corroborate the suggestion that land transportation (TRA) is the

leading contributor to NINC, with a global average of 44%. In the USA and Mexico, the TRA contributions are very high, 58.8% and 73.4%, respectively. In Western Europe the contributions vary between 50% and 65%. In India and Nepal, where the burning of solid fuels in households is ubiquitous (Lelieveld *et al* 2015, Chowdhury *et al* 2019), this source sector is prevalent (~25%). In China the industrial sector and power generation from fossil fuels together contribute ~35% to NINC. In central Africa, biomass burning is the single largest contributor to NINC (>70%). Our results suggest that targeted air quality control measures and tightening of the WHO guideline for ambient NO<sub>2</sub> are needed to substantially reduce the burden of new asthma incident cases in children and adolescents.

#### Data availability statement

Data on incidence of asthma from exposure to NO<sub>2</sub> produced by this study are disseminated by country through SI Data1. Spatial raster of NO<sub>2</sub><sup>F</sup> and data of incidence of asthma from exposure to NO<sub>2</sub> by provinces will be made available upon reasonable request to the authors.

The data that support the findings of this study are available upon reasonable request from the authors.

#### Acknowledgments

The authors acknowledge Dr Perry Hystad, Oregon State University, for maintaining the repository of NO<sub>2</sub> land use regression model data (<https://health.oregonstate.edu/labs/spatial-health/resources-equipment>). VK acknowledges the Alexander von Humboldt foundation postdoctoral fellowship.

#### Conflict of interest

The authors declare no competing interests.

#### ORCID iDs

Sourangsu Chowdhury  <https://orcid.org/0000-0002-4600-2226>

Andy Haines  <https://orcid.org/0000-0002-8053-4605>

Klaus Klingmüller  <https://orcid.org/0000-0002-8425-8150>

Vinod Kumar  <https://orcid.org/0000-0002-8405-3470>

Andrea Pozzer  <https://orcid.org/0000-0003-2440-6104>

Chandra Venkataraman  <https://orcid.org/0000-0002-2280-3360>

Jos Lelieveld  <https://orcid.org/0000-0001-6307-3846>

## References

- Achakulwisut P, Brauer M, Hystad P and Anenberg S C 2019 Global, national, and urban burdens of paediatric asthma incidence attributable to ambient NO<sub>2</sub> pollution: estimates from global datasets *Lancet Planet. Health* **3** e166–78
- Anderson H R, Favarato G and Atkinson R W 2013 Long-term exposure to air pollution and the incidence of asthma: meta-analysis of cohort studies *Air Qual. Atmos. Health* **6** 47–56
- Anenberg S C et al 2018 Estimates of the global burden of ambient PM<sub>2.5</sub>, ozone, and NO<sub>2</sub> on asthma incidence and emergency room visits *Environ. Health Perspect.* **126** 1–14
- Boersma K F et al 2018 Improving algorithms and uncertainty estimates for satellite NO<sub>2</sub> retrievals: results from the quality assurance for the essential climate variables (QA4ECV) project *Atmos. Meas. Tech.* **11** 6651–78
- Boersma K F, Vinken G C M and Tournadre J 2015 Ships going slow in reducing their NO<sub>x</sub> emissions: changes in 2005–2012 ship exhaust inferred from satellite measurements over Europe *Environ. Res. Lett.* **10** 74007
- Bowatte G, Lodge C, Lowe A J, Erbas B, Perret J, Abramson M J, Matheson M and Dharmage S C 2015 The influence of childhood traffic-related air pollution exposure on asthma, allergy and sensitization: a systematic review and a meta-analysis of birth cohort studies *Allergy* **70** 245–56
- Brandt E B et al 2015 Exposure to allergen and diesel exhaust particles potentiates secondary allergen-specific memory responses, promoting asthma susceptibility *J. Allergy Clin. Immunol.* **136** 295–303.e7
- Chowdhury S, Dey S, Guttikunda S, Pillarisetti A, Smith K R and Di Girolamo L 2019 Indian annual ambient air quality standard is achievable by completely mitigating emissions from household sources *Proc. Natl Acad. Sci. USA* **116** 10711–6
- Chowdhury S, Pozzer A, Dey S, Klingmueller K and Lelieveld J 2020 Changing risk factors that contribute to premature mortality from ambient air pollution between 2000 and 2015 *Environ. Res. Lett.* **15** 074010
- De Meij A, Pozzer A, Pringle K J, Tost H and Lelieveld J 2012 EMAC model evaluation and analysis of atmospheric aerosol properties and distribution with a focus on the Mediterranean region *Atmos. Res.* **114–115** 38–69
- Degraeuwe B, Pisoni E, Peduzzi E, De Meij A, Monforti-Ferrario F, Bodis K, Mascherpa A, Astorga-Llorens C, Thunis P and Vignati E 2019 *Urban NO<sub>2</sub> Atlas UR 29943 EN* (Luxembourg: Publications Office of the European Union) (<https://ec.europa.eu/jrc/en/publication/euro-scientific-and-technical-research-reports/urban-no2-atlas>)
- Ding J, Miyazaki K, Van Der A R R, Mijling B, Kurokawa J I, Cho S Y, Janssens-Maenhout G, Zhang Q, Liu F and Felicitas Levelt P 2017 Intercomparison of NO<sub>x</sub> emission inventories over East Asia *Atmos. Chem. Phys.* **17** 10125–41
- Florczyk A J et al 2019 GHSL data package 2019 *EUR 29788 EN* (Luxembourg: Publications Office of the European Union) ([https://ghsl.jrc.ec.europa.eu/documents/GHSL\\_Data\\_Package\\_2019.pdf?t=1478q532234372](https://ghsl.jrc.ec.europa.eu/documents/GHSL_Data_Package_2019.pdf?t=1478q532234372))
- Foreman K J et al 2018 Forecasting life expectancy, years of life lost, and all-cause and cause-specific mortality for 250 causes of death: reference and alternative scenarios for 2016–40 for 195 countries and territories *Lancet* **392** 2052–90
- Friedman M S, Powell K E, Hutwagner L, Graham L M and Teague W G 2001 Impact of changes in transportation and commuting behaviors during the 1996 summer olympic games in Atlanta on air quality and childhood asthma *J. Am. Med. Assoc.* **285** 897–905
- Fukuto J M, Carrington S J, Tantillo D J, Harrison J G, Ignarro L J, Freeman B A, Chen A and Wink D A 2012 Small molecule signaling agents: the integrated chemistry and biochemistry of nitrogen oxides, oxides of carbon, dioxygen, hydrogen sulfide, and their derived species *Chem. Res. Toxicol.* **25** 769–93
- Gasana J, Dillikar D, Mendy A, Forno E and Ramos Vieira E 2012 Motor vehicle air pollution and asthma in children: a meta-analysis *Environ. Res.* **117** 36–45
- Gauderman W J et al 2004 The effect of air pollution on lung development from 10 to 18 years of age *N. Engl. J. Med.* **351** 1057–67
- Gehring U et al 2013 Air pollution exposure and lung function in children: the ESCAPE project *Environ. Health Perspect.* **121** 1357–64
- Gehring U, Wijga A H, Koppelman G H, Vonk J M, Smit H A and Brunekreef B 2020 Air pollution and the development of asthma from birth until young adulthood *Eur. Respir. J.* **56** 2000147
- Gerstle R W, Cuffe S T, Orning A A and Schwartz C H 1965 Air pollutant emissions from coal-fired power plants, report no. 2 *J. Air Pollut. Control Assoc.* **15** 59–64
- Global Asthma Network 2018 The global asthma report 2018 (Auckland: Global Asthma Network) ([www.globalasthmanetwork.org](http://www.globalasthmanetwork.org))
- Guarnieri M and Balmes J R 2014 Outdoor air pollution and asthma *Lancet* **383** 1581–92
- Hoesly R M et al 2018 Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS) *Geosci. Model Dev.* **11** 369–408
- Holst G J, Pedersen C B, Thygesen M, Brandt J, Geels C, Bønløkke J H and Sigsgaard T 2020 Air pollution and family related determinants of asthma onset and persistent wheezing in children: nationwide case-control study *BMJ* **370** m2791
- Jacquemin B et al 2009 Home outdoor NO<sub>2</sub> and new onset of self-reported asthma in adults *Epidemiology* **20** 119–26
- Jaegle L, Steinberger L, Martin R V and Chance K 2005 Global partitioning of NO<sub>x</sub> sources using satellite observations: relative roles of fossil fuel combustion, biomass burning and soil emissions *Faraday Discuss.* **130** 407–23
- Janssens-Maenhout G et al 2015 HTAP-v2.2: a mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution *Atmos. Chem. Phys.* **15** 11411–32
- Jerrett M et al 2008 Traffic-related air pollution and asthma onset in children: a prospective cohort study with individual exposure measurement *Environ. Health Perspect.* **116** 1433–8
- Jöckel P et al 2016 Earth system chemistry integrated modelling (ESCiMo) with the modular earth submodel system (MESSy) version 2.51 *Geosci. Model Dev.* **9** 1153–200
- Jöckel P, Kerkweg A, Pozzer A, Sander R, Tost H, Riede H, Baumgaertner A, Gromov S and Kern B 2010 Development cycle 2 of the modular earth submodel system (MESSy2) *Geosci. Model Dev.* **3** 717–52
- Kaiser J W et al 2012 Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power *Biogeosciences* **9** 527–54
- Kerkweg A, Sander R, Tost H and Jöckel P 2006 Technical note: implementation of prescribed (OFFLEM), calculated (ONLEM), and pseudo-emissions (TNUDGE) of chemical species in the modular earth submodel system (MESSy) *Atmos. Chem. Phys.* **6** 3603–9
- Khreis H, Cirach M, Mueller N, De Hoogh K, Hoek G, Nieuwenhuijsen M J and Rojas-Rueda D 2019 Outdoor air pollution and the burden of childhood asthma across Europe *Eur. Respir. J.* **54** 1802194
- Khreis H, Kelly C, Tate J, Parslow R, Lucas K and Nieuwenhuijsen M 2017 Exposure to traffic-related air pollution and risk of development of childhood asthma: a systematic review and meta-analysis *Environ. Int.* **100** 1–31
- Kim Y-M, Kim Y-S, Jeon S G and Kim Y-K 2013 Immunopathogenesis of allergic asthma: more than the Th<sub>2</sub> hypothesis *Allergy Asthma Immunol. Res.* **5** 189–96
- Kudo M, Ishigatsubo Y and Aoki I 2013 Pathology of asthma *Frontiers Microbiol.* **4** 1–16

- Larkin A, Geddes J A, Martin R V, Xiao Q, Liu Y, Marshall J D, Brauer M and Hystad P 2017 Global land use regression model for nitrogen dioxide air pollution *Environ. Sci. Technol.* **51** 6957–64
- Lelieveld J, Evans J S, Fnais M, Giannadaki D and Pozzer A 2015 The contribution of outdoor air pollution sources to premature mortality on a global scale *Nature* **525** 367–71
- Lelieveld J, Klingmueller K, Pozzer A, Burnett R T, Haines A and Ramanathan V 2019a Effects of fossil fuel and total anthropogenic emission removal on public health and climate *Proc. Natl Acad. Sci. USA* **116** 7192–7
- Lelieveld J, Klingmueller K, Pozzer A, Poeschl U, Fnais M, Daiber A and Muenzel T 2019b Cardiovascular disease burden from ambient air pollution in Europe reassessed using novel hazard ratio functions *Eur. Heart J.* **40** 1590–6
- Melchiorri M, Florczyk A J, Freire S, Schiavina M, Pesaresi M and Kemper T 2018 Unveiling 25 years of planetary urbanization with remote sensing: perspectives from the global human settlement layer *Remote Sens.* **10** 1–19
- Modig L, Torén K, Janson C, Jarvholm B and Forsberg B 2009 Vehicle exhaust outside the home and onset of asthma among adults *Eur. Respir. J.* **33** 1261–7
- Murray C J L et al 2020 Global burden of 87 risk factors in 204 countries and territories, 1990–2019: a systematic analysis for the global burden of disease study 2019 *Lancet* **396** 1223–49
- Nishimura K K et al 2013 Early-life air pollution and asthma risk in minority children: the GALA II and SAGE II studies *Am. J. Respir. Crit. Care Med.* **188** 309–18
- O'Connor G T et al 2008 Acute respiratory health effects of air pollution on children with asthma in US inner cities *J. Allergy Clin. Immunol.* **121** 1133–1139.e1
- Pozzer A, De Meij A, Pringle K J, Tost H, Doering U M, Van Aardenne J and Lelieveld J 2012 Distributions and regional budgets of aerosols and their precursors simulated with the EMAC chemistry-climate model *Atmos. Chem. Phys.* **12** 961–87
- Pozzer A, Jöckel P and Van Aardenne J 2009 The influence of the vertical distribution of emissions on tropospheric chemistry *Atmos. Chem. Phys.* **9** 9417–32
- Sander R et al 2011 The atmospheric chemistry box model CAABA/MECCA-3.0 *Geosci. Model Dev.* **4** 373–80
- Sander R et al 2019 The community atmospheric chemistry box model CAABA/MECCA-4.0 *Geosci. Model Dev.* **12** 1365–85
- Sander R, Kerkweg A, Jöckel P and Lelieveld J 2005 Technical note: the new comprehensive atmospheric chemistry module MECCA *Atmos. Chem. Phys.* **5** 445–50
- Seinfeld J H and Pandis S N 2016 *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change* 3rd edn (New York: Wiley)
- Shima M, Nitta Y, Ando M and Adachi M 2002 Effects of air pollution on the prevalence and incidence of asthma in children *Arch. Environ. Health* **57** 529–35
- Soriano J B et al 2017 Global, regional, and national deaths, prevalence, disability-adjusted life years, and years lived with disability for chronic obstructive pulmonary disease and asthma, 1990–2015: a systematic analysis for the global burden of disease study 2015 *Lancet Respir. Med.* **5** 691–706
- Stanaway J D et al 2018 Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the global burden of disease study 2017 *Lancet* **392** 1923–94
- Stevens E L, Rosser F, Forno E, Peden D and Celedón J C 2019 Can the effects of outdoor air pollution on asthma be mitigated? *J. Allergy Clin. Immunol.* **143** 2016–2018.e1
- Takenoue Y, Kaneko T, Miyamae T, Mori M and Yokota S 2012 Influence of outdoor NO<sub>2</sub> exposure on asthma in childhood: meta-analysis *Pediatr. Int.* **54** 762–9
- Tian H, Liu K, Hao J, Wang Y, Gao J, Qiu P and Zhu C 2013 Nitrogen oxides emissions from thermal power plants in china: current status and future predictions *Environ. Sci. Technol.* **47** 11350–7
- Tsimpidi A P, Karydis V A, Pozzer A, Pandis S N and Lelieveld J 2014 ORACLE (v1.0): module to simulate the organic aerosol composition and evolution in the atmosphere *Geosci. Model Dev.* **7** 3153–72
- Tsimpidi A P, Karydis V A, Pozzer A, Pandis S N and Lelieveld J 2018 ORACLE 2D (v2.0): an efficient module to compute the volatility and oxygen content of organic aerosol with a global chemistry–climate model *Geosci. Model Dev.* **11** 3369–89
- USEPA 2008 Integrated science assessment for oxides of nitrogen and sulfur–environmental criteria *Dec. 2008 EPA600R-08082F* **73** 75716–7 ([https://hero.epa.gov/index.cfm/reference/details/reference\\_id/157073](https://hero.epa.gov/index.cfm/reference/details/reference_id/157073))
- USEPA 2016 Integrated science assessment for oxides of nitrogen–health criteria (final report) *Wash. DC Contract No EPA600R-08071* (<https://www.epa.gov/isa/integrated-science-assessment-isa-nitrogen-dioxide-health-criteria>)
- Venkataraman C, Bhushan M, Dey S, Ganguly D, Gupta T, Habib G, Kesarkar A, Phuleria H and Raman R S 2020 Indian network project on carbonaceous aerosol emissions, source apportionment and climate impacts (COALESCE) *Bull. Am. Meteorol. Soc.* **101** E1052–68
- WHO 2006 WHO Air quality guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide: global update 2005: summary of risk assessment (Geneva: World Health Organization) pp 1–22
- Wijga A H et al 2014 Cohort profile: the prevention and incidence of asthma and Mite Allergy (PIAMA) birth cohort *Int. J. Epidemiol.* **43** 527–35