Faculty of Science and Engineering

School of Biological and Marine Sciences

2013-02-26

Global environmental changes: setting priorities for Latin American coastal habitats.

Turra, A

http://hdl.handle.net/10026.1/1463

10.1111/gcb.12186 Glob Chang Biol

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

COMMENTARY

Global environmental changes: setting priorities for Latin American coastal habitats

ALEXANDER TURRA¹, ALDO CR ÓQUER², ALVAR CARRANZA³, ANDR ÉS MANSILLA⁴, ARSENIO J. ARECES⁵, CAMILO WERLINGER⁶, CARLOS MART ÍNEZ-BAY ÓN⁵, CRISTINA A.G. NASSAR⁷, ESTELA PLASTINO⁸, EVANGELINA SCHWINDT⁹, FABRIZIO SCARABINO¹⁰, FUNGYI CHOW⁸, FÉLIX L. FIGUEROA¹¹, FL ÁVIO BERCHEZ⁸, JASON M. HALL-SPENCER¹², LUIS A. SOTO¹³, MARCOS S. BUCKERIDGE⁸, MARGARETH S. COPERTINO¹⁴, MARIA TEREZA SZ ÉCHY¹⁵, NATALIA GHILARDI-LOPES¹⁶, PAULO HORTA¹⁷, RICARDO COUTINHO¹⁸, SIMONETTA FRASCHETTI¹⁹ and ZELINDA M. A. N. LEÃO²⁰

¹Departamento de Oceanografia Biológica, Instituto Oceanográfico, Universiade de São Paulo, São Paulo, Brazil, ²Departamento de Estudios Ambientales, Simon Bolivar University, Sartenejas, ³Polo de Desarrollo Universitario 'Grupo de Investigación y formación de recursos humanos en biodiversidad' Centro Universitario Regional Este – CURE, Universidad de la República Oriental del Uruguay, Sede Maldonado, Uruguay, ⁴Laboratorio de Macroalgas Antárticas y Subantárticas, Universidad de Magallanes and Instituto de Ecología y Biodiversidad, Punta Arenas Chile, ⁵Instituto de Oceanologia, Universidade de La Habana, Cuba, ⁶Facultad de Ciencias Naturales y Oceanograficas, Departamento de Oceanografia, Universidad de Concepcion, Concepcion, Chile, ⁷Departamento de Botânica, Instituto de Biologia, Universidade Federal do Rio de Janeiro, Brazil, ⁸Departamento de Botânica, Instituto de Biociências, Universidade de São Paulo, São Paulo, Brazil, ⁹Grupo de Ecología en Ambientes Costeros, Centro Nacional Patagonico, Puerto Madryn, Argentina, ¹⁰Dirección Nacional de Recursos Acuático and Museo Nacional de Historia Natural, Montevideo, Uruguay, ¹¹Ecology Department, Faculty of Sciences, Málaga University, Málaga, Spain, ¹²Marine Biology and Ecology Research Centre, University of Plymouth, Plymouth, UK, ¹³Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, México, ¹⁴Instituto de Oceanografia Universidade Federal do Rio Grande (FURG) Rio Grande, Brazil, ¹⁵Departamento de Botânica, Universidade Federal do Rio de Janeiro, Brazil, ¹⁶Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, ¹⁷Departamento de Botânica, Centro de Ciências Biológicas, Universidade Federal de Santa Catarina, Brazil, ¹⁸Instituto de Estudos do Mar Almirante Paulo Moreira, Arraial do Cabo, Brazil, ¹⁹Laboratory of Marine Biology, Department of Biological and Environmental Science and Technologies, University of Salento, ²⁰Departamento de 1 Sedimentologia, Instituto de Geociências, Universidade Federal da Bahia, Brazil

Abstract

As the effects of the Global Climate Changes on the costal regions of Central and South Americas advance, there is proportionally little research being made to understand such impacts. This commentary puts forward a series of propositions of strategies to improve performance of Central and South American science and policy making in order to cope with the future impacts of the Global Climate Changes in their coastal habitats.

Keywords: benthic ecology, climate impacts, habitat mapping, long-term monitoring, marine biodiversity

Received 29 August 2012 and accepted 13 February 2013

The need for a science-policy agenda in Central and South America

The Intergovernmental Panel for Climate Change (IPCC) reports that Global Environmental Changes (GEC) are occurring quicker than at any other time over the last 25 million years and impacting upon marine environments (Bellard *et al.*, 2012). There is overwhelming evidence showing that GEC are affecting both

Correspondence: Marcos S. Buckeridge, tel. ???, fax ???, 2 e-mail: msbuck@usp.br the quality and quantity of the goods and services provided by a wide range of marine ecosystems.

To discuss regional preparedness for global environmental changes, a workshop was held in Ilhabela, Brazil (22–26 April 2012) entitled 'Evaluating the Sensitivity of Central and South American Benthic Communities to Global Environmental Changes' that drew together scientists from ten Latin American and three European countries. Our analysis revealed critical knowledge gaps that hinder policy-making and assessments for the forthcoming IPCC Report (AR5, 2013–2014). We developed key recommendations on how to foster the

© 2013 Blackwell Publishing Ltd

	G	С	B	1	2	1	8	6	R	Dispatch: 6.3.13	Journal: GCB	CE: Sangeetha
Journal Name				Manuscript No.					D	Author Received:	No. of pages: 5	PE: Eswari

development of a regional science-policy agenda to meet urgent demand for sound scientific advice in the face of rapid changes to marine coastal ecosystems in Latin America.

Threats to ecologically and socio-economically important coastal habitats in Latin America

Central and South America is the home of 1/3 of the world's most biodiverse countries, and is one of the most urbanized regions in the world (Unep, 2011). Besides regional heterogeneity, and significant variation in size and economic development, the 33 countries of the region have relatively young democracies that face a number of common political, social-economic, environmental, and science-policy issues. The marine habitats are of fundamental importance for the approximately 610 million coastal residents, but the need to develop sustainable coastal management occurs at a time of rapidly changing climate coupled with social upheaval such as uncontrolled urbanization and social inequality.

Latin America marine realms include a wide range of benthic ecosystems, many of which are unique and constitute hotspots of biodiversity (Miloslavich *et al.*, 2011). These include the kelp forests on the Cape Horn Biosphere Reserve (Fig. 1; Rozzi *et al.*, 2012), the huge rhodolith beds along the Tropical Southwestern Atlantic coast (Berchez *et al.*, 2009; Amado-Filho *et al.*, 2012), the large blue carbon ecosystems, formed by tropical mangroves and seagrass beds (Copertino, 2011)and the highly biodiverse coral reefs of the Tropical Atlantic, with their large number of endemic species (Leão *et al.*, 2003). Therefore, major efforts to protect these marine




Fig. 1 A kelp forest in the Cape Horn Biosphere Reserve, Chile. This habitat is extremely important as a CO₂ sink and for fisheries. Centolla crab *Lithodes santolla* growing on *Macrocystis pyrifera* at the Capitan Aracena Island, Magallanes (Photo: Mathias Hüne).

habitats are essential. Fundamental regional economic activities, such as fisheries, with the world's highest average annual growth in the period 1970–2008 (21.1% as reported by Salas *et al.*, 2011), and tourism, with an 8.6–13.9% total contribution to gross domestic product (Wttc, 2012), depend on marine environmental quality.

Multiple human impacts endanger Latin America coastal habitats. Changes in the composition and distribution of sensitive habitats are already occurring (Martins et al., 2012), with highly impacted sites in the Eastern Caribbean, and medium to highly impacted zones around almost the entire continent (Halpern et al., 2008). Without timely action the situation will steadily deteriorate. Bleaching and diseases in coral reefs (Fig. 2), both linked to ocean warming, are becoming an increasing problem (Wilkinson & Souter, 2008). Kelp forests have proven to be highly susceptible to temperature and current changes (Wernberg et al., 2011) and ocean acidification not only threatens to degrade the world's largest rhodolith beds along the Brazilian coast (Amado-Filho et al., 2012) but also to seriously reduce the ability of edible shellfish, such as mussels and oysters, to produce shells, thereby threatening local aquaculture activities and food security. Extreme events, such as cyclones, are occurring with greater frequency (Emanuel, 2005), thereby impacting coastal habitats, with particular severity in the SE Atlantic coast. Moreover, harmful algal blooms, partially related to temperature increase, have negative impacts on the quality of coastal areas as a whole.

Gaps in scientific knowledge

Concerted efforts to understand the effects of GEC on Latin America coastal habitats lag behind other regions



Fig. 2 A coral bleaching event. Bleaching events in the Caribbean Sea are becoming more frequent and severe (Photo: Aldo Croquer).

worldwide, leaving society ill-prepared to cope with future changes. The paucity of time-series data in the southern hemisphere is especially acute in developing countries (Rosenzweig *et al.*, 2008). Less than 5% of the participants in the Second International Symposium on the Effects of Climate Change on the World's Oceans (Korea, May 2012) were from C&SA, exemplifying the low priority afforded to the issue in the regional scientific agenda. In short, baseline, monitoring and detailed forecast studies are insufficient for a specific understanding of detrimental GEC effects in the region. This has arisen due to a lack of scientific incentives and a dearth of efforts at the science-policy interface across the entire Latin America region.

Baselines

18

Integrated baseline studies are required to assess seabed-habitat distribution and quality, as well as human threats and risks associated with local and regional climate change scenarios. National support, within a multinational strategy, will be essential for systematic habitat mapping that should include geomorphological and ecological features at different spatial scales, using standardized approaches, to facilitate spatial and temporal comparisons, as well as the organization and dissemination of information. This will allow identification of biodiversity hot-spots, habitats of high value in terms of ecosystem services, and areas most vulnerable and less resilient to local anthropogenic impacts and GEC. It is imperative to take into account the potential synergies deriving from the interaction of multistressors, as the effects of GEC will differ according to the different combinations of threats. This information would also be important as a base for marine spatial planning strategies. For this issue, efforts should be targeted to the less studied ecosystems and regions, such as the Cape Horn Biosphere Reserve and the Brazilian rhodolith beds.

Monitoring

A strategic array of physical and biological monitoring stations is an urgent requirement in C&SA, to fill critical knowledge-gaps, and provide an early warning system of GEC on coastal communities. The systematic application of monitoring protocols to each habitat, scale, and level of organization, as well as to the various oceanographic conditions, is essential for documenting habitat degradation, carbon sinks, the reduction of primary and secondary production, and habitat destruction, fragmentation or loss, as well as biological invasion, and regime shifts. Thus, support for longterm time-series data collection through national and international networks is required using rigorous standardized protocols. The Monitoring Network for Coastal Benthic Habitats (ReBentos), to date one of the main extensive networks implemented in Latin America for monitoring marine habitats, groups together around 100 researchers starting to apply standardized protocols to both soft and hard substrata, viz., rocky shores, coral reefs, rhodolith beds, mangroves, salt marshes, estuaries, and sandy beaches, at stations distributed all along the Brazilian Coast. The South American Research Group for Coastal Ecosystems (SARCE), comprising 108 sampling localities, is another example. Efforts should be made to spread these tried and tested schemes to other countries in the region, through combining procedures and efforts with local projects already under way, and building an open access data-base to provide information for local, regional, and global habitat health evaluation and forecasts, this including the Regular Process of the United Nations for Global Reporting and Assessment of the State of the Marine Environment. Initial efforts should be centered on locations already undergoing immediate damaging pressures, such as the Caribbean Coral Reefs, the SE Atlantic rocky shores, or the southernmost kelp forests.

Forecasts

The absence of baseline studies seriously compromises reliable forecasting in Latin America. There is an urgent need to refining regional and local scenarios of threats related to GEC, to assess the uncertainties, risks, and thresholds at organism and ecosystem levels. Not only the identification and quantification of carbon sinks and cycling processes but also experimental and modeling approaches, are key challenges to forecasting future changes.

A scientific-support policy

The challenges are so great that collaborative efforts among institutions at national and international levels are essential. Most of the present initiatives in Latin America are national, for example, Brazilian Network for Blobal Climate Changes (Rede CLIMA), or bilateral, for example, the CNPq–CONICET Brazil & Argentina funding support. Efforts should be centered on networking the knowledge-base across disciplines and among all Latin American countries. Besides strengthening the support of national science funding agencies to studies focused on GEC, multilateral international agreements are also required. The incentive of capacitybuilding efforts at undergraduate and graduate levels, and of habitat mapping and the evaluation of GEC effects, is mandatory, as is stimulation of the formation and recruitment of interdisciplinary capacities in marine and human sciences, and technology, at a continental level. There is also a need to stimulate a better and wider communication of GEC to society as a whole, through innovative educational approaches and efficient scientific-outreach efforts, with focus on the Latin American marine environments, which, by leading to greater public involvement, could thus increase political interest.

Political and governance issues

Urgency requires the immediate establishment of a collaborative framework, to so induce a systematic and integrated spatial planning process for the sustainable use of marine biodiversity and other resources in C&SA. This would include joint efforts to identify and give precedence to the most pressing issues related to GEC and coastal habitats. There is the need for a more pro-active engagement of Latin American governments and sector-ministries, as well as the recruitment of socio-economic stakeholders, in a co-management effort regarding GEC and sustainable development, with a more evident focus on the sea. The delimitation of marine protected areas to reach the 10% goals established during the COP-10 - Convention of Biological Diversity - is a priority. Even in Brazil, the most protected country of the region (Halpern et al., 2012), only 1.5% of the exclusive economic zone is protected and nearly 9% of priority areas for marine conservation have already been ceded to oil companies for offshore exploitation (Scarano et al., 2012). The establishment of national councils, such as the Brazilian Inter-ministry Commission for Marine Resources, or processes, such as the National Science, Technology and Innovation Conferences in Brazil, could be considered as models to be followed. Initiatives should emerge as political efforts, under the responsibility of those countries already undertaking successful experiences, or those in better economic conditions, such as Argentina, Brazil, Chile, and Mexico. International articulation efforts should be reinforced by establishing formal mandates and securing resources for leadership institutions and initiatives, such as the Intergovernmental Oceanographic Commission at UNESCO, to consolidate South-to-North, and significantly increase South-to-South collaboration, thereby also benefiting other areas, such as Africa, India, and Southeast Asia.

Science and policy-making under an integrated perspective

Nations should improve communication between policy makers and scientists, to the point that new policies can

be based on the best available evidence, and scientific studies widened to include the most policy-relevant questions. The UN Regular Process, IPCC, IPBES (Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services), and the Future Earth initiatives should be considered as opportunities for the international integration of this agenda. Finally, once both IPBES and IPCC undertake the regular and timely assessment of knowledge, thereby identifying and giving precedence to the key scientific information needed for policymakers, priority on issues regarding climate change in the IPBES agenda, and those on marine habitats, biodiversity, and ecosystem services in the IPCC, becomes mandatory.

Operational agenda for the near future

As it is impossible to address all the issues simultaneously, multicriteria analysis becomes necessary. This would include the survey and analysis of existing data, to thus facilitate the identification of priorities for urgent action. Our main recommendations include sensitivity analysis on an eco-regional scale, thence addressing vulnerability to GEC on a habitat basis. This would include ecological (e.g., geographical distribution and associated biodiversity) and socio-economic (e.g., the economic evaluation of ecosystem goods and services) aspects. Local forecasting, based on the downscaling of available GEC scenarios, geographical distribution, conservation status, and the likely response of different habitats in different eco-regions, as well as the evaluation of potential ecological and socio-economic impacts should be a first step. Once having established the geographical scope of priority issues, both scientists and policy-makers should work together, by searching for the most effective governance setting to design and implement adaptation and/or mitigation schedules, either at international, national, or local levels.

Acknowledgements

Supported by FAPESP (2011/22074-0) and ReBentos (CNPq/ FAPESP). English revision by Ramon Arthur Clark. Authors thank the comments on the manuscript made by the colleague Fabio Scarano (The Nature Conservancy).

References

- Amado-Filho GM, Moura RL, Bastos AC et al. (2012) Rhodolith beds are major CaCO3 bio-factories in the tropical South West Atlantic. PLoS ONE, 7, e35171.
- Bellard C, Bertelsmeier C, Leadley P, Thuiller W, Courchamp F (2012) Effects of climate change on the future of biodiversity. *Ecology Letters*, 15, 365–377.
- Berchez FAS, Tiago CG, Rosso S, Dias G, Oliveira EC (2009) Structure of a coralline algal bed on southeastern Brazil. Brazilian Journal of Ecology, 13, 49–57.
- Copertino MS (2011) Add coastal vegetation to the climate critical list. *Nature*, **473**, 255.

Emanuel K (2005) Increasing destructiveness of tropical cyclones over the past 30 years. *Nature*, **436**, 686–688.

Halpern BS, Walbridge S, Selkoe KA et al. (2008) Global map of human impact on marine ecosystems. Science, 319, 948–952.

Halpern BS, Longo C, Hardy D et al. (2012) An index to assess the health and benefits of the global ocean. Nature, 488, 615–620.

- Leão ZMAN, Kikuchi RKP, Testa V (2003) Corals and coral reefs of Brazil. Latin American Coral Reefs, 13, 9–52.
- Martins CDL, Arantes N, Faveri C et al. (2012) The impact of coastal urbanization on the structure of phytobenthic communities in southern Brazil. Marine Pollution Bulletin, 64, 772–778.
- Miloslavich P, Klein E, Díaz JM et al. (2011) Marine biodiversity in the Atlantic and Pacific Coasts of South America: knowledge and gaps. PLoS ONE, 6, e14631.
- Rosenzweig C, Karoly D, Vicarelli M et al. (2008) Attributing physical and biological impacts to anthropogenic climate change. Nature, 453, 353–357.

- Rozzi R, Armesto JJ, Gutiérrez JR et al. (2012) Integrating ecology and environmental ethics: Earth stewardship in the southern end of the Americas. BioScience, 62, 226–236.
- Salas S, Chuenpagdee R, Charles A, Seijo JC (eds) (2011) Coastal Fisheries of Latin America and the Caribbean, ???, Rome.
- Scarano F, Guimarães A, Silva JMD (2012) Rio+20: Lead by example. Nature, 486, 25-26.
- Unep (ed.) (2011) Annual Report 2010: A Year in Review. United Nations Environment Programme, Nairobi.
- Wernberg T, Russell BD, Moore PJ et al. (2011) Impacts of climate change in a global hotspot for temperate marine biodiversity and ocean warming. Journal of Experimental Marine Biology and Ecology, 400, 7–16.
- Wilkinson C, Souter D (eds.) (2008) Status of Caribbean Coral Reefs After Bleaching and Hurricanes in 2005. GCRMN, Townsville.
- Wttc (ed.) (2012) Travel & Tourism Economic Impact 2012–Caribbean, World Travel & Tourism Council, London.

53

Author Query Form

Journal: GCB Article: 12186

Dear Author,

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers on the query sheet if there is insufficient space on the page proofs. Please write clearly and follow the conventions shown on the attached corrections sheet. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication.


Many thanks for your assistance.

Query reference	Query	Remarks
1	AUTHOR: Please provide full address details for all affiliations.	
2	AUTHOR: Please provide telephone number and fax number for corresponding author.	
3	AUTHOR: Please provide the publisher name for reference Salas et al. (2011).	

WILEY-BLACKWELL

USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION

Required software to e-Annotate PDFs: Adobe Acrobat Professional or Adobe Reader (version 7.0 or above). (Note that this document uses screenshots from Adobe Reader X) The latest version of Acrobat Reader can be downloaded for free at: http://get.adobe.com/uk/reader/



be

nc

Blanchard and Kiyotaki (1987), erfect competition in general equilib ts of aggregate demand and supply lassical framework assuming monop een an evogenous number of firms

3. Add note to text Tool – for highlighting a section to be changed to bold or italic.

important works on energy of onire

M henceforth)¹ we open the 'black h



Highlights text in yellow and opens up a text box where comments can be entered.

How to use it

nain compo

level, are exc

- Highlight the relevant section of text.
- Click on the Add note to text icon in the Annotations section.


4. Add sticky note Tool – for making notes at specific points in the text.



Marks a point in the proof where a comment needs to be highlighted.


How to use it

- Click on the Add sticky note icon in the Annotations section.
- Click at the point in the proof where the comment should be inserted
- Type instruction on what should be changed regarding the text into the yellow box that appears.




- Type the comment into the yellow box that appears.

iand and supply snocks. Most of



WILEY-BLACKWELL


USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION






How to use it

- Click on one of the shapes in the Drawing Markups section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, move the cursor over the shape until an arrowhead appears.
- Double click on the shape and type any text in the red box that appears.



Allows shapes, lines and freeform annotations to be drawn on proofs and for comment to be made on these marks..



For further information on how to annotate proofs, click on the Help menu to reveal a list of further options:

