



Global Flood Exposure from Different Sized Rivers

Mark V Bernhofen¹, Mark A Trigg¹, P Andrew Sleigh¹, Christopher C Sampson², Andrew M Smith²

¹School of Civil Engineering, University of Leeds, LS2 9JT, United Kingdom

²Fathom, Square Works, 17-18 Berkeley Square, BS8 1HB, United Kingdom

5 *Correspondence to:* Mark V Bernhofen (cn13mvp@leeds.ac.uk)

Abstract. There is now a wealth of data to calculate global flood exposure. Available datasets differ in detail and representation of both global population distribution and global flood hazard. Previous studies of global flood risk have used datasets interchangeably without addressing the impacts using different datasets could have on exposure estimates. By calculating flood exposure to different sized rivers using a model independent geomorphological approach, we show that
10 limits placed on the size of river represented in global flood models result in global flood exposure estimates that differ by greater than a factor of 2. The choice of population dataset is found to be equally important and can have enormous impacts on national flood exposure estimates. Up-to-date, high resolution population data is vital for accurately representing exposure to smaller rivers and will be key in improving the global flood risk picture. Our results inform the appropriate application of these datasets and where further development and research is needed.

15 1 Introduction

River floods are amongst the most frequent and damaging natural disasters globally (Wallemacq et al., 2015). Considerable effort has gone into understanding global river flooding over the last decade, and a number of global flood models (GFMs) have been developed concurrently (Yamazaki et al., 2011, Pappenberger et al., 2012, Winsemius et al., 2013, Rudari et al., 2015, Sampson et al., 2015, Dottori et al., 2016). The usefulness of these GFMs was initially limited to coarse scale flood
20 risk assessments (Ward et al., 2015), largely due to global-scale data limitations. However, the incorporation of higher accuracy terrain data available at the national level has shown that their modelling frameworks are also suited to identifying more localized risk when utilising local data (Wing et al., 2017). Previous studies comparing GFMs have shown there is disagreement between the global flood extents (Trigg et al., 2016b, Bernhofen et al., 2018b, Aerts et al., 2020). This disagreement between GFMs stems from different model structures and methods. One key difference between the models,
25 which has not yet been explored, is the size of their river networks. The models have different river size thresholds at which they simulate fluvial events. These thresholds determine the number, and size, of rivers represented in GFMs. The size of a model's river network is contingent on both the quality of the model input datasets and the computational efficiency of the model. It also influences estimates of global flood exposure, as larger river networks result in higher simulated flood volumes and potential exposure. As Remote Sensing (RS) technologies continue to advance, so will the granularity at which
30 rivers can be represented globally. Smaller rivers, previously unrepresented in coarse global datasets, will be able to be studied and modelled at large scales; potentially reframing current global flood exposure estimates. Limited work has been



dedicated to the investigation of the human interaction with rivers of different size (Kummu et al., 2011). Understanding this interaction globally, particularly with respect to river flooding, will inform us about the completeness of current global flood exposure studies and identify where further study and development is needed.

35 A comprehensive understanding of flood risk requires information about the hazard, what or who is exposed, and their vulnerability. Exposure could include damages (both direct and indirect), exposed gross domestic product (GDP), exposed assets, and most commonly: exposed people (Ward et al., 2020). Identifying flood exposed populations usually involves intersecting a flood hazard map with a population map. The methods and inputs used to produce population datasets differ, and so does their intended use (Leyk et al., 2019). Recently released population maps, which utilize commercial RS data and
40 are an order of magnitude more resolved than existing population datasets (Tiecke, 2017) are already being used for disaster preparedness and response (Facebook, 2019). However, our current understanding of global flood exposure is based on existing global population datasets, and these datasets have been used interchangeably in global studies (Tanoue et al., 2016, Jongman et al., 2012, Dottori et al., 2018) with little comment about their relative merit. The credibility of existing global flood exposure estimates in light of new, more detailed, population data and the implications of their interchangeable use in
45 studies of global flood exposure needs to be explored. A recent study by Smith et al. (2019) reported large disagreement between flood exposure estimates calculated in 18 developing countries using three different population datasets. The identification of population data as one of the chief sources of uncertainty in global flood exposure studies warrants further investigation at the global scale.

Improving global flood exposure calculations necessitates a better understanding of both GFMs and the population data their
50 flood hazard maps are intersected with. Specifically, river network size in GFMs and its impact on flood exposure estimates needs to be explored. The number of rivers modelled in GFMs differs by several orders of magnitude depending on the model used. The size of a GFM's river network is limited by the resolution of input data, such as the underlying digital elevation model (DEM) and climatology (Dottori et al., 2016); or by the computational efficiency of the model, as the introduction of smaller rivers exponentially increases the modelled domain. Differences in river network size mean
55 potentially exposed people and infrastructure are missed in global flood risk assessments. This missed exposure has not yet been quantified at the global scale. Advances in RS and computational capacity will enable more rivers to be represented in future iterations of GFMs. But what coverage should modellers be aiming for globally? Understanding where the representation of smaller rivers is most needed (areas of higher exposure) can inform model developers where to prioritize model development and inform model users about the limitations of currently available models. The population data used in
60 global flood exposure calculations varies just as much as the GFMs do. Recent advances in population data, providing more detail and employing new modelling techniques, have been shown to drastically reduce flood exposure estimates in developing countries (Smith et al., 2019). Understanding how both new and existing population datasets differ in their resulting exposure estimates, both regionally and within the hierarchy of the river network, can inform users about the most appropriate population dataset to use.



65 To explicitly explore the impact of river network size on global flood exposure estimates, we use a geomorphological
measure of a river's flood susceptibility, which is independent from current GFMs and the additional uncertainties their
different model structures bring. Fluvial processes contribute to the evolution of a landscape over time. The erosional action
of flowing water has shaped the terrain of drainage basins to reflect the historical flow of water through them. Geomorphological
approaches to mapping river flood susceptibility rely on the concept that the cumulative
70 hydrogeomorphic effect of past flood events, evident in topography data, is indicative of a river's propensity to flood. Such
approaches to flood mapping have been applied over a number of scales: from local (Nardi et al., 2006, Nobre et al., 2016,
Dodov and Foufoula-Georgiou, 2006), to national (Jafarzadegan et al., 2018), to regional (Lugeri et al., 2010) and global
(Nardi et al., 2019). The computational efficiency of hydrogeomorphic flood mapping, coupled with its reliance on only
terrain data as input, make it useful for a 'first look' global scale analysis; intended to inform future development of higher
75 accuracy hydrological flood mapping (Di Baldassarre et al., 2020).

Our geomorphological approach to mapping a river's flood susceptibility, herein referred to as the River Flood Susceptibility
Map (RFSM) is based on new topography data (Yamazaki et al., 2017), which incorporates crowdsourced information to
better represent the locations of rivers and streams (Yamazaki et al., 2019). Validation of our calibrated methodology
(outlined in detail in the Supplementary Material) shows that the RFSM better replicates GFM hazard maps in Africa than an
80 existing global geomorphological approach (Nardi et al., 2019). We also show that the RFSM performs similarly to the best
GFMs (Dottori et al., 2016, Sampson et al., 2015, Yamazaki et al., 2011) when validated against historical flood events
(Bernhofen et al., 2018b). The RFSM allows us to easily discretize the flood map into different river sizes (independently of
GFMs). We investigate the human interface with these different size rivers using three population datasets. Facebook's High
Resolution Settlement Layer (HRSL), ([https://dataforgood.fb.com/docs/high-resolution-population-density-maps-
85 demographic-estimates-documentation/](https://dataforgood.fb.com/docs/high-resolution-population-density-maps-demographic-estimates-documentation/)) (1 arc-second, ~30 m resolution at the equator) (Tiecke, 2017) which is currently
only available in 168 countries globally, and two population datasets used extensively in previous studies of global flood
risk: the Global Human Settlement Population (GHS-POP) ([http://doi.org/10.2905/0C6B9751-A71F-4062-830B-
43C9F432370F](http://doi.org/10.2905/0C6B9751-A71F-4062-830B-43C9F432370F)) (9 arc-second, ~250 m resolution at the equator) (Freire et al., 2015) and WorldPop
(<https://dx.doi.org/10.5258/SOTON/WP00645>) (3 arc-second, ~90 m resolution at the equator) (Stevens et al., 2015, Lloyd
90 et al., 2019). We present a global picture of flood exposure to different size rivers, both in the present day, and how it has
changed over the past 40 years. We then compare the flood exposure calculated using different population layers, exploring
the implications this has on national level flood exposure estimates and examine the impact that river size has on any
disagreement. Finally, we address the size of rivers represented in GFMs specifically and investigate how their chosen river
network size impacts both global and national flood exposure estimates and what implications this has for previously
95 published global flood risk assessments.



2 Methods


2.1 Mapping River Flood Susceptibility

We use a geomorphological approach to mapping river flood susceptibility, which is independent from the global flood models (GFM). Previous GFM comparison studies found that multiple aspects of model structure contributed towards disagreement (Trigg et al., 2016b, Bernhofen et al., 2018b, Aerts et al., 2020). Using a geomorphological approach, we are able to explore just one aspect of disagreement: river network size. This approach allows us to explore all stream scales as drainage paths can be identified from the terrain alone. It is not influenced by the structure of the different GFMs and does not have the same computational restraints as a global hydrodynamic model. This approach is different from the GFMs in that it does not measure the flood extent for a given return period flood, but rather a river and surrounding location's static susceptibility to flooding.

The method used for delineating a river's flood susceptibility is based on the Height Above Nearest Drainage (HAND) methodology developed by Nobre et al. (2011). This hydrogeomorphic approach, requiring only terrain data as input, is computationally efficient, and can be easily modified to produce auxiliary data layers.

Our method, referred to as the River Flood Susceptibility Map (RFSM) (Bernhofen et al., 2021), is illustrated in *Figure 1* and takes three gridded datasets as input: a digital elevation model (DEM), its derived drainage directions, and its upstream drainage area (UDA). We use MERIT hydro data (Yamazaki et al., 2019), a hydrography dataset based on the error improved SRTM DEM: MERIT DEM (Yamazaki et al., 2017). MERIT Hydro is an improvement on previously available global hydrography datasets such as HYDROSHEDs (Lehner et al., 2008) in terms of both spatial coverage and its representation of small streams. Its improved representation of small streams is enabled by its incorporation of global water body data and crowdsourced Open Street Map river data. This makes it particularly suited to this study, where we are interested in examining the flood susceptibility of rivers down to the smallest streams.

The river network is extracted from the upstream drainage area dataset by specifying a minimum threshold river size (in units of UDA). Identifying the headwater of a river is no trivial task, with regional and climatic factors playing a part (Montgomery and Dietrich, 1988, Tarboton et al., 1991). Previous work exploring optimal initiation thresholds for geomorphological floodplain mapping found that DEMs with a resolution of 1 arc second (~30 m) could use initiation thresholds less than 10 km² UDA. In the same study, a 3 arc second (~90 m) resolution DEM was used with a 100 km² UDA threshold (Annis et al., 2019). The MERIT Hydro data we use in this study has a resolution of 3 arc seconds (~90 m). But its incorporation of crowdsourced river data has optimized its representation of small streams and rivers. As such, we use a globally consistent river initiation threshold of 10 km² UDA for the RFSM. This is a large assumption, as in some locations globally there will be no visible channel at this location. However, we argue that removing areas of potential exposure to avoid overprediction in some areas goes against the premise of this study, which is to explore and identify 'missed' areas of exposure. The exposure calculations for small streams should therefore be interpreted with these limitations in mind.




130 **Figure 1. Illustrative example of the method for mapping river flood susceptibility. (a) User defined input parameters include the**
 minimum river size (5 km² used for illustrative purposes rather than 10 km² used in the study) and the height limits H_n for each
 stream order. Dataset inputs include a digital elevation model (DEM), flow direction grid, and an upstream drainage area grid
 (represented on a 10x10 1 km² grid for illustrative purposes). Rivers (as defined by the minimum river size threshold) are
 classified into Strahler stream orders. (b) Each Strahler stream order is processed separately using the Height Above Nearest
 Drainage (HAND) method and then the layers are combined. In areas of overlap the values for the highest order streams are
 135 retained. (c) Two outputs are produced: a map of the drainage area of the nearest flooded river and a map of the Strahler order of
 the nearest flooded river.



Once the river network has been extracted, the rivers in the network are classified based on their Strahler stream orders (Strahler, 1957). The Strahler stream order is a dimensionless indicator of the magnitude of the river based on its hierarchy within the drainage basin. Our method requires the user to assign each order river a maximum height above nearest drainage, H_n (see *Figure 1a*). Each order's H_n value is calibrated against reference flood maps across the world. To account for climatic variability in a river's flood susceptibility (Smith et al., 2015), we split the globe into five simplified Köppen-Geiger climate zones (*Figure 2*): Tropical, Arid, Temperate, Continental and Polar. Polar regions are excluded from our analysis as these regions are dominated by glacial not fluvial processes (Chen et al., 2019). The RFSM has uniquely calibrated H_n values in each of the four climate zones. We use 19 reference flood maps to calibrate the RFSM. These maps span 5 different continents across all four climate zones considered. Reference flood maps are a mixture of national, continental, and global flood hazard maps. To maintain consistency across the calibration data, we use 100-year return period flood hazard maps. We use a combination of national, continental, and global flood hazard maps for calibration in each climate zone. This is to ensure that there is sufficient calibration data for each Strahler order river, as only the national flood hazard data captures flooding for low order rivers. The final H_n values for each climate zone are shown in *Figure 2*. More detailed information on the calibration of the RFSM on each of the calibration basins can be found in the Supplementary Material.

Once H_n values for each order have been assigned, each stream order is processed separately (*Figure 1b*), and then merged together. In areas of overlap, the highest order stream retains the values. Two datasets are produced as output: a map of the flooded river's upstream drainage area, and a map of the flooded river's Strahler stream order. Illustrations of these two outputs are shown in *Figure 1c*.



Order	Maximum H_n (m)			
	Tr	Ar	Te	Co
1	1	0	0	0
2	2	0	1	0
3	4	0	2	2
4	6	1	4	4
5	8	3	5	5
6	8	3	7	6
7	9	5	9	8
8	10	5	10	9
9	10	12	10	10
10	10	13	13	11
11	13	14	-	12

155

Figure 2. The globe split into simplified Köppen Geiger climate zones and the final maximum Heights Above Nearest Drainage (H_n) for each Strahler stream order in the four climate zones considered (Polar regions are excluded from the analysis)



- 160 The RFSM was validated against both existing GFMs and observed flood events. Validation against GFMs was carried out for the whole of the African continent using the aggregated output of 6 GFMs from a previous model intercomparison study (Trigg et al., 2016a). To assess the credibility of the RFSM, it was also validated alongside an existing global geomorphological floodplain map (Nardi et al., 2019). The results of the GFM validation show that the RFSM produces credible flood extents when compared with existing GFM outputs in Africa. The RFSM correctly captures over 90% of high
- 165 agreement flood zones (where at least 5 out of 6 GFMs agree) in 7 of the 8 major drainage basins in Africa. In the East African basin, the RFSM captures 87% of this high agreement flood zone. Comparing commonly used measure of fit scores for the RFSM and the existing global geomorphological floodplain map, the RFSM scores higher in all the major drainage basins in Africa except for North Africa (where both maps score poorly due to the Sahara desert). The RFSM was also validated against observed flood events in Nigeria and Mozambique. The 2012 flooding in Nigeria and the 2007 floods in
- 170 Mozambique affected four million people and over one hundred thousand people respectively (Bernhofen et al., 2018b). Validation data for both these flood events used in a previous GFM validation comparison study (Bernhofen et al., 2018a) was also used to validate the RFSM. Validation of the RFSM against these historical flood events show that it performs similarly to the best performing GFMs. Further detail about the validation of the RFSM can be found in the Supplementary Material.
- 175 It is important to note, our methodology does not account for flood protection measures and cannot communicate the probability of flooding in any location. It consistently represents a river's flood susceptibility based on the surrounding terrain alone. The method's intended use is as a global 'first look' analysis to inform future model development and use.



2.3 Measuring Exposure

180 We investigate the human exposure to river flood susceptibility. Human exposure is herein defined as the intersection of our
flood susceptibility map and a spatially distributed population layer. Three population datasets are used to measure exposure:
Facebook's High Resolution Settlement Layer (HRSL) ([https://dataforgood.fb.com/docs/high-resolution-population-density-
185 maps-demographic-estimates-documentation/](https://dataforgood.fb.com/docs/high-resolution-population-density-maps-demographic-estimates-documentation/)) (Facebook and CIESIN, 2016), The European Commission Joint Research
Centre's Global Human Settlement Population (GHS-POP) ([http://doi.org/10.2905/0C6B9751-A71F-4062-830B-
43C9F432370F](http://doi.org/10.2905/0C6B9751-A71F-4062-830B-43C9F432370F)) (Schiavina, 2019), and WorldPop (<https://dx.doi.org/10.5258/SOTON/WP00645>) (Stevens et al., 2015).
These population datasets all use the same initial input census data, from GPWv4 (Center for International Earth Science
Information Network - CIESIN - Columbia University, 2016), but their methods for allocating the population across gridded
cells differ. Facebook's HRSL is the only dataset of the three lacking full global coverage (at the time of writing 168
countries have been mapped). It is also the most recent, with work ongoing to map the remaining countries. HRSL uses
190 ultra-high resolution commercial satellite imagery (~50 cm resolution) and convolutional neural networks to detect
individual buildings at the country level (Tiecke, 2017). Subnational census data is then proportionally allocated to the
identified buildings at 1 arc second resolution (~30 m at the equator).

Similarly to the HRSL in methodology, JRC's GHS-POP dataset identifies built up areas from Landsat imagery and
proportionally allocates census data to the built up areas (Freire et al., 2015). In regions where no settlements can be
195 identified, but where census data indicates there is a population, the population is evenly distributed across the census area
using areal weighting (Freire et al., 2016). This can occur in some rural areas, where small settlements are not captured by
the Landsat imagery. Despite being coarser in spatial resolution at 9 arc seconds (~250 m at the equator), GHS-POP
provides consistent multi-temporal population estimates (1975-1990-2000-2015) allowing for accurate analyses over time
(Freire et al., 2020).

200 Unlike the other two population datasets, which evenly spread census data over identified settlements, WorldPop uses a
complex model to disaggregate population over an area (Leyk et al., 2019). It uses a random forest model and a number of
ancillary datasets to dynamically weight the distribution of census data over a 3 arc second (~90 m at the equator) gridded
area (Stevens et al., 2015).

Exposure calculations necessitate uniformity between the intersecting datasets in terms of spatial resolution. As such, the
205 GHS-POP layer was resampled from 9 arc second resolution and the population evenly distributed to a 3 arc second
resolution grid to allow for analysis with a flood map of the same resolution. Conversely, for the HRSL exposure
calculations the RFSM was resampled from 3 arc second to 1 arc second resolution. When comparing the exposure results
between population datasets the epoch used for comparison was 2015. National population totals for the HRSL and
WorldPop datasets were scaled relative to GHS-POP 2015 national population totals.



3 Results and Discussion

3.1 Global Exposure to Different Sized Rivers

Rivers were classified into 6 different sizes, expressed in upstream drainage area (UDA) (km²), with the ranges increasing in powers of 10. River classifications based on UDA, depicted in *Figure 3b* for Nigeria, were as follows: stream (10-100 km²),
215 small river (100-1,000 km²), medium river (1,000-10,000 km²), medium-large river (10,000-100,000 km²), large river (100,000-1,000,000 km²), and huge river (>1,000,000 km²).

Flood exposure is first calculated using the GHS-POP layer. Globally, we find 1.94 billion people are susceptible to flooding from rivers with a UDA greater than 10 km². Breaking this down by continent, Asia's flood exposure is 1.49 billion, Africa's is 203 million, Europe's is 104 million, North America's is 81 million, South America's is 59 million, and Oceania's is 3.5
220 million. Splitting global flood exposure by river size, of the total exposed: 18.2% are from streams, 26.4% from small rivers, 23.7% from medium rivers, 17.2% from medium-large rivers, 8.4% from large rivers, and 6.1% from huge rivers. Asia makes up over 75% of the total global flood exposure, the majority of this amount coming from India and China, which are by far the two most exposed countries (see *Figure 3a*). Roughly half of India's flood exposure is from streams and small rivers. Comparably, in China, this figure is closer to a third. This is likely due to the degree of urbanisation in both countries;
225 the percentage of China's urban population is double that of India's (WorldBank, 2018). Urban areas are disproportionately located on large rivers due to the historical tendency for settlements to form in areas fertile for farming and convenient for transport (McCool et al., 2009). As such, a greater proportion of flood exposure in China comes from larger rivers, whereas in India, a greater proportion comes from rural exposure to smaller rivers. Rivers classified as 'huge' are only found in some countries, but often they pose a large proportion of the national flood risk. For example, the Brahmaputra in Bangladesh and
230 the Nile in Egypt and Sudan are responsible for just under half of the national flood exposure in their respective countries.

To identify countries with the most acute flood risk, exposure was normalized against total national population (*Figure 3c*). Suriname has the highest normalized exposure, with 894 people exposed per 1000. The country's low elevation relief, and its capital city situated on the banks of the Suriname river near its outlet into the Atlantic Ocean, makes Suriname particularly vulnerable to flooding (WorldBank, 2019). Four of the top 10 most 'normally' exposed countries are in south or south east
235 Asia. These include: Bangladesh, Cambodia, Thailand, and Vietnam. Flooding in these countries is severe and annual, normally occurring each year during the monsoon season. In Europe, the Netherlands has a high normalized exposure, 738 exposed per 1000. The Netherlands has a long history of flooding due to its low elevation, flat terrain, and high population density. It also has the most advanced flood defence systems in the world, designed to contain river water levels with a probability of occurrence once every 1250 years (Stokkom et al., 2005). Geomorphological approaches to flood mapping,
240 such as the RFSM, cannot model probabilities of occurrence; and are therefore unable to represent flood prevention measures and distinguish defended and undefended floodplain zones. Much of the exposed population in the Netherlands, as well as other countries with flood protection, reside in the defended area of a floodplain. This does not eliminate their risk of



flooding; just reduces the probability of it. The severity of a flood event when defences fail can be catastrophic, resulting in high velocity flows and rapid inundation with little to no warning.

245 The results presented in this section, and *Figure 3*, were calculated using the GHS-POP layer. The results of this analysis calculated using the other two population datasets (WorldPop and HRSL) are included in the Supplementary Material. Global exposure calculated using the WorldPop layer is just over 2 Billion, roughly 81 Million larger than the global figure calculated using GHS-POP. The proportion of global exposure per river size for WorldPop is as follows: 18.5% from streams, 26% from small rivers, 23.9% from medium rivers, 17% from medium-large rivers, 8.6% from large rivers, and 6%
250 from huge rivers. Each of these is within a percentage point of the GHS-POP calculated figures. The top 50 countries, both in terms of total population and normalized population exposed, remain largely the same when using WorldPop data, although their order changes slightly. It's not possible to compare these global results with HRSL calculated exposure, as it does not yet have global coverage. We explore the implications of using different population datasets for flood exposure calculations in greater detail later in this paper.

255




Figure 3. GHS-POP calculated flood exposure. (a) Top 50 most exposed countries in terms of total flood exposure. (b) The river size classifications visualized in Nigeria. (c) Top 50 most exposed countries in terms of normalized flood exposure (normalized to country's total population)




260 3.2 Exposure Change from 1975 – 2015

An advantage of both the GHS-POP and WorldPop datasets is their population estimates across different time scales, allowing for exposure analysis over time. WorldPop has annual population maps from 200-2020 and GHS-POP has population estimates across four epochs: 1975, 1990, 2000, 2015. Here, using GHS-POP's multitemporal population layers, we calculate exposure change over a period of 40 years. Normalized flood exposure estimates were calculated for the years 265 1975, 1990, 2000, and 2015. Globally, total flood exposure grew between 1975 and 2015 from 257 people per 1000 to 265 people per 1000. Interestingly, in both Tropical and Arid climates total flood exposure over this 40 year period grew by 11 people per 1000; but in Temperate and Continental climates total flood exposure decreased by 4 and 10 people per 1000, respectively. Developing countries are largely located in tropical and arid climates, conversely, developed economies are prevalent in temperate and continental climates. These findings correspond with previous work done by Jongman et al. 270 (2012), which found developing countries had the largest increases in exposure relative to population growth in the period 1970-2010. Exposure over the period 1975-2015 increased for streams, medium-large, large and huge rivers. There were slight reductions in exposure for small and medium sized rivers.

Exposure changes at the national level are depicted in *Figure 4*. The highest increase in overall flood exposure was seen in Nepal and French Guinea. In both countries, the proportion of exposed population grew by 200 people per 1000 in the period 275 1975-2015. In French Guinea, this sudden increase is largely due to the population growth of Saint-Laurent-du-Maroni, a town situated on the banks of the Maroni river. From 1975-2015 the town's population grew 1800% compared with the national population growth of 360%. In Nepal, one of the top 10 fastest urbanizing countries in the world (Bakrania, 2015), the flood exposure growth is a result of this fast urbanisation in cities such as Kathmandu, which is dissected by eight different rivers. An exposure decrease of 172 people per 1000 was seen in South Sudan.. This is due to the growth of urban 280 areas outside the Sudd swamp in cities such as Juba, Yei, Yambio, Nzara, and Wao. South Sudan has been hit by devastating floods in the past year, which displaced over 800,000 people (OCHA, 2020). Had relative population exposure in South Sudan grown, rather than shrunk, the recent flooding could have been even worse.



All Rivers (UDA >10 km²)





Change in normalized exposure 1975-2015 (p. per 1000)





Streams (UDA 10-100 km²)

Small Rivers (UDA 100-1,000 km²)





Medium Rivers (UDA 1,000-10,000 km²)

Medium-Large Rivers (UDA 10,000-100,000 km²)



Large Rivers (UDA 100,000-1,000,000 km²)

Huge Rivers (UDA >1,000,000 km²)



285 **Figure 4. Country level river flood exposure change from 1975-2015 calculated using GHS-POP. River size expressed in Upstream Drainage Area (UDA)**



3.3 Variation in Exposure

Exposure differences arising from the use of different population layers were calculated for the 168 countries where all three population datasets are available (*Figure 5*) (see Supplementary Material for a list of the missing countries). In the countries examined, the exposure calculated with WorldPop data was the highest (270 exposed per 1000), followed by GHS-POP (256
290 exposed per 1000), HRSL exposure was the lowest (235 exposed per 1000). These findings correlate with a previous study by Smith et al. (2019) which found WorldPop data overestimated flood exposure compared to HRSL data in each of the 18 developing countries examined.


Differences in calculated exposure across the river sizes are shown in *Figure 5b*. Exposure differences were most pronounced for smaller rivers (streams, small, and medium rivers), while there was almost no exposure difference for the
295 largest river class (huge). The overall trend across all river sizes consistently shows that WorldPop estimated the highest exposure, followed by GHS-POP, and then HRSL with the lowest.

The methods for producing the three population layers can go some way towards explaining the differences in calculated exposure, these methods are visualized in *Figure 6*, where we qualitatively compare the settlement distribution of the three population datasets along the Likuala-aux-Herbes river in the Republic of Congo. WorldPop's population distribution
300 algorithm spreads some residual population across the grid in areas where no settlements have been identified. This is done under the assumption that not all 'built up' areas will be picked up in the satellite imagery (TReNDS, 2020). When intersected with a flood extent, this residual population spread can lead to overprediction of flood exposure in rural areas with respect to the other two population datasets. Conversely, the approach implemented by both GHS-POP and HRSL
(which spread census data only over identified 'built up' areas) is more sensitive to omission and commission errors arising
305 from the classification of settlements (Palacios-Lopez et al., 2019). For example, undetected settlements outside the flood extent would result in artificially higher flood exposure estimates as the underlying census data is only spread across the identified settlements (an incorrectly greater proportion of which are now identified as within the flood extent). Similarly, commission errors (false positives) are common in sandy or rocky landscapes and often occur in coastal areas or along riverbanks. Commission and omission errors can lead to either artificial increases or decreases in flood exposure estimates,
310 depending on the location of these errors with respect to the flood extent.

The resolution of the population layers should also be considered. GHS-POP's fairly coarse (9 arc second) resolution means that in some areas where the potential for flooding (or not) falls within the resolution of a 9 arc second grid cell, the settlement's avoidance (or not) of the flood risk cannot be accurately represented. This effect can be reduced by upsampling and proportionally reallocating the population to a grid that matches the resolution of the flooded data, as we have done in
315 this study. Similarly, the spatial resolution of the underlying satellite imagery should be considered. Both GHS-POP and WorldPop identify settlements using Landsat imagery at 30 m resolution, while HRSL identifies settlements using DigitalGlobe imagery at 0.5 m resolution. Previous work by (Tiecke, 2017) showed that HRSL was able to identify



buildings missed by GHS-POP, highlighting the importance of high-resolution imagery for comprehensive building classification.



320

Figure 5. HRSL, WorldPop, and GHS-POP flood exposure comparison in 169 countries. (a) Comparison of the total normalized flood exposure between the three population datasets in all available countries. (b) How the calculated exposure figures differ per river size classifications. (c) Country level statistics for average normalized exposure and the sensitivity of the exposure calculation to the choice of population dataset (measured in range of calculated exposures.). The higher up the X axis and Y axis, the greater the average exposure and sensitivity to the choice of population dataset, respectively.

325




The use of different population datasets had a negligible effect on exposure estimates for the huge river class. Large settlements tend to form around rivers of this size, and on coastlines where rivers of this size drain. Large urban areas are easily identifiable from remote sensing data, which means the population distribution (and resulting exposure estimates) for these urban centres show less variation between the datasets. Conversely, non-urban flood exposure estimates to smaller
330 rivers show greater sensitivity to the choice of population layer. This is because the approach to non-urban population mapping between the three datasets differ. WorldPop, as mentioned previously, distributes administrative level census data across all 3 arcsecond pixels in order to mitigate the impacts of potential omission and commission errors in the settlement data. This approach leads to some overestimation in rural populations (Smith et al., 2019, Wardrop et al., 2018). GHS-POP, which distributes census data over Landsat identified settlements (and in non built-up areas distributes population at the
335 census unit by areal weighting), tends to underestimate rural populations. (Liu et al., 2020, Leyk et al., 2019). HRSL's use of ultra-high resolution satellite imagery has been shown in previous studies to accurately identify rural settlements (Tiecke, 2017, Smith et al., 2019). However, there is still significant uncertainty in the underlying census data and the method of proportional allocation used to distribute the census data is relatively crude.

Calculating the general trends of exposure between the population layers is useful for making broad conclusions about the
340 suitability of a population layer. Understanding the variations of the data at the country level leads to more actionable information about the appropriate use of different population layers. The disagreement between the population layer exposure estimates for each country varies significantly (*Figure 5c*). In the three countries with the highest exposure disagreement (Belize, The Republic of Congo, and Guinea-Bissau) WorldPop estimates of exposure are far greater than either HRSL or GHS-POP estimates. In Belize, a country with large areas of inundated wetlands, WorldPop estimates
345 135,000 people exposed, while GHS-POP and HRSL estimate 70,000 and 80,000 exposed, respectively. In the Republic of Congo, a country with large areas of floodplain, WorldPop estimates 1.3 Million people exposed and GHS-POP and HRSL estimate 810,000 and 780,000 exposed respectively. WorldPop's method of distributing the population over a large area results in significant overestimation compared with HRSL or GHS-POP in these rural inundated areas. This is evident in
350 *Figure 6*, where the population distributions are qualitatively compared with manually identified settlements in a small region in the Republic of Congo. WorldPop's distribution of population results in a larger area of exposure compared with the other two datasets and the manually identified settlements. This then leads to overestimation in regions with large areas of flood extent. This can be seen in greater detail in *Figure 7* for Guinea-Bissau. In Guinea-Bissau, GHS-POP and HRSL (which estimate exposures of 180,000 and 160,000 respectively) identify settlements largely situated outside the floodplains ('dry' cells in red). Comparatively, WorldPop's residual population spread leads to far more 'wet' population cells and an
355 estimate of exposure (480,000) more than double that of the other two population layers. The exposure disagreement in these three countries is compounded by the relatively large areas of inundation in each country. The percentage inundated area is 25%, 30%, and 26% for Guinea-Bissau, Belize, and The Republic of Congo, respectively. In comparison, the percentage of populated area defined by the population layers is less than 5% for GHS-POP and HRSL, but more than 95% for WorldPop in each of the three countries. As exposure in this study is defined as the intersection of the flooded area and the populated



360 area, it is understandable that WorldPop's exposure estimates are more sensitive to the area of inundation. This is evident
when examining a country with high exposure disagreement but with a comparatively smaller area of inundation. In Bosnia
and Herzegovina (*Figure 7*), the percentage of flooded area is just 9% and the GHS-POP layer estimates far greater exposure
(1 Million) than either WorldPop (680,000) or HRSL (610,000). Here, where much of the exposure occurs near the banks of
the rivers, the coarse spatial resolution of GHS-POP is less able to precisely locate settlements situated just outside the
365 floodplain. As a result, more populated cells are flagged 'at risk' compared to the higher resolution HRSL layer.




370 **Figure 6.** Qualitative comparison of settlement distributions on the Likouala-aux-Herbes river in the Republic of Congo. (a) RFSM flood
extent along with manually identified settlements from high resolution Google Earth satellite imagery. (b) HRSL settlement distribution.
(c) WorldPop settlement distribution (resampled to 1 arc second for comparison). (d) GHS-POP settlement distribution (resampled to 1 arc
second for comparisons). Map Data: © Google, Maxar Technologies 2021

These results have shown that the use of different population layers can lead to vastly different flood exposure estimates
because of inherent differences in their spatial resolutions, methods used, and assumptions made to produce them. Our
comparative analysis has identified in which countries exposure calculations are sensitive to the choice of population layer
and shed light on some of the reasons for exposure disagreement. However, there is a limit to the conclusions that can be
375 drawn from comparative analyses alone, and there is an urgent gap for more studies which validate the accuracy of these
population layers using ground-truthed data.

It would be imprudent to definitively recommend one population dataset for use in flood exposure studies without extensive
comparative global validation. However, previous studies have shown that HRSL performs better than existing population
datasets at mapping reference building footprints, especially in rural areas (Tiecke, 2017, Smith et al., 2019). Our results also
380 point to some of the benefits of using HRSL. Its settlement identification method for population distribution avoids exposure
overprediction common in other population data and its high resolution can better capture the accurate location of
settlements. Despite this, HRSL shouldn't be considered a catchall dataset for flood exposure. Its high resolution may limit
its use in certain situations due to computational restraints. Similarly, in studies of flood risk over time population data with
multiple temporal epochs, such as GHS-POP or WorldPop, are better suited. The results we present in this section, and
385 *Figure 5*, are intended to inform users of these population datasets about their appropriate use. In countries with high



exposure disagreement, the choice of population dataset for flood exposure should be carefully considered, and further accuracy assessments of the population layers are recommended.



390 **Figure 7.** Comparison of population datasets and their intersection with the flood extent in Bosnia and Herzegovina and Guinea-Bissau. The top windows show the River Flood Susceptibility Map (RFSM) for each country, split into the different river size



categories. The remaining windows show the three different population maps and their intersection with the flood maps. Blue cells indicate the population cell is wet (W) or inundated. Yellow-Red cells indicate the population cell is dry (D) or not inundated. All population datasets in this figure were resampled to 30 m resolution for comparison.

3.4 Relevance to Global Flood Models

395 The minimum size of river represented in Global Flood risk Models (GFMs) varies (see *Table 1*), with minimum river size thresholds ranging between 50-5000 km² UDA, three orders of magnitude. River network size can be limited by the granularity of input data such as rainfall (Dottori et al., 2016), or by the computational demand of modelling floods at the global scale.

Table 1. Global Flood Model River Representation

Minimum River Size (upstream drainage area)	Global Flood Risk Model	River Sizes Modelled (P = Partial)
50 km ²	Fathom (Sampson et al., 2015)	Stream (P), Small, Medium, Medium-Large, Large, Huge
500 km ²	ECMWF (Pappenberger et al., 2012) and U-Tokyo (Yamazaki et al., 2011)	Small (P), Medium, Medium-Large, Large, Huge
1000 km ²	CIMA-UNEP (Rudari et al., 2015)	Medium, Medium-Large, Large, Huge
5000 km ²	JRC (Dottori et al., 2016)	Medium (P), Medium-Large, Large, Huge

400

Differences in river network size between GFMs undoubtedly lead to differences in global flood exposure estimates. These differences can be even more pronounced at the national level, where GFMs have been used to inform disaster risk management (Ward et al., 2015). Flood exposure was calculated for the different GFM river thresholds using the GHS-POP layer. Globally, we found that exposure estimates between the river threshold which results in the largest river network (>50 km² UDA), and the river threshold which results in the smallest river network (>5000 km² UDA), differ by over a factor of 2. If the size of the river network was further increased by reducing the river threshold to 10 km² UDA (below current GFM representation), the exposed population captured increases by 13%.

405

At the national level, in countries such as Suriname, The Republic of Congo, and Egypt, the greatest proportion of flood risk is posed by rivers with a UDA of 5000 km² or greater. In these countries, GFMs could be used interchangeably.

410

Understanding what size rivers pose a significant flood risk is key to accurately representing national flood risk. In Benin, for example, the estimated flood exposure when a 5000 km² UDA threshold is applied is 0.49 million people. When the threshold is reduced to 1000 km² UDA, the estimated exposure increases to 1.8 million people. Some countries do not have large rivers flowing through them, and the flood risk will result entirely from smaller rivers. Often these are island nations, such as in Jamaica or Trinidad and Tobago, where all flood risk is from rivers smaller than UDA 1000 km². However, in

415

Andorra for example, a landlocked country, to capture any flood exposure, a 50 km² UDA threshold is needed.

To aid national level flood risk practitioners in their choice of GFM, we calculated the minimum river threshold required to capture a given percentage of the largest river network's (>50 km² UDA) national exposure. Exposure percentages ranging



from 10-90% were calculated for each of the three population datasets used in this study and mapped for each nation, globally. All 27 maps are included in the Supplementary Material. *Figure 8*, which shows the minimum river threshold required to capture at least 50% of possible GHS-POP exposure illustrates these results. The map shows that while in some countries GFMs could be used interchangeably, in others, the size of the river network could significantly impact national flood exposure estimates.




Figure 8. What size river threshold is required to capture at least 50% of the country's total GFM flood (>50 km²) exposure. Calculated using GHS-POP population layer. Results could inform the choice of Global Flood Model used at the national level.

It is difficult to exhaustively compare global flood exposure estimates from previous GFM studies as often exposure is expressed differently (e.g. expected annual exposure (EAE) vs. exposure to a return period flood) and sometimes global exposure is not reported at all. In the comparable studies, there is significant variation in global flood exposure estimates. In Ward et al. (2013) global EAE was calculated at 169 Million. This figure is almost triple the 58 Million calculated by Dottori et al. (2018) and the 54 Million calculated by Alfieri et al. (2017). In studies reporting exposure to a 100 year flood, Hirabayashi et al. (2013) estimate 847 Million people exposed and Jongman et al. (2012) estimate 805 Million exposed. The need for independent model comparison studies was met by Trigg et al. (2016b) and Aerts et al. (2020) who compared GFM output in Africa and China respectively. These studies compared the output of multiple GFMs, finding large disagreement between the modelled flood extents. Both studies also found large variations in calculated exposure. However, differences in exposure calculated by the GFMs were found to be influenced just as much by different model forcing and resolution as by differences in river network size. Uncertainty in GFMs needs to be explored across the model cascade to



identify where the models need to improve. Studies such as Zhou et al. (2020), which explores uncertainty in model forcing; and this study, which explores uncertainties in river network size, are important steps in directing future model development. Granularity of input data is the main obstacle to increasing river network size in GFMs. The terrain data in all these models, which strongly influences their performance, is derived from the Shuttle Radar and Topography Mission (SRTM), a mission over two decades old (Farr et al., 2007). New, 1 arc second resolution (~30 m at the equator) global DEMs have recently been released by both the National and Aeronautics and Space Administration (NASA) and the European Space Agency (ESA). The ESA DEM is particularly important as its elevation is based on newer satellite data from TanDEM-X. A new method for deriving an elevation map from satellite images has also been developed by Google, capable of generating DEMs at 1m resolution (Nevo, 2019). Whether its terrain or climatology data, new and improved methods are constantly being developed and better datasets are being released. There is scope in the near future for increasing river network size in GFMs. This comes at a computational cost, however; whether it's the use of a higher resolution DEM or the exponential increase in number of rivers to model when the threshold river size is reduced. Understanding where the representation of smaller rivers is needed most, namely in areas of high exposure, would streamline the future development of GFMs, targeting improvements in areas where flood risk is highest.

4 Conclusions

This study has presented the first global picture of flood exposure categorised by different sized rivers. We introduced a simple geomorphological approach to delineating a river's flood susceptibility, which is suitable for global scale 'first look' studies such as this and importantly, allows an assessment of river network size independent of global flood model structural and computational limitations. We find that over 75% of the global flood exposure is in Asia, with China and India making up a significant proportion of this total. Streams (UDA 10-100 km²) and small rivers (UDA 100-1000 km²) are responsible for over half of India's flood risk. At the global scale, these rivers contribute to 45% of total flood exposure, emphasizing the importance of the incorporation of these smaller rivers into global flood risk studies. We find that large increases and decreases in flood exposure over the last forty years are a result of urbanisation, either inside the flood risk zone or outside of it. The effect that the choice of population dataset had on exposure calculations differed between countries. Globally, this effect was most pronounced on smaller rivers, suggesting future studies that incorporate these smaller rivers should be careful in their choice of population data. Global flood models, the current tools for examining global flood risk, differ significantly in the size of their river networks. We found that the global flood exposure estimates differed by greater than a factor of 2 when calculated using the GFM river threshold which results in the largest river network (UDA >50 km²) compared to the river threshold which results in the smallest river network (UDA >5000 km²). These differences were often more pronounced at the national level.

The results of this study are intended to inform both the developers and users of global river flood models. Consideration of river network size, and how this relates to exposure, is imperative to having a comprehensive picture of flood risk. Increasing



the size of the river network comes with both data and computational restraints. Doubling the resolution of the models (from
470 1km to 90 m to 30 m) requires an order of magnitude increase in computing power. Finer resolution grids are imperative for
representing small streams accurately. This has big implications for models currently operating at coarse resolution.
Modelling smaller rivers requires not only detailed high-resolution data, but also efficient modelling structures capable of
running at higher resolutions. Understanding where the representation of small rivers is needed most (areas of high
exposure) can focus future model development. Similarly, accurate flood exposure estimates necessitate accurate population
475 data. We have shown that the choice of population data used in exposure calculations can have an enormous impact on flood
exposure estimates and we have identified in which countries this disagreement is most extreme and have identified some of
the reasons for this. Flood risk practitioners should use these results as guidance about which population layer is best suited
for their locality and use. There is need for further research in this area, incorporating more population data, as these layers
play such an integral role in flood exposure calculations. In addition to more comparative analyses, there is also an urgent
480 need for this population data to be validated at the global scale with actual data collected on the ground. Only then can
definitive conclusions be drawn about the appropriate use of different population datasets. The selection of GFM available
to the end user is large and increasing. However, differences in the size of river networks between the models can have a
significant impact on flood exposure estimates. While available GFMs could be used interchangeably in some countries, in
others, discrepancies in river network size would lead to vastly different national flood exposure estimates. The results of
485 this study should help to inform GFM users about the appropriate choice of GFM for their country of interest.

Author Contributions

MVB and MAT conceived of the study. MB designed and carried out the analysis. MAT PAS CCS AMS supervised the project. MVB drafted the manuscript. All authors contributed towards the discussion and editing of the manuscript.

Competing Interests

490 The authors declare that they have no conflict of interest.

Acknowledgements

MVB was funded by the UK National Environmental Research Council Grant NE/R008949/1 and iCASE funding from Fathom Global. This work was also supported by the Water Security and Sustainable Development Hub funded by the UK Research and Innovation's Global Challenges Research Fund (GCRF) [grant number: ES/S008179/1]. This work was
495 undertaken on ARC3, part of the High-Performance Computing facilities at the University of Leeds, UK. The authors would like to thank the members of the Global Flood Partnership, who have helped to shape this research through discussion and feedback at numerous GFP workshops.



Data Availability

All the data used in this study is freely available to download. The River Flood Susceptibility Maps are available from the
500 University of Leeds at <https://doi.org/10.5518/947>. Facebook's High Resolution Settlement Layer can be downloaded by
following the instructions in this link [https://dataforgood.fb.com/docs/high-resolution-population-density-maps-
demographic-estimates-documentation/](https://dataforgood.fb.com/docs/high-resolution-population-density-maps-demographic-estimates-documentation/). The GHS-POP data can be downloaded here [http://doi.org/10.2905/0C6B9751-
A71F-4062-830B-43C9F432370F](http://doi.org/10.2905/0C6B9751-A71F-4062-830B-43C9F432370F). The WorldPop data can be downloaded here
<https://dx.doi.org/10.5258/SOTON/WP00645>.

505

References

- AERTS, J. P. M., UHLEMANN-ELMER, S., EILANDER, D. & WARD, P. J. 2020. Comparison of estimates of global flood models for
flood hazard and exposed gross domestic product: a China case study. *Nat. Hazards Earth Syst. Sci.*, 20, 3245-3260.
- 510 ALFIERI, L., BISSELINK, B., DOTTORI, F., NAUMANN, G., DE ROO, A., SALAMON, P., WYSER, K. & FEYEN, L. 2017. Global
projections of river flood risk in a warmer world. *Earths Future*, 5, 171-182.
- ANNIS, A., NARDI, F., MORRISON, R. R. & CASTELLI, F. 2019. Investigating hydrogeomorphic floodplain mapping performance
with varying DTM resolution and stream order. *Hydrological Sciences Journal*, 64, 525-538.
- 515 BAKRANIA, S. 2015. Urbanisation and urban growth in Nepal. *GSDRC Helpdesk Research Report 1294*. Birmingham, UK: GSDRC
University of Birmingham.
- BERNHOFEN, M. V., TRIGG, M. A., SLEIGH, P. A., SAMPSON, C. & SMITH, A. 2021. River Flood Susceptibility Map.
- BERNHOFEN, M. V., WHYMAN, C., TRIGG, M. A., SLEIGH, P. A., SMITH, A., SAMPSON, C., YAMAZAKI, D., WARD, P.,
RUDARI, R., PAPPENBERGER, F., DOTTORI, F., SALAMON, P. & WINSEMIUS, H. C. 2018a. Analysis inputs for
validation of six global flood models against observed flood events in Nigeria and Mozambique. Research Data Leeds
520 Repository.
- BERNHOFEN, M. V., WHYMAN, C., TRIGG, M. A., SLEIGH, P. A., SMITH, A. M., SAMPSON, C. C., YAMAZAKI, D., WARD, P.
J., RUDARI, R., PAPPENBERGER, F., DOTTORI, F., SALAMON, P. & WINSEMIUS, H. C. 2018b. A first collective
validation of global fluvial flood models for major floods in Nigeria and Mozambique. *Environmental Research Letters*, 13.
- 525 CENTER FOR INTERNATIONAL EARTH SCIENCE INFORMATION NETWORK - CIESIN - COLUMBIA UNIVERSITY 2016.
Gridded Population of the World, Version 4 (GPWv4): Population Count. Palisades, NY: NASA Socioeconomic Data and
Applications Center (SEDAC).
- CHEN, S.-A., MICHAELIDES, K., GRIEVE, S. W. D. & SINGER, M. B. 2019. Aridity is expressed in river topography globally. *Nature*.
- DI BALDASSARRE, G., NARDI, F., ANNIS, A., ODONGO, V., RUSCA, M. & GRIMALDI, S. 2020. Brief communication: Comparing
hydrological and hydrogeomorphic paradigms for global flood hazard mapping. *Nat. Hazards Earth Syst. Sci.*, 20, 1415-1419.
- 530 DODOV, B. A. & FOUFOULA-GEORGIOU, E. 2006. Floodplain morphometry extraction from a high-resolution digital elevation
model: a simple algorithm for regional analysis studies. *IEEE Geoscience and Remote Sensing Letters*, 3, 410-413.
- DOTTORI, F., SALAMON, P., BIANCHI, A., ALFIERI, L., HIRPA, F. A. & FEYEN, L. 2016. Development and evaluation of a
framework for global flood hazard mapping. *Advances in Water Resources*, 94, 87-102.
- 535 DOTTORI, F., SZEWCZYK, W., CISCAR, J. C., ZHAO, F., ALFIERI, L., HIRABAYASHI, Y., BIANCHI, A., MONGELLI, I.,
FRIELER, K., BETTS, R. A. & FEYEN, L. 2018. Increased human and economic losses from river flooding with anthropogenic
warming. *Nature Climate Change*, 8, 781-+.
- FACEBOOK. 2019. *Data for Good: Population Density Maps* [Online]. Available: [https://dataforgood.fb.com/tools/population-density-
maps/](https://dataforgood.fb.com/tools/population-density-maps/) [Accessed December 2020].
- 540 FACEBOOK & CIESIN 2016. High Resolution Settlement Layer (HRSL). In: UNIVERSITY, F. C. L. A. C. F. I. E. S. I. N.-C.-C. (ed.).
FARR, T. G., ROSEN, P. A., CARO, E., CRIPPEN, R., DUREN, R., HENSLEY, S., KOBRICK, M., PALLER, M., RODRIGUEZ, E.,
ROTH, L., SEAL, D., SHAFFER, S., SHIMADA, J., UMLAND, J., WERNER, M., OSKIN, M., BURBANK, D. & ALSDORF,
D. 2007. The shuttle radar topography mission. *Reviews of Geophysics*, 45, 33.



- FREIRE, S., KEMPER, T., PESARESI, M., FLORCZYK, A. & SYRRIS, V. Combining GHSL and GPW to improve global population mapping. 2015 IEEE International Geoscience and Remote Sensing Symposium (IGARSS), 26-31 July 2015. 2541-2543.
- 545 FREIRE, S., SCHIAVINA, M., FLORCZYK, A. J., MACMANUS, K., PESARESI, M., CORBANE, C., BORKOVSKA, O., MILLS, J., PISTOLESI, L., SQUIRES, J. & SLIUZAS, R. 2020. Enhanced data and methods for improving open and free global population grids: putting 'leaving no one behind' into practice. *International Journal of Digital Earth*, 13, 61-77.
- FRIERE, S., MACMANUS, K., PESARESI, M., DOXSEY-WHITFIELD, E. & MILLS, J. 2016. *Development of new open and free multi-temporal global population grids at 250 m resolution*.
- 550 HIRABAYASHI, Y., MAHENDRAN, R., KOIRALA, S., KONOSHIMA, L., YAMAZAKI, D., WATANABE, S., KIM, H. & KANAE, S. 2013. Global flood risk under climate change. *Nature Climate Change*, 3, 816-821.
- JAFARZADEGAN, K., MERWADE, V. & SAKSENA, S. 2018. A geomorphic approach to 100-year floodplain mapping for the Conterminous United States. *Journal of Hydrology*, 561, 43-58.
- 555 JONGMAN, B., WARD, P. J. & AERTS, J. 2012. Global exposure to river and coastal flooding: Long term trends and changes. *Global Environmental Change-Human and Policy Dimensions*, 22, 823-835.
- KUMMU, M., DE MOEL, H., WARD, P. J. & VARIS, O. 2011. How Close Do We Live to Water? A Global Analysis of Population Distance to Freshwater Bodies. *Plos One*, 6, 13.
- LEHNER, B., VERDIN, K. & JARVIS, A. 2008. New Global Hydrography Derived From Spaceborne Elevation Data. *Eos, Transactions American Geophysical Union*, 89, 93-94.
- 560 LEYK, S., GAUGHAN, A. E., ADAMO, S. B., DE SHERBININ, A., BALK, D., FREIRE, S., ROSE, A., STEVENS, F. R., BLANKESPOOR, B., FRYE, C., COMENETZ, J., SORICHETTA, A., MACMANUS, K., PISTOLESI, L., LEVY, M., TATEM, A. J. & PESARESI, M. 2019. The spatial allocation of population: a review of large-scale gridded population data products and their fitness for use. *Earth Syst. Sci. Data*, 11, 1385-1409.
- LIU, F., WANG, S., XU, Y., YING, Q., YANG, F. & QIN, Y. 2020. Accuracy assessment of Global Human Settlement Layer (GHSL) built-up products over China. *PLOS ONE*, 15, e0233164.
- 565 LLOYD, C. T., CHAMBERLAIN, H., KERR, D., YETMAN, G., PISTOLESI, L., STEVENS, F. R., GAUGHAN, A. E., NIEVES, J. J., HORNBY, G., MACMANUS, K., SINHA, P., BONDARENKO, M., SORICHETTA, A. & TATEM, A. J. 2019. Global spatio-temporally harmonised datasets for producing high-resolution gridded population distribution datasets. *Big Earth Data*, 3, 108-139.
- 570 LUGERI, N., KUNDZEWICZ, Z. W., GENOVESE, E., HOCHRAINER, S. & RADZIEJEWSKI, M. 2010. River flood risk and adaptation in Europe—assessment of the present status. *Mitigation and Adaptation Strategies for Global Change*, 15, 621-639.
- MCCOOL, S. F., CLARK, R. N. & STANKEY, G. H. 2009. *Water and People: Challenges at the Interface of Symbolic and Utilitarian Values*, DIANE Publishing.
- MONTGOMERY, D. R. & DIETRICH, W. E. 1988. Where do channels begin? 336, 232.
- 575 NARDI, F., ANNIS, A., DI BALDASSARRE, G., VIVONI, E. R. & GRIMALDI, S. 2019. GFPLAIN250m, a global high-resolution dataset of Earth's floodplains. *Scientific Data*, 6, 180309.
- NARDI, F., VIVONI, E. R. & GRIMALDI, S. 2006. Investigating a floodplain scaling relation using a hydrogeomorphic delineation method. 42.
- 580 NEVO, S. 2019. An Inside Look at Flood Forecasting. *Google AI Blog* [Online]. Available from: <https://ai.googleblog.com/2019/09/an-inside-look-at-flood-forecasting.html>.
- NOBRE, A. D., CUARTAS, L. A., HODNETT, M., RENNO, C. D., RODRIGUES, G., SILVEIRA, A., WATERLOO, M. & SALESKA, S. 2011. Height Above the Nearest Drainage - a hydrologically relevant new terrain model. *Journal of Hydrology*, 404, 13-29.
- NOBRE, A. D., CUARTAS, L. A., MOMO, M. R., SEVERO, D. L., PINHEIRO, A. & NOBRE, C. A. 2016. HAND contour: a new proxy predictor of inundation extent. *Hydrological Processes*, 30, 320-333.
- 585 OCHA 2020. Sudan: Floods Flash Update No. 8. OCHA.
- PALACIOS-LOPEZ, D., BACHOFER, F., ESCH, T., HELDENS, W., HIRNER, A., MARCONCINI, M., SORICHETTA, A., ZEIDLER, J., KUENZER, C., DECH, S., TATEM, A. J. & REINARTZ, P. 2019. New Perspectives for Mapping Global Population Distribution Using World Settlement Footprint Products. 11, 6056.
- PAPPENBERGER, F., DUTRA, E., WETTERHALL, F. & CLOKE, H. L. 2012. Deriving global flood hazard maps of fluvial floods through a physical model cascade. *Hydrology and Earth System Sciences*, 16, 4143-4156.
- 590 RUDARI, R., SILVESTRO, F., CAMPO, L., REBORA, N., BONI, G. & HEROLD, C. 2015. IMPROVEMENT OF THE GLOBAL FLOOD MODEL FOR THE GAR 2015.
- SAMPSON, C. C., SMITH, A. M., BATES, P. B., NEAL, J. C., ALFIERI, L. & FREER, J. E. 2015. A high-resolution global flood hazard model. *Water Resources Research*, 51, 7358-7381.
- 595 SCHIAVINA, M. F., S.; MACMANUS, K 2019. GHS population grid multitemporal (1975, 1990, 2000, 2015). European Commission, Joint Research Centre (JRC).
- SMITH, A., BATES, P. D., WING, O., SAMPSON, C., QUINN, N. & NEAL, J. 2019. New estimates of flood exposure in developing countries using high-resolution population data. *Nature Communications*, 10, 1814.



- 600 SMITH, A., SAMPSON, C. & BATES, P. 2015. Regional flood frequency analysis at the global scale. *Water Resources Research*, 51, 539-553.
- STEVENS, F. R., GAUGHAN, A. E., LINARD, C. & TATEM, A. J. 2015. Disaggregating Census Data for Population Mapping Using Random Forests with Remotely-Sensed and Ancillary Data. *Plos One*, 10, 22.
- STOKKOM, H., SMITS, A. & LEUVEN, R. S. E. W. 2005. Flood defense in The Netherlands: A New Era, A New Approach. *Water International - WATER INT*, 30, 76-87.
- 605 STRAHLER, A. N. 1957. Quantitative analysis of watershed geomorphology. *Eos, Transactions American Geophysical Union*, 38, 913-920.
- TANOUE, M., HIRABAYASHI, Y. & IKEUCHI, H. 2016. Global-scale river flood vulnerability in the last 50 years. *Scientific Reports*, 6, 36021.
- TARBOTON, D. G., BRAS, R. L. & RODRIGUEZ-ITURBE, I. J. H. P. 1991. On the extraction of channel networks from digital elevation data. 5, 81.
- 610 TIECKE, T. G. L., XIANMING ZHANG, AMY GROS, ANDREAS LI, NAN YETMAN, GREGORY KILIC, TALIP MURRAY, SIOBHAN BLANKESPOOR, BRIAN PRYDZ, ESPEN B. DANG, HAI-ANH H. 2017. *Mapping the World Population One Building at a Time*.
- TRENDS 2020. Leaving no one off the map: a guide for gridded population data for sustainable development. Thematic Research Network on Data and Statistics.
- 615 TRIGG, M. A., BIRCH, C. E., NEAL, J., BATES, P., SMITH, A., SAMPSON, C., YAMAZAKI, D., HIRABAYASHI, Y., PAPPENBERGER, F., DUTRA, E., WARD, P. J., WINSEMIUS, H. C., SALAMON, P., DOTTORI, F., RUDARI, R., KAPPES, M. S., SIMPSON, A., HADZILACOS, G. & FEWTRELL, T. 2016a. Aggregated fluvial flood hazard output for six Global Flood Models for the African Continent. Research Data Leeds Repository.
- 620 TRIGG, M. A., BIRCH, C. E., NEAL, J. C., BATES, P. D., SMITH, A., SAMPSON, C. C., YAMAZAKI, D., HIRABAYASHI, Y., PAPPENBERGER, F., DUTRA, E., WARD, P. J., WINSEMIUS, H. C., SALAMON, P., DOTTORI, F., RUDARI, R., KAPPES, M. S., SIMPSON, A. L., HADZILACOS, G. & FEWTRELL, T. J. 2016b. The credibility challenge for global fluvial flood risk analysis. *Environmental Research Letters*, 11, 10.
- WALLEMACQ, P., GUHA-SAPIR, D., MCCLEAN, D., CRED & UNISDR 2015. *The Human Cost of Natural Disasters - A global perspective*.
- 625 WARD, P. J., BLAUHUT, V., BLOEMENDAAL, N., DANIELL, J. E., DE RUITER, M. C., DUNCAN, M. J., EMBERSON, R., JENKINS, S. F., KIRSCHBAUM, D., KUNZ, M., MOHR, S., MUIS, S., RIDDELL, G. A., SCHÄFER, A., STANLEY, T., VELDKAMP, T. I. E. & WINSEMIUS, H. C. 2020. Review article: Natural hazard risk assessments at the global scale. *Nat. Hazards Earth Syst. Sci.*, 20, 1069-1096.
- 630 WARD, P. J., JONGMAN, B., SALAMON, P., SIMPSON, A., BATES, P., DE GROEVE, T., MUIS, S., DE PEREZ, E. C., RUDARI, R., TRIGG, M. A. & WINSEMIUS, H. C. 2015. Usefulness and limitations of global flood risk models. *Nature Climate Change*, 5, 712-715.
- WARD, P. J., JONGMAN, B., WEILAND, F. S., BOUWMAN, A., VAN BEEK, R., BIERKENS, M. F. P., LIGTVOET, W. & WINSEMIUS, H. C. 2013. Assessing flood risk at the global scale: model setup, results, and sensitivity. *Environmental Research Letters*, 8, 10.
- 635 WARDROP, N. A., JOCHEM, W. C., BIRD, T. J., CHAMBERLAIN, H. R., CLARKE, D., KERR, D., BENGTTSSON, L., JURAN, S., SEAMAN, V. & TATEM, A. J. 2018. Spatially disaggregated population estimates in the absence of national population and housing census data. 115, 3529-3537.
- WING, O. E. J., BATES, P. D., SAMPSON, C. C., SMITH, A. M., JOHNSON, K. A. & ERICKSON, T. A. 2017. Validation of a 30 m resolution flood hazard model of the conterminous United States. *Water Resources Research*, 53, 7968-7986.
- 640 WINSEMIUS, H. C., VAN BEEK, L. P. H., JONGMAN, B., WARD, P. J. & BOUWMAN, A. 2013. A framework for global river flood risk assessments. *Hydrology and Earth System Sciences*, 17, 1871-1892.
- WORLD BANK 2018. United Nation's Population Division's World Urbanization Prospects: 2018 Revision. In: WORLD BANK (ed.).
- WORLD BANK. 2019. *Over 350,000 Surinamese to Benefit from Flood Risk Management* [Online]. reliefweb. Available: <https://reliefweb.int/report/suriname/over-350000-surinamese-benefit-flood-risk-management> [Accessed].
- 645 YAMAZAKI, D., IKESHIMA, D., SOSA, J., BATES, P. D., ALLEN, G. H. & PAVELSKY, T. M. 2019. MERIT Hydro: A High-Resolution Global Hydrography Map Based on Latest Topography Dataset. *Water Resources Research*, 55, 5053-5073.
- YAMAZAKI, D., IKESHIMA, D., TAWATARI, R., YAMAGUCHI, T., O'LOUGHLIN, F., NEAL, J. C., SAMPSON, C. C., KANAE, S. & BATES, P. D. 2017. A high-accuracy map of global terrain elevations. *Geophysical Research Letters*, 44, 5844-5853.
- 650 YAMAZAKI, D., KANAE, S., KIM, H. & OKI, T. 2011. A physically based description of floodplain inundation dynamics in a global river routing model. *Water Resources Research*, 47, 21.
- ZHOU, X., MA, W., ECHIZENYA, W. & YAMAZAKI, D. 2020. Uncertainty in flood frequency analysis of hydrodynamic model simulations. *Nat. Hazards Earth Syst. Sci. Discuss.*, 2020, 1-31.