

Published in final edited form as:

Nat Ecol Evol. 2018 July ; 2(7): 1042–1043. doi:10.1038/s41559-018-0573-8.

Global gaps in soil biodiversity data

Erin K. Cameron^{1,2,*}, Inês S. Martins^{2,3}, Patrick Lavelle^{4,5,6}, Jérôme Mathieu^{7,8}, Leho Tedersoo⁹, Felix Gottschall^{2,10}, Carlos A. Guerra^{2,3}, Jes Hines^{2,10}, Guillaume Patoine^{2,11}, Julia Siebert^{2,10}, Marten Winter², Simone Cesarz^{2,10}, Manuel Delgado-Baquerizo^{12,13}, Olga Ferlian^{2,10}, Noah Fierer^{12,14}, Holger Kreft¹⁵, Thomas E. Lovejoy¹⁶, Luca Montanarella¹⁷, Alberto Orgiazzi¹⁷, Henrique M. Pereira^{2,3,18}, Helen R. P. Phillips^{2,10}, Josef Settele^{2,19,20}, Diana H. Wall^{21,22}, and Nico Eisenhauer^{2,10}

¹Faculty of Biological and Environmental Sciences, PO Box 65, FI 00014, University of Helsinki, Finland ²German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103, Leipzig, Germany ³Institute of Biology, Martin Luther University Halle-Wittenberg, Am Kirchtor 1, 06108, Halle (Saale), Germany ⁴UPMC Université Paris 06, iEES Paris, 32 Av. Henri Varagnat, 93143 Bondy Cedex, France ⁵IRD, iEES Paris, Centre IRD Ile de France, 32 Av. Henri Varagnat, 93143 Bondy Cedex, France ⁶Centro Internacional de Agricultura Tropical (CIAT), TSBF_LAC, ap aereo 6713, Cali, Colombia ⁷Sorbonne Universités, UPMC Univ. Paris 06, IRD, CNRS, INRA, UPEC, University Paris Diderot ⁸Institute of Ecology and Environmental Sciences, iEES Paris, 4 place Jussieu, 75005 Paris, France ⁹Natural History Museum, University of Tartu, 14A Ravila, 50411 Tartu, Estonia ¹⁰Institute of Biology, Leipzig University, Deutscher Platz 5e, 04103, Leipzig, Germany ¹¹Center for Environmental Research and Technology, General and Theoretical Ecology, University of Bremen, Leobener Str, 28359 Bremen, Germany ¹²Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO 80309 ¹³Departamento de Biología y Geología, Física y Química Inorgánica, Escuela Superior de Ciencias Experimentales y Tecnología, Universidad Rey Juan Carlos, Calle Tulipán Sin Número, Móstoles 28933, Spain ¹⁴Department of Ecology and Evolutionary Biology, University of Colorado, Boulder CO 80309 ¹⁵Biodiversity, Macroecology & Biogeography, University of Goettingen, Göttingen, Germany ¹⁶Department of Environmental Science and Policy, George Mason University, Fairfax, VA 22030, USA ¹⁷European Commission, Joint Research Centre (JRC), Sustainable Resources Directorate, Land Resources Unit, Ispra, Italy ¹⁸Infraestruturas de Portugal Biodiversity Chair, CiBio/InBIO, Universidade do Porto, 4485-661, Vairão, Portugal ¹⁹Helmholtz Centre for Environmental Research, UFZ, Department of Community Ecology, Theodor-Lieser-Str. 4, 06120 Halle, Germany ²⁰Institute of Biological Sciences, University of the Philippines Los Baños, College, 4031 Laguna, Philippines ²¹Department of Biology, Colorado State University, Fort Collins, CO, USA ²²School of Global Environmental Sustainability, Colorado State University, Fort Collins, CO, USA

*Correspondence to: erin.cameron@helsinki.fi.

To the Editor

Soil biodiversity represents a major terrestrial biodiversity pool, supports key ecosystem services, and is under pressure from human activities¹. Yet, soil biodiversity has been neglected from many global biodiversity assessments and policies. This omission is undoubtedly related to the paucity of comprehensive information on soil biodiversity, particularly on larger spatial scales. Information on belowground species distributions, population trends, endemism, and threats to belowground diversity is important for conservation prioritization, but is practically nonexistent. As a consequence, much of our understanding of global macroecological patterns in biodiversity, as well as mapping of global biodiversity hotspots, has been based on aboveground taxa (such as plants²) and has not considered the functionally important, but less visible, biodiversity found in soil.

We mapped the study sites from existing global datasets on soil biodiversity (soil macrofauna³, fungi⁴, and bacteria⁵) to examine key data gaps worldwide (Fig. 1). Our map indicates significant gaps in soil biodiversity data remain across northern latitudes, including most of Russia and Canada. Data are also lacking from much of central Asia and central Africa (for example, the Sahara desert), as well as many tropical regions. The higher density of soil biodiversity sampling sites in Europe and the United States is similar to patterns observed for data on terrestrial bird, mammal, and amphibian species⁶, as well as plants⁷. Yet, in such aboveground datasets, the gaps in understudied regions are much less pronounced than in the soil biodiversity datasets shown here. The comparative lack of soil biodiversity data across these regions limits our ability to examine global macroecological patterns and to quantify potential mismatches between aboveground and soil biodiversity. The potential for such mismatches (areas with high aboveground diversity, but low soil biodiversity, or vice versa) may be substantial, as some evidence suggests that plant species richness declines more rapidly towards the North Pole than fungal species richness, which reaches a plateau⁴.

Soil ecologists are increasingly conducting their own large-scale assessments (such as the African Soil Microbiology Project⁸) and additional databases on soil biodiversity are beginning to be developed to address knowledge gaps⁹, in part through the Global Soil Biodiversity Initiative. However, increased efforts to fill these gaps and to compile additional global datasets on other soil taxa (for example, mesofauna) are needed to allow more detailed analyses of soil biodiversity at broad spatial scales. Of major concern is the lack of a global consensus on sampling strategies and methodological approaches to assess soil biodiversity, which in many cases, makes it challenging to compare datasets directly. Furthermore, greater cooperation with conservation biologists and policy makers is needed to better integrate soil biodiversity into global policies. For instance, soil biodiversity should be more explicitly considered in the post-2020 global biodiversity framework¹⁰ that will follow the Strategic Plan for Biodiversity 2011-2020 and in future assessments of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services¹¹.

These evident gaps in soil biodiversity data restrict our ability to develop policies to protect soil biodiversity. We argue that addressing these data gaps will ultimately benefit human

well-being¹ and provide an impetus for increased policy-relevant research on soil biodiversity.

Acknowledgements

We thank all contributors to the Macrofauna database, especially Emmanuel Laped, George Brown, and Carlos Fragoso. We thank Mar Cabeza for helpful suggestions on an earlier version of this manuscript. We acknowledge support by the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig (funded by the German Research Foundation, FZT 118), the European Research Council (ERC Starting Grant 677232, ECOWORM), the Academy of Finland (285882), and the Natural Sciences and Engineering Research Council of Canada (postdoctoral fellowship to E.K.C.).

References

1. Orgiazzi, A., et al., editors. Global Soil Biodiversity Atlas. Publications Office of the European Union; 2016.
2. Kreft H, Jetz W. Proc Natl Acad Sci. 2007; 104:5925–5930. [PubMed: 17379667]
3. Mathieu, J., Lavelle, P. Global Soil Biodiversity Atlas. Orgiazzi, A., et al., editors. Publications Office of the European Union; 2016.
4. Tedersoo L, et al. Science. 2014; 346:1256688. [PubMed: 25430773]
5. Delgado-Baquerizo M, et al. Science. 2018; 359:320–325. [PubMed: 29348236]
6. Meyer C, et al. Nature Communications. 2015; 6:8221.
7. Meyer C, et al. Ecology Letters. 2016; 19:992–1006. [PubMed: 27250865]
8. Wild S, et al. Nat News. 2016; 539:152.
9. Ramirez KS, et al. Front Ecol Evol. 2015; 3:91.
10. CBD. 2017. <https://www.cbd.int/doc/press/2017/pr-2017-12-16-sbstta-en.pdf>
11. IPBES. Global assessment. <https://www.ipbes.net/deliverables/2c-global-assessment>




Fig. 1. Global distribution of sampling sites for soil bacteria⁵ (yellow; n = 237), soil macrofauna³ (red; n = 2163), and soil fungi⁴ (blue; n = 326).