

GLUCOSE CHALLENGE STIMULATES REACTIVE OXYGEN SPECIES (ROS) GENERATION BY LEUCOCYTES

PRIYA MOHANTY, WAEL HAMOUDA, RAJESH GARG, AHMAD ALJADA, HUSAM GHANIM, PARESH DANDONA

Division of Endocrinology, Diabetes and Metabolism; State University of New York at Buffalo and Kaleida Health, 3 Gates Circle, Buffalo, NY 14209

Abstract

Diabetes mellitus is associated with increased ROS generation, oxidative injury and obesity. To elucidate the relationship between nutrition and ROS generation, we have investigated the effect of glucose challenge on ROS generation by leucocytes, p47^{phox} protein, a key protein in the enzyme NADPH oxidase and α -tocopherol levels. Blood samples were drawn from 14 normal subjects prior to, at 1, 2 and 3 h following ingestion of 75 g glucose. ROS generation by polymorphonuclear leucocytes (PMNL) and mononuclear cells (MNC) increased to a peak of $244 \pm 42\%$ and $233 \pm 34\%$ of the basal respectively at 2h. The levels of p47^{phox} in MNC homogenates increased significantly at 2 h and 3 h after glucose intake. α -Tocopherol levels decreased significantly at 1 h, 2 h and 3 h. We conclude that glucose intake stimulates ROS generation and p47^{phox} of NADPH oxidase; increases oxidative load and causes a fall in α -tocopherol concentration.

Introduction

Diabetes mellitus is associated with an increase in ROS generation by mononuclear cells¹ and an increased oxidative load resulting in oxidative damage to lipids², proteins³ and DNA⁴. Acute hyperglycemia has been shown to result in an increase in blood pressure, which is prevented by antioxidants⁵; this suggests that acute hyperglycemia probably causes increased generation of ROS and thus a diminished bioavailability of nitric oxide (NO)⁵⁻⁷. We have also shown that ROS generation by leucocytes falls after weight loss (unpublished data). This raises the possibility that nutrition may modulate ROS generation. These facts suggest that chronic hyperglycemia (diabetes mellitus) and acute hyperglycemia may result in increased ROS generation (unpublished data).

We therefore, embarked on a study to investigate the effect of glucose challenge on leukocytic ROS generation. We also investigated the levels of p47^{phox}, the key protein component of NADPH oxidase⁸⁻⁹ to determine whether the NADPH units increase in number following glucose challenge. Plasma concentration of thiobarbituric acid reacting substances (TBARS) was measured since it is an index of lipid peroxidation.

Subjects, Materials and Methods

Fourteen normal subjects (7 males and 7 females; age range: 25-42 years; weight range: 53-80 kg; mean BMI: $24.2 \pm 0.6 \text{ kg/m}^2$) participated in the study. They were given 75 g of glucose dissolved in 300 ml water (Glucola) to drink over 5 minutes. Controls (n=6) drank 300 ml water containing saccharin. Blood samples were obtained at 0,1,2 and 3h. PMNL and MNC isolation and ROS generation were carried out as previously described¹⁰. p47^{phox} was measured by Western blotting using anti-

p47^{phox} antibody from Transduction Labs (Lexington, KY). TBARS were measured as described by Ohkawa et al¹¹. α -Tocopherol was measured in plasma by HPLC as described previously¹². Insulin levels were determined using an ELISA kit from Diagnostic Systems Laboratories Inc. (Webster, TX).

Statistical analysis was carried out by one way analysis of variance (ANOVA) for repeated measures. The data on ROS generation were analyzed by reducing the basal readings to 100%. Values are expressed as mean \pm S.E.

Results

The mean plasma glucose and insulin concentrations peaked at 1h to $108.5 \pm 9.4 \text{ mg/dl}$ and $66.6 \mu\text{U/ml}$ (Table 1). The mean ROS generation by PMNL in the fasting state (basal) was $117 \pm 19 \text{ mV}$ (100%). It increased to a peak of $244 \pm 42\%$ over the basal (Figure 1)($p < 0.005$). The mean ROS generation by MNC in the fasting state (basal) was $213 \pm 40 \text{ mV}$ (100%); it increased to a peak of $233 \pm 34\%$ over the basal at 2h (Figure 2)($p < 0.001$). There was no change in ROS generation by leucocytes following a drink of 300 ml water or saccharine containing water. The levels of p47^{phox} in MNC homogenates, representative of NADPH oxidase, did not alter at 1 h, but increased significantly at 2 h and 3 h after glucose intake (Figure 3) with a peak effect at 2 h.

TBARS increased from a basal level of $1.07 \pm 0.10 \mu\text{mol/L}$ (100%) to $116.7 \pm 7.2\%$ of basal at 1 h, $120.6 \pm 8.3\%$ of basal at 2 h and $108 \pm 12.9\%$ of basal at 3 h ($p = 0.065$) (Figure 4). The mean fasting α -tocopherol level was $8.7 \pm 0.4 \mu\text{g/ml}$ (basal, 100%). α -tocopherol levels fell to $99\% \pm 0.4\%$, $96.1\% \pm 0.6\%$ and $96 \pm 0.6\%$ of the basal at 1, 2 and 3 h respectively ($p < 0.001$) (figure 5).




Figure 1: ROS generation by PMNL prior to and following glucose challenge (75 g) at 1, 2 and 3 h. Results are expressed as % increase over fasting basal (100%). Note that the ROS generation increased significantly at 1 h and increased further at 2 h and then decreased at 3 h.




Figure 2: ROS generation by MNC prior to and following glucose challenge (75 g) at 1, 2 and 3 h. Results are expressed as % increase over fasting basal (100%). Note that the ROS generation is increased significantly at 1 h and increased further at 2 h and then declined at 3 h.




Figure 3: A) Western Blot showing the relative expression of p47^{phox} in MNC. Note that p47^{phox} expression is induced at 1 h after glucose intake. Maximum induction is observed at 2 h. This blot is a representative of four different experiments with four different subjects. B) Densitometric semi-quantitative analysis of p47^{phox} protein levels in MNC. Note the significant increase at 2 h ($p < 0.05$).




Figure 4: Change in TBARS at 1, 2 and 3 h after glucose challenge. Results are expressed as mean \pm S.E. increase over the basal value (100%). Note that the increase at 1 h and 2 h was significant with paired t-test.




Figure 5: Change in α -tocopherol levels at 1, 2 and 3 h after glucose challenge. Values are expressed as mean \pm S.E. % of baseline, baseline being normalized as 100%. Note that the fall in α -tocopherol was significant at 2 h and 3 h.

Discussion

Our data demonstrate clearly for the first time that a challenge with glucose results consistently in an increase in ROS generation by leucocytes in normal subjects. No increase was observed following the intake of 300 ml of water containing saccharine. There was also an increase in the expression of p47^{phox}, the key protein component of NADPH oxidase. The increase in p47^{phox} was observed at 2 and 3 h with no significant change at 1 h. Since ROS generation by MNC increased at 1 h, while p47^{phox} increased only at 2 and 3 h, it would appear that the initial increase in ROS generation is a function of increased activity of pre-existent NADPH oxidase units, while the further increases at 2 and 3 h may receive a contribution from the formation of additional new NADPH oxidase units. Clearly, there is no direct quantitative relationship between p47^{phox} expression and ROS generation; however, there is a qualitative relationship between the two and thus, the increase of 70% in p47^{phox} reflecting an increase in the enzyme NADPH oxidase, may contribute to increased superoxide generation. It is possible that glucose is an important modulator of p47^{phox} gene expression.

There is a recent demonstration of a depletion in antioxidant reserve following an oral glucose tolerance test in normal subjects and in patients with type 2 diabetes^{13,14}. The previous studies did not investigate ROS generation, although the depletion of antioxidant reserve (α -tocopherol) is consistent with an increase in ROS generation. Our study demonstrates increased ROS generation by PMNL and MNC, it is also possible that increased ROS generation is more generalized. Glucose and/or food intake stimulate the metabolic rate¹⁵ and therefore probably stimulate metabolic activity at the cellular and mitochondrial level, which may result in increased ROS generation. Triiodothyronine administration also stimulates metabolism and has recently been shown by us to increase ROS generation by PMNL and MNC¹⁶.

We have recently demonstrated that carvedilol (a β blocker) administration to normal subjects for 1 week results in a significant decrease in ROS generation by PMNL and MNC with a parallel reduction in oxidative damage to amino acids¹⁷. Thus, β -adrenergic modulation of NADPH oxidase is a possibility. It is therefore possible that nor-epinephrine, known to be released following glucose intake, may stimulate ROS generation by leucocytes. Another leucocyte membrane enzyme, Na-K ATPase, has previously been shown by us to be stimulated by glucose challenge¹⁸ and T₃ administration¹⁹ and to be inhibited by β -blockade²⁰.

The possibility that insulin may play a role in the stimulation of ROS generation also has to be considered.

MNC are known to have insulin receptors²¹ but glucose transport in MNC appears to be primarily mediated by GLUT-1 rather than GLUT-4²². There is at least one study demonstrating insulin stimulated GLUT-1 mediated glucose transport by MNC²². On the other hand, insulin may have a direct effect on ROS generation. Indeed, we have now shown that insulin infusion inhibits ROS generation by leucocytes²³. Thus, it is unlikely that insulin contributed to the stimulation of ROS generation following glucose challenge.

The small (4%) but significant fall in α -tocopherol concentration following a 75 g glucose (\approx 300 k calories) challenge is consistent with an increase in oxidative load due to increased ROS generation. All subjects tested demonstrated this fall. These data are consistent with those of Ceriello et al¹³, who demonstrated a similar fall in α -tocopherol after glucose challenge. It is unlikely that these small changes affect ROS generation, since much larger changes (180%) in concentrations of α -tocopherol associated with large doses of this vitamin have been shown to alter ROS (60% inhibition) generation over a prolonged period of vitamin E administration²⁴.

The link between nutrition and oxidative damage is an important one since oxidative damage of lipids, the LDL particle in particular, may contribute to atherosclerosis²⁵. Oxidized LDL, which is internalized via the scavenger receptor of the circulating monocytes and the subendothelial macrophage which form the foam cells; and the latter cells form the fatty streak, the initial lesion of atherosclerosis²⁶. Our work opens the way for the investigation of the effect of other macronutrients on ROS generation and oxidative damage.

In conclusion, glucose challenge induces a consistent increase in ROS generation by leucocytes, an increase in p47^{phox}, an integral protein component of NADPH oxidase, and a decrease in α -tocopherol concentrations in plasma.

The authors acknowledge the support of the William G. McGowan Charitable Fund (Washington, DC).

References:

1. Hiramatsu K, Arimori S. Increased superoxide production by mononuclear cells of patients with hypertriglyceridemia and diabetes. *Diabetes* 1988; 37:832-7.
2. Jain SK. Hyperglycemia can cause membrane lipid peroxidation and osmotic fragility in human red blood cells. *J Biol Chem* 1989; 264:340-5.
3. Aljada A, Thusi K, Armstrong D, Nicotera T, Dandona P. Increased carbonylation of proteins in diabetes. *Diabetes* 1995; 44 suppl: I 13A:266-333.

4. Dandona P, Thusu K, Snyder B, Makowski J, Armstrong D, Nicotera T. Oxidative damage to DNA in diabetes mellitus. *Lancet* 1996; 347:444-5.

5. Marfella R, Giovanni V, Acampora R, La Marca C, Giunta R, Lucarelli C, Paolisso G, Ceriello A, Giugliano D. Glutathione reverses systemic hemodynamic changes induced by acute hyperglycemia in healthy subjects. *Amer J Physiol* 1995; 01 93 -1849; E I 167-E 1173.

6. Giugliano D, Marfella R, Coppola L, Verrazzo G, Acampora R, Giunta R, Nappo F, Lucarelli C, D'Onofrio F. Vascular effects of acute hyperglycemia in humans are reversed by L-Arginine. *Circulation* 1997; 95:1783-90.

7. Paolisso G, Giugliano D. Oxidative stress and insulin action: is there a relationship? *Diabetologia* 1996; 39:357-263.

8. Serra MC, Bellavite P, Davoli A, Bannister JV, Rossi F. Isolation from neutrophil membranes of a complex containing active NADPH oxidase and cytochrome b-245. *Biochim Biophys Acta* 1984;788:138-46.

9. Heyworth PG, Cumutte JT, Nauseef WM, Volpp BD, Pearson DW, Rosen H, Clark RA. Neutrophil nicotinamide adenine dinucleotide phosphate oxidase assembly. Translocation of p47^{phox} and p67^{phox} requires interaction between p47^{phox} and cytochrome b558. *J Clin Invest* 1991; 87(1):352-6.

10. Dandona P, Thusu K, Hafeez R, Abdel-Rahman E, Chaudhuri A. Effect of hydrocortisone on oxygen free radical generation by mononuclear cells. *Metabolism* 1998;47:788-91.

11. Okhawa H, Ohishi N, Yagi K: Reaction of linoleic acid hydroperoxide with thiobarbituric acid. *J Lipid Res* 1978; 19:1053-1057.

12. Browne RW, Armstrong D: Simultaneous determination of serum retinol, tocopherols and carotenoids by HPLC. *Methods Mol Biol* 1998; 108:269-275.

13. Ceriello A, Bortolotti N, Crescentini A et al: Antioxidant defences are reduced during oral glucose tolerance test in normal and non-insulin-dependent subjects. *Eur J Clin Invest* 1998;28:329-33.

14. Ceriello A, Bortolotti N, Motz E et al. Meal-generated oxidative stress in type2 diabetic patients. *Diabetes Care* 1998;21:1529-33.

15. Wells S, Lilavivathana V, Campbell RG. Increased plasma norepinephrine concentrations and metabolic rates following glucose ingestion in man. *Metabolism* 1980;29:806-9.

16. Magsino C, Hamouda W, Afzal A, Aljada A, Dandona P. The effect of triiodothyronine on reactive oxygen species generation by leucocytes, indices of oxidative damage and antioxidant reserve. *Metabolism* 2000 (in press).

17. Dandona P, Karne R, Ghanim H, Hamouda W, Aljada A, Magsino C. Carvedilol inhibits oxygen species (ROS)

generation by leukocytes and oxidative damage to amino acids. *Circulation* 2000; 101:122-124.

18. Turaihi K, Baron DN, Dandona P. Effect of glucose intake on human leucocyte 86Rb influx and [3H]-ouabain binding. *Metabolism* 1988;37:171-4.

19. Turaihi K, Khan FA, Baron DN, Dandona P. Effect of short term triiodothyronine administration on human leucocyte Rb(k) influx and Na efflux. *J Clin Endocrinol Metab* 1987; 65(5):1031-4.

20. Turaihi K, Perelman M, Baron DN, Dandona P. Effect of beta-blockade and subsequent triiodothyronine administration on human leucocyte Na-K ATPase. *Metabolism* 1988; 37(5):499-501.

21. Flier JS, Minaker KL, Landsberg L, Young JB, Pallotta J, Rowe JW. Impaired in vivo insulin clearance with severe target-cell resistance to insulin. *Diabetes* 1982;31:132-5.

22. Daneman D, Zinman B, Elliott ME, Bilan PJ, Klip A. Insulin-stimulated glucose transport in circulating mononuclear cells from nondiabetic and IDDM subjects. *Diabetes* 1992;41:227-34.

23. Dandona P, Aljada A, Mohanty P, Ghanim H, Hamouda W, Assian E, Chowhan S. Insulin inhibits ROS generation, p47^{phox} and NFκB in mononuclear cells and reduces plasma sICAM-1 and MCP-1 in obese subjects. *Endo 2000, the 82nd Annual Meeting of the Endocrine Society* (abstract).

24. Devaraj S, Li D, Jialal I. The effects of alpha tocopherol supplementation on monocyte function. Decreased lipid oxidation, interleukin 1 beta secretion and monocyte adhesion to endothelium. *J Clin Invest* 1996; 98:756-763.

25. Steinberg D, Parathasarathy S, Carew TE, Khoo JC, Witztum JL. Beyond cholesterol. Modifications of low-density lipoprotein that increases its atherogenicity. *N Eng J Med* 1989;320:915-24.

26. Ross R. Atherosclerosis-an inflammatory disease. *N Engl J Med* 1999;340:115-126.

	0 hour	1 hour	2 hours	3 hours
Glucose (mg/dl)	83 ± 7.8	108 ± 9.4	107 ± 8.8	70 ± 3.4
Triglycerides (mg/dl)	142 ± 28	136 ± 22	133 ± 22	151 ± 50
Cholesterol (mg/dl)	195 ± 11	191 ± 10	189 ± 10	197 ± 1
Insulin (µU/ml)	7.9 ± 1.8	66.6 ± 15.7	32.8 ± 10.3	16.5 ± 9

Table 1: Glucose, triglyceride, cholesterol, and insulin concentrations prior to and following glucose intake.