

Incubation with ethanol, which is readily oxidized, also causes increases in these enzyme activities. These increases occur more quickly than with acetate or pyruvate, probably because ethanol can, in being oxidized to acetate, serve as an immediate source of energy (Maitra & Estabrook, 1967).

Thus this yeast has an active tricarboxylic acid cycle and oxidative pathways. However, there is a lag before acetate is oxidized linearly. Inhibition of cytoplasmic protein synthesis completely prevents adaptation, and the activities of a number of relevant enzymes, especially isocitrate lyase, increase during adaptation. Consequently it is probable that the activity of some enzyme or enzymes, such as those above, is directly or indirectly limiting for acetate oxidation but not for the oxidation of ethanol or pyruvate.

Gosling, J. P. & Duggan, P. F. (1968). *Biochem. J.* **110**, 19P.

Maitra, P. K. & Estabrook, R. W. (1967). *Arch. Biochem. Biophys.* **121**, 129.

Roodyn, D. B. & Wilkie, D. (1968). *The Biogenesis of Mitochondria*. London: Methuen and Co. Ltd.

Siegel, M. R. & Sisler, H. D. (1965). *Biochim. biophys. Acta*, **103**, 558.

### Glycopeptide Linkages in a Phosphomannan Peptide from Yeast Cell Walls

By T. N. CAWLEY and R. LETTERS. (*Department of Biochemistry, University College, Dublin, and The Research Laboratory, Arthur Guinness, Son and Co. (Dublin) Ltd., St James's Gate, Dublin 8, Irish Republic*)

Glycopeptides containing mannose have been extracted from yeast cell walls by anhydrous ethylenediamine (Korn & Northcote, 1960). Subsequent purification (Sentandreu & Northcote, 1968) by treatment with Pronase and gel filtration yielded a high-molecular-weight mannan-peptide rich in serine and threonine. Sentandreu & Northcote (1968) were able to demonstrate, from their investigations of the  $\beta$ -elimination reaction of the glycopeptide in dilute alkali, that one of the linkages between carbohydrate and amino acids is an *O*-mannosyl bond to serine and threonine. The present report deals with similar investigations on the glycopeptide linkages of a mannan-peptide that was liberated directly from isolated yeast cell walls by Pronase digestion (Cawley & Letters, 1968).

Amino acid analyses of yeast cell wall, and of the phosphoglycopeptide released from it by Pronase treatment, show that Pronase treatment results in an enrichment of serine and threonine in the phosphoglycopeptide. Treatment of the phosphoglycopeptide (70 mg.) with 0.1 M-KOH (10 ml.) at room temperature for 24 hr. results in a loss of 75%

of the serine and 79% of the threonine initially present. These losses are due to a  $\beta$ -elimination reaction that produces dehydro derivatives of serine and threonine with a characteristic u.v. absorption at 240 nm. During the elimination reaction a steady increase in the extinction at this wavelength was observed. When the phosphoglycopeptide (100 mg.) was treated with sodium borohydride (30 mg.) in 0.1 M-KOH (10 ml.) for 24 hr. at room temperature, and then hydrolysed to liberate amino acids, we were able to detect a new amino acid, identified as  $\alpha$ -aminobutyric acid by high-voltage paper electrophoresis. This provides confirmatory evidence for formation of dehydrothreonine during the  $\beta$ -elimination. The decrease in the proportion of serine and threonine in the glycopeptide after treatment with alkaline borohydride and the simultaneous increase in the proportion of alanine together with the appearance of  $\alpha$ -aminobutyric acid indicate that serine and threonine are involved in linkages through their hydroxyl groups. The alkoxide moiety eliminated from the phosphoglycopeptide by 0.1 M-KOH was found to be a mixture of free mannose and mannose oligosaccharides, thus showing that the serine and threonine hydroxyl groups are involved in glycosidic linkages. Yeast mannan, which can be obtained by alkali treatment of cell wall, is found to be rich in glucosamine. However, Pronase digestion of the cell wall does not solubilize glucosamine but concentrates it in the insoluble residue. The residue obtained after continuous digestion with Pronase for 7 days contained 0.2% nitrogen. An amino acid analysis showed that alanine and glucosamine each accounted for approx. 10% of the total  $\alpha$ -amino nitrogen, and aspartic acid, glutamic acid and serine each accounted for a further 8%.

These results show that the phosphomannan-peptide released by direct treatment of yeast cell walls with Pronase contains glycopeptide linkages that are similar to those found in the glycopeptide investigated by Sentandreu & Northcote (1968). Therefore the direct treatment of isolated yeast cell walls with Pronase provides a simpler and quicker method for obtaining large amounts of phosphomannan-peptide for chemical studies.

We thank Dr K. Morgan for the amino acid analyses. T.N.C. acknowledges the receipt of a Guinness Research Scholarship.

Cawley, T. N. & Letters, R. (1968). *Biochem. J.* **110**, 9P.

Korn, E. D. & Northcote, D. H. (1960). *Biochem. J.* **75**, 12.

Sentandreu, R. & Northcote, D. H. (1968). *Biochem. J.* **109**, 419.