

THE GOLDBERGER-TREIMAN RELATION AND
CHIRAL SYMMETRY BREAKING*

H. F. Jones†

Stanford Linear Accelerator Center
Stanford University, Stanford, California 94305

and

M. D. Scadron

Physics Department, University of Arizona
Tucson, Arizona 85721

ABSTRACT

We review the dispersion-theory calculation of the Goldberger-Treiman discrepancy $\Delta \equiv 1 - mg_A / (f_\pi g)$. Our estimate is still some way from the experimental value. The latter and its SU(3) counterparts can be used to determine the chiral-symmetry-breaking parameter c appearing in the $(3, \bar{3}) + (\bar{3}, 3)$ Hamiltonian $u_0 + cu_8$. We find $c \approx -1$ in agreement with its determination from phenomenological σ terms.

(Submitted to Phys. Rev.)

* Work supported in part by the U.S. Atomic Energy Commission and by the U.S. National Science Foundation, grant number GP 43714, respectively.

† On leave of absence from Physics Department, Imperial College, London SW7, England.

I. INTRODUCTION AND FORMALISM

The validity or otherwise of the proposed $(3, \bar{3}) + (\bar{3}, 3)$ model¹ for the breaking of the chiral $SU(3) \times SU(3)$ symmetry can be tested in various ways. In a previous paper² we have compared the predictions of the model against phenomenological σ terms, concluding that the generally large values of these quantities can be reconciled with the model only if the parameter c occurring in the symmetry-breaking Hamiltonian $H' = u_0 + cu_8$ takes the value $c \approx -1$, rather than the value $c \approx -1.25$ derived from the pseudoscalar meson mass formula.

An alternative means of probing the value of c is provided by the Goldberger-Treiman (G-T) relation³ and its $SU(3)$ counterparts. This is connected to the above determination in two ways. First, the discrepancy between the experimental value $f_\pi \approx 93$ MeV found from $\pi \rightarrow \mu\nu$ decay⁴ and the theoretical prediction $f_\pi^{GT} = m_N g_A / g \approx 87$ MeV can be reduced slightly if the coupling of the $\sigma(\epsilon)$ particle to pions is large, which, in the σ model, directly follows from the large value⁵ (≈ 70 MeV) of the $\pi N \sigma$ term $\sigma(\pi N)$. Second, the $SU(3)$ G-T discrepancies afford an independent determination of c provided that the hyperon coupling constants are known. In fact, as we shall see, the determination is rather insensitive to $SU(3)$ breaking of these couplings and indeed leads to $c \approx -1$, but cannot credibly be stretched to accommodate the canonical value $c \approx -1.25$.

Both of these aspects of the G-T relation have been discussed in the literature before. However, previous evaluations of the G-T discrepancy by means of dispersion relations^{6,7} are marred by errors and do not use the connection between the σ meson coupling constants and the $\pi N \sigma$ term. Our treatment of the $SU(3)$ relations is more detailed than that of Dashen⁸ and has the advantage of a now clearer experimental situation as regards the hyperon coupling constants.

The objects of our study will be the G-T discrepancies Δ , which for πNN is defined as⁶

$$\Delta_{\pi NN} = 1 - \frac{mg_A}{f_\pi g} \quad (1)$$

Numerically this discrepancy is found to be

$$\Delta_{\pi NN} = 0.058 \pm 0.013 \quad (2)$$

using the values^{9, 10} $m \equiv m_N = 938.9$ MeV, $g_A \equiv g_A(0) = 1.25$, $f_\pi = 93.0$ MeV, and¹¹ $g \equiv g_{\pi NN} = 13.40$.

The formalism for the first aspect of our work has been set up by Pagels.⁶ One begins by defining the nucleon-nucleon matrix element of the axial-vector current as^{12, 13}

$$\langle N(p') | A_\mu^j(x) | N(p) \rangle = i \bar{u}(p') \left(\frac{1}{2} \tau^j \right) \left[g_A(q^2) \gamma_\mu + h_A(q^2) q_\mu \right] \gamma_5 u(p) e^{-iq \cdot x} \quad (3)$$

where $q = p' - p$.

Taking the divergence one has

$$\langle N(p') | \partial \cdot A^j(0) | N(p) \rangle = D(q^2) \bar{u}(p') \tau^j \gamma_5 u(p) \quad (4)$$

where

$$D(q^2) = m g_A(q^2) + \frac{1}{2} q^2 h_A(q^2) \quad (5)$$

which can be separated into a contribution from the pion pole plus the background:

$$D(q^2) = D_{\text{pole}}(q^2) + \bar{D}(q^2) \quad (6)$$

Evaluating the pole term at $q^2 = 0$ leads to

$$D(0) = m g_A = f_\pi g + \bar{D}(0) \quad (7)$$

One then assumes an unsubtracted dispersion relation for $\bar{D}(q^2)$:

$$\bar{D}(0) = \frac{1}{\pi} \int_{9\mu^2}^{\infty} \text{Im } \bar{D}(t) \frac{dt}{t} \quad (8)$$

The PCAC relation, $\partial \cdot A = \mu^2 f_\pi \phi$, connects $\bar{D}(q^2)$ to the pionic form factor $K(q^2)$ of the nucleon, leading to

$$\text{Im } \bar{D}(t) = \mu^2 f_\pi g \frac{\text{Im } K(t)}{\mu^2 - t}, \quad (9)$$

where $K(t)$ is normalized to unity at $q^2 \equiv t = \mu^2$.

Thus Eq. (8) can be rewritten as

$$\bar{D}(0) = f_\pi g \frac{\mu^2}{\pi} \int_{9\mu^2}^{\infty} \frac{dt}{t(\mu^2 - t)} \text{Im } K(t), \quad (10)$$

and Eq. (7) as

$$\Delta_{\pi NN} \equiv 1 - \frac{m g_A}{f_\pi g} = \frac{\mu^2}{\pi} \int_{9\mu^2}^{\infty} \frac{dt}{t(t - \mu^2)} \text{Im } K(t). \quad (11)$$

This can be interpreted as a once-subtracted dispersion relation in the pionic form factor of the nucleon, Δ being thus a measure of the variance of this quantity between $q^2 = \mu^2$ and $q^2 = 0$:

$$\Delta_{\pi NN} = K(\mu^2) - K(0) \quad (12)$$

The contributions to the dispersion integral will be examined in Section II, where the differences with previous work will be carefully pointed out. The value of the σ term $\sigma(\pi N)$ and the form of the low-energy approximation to the $L_t = 0$ πN amplitude will prove to be important, as discussed in Section III.

The second aspect of our work has been discussed briefly by Dashen.^{8,14} Here one evaluates the non-pole part of $D(0)$ in Eq. (7) in terms of the structure of the chiral-symmetry breaking Hamiltonian:

$$\bar{D}(0) = \frac{1}{2} \langle N, \partial \cdot \bar{A}^3(0) | N \rangle = - \frac{1}{2} i \langle N | [Q_5^3, H'] | N \rangle. \quad (13)$$

In the $(3, \bar{3}) + (\bar{3}, 3)$ model H' takes the form

$$H' = u_0 + cu_8 \quad (14)$$

and the relevant commutators in Eq. (13) are

$$[Q_5^i, u^j] = -i d^{ijk} v_k \quad (15)$$

Thus, from this point of view

$$\bar{D}(0) = - \frac{\sqrt{2+c}}{2\sqrt{3}} (\bar{v}_\pi)_{NN} \quad (16)$$

and so

$$\Delta_{\pi NN} = \frac{1}{f_\pi g} \frac{\sqrt{2+c}}{2\sqrt{3}} (\bar{v}_\pi)_{NN} \quad (17)$$

The SU(3) analogues of this relation can be used to eliminate the unknown quantity $(\bar{v}_\pi)_{NN}$ in favor of hyperon discrepancy functions $\Delta_{KN\Sigma}$, Δ_{KNA} . This analysis is discussed in Section IV, where, in particular, the insensitivity of the derived relation to SU(3) breaking is pointed out.

Following these calculations a discussion of the experimental discrepancy of $\Delta_{\pi NN}$ and the status of the GMOR model is given in Section V.

II. SATURATION OF THE DISPERSION RELATION

We now re-examine estimates^{6,7} of $\Delta_{\pi NN}$ obtained by saturating (11) with the lowest lying intermediate states. Barring the existence of a heavy pion or a large 3π enhancement,¹⁵ it has been shown¹⁶ that the contribution from the 3π intermediate state is completely negligible. Furthermore the high-energy contribution, i. e. the contribution from values of $t > 4 m^2$, has been shown¹⁷ to be bounded by 0.01. Thus we can restrict our attention to the resonant $\rho\pi$ and $\sigma\pi$ intermediate states and take the upper limit of integration in Eq. (11) as $4 m^2$. The contributions are shown diagrammatically in Fig. 1.




FIG. 1

In the absence of a complete representation of the right hand amplitudes $\pi\sigma \rightarrow N\bar{N}$ and $\pi\rho \rightarrow N\bar{N}$, we can only attempt an estimate thereof using s-, u- and t-channel pole terms.

We consider first, then, the s- and u-channel nucleon poles, leading to the contributions shown diagrammatically in Figs. 2(a) and (b).




FIG. 2a




FIG. 2b

These poles are the first approximations to a fixed- t dispersion calculation of the right-hand amplitude; in this spirit their couplings are to be taken as constants, as previous authors have done.^{6,7}

A. $\rho\pi$ Intermediate State

The couplings necessary for this calculation are defined by the following phenomenological Hamiltonians.

$$H_{\pi NN} = g_{\pi NN} \bar{N} \tau_i \gamma_5 N \Pi^i \quad (18)$$

$$H_{\rho\pi\pi} = g_{\rho\pi\pi} i \epsilon_{ijk} \Pi^{*i} (q + q')^\mu \rho_\mu^j \Pi^k \quad (19)$$

$$H_{\rho NN} = g_{\rho NN} \bar{N}(p') \left(\frac{1}{2} \tau_i \right) \left[F_1^V \gamma^\mu + F_2^V i \sigma_{\mu\nu} (p' - p)^\nu / 2m \right] \rho^{i,\mu} N(p) \quad (20)$$

where, by vector dominance, F_1^V , F_2^V are respectively the Dirac and Pauli isovector couplings, with the values $F_1^V = 1$, $F_2^V = \kappa_V = 3.7$, and $g_{\rho\pi\pi} \approx g_{\rho NN} \equiv g_\rho$, with $g_\rho^2/4\pi \approx 2.8$, as determined by the width⁹ $\Gamma_{\rho\pi\pi} \approx 150$ MeV.

The contribution to the absorptive part of K is then calculated to be¹⁸, for $(m_\rho + \mu)^2 \leq t \leq 4m^2$,

$$\text{Im } K_{\rho\pi}^N(t) = \frac{g_\rho^2}{4\pi} \left[\frac{Q_\rho}{\sqrt{t}} \frac{t - m_\rho^2 - \mu^2}{m_\rho^2} + \frac{\kappa_V Q_\rho^2 - \mu^2}{P \sqrt{t}} \tan^{-1} A_\rho \right], \quad (21)$$

where

$$Q_\rho = \frac{((t - (m_\rho - \mu)^2) (t - (m_\rho + \mu)^2))^{1/2}}{2\sqrt{t}} \quad (22)$$

$$P = \frac{1}{2} (4m^2 - t)^{1/2} \quad (23)$$

and

$$A_\rho = \frac{4 P Q_\rho}{t - m_\rho^2 - \mu^2} \quad (24)$$

When inserted into the dispersion integral and evaluated numerically, this gives a contribution to Δ

$$\Delta^N(\rho\pi) = 0.015 \quad , \quad (25)$$

in agreement with the calculation of Braathen.⁷

The inclusion of the pion pole, Fig. 3,




FIG. 3

leads to a contribution to the absorptive part

$$\text{Im } K_{\rho\pi}^S(t) = - \frac{g_\rho^2}{4\pi} \cdot \frac{4 Q_\rho^3 \sqrt{t}}{m_\rho^2 (t - \mu^2)} \quad (26)$$

and to Δ

$$\Delta^S(\rho\pi) = - 0.0055 \quad (27)$$

B. $\sigma\pi$ Intermediate State

In the calculation of the contribution from this intermediate state, we will differ in some important respects from previous treatments. Let us, however, first calculate the absorptive part for a 0^+ σ -particle of arbitrary mass m_σ and couplings $g_{\sigma NN}$, $g_{\sigma\pi\pi}$ defined by the phenomenological Hamiltonians

$$H_{\sigma NN} = g_{\sigma NN} \bar{N} N \phi_\sigma \quad (28)$$

$$H_{\sigma\pi\pi} = g_{\sigma\pi\pi} \Pi^{*i} \delta_{ij} \Pi^j \phi_\sigma \quad (29)$$

From Figs. 2a and 2b we obtain an absorptive part

$$\text{Im } K_{\sigma\pi}^N(t) = \frac{g_{\sigma\pi\pi} g_{\sigma NN}}{4\pi} \cdot \frac{m}{t} (t - m_\sigma^2 + \mu^2) \cdot \frac{1}{2 P\sqrt{t}} \tan^{-1} A_\sigma \quad (30)$$

As discussed in the next Section we will use the values of the couplings $g_{\sigma\pi\pi}$, $g_{\sigma NN}$ as given in the σ model,¹⁹ viz.

$$g_{\sigma NN} = g \quad (31)$$

$$g_{\sigma\pi\pi} = \frac{g}{2m} (m_\sigma^2 - \mu^2), \quad (32)$$

and a mass $m_\sigma = 500$ MeV, somewhat lower than the mass of the physical ϵ particle.⁹ We choose this value to agree with the pion-nucleon σ term and in recognition of the fact that as far as the σ particle is concerned we are not including closed loops, which renormalize the mass upwards.²⁰

With these parameters the calculated contribution to Δ is²¹

$$\Delta^N(\sigma\pi) = 0.01 \quad (33)$$

The self-energy diagram, Fig. 3, gives an absorptive part¹⁸

$$\text{Im } K_{\sigma\pi}^S(t) = - \frac{g_{\sigma\pi\pi}^2}{4\pi} \frac{t - m_\sigma^2 + \mu^2}{4t(t - \mu^2)}, \quad (34)$$

which gives a negligible contribution to Δ when inserted in the dispersion integral, Eq. (11).

C. Resonant Contributions

So far our discussion has been limited to the inclusion of nucleon poles in the s- and u-channels of the amplitudes $\pi\pi \rightarrow N\bar{N}$ and $\pi\sigma \rightarrow N\bar{N}$. In the absence of a reliable method of estimating the continuum contribution, we can consider the inclusion of higher mass resonances in these channels. The structure of

the resonance spectrum is such that these resonances do not contribute as much as might have been expected. The reasons for this are as follows:

(i) For the $\rho\pi$ intermediate state, the $\Delta(1236)$ contribution is suppressed (apart from the mass factor) by the M1 nature of the $\rho N\Delta$ vertex near resonance.²² Higher resonant contributions will be suppressed primarily because of the mass factor, and as an optimistic estimate we may hope for at most

$$\Delta_{(\rho\pi)}^{\text{proper}} \approx 0.02$$

for the proper $\rho\pi$ contribution.

(ii) For the $\sigma\pi$ intermediate state $\Delta(1236)$ is excluded by isospin invariance. Higher resonances such as $N'(1470)$, $N'(1520)$, $N'(1535)$ etc. appear⁹ to be rather weakly coupled to the ϵN system. Hence, assuming that the combined resonance contributions add, it seems that the most we can hope for is

$$\Delta(\sigma\pi) \approx 0.015$$

D. Strange Particle Intermediate States

The next highest mass intermediate states would be the strange counterparts of (A) and (B), viz. $K^*\bar{K}$ and $\kappa\bar{K}$ respectively. The couplings of K^* and K can be related by SU(3) to those of ρ and π , giving an overall Clebsch-Gordon factor ≈ 0.4 for the Dirac coupling and 0.2 for the Pauli coupling. Combined with a large reduction factor from higher masses in the dispersion relation, the effect is to make the $K^*\bar{K}$ contribution negligible. Since the $\sigma(\epsilon)$ is predominantly a unitary singlet,²³ the couplings of the κ and the σ cannot be so directly related, but barring an unforeseen enhancement, the mass effect in the dispersion relation should have the same result of also making the $\kappa\bar{K}$ contribution negligible.

To sum up, our calculated contributions to Δ , Eqs. (25), (27) and (33), amount to 0.02. Our value for $\Delta(\rho\pi)$ agrees with that given by Braathen (Ref. 7, Table 5) and is greater than that of Pagels (Ref. 16, footnote 9) because of the inclusion of the Pauli term and the use of a larger value of g_{ρ}^2 . Our value for $\Delta(\sigma\pi)$ is positive and greater than the $\Delta(\epsilon\pi)$ of Braathen²⁴ because of a different evaluation of the $\sigma\pi$ contribution, as discussed in the next section.

In addition to the calculated contributions we have looked at possible resonance contributions and estimated that they may amount to some 0.01. Finally, there is the high energy contribution, from states with invariant masses greater than $2m$, which, given the assumptions of Pagels,^{6,17} is bounded by 0.01. It is interesting, however, that a reasonable cutoff of 2 GeV leads to a $\rho\pi$ high energy contribution⁷ to Δ of 0.028.

In spite of the increase over previous estimates, with a possible total contribution of, say, 0.035, we are still some way from the present experimental value, Eq. (2). However, the discrepancy is small enough that one may hope for an ultimate reconciliation. It may be that the high energy bound of Ref. 6, and the assumptions made therein, holds the key to the puzzle.

III. THE σ MODEL AND THE $\pi N \sigma$ TERM

The parameters of the σ meson used in Section II are intimately related to the value of the $\pi N \sigma$ term,

$$\sigma(\pi N) \equiv \langle N | [Q_5^\pi, i \partial \cdot A^\pi] | N \rangle . \quad (35)$$

This can be obtained by an extrapolation of the background (non-pole) forward isotopic-even πN amplitude

$$\bar{F}^{(+)}(\nu, t) = \bar{A}^{(+)} + \nu \bar{B}^{(+)} - g^2/m \quad (36)$$

to the on-shell sub-threshold point²⁵ $\nu = 0, t = 2\mu^2$:

$$\bar{F}^{(+)}(0, 2\mu^2) = \frac{\sigma(\pi N)}{f_\pi^2} + O(\mu^4) \quad (37)$$

or to the double off-shell point⁸ $\nu = 0, t = 0$ with $q^2 = q'^2 = 0$:

$$\bar{F}^{(+)}(0, 0; q^2 = q'^2 = 0) = - \frac{\sigma(\pi N)}{f_\pi^2} \quad (38)$$

Phenomenological analyses using both these methods seem to be converging^{2,5,26} on a value of $\sigma(\pi N) \approx 70$ MeV.

The final term in Eq. (36), which arises from the difference between the nucleon pole contributions as evaluated in dispersion theory in contrast to field theory, is crucial to what follows, and has been noted by several authors. In the presence of this term the t-channel " σ dominance" model for $\bar{F}^{(+)}$ reads²⁷

$$\bar{F}^{(+)}(\nu, t) \approx \frac{2 g_{\sigma NN} g_{\sigma \pi \pi}}{m_\sigma^2 - t} - \frac{g_\sigma^2}{m} \quad (39)$$

The σ model couplings (31) and (32) then ensure that $\overline{F}^{(+)}(\nu, t)$ vanishes at the on-shell point $\nu = 0, t = \mu^2$:

$$\overline{F}^{(+)}(0, \mu^2) \approx 0 \quad (40)$$

which is a good approximation²⁶ to the required vanishing²⁸ of the amplitude at the off-shell Adler consistency point $\nu = 0, t = \mu^2, q^2 = 0, q'^2 = \mu^2$. Thus we may have some confidence in the couplings (31) and (32), or at least in their product, which is all that enters into the calculation of $\Delta(\sigma\pi)$.

Parenthetically, we may remark that a similar situation obtains with the isotopic-odd forward scattering amplitude and the ρ couplings. The t-channel " ρ dominance" model²⁹ for $\overline{F}^{(-)} = \overline{A}^{(-)} + \nu \overline{B}^{(-)}$ implies

$$\nu^{-1} \overline{F}^{(-)}(\nu, t) \approx \frac{g_{\rho\pi\pi} g_{\rho NN}}{m_{\rho}^2 - t} - \frac{g_{\rho}^2}{2m^2} \quad (41)$$

The value of g_{ρ} we have used in Section II, which is in fact that given by the KSFR relation³⁰

$$\frac{g_{\rho}^2}{m_{\rho}^2} = \frac{1}{2 f_{\pi}^2} \quad (42)$$

is such as to cast (41) into the form of the Adler-Weisberger relation³¹ with the G-T value for g_A . Once the $0(\mu^2)$ background is taken into account, it can be shown that (41) is in good agreement with the low energy data for various values of ν and t near threshold.²⁶

Returning to the question of the σ couplings, we note that the difference between our evaluation of the $\sigma\pi$ contribution and that of Pagels⁶ derives from the extra term in Eq. (39), in the absence of which the values of the πN scattering lengths constrained $g_{\sigma NN} g_{\sigma\pi\pi}$ to be small. Since $g_{\sigma\pi\pi}$ was regarded as

fixed by the width $\Gamma_{\sigma \rightarrow \pi\pi}$, the net result was to suppress $g_{\sigma NN}$ and hence the triangle graph contributions, leaving only a small (negative) self-energy contribution.

Our attitude is to keep the σ model relations (31) and (32) for the couplings and then allow the value of the σ term to determine the effective mass to be used in the calculation. The connection is provided by confronting (37) with (39), which gives³²

$$\sigma(\pi N) \approx g_A^2 (\mu/m_\sigma)^2 m \quad (43)$$

where, to this order of accuracy we have used the G-T relation itself and have neglected the pion mass relative to the σ mass.

There is some ambiguity at this stage. Since we are using the lowest-order unrenormalized³³ couplings for σ , we should perhaps for consistency take $g_A = 1$ in Eq. (43). An input σ term of the order of 70 MeV would then lead to $m_\sigma^2 \approx 13\mu^2$, or $m_\sigma \approx 500$ MeV. Given this value of m_σ , the σ model relations then give

$$g_{\sigma\pi\pi} \approx 12\mu, \quad (44)$$

while $g_{\sigma NN}$ as always is constrained to have the value g .

In principle (44) can be compared directly with experiment in the form of the σ width.^{34,35} Unfortunately the experimental situation is rather confused, we have the above-mentioned renormalization ambiguity, and the narrow width approximation is not obviously a very appropriate one.

We wish, however, to make the following observation. According to the narrow-width approximation (NWA) the σ width is given by³⁴

$$\Gamma_{\sigma \rightarrow \pi\pi} = \frac{3}{2} \frac{g_{\sigma\pi\pi}^2}{4\pi m_{\sigma}^2} (m_{\sigma}^2 - 4\mu^2)^{1/2} \quad (45)$$

which, for $m_{\sigma}^2 \approx 13\mu^2$, has the value

$$\Gamma_{\sigma \rightarrow \pi\pi} \approx 4\mu \approx 550 \text{ MeV} \quad (46)$$

within the range of experiment.³⁶ Although the use of the NWA is not a priori very reasonable for a width of this magnitude, the general feature of this approximation is that a coupling larger than the physical one is needed to obtain agreement with a given physical width. (For example, the $\pi N \Delta$ coupling derived from the Δ width in NWA is 40% greater than the physical coupling.³⁶) Thus the neglect in (43) of the factor g_A^2 , the G-T approximation, and the neglect of μ^2 compared to m_{σ}^2 , which would have led to a (factor of 2) larger physical coupling constant, to some extent takes account of this effect. We therefore take the attitude of using the coupling constant (44) and at the same time ignoring finite width effects.

With $\sigma(\pi N) \approx 70 \text{ MeV}$, leading, as we have seen, to $\Gamma_{\sigma} \approx m_{\sigma} \approx 500 \text{ MeV}$, the σ model may be taken as a reasonable approximation to reality. By way of contrast we note that if $\sigma(\pi N)$ were very small, say $\sigma(\pi N) \approx 15 \text{ MeV}$, then the σ model would lead to $m_{\sigma} \approx 1100 \text{ MeV}$, $g_{\sigma\pi\pi} \approx 30\mu$ and $\Gamma_{\sigma} \approx 2000 \text{ MeV}$, and would not then be physically relevant.

The difference of our σ contribution from Braathen's ϵ contribution lies in the fact that we have taken $g_{\sigma NN}$ to be fixed by the σ model and determined m_{σ} from the pion-nucleon σ term, leading to a not unreasonable value for Γ_{σ} .

Braathen, on the other hand, took m_{ϵ} and Γ_{ϵ} as input, leading to a much-reduced value for $g_{\epsilon NN}$. Comparison of the resulting Δ 's shows the sensitivity of the calculation to such factors.

IV. SU(3) × SU(3) CHIRAL SYMMETRY BREAKING

We now turn to the use of the G-T relation to probe the nature of the symmetry breaking Hamiltonian H' . In particular, if this is assumed to have the $(3, \bar{3}) + (\bar{3}, 3)$ form $H' = u_0 + cu_8$, the G-T relation provides an independent determination of the value of c .

Equation (17) on its own will not suffice, since we have no way of evaluating $(\bar{v}_\pi)_{NN}$. The method of procedure will therefore be to eliminate this quantity by means of the SU(3) generalizations of Eq. (17). Defining the $K^+ p \Lambda$ and $K^+ p \Sigma^0$ discrepancies as

$$\Delta_{K\Lambda} \equiv 1 - \frac{\frac{1}{2} (m_\Lambda + m_N) g_A^{K\Lambda}}{f_K g_{K\Lambda}} \quad (47)$$

and

$$\Delta_{K\Sigma} = 1 - \frac{\frac{1}{2} (m_\Sigma + m_N) g_A^{K\Sigma}}{f_K g_{K\Sigma}}, \quad (48)$$

we have³⁷, as analogues of (17),

$$\Delta_{K\Lambda} = \frac{1}{f_K g_{K\Lambda}} \frac{\sqrt{2} - 1/2 c}{2\sqrt{3}} (\bar{v}_K)_{\Lambda} \quad (49)$$

$$\Delta_{K\Sigma} = \frac{1}{f_K g_{K\Sigma}} \frac{\sqrt{2} - 1/2 c}{2\sqrt{3}} (\bar{v}_K)_{\Sigma} \quad (50)$$

The matrix elements of \bar{v} appearing in (49) and (50) are related in SU(3) according to

$$(\bar{v}_P)_{fi} = 2(\bar{v}_\pi)_{NN} (d_v d_{Pfi} - f_v f_{Pfi}) \quad (51)$$

in which we have an unknown f_V ($d_V = 1 - f_V$). This, however, may be eliminated between the two equations, leading to the relation^{8,14,38}

$$\frac{\sqrt{2} + c}{\sqrt{2} - \frac{1}{2}c} = \frac{2f_\pi}{f_K} \frac{g_{\pi NN} \Delta_{\pi NN}}{-\sqrt{3} g_{KNA} \Delta_{KNA} + g_{KN\Sigma} \Delta_{KN\Sigma}} \quad (52)$$

If one further assumes exact kaon PCAC in the operator sense, the left-hand side of (52) becomes $f_\pi m_\pi^2 / f_K m_K^2$, in agreement with GMOR. In this case (52) is then equivalent to the sum rules derived by Dashen and Weinstein.^{14,8} We prefer, however, to follow the philosophy of Furlan and Paver¹⁴ and regard (52) as the baryon analogue of the GMOR meson mass formula from which we can independently derive the value of c .

The procedure we will adopt in analyzing (52) is first of all to assume SU(3) invariance for the baryon couplings to mesons and axial currents,³⁹ viz.

$$g_{Pfi} = 2g \left(d_P d_{Pfi} - f_P f_{Pfi} \right) \quad (53)$$

and

$$g_A^{Pfi} = 2g_A \left(d_A d_{Pfi} - f_A f_{Pfi} \right), \quad (54)$$

with $d_P + f_P = d_A + f_A = 1$, and then afterwards to examine the effects of SU(3) breaking.

In the case of exact SU(3), Eq. (52) can be recast in the form

$$-\frac{\sqrt{2} + c}{c} = \frac{3}{2} \Delta_{\pi NN} \left[\left(\frac{f_K}{f_\pi} - 1 \right) - \frac{mg_A}{f_\pi g} \left(\frac{m_\Sigma + m_\Lambda - 2m_N}{4m_N} - f_A \frac{m_\Sigma - m_\Lambda}{2m_N} \right) \right]^{-1} \quad (55)$$

Remarkably we see that f_P does not appear here, i. e., that the relation is independent⁴⁰ of the d/f ratio for the strong coupling constants once these are assumed to obey SU(3). Moreover the dependence on the parameter f_A is

greatly suppressed by the factor $(m_\Sigma - m_\Lambda)$. The remaining terms in the right-hand denominator are all SU(3) breaking effects, and are of such a magnitude measured against $\Delta_{\pi NN}$ that the value of c derived from (55) is considerably different from the canonical GMOR value of -1.25 .

A single angle fit to hyperon semileptonic decays gives³⁹

$$d_A/f_A = 1.7 ,$$

corresponding to $f_A = 0.37$, while, with the aid of the nonrenormalization theorem⁴² for K_{l3} decay, f_K/f_π can be determined from the value⁴³ of $f_K/(f_\pi f_\pi(0))$ to be $f_K/f_\pi = 1.24 \pm 0.02$. For $\Delta_{\pi NN}$ we insert the experimental value given in Eq. (2) and obtain

$$c_{GT} = -0.9 \pm 0.1 \quad (56)$$

in remarkable agreement with the value obtained by an examination of σ terms.² The degree of deviation from the GMOR prediction is more fairly measured by the quantity $(\sqrt{2+c})/(\sqrt{2-\frac{1}{2}c})$, which, from (56), is

$$\left(\frac{\sqrt{2+c}}{\sqrt{2-\frac{1}{2}c}} \right)_{GT} = 0.3 \pm .06 \quad (57)$$

compared with the expected value¹ $m_\pi^2/m_K^2 \approx 0.08$, or⁸ $f_\pi m_\pi^2/f_K m_K^2 \approx 0.06$.

We now go back to examine our assumption of SU(3) invariance for the various couplings g_{Pfi} , g_A^{Pfi} .

In the general case (52) can be written as

$$\frac{\sqrt{2+c}}{\sqrt{2-\frac{1}{2}c}} = \frac{\Delta_{\pi NN}}{\frac{f_K}{f_\pi} A - \frac{mg_A}{f_\pi g} \left[\left(\frac{m_\Sigma + m_\Lambda + 2m_N}{4m_N} \right) B - \left(\frac{m_\Sigma - m_\Lambda}{2m_N} \right) C \right]} \quad (58)$$

where

$$A = (\pm\sqrt{3}g_{KNA} + g_{KN\Sigma})/2g \quad (59)$$

$$B = \left(-\sqrt{3} g_A^{\text{KNA}} + g_A^{\text{KN}\Sigma} \right) / 2g_A \quad (60)$$

and

$$C = -\left(\sqrt{3} g_A^{\text{KNA}} + g_A^{\text{KN}\Sigma} \right) / 4g_A, \quad (61)$$

which, in the SU(3) limit we have just been discussing, take on the values

$$A = B = 1 \text{ and } C = f_A.$$

The experimental situation for the strong coupling constants appearing in (59) is as follows.¹⁰ All estimates of $g_{\text{KN}\Sigma}^2/4\pi$ have been that this quantity is small, of the order of 2. Modern methods of extracting the coupling constants from KN scattering evaluate the combination $g_Y^2 \equiv (g_{\text{KNA}}^2 + 0.84 g_{\text{KN}\Sigma}^2)/4\pi$. In the 1973 compilation¹⁰ the most reliable value was considered to be⁴⁴ $g_Y^2 = 12.2 \pm 2.5$; however, the value $g_Y^2 = 15.2 \pm 2.3$ obtained by the Cutkosky conformal mapping technique⁴⁵ was not mentioned. Furthermore, a recent experiment⁴⁶ indicates that $g_Y^2 = 19.8 \pm 4.4$ within the zero range approximation, thus negating the original zero range estimates⁴⁷ of $g_Y^2 \approx 4$. Hence, at the present time, we believe that fair estimates of the coupling constants are

$$\begin{aligned} g_{\text{KNA}}^2/4\pi &= 14 \pm 2 \\ g_{\text{KN}\Sigma}^2/4\pi &= 2 \pm 1 \end{aligned} \quad (62)$$

in reasonable agreement with the original evaluation of Kim.⁴⁸ Moreover these values would predict $g_{\pi\Lambda\Sigma}^2/4\pi \approx 9$ by SU(3), compared to the values 10.9 ± 0.3 , 12.9 ± 0.8 obtained phenomenologically^{49,50} from $\Lambda\pi$ partial-wave dispersion relations. Thus we will certainly believe SU(3) to the extent that it determines the signs of g_{KNA} and $g_{\text{KN}\Sigma}$ and hence deduce from (62) that

$$\begin{aligned} g_{\text{KNA}} &= -13 \pm 1 \\ g_{\text{KN}\Sigma} &= 5 \pm 1 \end{aligned} \quad (63)$$

Substituting these values into (59) we find

$$A = 1 \pm 0.1 \quad (64)$$

The situation with regard to the SU_3 -Cabibbo theory of hyperon semi-leptonic decays has also improved recently.⁴¹ The $e-\nu$ correlation measurements in $\Lambda \rightarrow pe\bar{\nu}$ lead to the axial vector-to-vector ratio of $g_1/f_1 \sim 0.7$ with 10% error, which is significantly larger than previously reported. Since the Cabibbo estimate of g_1/f_1 is $(d_A + \frac{1}{3} f_A) g_A \approx 0.72$, using $d_A/f_A \approx 1.7$ which is obtained from the remaining hyperon decays,³⁹ we can evaluate (60) as

$$B = 1 \pm 0.1 \quad (65)$$

As previously noted in the symmetry limit, the contribution of C in (58) is greatly suppressed by the factor $(m_\Sigma - m_\Lambda)$, so that we may take $C = f_A$ without serious error.

With these values (58) gives

$$\frac{\sqrt{2+c}}{\sqrt{2-\frac{1}{2}c}} = \frac{0.058 \pm 0.013}{0.21 \pm 0.16} \quad (66)$$

While the error in the denominator of (66) is large, it is clear that the GMOR value of $0.06 - 0.08$ for $(\sqrt{2+c})/(\sqrt{2-\frac{1}{2}c})$ is still remote. We therefore arrive at the conclusion, bearing out the misgivings expressed in Ref. 8, that, given a symmetry-breaking Hamiltonian of the form $u_0 + cu_8$, the value of c derived from the Goldberger-Treiman relation cannot be -1.25 , but instead is close to -1 , in agreement with the value obtained from phenomenological analyses of σ terms.

V. DISCUSSION AND CONCLUSIONS

We have discussed two aspects of the Goldberger-Treiman relation: the attempt to account for the experimental πN discrepancy $\Delta_{\pi NN}$ by means of dispersion relations, and the use of this quantity and its SU(3) counterparts to test the nature of the chiral-symmetry-breaking Hamiltonian.

The narrowing of the gap between the experimental and theoretical values of $\Delta_{\pi NN}$ arises from two factors: first, a not insignificant contribution from the $\sigma\pi$ intermediate states, ultimately connected to the large value of the $\pi N \sigma$ term, and second, a shift in the experimental parameters all in the direction of reducing the experimental value of $\Delta_{\pi NN}$. Thus g_A has increased in time to the currently accepted value⁹ 1.25 ± 0.01 . The πN coupling constant g has tended to decrease with time, and indeed the most recent and most accurate determination⁵¹ $g = 13.40 \pm 0.08$, which we used in arriving at Eq. (2) is appreciably less than the more generally quoted¹⁰ value $g \approx 13.6$. The nucleon mass⁵² has not, of course, varied to any appreciable extent, but the value to be used in Eq. (2) is actually the average of the neutron and proton masses, namely 938.9 MeV. From the π^+ lifetime⁹ $\tau = (2.6024 \pm 0.024) \times 10^{-8}$ sec and using the value¹² $G = (1.026 \pm 0.001) \times 10^{-5} m_p^{-2}$, one can derive

$$f_{\pi} \cos \theta_A = (90.778 \pm 0.042) \text{ MeV} \quad (67)$$

Given $f_K/f_{\pi} = 1.24 \pm 0.02$, the ratio of $K_{\mu 2}$ to $\pi_{\mu 2}$ rates implies

$$\cos \theta_A = 0.976 \pm 0.001$$

which leads to the value quoted in Section I along with its error:

$$f_{\pi} = (93.0 \pm 1) \text{ MeV} \quad (68)$$

including the possible 1% radiative corrections to $\pi^+ \rightarrow \mu^+ \nu$.

These are the values we have used in Eq. 2 to obtain $\Delta_{\pi NN}^{\text{exp}}$ together with its error, which, as we have seen, is large enough to make an ultimate agreement between theory and experiment by no means improbable. The change from a few years ago⁶ arises from an increased theoretical estimate and a trend in the experimental numbers all in the direction of decreasing $\Delta_{\pi NN}^{\text{exp}}$.

As far as the second aspect is concerned, we have shown in the present paper and in Ref. 2 that two of the measures of $SU(2) \times SU(2)$ breaking noted by Li and Pagels,⁵³ namely

$$\Delta_{\pi NN} \equiv 1 - \frac{m g_A}{f_\pi g} \approx 0.058 \quad (69)$$

and

$$\frac{\sigma(\pi N)}{m} \approx \frac{(70 \text{ MeV})}{(939 \text{ MeV})} \approx 0.075, \quad (70)$$

can be explained by the same chiral-symmetry-breaking mechanism $H' = u_0 + c u_8$ with $c \approx -1$.

The third measure,

$$\frac{m_\pi^2}{m_K^2} \approx 0.077, \quad (71)$$

seems difficult to reconcile with the first two. In the standard GMOR version of $SU(3) \times SU(3)$, c is related to the masses by soft pion, kaon and eta PCAC according to

$$c_{\text{GMOR}} \approx -\sqrt{2} \left(\frac{m_K^2 - m_\pi^2}{m_K^2 + \frac{1}{2}m_\pi^2} \right) \approx -1.25 \quad (72)$$

The validity of the extrapolations involved in kaon and eta PCAC can certainly be questioned, but the same value can be derived either by SU(3) symmetry⁵⁴ for the quantities $\langle 0 | v_i | P_i \rangle$ or⁵⁵ by SU(3) symmetry of the m_P^2 and only pion PCAC. There thus appears to be a rather fundamental clash between the first two derivations of c and the mass formula, which we have attempted² to resolve in terms of some degree of SU(3) breaking of $\langle 0 | v_i | P_i \rangle$ and of the vacuum.⁵⁶

If this resolution is not found convincing (and certainly, as we have shown, the value $c = -1.25$ is in violent disagreement with the SU(3) G-T relation (52), whose derivation is on very firm ground according to the standard approach), the conclusion to be drawn is that the irreducible $(3, \bar{3}) + (\bar{3}, 3)$ representation is not compatible with the data. The most plausible alternative, a reducible representation with a small admixture⁵⁷ of $(8, 8)$, has yet to be confronted with all the data in a systematic way.

ACKNOWLEDGMENTS

The first author would like to thank Prof. S. D. Drell for his hospitality at SLAC and Prof. E. Golowich for helpful discussions. The second author gratefully acknowledges conversations with Dr. L. R. Thebaud, Prof. G. Furlan, Dr. P. H. Weisz and Dr. H. Pagels.

REFERENCES

1. M. Gell-Mann, R. J. Oakes and B. Renner (GMOR), Phys. Rev. 175, 2195 (1968); S. Glashow and S. Weinberg, Phys. Rev. Letters 20, 224 (1968).
2. M. D. Scadron and H. F. Jones, Phys. Rev., to be published.
3. M. L. Goldberger and S. B. Treiman, Phys. Rev. 111, 354 (1958).
4. This value will be discussed in Section V.
5. E. Reya, Mainz preprint MZ-TH-73/1 (Table I); H. Nielsen and G. C. Oades, Nucl. Phys. B72, 310 (1974).
6. H. Pagels, Phys. Rev. 179, 1337 (1969).
7. H. J. Braathen, Nucl. Phys. B44, 93 (1972).
8. R. Dashen, Proceedings of LIV th. International School of Physics "Enrico Fermi" 1971, ed. R. Gatto (Academic Press, 1972).
9. Particle Data Group, Review of Particle Properties (Phys. Letters 50B, April 1974).
10. H. Pilkuhn et al., Compilation of Coupling Constants and Low Energy Parameters (Nucl. Phys. B65, 460 (1973)).
11. The choice of g will be discussed in Section V.
12. See, e.g., V. de Alfaro, S. Fubini, G. Furlan and C. Rossetti, Currents in Hadron Physics (North Holland Publishing Co., 1973).
13. Our notation differs from Ref. 12 only in that our γ_5 is a factor of $(-i)$ times the γ_5 defined therein.
14. See also R. Dashen and M. Weinstein, Phys. Rev. 188, 2330 (1969); G. Furlan and N. Paver, Invited talk given at "Triangle Seminar," Vienna, December 1973.
15. R. A. Coleman and J. W. Moffat, Phys. Rev. 186, 1635 (1969).

16. H. Pagels and A. Zepeda, Phys. Rev. D5, 3262 (1972).
17. The bound was originally given in Ref. 6 as 0.004, but this was later corrected by Braathen (Ref. 7).
18. This agrees with the absorptive part given in Ref. 7, modulo misprints therein.
19. M. Gell-Mann and M. Lévy, Nuovo Cimento 16, 705 (1960).
20. J. L. Basdevant and B.W. Lee, Phys. Rev. D2, 1680 (1970).
21. This was much to our surprise, as our absorptive part is twice that given in Ref. 7, yet the contribution to Δ turns out to be half that quoted therein.
22. However, the Δ contribution can eventually be shown to be positive due to the sign of the Δ coupling constants as determined by M. D. Scadron and L. R. Thebaud, Phys. Rev. D8, 2190 (1973); see Eq. (40).
23. See, e.g., D. Morgan, Rutherford Laboratory preprint RL-74-063 (April, 1974).
24. We believe the value 0.023 quoted by Braathen for $\Delta(\sigma\pi)$ to be incorrect for the reasons stated in Ref. 21.
25. T. P. Cheng and R. Dashen, Phys. Rev. Letters 26, 594 (1971); Y. C. Liu and J. A. M. Vermaseren, Phys. Rev. D8, 1602 (1973).
26. M. D. Scadron and L. R. Thebaud, Phys. Letters 46B, 257 (1973); Phys. Rev. D9, 1544 (1974).
27. S. Gasiorowicz and D. A. Geffen, Rev. Mod. Phys. 41, 531 (1969); H. Schnitzer, Phys. Rev. D5, 1490 (1972), App. A.
28. S. L. Adler, Phys. Rev. 137, B 1022 (1965).
29. J. J. Sakurai, Phys. Rev. Letters 17, 552 (1966).

30. K. Kawarabayashi and M. Suzuki, Phys. Rev. Letters 16, 255 (1966);
Riazuddin and Fayazuddin, Phys. Rev. 147, 1071 (1966).
31. S. L. Adler, Phys. Rev. Letters 14, 1051 (1965); W. I. Weisberger, *ibid.*
p. 1047.
32. See, e.g., L. Turner and M. G. Olsson, Phys. Rev. D6, 3522 (1972).
33. For a discussion of g_A and the G-T relation in the renormalized σ model
see H. J. Strubbe, Nucl. Phys. B28, 299 (1972).
33. J. L. Petersen and J. Pišút, Nucl. Phys. B38, 207 (1972). These authors
find that the ratio $g_{\sigma NN}/g_{\sigma\pi\pi}$ as given by the σ model is satisfied with
some sets of $\pi\pi$ phase shifts, but not with others.
35. J. C. Le Guillou, A. Morel and H. Navelet, Nuovo Cimento 5A, 659 (1971).
36. G. Höhler, H. P. Jakob and R. Strauss, Nucl. Phys. B39, 237 (1972).
37. The derivation of these equations involves, of course, kaon PCAC. In
general we believe that this should be used with caution, but is probably
valid here, since the kaon extrapolation is small on the mass scale set
by the external baryon states. In the GMOR/Dashen philosophy, kaon
PCAC is on an essentially equal footing with pion PCAC.
38. The Clebsch-Gordon factors in Eq. (52) differ somewhat from those in
the corresponding expressions in Refs. 8, 14 due to a different choice of
charge modes, which we have explicitly stated above Eq. (47).
39. L. -M. Chounet, J. -M. Gaillard and M. K. Gaillard, Phys. Rep. 4, 199
(1972).
40. Of the four possible expressions for $(\sqrt{2} + c)/(\sqrt{2} - \frac{1}{2}c)$ involving $\Delta_{\pi NN}$,
 $\Delta_{\pi\Sigma\Lambda}$, $\Delta_{KN\Lambda}$ and $\Delta_{KN\Sigma}$, Eq. (52) is unique in this respect, and is there-
fore the least sensitive to the effects of SU(3) breaking of the strong
coupling constants.

41. CERN-Heidelberg and Argonne-Chicago groups, Proceedings of the XVII International Conference on High Energy Physics, Chicago-Batavia, Ill., 1972.
42. M. Ademollo and R. Gatto, Phys. Rev. Letters 13, 264 (1965);
P. Langacker and H. Pagels, *ibid.* 30, 630 (1973).
43. See, e. g. , Ref. 12 or Ref. 39.
44. E. Pietarinen and C. P. Knudsen, Nucl. Phys. B57, 637 (1973).
45. R. C. Miller et al. , Nucl. Phys. B37, 401 (1972).
46. P. Baillon et al. , CERN preprint CERN/D. Ph. II/PHYS 74-6 (January 1974).
47. See, e. g. , B. R. Martin and M. Sakitt, Phys. Rev. 183, 1352 (1969).
Their errors are, however, large and therefore not inconsistent with $g_Y^2 \sim 15$.
48. J. K. Kim, Phys. Rev. Letters 19, 1079 (1967).
49. B. Martin, Phys. Rev. 138B, 1136 (1965).
50. J. Engels, Nucl. Phys. B51, 269 (1973).
51. D. V. Bugg, A. A. Carter and J. R. Carter, Phys. Letters 44B, 278 (1973).
52. From the viewpoint of a once-subtracted dispersion relation for $K(q^2)$, Eq. (12), the nucleon mass entering the G-T discrepancy is the physical and not the chiral-limiting nucleon mass. It should be pointed out, however, that the $(3, \bar{3})$ values $c \approx -1.0$ and $(u_0/u_8)_N \approx 1$ (see Refs. 2, 26) imply that $m_N - m_0 = (u_0 + cu_8)_N \approx 0$, so that the physical nucleon and chiral-limiting nucleon masses are essentially identical.

53. L. F. Li and H. Pagels, Phys. Rev. Letters 27, 1089 (1971); Phys. Rev. D5, 1509 (1972).
54. P. Langacker and H. Pagels, Phys. Rev. D8, 4595 (1973).
55. J. Ellis, Nucl. Phys. B13, 153 (1969).
56. To counteract the derivation of Ref. 55, there would have to be some SU(3) breaking of the m_p^2 .
57. A. Sirlin and M. Weinstein, Phys. Rev. D6, 3588 (1972);
A. McDonald, S. P. Rosen and T.K. Kuo, Phys. Letters 40B, 675 (1972);
A. M. Harun ar Rashid, Nuovo Cimento 17A, 288 (1972);
D. T. Cornwell, H. Genz, H. Steiner and S. Eliezer, Karlsruhe pre-print TKP 2/74 (February 1974).

FIGURE CAPTIONS

1. $\rho\pi$ and $\sigma\pi$ intermediate state contributions to Δ .
- 2a and 2b. s- and u-channel nucleon pole diagrams.
3. t-channel pion pole diagram.




FIG. 1




FIG. 2a




FIG. 2b




FIG. 3