

## GORENSTEIN TORIC SINGULARITIES AND CONVEX POLYTOPES

ATSUSHI NOMA

(Received July 9, 1990, revised January 4, 1991)

**Introduction.** Let  $X$  be a normal projective variety over a field  $k$  and  $D$  an *ample  $\mathbf{Q}$ -divisor*, i.e., a rational coefficient Weil divisor such that  $bD$  is an ample Cartier divisor for some positive integer  $b$ . We consider a normal graded ring  $R(X, D)$  defined by  $R(X, D) = \bigoplus_{n \geq 0} H^0(X, \mathcal{O}_X(nD))t^n$ . Here  $t$  is an indeterminate and  $\mathcal{O}_X(nD)$  are the sheaves defined by  $\Gamma(U, \mathcal{O}_X(nD)) = \{f \in K(X); \operatorname{div}_U(f) + nD|_U \geq 0\}$  for each open set  $U$  of  $X$ . We are interested in finding a criterion for a normal projective variety  $X$  to have an ample  $\mathbf{Q}$ -divisor  $D$  with  $R(X, D)$  Gorenstein. Concerning this problem, see also [1, Chapter 5], [10]. Here we discuss this problem when  $X$  is a projective torus embedding defined over  $k$ .

Our main results are the following:

**COROLLARY 2.5.** *Let  $X$  be a projective torus embedding and  $D$  an ample Cartier divisor. Then  $R(X, D)$  is Gorenstein if and only if the canonical sheaf  $\omega_X$  on  $X$  is isomorphic to an invertible sheaf  $\mathcal{O}_X(-\delta D)$  for a positive integer  $\delta$ .*

**THEOREM 2.6.** *Every projective torus embedding  $X$  has an ample  $\mathbf{Q}$ -divisor  $D$  stable under the torus action such that  $R(X, D)$  is Gorenstein.*

To obtain these results, we proceed as follows: First, given a *rational convex  $r$ -polytope*  $P$  in  $\mathbf{R}^r$  (i.e., an  $r$ -dimensional convex polytope whose vertices have rational coordinates in  $\mathbf{R}^r$ ), we construct a pair of projective torus embedding  $X(P)$  over  $k$  and an ample  $\mathbf{Q}$ -divisor  $D(P)$  (Proposition 1.3) following [7, Chapter 2], so that  $R(X(P), D(P))$  is isomorphic to the normal semigroup  $k$ -algebra

$$R(P) = \bigoplus_{n \geq 0} \left\{ \sum_{m \in nP \cap \mathbf{Z}^r} ke(m) \right\} t^n.$$

Here  $t$  is an indeterminate and  $e$  is the isomorphism from  $\mathbf{Z}^r$  ( $\subset \mathbf{R}^r$ ) into the Laurent polynomial ring  $k[X_1^{\pm 1}, \dots, X_r^{\pm 1}]$  sending  $(m_1, \dots, m_r)$  to  $X_1^{m_1} \cdots X_r^{m_r}$ . Thus  $X(P)$  is isomorphic to  $\operatorname{Proj}(R(P))$  (Proposition 1.5). Conversely, it turns out that every pair of projective torus embedding  $X$  and a  $T$ -stable ample  $\mathbf{Q}$ -divisor  $D$  on  $X$  is obtained from a rational convex  $r$ -polytope in  $\mathbf{R}^r$  in this way (Proposition 1.3). On the other hand, since  $R(X(P), D(P))$  ( $\simeq R(P)$ ) is Cohen-Macaulay (cf. [4]), we can apply the criterion [10, Corollary 2.9] for the Gorenstein property to  $R(X(P), D(P))$ . Therefore we have Proposition 2.2, which is a criterion for  $R(X(P), D(P)) \simeq R(P)$  to be Gorenstein in terms

of  $D(P)$  on  $X(P)$  as well as the maximal faces of  $P$ . This yields another proof for a theorem of Hibi [3]. As immediate consequences of Proposition 2.2, we get our main results.

I should like to thank Professor Takayuki Hibi for his lecture at Tsuda College in October, 1989, from which this material stemmed. Also, I should like to thank Professor Kei-ichi Watanabe for valuable suggestions and kind advice.

**Preliminaries.**

(0.1)  $[a]$  denotes the greatest integer not greater than  $a \in \mathbf{R}$ .  $\lceil a \rceil$  denotes  $-[-a]$  for  $a \in \mathbf{R}$ .

(0.2) For notion of torus embeddings, we refer the reader to [7]. All torus embeddings will be defined over a fixed field  $k$ . Let  $T$  be an  $r$ -dimensional algebraic torus  $\text{Spec}(k[X_1^{\pm 1}, \dots, X_r^{\pm 1}])$  over  $k$ . Let  $M, N$  be the group of characters and one-parameter subgroups, respectively. By  $e(m)$ , we denote the Laurent monomial corresponding to a character  $m$ . Namely  $e(m) = X_1^{m_1} \cdots X_r^{m_r}$  for  $m = (m_1, \dots, m_r)$ . Set  $M_{\mathbf{R}} = M \otimes_{\mathbf{Z}} \mathbf{R}$  and  $N_{\mathbf{R}} = N \otimes_{\mathbf{Z}} \mathbf{R}$ . Let  $\langle, \rangle : M_{\mathbf{R}} \times N_{\mathbf{R}} \rightarrow \mathbf{R}$  represent the natural non-degenerate pairing. For a complete fan  $\Delta$  of  $N$ ,  $\Delta(i)$  denotes the  $i$ -dimensional cones of  $\Delta$ . A one-dimensional cone  $\rho \in \Delta(1)$  is generated by a unique primitive integral vector  $n(\rho)$  ([7, p. 24]). We denote by  $\text{SF}(N, \Delta)$  the additive group consisting of  $\Delta$ -linear support functions ([7, p. 66]). Set  $\text{SF}(N, \Delta, \mathcal{Q}) = \text{SF}(N, \Delta) \otimes_{\mathbf{Z}} \mathcal{Q}$ . Its elements are also called  $\Delta$ -linear support functions. Then we have two injections  $M \rightarrow \text{SF}(N, \Delta)$  sending  $m$  to  $\langle m, \rangle$ , and  $\text{SF}(N, \Delta) \rightarrow \mathbf{Z}^{\Delta(1)}$  sending  $h$  to  $(h(n(\rho)))_{\rho \in \Delta(1)}$ . Let  $X$  be a complete torus embedding  $T_N \text{emb}(\Delta)$ . By  $T\text{Div}(X)$ ,  $T\text{CDiv}(X)$  and  $\text{PDiv}(X)$ , we denote the groups of  $T$ -stable Weil divisors,  $T$ -stable Cartier divisors and principal divisors on  $X$ , respectively. Also, by  $T\text{Div}(X, \mathcal{Q})$  (resp.  $T\text{CDiv}(X, \mathcal{Q})$ ), we denote the group of  $T$ -stable  $\mathcal{Q}$ -divisors (resp.  $T$ -stable  $\mathcal{Q}$ -Cartier divisors). Namely  $T\text{Div}(X, \mathcal{Q}) = T\text{Div}(X) \otimes_{\mathbf{Z}} \mathcal{Q}$  and  $T\text{CDiv}(X, \mathcal{Q}) = T\text{CDiv}(X) \otimes_{\mathbf{Z}} \mathcal{Q}$ . The one-dimensional cones  $\rho$  of  $\Delta(1)$  are in one-to-one correspondence with the irreducible  $T$ -stable closed subvarieties  $V(\rho)$  of codimension one in  $X$ . Therefore the map  $\mathbf{Z}^{\Delta(1)} \rightarrow T\text{Div}(X)$  sending  $g$  to  $D_g = -\sum_{\rho \in \Delta(1)} g_{\rho} \cdot V(\rho)$  is a bijection, and induces two isomorphisms of groups,  $\text{SF}(N, \Delta) \rightarrow T\text{CDiv}(X)$  and  $M \rightarrow \text{PDiv}(X) \cap T\text{CDiv}(X)$ . As a result, we have two commutative diagrams (cf. [7, §2.1]):

$$\begin{array}{ccccc}
 M & \longrightarrow & \text{SF}(N, \Delta) & \longrightarrow & \mathbf{Z}^{\Delta(1)} \\
 \downarrow & & \downarrow & & \downarrow \\
 \text{PDiv}(X) \cap T\text{CDiv}(X) & \longrightarrow & T\text{CDiv}(X) & \longrightarrow & T\text{Div}(X)
 \end{array}$$
  

$$\begin{array}{ccc}
 \text{SF}(N, \Delta, \mathcal{Q}) & \longrightarrow & \mathcal{Q}^{\Delta(1)} \\
 \downarrow & & \downarrow \\
 T\text{CDiv}(X, \mathcal{Q}) & \longrightarrow & T\text{Div}(X, \mathcal{Q}) .
 \end{array}$$

**1. Rational polytopes and projective torus embeddings.**

LEMMA 1.1. *Let  $X = T_N \text{emb}(\Delta)$  be an  $r$ -dimensional complete torus embedding. For  $g \in \mathcal{Q}^{\Delta(1)}$ , the set  $\square_g = \{m \in M_{\mathbf{R}}; \langle m, n(\rho) \rangle \geq g_\rho \text{ for all } \rho \in \Delta(1)\}$  is a (possibly empty) convex polytope in  $M_{\mathbf{R}}$ . The set  $H^0(X, \mathcal{O}_X(D_g))$  of global sections of the divisorial  $\mathcal{O}_X$ -module  $\mathcal{O}_X(D_g)$  is the finite dimensional  $k$ -vector space with  $\{e(m); m \in M \cap \square_g\}$  as a basis. Let  $\text{div}(e(m)) + D_g = \sum_{\rho \in \Delta(1)} a_\rho \cdot V(\rho)$  for an element  $m \in M$ . Then  $m$  is in  $\text{int}(\square_g)$  if and only if the coefficient  $a_\rho$  for each  $\rho \in \Delta(1)$  is a positive rational number. Here  $\text{int}(\square_g)$  denotes the interior of the convex polytope  $\square_g$ .*

PROOF. The first part is the same as that in the case of  $g \in \mathbf{Z}^{\Delta(1)}$  (cf. [7, Lemma 2.3]). Since  $n(\rho)$  is a primitive vector and the pairing  $\langle \cdot, \cdot \rangle$  is non-degenerate, we have  $\square_g \cap M = \square_{[g]} \cap M$ , where  $[g]$  denotes the integral vector  $([g_\rho])_{\rho \in \Delta(1)}$ . On the other hand, we have  $\mathcal{O}_X(D_g) = \mathcal{O}_X(D_{[g]})$  by definition. Hence we may assume that  $g \in \mathbf{Z}^{\Delta(1)}$ . In this case, the assertion follows from [5, p. 42, Theorem] (cf. [7, Lemma 2.3]). The rest is obvious.

Recall that a  $\Delta$ -linear support function  $h \in \text{SF}(N, \Delta, \mathcal{Q})$  is said to be *strictly upper convex* with respect to  $\Delta$  if  $h$  is upper convex, namely  $h(n) + h(n') \leq h(n + n')$  for all  $n, n' \in N_{\mathbf{R}}$ , and  $\Delta$  is the coarsest among the fans  $\Delta'$  in  $N$  for which  $h$  is  $\Delta'$ -linear (cf. [7, p. 82]).

LEMMA 1.2. *Let  $X = T_N \text{emb}(\Delta)$  be an  $r$ -dimensional complete torus embedding and  $h \in \text{SF}(N, \Delta, \mathcal{Q})$ . Then  $D_h$  is an ample  $\mathcal{Q}$ -divisor if and only if  $h$  is strictly upper convex with respect to  $\Delta$ .*

PROOF. This easily follows from [7, Corollary 2.14]. q.e.d.

PROPOSITION 1.3. *Let  $P$  be a rational convex  $r$ -polytope in  $M_{\mathbf{R}} = \mathbf{R}^r$  and  $h_P: N_{\mathbf{R}} \rightarrow \mathbf{R}$  the support function for  $P$  defined by  $h_P(n) = \inf\{\langle m, n \rangle; m \in P\}$ . Then there exists a unique finite complete fan  $\Delta_P$  in  $N$  such that  $h_P$  is strictly upper convex  $\Delta_P$ -linear with respect to  $\Delta_P$ . We denote the corresponding  $r$ -dimensional projective torus embedding  $T_N \text{emb}(\Delta_P)$  and the ample  $T$ -stable  $\mathcal{Q}$ -divisor  $D_{h_P}$  by  $X(P)$  and  $D(P)$ . Conversely, every pair of a projective torus embedding and a  $T$ -stable ample  $\mathcal{Q}$ -divisor on it is obtained from a rational convex  $r$ -polytope in  $M_{\mathbf{R}}$  in this way.*

PROOF. The first part follows from [7, Theorem A.18 and Corollary A.19]. Then, by (1.2),  $D(P)$  is a  $T$ -stable ample  $\mathcal{Q}$ -divisor on  $X(P)$ . Conversely, given a projective torus embedding  $X$  with a  $T$ -stable ample  $\mathcal{Q}$ -divisor  $D$ , there exist a complete fan  $\Delta$  and a strictly upper convex  $\Delta$ -linear support function  $h \in \text{SF}(N, \Delta, \mathcal{Q})$  with  $X = T_N \text{emb}(\Delta)$  and  $D = D_h$ , by (1.2). Set  $\square_h = \{u \in M_{\mathbf{R}}; \langle u, n(\rho) \rangle \geq h(n(\rho)) \text{ for all } \rho \in \Delta(1)\}$ . By construction and [7, Theorem A.18 and Corollary A.19], we have  $X = X(\square_h)$  and  $D = D(\square_h)$ . q.e.d.

REMARK 1.4. In (1.3),  $D(P)$  is a Cartier divisor if and only if  $P$  is an *integral*

convex  $r$ -polytope, namely, a convex polytope whose vertices have integral coordinates.  $D(P)$  is a Weil divisor if and only if  $P$  is *facet-reticular*, that is, each supporting hyperplane carried by a *facet* (maximal face) of  $P$  contains an element of  $M$ .

PROPOSITION 1.5. *Let  $P$  be a rational convex  $r$ -polytope in  $M_{\mathbf{R}}$ . Then the graded semigroup ring*

$$R(P) = \bigoplus_{n \geq 0} \left\{ \sum_{m \in nP \cap M} ke(m) \right\} t^n$$

is isomorphic as a graded  $k$ -algebra to the graded ring  $R(X(P), D(P))$  associated with the projective torus embedding  $X(P)$  and the  $T$ -stable ample  $\mathcal{Q}$ -divisor  $D(P)$ . Consequently,  $\text{Proj}(R(P))$  is isomorphic to  $X(P)$  and the sheaf  $\mathcal{O}_{X(P)}(n) := R(P)(n)^\sim$  on  $\text{Proj}(R(P))$  corresponds via this isomorphism to  $\mathcal{O}_{X(P)}(nD(P))$  for all  $n \in \mathbf{Z}$ .

PROOF. Since  $\square_{nh_P} = nP$  and  $D(nP) = D_{nh_P}$  for all  $n \in \mathbf{N}$ , we have

$$H^0(X(P), \mathcal{O}_{X(P)}(nD(P))) = \sum_{m \in nP \cap M} ke(m)$$

by (1.1). This implies that  $R(P) \simeq R(X(P), D(P))$ . The rest follows from a standard argument in the theory of Demazure's construction (cf. [10, Lemma 2.1]). q.e.d.

COROLLARY 1.6. *For an  $r$ -dimensional projective torus embedding  $X = T_{\mathbf{N}}\text{emb}(\Delta)$  and a strictly upper convex  $\Delta$ -linear support function  $h \in \text{SF}(\mathbf{N}, \Delta, \mathcal{Q})$ , we have:*

- (a)  $\dim_k H^0(X, \mathcal{O}_X(nD_h)) = \begin{cases} \#(n\square_h \cap M) & \text{if } n \geq 0 \\ 0 & \text{if } n < 0; \end{cases}$
- (b)  $\dim_k H^i(X, \mathcal{O}_X(nD_h)) = 0$  for  $0 < i < r$  and all  $n \in \mathbf{Z}$ ;
- (c)  $\dim_k H^r(X, \mathcal{O}_X(nD_h)) = \begin{cases} 0 & \text{if } n \geq 0 \\ \#(\text{int}((-n)\square_h) \cap M) & \text{if } n < 0. \end{cases}$

PROOF. (a) follows from (1.1). Since  $R(X, D_h)$  is a normal numerical semigroup ring by (1.3) and (1.5),  $R(X, D_h)$  is normal and Cohen-Macaulay by a theorem of Hochster [4]. Therefore, (b) follows from [10, Corollary 2.4]. By the Serre duality, we have  $\text{Hom}_k(H^r(X, \mathcal{O}_X(nD_h)), k) \simeq H^0(X, \mathcal{O}_X(-[nD_h] + K_X))$ , where  $K_X$  denotes a canonical divisor on  $X$ . Since  $K_X = -\sum_{\rho \in \Delta(1)} V(\rho)$  (cf. [5, p. 29]), (c) follows from (1.1). q.e.d.

REMARK 1.7. Let  $P$  be a rational convex  $r$ -polytope in  $\mathbf{R}^r$  and  $m = \min\{i \in \mathbf{N}; i > 0 \text{ and } iP \text{ is integral}\}$ . By (1.3), (1.5) and (1.6), we have  $\#(nP \cap \mathbf{Z}^r) = \chi(X(P), \mathcal{O}_{X(P)}(nD(P)))$  for  $n \geq 0$  and  $\#(\text{int}((-n)P) \cap \mathbf{Z}^r) = (-1)^r \chi(X(P), \mathcal{O}_{X(P)}(nD(P)))$  for  $n < 0$ , where  $\chi(X(P), \mathcal{O}_{X(P)}(nD(P))) := \sum_{j=0}^r (-1)^j \dim_k H^j(X(P), \mathcal{O}_{X(P)}(nD(P)))$ . By a result due to Snapper and Kleiman, for every  $d \in \mathbf{Z}$ , there exists a polynomial  $P_d(\lambda)$  with coefficients in  $\mathcal{Q}$  such

that  $\chi(X(P), \mathcal{O}_{X(P)}((d+m\lambda)D(P))) = P_d(\lambda)$ . Thus we recover the reciprocity theorem for Ehrhart quasi-polynomials. (See, for example, [7, Proposition 2.24], [9, (4.6)]).

**2. Criteria for Gorenstein property.**

LEMMA 2.1. *Let  $\Delta$  be a complete fan in  $N$  and  $h \in \text{SF}(N, \Delta, \mathcal{Q})$  a strictly upper convex  $\Delta$ -linear support function. Set  $\square_h = \{u \in M_{\mathbf{R}}; \langle u, n(\rho) \rangle \geq h(n(\rho)) \text{ for each } \rho \in \Delta(1)\}$ . Suppose that  $h$  has negative values except at the origin, or equivalently,  $\square_h$  contains the origin in its interior. Then the set of vertices of the polar convex polyhedral set  $(\square_h)^\circ := \{v \in N_{\mathbf{R}}; \langle u, v \rangle \geq -1 \text{ for all } u \in \square_h\}$  for  $\square_h$  is  $\{-1/h(n(\rho))n(\rho); \rho \in \Delta(1)\}$ .*

PROOF. By [7, Corollary A.19], there exists a bijection from  $\Delta(1)$  to the set  $\mathcal{F}^{r-1}(\square_h)$  of  $(r-1)$ -dimensional faces of  $\square_h$  sending  $\rho \in \Delta(1)$  to  $Q_\rho := \{u \in \square_h; \langle u, n(\rho) \rangle = h(n(\rho))\}$ . Also, by [7, Proposition A.17], there exists a bijection from  $\mathcal{F}^{r-1}(\square_h)$  to the set of vertices of  $(\square_h)^\circ$  sending an  $(r-1)$ -dimensional face  $Q$  to  $Q^* := \{v \in (\square_h)^\circ; \langle u, v \rangle = -1 \text{ for all } u \in Q\}$ . Then we observe that  $(Q_\rho)^*$  is  $-1/h(n(\rho))n(\rho)$ . q.e.d.

For a Noetherian graded ring  $R$  with the canonical module  $K_R$  of  $R$ , we consider the integer  $a(R)$  defined by  $a(R) = -\min\{m \in \mathbf{Z}; (K_R)_m \neq 0\}$ . For details concerning this integer, see [1, p. 194].

PROPOSITION 2.2 (cf. [2], [3]). *For a rational convex  $r$ -polytope  $P$  in  $M_{\mathbf{R}} = \mathbf{R}^r$  with  $M = \mathbf{Z}^r$  and a positive integer  $\delta$ , the following are equivalent:*

- (a) *The semigroup ring  $R(P)$  over  $k$  is a Gorenstein ring with  $a(R(P)) = -\delta$ .*
- (b) *The projective torus embedding  $X(P) = T_N \text{emb}(\Delta_P)$  over  $k$ , and the ample  $\mathcal{Q}$ -divisor  $D(P) = \sum_{\rho \in \Delta_P(1)} (p_\rho/q_\rho) \cdot V(\rho)$  ( $q_\rho > 0$ ,  $p_\rho$  and  $q_\rho$  are coprime) satisfy the following:*
  - (b1) *There exist a positive integer  $r_\rho$  for each  $\rho \in \Delta_P(1)$  and a character  $m \in M$  such that*

$$\delta D(P) + \text{div}(e(m)) = \sum_{\rho \in \Delta_P(1)} (1/r_\rho) \cdot V(\rho);$$

- (b2)  *$\delta$  and  $q_\rho$  are coprime for each  $\rho \in \Delta_P(1)$ .*
- (c) *(Hibi's condition)  $P$  satisfies the following:*
  - (c1) *There exists a character  $m \in M$  such that the polar polyhedral set  $(\delta P - m)^\circ := \{v \in N_{\mathbf{R}}; \langle u, v \rangle \geq -1 \text{ for all } u \in \delta P - m\}$  for  $\delta P - m := \{\delta p - m \in M_{\mathbf{R}}; p \in P\}$  is an integral convex  $r$ -polytope;*
  - (c2) *The convex hull  $\tilde{P}$  of the set  $\{(u, 0) \in M_{\mathbf{R}} \times \mathbf{R}; u \in P\} \cup \{(0, \dots, 0, 1/\delta)\}$  in  $M_{\mathbf{R}} \times \mathbf{R}$  is facet-reticular (cf. (1.4)).*

PROOF. (a) $\Leftrightarrow$ (b): By (1.5),  $R(P)$  is isomorphic to  $R(X(P), D(P))$  and, therefore,  $R(X(P), D(P))$  is Cohen-Macaulay (cf. [4]). Since a canonical divisor  $K_{X(P)}$  on  $X(P)$  is  $-\sum_{\rho \in \Delta_P(1)} V(\rho)$  (cf. [5, p. 29, Theorem 9, III.d]), it follows from [10, Corollary 2.9] that  $R(P)$  is a Gorenstein ring with  $a(R(P)) = -\delta$  if and only if there exists a character

$m \in M$  such that  $\delta D(P) + \text{div}(e(m)) = \sum_{\rho \in \Delta_P(1)} (1/q_\rho) \cdot V(\rho)$ . Note that a semi-invariant rational function  $f \in K(X(P))^*$  is a scalar multiple of a character  $m \in M$ .

Suppose (a) holds. By the preceding remark, we have the relation above and, therefore, (b1) holds. Rewriting this relation, we have  $\text{div}(e(m)) = \sum_{\rho \in \Delta_P(1)} \{(1 - \delta p_\rho)/q_\rho\} \cdot V(\rho)$ . Hence  $(1 - \delta p_\rho)/q_\rho$  is an integer and, therefore,  $\delta$  and  $q_\rho$  are coprime for each  $\rho \in \Delta_P(1)$ .

Conversely, suppose (b) holds. By the preceding remark, we claim that  $r_\rho = q_\rho$  for each  $\rho \in \Delta_P(1)$ . Since  $r_\rho$  is a factor of  $q_\rho$ ,  $b_\rho := (q_\rho/r_\rho)$  is a positive integer. Then, by (b1),  $(b_\rho - \delta p_\rho)/(r_\rho b_\rho)$  is an integer and, therefore,  $b_\rho$  is a factor of  $\delta p_\rho$ . Hence we have  $b_\rho = 1$  for each  $\rho \in \Delta_P(1)$  as required, because neither  $\delta$  nor  $p_\rho$  has any common factor with  $q_\rho$ .

(b1)  $\Rightarrow$  (c1): Set  $g = \delta h_P - m \in \text{SF}(N, \Delta_P, \mathbf{Q})$ . Since  $D_g = \delta D(P) + \text{div}(e(m))$  and  $D_g$  is ample,  $g$  is strictly upper convex and  $g(n(\rho)) = -(1/r_\rho)$  for each  $\rho \in \Delta_P(1)$ . Therefore, by (2.1), the set of vertices of the polar convex polyhedral set  $(\square_g)^\circ$  is  $\{r_\rho n(\rho); \rho \in \Delta_P(1)\}$  ( $= \{-(1/g(n(\rho)))n(\rho); \rho \in \Delta_P(1)\}$ ). On the other hand, we have  $\square_g = \delta P - m$  by definition. Therefore  $(\delta P - m)^\circ$  is an integral convex polytope.

(c1)  $\Rightarrow$  (b1): Set  $g = \delta h_P - m \in \text{SF}(N, \Delta_P, \mathbf{Q})$ . Since  $g$  is strictly upper convex with respect to  $\Delta_P$  and  $O \in \text{int}(\delta P - m)$ , it follows from (2.1) that the vertex set of  $(\delta P - m)^\circ$  is  $\{-(1/g(n(\rho)))n(\rho); \rho \in \Delta_P(1)\}$ . Hence, by assumption,  $-(1/g(n(\rho)))n(\rho)$  is an integral vector. Since  $n(\rho)$  is a primitive integral vector and  $g \in \text{SF}(N, \Delta_P, \mathbf{Q})$  is negative-valued,  $r_\rho := -1/(g(n(\rho)))$  is a positive integer for each  $\rho \in \Delta_P(1)$  and  $\delta D(P) + \text{div}(e(m)) = D_g = \sum_{\rho} (1/r_\rho) \cdot V(\rho)$ .

(b2)  $\Leftrightarrow$  (c2): Since the supporting hyperplane carried by the facet of  $P$  corresponding to  $\rho \in \Delta_P(1)$  is  $H_\rho = \{u \in M_{\mathbf{R}}; \langle u, n(\rho) \rangle = h_\rho(n(\rho))\}$ , the supporting hyperplane carried by a facet of  $\tilde{P}$  is of the form  $\tilde{H}_\rho := \{(u, x) \in M_{\mathbf{R}} \times \mathbf{R}; \delta x + (1/h_\rho(n(\rho)))\langle u, n(\rho) \rangle = 1\}$  or  $\{(u, 0) \in M_{\mathbf{R}} \times \mathbf{R}\}$ . Since  $h_\rho(n(\rho)) = -(p_\rho/q_\rho)$  and  $n(\rho)$  is a primitive vector,  $\delta$  and  $q_\rho$  are coprime if and only if  $\tilde{H}_\rho \cap (M \times \mathbf{Z})$  is non-empty. q.e.d.

REMARK 2.3. The equivalence between the conditions (a) and (c) in (2.2) is originally due to Hibi [3]. Combining the equivalence between (a) and (c) in (2.2) and a theorem of Stanley [8, Theorem 4.4], we get another proof for theorems of Hibi [2], [3]. Our proof makes clear why the condition (c2) in (2.2) is needed, in terms of Demazure's construction. Indeed, let  $R(X, D)$  be a Cohen-Macaulay graded ring obtained from a normal projective variety  $X$  and an ample  $\mathbf{Q}$ -divisor  $D = \sum_V (p_V/q_V) \cdot V$ , with  $V$  running through irreducible subvarieties of codimension 1, where  $q_V > 0$  and  $p_V, q_V$  are coprime for each  $V$ . Then it follows from [10, Corollary 2.9] that  $R(X, D)$  is Gorenstein if the Veronese subring  $R(X, D)^{(d)}$  of order  $d$  is Gorenstein for an integer  $d$  such that  $a(R(X, D)) \equiv 0 \pmod{d}$  and that  $d$  and  $q_V$  are coprime for each  $V$ .

COROLLARY 2.4 (cf. [7, (2.20)]). *For a rational convex  $r$ -polytope  $P$  in  $M_{\mathbf{R}} = \mathbf{R}^r$  with  $M = \mathbf{Z}^r$  and a positive integer  $\delta$ , the following are equivalent:*

(a)  *$P$  is integral and there exists a character  $m \in M$  such that the polar polyhedral set  $(\delta P - m)^\circ$  for  $\delta P - m$  is an integral convex  $r$ -polytope;*

(b) The  $\mathcal{Q}$ -divisor  $D(P)$  on the projective torus embedding  $X(P)$  is an ample Cartier divisor. The invertible sheaf  $\mathcal{O}_{X(P)}(-\delta D(P))$  is isomorphic to the canonical sheaf  $\omega_{X(P)}$ .

PROOF. It follows from (1.4) and (2.2) that (a) holds if and only if  $D(P)$  is a Cartier divisor and there exists a character  $m \in M$  such that  $\delta D(P) + \text{div}(e(m)) = \sum_{\rho \in \Delta_P(1)} V(\rho)$ . Since a canonical divisor  $K_{X(P)}$  on  $X(P)$  is  $-\sum_{\rho \in \Delta_P(1)} V(\rho)$ , (a) is equivalent to (b).  
q.e.d.

Since every Cartier divisor on a complete torus embedding is linearly equivalent to a  $T$ -stable Cartier divisor (cf. [6, Proposition 6.1]), we have:

COROLLARY 2.5. Let  $X$  be a projective torus embedding and  $D$  an ample Cartier divisor. Then  $R(X, D)$  is Gorenstein if and only if the canonical sheaf  $\omega_X$  on  $X$  is isomorphic to an invertible sheaf  $\mathcal{O}_X(-\delta D)$  for a positive integer  $\delta$ .

THEOREM 2.6. Every projective torus embedding  $X$  has a  $T$ -stable ample  $\mathcal{Q}$ -divisor  $D$  such that  $R(X, D)$  is a Gorenstein ring with  $a(R(X, D)) = -1$ .

PROOF. By assumption,  $X = T_N \text{emb}(\Delta)$  has a  $T$ -stable ample Cartier divisor  $E$  of the form  $E = \sum_{\rho \in \Delta(1)} a_\rho \cdot V(\rho)$ ,  $a_\rho > 0$ . Set  $c = \text{L.C.M.}\{a_\rho; \rho \in \Delta(1)\}$  and  $D = (1/c)E$ . By (1.3),  $(X, D)$  corresponds to a rational polytope  $P$  in  $M_{\mathbf{R}}$ . Then, by (1.5) and (2.2),  $R(X, D)$  is a Gorenstein ring with  $a(R(X, D)) = -1$ , as required.  
q.e.d.

#### REFERENCES

- [ 1 ] S. GOTO AND K.-I. WATANABE, On graded rings, I, J. Math. Soc. Japan 30 (1978), 179–213.
- [ 2 ] T. HIBI, Dual polytopes of rational convex polytopes, preprint.
- [ 3 ] T. HIBI, A combinatorial self-reciprocity theorem for Ehrhart quasi-polynomials of rational convex polytopes, preprint.
- [ 4 ] M. HOCHSTER, Rings of invariants of tori, Cohen-Macaulay rings generated by monomials, and polytopes, Ann. of Math. 96 (1972), 318–337.
- [ 5 ] G. KEMPF, F. KNUDSEN, D. MUMFORD AND B. SAINT-DONAT, Toroidal embeddings I, Lecture Notes in Math. 339, Springer-Verlag, Berlin/New York, 1973.
- [ 6 ] T. ODA, Lectures on Torus Embeddings and Applications, Tata Inst. Fund. Res. Lectures on Math. and Phys. 58, Bombay, 1978.
- [ 7 ] T. Oda, Convex Bodies and Algebraic Geometry, Erg. der Math. (3) 15, Springer-Verlag, Berlin/New York, 1988.
- [ 8 ] R. P. STANLEY, Hilbert functions of graded algebras, Adv. in Math. 28 (1978), 57–83.
- [ 9 ] R. P. STANLEY, Enumerative Combinatorics, Volume I, Wadsworth, Monterey, Calif., 1986.
- [ 10 ] K.-I. WATANABE, Some remarks concerning Demazure's construction of normal graded rings, Nagoya Math. J. 83 (1981), 203–211.

DEPARTMENT OF MATHEMATICS  
SCHOOL OF SCIENCE AND ENGINEERING  
WASEDA UNIVERSITY  
OHKUBO, SHINJUKU, TOKYO 169  
JAPAN

