

University of New Mexico

UNM Digital Repository

Mathematics and Statistics Faculty and Staff
Publications

Academic Department Resources

11-2017


Graph Structures in Bipolar Neutrosophic Environment

Florentin Smarandache

Muhammad Akram

Muzzamal Sitara

Follow this and additional works at: https://digitalrepository.unm.edu/math_fsp

 Part of the [Algebraic Geometry Commons](#), [Analysis Commons](#), [Discrete Mathematics and Combinatorics Commons](#), [Geometry and Topology Commons](#), and the [Logic and Foundations Commons](#)

Article

Graph Structures in Bipolar Neutrosophic Environment

Muhammad Akram ^{1,*} , Muzzamal Sitara ¹ and Florentin Smarandache ^{2,*} 

¹ Department of Mathematics, University of the Punjab, New Campus, Lahore 54590, Pakistan; muzzamalsitara@gmail.com

² Mathematics & Science Department, University of New Mexico, 705 Gurley Ave., Gallup, NM 87301, USA

* Correspondence: m.akram@pucit.edu.pk (M.A.); fsmarandache@gmail.com (F.S.); Tel.: +92-42-99231241 (M.A.)

Received: 14 September 2017; Accepted: 27 October 2017; Published: 6 November 2017

Abstract: A bipolar single-valued neutrosophic (BSVN) graph structure is a generalization of a bipolar fuzzy graph. In this research paper, we present certain concepts of BSVN graph structures. We describe some operations on BSVN graph structures and elaborate on these with examples. Moreover, we investigate some related properties of these operations.

Keywords: graph structure; bipolar single-valued neutrosophic (BSVN) graph structure

MSC: 03E72; 05C72; 05C78; 05C99

1. Introduction

Fuzzy graphs are mathematical models for dealing with combinatorial problems in different domains, including operations research, optimization, computer science and engineering. In 1965, Zadeh [1] proposed fuzzy set theory to deal with uncertainty in abundant meticulous real-life phenomena. Fuzzy set theory is affluently applicable in real-time systems consisting of information with different levels of precision. Subsequently, Atanassov [2] introduced the idea of intuitionistic fuzzy sets in 1986. However, for many real-life phenomena, it is necessary to deal with the implicit counter property of a particular event. Zhang [3] initiated the concept of bipolar fuzzy sets in 1994. Evidently bipolar fuzzy sets and intuitionistic fuzzy sets seem to be similar, but they are completely different sets. Bipolar fuzzy sets have large number of applications in image processing and spatial reasoning. Bipolar fuzzy sets are more practical, advantageous and applicable in real-life phenomena. However, both bipolar fuzzy sets and intuitionistic fuzzy sets cope with incomplete information, because they do not consider indeterminate or inconsistent information that clearly appears in many systems of different fields, including belief systems and decision-support systems. Smarandache [4] introduced neutrosophic sets as a generalization of fuzzy sets and intuitionistic fuzzy sets. A neutrosophic set has three constituents: truth membership, indeterminacy membership and falsity membership, for which each membership value is a real standard or non-standard subset of the unit interval $[0^-, 1^+]$. In real-life problems, neutrosophic sets can be applied more appropriately by using the single-valued neutrosophic sets defined by Smarandache [4] and Wang et al. [5]. Deli et al. [6] considered bipolar neutrosophic sets as a generalization of bipolar fuzzy sets. They also studied some operations and applications in decision-making problems.

On the basis of Zadeh's fuzzy relations [7], Kauffman defined fuzzy graphs [8]. In 1975, Rosenfeld [9] discussed a fuzzy analogue of different graph-theoretic ideas. Later on, Bhattacharya [10] gave some remarks on fuzzy graphs in 1987. Akram [11] first introduced the notion of bipolar fuzzy graphs. In 2011, Dinesh and Ramakrishnan [12] studied fuzzy graph structures and discussed their properties. In 2016, Akram and Akmal [13] proposed the concept of bipolar fuzzy graph structures. Certain concepts on graphs have been discussed in [14–19]. Ye [20–22] considered several applications

of single-valued neutrosophic sets. In this research paper, we present certain concepts of bipolar single-valued neutrosophic graph structures (BSVNGSs). We introduce some operations on BSVNGSs and elaborate on these with examples. Moreover, we investigate some relevant and remarkable properties of these operators.

We have used standard definitions and terminologies in this paper. For other notations, terminologies and applications not mentioned in the paper, the readers are referred to [23–29].

2. Bipolar Single-Valued Neutrosophic Graph Structures

Definition 1. [4] A neutrosophic set N on a non-empty set V is an object of the form

$$N = \{(v, T_N(v), I_N(v), F_N(v)) : v \in V\}$$

where $T_N, I_N, F_N : V \rightarrow [0^-, 1^+]$ and there is no restriction on the sum of $T_N(v)$, $I_N(v)$ and $F_N(v)$ for all $v \in V$.

Definition 2. [5] A single-valued neutrosophic set N on a non-empty set V is an object of the form

$$N = \{(v, T_N(v), I_N(v), F_N(v)) : v \in V\}$$

where $T_N, I_N, F_N : V \rightarrow [0, 1]$ and the sum of $T_N(v)$, $I_N(v)$ and $F_N(v)$ is confined between 0 and 3 for all $v \in V$.

Definition 3. [23] A BSVN set on a non-empty set V is an object of the form

$$B = \{(v, T_B^P(v), I_B^P(v), F_B^P(v), T_B^N(v), I_B^N(v), F_B^N(v)) : v \in V\}$$

where $T_B^P, I_B^P, F_B^P : V \rightarrow [0, 1]$ and $T_B^N, I_B^N, F_B^N : V \rightarrow [-1, 0]$. The positive values $T_B^P(v)$, $I_B^P(v)$ and $F_B^P(v)$ denote the truth, indeterminacy and falsity membership values of an element $v \in V$, whereas negative values $T_B^N(v)$, $I_B^N(v)$ and $F_B^N(v)$ indicate the implicit counter property of truth, indeterminacy and falsity membership values of an element $v \in V$.

Definition 4. [23] A BSVN graph on a non-empty set V is a pair $G = (B, R)$, where B is a BSVN set on V and R is a BSVN relation in V such that

$$\begin{aligned} T_R^P(b, d) &\leq T_B^P(b) \wedge T_B^P(d), & I_R^P(b, d) &\leq I_B^P(b) \wedge I_B^P(d), & F_R^P(b, d) &\leq F_B^P(b) \vee F_B^P(d), \\ T_R^N(b, d) &\geq T_B^N(b) \vee T_B^N(d), & I_R^N(b, d) &\geq I_B^N(b) \vee I_B^N(d), & F_R^N(b, d) &\geq F_B^N(b) \wedge F_B^N(d) \end{aligned}$$

for all $b, d \in V$.

We now define the BSVNGS.

Definition 5. [30] A BSVNGS of a graph structure $\check{G}_s = (V, V_1, V_2, \dots, V_m)$ is denoted by $\check{G}_{bn} = (B, B_1, B_2, \dots, B_m)$, where $B = \langle b, T^P(b), I^P(b), F^P(b), T^N(b), I^N(b), F^N(b) \rangle$ is a BSVN set on the set V and $B_k = \langle (b, d), T_k^P(b, d), I_k^P(b, d), F_k^P(b, d), T_k^N(b, d), I_k^N(b, d), F_k^N(b, d) \rangle$ are the BSVN sets on V_k such that

$$\begin{aligned} T_k^P(b, d) &\leq \min\{T^P(b), T^P(d)\}, & I_k^P(b, d) &\leq \min\{I^P(b), I^P(d)\}, & F_k^P(b, d) &\leq \max\{F^P(b), F^P(d)\}, \\ T_k^N(b, d) &\geq \max\{T^N(b), T^N(d)\}, & I_k^N(b, d) &\geq \max\{I^N(b), I^N(d)\}, & F_k^N(b, d) &\geq \min\{F^N(b), F^N(d)\} \end{aligned}$$

for all $b, d \in V$. Note that $0 \leq T_k^P(b, d) + I_k^P(b, d) + F_k^P(b, d) \leq 3$, $-3 \leq T_k^N(b, d) + I_k^N(b, d) + F_k^N(b, d) \leq 0$ for all $(b, d) \in V_k$, and (b, d) represents an edge between two vertices b and d . In this paper we use bd in place of (b, d) .

Example 1. Consider a graph structure $\check{G}_s = (V, V_1, V_2, V_3)$ such that $V = \{b_1, b_2, b_3, b_4, b_5, b_6, b_7, b_8\}$, $V_1 = \{b_1b_2, b_2b_7, b_4b_8, b_6b_8, b_5b_6, b_3b_4\}$, $V_2 = \{b_1b_5, b_5b_7, b_3b_6, b_7b_8\}$, and $V_3 = \{b_1b_3, b_2b_4\}$. Let B be a BSVN set on V given in Table 1 and B_1, B_2 and B_3 be BSVN sets on V_1, V_2 and V_3 , respectively, given in Table 2.

Table 1. Bipolar single-valued neutrosophic (BSVN) set B on vertex set V .

B	b_1	b_2	b_3	b_4	b_5	b_6	b_7	b_8
T^P	0.5	0.4	0.4	0.5	0.3	0.4	0.5	0.3
I^P	0.4	0.3	0.4	0.4	0.2	0.4	0.5	0.4
F^P	0.6	0.5	0.4	0.6	0.4	0.7	0.4	0.5
T^N	-0.5	-0.4	-0.4	-0.5	-0.3	-0.4	-0.5	-0.3
I^N	-0.4	-0.3	-0.4	-0.4	-0.2	-0.4	-0.5	-0.4
F^N	-0.6	-0.5	-0.4	-0.6	-0.4	-0.7	-0.4	-0.5

Table 2. BSVN sets B_1, B_2 and B_3 .

B_1	b_1b_2	b_2b_7	b_4b_8	b_6b_8	b_5b_6	b_3b_4	B_2	b_1b_5	b_5b_7	b_3b_6	b_7b_8	B_3	b_1b_3	b_2b_4
T^P	0.4	0.4	0.3	0.3	0.3	0.4	T^P	0.3	0.3	0.4	0.3	T^P	0.4	0.4
I^P	0.3	0.3	0.4	0.4	0.2	0.4	I^P	0.2	0.2	0.4	0.4	I^P	0.4	0.3
F^P	0.6	0.5	0.6	0.7	0.7	0.6	F^P	0.6	0.4	0.7	0.5	F^P	0.6	0.6
T^N	-0.4	-0.4	-0.3	-0.3	-0.3	-0.4	T^N	-0.3	-0.3	-0.4	-0.3	T^N	-0.4	-0.4
I^N	-0.3	-0.3	-0.4	-0.4	-0.2	-0.4	I^N	-0.2	-0.2	-0.4	-0.4	I^N	-0.4	-0.3
F^N	-0.6	-0.5	-0.6	-0.7	-0.7	-0.6	F^N	-0.6	-0.4	-0.7	-0.5	F^N	-0.6	-0.6

By direct calculations, it is easy to show that $\check{G}_{bn} = (B, B_1, B_2, B_3)$ is a BSVNGS. This BSVNGS is shown in Figure 1. Generated with LaTeXDraw 2.0.8 on Saturday March 11 20:30:24 PKT 2017.




Figure 1. A BSVN graph structure.

Definition 6. A BSVNGS $\check{G}_{bn} = (B, B_1, B_2, \dots, B_m)$ is called a B_k -cycle if $(supp(B), supp(B_1), supp(B_2), \dots, supp(B_m))$ is a B_k -cycle.

Example 2. Consider a BSVNGS $\check{G}_{bn} = (B, B_1, B_2)$ as shown in Figure 2.




Figure 2. A BSVN B_1 -cycle.

\check{G}_{bn} is a B_1 -cycle, as $(supp(B), supp(B_1), supp(B_2))$ is a B_1 -cycle, that is, b_2 - b_3 - b_4 - b_5 - b_2 .

Definition 7. A BSVNGS $\check{G}_{bn} = (B, B_1, B_2, \dots, B_m)$ is a BSVN fuzzy B_k -cycle (for any k) if \check{G}_{bn} is a B_k -cycle and no unique B_k -edge bd exists in \check{G}_{bn} , such that $T_{B_k}^P(bd) = \min\{T_{B_k}^P(ef) : ef \in B_k = supp(B_k)\}$, $I_{B_k}^P(bd) = \min\{I_{B_k}^P(ef) : ef \in B_k = supp(B_k)\}$, $F_{B_k}^P(bd) = \max\{F_{B_k}^P(ef) : ef \in B_k = supp(B_k)\}$, $T_{B_k}^N(bd) = \max\{T_{B_k}^N(ef) : ef \in B_k = supp(B_k)\}$, $I_{B_k}^N(bd) = \max\{I_{B_k}^N(ef) : ef \in B_k = supp(B_k)\}$ or $F_{B_k}^N(bd) = \min\{F_{B_k}^N(ef) : ef \in B_k = supp(B_k)\}$.

Example 3. Consider a BSVNGS $\check{G}_{bn} = (B, B_1, B_2)$ as depicted in Figure 3.




Figure 3. A BSVN fuzzy B_2 -cycle.

$T_{B_2}^P(bd) = \min\{T_{B_2}^P(ef) : ef \in B_2 = supp(B_2)\}$, $I_{B_2}^P(bd) = \min\{I_{B_2}^P(ef) : ef \in B_2 = supp(B_2)\}$, $F_{B_2}^P(bd) = \max\{F_{B_2}^P(ef) : ef \in B_2 = supp(B_2)\}$, $T_{B_2}^N(bd) = \max\{T_{B_2}^N(ef) : ef \in B_2 = supp(B_2)\}$, $I_{B_2}^N(bd) = \max\{I_{B_2}^N(ef) : ef \in B_2 = supp(B_2)\}$ or $F_{B_2}^N(bd) = \min\{F_{B_2}^N(ef) : ef \in B_2 = supp(B_2)\}$.

Definition 8. A sequence of vertices (distinct) in a BSVNGS $\check{G}_{bn} = (B, B_1, B_2, \dots, B_m)$ is called a B_k -path, that is, b_1, b_2, \dots, b_m , such that $b_{k-1}b_k$ is a BSVN B_k -edge, for all $k = 2, \dots, m$.

Example 4. Consider a BSVNGS $\check{G}_{bn} = (B, B_1, B_2)$ as represented in Figure 4.




Figure 4. A BSVN B_2 -path.

In this BSVNGS, the sequence of distinct vertices b_1, b_4, b_3, b_2 is a BSVN B_2 -path.

Definition 9. Let $\check{G}_{bn} = (B, B_1, B_2, \dots, B_m)$ be a BSVNGS. The positive truth strength $T^P.P_{B_k}$, positive falsity strength $F^P.P_{B_k}$, and positive indeterminacy strength $I^P.P_{B_k}$ of a B_k -path, $P_{B_k} = b_1, b_2, \dots, b_n$, are defined as

$$T^P.P_{B_k} = \bigwedge_{h=2}^n [T_{B_k}^P(b_{h-1}b_h)], I^P.P_{B_k} = \bigwedge_{h=2}^n [I_{B_k}^P(b_{h-1}b_h)], F^P.P_{B_k} = \bigvee_{h=2}^n [F_{B_k}^P(b_{h-1}b_h)]$$

Similarly, the negative truth strength $T^N.P_{B_k}$, negative falsity strength $F^N.P_{B_k}$, and negative indeterminacy strength $I^N.P_{B_k}$ of a B_k -path are defined as

$$T^N.P_{B_k} = \bigvee_{h=2}^n [T_{B_k}^N(b_{h-1}b_h)], I^N.P_{B_k} = \bigvee_{h=2}^n [I_{B_k}^N(b_{h-1}b_h)], F^N.P_{B_k} = \bigwedge_{h=2}^n [F_{B_k}^N(b_{h-1}b_h)]$$

Example 5. Consider a BSVNGS $\check{G}_{bn} = (B, B_1, B_2, B_3)$ as shown in Figure 5.




Figure 5. A bipolar single-valued neutrosophic graph structure (BSVNGS) $\check{G}_{bn} = (B, B_1, B_2, B_3)$.

In this BSVNGS, there is a B_1 -path, that is, $P_{B_1} = b_5, b_6, b_8, b_4, b_3$. Thus, $T^N.P_{B_1} = -0.2$, $I^N.P_{B_1} = -0.2$, $F^N.P_{B_2} = -0.6$, $T^P.P_{B_1} = 0.2$, $I^P.P_{B_1} = 0.2$ and $F^P.P_{B_2} = 0.6$.

Definition 10. Let $\check{G}_{bn} = (B, B_1, B_2, \dots, B_m)$ be a BSVNGS. Then

- The B_k -positive strength of connectedness of truth between two nodes b and d is defined by $T_{B_k}^{P\infty}(bd) = \bigvee_{l \geq 1} \{T_{B_k}^{Pl}(bd)\}$, such that $T_{B_k}^{Pl}(bd) = (T_{B_k}^{Pl-1} \circ T_{B_k}^{P1})(bd)$ for $l \geq 2$ and $T_{B_k}^{P2}(bd) = (T_{B_k}^{P1} \circ T_{B_k}^{P1})(bd) = \bigvee_y (T_{B_k}^{P1}(by) \wedge T_{B_k}^{P1}(yd))$.
- The B_k -positive strength of connectedness of indeterminacy between two nodes b and d is defined by $I_{B_k}^{P\infty}(bd) = \bigvee_{l \geq 1} \{I_{B_k}^{Pl}(bd)\}$, such that $I_{B_k}^{Pl}(bd) = (I_{B_k}^{Pl-1} \circ I_{B_k}^{P1})(bd)$ for $l \geq 2$ and $I_{B_k}^{P2}(bd) = (I_{B_k}^{P1} \circ I_{B_k}^{P1})(bd) = \bigvee_y (I_{B_k}^{P1}(by) \wedge I_{B_k}^{P1}(yd))$.
- The B_k -positive strength of connectedness of falsity between two nodes b and d is defined by $F_{B_k}^{P\infty}(bd) = \bigwedge_{l \geq 1} \{F_{B_k}^{Pl}(bd)\}$, such that $F_{B_k}^{Pl}(bd) = (F_{B_k}^{Pl-1} \circ F_{B_k}^{P1})(bd)$ for $l \geq 2$ and $F_{B_k}^{P2}(bd) = (F_{B_k}^{P1} \circ F_{B_k}^{P1})(bd) = \bigwedge_y (F_{B_k}^{P1}(by) \vee F_{B_k}^{P1}(yd))$.
- The B_k -negative strength of connectedness of truth between two nodes b and d is defined by $T_{B_k}^{N\infty}(bd) = \bigwedge_{l \geq 1} \{T_{B_k}^{Nl}(bd)\}$, such that $T_{B_k}^{Nl}(bd) = (T_{B_k}^{Nl-1} \circ T_{B_k}^{N1})(bd)$ for $l \geq 2$ and $T_{B_k}^{N2}(bd) = (T_{B_k}^{N1} \circ T_{B_k}^{N1})(bd) = \bigwedge_y (T_{B_k}^{N1}(by) \vee T_{B_k}^{N1}(yd))$.
- The B_k -negative strength of connectedness of indeterminacy between two nodes b and d is defined by $I_{B_k}^{N\infty}(bd) = \bigwedge_{l \geq 1} \{I_{B_k}^{Nl}(bd)\}$, such that $I_{B_k}^{Nl}(bd) = (I_{B_k}^{Nl-1} \circ I_{B_k}^{N1})(bd)$ for $l \geq 2$ and $I_{B_k}^{N2}(bd) = (I_{B_k}^{N1} \circ I_{B_k}^{N1})(bd) = \bigwedge_y (I_{B_k}^{N1}(by) \vee I_{B_k}^{N1}(yd))$.
- The B_k -negative strength of connectedness of falsity between two nodes b and d is defined by $F_{B_k}^{N\infty}(bd) = \bigvee_{l \geq 1} \{F_{B_k}^{Nl}(bd)\}$, such that $F_{B_k}^{Nl}(bd) = (F_{B_k}^{Nl-1} \circ F_{B_k}^{N1})(bd)$ for $l \geq 2$ and $F_{B_k}^{N2}(bd) = (F_{B_k}^{N1} \circ F_{B_k}^{N1})(bd) = \bigvee_y (F_{B_k}^{N1}(by) \wedge F_{B_k}^{N1}(yd))$.

Definition 11. Let $\check{G}_{bn} = (B, B_1, B_2, \dots, B_m)$ be a BSVNGS and “ b ” be a node in \check{G}_{bn} . Let $(B', B'_1, B'_2, \dots, B'_m)$ be a BSVN subgraph structure of \check{G}_{bn} induced by $B \setminus \{b\}$ such that $\forall e \neq b, f \neq b$

$$\begin{aligned}
 T_{B'_k}^P(b) &= I_{B'_k}^P(b) = F_{B'_k}^P(b) = T_{B'_k}^P(be) = I_{B'_k}^P(be) = F_{B'_k}^P(be) = 0 \\
 T_{B'_k}^N(b) &= I_{B'_k}^N(b) = F_{B'_k}^N(b) = T_{B'_k}^N(be) = I_{B'_k}^N(be) = F_{B'_k}^N(be) = 0 \\
 T_{B'_k}^P(e) &= T_B^P(e), I_{B'_k}^P(e) = I_B^P(e), F_{B'_k}^P(e) = F_B^P(e), T_{B'_k}^N(e) = T_B^N(e), I_{B'_k}^N(e) = I_B^N(e), F_{B'_k}^N(e) = F_B^N(e) \\
 T_{B'_k}^P(e) &= T_{B'_k}^P(e), I_{B'_k}^P(e) = I_{B'_k}^P(e), F_{B'_k}^P(e) = F_{B'_k}^P(e), T_{B'_k}^N(e) = T_{B'_k}^N(e), I_{B'_k}^N(e) = I_{B'_k}^N(e), F_{B'_k}^N(e) = F_{B'_k}^N(e) \\
 F_{B'_k}^N(e) &= F_{B'_k}^N(e), \forall \text{ edges } be, ef \in \check{G}_{bn}
 \end{aligned}$$

Then b is a BSVN fuzzy B_k cut-vertex if $T_{B_k}^{P\infty}(ef) > T_{B'_k}^{P\infty}(ef)$, $I_{B_k}^{P\infty}(ef) > I_{B'_k}^{P\infty}(ef)$, $F_{B_k}^{P\infty}(ef) > F_{B'_k}^{P\infty}(ef)$, $T_{B_k}^{N\infty}(ef) < T_{B'_k}^{N\infty}(ef)$, $I_{B_k}^{N\infty}(ef) < I_{B'_k}^{N\infty}(ef)$ and $F_{B_k}^{N\infty}(ef) < F_{B'_k}^{N\infty}(ef)$, for some $e, f \in B \setminus \{b\}$. Note that vertex b is a BSVN fuzzy $B_k - T^P$ cut-vertex if $T_{B_k}^{P\infty}(ef) > T_{B'_k}^{P\infty}(ef)$, it is a BSVN fuzzy $B_k - I^P$ cut-vertex if $I_{B_k}^{P\infty}(ef) > I_{B'_k}^{P\infty}(ef)$, and it is a BSVN fuzzy $B_k - F^P$ cut-vertex if $F_{B_k}^{P\infty}(ef) > F_{B'_k}^{P\infty}(ef)$. Moreover, vertex b is a BSVN fuzzy $B_k - T^N$ cut-vertex if $T_{B_k}^{N\infty}(ef) < T_{B'_k}^{N\infty}(ef)$, it is a BSVN fuzzy $B_k - I^N$ cut-vertex if $I_{B_k}^{N\infty}(ef) < I_{B'_k}^{N\infty}(ef)$ and it is a BSVN fuzzy $B_k - F^N$ cut-vertex if $F_{B_k}^{N\infty}(ef) < F_{B'_k}^{N\infty}(ef)$.

Example 6. Consider a BSVNGS $\check{G}_{bn} = (B, B_1, B_2)$ as depicted in Figure 6, and let $\check{G}'_{bh} = (B', B'_1, B'_2)$ be a BSVN subgraph structure of the BSVNGS \check{G}_{bn} , which is obtained through deletion of vertex b_2 .




Figure 6. A BSVNGS $\check{G}_{bn}^x = (B, B_1, B_2)$.

The vertex b_2 is a BSVN fuzzy $B_1 - I^P$ cut-vertex and a BSVN fuzzy $B_1 - I^N$ cut-vertex, because $I_{B_1}^{P\infty}(b_2b_5) = 0$, $I_{B_1}^{P\infty}(b_2b_5) = 0.5$, $I_{B_1}^{P\infty}(b_4b_3) = 0.7$, $I_{B_1}^{P\infty}(b_4b_3) = 0.7$, $I_{B_1}^{P\infty}(b_3b_5) = 0.3$, $I_{B_1}^{P\infty}(b_3b_5) = 0.4$, $I_{B_1}^{N\infty}(b_2b_5) = 0 > -0.5 = I_{B_1}^{N\infty}(b_2b_5)$, $I_{B_1}^{N\infty}(b_4b_3) = 0.7 = I_{B_1}^{N\infty}(b_4b_3)$ and $I_{B_1}^{N\infty}(b_3b_5) = -0.3 > -0.4 = I_{B_1}^{N\infty}(b_3b_5)$.

Definition 12. Suppose $\check{G}_{bn} = (B, B_1, B_2, \dots, B_m)$ is a BSVNGS and bd is a B_k -edge. Let $(B', B'_1, B'_2, \dots, B'_m)$ be a BSVN fuzzy spanning subgraph structure of \check{G}_{bn} , such that

$$\begin{aligned} T_{B'_k}^P(bd) = 0 = I_{B'_k}^P(bd) = F_{B'_k}^P(bd), T_{B'_k}^N(bd) = 0 = I_{B'_k}^N(bd) = F_{B'_k}^N(bd) \\ T_{B'_k}^P(gh) = T_{B_k}^P(gh), I_{B'_k}^P(gh) = I_{B_k}^P(gh) \\ F_{B'_k}^P(bd) = F_{B_k}^P(bd), T_{B'_k}^N(gh) = T_{B_k}^N(gh), I_{B'_k}^N(gh) = I_{B_k}^N(gh), F_{B'_k}^N(bd) = F_{B_k}^N(bd), \forall \text{ edges } gh \neq bd \end{aligned}$$

Then bd is a BSVN fuzzy B_k -bridge if $T_{B'_k}^{P\infty}(ef) > T_{B_k}^{P\infty}(ef)$, $I_{B'_k}^{P\infty}(ef) > I_{B_k}^{P\infty}(ef)$, $F_{B'_k}^{P\infty}(ef) > F_{B_k}^{P\infty}(ef)$, $T_{B'_k}^{N\infty}(ef) < T_{B_k}^{N\infty}(ef)$, $I_{B'_k}^{N\infty}(ef) < I_{B_k}^{N\infty}(ef)$ and $F_{B'_k}^{N\infty}(ef) < F_{B_k}^{N\infty}(ef)$, for some $e, f \in V$. Note that bd is a BSVN fuzzy $B_k - T^P$ bridge if $T_{B'_k}^{P\infty}(ef) > T_{B_k}^{P\infty}(ef)$, it is a BSVN fuzzy $B_k - I^P$ bridge if $I_{B'_k}^{P\infty}(ef) > I_{B_k}^{P\infty}(ef)$ and it is a BSVN fuzzy $B_k - F^P$ bridge if $F_{B'_k}^{P\infty}(ef) > F_{B_k}^{P\infty}(ef)$. Moreover, bd is a BSVN fuzzy $B_k - T^N$ bridge if $T_{B'_k}^{N\infty}(ef) < T_{B_k}^{N\infty}(ef)$, it is a BSVN fuzzy $B_k - I^N$ bridge if $I_{B'_k}^{N\infty}(ef) < I_{B_k}^{N\infty}(ef)$ and it is a BSVN fuzzy $B_k - F^N$ bridge if $F_{B'_k}^{N\infty}(ef) < F_{B_k}^{N\infty}(ef)$.

Example 7. Consider a BSVNGS $\check{G}_{bn} = (B, B_1, B_2)$ as depicted in Figure 6 and $\check{G}'_{bs} = (B', B'_1, B'_2)$, a BSVN spanning subgraph structure of the BSVNGS \check{G}_{bn} obtained by deleting B_1 -edge (b_2b_5) and that is shown in Figure 7.




Figure 7. A BSVNGS $\check{G}'_{bs} = (B', B'_1, B'_2)$.

This edge (b_2b_5) is a BSVN fuzzy B_1 -bridge, as $T_{B'_1}^{P\infty}(b_2b_5) = 0.3$, $T_{B_1}^{P\infty}(b_2b_5) = 0.4$, $I_{B'_1}^{P\infty}(b_2b_5) = 0.3$, $I_{B_1}^{P\infty}(b_2b_5) = 0.4$, $F_{B'_1}^{P\infty}(b_2b_5) = 0.4$, $F_{B_1}^{P\infty}(b_2b_5) = 0.5$, $T_{B'_1}^{N\infty}(b_2b_5) = -0.3 > -0.4 = T_{B_1}^{N\infty}(b_2b_5)$, $I_{B'_1}^{N\infty}(b_2b_5) = -0.3 > -0.4 = I_{B_1}^{N\infty}(b_2b_5)$, and $F_{B'_1}^{N\infty}(b_2b_5) = -0.4 > -0.5 = F_{B_1}^{N\infty}(b_2b_5)$.

Definition 13. A BSVNGS $\check{G}_{bn} = (B, B_1, B_2, \dots, B_m)$ is a B_k -tree if $(supp(B), supp(B_1), supp(B_2), \dots, supp(B_m))$ is a B_k -tree. Alternatively, \check{G}_{bn} is a B_k -tree if \check{G}_{bn} has a subgraph induced by $supp(B_k)$ that forms a tree.

Example 8. Consider the BSVNGS $\check{G}_{bn} = (B, B_1, B_2)$ as depicted in Figure 8.




Figure 8. A BSVN B_2 -tree.

This BSVNGS $\check{G}_{bn} = (B, B_1, B_2)$ is a B_2 -tree, as $(supp(B), supp(B_1), supp(B_2))$ is a B_2 -tree.

Definition 14. A BSVNGS $\check{G}_{bn} = (B, B_1, B_2, \dots, B_m)$ is a BSVN fuzzy B_k -tree if \check{G}_{bn} has a BSVN fuzzy spanning subgraph structure $\check{H}_{bn} = (B', B'_1, B'_2, \dots, B'_m)$ such that for all B_k -edges, bd not in \check{H}_{bn} :

1. \check{H}_{bn} is a B'_k -tree.

2. $T_{B_k}^P(bd) < T_{B'_k}^{P\infty}(bd)$, $I_{B_k}^P(bd) < I_{B'_k}^{P\infty}(bd)$, $F_{B_k}^P(bd) < F_{B'_k}^{P\infty}(bd)$, $T_{B_k}^N(bd) > T_{B'_k}^{N\infty}(bd)$, $I_{B_k}^N(bd) > I_{B'_k}^{N\infty}(bd)$, and $F_{B_k}^N(bd) > F_{B'_k}^{N\infty}(bd)$.

In particular, \check{G}_{bn} is a BSVN fuzzy $B_k - T^P$ tree if $T_{B_k}^P(bd) < T_{B'_k}^{P\infty}(bd)$, it is a BSVN fuzzy $B_k - I^P$ tree if $I_{B_k}^P(bd) < I_{B'_k}^{P\infty}(bd)$, and it is a BSVN fuzzy $B_k - F^P$ tree if $F_{B_k}^P(bd) > F_{B'_k}^{P\infty}(bd)$. Moreover, \check{G}_{bn} is a BSVN fuzzy $B_k - T^N$ tree if $T_{B_k}^N(bd) > T_{B'_k}^{N\infty}(bd)$, it is a BSVN fuzzy $B_k - I^N$ tree if $I_{B_k}^N(bd) > I_{B'_k}^{N\infty}(bd)$, and it is a BSVN fuzzy $B_k - F^N$ tree if $F_{B_k}^N(bd) < F_{B'_k}^{N\infty}(bd)$.

Example 9. Consider the BSVNGS $\check{G}_{bn} = (B, B_1, B_2)$ as depicted in Figure 9.




Figure 9. A BSVN fuzzy B_1 -tree.

It is B_2 -tree, rather than a B_1 -tree. However, it is a BSVN fuzzy B_1 -tree, because it has a BSVN fuzzy spanning subgraph (B', B'_1, B'_2) as a B'_1 -tree, which is obtained through the deletion of the B_1 -edge b_2b_5 from \check{G}_{bn} . Moreover, $T_{B'_1}^{P\infty}(b_2b_5) = 0.3$, $T_{B'_1}^P(b_2b_5) = 0.2$, $I_{B'_1}^{P\infty}(b_2b_5) = 0.3$, $I_{B'_1}^P(b_2b_5) = 0.1$, $F_{B'_1}^{P\infty}(b_2b_5) = 0.4$, $F_{B'_1}(b_2b_5) = 0.5$. $T_{B'_1}^{N\infty}(b_2b_5) = -0.3 < -0.2 = T_{B'_1}^N(b_2b_5)$, $I_{B'_1}^{N\infty}(b_2b_5) = -0.3 < -0.1 = I_{B'_1}^N(b_2b_5)$ and $F_{B'_1}^{N\infty}(b_2b_5) = -0.4 > -0.5 = F_{B'_1}(b_2b_5)$.

Now we define the operations on BSVNGSs.

Definition 15. Let $\check{G}_{b1} = (B_1, B_{11}, B_{12}, \dots, B_{1m})$ and $\check{G}_{b2} = (B_2, B_{21}, B_{22}, \dots, B_{2m})$ be two BSVNGSs. The Cartesian product of \check{G}_{b1} and \check{G}_{b2} , denoted by

$$\check{G}_{b1} \times \check{G}_{b2} = (B_1 \times B_2, B_{11} \times B_{21}, B_{12} \times B_{22}, \dots, B_{1m} \times B_{2m})$$

is defined as

$$(i) \quad \begin{cases} T_{(B_1 \times B_2)}^P(bd) = (T_{B_1}^P \times T_{B_2}^P)(bd) = T_{B_1}^P(b) \wedge T_{B_2}^P(d) \\ I_{(B_1 \times B_2)}^P(bd) = (I_{B_1}^P \times I_{B_2}^P)(bd) = I_{B_1}^P(b) \wedge I_{B_2}^P(d) \\ F_{(B_1 \times B_2)}^P(bd) = (F_{B_1}^P \times F_{B_2}^P)(bd) = F_{B_1}^P(b) \vee F_{B_2}^P(d) \end{cases}$$

$$\begin{aligned}
 \text{(ii)} \quad & \begin{cases} T_{(B_1 \times B_2)}^N(bd) = (T_{B_1}^N \times T_{B_2}^N)(bd) = T_{B_1}^N(b) \vee T_{B_2}^N(d) \\ I_{(B_1 \times B_2)}^N(bd) = (I_{B_1}^N \times I_{B_2}^N)(bd) = I_{B_1}^N(b) \vee I_{B_2}^N(d) \\ F_{(B_1 \times B_2)}^N(bd) = (F_{B_1}^N \times F_{B_2}^N)(bd) = F_{B_1}^N(b) \wedge F_{B_2}^N(d) \end{cases} \\
 & \text{for all } (bd) \in V_1 \times V_2, \\
 \text{(iii)} \quad & \begin{cases} T_{(B_{1k} \times B_{2k})}^P(bd_1)(bd_2) = (T_{B_{1k}}^P \times T_{B_{2k}}^P)(bd_1)(bd_2) = T_{B_{1k}}^P(b) \wedge T_{B_{2k}}^P(d_1d_2) \\ I_{(B_{1k} \times B_{2k})}^P(bd_1)(bd_2) = (I_{B_{1k}}^P \times I_{B_{2k}}^P)(bd_1)(bd_2) = I_{B_{1k}}^P(b) \wedge I_{B_{2k}}^P(d_1d_2) \\ F_{(B_{1k} \times B_{2k})}^P(bd_1)(bd_2) = (F_{B_{1k}}^P \times F_{B_{2k}}^P)(bd_1)(bd_2) = F_{B_{1k}}^P(b) \vee F_{B_{2k}}^P(d_1d_2) \end{cases} \\
 \text{(iv)} \quad & \begin{cases} T_{(B_{1k} \times B_{2k})}^N(bd_1)(bd_2) = (T_{B_{1k}}^N \times T_{B_{2k}}^N)(bd_1)(bd_2) = T_{B_{1k}}^N(b) \vee T_{B_{2k}}^N(d_1d_2) \\ I_{(B_{1k} \times B_{2k})}^N(bd_1)(bd_2) = (I_{B_{1k}}^N \times I_{B_{2k}}^N)(bd_1)(bd_2) = I_{B_{1k}}^N(b) \vee I_{B_{2k}}^N(d_1d_2) \\ F_{(B_{1k} \times B_{2k})}^N(bd_1)(bd_2) = (F_{B_{1k}}^N \times F_{B_{2k}}^N)(bd_1)(bd_2) = F_{B_{1k}}^N(b) \wedge F_{B_{2k}}^N(d_1d_2) \end{cases} \\
 & \text{for all } b \in V_1, (d_1d_2) \in V_{2k}, \text{ and} \\
 \text{(v)} \quad & \begin{cases} T_{(B_{1k} \times B_{2k})}^P(b_1d)(b_2d) = (T_{B_{1k}}^P \times T_{B_{2k}}^P)(b_1d)(b_2d) = T_{B_{2k}}^P(d) \wedge T_{B_{1k}}^P(b_1b_2) \\ I_{(B_{1k} \times B_{2k})}^P(b_1d)(b_2d) = (I_{B_{1k}}^P \times I_{B_{2k}}^P)(b_1d)(b_2d) = I_{B_{2k}}^P(d) \wedge I_{B_{1k}}^P(b_1b_2) \\ F_{(B_{1k} \times B_{2k})}^P(b_1d)(b_2d) = (F_{B_{1k}}^P \times F_{B_{2k}}^P)(b_1d)(b_2d) = F_{B_{2k}}^P(d) \vee F_{B_{1k}}^P(b_1b_2) \end{cases} \\
 \text{(vi)} \quad & \begin{cases} T_{(B_{1k} \times B_{2k})}^N(b_1d)(b_2d) = (T_{B_{1k}}^N \times T_{B_{2k}}^N)(b_1d)(b_2d) = T_{B_{2k}}^N(d) \vee T_{B_{1k}}^N(b_1b_2) \\ I_{(B_{1k} \times B_{2k})}^N(b_1d)(b_2d) = (I_{B_{1k}}^N \times I_{B_{2k}}^N)(b_1d)(b_2d) = I_{B_{2k}}^N(d) \vee I_{B_{1k}}^N(b_1b_2) \\ F_{(B_{1k} \times B_{2k})}^N(b_1d)(b_2d) = (F_{B_{1k}}^N \times F_{B_{2k}}^N)(b_1d)(b_2d) = F_{B_{2k}}^N(d) \wedge F_{B_{1k}}^N(b_1b_2) \end{cases} \\
 & \text{for all } d \in V_2, (b_1b_2) \in V_{1k}.
 \end{aligned}$$

Example 10. Consider $\check{G}_{b_1} = (B_1, B_{11}, B_{12})$ and $\check{G}_{b_2} = (B_2, B_{21}, B_{22})$ as BSVNGSs of GSs $\check{G}_{s_1} = (V_1, V_{11}, V_{12})$ and $\check{G}_{s_2} = (V_2, V_{21}, V_{22})$, respectively, as depicted in Figure 10, where $V_{11} = \{b_1b_2\}$, $V_{12} = \{b_3b_4\}$, $V_{21} = \{d_1d_2\}$, and $V_{22} = \{d_2d_3\}$.




Figure 10. Two BSVNGSs \check{G}_{b_1} and \check{G}_{b_2} .

The Cartesian product of \check{G}_{b_1} and \check{G}_{b_2} , defined as $\check{G}_{b_1} \times \check{G}_{b_2} = \{B_1 \times B_2, B_{11} \times B_{21}, B_{12} \times B_{22}\}$, is depicted in Figures 11 and 12.




Figure 11. $\check{G}_{b1} \times \check{G}_{b2}$.




Figure 12. $\check{G}_{b1} \times \check{G}_{b2}$.

Theorem 1. The Cartesian product $\check{G}_{b1} \times \check{G}_{b2} = (B_1 \times B_2, B_{11} \times B_{21}, B_{12} \times B_{22}, \dots, B_{1m} \times B_{2m})$ of two BSVNGSs of GSs \check{G}_{s1} and \check{G}_{s2} is a BSVNGS of $\check{G}_{s1} \times \check{G}_{s2}$.

Proof. Consider two cases:

Case 1. For $b \in V_1, d_1d_2 \in V_{2k}$,

$$\begin{aligned} T_{(B_{1k} \times B_{2k})}^P((bd_1)(bd_2)) &= T_{B_1}^P(b) \wedge T_{B_{2k}}^P(d_1d_2) \\ &\leq T_{B_1}^P(b) \wedge [T_{B_2}^P(d_1) \wedge T_{B_2}^P(d_2)] \\ &= [T_{B_1}^P(b) \wedge T_{B_2}^P(d_1)] \wedge [T_{B_1}^P(b) \wedge T_{B_2}^P(d_2)] \\ &= T_{(B_1 \times B_2)}^P(bd_1) \wedge T_{(B_1 \times B_2)}^P(bd_2) \end{aligned}$$

$$\begin{aligned} T_{(B_{1k} \times B_{2k})}^N((bd_1)(bd_2)) &= T_{B_1}^N(b) \vee T_{B_{2k}}^N(d_1d_2) \\ &\geq T_{B_1}^N(b) \vee [T_{B_2}^N(d_1) \vee T_{B_2}^N(d_2)] \\ &= [T_{B_1}^N(b) \vee T_{B_2}^N(d_1)] \vee [T_{B_1}^N(b) \vee T_{B_2}^N(d_2)] \\ &= T_{(B_1 \times B_2)}^N(bd_1) \vee T_{(B_1 \times B_2)}^N(bd_2) \end{aligned}$$

$$\begin{aligned}
 I_{(B_{1k} \times B_{2k})}^P((bd_1)(bd_2)) &= I_{B_1}^P(b) \wedge I_{B_{2k}}^P(d_1d_2) \\
 &\leq I_{B_1}^P(b) \wedge [I_{B_2}^P(d_1) \wedge I_{B_2}^P(d_2)] \\
 &= [I_{B_1}^P(b) \wedge I_{B_2}^P(d_1)] \wedge [I_{B_1}^P(b) \wedge I_{B_2}^P(d_2)] \\
 &= I_{(B_1 \times B_2)}^P(bd_1) \wedge I_{(B_1 \times B_2)}^P(bd_2)
 \end{aligned}$$

$$\begin{aligned}
 I_{(B_{1k} \times B_{2k})}^N((bd_1)(bd_2)) &= I_{B_1}^N(b) \vee I_{B_{2k}}^N(d_1d_2) \\
 &\geq I_{B_1}^N(b) \vee [I_{B_2}^N(d_1) \vee I_{B_2}^N(d_2)] \\
 &= [I_{B_1}^N(b) \vee I_{B_2}^N(d_1)] \vee [I_{B_1}^N(b) \vee I_{B_2}^N(d_2)] \\
 &= I_{(B_1 \times B_2)}^N(bd_1) \vee I_{(B_1 \times B_2)}^N(bd_2)
 \end{aligned}$$

$$\begin{aligned}
 F_{(B_{1k} \times B_{2k})}^P((bd_1)(bd_2)) &= F_{B_1}^P(b) \vee F_{B_{2k}}^P(d_1d_2) \\
 &\leq F_{B_1}^P(b) \vee [F_{B_2}^P(d_1) \vee F_{B_2}^P(d_2)] \\
 &= [F_{B_1}^P(b) \vee F_{B_2}^P(d_1)] \vee [F_{B_1}^P(b) \vee F_{B_2}^P(d_2)] \\
 &= F_{(B_1 \times B_2)}^P(bd_1) \vee F_{(B_1 \times B_2)}^P(bd_2)
 \end{aligned}$$

$$\begin{aligned}
 F_{(B_{1k} \times B_{2k})}^N((bd_1)(bd_2)) &= F_{B_1}^N(b) \wedge F_{B_{2k}}^N(d_1d_2) \\
 &\geq F_{B_1}^N(b) \wedge [F_{B_2}^N(d_1) \wedge F_{B_2}^N(d_2)] \\
 &= [F_{B_1}^N(b) \wedge F_{B_2}^N(d_1)] \wedge [F_{B_1}^N(b) \wedge F_{B_2}^N(d_2)] \\
 &= F_{(B_1 \times B_2)}^N(bd_1) \wedge F_{(B_1 \times B_2)}^N(bd_2)
 \end{aligned}$$

for $bd_1, bd_2 \in V_1 \times V_2$.

Case 2. For $b \in V_2, d_1d_2 \in V_{1k}$,

$$\begin{aligned}
 T_{(B_{1k} \times B_{2k})}^P((d_1b)(d_2b)) &= T_{B_2}^P(b) \wedge T_{B_{1k}}^P(d_1d_2) \\
 &\leq T_{B_2}^P(b) \wedge [T_{B_1}^P(d_1) \wedge T_{B_1}^P(d_2)] \\
 &= [T_{B_2}^P(b) \wedge T_{B_1}^P(d_1)] \wedge [T_{B_2}^P(b) \wedge T_{B_1}^P(d_2)] \\
 &= T_{(B_1 \times B_2)}^P(d_1b) \wedge T_{(B_1 \times B_2)}^P(d_2b)
 \end{aligned}$$

$$\begin{aligned}
 T_{(B_{1k} \times B_{2k})}^N((d_1b)(d_2b)) &= T_{B_2}^N(b) \vee T_{B_{1k}}^N(d_1d_2) \\
 &\geq T_{B_2}^N(b) \vee [T_{B_1}^N(d_1) \vee T_{B_1}^N(d_2)] \\
 &= [T_{B_2}^N(b) \vee T_{B_1}^N(d_1)] \vee [T_{B_2}^N(b) \vee T_{B_1}^N(d_2)] \\
 &= T_{(B_1 \times B_2)}^N(d_1b) \vee T_{(B_1 \times B_2)}^N(d_2b)
 \end{aligned}$$

$$\begin{aligned}
 I_{(B_{1k} \times B_{2k})}^P((d_1b)(d_2b)) &= I_{B_2}^P(b) \wedge I_{B_{1k}}^P(d_1d_2) \\
 &\leq I_{B_2}^P(b) \wedge [I_{B_1}^P(d_1) \wedge I_{B_1}^P(d_2)] \\
 &= [I_{B_2}^P(b) \wedge I_{B_1}^P(d_1)] \wedge [I_{B_2}^P(b) \wedge I_{B_1}^P(d_2)] \\
 &= I_{(B_1 \times B_2)}^P(d_1b) \wedge I_{(B_1 \times B_2)}^P(d_2b)
 \end{aligned}$$

$$\begin{aligned}
 I_{(B_{1k} \times B_{2k})}^N((d_1b)(d_2b)) &= I_{B_2}^N(b) \vee I_{B_{1k}}^N(d_1d_2) \\
 &\geq I_{B_2}^N(b) \vee [I_{B_1}^N(d_1) \vee I_{B_1}^N(d_2)] \\
 &= [I_{B_2}^N(b) \vee I_{B_1}^N(d_1)] \vee [I_{B_2}^N(b) \vee I_{B_1}^N(d_2)] \\
 &= I_{(B_1 \times B_2)}^N(d_1b) \vee I_{(B_1 \times B_2)}^N(d_2b)
 \end{aligned}$$

$$\begin{aligned}
 F_{(B_{1k} \times B_{2k})}^P((d_1b)(d_2b)) &= F_{B_2}^P(b) \vee F_{B_{1k}}^P(d_1d_2) \\
 &\leq F_{B_2}^P(b) \vee [F_{B_1}^P(d_1) \vee F_{B_1}^P(d_2)] \\
 &= [F_{B_2}^P(b) \vee F_{B_1}^P(d_1)] \vee [F_{B_2}^P(b) \vee F_{B_1}^P(d_2)] \\
 &= F_{(B_1 \times B_2)}^P(d_1b) \vee F_{(B_1 \times B_2)}^P(d_2b)
 \end{aligned}$$

$$\begin{aligned}
 F_{(B_{1k} \times B_{2k})}^N((d_1b)(d_2b)) &= F_{B_2}^N(b) \wedge F_{B_{1k}}^N(d_1d_2) \\
 &\geq F_{B_2}^N(b) \wedge [F_{B_1}^N(d_1) \wedge F_{B_1}^N(d_2)] \\
 &= [F_{B_2}^N(b) \wedge F_{B_1}^N(d_1)] \wedge [F_{B_2}^N(b) \wedge F_{B_1}^N(d_2)] \\
 &= F_{(B_1 \times B_2)}^N(d_1b) \wedge F_{(B_1 \times B_2)}^N(d_2b)
 \end{aligned}$$

for $d_1b, d_2b \in V_1 \times V_2$.

Both cases hold for all $k \in \{1, 2, \dots, m\}$. This completes the proof. \square

Definition 16. Let $\check{G}_{b_1} = (B_1, B_{11}, B_{12}, \dots, B_{1m})$ and $\check{G}_{b_2} = (B_2, B_{21}, B_{22}, \dots, B_{2m})$ be two BSVNGSs. The cross product of \check{G}_{b_1} and \check{G}_{b_2} , denoted by

$$\check{G}_{b_1} * \check{G}_{b_2} = (B_1 * B_2, B_{11} * B_{21}, B_{12} * B_{22}, \dots, B_{1m} * B_{2m})$$

is defined as

$$\begin{aligned}
 \text{(i)} \quad &\left\{ \begin{aligned} T_{(B_1 * B_2)}^P(bd) &= (T_{B_1}^P * T_{B_2}^P)(bd) = T_{B_1}^P(b) \wedge T_{B_2}^P(d) \\ I_{(B_1 * B_2)}^P(bd) &= (I_{B_1}^P * I_{B_2}^P)(bd) = I_{B_1}^P(b) \wedge I_{B_2}^P(d) \\ F_{(B_1 * B_2)}^P(bd) &= (F_{B_1}^P * F_{B_2}^P)(bd) = F_{B_1}^P(b) \vee F_{B_2}^P(d) \end{aligned} \right. \\
 \text{(ii)} \quad &\left\{ \begin{aligned} T_{(B_1 * B_2)}^N(bd) &= (T_{B_1}^N * T_{B_2}^N)(bd) = T_{B_1}^N(b) \vee T_{B_2}^N(d) \\ I_{(B_1 * B_2)}^N(bd) &= (I_{B_1}^N * I_{B_2}^N)(bd) = I_{B_1}^N(b) \vee I_{B_2}^N(d) \\ F_{(B_1 * B_2)}^N(bd) &= (F_{B_1}^N * F_{B_2}^N)(bd) = F_{B_1}^N(b) \wedge F_{B_2}^N(d) \end{aligned} \right.
 \end{aligned}$$

for all $(bd) \in V_1 \times V_2$, and

$$\begin{aligned}
 \text{(iii)} \quad &\left\{ \begin{aligned} T_{(B_{1k} * B_{2k})}^P(b_1d_1)(b_2d_2) &= (T_{B_{1k}}^P * T_{B_{2k}}^P)(b_1d_1)(b_2d_2) = T_{B_{1k}}^P(b_1b_2) \wedge T_{B_{2k}}^P(d_1d_2) \\ I_{(B_{1k} * B_{2k})}^P(b_1d_1)(b_2d_2) &= (I_{B_{1k}}^P * I_{B_{2k}}^P)(b_1d_1)(b_2d_2) = I_{B_{1k}}^P(b_1b_2) \wedge I_{B_{2k}}^P(d_1d_2) \\ F_{(B_{1k} * B_{2k})}^P(b_1d_1)(b_2d_2) &= (F_{B_{1k}}^P * F_{B_{2k}}^P)(b_1d_1)(b_2d_2) = F_{B_{1k}}^P(b_1b_2) \vee F_{B_{2k}}^P(d_1d_2) \end{aligned} \right. \\
 \text{(iv)} \quad &\left\{ \begin{aligned} T_{(B_{1k} * B_{2k})}^N(b_1d_1)(b_2d_2) &= (T_{B_{1k}}^N * T_{B_{2k}}^N)(b_1d_1)(b_2d_2) = T_{B_{1k}}^N(b_1b_2) \vee T_{B_{2k}}^N(d_1d_2) \\ I_{(B_{1k} * B_{2k})}^N(b_1d_1)(b_2d_2) &= (I_{B_{1k}}^N * I_{B_{2k}}^N)(b_1d_1)(b_2d_2) = I_{B_{1k}}^N(b_1b_2) \vee I_{B_{2k}}^N(d_1d_2) \\ F_{(B_{1k} * B_{2k})}^N(b_1d_1)(b_2d_2) &= (F_{B_{1k}}^N * F_{B_{2k}}^N)(b_1d_1)(b_2d_2) = F_{B_{1k}}^N(b_1b_2) \wedge F_{B_{2k}}^N(d_1d_2) \end{aligned} \right.
 \end{aligned}$$

for all $(b_1b_2) \in V_{1k}, (d_1d_2) \in V_{2k}$.

Example 11. The cross product of BSVNGSs \check{G}_{b_1} and \check{G}_{b_2} shown in Figure 10 is defined as $\check{G}_{b_1} * \check{G}_{b_2} = \{B_1 * B_2, B_{11} * B_{21}, B_{12} * B_{22}\}$ and is depicted in Figure 13.




Figure 13. $\check{G}_{b_1} \times \check{G}_{b_2}$.

Theorem 2. The cross product $\check{G}_{b_1} * \check{G}_{b_2} = (B_1 * B_2, B_{11} * B_{21}, B_{12} * B_{22}, \dots, B_{1m} * B_{2m})$ of two BSVNSGSs of GSs \check{G}_{s_1} and \check{G}_{s_2} is a BSVNSGS of $\check{G}_{s_1} * \check{G}_{s_2}$.

Proof. For $b_1 b_2 \in V_{1k}, d_1 d_2 \in V_{2k}$,

$$\begin{aligned} T_{(B_{1k} * B_{2k})}^P((b_1 d_1)(b_2 d_2)) &= T_{B_{1k}}^P(b_1 b_2) \wedge T_{B_{2k}}^P(d_1 d_2) \\ &\leq [T_{B_1}^P(b_1) \wedge T_{B_1}^P(b_2) \wedge [T_{B_2}^P(d_1) \wedge T_{B_2}^P(d_2)]] \\ &= [T_{B_1}^P(b_1) \wedge T_{B_2}^P(d_1)] \wedge [T_{B_1}^P(b_2) \wedge T_{B_2}^P(d_2)] \\ &= T_{(B_1 * B_2)}^P(b_1 d_1) \wedge T_{(B_1 * B_2)}^P(b_2 d_2) \end{aligned}$$

$$\begin{aligned} T_{(B_{1k} * B_{2k})}^N((b_1 d_1)(b_2 d_2)) &= T_{B_{1k}}^N(b_1 b_2) \vee T_{B_{2k}}^N(d_1 d_2) \\ &\geq [T_{B_1}^N(b_1) \vee T_{B_1}^N(b_2) \vee [T_{B_2}^N(d_1) \vee T_{B_2}^N(d_2)]] \\ &= [T_{B_1}^N(b_1) \vee T_{B_2}^N(d_1)] \vee [T_{B_1}^N(b_2) \vee T_{B_2}^N(d_2)] \\ &= T_{(B_1 * B_2)}^N(b_1 d_1) \vee T_{(B_1 * B_2)}^N(b_2 d_2) \end{aligned}$$

$$\begin{aligned} I_{(B_{1k} * B_{2k})}^P((b_1 d_1)(b_2 d_2)) &= I_{B_{1k}}^P(b_1 b_2) \wedge I_{B_{2k}}^P(d_1 d_2) \\ &\leq [I_{B_1}^P(b_1) \wedge I_{B_1}^P(b_2) \wedge [I_{B_2}^P(d_1) \wedge I_{B_2}^P(d_2)]] \\ &= [I_{B_1}^P(b_1) \wedge I_{B_2}^P(d_1)] \wedge [I_{B_1}^P(b_2) \wedge I_{B_2}^P(d_2)] \\ &= I_{(B_1 * B_2)}^P(b_1 d_1) \wedge I_{(B_1 * B_2)}^P(b_2 d_2) \end{aligned}$$

$$\begin{aligned} I_{(B_{1k} * B_{2k})}^N((b_1 d_1)(b_2 d_2)) &= I_{B_{1k}}^N(b_1 b_2) \vee I_{B_{2k}}^N(d_1 d_2) \\ &\geq [I_{B_1}^N(b_1) \vee I_{B_1}^N(b_2) \vee [I_{B_2}^N(d_1) \vee I_{B_2}^N(d_2)]] \\ &= [I_{B_1}^N(b_1) \vee I_{B_2}^N(d_1)] \vee [I_{B_1}^N(b_2) \vee I_{B_2}^N(d_2)] \\ &= I_{(B_1 * B_2)}^N(b_1 d_1) \vee I_{(B_1 * B_2)}^N(b_2 d_2) \end{aligned}$$

$$\begin{aligned}
 F_{(B_{1k} * B_{2k})}^P((b_1 d_1)(b_2 d_2)) &= F_{B_{1k}}^P(b_1 b_2) \vee F_{B_{2k}}^P(d_1 d_2) \\
 &\leq [F_{B_1}^P(b_1) \vee F_{B_1}^P(b_2)] \vee [F_{B_2}^P(d_1) \vee F_{B_2}^P(d_2)] \\
 &= [F_{B_1}^P(b_1) \vee F_{B_2}^P(d_1)] \vee [F_{B_1}^P(b_2) \vee F_{B_2}^P(d_2)] \\
 &= F_{(B_1 * B_2)}^P(b_1 d_1) \vee F_{(B_1 * B_2)}^P(b_2 d_2)
 \end{aligned}$$

$$\begin{aligned}
 F_{(B_{1k} * B_{2k})}^N((b_1 d_1)(b_2 d_2)) &= F_{B_{1k}}^N(b_1 b_2) \wedge F_{B_{2k}}^N(d_1 d_2) \\
 &\geq [F_{B_1}^N(b_1) \wedge F_{B_1}^N(b_2)] \wedge [F_{B_2}^N(d_1) \wedge F_{B_2}^N(d_2)] \\
 &= [F_{B_1}^N(b_1) \wedge F_{B_2}^N(d_1)] \wedge [F_{B_1}^N(b_2) \wedge F_{B_2}^N(d_2)] \\
 &= F_{(B_1 * B_2)}^N(b_1 d_1) \wedge F_{(B_1 * B_2)}^N(b_2 d_2)
 \end{aligned}$$

where $b_1 d_1, b_2 d_2 \in V_1 * V_2$ and $h \in \{1, 2, \dots, m\}$. \square

Definition 17. Let $\check{G}_{b_1} = (B_1, B_{11}, B_{12}, \dots, B_{1m})$ and $\check{G}_{b_2} = (B_2, B_{21}, B_{22}, \dots, B_{2m})$ be two BSVNGSs. The composition of \check{G}_{b_1} and \check{G}_{b_2} , denoted by

$$\check{G}_{b_1} \circ \check{G}_{b_2} = (B_1 \circ B_2, B_{11} \circ B_{21}, B_{12} \circ B_{22}, \dots, B_{1m} \circ B_{2m})$$

is defined as

$$\begin{aligned}
 \text{(i)} \quad & \left\{ \begin{aligned} T_{(B_1 \circ B_2)}^P(bd) &= (T_{B_1}^P \circ T_{B_2}^P)(bd) = T_{B_1}^P(b) \wedge T_{B_2}^P(d) \\ I_{(B_1 \circ B_2)}^P(bd) &= (I_{B_1}^P \circ I_{B_2}^P)(bd) = I_{B_1}^P(b) \wedge I_{B_2}^P(d) \\ F_{(B_1 \circ B_2)}^P(bd) &= (F_{B_1}^P \circ F_{B_2}^P)(bd) = F_{B_1}^P(b) \vee F_{B_2}^P(d) \end{aligned} \right. \\
 \text{(ii)} \quad & \left\{ \begin{aligned} T_{(B_1 \circ B_2)}^N(bd) &= (T_{B_1}^N \circ T_{B_2}^N)(bd) = T_{B_1}^N(b) \vee T_{B_2}^N(d) \\ I_{(B_1 \circ B_2)}^N(bd) &= (I_{B_1}^N \circ I_{B_2}^N)(bd) = I_{B_1}^N(b) \vee I_{B_2}^N(d) \\ F_{(B_1 \circ B_2)}^N(bd) &= (F_{B_1}^N \circ F_{B_2}^N)(bd) = F_{B_1}^N(b) \wedge F_{B_2}^N(d) \end{aligned} \right.
 \end{aligned}$$

for all $(bd) \in V_1 \times V_2$,

$$\begin{aligned}
 \text{(iii)} \quad & \left\{ \begin{aligned} T_{(B_{1k} \circ B_{2k})}^P(bd_1)(bd_2) &= (T_{B_{1k}}^P \circ T_{B_{2k}}^P)(bd_1)(bd_2) = T_{B_1}^P(b) \wedge T_{B_{2k}}^P(d_1 d_2) \\ I_{(B_{1k} \circ B_{2k})}^P(bd_1)(bd_2) &= (I_{B_{1k}}^P \circ I_{B_{2k}}^P)(bd_1)(bd_2) = I_{B_1}^P(b) \wedge I_{B_{2k}}^P(d_1 d_2) \\ F_{(B_{1k} \circ B_{2k})}^P(bd_1)(bd_2) &= (F_{B_{1k}}^P \circ F_{B_{2k}}^P)(bd_1)(bd_2) = F_{B_1}^P(b) \vee F_{B_{2k}}^P(d_1 d_2) \end{aligned} \right. \\
 \text{(iv)} \quad & \left\{ \begin{aligned} T_{(B_{1k} \circ B_{2k})}^N(bd_1)(bd_2) &= (T_{B_{1k}}^N \circ T_{B_{2k}}^N)(bd_1)(bd_2) = T_{B_1}^N(b) \vee T_{B_{2k}}^N(d_1 d_2) \\ I_{(B_{1k} \circ B_{2k})}^N(bd_1)(bd_2) &= (I_{B_{1k}}^N \circ I_{B_{2k}}^N)(bd_1)(bd_2) = I_{B_1}^N(b) \vee I_{B_{2k}}^N(d_1 d_2) \\ F_{(B_{1k} \circ B_{2k})}^N(bd_1)(bd_2) &= (F_{B_{1k}}^N \circ F_{B_{2k}}^N)(bd_1)(bd_2) = F_{B_1}^N(b) \wedge F_{B_{2k}}^N(d_1 d_2) \end{aligned} \right.
 \end{aligned}$$

for all $b \in V_1, (d_1 d_2) \in V_{2k}$,

$$\begin{aligned}
 \text{(v)} \quad & \left\{ \begin{aligned} T_{(B_{1k} \circ B_{2k})}^P(b_1 d)(b_2 d) &= (T_{B_{1k}}^P \circ T_{B_{2k}}^P)(b_1 d)(b_2 d) = T_{B_2}^P(d) \wedge T_{B_{1k}}^P(b_1 b_2) \\ I_{(B_{1k} \circ B_{2k})}^P(b_1 d)(b_2 d) &= (I_{B_{1k}}^P \circ I_{B_{2k}}^P)(b_1 d)(b_2 d) = I_{B_2}^P(d) \wedge I_{B_{1k}}^P(b_1 b_2) \\ F_{(B_{1k} \circ B_{2k})}^P(b_1 d)(b_2 d) &= (F_{B_{1k}}^P \circ F_{B_{2k}}^P)(b_1 d)(b_2 d) = F_{B_2}^P(d) \vee F_{B_{1k}}^P(b_1 b_2) \end{aligned} \right. \\
 \text{(vi)} \quad & \left\{ \begin{aligned} T_{(B_{1k} \circ B_{2k})}^N(b_1 d)(b_2 d) &= (T_{B_{1k}}^N \circ T_{B_{2k}}^N)(b_1 d)(b_2 d) = T_{B_2}^N(d) \vee T_{B_{1k}}^N(b_1 b_2) \\ I_{(B_{1k} \circ B_{2k})}^N(b_1 d)(b_2 d) &= (I_{B_{1k}}^N \circ I_{B_{2k}}^N)(b_1 d)(b_2 d) = I_{B_2}^N(d) \vee I_{B_{1k}}^N(b_1 b_2) \\ F_{(B_{1k} \circ B_{2k})}^N(b_1 d)(b_2 d) &= (F_{B_{1k}}^N \circ F_{B_{2k}}^N)(b_1 d)(b_2 d) = F_{B_2}^N(d) \wedge F_{B_{1k}}^N(b_1 b_2) \end{aligned} \right.
 \end{aligned}$$

for all $d \in V_2, (b_1 b_2) \in V_{1k}$, and

$$\begin{aligned}
 \text{(vii)} \quad & \left\{ \begin{aligned} T_{(B_{1k} \circ B_{2k})}^P(b_1 d_1)(b_2 d_2) &= (T_{B_{1k}}^P \circ T_{B_{2k}}^P)(b_1 d_1)(b_2 d_2) = T_{B_{1k}}^P(b_1 b_2) \wedge T_{B_2}^P(d_1) \wedge T_{B_2}^P(d_2) \\ I_{(B_{1k} \circ B_{2k})}^P(b_1 d_1)(b_2 d_2) &= (I_{B_{1k}}^P \circ I_{B_{2k}}^P)(b_1 d_1)(b_2 d_2) = I_{B_{1k}}^P(b_1 b_2) \wedge I_{B_2}^P(d_1) \wedge I_{B_2}^P(d_2) \\ F_{(B_{1k} \circ B_{2k})}^P(b_1 d_1)(b_2 d_2) &= (F_{B_{1k}}^P \circ F_{B_{2k}}^P)(b_1 d_1)(b_2 d_2) = F_{B_{1k}}^P(b_1 b_2) \vee F_{B_2}^P(d_1) \vee F_{B_2}^P(d_2) \end{aligned} \right.
 \end{aligned}$$

$$(viii) \begin{cases} T_{(B_{1k} \circ B_{2k})}^N(b_1 d_1)(b_2 d_2) = (T_{B_{1k}}^N \circ T_{B_{2k}}^N)(b_1 d_1)(b_2 d_2) = T_{B_{1k}}^N(b_1 b_2) \vee T_{B_2}^N(d_1) \vee T_{B_2}^N(d_2) \\ I_{(B_{1k} \circ B_{2k})}^N(b_1 d_1)(b_2 d_2) = (I_{B_{1k}}^N \circ I_{B_{2k}}^N)(b_1 d_1)(b_2 d_2) = I_{B_{1k}}^N(b_1 b_2) \vee I_{B_2}^N(d_1) \vee I_{B_2}^N(d_2) \\ F_{(B_{1k} \circ B_{2k})}^N(b_1 d_1)(b_2 d_2) = (F_{B_{1k}}^N \circ F_{B_{2k}}^N)(b_1 d_1)(b_2 d_2) = F_{B_{1k}}^N(b_1 b_2) \wedge F_{B_2}^N(d_1) \wedge F_{B_2}^N(d_2) \end{cases}$$

for all $(b_1 b_2) \in V_{1k}, (d_1 d_2) \in V_{2k}$ such that $d_1 \neq d_2$.

Example 12. The composition of BSVNGSs \check{G}_{b1} and \check{G}_{b2} shown in Figure 10 is defined as $\check{G}_{b1} \circ \check{G}_{b2} = \{B_1 \circ B_2, B_{11} \circ B_{21}, B_{12} \circ B_{22}\}$ and is depicted in Figures 14 and 15.




Figure 14. $\check{G}_{b1} \circ \check{G}_{b2}$.




Figure 15. $\check{G}_{b1} \circ \check{G}_{b2}$.

Theorem 3. The composition $\check{G}_{b1} \circ \check{G}_{b2} = (B_1 \circ B_2, B_{11} \circ B_{21}, B_{12} \circ B_{22}, \dots, B_{1m} \circ B_{2m})$ of two BSVNGSs of GSs \check{G}_{s1} and \check{G}_{s2} is a BSVNGS of $\check{G}_{s1} \circ \check{G}_{s2}$.

Proof. Consider three cases:

Case 1. For $b \in V_1, d_1d_2 \in V_{2k}$,

$$\begin{aligned} T_{(B_{1k} \circ B_{2k})}^P((bd_1)(bd_2)) &= T_{B_1}^P(b) \wedge T_{B_{2k}}^P(d_1d_2) \\ &\leq T_{B_1}^P(b) \wedge [T_{B_2}^P(d_1) \wedge T_{B_2}^P(d_2)] \\ &= [T_{B_1}^P(b) \wedge T_{B_2}^P(d_1)] \wedge [T_{B_1}^P(b) \wedge T_{B_2}^P(d_2)] \\ &= T_{(B_1 \circ B_2)}^P(bd_1) \wedge T_{(B_1 \circ B_2)}^P(bd_2) \end{aligned}$$

$$\begin{aligned} T_{(B_{1k} \circ B_{2k})}^N((bd_1)(bd_2)) &= T_{B_1}^N(b) \vee T_{B_{2k}}^N(d_1d_2) \\ &\geq T_{B_1}^N(b) \vee [T_{B_2}^N(d_1) \vee T_{B_2}^N(d_2)] \\ &= [T_{B_1}^N(b) \vee T_{B_2}^N(d_1)] \vee [T_{B_1}^N(b) \vee T_{B_2}^N(d_2)] \\ &= T_{(B_1 \circ B_2)}^N(bd_1) \vee T_{(B_1 \circ B_2)}^N(bd_2) \end{aligned}$$

$$\begin{aligned} I_{(B_{1k} \circ B_{2k})}^P((bd_1)(bd_2)) &= I_{B_1}^P(b) \wedge I_{B_{2k}}^P(d_1d_2) \\ &\leq I_{B_1}^P(b) \wedge [I_{B_2}^P(d_1) \wedge I_{B_2}^P(d_2)] \\ &= [I_{B_1}^P(b) \wedge I_{B_2}^P(d_1)] \wedge [I_{B_1}^P(b) \wedge I_{B_2}^P(d_2)] \\ &= I_{(B_1 \circ B_2)}^P(bd_1) \wedge I_{(B_1 \circ B_2)}^P(bd_2) \end{aligned}$$

$$\begin{aligned} I_{(B_{1k} \circ B_{2k})}^N((bd_1)(bd_2)) &= I_{B_1}^N(b) \vee I_{B_{2k}}^N(d_1d_2) \\ &\geq I_{B_1}^N(b) \vee [I_{B_2}^N(d_1) \vee I_{B_2}^N(d_2)] \\ &= [I_{B_1}^N(b) \vee I_{B_2}^N(d_1)] \vee [I_{B_1}^N(b) \vee I_{B_2}^N(d_2)] \\ &= I_{(B_1 \circ B_2)}^N(bd_1) \vee I_{(B_1 \circ B_2)}^N(bd_2) \end{aligned}$$

$$\begin{aligned} F_{(B_{1k} \circ B_{2k})}^P((bd_1)(bd_2)) &= F_{B_1}^P(b) \vee F_{B_{2k}}^P(d_1d_2) \\ &\leq F_{B_1}^P(b) \vee [F_{B_2}^P(d_1) \vee F_{B_2}^P(d_2)] \\ &= [F_{B_1}^P(b) \vee F_{B_2}^P(d_1)] \vee [F_{B_1}^P(b) \vee F_{B_2}^P(d_2)] \\ &= F_{(B_1 \circ B_2)}^P(bd_1) \vee F_{(B_1 \circ B_2)}^P(bd_2) \end{aligned}$$

$$\begin{aligned} F_{(B_{1k} \circ B_{2k})}^N((bd_1)(bd_2)) &= F_{B_1}^N(b) \wedge F_{B_{2k}}^N(d_1d_2) \\ &\geq F_{B_1}^N(b) \wedge [F_{B_2}^N(d_1) \wedge F_{B_2}^N(d_2)] \\ &= [F_{B_1}^N(b) \wedge F_{B_2}^N(d_1)] \wedge [F_{B_1}^N(b) \wedge F_{B_2}^N(d_2)] \\ &= F_{(B_1 \circ B_2)}^N(bd_1) \wedge F_{(B_1 \circ B_2)}^N(bd_2) \end{aligned}$$

for $bd_1, bd_2 \in V_1 \circ V_2$.

Case 2. For $b \in V_2, d_1d_2 \in V_{1k}$,

$$\begin{aligned} T_{(B_{1k} \circ B_{2k})}^P((d_1b)(d_2b)) &= T_{B_2}^P(b) \wedge T_{B_{1k}}^P(d_1d_2) \\ &\leq T_{B_2}^P(b) \wedge [T_{B_1}^P(d_1) \wedge T_{B_1}^P(d_2)] \\ &= [T_{B_2}^P(b) \wedge T_{B_1}^P(d_1)] \wedge [T_{B_2}^P(b) \wedge T_{B_1}^P(d_2)] \\ &= T_{(B_1 \circ B_2)}^P(d_1b) \wedge T_{(B_1 \circ B_2)}^P(d_2b) \end{aligned}$$

$$\begin{aligned}
 T_{(B_{1k} \circ B_{2k})}^N((d_1b)(d_2b)) &= T_{B_2}^N(b) \vee T_{B_{1k}}^N(d_1d_2) \\
 &\geq T_{B_2}^N(b) \vee [T_{B_1}^N(d_1) \vee T_{B_1}^N(d_2)] \\
 &= [T_{B_2}^N(b) \vee T_{B_1}^N(d_1)] \vee [T_{B_2}^N(b) \vee T_{B_1}^N(d_2)] \\
 &= T_{(B_1 \circ B_2)}^N(d_1b) \vee T_{(B_1 \circ B_2)}^N(d_2b)
 \end{aligned}$$

$$\begin{aligned}
 I_{(B_{1k} \circ B_{2k})}^P((d_1b)(d_2b)) &= I_{B_2}^P(b) \wedge I_{B_{1k}}^P(d_1d_2) \\
 &\leq I_{B_2}^P(b) \wedge [I_{B_1}^P(d_1) \wedge I_{B_1}^P(d_2)] \\
 &= [I_{B_2}^P(b) \wedge I_{B_1}^P(d_1)] \wedge [I_{B_2}^P(b) \wedge I_{B_1}^P(d_2)] \\
 &= I_{(B_1 \circ B_2)}^P(d_1b) \wedge I_{(B_1 \circ B_2)}^P(d_2b)
 \end{aligned}$$

$$\begin{aligned}
 I_{(B_{1k} \circ B_{2k})}^N((d_1b)(d_2b)) &= I_{B_2}^N(b) \vee I_{B_{1k}}^N(d_1d_2) \\
 &\geq I_{B_2}^N(b) \vee [I_{B_1}^N(d_1) \vee I_{B_1}^N(d_2)] \\
 &= [I_{B_2}^N(b) \vee I_{B_1}^N(d_1)] \vee [I_{B_2}^N(b) \vee I_{B_1}^N(d_2)] \\
 &= I_{(B_1 \circ B_2)}^N(d_1b) \vee I_{(B_1 \circ B_2)}^N(d_2b)
 \end{aligned}$$

$$\begin{aligned}
 F_{(B_{1k} \circ B_{2k})}^P((d_1b)(d_2b)) &= F_{B_2}^P(b) \vee F_{B_{1k}}^P(d_1d_2) \\
 &\leq F_{B_2}^P(b) \vee [F_{B_1}^P(d_1) \vee F_{B_1}^P(d_2)] \\
 &= [F_{B_2}^P(b) \vee F_{B_1}^P(d_1)] \vee [F_{B_2}^P(b) \vee F_{B_1}^P(d_2)] \\
 &= F_{(B_1 \circ B_2)}^P(d_1b) \vee F_{(B_1 \circ B_2)}^P(d_2b)
 \end{aligned}$$

$$\begin{aligned}
 F_{(B_{1k} \circ B_{2k})}^N((d_1b)(d_2b)) &= F_{B_2}^N(b) \wedge F_{B_{1k}}^N(d_1d_2) \\
 &\geq F_{B_2}^N(b) \wedge [F_{B_1}^N(d_1) \wedge F_{B_1}^N(d_2)] \\
 &= [F_{B_2}^N(b) \wedge F_{B_1}^N(d_1)] \wedge [F_{B_2}^N(b) \wedge F_{B_1}^N(d_2)] \\
 &= F_{(B_1 \circ B_2)}^N(d_1b) \wedge F_{(B_1 \circ B_2)}^N(d_2b)
 \end{aligned}$$

for $d_1b, d_2b \in V_1 \circ V_2$.

Case 3. For $(b_1b_2) \in V_{1k}, (d_1d_2) \in V_{2k}$ such that $d_1 \neq d_2$,

$$\begin{aligned}
 T_{(B_{1k} \circ B_{2k})}^P((b_1d_1)(b_2d_2)) &= T_{B_{1k}}^P(b_1b_2) \wedge T_{B_2}^P(d_1) \wedge T_{B_2}^P(d_2) \\
 &\leq [T_{B_1}^P(b_1) \wedge T_{B_1}^P(b_2)] \wedge [T_{B_2}^P(d_1) \wedge T_{B_2}^P(d_2)] \\
 &= [T_{B_1}^P(b_1) \wedge T_{B_2}^P(d_1)] \wedge [T_{B_1}^P(b_2) \wedge T_{B_2}^P(d_2)] \\
 &= T_{(B_1 \circ B_2)}^P(b_1d_1) \wedge T_{(B_1 \circ B_2)}^P(b_2d_2)
 \end{aligned}$$

$$\begin{aligned}
 T_{(B_{1k} \circ B_{2k})}^N((b_1d_1)(b_2d_2)) &= T_{B_{1k}}^N(b_1b_2) \vee T_{B_2}^N(d_1) \vee T_{B_2}^N(d_2) \\
 &\geq [T_{B_1}^N(b_1) \vee T_{B_1}^N(b_2)] \vee [T_{B_2}^N(d_1) \vee T_{B_2}^N(d_2)] \\
 &= [T_{B_1}^N(b_1) \vee T_{B_2}^N(d_1)] \vee [T_{B_1}^N(b_2) \vee T_{B_2}^N(d_2)] \\
 &= T_{(B_1 \circ B_2)}^N(b_1d_1) \vee T_{(B_1 \circ B_2)}^N(b_2d_2)
 \end{aligned}$$

$$\begin{aligned}
 I_{(B_{1k} \circ B_{2k})}^P((b_1 d_1)(b_2 d_2)) &= I_{B_{1k}}^P(b_1 b_2) \wedge I_{B_2}^P(d_1) \wedge I_{B_2}^P(d_2) \\
 &\leq [I_{B_1}^P(b_1) \wedge I_{B_1}^P(b_2) \wedge [I_{B_2}^P(d_1) \wedge I_{B_2}^P(d_2)]] \\
 &= [I_{B_1}^P(b_1) \wedge I_{B_2}^P(d_1)] \wedge [I_{B_1}^P(b_2) \wedge I_{B_2}^P(d_2)] \\
 &= I_{(B_1 \circ B_2)}^P(b_1 d_1) \wedge I_{(B_1 \circ B_2)}^P(b_2 d_2)
 \end{aligned}$$

$$\begin{aligned}
 I_{(B_{1k} \circ B_{2k})}^N((b_1 d_1)(b_2 d_2)) &= I_{B_{1k}}^N(b_1 b_2) \vee I_{B_2}^N(d_1) \vee I_{B_2}^N(d_2) \\
 &\geq [I_{B_1}^N(b_1) \vee I_{B_1}^N(b_2) \vee [I_{B_2}^N(d_1) \vee I_{B_2}^N(d_2)]] \\
 &= [I_{B_1}^N(b_1) \vee I_{B_2}^N(d_1)] \vee [I_{B_1}^N(b_2) \vee I_{B_2}^N(d_2)] \\
 &= I_{(B_1 \circ B_2)}^N(b_1 d_1) \vee I_{(B_1 \circ B_2)}^N(b_2 d_2)
 \end{aligned}$$

$$\begin{aligned}
 F_{(B_{1k} \circ B_{2k})}^P((b_1 d_1)(b_2 d_2)) &= F_{B_{1k}}^P(b_1 b_2) \vee F_{B_2}^P(d_1) \vee F_{B_2}^P(d_2) \\
 &\leq [F_{B_1}^P(b_1) \vee F_{B_1}^P(b_2) \vee [F_{B_2}^P(d_1) \vee F_{B_2}^P(d_2)]] \\
 &= [F_{B_1}^P(b_1) \vee F_{B_2}^P(d_1)] \vee [F_{B_1}^P(b_2) \vee F_{B_2}^P(d_2)] \\
 &= F_{(B_1 \circ B_2)}^P(b_1 d_1) \vee F_{(B_1 \circ B_2)}^P(b_2 d_2)
 \end{aligned}$$

$$\begin{aligned}
 F_{(B_{1k} \circ B_{2k})}^N((b_1 d_1)(b_2 d_2)) &= F_{B_{1k}}^N(b_1 b_2) \wedge F_{B_2}^N(d_1) \wedge F_{B_2}^N(d_2) \\
 &\geq [F_{B_1}^N(b_1) \wedge F_{B_1}^N(b_2) \wedge [F_{B_2}^N(d_1) \wedge F_{B_2}^N(d_2)]] \\
 &= [F_{B_1}^N(b_1) \wedge F_{B_2}^N(d_1)] \wedge [F_{B_1}^N(b_2) \wedge F_{B_2}^N(d_2)] \\
 &= F_{(B_1 \circ B_2)}^N(b_1 d_1) \wedge F_{(B_1 \circ B_2)}^N(b_2 d_2)
 \end{aligned}$$

where $b_1 d_1, b_2 d_2 \in V_1 \circ V_2$.

All cases are satisfied for all $k \in \{1, 2, \dots, m\}$. \square

3. Conclusions

The notion of bipolar fuzzy graphs is applicable in several domains of engineering, expert systems, pattern recognition, signal processing, neural networks, medical diagnosis and decision-making. BSVNGSs show more flexibility, compatibility and precision for a system than single-valued neutrosophic graph structures. In this research paper, we introduced certain concepts of BSVNGSs and elaborated on them with suitable examples. Further, we defined some operations on BSVNGSs and investigated some relevant properties of these operations. We intend to generalize our research of fuzzification to (1) concepts of BSVN soft graph structures, (2) concepts of BSVN rough fuzzy graph structures, (3) concepts of BSVN fuzzy soft graph structures, and (4) concepts of BSVN rough fuzzy soft graph structures.

Acknowledgments: The authors are thankful to referees for their valuable comments and suggestion.

Author Contributions: Muhammad Akram, Muzzamal Sitara and Florentin Smarandache conceived and designed the experiments; Muhammad Akram performed the experiments; Muhammad Akram and Muzzamal Sitara analyzed the data; Florentin Smarandache contributed reagents/materials/analysis tools; Muzzamal Sitara wrote the paper.

Conflicts of Interest: The authors declare no conflict of interest regarding the publication of this research paper.

References

1. Zadeh, L.A. Fuzzy sets. *Inf. Control* **1965**, *8*, 338–353.
2. Atanassov, K. Intuitionistic fuzzy sets. *Fuzzy Sets Syst.* **1986**, *20*, 87–96.

3. Zhang, W.-R. Bipolar fuzzy sets and relations: A computational framework for cognitive modeling and multiagent decision analysis. In Proceedings of the Industrial Fuzzy Control and Intelligent Systems Conference, and the NASA Joint Technology Workshop on Neural Networks and Fuzzy Logic Fuzzy Information Processing Society Biannual Conference, San Antonio, TX, USA, 18–21 December 1994; pp. 305–309.
4. Smarandache, F. *Neutrosophy Neutrosophic Probability, Set, and Logic*; Amer Res Press: Rehoboth, MA, USA, 1998.
5. Wang, H.; Smarandache, F.; Zhang, Y.Q.; Sunderraman, R. Single valued neutrosophic sets. *Multispace Multistruct.* **2010**, *4*, 410–413.
6. Deli, I.; Ali, M.; Smarandache, F. Bipolar neutrosophic sets and their application based on multi-criteria decision making problems. In Proceedings of the International Conference IEEE Advanced Mechatronic Systems (ICAMechS), Beijing, China, 22–24 August 2015; pp. 249–254.
7. Zadeh, L.A. Similarity relations and fuzzy orderings. *Inf. Sci.* **1971**, *3*, 177–200.
8. Kauffman, A. *Introduction to the Theory of Fuzzy Subsets*; Academic Press: Cambridge, MA, USA, 1973; Volume 1.
9. Rosenfeld, A. Fuzzy graphs. In *Fuzzy Sets and Their Applications*; Zadeh, L.A., Fu, K.S., Eds.; Academic Press: New York, NY, USA, 1975; pp. 77–95.
10. Bhattacharya, P. Some remarks on fuzzy graphs. *Pattern Recognit. Lett.* **1987**, *6*, 297–302.
11. Akram, M. Bipolar fuzzy graphs. *Inf. Sci.* **2011**, *181*, 5548–5564.
12. Dinesh, T.; Ramakrishnan, T.V. On generalised fuzzy graph structures. *Appl. Math. Sci.* **2011**, *5*, 173–180.
13. Akram, M.; Akmal, R. Application of bipolar fuzzy sets in graph structures. *Appl. Comput. Intell. Soft Comput.* **2016**, *2016*, doi:10.1155/2016/5859080.
14. Akram, M.; Akmal, R. Certain operations on bipolar fuzzy graph structures. *Appl. Appl. Math.* **2016**, *11*, 463–488.
15. Akram, M.; Shahzadi, S. Representation of graphs using intuitionistic neutrosophic soft sets. *J. Math. Anal.* **2016**, *7*, 31–53.
16. Akram, M.; Shahzadi, G. Operations on single-valued neutrosophic graphs. *J. Uncertain Syst.* **2017**, *11*, 176–196.
17. Broumi, S.; Smarandache, F.; Talea, M.; Bakali, A. An introduction to bipolar single valued neutrosophic graph theory. *Appl. Mech. Mater.* **2016**, *841*, 184–191.
18. Dhavaseelan, R.; Vikramaprasad, R.; Krishnaraj, V. Certain types of neutrosophic graphs. *Int. J. Math. Sci. Appl.* **2015**, *5*, 333–339.
19. Dinesh, T. A study on graph structures Incidence Algebras and Their Fuzzy Analogues. Ph.D. Thesis, Kannur University, Kannur, India, 2011.
20. Ye, J. Single-valued neutrosophic minimum spanning tree and its clustering method. *J. Intell. Syst.* **2014**, *23*, 311–324.
21. Ye, J. Improved correlation coefficients of single-valued neutrosophic sets and interval neutrosophic sets for multiple attribute decision making. *J. Intell. Fuzzy Syst.* **2014**, *27*, 2453–2462.
22. Ye, J. A multicriteria decision-making method using aggregation operators for simplified neutrosophic sets. *J. Intell. Fuzzy Syst.* **2014**, *26*, 2459–2466.
23. Akram, M.; Sarwar, M. Novel multiple criteria decision making methods based on bipolar neutrosophic sets and bipolar neutrosophic graphs. *Ital. J. Pure Appl. Math.* **2017**, *38*, 368–389.
24. Majumdar, P.; Samanta, S.K. On similarity and entropy of neutrosophic sets. *J. Intell. Fuzzy Syst.* **2014**, *26*, 1245–1252.
25. Mordeson, J.N.; Chang-Shyh, P. Operations on fuzzy graphs. *Inf. Sci.* **1994**, *79*, 159–170.
26. Mordeson, J.N.; Nair, P.S. *Fuzzy Graphs and Fuzzy Hypergraphs*; Physica Verlag: Heidelberg, Germany, 1998; Second Edition 2001.
27. Parvathi, R.; Karunambigai, M.G.; Atanassov, K.T. Operations on intuitionistic fuzzy graphs. In Proceedings of the IEEE International Conference on Fuzzy Systems, Jeju Island, Korea, 20–24 August 2009; pp. 1396–1401.
28. Peng, J.J.; Wang, J.Q.; Zhang, H.Y.; Chen, X.H. An outranking approach for multi-criteria decision-making problems with simplified neutrosophic sets. *Appl. Soft Comput.* **2014**, *25*, 336–346.
29. Sampathkumar, E. Generalized graph structures. *Bull. Kerala Math. Assoc.* **2006**, *3*, 65–123.
30. Akram, M.; Sitara, M. Bipolar neutrosophic graph structures. *J. Indones. Math. Soc.* **2017**, *23*, 55–76.

