

Gravity and the crossed product

Edward Witten

*School of Natural Sciences, Institute for Advanced Study,
1 Einstein Drive, Princeton, NJ 08540 U.S.A.*

E-mail: witten@ias.edu

ABSTRACT: Recently Leutheusser and Liu [1, 2] identified an emergent algebra of Type III_1 in the operator algebra of $\mathcal{N} = 4$ super Yang-Mills theory for large N . Here we describe some $1/N$ corrections to this picture and show that the emergent Type III_1 algebra becomes an algebra of Type II_∞ . The Type II_∞ algebra is the crossed product of the Type III_1 algebra by its modular automorphism group. In the context of the emergent Type II_∞ algebra, the entropy of a black hole state is well-defined up to an additive constant, independent of the state. This is somewhat analogous to entropy in classical physics.

KEYWORDS: $1/N$ Expansion, AdS-CFT Correspondence, Black Holes

ARXIV EPRINT: [2112.12828](https://arxiv.org/abs/2112.12828)

Contents

1	Introduction	1
2	The large N limit	2
3	The $1/N$ expansion and the crossed product	9
3.1	Beyond the large N limit	9
3.2	The crossed product	14
3.3	Classical-quantum states and the modular operator	17
3.4	Traces	19
3.5	Density matrices and entropies	21
4	Other conserved charges	23

1 Introduction

Recently Leutheusser and Liu [1, 2] studied the operator algebra of $\mathcal{N} = 4$ super Yang-Mills theory in a novel way, arguing that in the strict large N limit, at a temperature above the Hawking-Page transition, there is an emergent operator algebra that is a von Neumann algebra of Type III₁. In the context of the thermofield double state, which is dual to a two-sided eternal black hole [3], the emergent Type III₁ algebra and its restrictions to suitable regions of the boundary (or bulk) led to an interesting new perspective on black hole physics.

In the present article, certain $1/N^2$ corrections to this picture will be analyzed. In the bulk dual of $\mathcal{N} = 4$ super Yang-Mills theory, Newton’s constant G is proportional to $1/N^2$, so including corrections of order $1/N^2$ in the boundary theory is equivalent to including corrections of order G in the bulk. In the strict large N limit, the algebra of observables on the right or left side of the black hole horizon contains a central generator, related to the black hole mass or horizon area. With $\mathcal{O}(G)$ corrections included, that is no longer the case and the right and left algebras become “factors” in von Neumann algebra language — that is, algebras with trivial center, analogous to a simple Lie group. More specifically, the correction of order G or $1/N^2$ that we analyze deforms the Type III₁ algebra of the large N limit into a factor of Type II_∞.

The fact that the limiting large N algebra has a nontrivial central generator is the reason that a correction perturbative in $1/N$ can qualitatively change the algebra. There are certainly additional contributions in the $1/N$ expansion that we do not analyze in this article, but once the center has been eliminated, we do not expect further perturbative corrections to change the nature of the algebra. Nonperturbative corrections are another story, of course, since if N is set to a definite integer, the algebra should be of Type I.

Mathematically, the Type II_∞ algebra arises as the “crossed product” of the Type III_1 algebra of the strict large N limit by its group of modular automorphisms. The crossed product construction has the surprising property of not depending on the cyclic separating state that is used to define it. This fact has been important in the mathematical theory of Type III_1 algebras [4–6] and is interesting physically because it might be a step toward understanding the algebra of observables outside a black hole horizon in a “background-independent” way.

Deforming the algebra of operators exterior to a black hole from Type III to Type II brings black hole physics closer to the standard framework of quantum mechanics. The operator algebra of an ordinary quantum system is of Type I. Type I algebras have pure states, as well as other familiar quantum concepts such as density matrices and von Neumann entropies. If in a bulk language, one could describe the algebra of observables exterior to a black hole as an algebra of Type I, the black hole information problem would very likely be solved. The algebra of observables of a local region in quantum field theory is of Type III. A Type III algebra does not have pure states, density matrices, or von Neumann entropies. The fact that the algebra of observables outside the horizon of a black hole is of Type III in the limit $G \rightarrow 0$ is one manifestation of the black hole information problem. Type II algebras are intermediate between the two cases. A Type II algebra does not have pure states, but it does have density matrices and von Neumann entropies. Describing the operators outside the black hole horizon by an algebra of Type II means that we do not understand the microstates of a black hole, but we do have a framework to analyze the black hole entropy (up to an overall additive constant, as will be explained).

In section 2 of this article, we will review aspects of the work of Liu and Leutheusser [1, 2], explaining some things in a way that will be helpful later. In section 3, we introduce the crossed product construction and the deformation of the emergent Type III_1 algebra of the strict large N limit to an algebra of Type II_∞ . We explain how to define a trace and a concept of entropy in the crossed product algebra.

As already explained, the deformation to a Type II algebra results from properly taking into account the black hole mass or area, which one can think of as a collective coordinate related to one of the conserved quantities that the black hole can carry, namely its energy. The black hole can carry other conserved charges, notably angular momentum and gauge charges, and a fuller description must include additional collective coordinates. As explained in section 4, this requires making a crossed product not just with the modular automorphism group but with a larger group of automorphisms. This more extended version of the crossed product, however, does not lead to a further qualitative change in the algebraic description.

Some background that might be helpful in understanding the present article can be found in [7, 8].

2 The large N limit

With a convenient normalization of the fields, the action of $\mathcal{N} = 4$ super Yang-Mills theory with gauge group $\text{SU}(N)$ is conveniently written $N\text{Tr } L$, where L is a gauge-invariant polynomial in the fields and their derivatives (with no explicit factors of N). With this




Figure 1. (a) A two-sided eternal black hole, with shaded regions to the future of the future horizon and the past of the past horizon. Operators on the right and left boundaries, labeled R and L , describe physics in the unshaded regions to the right or left of the black hole, respectively. Also sketched is a Cauchy hypersurface S that passes through the bifurcate horizon. The horizon divides S in a natural way into left and right portions S_ℓ and S_r . (b) A time band $B = (t_0, \infty)$ of the right boundary. According to [1, 2], in the large N limit, operators in B describe physics in the causal wedge of B , which is sketched in black.

normalization, correlation functions of any single-trace operator $\mathcal{W} = \text{Tr } W$, where W is a gauge-invariant polynomial in the fields and their derivatives (again with no explicit powers of N), have a simple large N scaling. In the vacuum or in a thermal state, \mathcal{W} has an expectation value of order N , a connected two-point function of order 1, and more generally a connected k -point function of order N^{2-k} . Therefore, to define a large N limit of the operator algebra, as analyzed by Leutheusser and Liu (LL) in [1, 2], we consider subtracted single-trace operators of the form $\mathcal{W} = \text{Tr } W - \langle \text{Tr } W \rangle$, with all possible choices of W . Since $\langle \text{Tr } W \rangle$ diverges for $N \rightarrow \infty$, the subtraction is necessary in order to get a large N limit.

For example, LL consider a thermal state at temperature T , on a spatial manifold S^3 . They are primarily interested in the case that T exceeds the Hawking-Page temperature T_{HP} , or equivalently that $\beta = 1/T$ is less than $\beta_{HP} = 1/T_{HP}$, because in this case a thermal ensemble on the boundary of AdS_5 is dual to a black hole in the interior. In fact, they consider a pair of copies of the theory, the “right” and “left” copies R and L , entangled in the thermofield double state Ψ_{TFD} at temperature $T > T_{HP}$, and dual to a two-sided eternal black hole [3]; see figure 1(a). For $T > T_{HP}$, the free energy of $\mathcal{N} = 4$ super Yang-Mills theory is of order N^2 for large N , with a T -dependent coefficient, just as in the case of QCD [9]. Likewise the terms of order N in the thermal expectation value $\langle \text{Tr } W \rangle$ (for various choices of W) are T -dependent in the high temperature phase, and therefore the algebra defined by LL depends explicitly on T . Similarly, if we introduce chemical potentials for other conserved charges that the black hole can carry — angular momentum and the charges associated with the R-symmetry group $\text{SU}(4)_R$ of $\mathcal{N} = 4$ super Yang-Mills — the large N limit of the algebra of single-trace operators depends on those chemical potentials.

As already remarked, the subtracted single-trace operators have Gaussian correlation functions: two-point functions of order 1, and k -point functions for $k > 2$ that vanish at $N = \infty$. Therefore, in the large N limit, commutators of single-trace operators (and their anticommutators, in the case of fermionic operators) are c -numbers. The matrix of commutators has a finite-dimensional kernel, as explained presently. For the moment we postpone discussion of the kernel and consider only the single-trace operators that

have nonzero commutators. Because those commutators are c -numbers, the subtracted single-trace operators describe a generalized free field theory in the large N limit, as explained by LL. From bounded functions of the operators of the generalized free field theory, we can make a von Neumann algebra \mathcal{A} . This is the emergent von Neumann algebra defined in [1, 2].

In the thermofield double state, we really have two copies of this algebra, the algebras $\mathcal{A}_{R,0}$ and $\mathcal{A}_{L,0}$ of the right and left system. (The subscript 0 means that we do not yet include the operators with trivial commutators; it will be removed when we include them.) LL give two explanations of the fact that $\mathcal{A}_{R,0}$ and $\mathcal{A}_{L,0}$ are von Neumann algebras of¹ Type III₁. One argument uses AdS/CFT duality. $\mathcal{A}_{R,0}$ and $\mathcal{A}_{L,0}$ are dual to the algebras $\mathcal{A}_{r,0}$ and $\mathcal{A}_{\ell,0}$ of low energy effective field theory outside the black hole horizon, on the right and left side respectively:

$$\mathcal{A}_{R,0} = \mathcal{A}_{r,0}, \quad \mathcal{A}_{L,0} = \mathcal{A}_{\ell,0}. \tag{2.1}$$

It will become clear that not including the central generators in $\mathcal{A}_{R,0}$ and $\mathcal{A}_{L,0}$ is dual to defining $\mathcal{A}_{r,0}$ and $\mathcal{A}_{\ell,0}$ in terms of low energy effective field theory in the field of a black hole of definite mass. When we extend $\mathcal{A}_{r,0}$ and $\mathcal{A}_{\ell,0}$ to let the mass vary, we will remove the subscript 0. Like the algebra of any local region in quantum field theory, $\mathcal{A}_{r,0}$ and $\mathcal{A}_{\ell,0}$ are of Type III₁. See footnote 8 below for a second explanation that the algebras are of Type III₁.

A black hole carries conserved charges — it has a mass or energy, and it can carry angular momentum or conserved gauge charges.² These charges are determined by the temperature and chemical potentials of the boundary theory. Since $\mathcal{A}_{L,0}$ and $\mathcal{A}_{R,0}$ depend on the temperature and the chemical potentials, likewise $\mathcal{A}_{\ell,0}$ and $\mathcal{A}_{r,0}$ depend on the black hole mass and other conserved charges. The mass and other conserved charges carried by the black hole are parameters of the classical black hole solution. One can think of those parameters as collective coordinates, which are shared between the left and right of the black hole and are not part of what is described by the left or right bulk algebras $\mathcal{A}_{\ell,0}$ and $\mathcal{A}_{r,0}$. $\mathcal{A}_{\ell,0}$ and $\mathcal{A}_{r,0}$ describe small fluctuations to the left or right of the black hole horizon, for particular values of the collective coordinates.

Of the conserved charges carried by the black hole, the one that is really important for a qualitative understanding is the mass or energy. We will postpone discussing the other conserved charges until section 4.

Since the black hole mass is not a generator of the bulk algebra $\mathcal{A}_{\ell,0}$ or $\mathcal{A}_{r,0}$, the gauge theory Hamiltonians H_L or H_R , which generate translations of the time coordinate t along the left or right boundaries, must not be part of the boundary algebra $\mathcal{A}_{L,0}$ or $\mathcal{A}_{R,0}$. The mechanism by which this comes about is straightforward. The Hamiltonian of $\mathcal{N} = 4$ super Yang-Mills theory, like the Lagrangian, is defined with an explicit factor of N :

$$H_R = \frac{N}{2g^2} \int_S d^3x \sqrt{g} \text{Tr} \left(F_{0i} F_{0j} g^{ij} + \dots \right), \tag{2.2}$$

¹An introduction to the different types of von Neumann algebra can be found, for example, in section 3 of [7] or section 6 of [8].

²In the context of $\mathcal{N} = 4$ super Yang-Mills theory, the conserved gauge charges are dual to the R-symmetry charges of the boundary theory.

where various additional terms are omitted. Above the Hawking-Page transition, because of the explicit factor of N , H_R has a thermal expectation value of order N^2 and also a connected two-point function of order N^2 . We can of course subtract the expectation value and consider the operator $H'_R = H_R - \langle H_R \rangle$. But because $\langle H'^2_R \rangle \sim N^2$, H'_R does not have a large N limit.³

If we simply divide H'_R by N , we get an operator with bounded fluctuations that does have a large N limit:

$$U = \frac{1}{N} H'_R. \tag{2.3}$$

The correlation functions of U have the same large N scaling as any other single-trace operator that is defined with no explicit power of N , so in particular U has a large N limit. However, in the large N limit, U is central. That is because, for any $\mathcal{V} \in \mathcal{A}_R$,

$$[U, \mathcal{V}] = \frac{1}{N} [H_R, \mathcal{V}] = -\frac{i}{N} \frac{\partial \mathcal{V}}{\partial t}. \tag{2.4}$$

Thus, at $N = \infty$, U becomes central and commutes $\mathcal{A}_{R,0}$. Since we defined $\mathcal{A}_{R,0}$ to consist only of the single-trace operators that have nontrivial commutators, U is not part of $\mathcal{A}_{R,0}$. Shortly, we will define an extended algebra \mathcal{A}_R with U as an additional (central) generator.

Because H_R is not part of $\mathcal{A}_{R,0}$, time translations are a group of outer automorphisms⁴ of that algebra. This fact is very important in the work of LL. Let B be a “time band” $t_0 < t < t_1$ in the right boundary, with $t_0, t_1 \in \mathbb{R} \cup \pm\infty$. Let $\mathcal{A}^B_{R,0}$ be the algebra generated by subtracted single-trace operators in B . Any local operator can be conjugated into the time band by a time translation. So if time translations were part of $\mathcal{A}_{R,0}$, $\mathcal{A}^B_{R,0}$ would simply coincide with $\mathcal{A}_{R,0}$. Since the generator of time translations is actually not part of $\mathcal{A}_{R,0}$, $\mathcal{A}^B_{R,0}$ is a proper subalgebra of $\mathcal{A}_{R,0}$. As explained by LL, the algebra $\mathcal{A}^B_{R,0}$ is dual to the bulk algebra of operators in the “causal wedge” of the time band B (figure 1(b)), and thus is of Type III₁, like the bulk algebra of any local region. If one backs away from the strict large N limit and considers an expansion in powers of $1/N$, then as asserted in [1, 2], the definition of $\mathcal{A}^B_{R,0}$ makes sense to all finite orders in $1/N$, since if $\phi(\vec{x}, t)$ is a local operator in a given time band, its time derivatives of any finite order are also local operators in the same time band.

Although the operators H_R and H_L do not have large N limits, the difference $\widehat{H} = H_R - H_L$ does have such a limit. Indeed, \widehat{H} annihilates the thermofield double state Ψ_{TFD} , so in particular it does not have divergent fluctuations in that state, in contrast to H_R and H_L . From a boundary point of view, the thermofield double Hilbert space \mathcal{H}_{TFD} is defined as the Hilbert space spanned by the states $\mathfrak{a}\Psi_{\text{TFD}}$ with $\mathfrak{a} \in \mathcal{A}_{R,0}$. (It can equally well be defined as the Hilbert space spanned by states $\mathfrak{a}\Psi_{\text{TFD}}$ with $\mathfrak{a} \in \mathcal{A}_{L,0}$; this gives the same

³This is analogous to the fact that in quantum statistical mechanics, the usual Hilbert space and Hamiltonian do not have an infinite volume limit (even if one subtracts a constant from the Hamiltonian so that its thermal expectation value vanishes). This is why the thermofield double was introduced [10]. It provides a way to describe the infinite volume limit in a Hilbert space. See for example [7].

⁴Similarly, in the large volume limit of quantum statistical mechanics, time translations are a group of outer automorphisms of the algebra of observables that acts on the thermofield double Hilbert space [10]; see [7] for an explanation.

Hilbert space.) From a bulk point of view, \mathcal{H}_{TFD} is more simply the Hilbert space obtained by quantizing the small fluctuations about the eternal black hole (for fixed values of the collective coordinates). The fact that $\hat{H} = H'_R - H'_L$ exists as an operator acting on \mathcal{H}_{TFD} but H'_R and H'_L do not shows that \mathcal{H}_{TFD} is not a simple tensor product⁵ of left and right Hilbert spaces acted on by $\mathcal{A}_{L,0}$ and $\mathcal{A}_{R,0}$. This nonfactorization of \mathcal{H}_{TFD} in the large N limit is also related to the fact that $\mathcal{A}_{L,0}$ and $\mathcal{A}_{R,0}$, defined in the large N limit, are not of Type I. If these algebras were of Type I, they would have irreducible representations \mathcal{H}_R and \mathcal{H}_L , and we could hope for a factorization $\mathcal{H}_{\text{TFD}} = \mathcal{H}_L \otimes \mathcal{H}_R$, similarly to what actually happens for large N below the Hawking-Page temperature. As the algebras are of Type III, there are no candidates for \mathcal{H}_L or \mathcal{H}_R .

What is the bulk dual of the fact that $H'_R = H_R - \langle H_R \rangle$ and $H'_L = H_L - \langle H_L \rangle$ do not have large N limits, but $\hat{H} = H'_R - H'_L$ does have such a limit? The eternal black hole has a Killing vector field V that generates time translations. V is future-directed timelike on the right side of the black hole outside the horizon and past-directed timelike on the left side of the black hole outside the horizon. The associated conserved charge is

$$\hat{h} = \int_S d\Sigma^\mu V^\mu T_{\mu\nu}, \tag{2.5}$$

where S is a Cauchy hypersurface of the bulk theory, and $T_{\mu\nu}$ is the energy-momentum tensor of the bulk fields (including the energy-momentum pseudo-tensor of the bulk gravitational fluctuations). The operator \hat{h} has a simple boundary dual

$$\hat{h} = \beta \hat{H}. \tag{2.6}$$

Eq. (2.6) reflects the fact that the vector field V reduces to $\beta \partial_t$ on the right boundary, and to $-\beta \partial_t$ on the left boundary. To try to split \hat{h} as the difference of a “right” part and a “left” part, one can proceed as follows. Choose the Cauchy hypersurface S to pass through the bifurcate horizon where the left and right exteriors of the eternal black hole meet (figure 1(a)), and write $S = S_\ell \cup S_r$, where S_ℓ and S_r are the left and right portions of S . Then define

$$\begin{aligned} h_r &= \int_{S_r} d\Sigma^\mu V^\mu T_{\mu\nu} \\ h_\ell &= - \int_{S_\ell} d\Sigma^\mu V^\mu T_{\mu\nu} \\ \hat{h} &= h_r - h_\ell. \end{aligned} \tag{2.7}$$

(A minus sign is included in the definition of h_ℓ because V is past-directed in the left region; with this minus sign included, h_ℓ formally propagates states towards the future.) These expressions formally split \hat{h} as a difference between right and left operators. The problem with the splitting is that, because of divergent fluctuations, h_r and h_ℓ do not make sense as operators. Actually, h_r and h_ℓ do make sense as quadratic forms; that is, for suitable states $\Psi, \Psi' \in \mathcal{H}_{\text{TFD}}$, the matrix elements $\langle \Psi' | h_r | \Psi \rangle$ and $\langle \Psi' | h_\ell | \Psi \rangle$ are well-defined. But h_r

⁵This fact has an analog in the thermodynamic limit of quantum statistical mechanics [10]; see [7].

and h_ℓ do not make sense as operators on \mathcal{H}_{TFD} , because for example, due to an ultraviolet divergence near the horizon, $|h_r|\Psi_{\text{TFD}}\rangle|^2 = \langle\Psi_{\text{TFD}}|h_r^2|\Psi_{\text{TFD}}\rangle = \infty$, so $h_r|\Psi_{\text{TFD}}\rangle$ is not a Hilbert space state. More generally, for any $\Psi \in \mathcal{H}_{\text{TFD}}$, $h_r\Psi$ is not square-integrable and so not in \mathcal{H}_{TFD} . All this is in parallel with properties of the boundary operators H'_R and H'_L , which likewise make sense as quadratic forms but, because of fluctuations, do not make sense as operators on \mathcal{H}_{TFD} .

The modular operator⁶ of the state Ψ_{TFD} for the algebra $\mathcal{A}_{R,0}$ is $\Delta = e^{-\beta\hat{H}}$. (Exchanging $\mathcal{A}_{R,0}$ and $\mathcal{A}_{L,0}$ reverses the sign of \hat{H} , so the modular operator of $\mathcal{A}_{L,0}$ for the state Ψ_{TFD} is $\Delta^{-1} = \exp(+\beta\hat{H})$.) One way to deduce the formula $\Delta = e^{-\beta\hat{H}}$ is to start with finite N , where the Hilbert space of the thermofield double is a simple tensor product $\mathcal{H}_L \otimes \mathcal{H}_R$, and the algebras of observables on the left and right factors are of Type I. In this description, the left and right Hamiltonians are $H_L = H \otimes 1$, $H_R = 1 \otimes H$, where H is the usual Hamiltonian on a single copy of the system. In such a Type I situation, a state $\Psi \in \mathcal{H}_L \otimes \mathcal{H}_R$ has reduced density matrices ρ_R and ρ_L for the right and left systems and the modular operator of Ψ for \mathcal{A}_R is a tensor product $\Delta = \rho_L^{-1} \otimes \rho_R$. In the case $\Psi = \Psi_{\text{TFD}}$, the density matrices are thermal: $\rho_R = \rho_L = e^{-\beta H}/Z$, where Z is the partition function, so $\Delta = \exp(-\beta H_R + \beta H_L) = \exp(-\beta\hat{H})$. Both Δ and \hat{H} have a large N limit, and, since the relation $\Delta = e^{-\beta\hat{H}}$ holds for every N , it automatically is valid in the large N limit. By contrast, the intermediate steps in the derivation involved operators ρ_R and ρ_L and Hilbert spaces \mathcal{H}_R and \mathcal{H}_L that do not have large N limits. The formula $\Delta = e^{-\beta\hat{H}}$ with $\hat{H}\Psi_{\text{TFD}} = 0$ implies

$$\Delta\Psi_{\text{TFD}} = \Psi_{\text{TFD}}. \tag{2.8}$$

Since $\beta\hat{H}$ is equal to the bulk operator \hat{h} , the bulk counterpart of the boundary statement $\Delta = e^{-\beta\hat{H}}$ is simply $\Delta = e^{-\hat{h}}$. The result $\Delta = e^{-\hat{h}}$ can also be deduced from a classical result of Bisognano and Wichman about Rindler space [11], or more exactly from the analog of this result for the eternal black hole.⁷

LL in [1, 2] exploited the relation $\Delta = e^{-\beta\hat{H}}$ or equivalently $\beta\hat{H} = -\log \Delta$ in the following way. Conjugation by $e^{iu\hat{H}}$, $u \in \mathbb{R}$ is called modular flow. Since \hat{H} generates time translations on the right boundary, modular flow for the algebra $\mathcal{A}_{R,0}$ with parameter u shifts the boundary time by $t \rightarrow t + \beta u$. Let B be the time band $t_0 < t < \infty$ (figure 1(b)). Modular flow for the algebra $\mathcal{A}_{R,0}$ with flow parameter $u > 0$ maps B to itself, and therefore it conjugates $\mathcal{A}_{R,0}^B$ to itself. Such a “half-sided modular inclusion” of von Neumann algebras⁸ has very strong implications, which LL use in proposing a way to probe behind the black hole horizon. One notable fact about their construction is the following. As LL explain, the half-sided modular inclusion in this problem can be seen to lowest order in G directly in the bulk description. Purely from a bulk point of view, it would not be clear that this structure persists beyond lowest order, when quantum fluctuations of the spacetime are

⁶See for example [8] for basic facts about modular operators.

⁷This analog was first described by Sewell [12], who also observed a close relation to Unruh’s thermal interpretation of Rindler space [13].

⁸Half-sided modular inclusions exist only for algebras of Type III₁ [14]. As observed in [1, 2], the existence of the half-sided modular inclusion $\mathcal{A}_{R,0}^B \subset \mathcal{A}_{R,0}$ gives an explanation of the Type III₁ nature of $\mathcal{A}_{R,0}$ that does not require duality with a bulk description.

taken into account. However, in the boundary theory, it is obvious that the definitions of $\mathcal{A}_{R,0}$ and $\mathcal{A}_{R,0}^B$ and the statements about a half-sided modular inclusion make sense to all orders in $1/N$. Hence from a bulk point of view, this structure exists to all orders in G .

The state Ψ_{TFD} for the algebra $\mathcal{A}_{R,0}$ is defined so that thermal expectation values are the same as expectation values in the state Ψ_{TFD} :

$$\langle \mathbf{a} \rangle = \langle \Psi_{\text{TFD}} | \mathbf{a} | \Psi_{\text{TFD}} \rangle, \quad \mathbf{a} \in \mathcal{A}_{R,0}. \quad (2.9)$$

The same formula holds for $\mathbf{a} \in \mathcal{A}_{L,0}$. Defined this way, Ψ_{TFD} describes expectation values of operators in $\mathcal{A}_{R,0}$ and $\mathcal{A}_{L,0}$, or their bulk duals $\mathcal{A}_{r,0}$ and $\mathcal{A}_{\ell,0}$, without taking into account the central operator U . Let us define an extended algebra \mathcal{A}_R that includes U . We define \mathcal{A}_R as a tensor product

$$\mathcal{A}_R = \mathcal{A}_{R,0} \otimes \mathcal{A}_U, \quad (2.10)$$

where \mathcal{A}_U is the abelian algebra of bounded functions of U . While $\mathcal{A}_{R,0}$ acts on \mathcal{H}_{TFD} , \mathcal{A}_R acts on $\widehat{\mathcal{H}}_{\text{TFD}} = \mathcal{H}_{\text{TFD}} \otimes L^2(\mathbb{R})$, where $L^2(\mathbb{R})$ is the space of square-integrable functions of U , with U acting by multiplication. An element of \mathcal{A}_R is a U -dependent element of $\mathcal{A}_{R,0}$ such as

$$\widehat{\mathbf{a}} = \int_{-\infty}^{\infty} du \mathbf{a}(u) e^{iuU}, \quad \mathbf{a}(u) \in \mathcal{A}_{R,0} \quad (2.11)$$

(here $\mathbf{a}(u)$ need not be a smooth function of u ; for example, we may have $\mathbf{a}(u) = \mathbf{a} \delta(u - u_0)$ for some \mathbf{a}, u_0 , corresponding to $\widehat{\mathbf{a}} = \mathbf{a} e^{iu_0 U}$). We can define an extended thermofield double state that describes thermal expectation values of elements of the extended algebra \mathcal{A}_R . To do this, we simply note that, as its connected k -point functions vanish in the large N limit for $k > 2$, the thermal correlation functions of U are Gaussian. In other words, there is a Gaussian function $g(U)$ such that for any k , $\langle U^k \rangle = \int_{-\infty}^{\infty} dU g(U) U^k$. By definition, $U = (H_R - \langle H_R \rangle)/N$ has mean 0; its variance is the heat capacity $\langle U^2 \rangle = -(1/N^2) \partial_\beta \langle H_R \rangle$. So the Gaussian function $g(U)$ is concretely

$$g(U) = \left(\frac{N^2}{2\pi |\partial_\beta \langle H_R \rangle|} \right)^{1/2} \exp \left(-\frac{N^2 U^2}{2 |\partial_\beta \langle H_R \rangle|} \right). \quad (2.12)$$

We can define an extended thermofield double state $\widehat{\Psi}_{\text{TFD}} = g(U)^{1/2} \Psi_{\text{TFD}}$ that captures thermal expectation values of operators in the extended algebra \mathcal{A}_R , in the sense that for $\widehat{\mathbf{a}}$ as in (2.11), we have

$$\langle \widehat{\mathbf{a}} \rangle = \langle \widehat{\Psi}_{\text{TFD}} | \widehat{\mathbf{a}} | \widehat{\Psi}_{\text{TFD}} \rangle = \int_{-\infty}^{\infty} dU g(U) \int_{-\infty}^{\infty} du e^{iuU} \langle \Psi_{\text{TFD}} | \mathbf{a}(u) | \Psi_{\text{TFD}} \rangle. \quad (2.13)$$

Because U is central in the algebra \mathcal{A}_R , the state $\widehat{\Psi}_{\text{TFD}}$ is what is called a classical-quantum state. In general, the definition of a classical-quantum state is as follows. Consider a bipartite quantum system AB with Hilbert space $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$. (In our application, $\mathcal{H}_{AB} = \widehat{\mathcal{H}}_{\text{TFD}}$, $\mathcal{H}_A = L^2(\mathbb{R})$ and $\mathcal{H}_B = \mathcal{H}_{\text{TFD}}$.) A classical-quantum density matrix on \mathcal{H}_{AB} is a density matrix of the form $\rho_{AB} = \sum_i \sqrt{p_i} |i\rangle_A \langle i|_A \otimes \rho_{i,B}$, for some basis $|i\rangle_A$ of \mathcal{H}_A , some density matrices $\rho_{i,B}$ on \mathcal{H}_B , and some positive numbers p_i with $\sum_i p_i = 1$. Equivalently, ρ_{AB} commutes with a maximal set of commuting operators of the A system (the diagonal

matrices in the basis $|i\rangle_A$). Because U is central, not only is $\widehat{\Psi}_{\text{TFD}}$ a classical-quantum state, but actually any state of the algebra \mathcal{A}_R is classical-quantum. When we deform away from the large N limit in section 3, there will no longer be a central element U , and the generic state will no longer be classical-quantum.

In this discussion, we extended the algebra $\mathcal{A}_{R,0}$ by adding another generator $U = H'_R/N$. What happens if we also define $U' = H'_L/N$ and add it to \mathcal{A}_L ? U' is not really “new” since $U - U' = \widehat{H}/N$, where \widehat{H} has a large N limit, so that \widehat{H}/N vanishes at $N = \infty$. So to get a proper arena for the action of all boundary operators in the large N limit, we only have to add one new generator U associated to fluctuations in energy (plus additional ones that are related in the same way to the angular momentum and conserved R-charges; see section 4). Thus to the algebras $\mathcal{A}_{L,0}$ and $\mathcal{A}_{R,0}$, we only add one new generator U which is in a sense “shared” between the two sides, rather as from a bulk point of view the horizon area A is visible from either side of the horizon and thus is “shared.” In the large N limit,

$$\mathcal{A}_L = \mathcal{A}_{L,0} \otimes \mathcal{A}_U, \tag{2.14}$$

in perfect parallel with eq. (2.10). With $1/N$ corrections included, matters are more subtle, as we will see in section 3, and it is no longer true that \mathcal{A}_L and \mathcal{A}_R have in common an operator such as U .

As explained in [1, 2], the algebra \mathcal{A}_R is of Type III in the large N limit. However, it is not a “factor” in von Neumann algebra language because its center does not just consist of the complex scalars \mathbb{C} . Rather, its center is the infinite-dimensional commutative algebra \mathcal{A}_U . It may seem somewhat nongeneric that \mathcal{A}_R has such a large center at $N = \infty$. We will learn in section 3 that $1/N$ corrections modify the algebra so that its center becomes trivial. In the process, the algebra will be deformed from being of Type III to being of Type II. Indeed, it will become a factor of Type II_∞ .

We conclude with a few facts that will be useful in section 3. First, if an algebra \mathcal{A} acts on a Hilbert space \mathcal{H} , a vector $\Psi \in \mathcal{H}$ is said to be cyclic for \mathcal{A} if \mathcal{H} is spanned by vectors $\mathbf{a}\Psi$, $\mathbf{a} \in \mathcal{A}$. It is said to be separating for \mathcal{A} if $\mathbf{a}\Psi = 0$, $\mathbf{a} \in \mathcal{A}$, implies $\mathbf{a} = 0$. The vector $\widehat{\Psi}_{\text{TFD}}$ is separating for \mathcal{A}_R because, as the thermal expectation value of any operator $\widehat{\mathbf{a}}^\dagger \widehat{\mathbf{a}}$ is always strictly positive for any $\widehat{\mathbf{a}} \neq 0$, eq. (2.13) implies that $\widehat{\mathbf{a}} \widehat{\Psi}_{\text{TFD}} \neq 0$ for all nonzero $\widehat{\mathbf{a}} \in \mathcal{A}_R$. $\widehat{\Psi}_{\text{TFD}}$ is also cyclic for \mathcal{A}_R because (1) \mathcal{H}_{TFD} was defined in a way that ensured that Ψ_{TFD} is cyclic for $\mathcal{A}_{R,0}$; (2) since the Gaussian function $g(U)$ is everywhere positive, any function $f(U)$ can be factored as $m(U)g^{1/2}(U)$ for some function $m(U)$, and therefore the state $g^{1/2}(U)$ is cyclic for the abelian algebra \mathcal{A}_U , acting on $L^2(\mathbb{R})$. The statement that the product state $\widehat{\Psi}_{\text{TFD}}$ is cyclic for $\mathcal{A}_R = \mathcal{A}_{R,0} \otimes \mathcal{A}_U$ is just a composite of these two statements.

3 The $1/N$ expansion and the crossed product

3.1 Beyond the large N limit

Our goal in this section is to understand $1/N$ corrections to the picture described in section 2. In section 2, we considered an algebra generated by single-trace operators with

their thermal expectation values subtracted, an example being $\mathcal{W} = \text{Tr } W - \langle \text{Tr } W \rangle$, where W is a polynomial in the fields and their derivatives. A typical element of the algebra was a complex linear combination of products of such operators. To depart from the strict large N limit, we have to modify the definition slightly so that coefficients are not complex numbers, but rather functions of N that have an asymptotic expansion in powers of $1/N^2$ around the $N = \infty$ limit. Thus an element of the algebra is a sum of elements

$$\sum_{k=0}^{\infty} a_k N^{-2k} \mathcal{W}_1 \mathcal{W}_2 \cdots \mathcal{W}_s, \quad a_k \in \mathbb{C}. \quad (3.1)$$

(The coefficients a_k depend on the choice of $\mathcal{W}_1, \dots, \mathcal{W}_s$. The associated von Neumann algebra is generated by bounded functions of such expressions.) Another way to say this is that we work not over \mathbb{C} but over a formal power series ring $\mathbb{C}[[1/N^2]]$. This is necessary because operator product coefficients and commutation relations of $\mathcal{N} = 4$ super Yang-Mills theory have nontrivial asymptotic expansions in powers of $1/N^2$. To define an algebra that is consistent with operator product expansions and commutators, we have to allow the operators to have N -dependent coefficients of this form.

In section 2, we described the algebra of bulk operators in the field of a black hole, in the large N limit, in terms of the left and right algebras $\mathcal{A}_{r,0}$ and $\mathcal{A}_{\ell,0}$ and a central generator U . In this description, the Hilbert space is $\widehat{\mathcal{H}}_{\text{TFD}} = \mathcal{H}_{\text{TFD}} \otimes L^2(\mathbb{R})$, where $\mathcal{A}_{r,0}$ and $\mathcal{A}_{\ell,0}$ act on \mathcal{H}_{TFD} , and $L^2(\mathbb{R})$ is the space of square-integrable functions of U , with U acting by multiplication. We also observed that the boundary operator H'_R/N makes sense in the large N limit as an element of the right algebra \mathcal{A}_R , and in the large N limit we identified H'_R/N with U .

Once we go beyond the $N = \infty$ limit, H'_R/N is no longer central. Rather, for any $\mathfrak{a} \in \mathcal{A}_R$, we have $[H'_R/N, \mathfrak{a}] = (-i/N)\partial_t \mathfrak{a}$. To identify a bulk dual of H'_R/N , we have to find a bulk operator that satisfies the same commutation relation. Such an operator presents itself, namely the operator $\frac{1}{\beta} \widehat{h}$, where \widehat{h} was defined in eq. (2.7). This operator satisfies the desired commutation relation, because \widehat{h} is the conserved charge associated to a Killing vector field that coincides with $\beta \partial_t$ on the right boundary. This suggests the identification

$$\frac{1}{N} H'_R = U + \frac{1}{\beta N} \widehat{h}. \quad (3.2)$$

Formally, one might be tempted to use here h_r/β rather than \widehat{h}/β , on the grounds that the two operators have the same commutation relations with operators that are to the right side of the black hole. However, as was explained earlier, h_r is not a well-defined operator. We will get more insight by expressing all statements in terms of operators that actually exist.

The left and right hand sides of eq. (3.2) have the desired commutation relations with elements of \mathcal{A}_R , but this does not determine the right hand side uniquely: without spoiling anything, we could add an arbitrary element⁹ $\mathfrak{b}_{\ell} \in \mathcal{A}_{\ell,0}$ or an arbitrary function $f(U)$. More generally, we could add any $\widehat{b} \in \mathcal{A}_{\ell,0} \otimes \mathcal{A}_U$. Thus we should ask whether eq. (3.2) should be

⁹We cannot add an element of $\mathcal{A}_{r,0}$, because such an element is noncentral and would contribute to the commutators with elements of $\mathcal{A}_{r,0}$.

corrected to

$$\frac{1}{N}H'_R \stackrel{?}{=} U + \frac{1}{N\beta}\hat{h} + \frac{1}{N}\hat{b} + \mathcal{O}(1/N^2), \quad (3.3)$$

where we also include the possibility of corrections in higher orders in $1/N$. The answer is that to reconstruct \mathcal{A}_R , there is no need to add any such corrections, because they can be removed by conjugation. Thus, we introduce the operator $\Pi = -\text{id}/dU$, acting on $L^2(\mathbb{R})$. Conjugation by $\exp(-i\hat{b}\Pi/N)$ will remove the \hat{b}/N term from the right hand side of eq. (3.3), and order by order in $1/N$, a similar conjugation would remove any possible higher order terms in $1/N$ that could be added in eq. (3.3) without spoiling the commutators with \mathcal{A}_R .

Since $\hat{h} = \beta(H'_R - H'_L)$, eq. (3.2) also implies

$$\frac{1}{N}H'_L = U. \quad (3.4)$$

Thus, order by order in $1/N$, we can interpret the boundary algebra \mathcal{A}_R as the algebra generated by $\mathcal{A}_{r,0}$ together with one more operator $U + \hat{h}/\beta N$, acting on $\mathcal{H}_{\text{TFD}} \otimes L^2(\mathbb{R})$. We will summarize this by saying that order by order in $1/N$, $\mathcal{A}_R = \mathcal{A}_{r,0} \rtimes \mathcal{A}_{U+\hat{h}/\beta N}$. Here $\mathcal{A}_{r,0} \rtimes \mathcal{A}_{U+\hat{h}/\beta N}$ is the algebra generated by $\mathcal{A}_{r,0}$ together with $U + \hat{h}/\beta N$, and the meaning of the symbol \rtimes is that $U + \hat{h}/\beta N$ does not commute with $\mathcal{A}_{r,0}$ but rather generates an outer automorphism of $\mathcal{A}_{r,0}$. At the von Neumann algebra level, we should consider not $U + \hat{h}/\beta N$ but bounded functions of this operator. However, we do not indicate this in the notation.

This description of \mathcal{A}_R , however, is only perturbative in $1/N$. It may not hold if we set N to a definite integer value such as 101, because the series of conjugations that was used to set $H'_R/N = U + \hat{h}/\beta N$ order by order may not converge at integer N . In fact, we expect that to happen, because the description just given leads to a Type II algebra, as we will see, but when N is an integer, \mathcal{A}_R is of Type I.

Going back to perturbation theory in $1/N$, the reader may observe an apparent difficulty. \mathcal{A}_R is supposed to commute with \mathcal{A}_L , which in the large N limit was identified with $\mathcal{A}_{\ell,0} \otimes \mathcal{A}_U$. Our proposal $\mathcal{A}_R = \mathcal{A}_{r,0} \rtimes \mathcal{A}_{U+\hat{h}/\beta N}$ is not consistent with that identification of \mathcal{A}_L . That is because, although $\mathcal{A}_{r,0}$ commutes with $\mathcal{A}_{\ell,0} \otimes \mathcal{A}_U$, $\mathcal{A}_{U+\hat{h}/\beta N}$ does not; $U + \hat{h}/\beta N$ commutes with U but not with $\mathcal{A}_{\ell,0}$. There is, however, a simple fix. Although $U + \hat{h}/\beta N$ does not commute with $\mathfrak{a}_\ell \in \mathcal{A}_{\ell,0}$, it does commute with $e^{i\Pi\hat{h}/\beta N}\mathfrak{a}_\ell e^{-i\Pi\hat{h}/\beta N}$. So we propose that \mathcal{A}_L is generated order by order in $1/N$ by U (or more exactly, by bounded functions of U) along with operators of the form $e^{i\Pi\hat{h}/\beta N}\mathfrak{a}_\ell e^{-i\Pi\hat{h}/\beta N}$. More succinctly, we describe this by writing $\mathcal{A}_L = e^{i\Pi\hat{h}/\beta N}\mathcal{A}_{\ell,0}e^{-i\Pi\hat{h}/\beta N} \rtimes \mathcal{A}_U$.

The underlying thermofield double state is symmetrical between \mathcal{A}_L and \mathcal{A}_R , but there is an apparent asymmetry between these descriptions of \mathcal{A}_L and \mathcal{A}_R . This asymmetry, however, can be removed by conjugating both algebras with $e^{-i\Pi\hat{h}/2N}$. This leads to

$$\begin{aligned} \mathcal{A}_R &= e^{-i\Pi\hat{h}/2\beta N}\mathcal{A}_{r,0}e^{i\Pi\hat{h}/2\beta N} \rtimes \mathcal{A}_{U+\hat{h}/2\beta N} \\ \mathcal{A}_L &= e^{i\Pi\hat{h}/2\beta N}\mathcal{A}_{\ell,0}e^{-i\Pi\hat{h}/2\beta N} \rtimes \mathcal{A}_{U-\hat{h}/2\beta N}. \end{aligned} \quad (3.5)$$

Since \widehat{h} is odd under the exchange of $\mathcal{A}_{\ell,0}$ and $\mathcal{A}_{r,0}$, these formulas treat \mathcal{A}_L and \mathcal{A}_R symmetrically.

However, the symmetrical formulas are slightly less convenient, so we instead will use the description

$$\begin{aligned}\mathcal{A}_R &= \mathcal{A}_{r,0} \rtimes \mathcal{A}_{U+\widehat{h}/\beta N} \\ \mathcal{A}_L &= e^{i\Pi\widehat{h}/\beta N} \mathcal{A}_{\ell,0} e^{-i\Pi\widehat{h}/\beta N} \rtimes \mathcal{A}_U,\end{aligned}\tag{3.6}$$

valid to all orders in $1/N$.

From this description, we can see at least heuristically that the $1/N$ corrections have deformed \mathcal{A}_R so that its center has become trivial. The central element U of \mathcal{A}_R has been deformed to $U + \widehat{h}/\beta N$, an element of \mathcal{A}_R that is not central. It is true that the deformed \mathcal{A}_R commutes with U , but U is not an element of \mathcal{A}_R , so it is not in the center of \mathcal{A}_R . Rather, U is an element of \mathcal{A}_L , which explains why it commutes with \mathcal{A}_R , since \mathcal{A}_L and \mathcal{A}_R commute. In section 3.2, we will understand more precisely that \mathcal{A}_R , as described in eq. (3.6), is a factor of Type II_∞ .

We have expressed eq. (3.6) in terms of the boundary algebras. However, duality tells us to identify these boundary algebras with bulk algebras \mathcal{A}_r and \mathcal{A}_ℓ :

$$\mathcal{A}_R = \mathcal{A}_r, \quad \mathcal{A}_L = \mathcal{A}_\ell.\tag{3.7}$$

Here \mathcal{A}_r is a bulk algebra of observables on the right side of the black hole that incorporates $\mathcal{A}_{r,0}$, the observable U that is central at $N = \infty$, and $1/N$ corrections, and \mathcal{A}_ℓ is an analogous algebra on the left side of the black hole.

One important fact about eq. (3.6) is that it implies that to all orders in $1/N$, the algebras \mathcal{A}_L and \mathcal{A}_R have a group of outer automorphisms that is not part of the exact theory. This is simply the abelian group of translations $U \rightarrow U + c$, $c \in \mathbb{R}$, generated by Π . To see that this is an outer automorphism of \mathcal{A}_R , observe that since the constant c is anyway part of the algebra $\mathcal{A}_{r,0}$, it does not matter whether we adjoin $U + \widehat{h}/\beta N$ or $U + c + \widehat{h}/\beta N$ to $\mathcal{A}_{r,0}$. Similarly, since c is anyway part of $e^{i\Pi\widehat{h}/\beta N} \mathcal{A}_{\ell,0} e^{-i\Pi\widehat{h}/\beta N}$, it does not matter if we adjoin U or $U + c$ to that algebra. It will turn out that there is a notion of entropy for the algebra \mathcal{A}_L or \mathcal{A}_R — in contrast to $\mathcal{A}_{r,0}$ and $\mathcal{A}_{\ell,0}$, for example — but this entropy is shifted by a constant if U is shifted. So the outer automorphism will mean that perturbatively in $1/N$, one can only define entropy differences.

An analog of eqs. (3.2) and (3.4) was expressed by Jafferis, Lewkowycz, Maldacena, and Suh (JLMS) [15] in terms of the “modular Hamiltonian” of the boundary theory, which is defined as minus the logarithm of the density matrix. The density matrix of the theory on the right boundary is $\rho_R = \frac{1}{Z} e^{-\beta H_R}$, so the modular Hamiltonian is $-\log \rho_R = \beta H_R + \log Z$. A similar formula holds for ρ_L . Here we view $\log Z$ as a c -number function of N ; in fact, $\log Z = -\beta \langle H_R \rangle + \langle S \rangle$, where $\langle S \rangle$ is the thermal entropy at inverse temperature β , also viewed as a c -number function of N . The JLMS formula for $\log \rho_R$ and $\log \rho_L$, which has

been important in understanding entanglement wedge reconstruction, reads

$$\begin{aligned}\beta H_R + \log Z &= h_r + \frac{A}{4G} \\ \beta H_L + \log Z &= h_\ell + \frac{A}{4G},\end{aligned}\tag{3.8}$$

where A is the area of the horizon, and a formal splitting $\hat{h} = h_r - h_\ell$ is used. Equivalently,

$$\begin{aligned}\beta H'_R &= h_r + \frac{A}{4G} - \langle S \rangle \\ \beta H'_L &= h_\ell + \frac{A}{4G} - \langle S \rangle.\end{aligned}\tag{3.9}$$

A comparison to eq. (3.4) shows that the relation is

$$\beta NU + \langle S \rangle = h_\ell + \frac{A}{4G}.\tag{3.10}$$

It is believed that only two linear combinations of the three operators h_r , h_ℓ , and $A/4G$, namely $h_r + A/4G$ and $h_\ell + A/4G$ (or equivalently \hat{h} and U) are actually well-defined quantum mechanically. This statement is related to the observation by Susskind and Uglum [16] that the generalized entropy of a black hole, namely $A/4G + S_{\text{out}}$ (where S_{out} is the entropy outside the horizon) is better-defined than either term is separately. If one believed that h_ℓ , h_r , and $A/4G$ are all separately well-defined, one would likely conclude that h_ℓ and h_r , being defined by integrals on the left and right of the horizon, are respectively elements of \mathcal{A}_ℓ and \mathcal{A}_r and therefore of \mathcal{A}_L and \mathcal{A}_R . Since H'_L and H'_R are also contained in \mathcal{A}_L and \mathcal{A}_R , one would then deduce that $A/4G$ is contained in both algebras and therefore is central in each. This is actually true to lowest order in G , but as we have seen not beyond lowest order.

In sections 3.2–3.5, we will explain some of the mathematical theory of the algebras that we encountered in eq. (3.6). These are algebras of a special type that has been important in the mathematical theory of von Neumann algebras of Type III. In explaining the theory, it is convenient to assume that we are working with ordinary functions of N rather than formal power series in $1/N$. If we were working with ordinary functions of N , it would not matter whether what we adjoin to $\mathcal{A}_{r,0}$ to build \mathcal{A}_R is $U + \hat{h}/\beta N$ or $\beta N(U + \hat{h}/\beta N) = \beta NU + \hat{h}$. So, setting

$$X = \beta NU,\tag{3.11}$$

we restate eq. (3.6) in the form

$$\mathcal{A}_R = \mathcal{A}_{r,0} \times \mathcal{A}_{\hat{h}+X}.\tag{3.12}$$

However, for the application to gravity, all formulas of sections 3.2–3.5 are supposed to be expanded in a formal power series in $1/N$ around $N = \infty$, since that was the setting for the derivation of eq. (3.6).

3.2 The crossed product

Suppose that a von Neumann algebra \mathcal{A} acts on a Hilbert space \mathcal{H} , and let T be a self-adjoint operator on \mathcal{H} that generates a group of automorphisms of \mathcal{A} . The condition for this is that

$$e^{iT_s} \mathbf{a} e^{-iT_s} \in \mathcal{A}, \quad \forall \mathbf{a} \in \mathcal{A}, s \in \mathbb{R}. \quad (3.13)$$

This gives an action of the additive group \mathbb{R} on \mathcal{A} by automorphisms. (This action is not always faithful since in general some elements of \mathbb{R} may act trivially.) The “crossed product” of \mathcal{A} by \mathbb{R} , sometimes denoted $\mathcal{A} \rtimes \mathbb{R}$, is defined as follows. Let $L^2(\mathbb{R})$ be the space of square-integrable functions of a real variable X . Then $\mathcal{A} \rtimes \mathbb{R}$ is the algebra that acts on $\mathcal{H} \otimes L^2(\mathbb{R})$ and is obtained by adjoining $T + X$, or more precisely bounded functions of $T + X$, to \mathcal{A} . Thus $\mathcal{A} \rtimes \mathbb{R}$ is generated, as an algebra, by operators $\mathbf{a} \otimes 1$, $\mathbf{a} \in \mathcal{A}$ and $e^{isT} \otimes e^{isX}$, $s \in \mathbb{R}$. Additively, $\mathcal{A} \rtimes \mathbb{R}$ is generated by operators $\mathbf{a} e^{isT} \otimes e^{isX}$, $\mathbf{a} \in \mathcal{A}$, $s \in \mathbb{R}$. To see that operators of the form $\mathbf{a} e^{isT} \otimes e^{isX}$ do form an algebra, observe that for $\mathbf{a}, \mathbf{b} \in \mathcal{A}$, $s, t \in \mathbb{R}$, we have

$$\left(\mathbf{a} e^{isT} \otimes e^{isX} \right) \left(\mathbf{b} e^{itT} \otimes e^{itX} \right) = \mathbf{a} e^{isT} \mathbf{b} e^{-isT} e^{i(s+t)T} \otimes e^{i(s+t)X}, \quad (3.14)$$

and since eq. (3.13) tells us that $e^{isT} \mathbf{b} e^{-isT} \in \mathcal{A}$, the right hand side is in $\mathcal{A} \rtimes \mathbb{R}$. One can often omit the tensor product symbol in formulas such as those of this paragraph without causing confusion.

The automorphism group that we have considered is said to be inner if $e^{isT} \in \mathcal{A}$, $s \in \mathbb{R}$; otherwise it is outer. In the case of an inner automorphism group, adjoining $T + X$ to \mathcal{A} is equivalent to just adjoining X , since bounded functions of T are already part of \mathcal{A} . So in this case, $\mathcal{A} \rtimes \mathbb{R}$ is just a tensor product $\mathcal{A} \otimes \mathcal{A}_X$, where \mathcal{A}_X is the commutative algebra of bounded functions of X . So the crossed product construction applied to an inner automorphism group does not give anything essentially new, and it certainly does not give a factor, since it gives an algebra with the large center \mathcal{A}_X . All automorphisms of a Type I factor are inner, but that is not true for Type II or Type III. The crossed product by \mathbb{R} is therefore potentially most interesting for Type II or Type III.

For the example that is important in the present article, let $\Psi \in \mathcal{H}$ be a cyclic separating vector for \mathcal{A} , and let $\Delta = e^{-\hat{h}}$ be the corresponding modular operator. The main theorem of Tomita-Takesaki theory says (in part) that Δ^{-is} is an automorphism of \mathcal{A} ; in other words, with $T = \hat{h}$, eq. (3.13) is satisfied. For an algebra of Type I, to prove this is a simple exercise with density matrices. In general, the proof is not so simple, but there is a relatively simple proof [21] in the case of most importance in physics, which is a hyperfinite algebra (an algebra that can be approximated by finite-dimensional matrix algebras). The automorphism group \mathbb{R}_Ψ generated by \hat{h} is called the modular automorphism group. Evidently, a restatement of eq. (3.12) is that $\mathcal{A}_R = \mathcal{A}_{r,0} \rtimes \mathbb{R}_{\Psi_{\text{TFD}}}$.

In the mathematical theory of Type III₁ factors [4–6], an important fact is that the crossed product of such a factor by its modular automorphism group \mathbb{R}_Ψ (for any Ψ) is a factor of Type II_∞. This result is relevant for us, because $\mathcal{A}_{r,0}$ is a Type III₁ factor, and therefore \mathcal{A}_R is a Type II_∞ factor. In particular, \mathcal{A}_R is a factor, and thus has trivial center, as was claimed in section 3.1. The Type II nature of \mathcal{A}_R will ultimately enable us to define

traces, density matrices, and entropies. A Type III₁ factor is actually the only type of von Neumann algebra whose crossed product with its modular automorphism group is a factor. So it is only because $\mathcal{A}_{r,0}$ is of Type III₁ that \mathcal{A}_R is a factor.

In the mathematical theory of Type III₁ algebras, it is essential that, despite appearances, for any von Neumann algebra \mathcal{A} with cyclic separating vector Ψ , the algebra $\mathcal{A} \rtimes \mathbb{R}_\Psi$ does not depend on¹⁰ Ψ , up to a unique equivalence. This means that when one studies $\mathcal{A} \rtimes \mathbb{R}_\Psi$, one is learning about properties of \mathcal{A} , not about properties of the pair \mathcal{A}, Ψ . Physically, the fact that $\mathcal{A} \rtimes \mathbb{R}_\Psi$ is essentially independent of Ψ is somewhat similar to a statement of background independence. We will call it state-independence. State-independence of $\mathcal{A} \rtimes \mathbb{R}_\Psi$ is proved using the Connes cocycle [5], which has had a number of interesting recent applications to quantum field theory and gravity [17–20].

As a first step toward the explanation, recall that if \mathcal{A} acts on \mathcal{H} with cyclic separating vector Ω , the Tomita operator S_Ω is defined by¹¹ $S_\Omega a\Omega = a^\dagger\Omega$, $a \in \mathcal{A}$. One also defines the modular operator $\Delta_\Omega = S_\Omega^\dagger S_\Omega$. More generally, if $\Phi \in \mathcal{H}$ is another vector, one defines the relative Tomita operator $S_{\Psi|\Omega}$ via the condition $S_{\Psi|\Omega} a\Omega = a^\dagger\Psi$. The relative modular operator is defined to be $\Delta_{\Psi|\Omega} = S_{\Psi|\Omega}^\dagger S_{\Psi|\Omega}$. Thus in particular

$$S_{\Psi|\Psi} = S_\Psi, \quad \Delta_{\Psi|\Psi} = \Delta_\Psi. \tag{3.15}$$

The Connes cocycle is defined as

$$u_{\Psi|\Omega}(t) = \Delta_{\Psi|\Omega}^{it} \Delta_\Omega^{-it} = \Delta_\Psi^{it} \Delta_{\Omega|\Psi}^{-it} \tag{3.16}$$

Clearly $u_{\Psi|\Omega}(t)$ is unitary for real t , and $u_{\Psi|\Psi}(t) = 1$. Some important additional properties of $u_{\Psi|\Omega}(t)$ are:

- $u_{\Psi|\Omega}(t) \in \mathcal{A}$.
- The two formulas for $u_{\Psi|\Omega}(t)$ given in eq. (3.16) are in fact equal.
- If Φ, Ω and Ψ are three cyclic separating states, then u satisfies a chain rule

$$u_{\Psi|\Omega}(t) u_{\Omega|\Phi}(t) = u_{\Psi|\Phi}(t) \tag{3.17}$$

and in particular

$$u_{\Psi|\Omega}(t) u_{\Omega|\Psi}(t) = 1. \tag{3.18}$$

The equivalence of the two formulas in eq. (3.16) actually amounts to this special case of the chain rule.

The proofs of these statements (due originally to Connes [5]) are explained in section 6 of [18].

¹⁰By contrast, if one more naively omits X and simply extends the algebra \mathcal{A} by adjoining bounded functions of \widehat{h}_Ψ (as opposed to functions of $\widehat{h}_\Psi + X$), the resulting algebra does depend on the choice of Ψ .

¹¹To be more precise, S_Ω is the closure of the operator defined by the following condition. The same remark applies for $S_{\Psi|\Omega}$ below, and similarly later to the Tomita operators of the commutant \mathcal{A}' . Some basic background on the Tomita operators and related matters can be found, for example, in [8]. In the present article, we use the most common convention for the relative Tomita and modular operators. In [8], a different choice was made for a reason explained in footnote 16 of that article.

Let \mathcal{A}' be the commutant of \mathcal{A} (the algebra of bounded operators on \mathcal{H} that commute with \mathcal{A}). If a vector is cyclic and separating for \mathcal{A} , then it is also cyclic and separating for \mathcal{A}' , so one can define operators $S'_\Psi, S'_{\Psi|\Omega}$ characterized by $S'_\Psi \mathbf{a}'\Omega = \mathbf{a}'^\dagger\Omega$, $S'_{\Psi|\Omega} \mathbf{a}'\Omega = \mathbf{a}'^\dagger\Psi$. Likewise, one defines $\Delta'_\Omega = S'^\dagger_\Omega S'_\Omega$, $\Delta'_{\Psi|\Omega} = S'_{\Psi|\Omega}{}^\dagger S'_{\Psi|\Omega}$. The Connes cocycle for \mathcal{A}' is defined by the obvious analog of eq. (3.16). However, one has $\Delta'_\Omega = \Delta_\Omega^{-1}$, $\Delta'_{\Psi|\Omega} = \Delta_{\Psi|\Omega}^{-1}$. So one can write the cocycle for \mathcal{A}' in terms of the modular operators for \mathcal{A} :

$$u'_{\Psi|\Omega}(t) = \Delta_{\Omega|\Psi}^{-it} \Delta_\Omega^{it} = \Delta_\Psi^{-it} \Delta_{\Psi|\Omega}^{it} \quad (3.19)$$

One has the same three properties as before: $u'_{\Psi|\Omega} \in \mathcal{A}'$; the two expressions for $u'_{\Psi|\Omega}$ are equal; given three states, u' satisfies a chain rule $u'_{\Psi|\Omega}(t)u'_{\Omega|\Phi}(t) = u'_{\Psi|\Phi}(t)$.

We now want to show that, given any two cyclic separating vectors Ω, Ψ , the two algebras $\mathcal{A} \rtimes \mathbb{R}_\Omega$ and $\mathcal{A} \rtimes \mathbb{R}_\Psi$ are conjugate in a natural fashion. To be more precise, setting $P = -\text{id}/dX$, the algebras are conjugate via the operator $u'_{\Psi|\Omega}(P)$ that we get by substituting t with P in the definition of $u'_{\Psi|\Omega}(t)$. (This substitution makes sense because P commutes with $u'_{\Psi|\Omega}(t)$ for all t , so we can define an operator $u'_{\Psi|\Omega}(P)$ that acts on a state with $P = t$ as $u'_{\Psi|\Omega}(t)$.) Thus, the claim is that

$$\mathcal{A} \rtimes \mathbb{R}_\Psi = u'_{\Psi|\Omega}(P)(\mathcal{A} \rtimes \mathbb{R}_\Omega)u'_{\Psi|\Omega}(P)^\dagger. \quad (3.20)$$

The chain rule implies that this construction leads to a unique conjugacy between the crossed product algebras for any two states; in other words, given three states Φ, Ω, Ψ , it does not matter if we conjugate directly from $\mathcal{A} \rtimes \mathbb{R}_\Phi$ to $\mathcal{A} \rtimes \mathbb{R}_\Psi$ according to eq. (3.20), or if we conjugate first from $\mathcal{A} \rtimes \mathbb{R}_\Phi$ to $\mathcal{A} \rtimes \mathbb{R}_\Omega$ and then to $\mathcal{A} \rtimes \mathbb{R}_\Psi$. This ability to select a *unique* conjugacy between the two algebras means that they are really physically the same rather than just being abstractly isomorphic.¹²

Since $u'_{\Psi|\Omega}$ is invertible, with inverse $u'_{\Omega|\Psi}$, to establish the conjugacy claimed in eq. (3.20), one just needs to show that $u'_{\Psi|\Omega}$ conjugates any of the generators of $\mathcal{A} \rtimes \mathbb{R}_\Omega$ to an element of $\mathcal{A} \rtimes \mathbb{R}_\Psi$. We will use the fact that $\mathcal{A} \rtimes \mathbb{R}_\Omega$ is generated by operators $\mathbf{a} \otimes 1$, $\mathbf{a} \in \mathcal{A}$, along with $e^{i\widehat{h}_\Omega s} \otimes e^{iXs} = \Delta_\Psi^{-is} \otimes e^{iXs}$, $s \in \mathbb{R}$. More informally, the generators are \mathbf{a} and $\Delta_\Omega^{-is} e^{iXs}$. Similarly, $\mathcal{A} \rtimes \mathbb{R}_\Psi$ is generated by operators \mathbf{a} and $\Delta_\Psi^{-is} e^{iXs}$. The conjugacy we want is trivial for the generators $\mathbf{a} \in \mathcal{A}$: since $u'_{\Psi|\Omega}(s) \in \mathcal{A}'$ for all s , and \mathcal{A} commutes with both \mathcal{A}' and P , $u'_{\Psi|\Omega}(P)$ commutes with \mathcal{A} . So we just need to show the desired conjugacy for the other generators $\Delta_\Omega^{-is} e^{isX}$:

$$u'_{\Psi|\Omega}(P)\Delta_\Omega^{-is} e^{isX} u'_{\Psi|\Omega}(P)^\dagger \in \mathcal{A} \rtimes \mathbb{R}_\Psi, \quad \forall s \in \mathbb{R}. \quad (3.21)$$

Since $[X, P] = i$, we have $e^{isX} u'_{\Psi|\Omega}(P)^\dagger = u'_{\Psi|\Omega}(P - s)^\dagger e^{isX}$. So

$$\begin{aligned} u'_{\Psi|\Omega}(P)\Delta_\Omega^{-is} e^{isX} u'_{\Psi|\Omega}(P)^\dagger &= u'_{\Psi|\Omega}(P)\Delta_\Omega^{-is} u'_{\Psi|\Omega}(P - s)^\dagger e^{isX} \\ &= \Delta_{\Omega|\Psi}^{-iP} \Delta_\Omega^{iP} \Delta_\Omega^{-is} \Delta_\Omega^{-iP+is} \Delta_{\Omega|\Psi}^{iP-is} e^{isX} \\ &= \Delta_{\Omega|\Psi}^{-is} e^{isX} = \Delta_\Psi^{-is} e^{isX} u_{\Psi|\Omega}(s). \end{aligned} \quad (3.22)$$

¹²Abstract isomorphism would have very little import. For example, in any quantum field theory, the algebra of observables in any topologically simple local region in any spacetime is believed to be a factor of Type III₁, so all these are isomorphic; but there is no useful sense in which they are “the same,” since there is no distinguished isomorphism between them.

Since $u_{\Psi|\Omega}(s) \in \mathcal{A} \subset \mathcal{A} \rtimes \mathbb{R}_\Psi$ and $\Delta_\Psi^{-is} e^{isX}$ is also a generator of $\mathcal{A} \rtimes \mathbb{R}_\Psi$, the final expression in eq. (3.22) is the product of two elements of $\mathcal{A} \rtimes \mathbb{R}_\Psi$, and therefore is an element of this algebra. This completes the proof that $u'_{\Psi|\Omega}(P)$ conjugates $\mathcal{A} \rtimes \mathbb{R}_\Omega$ to $\mathcal{A} \rtimes \mathbb{R}_\Psi$.

3.3 Classical-quantum states and the modular operator

In section 2, we encountered an algebra $\mathcal{A}_{r,0} \otimes \mathcal{A}_U$ with a large center. Because of this center, every state for this algebra is classical-quantum in a natural sense. For the crossed product algebra $\mathcal{A}_R = \mathcal{A}_{r,0} \rtimes \mathbb{R}_\Psi$ acting on $\widehat{\mathcal{H}} = \mathcal{H} \otimes L^2(\mathbb{R})$, that is no longer true, but a reasonable class of classical-quantum states are states of the form $\chi \otimes g^{1/2}(X)$, $\chi \in \mathcal{H}$, $g^{1/2}(X) \in L^2(\mathbb{R})$. It follows from the analysis of state independence in section 3.2 that even though the algebra $\mathcal{A}_{r,0} \rtimes \mathbb{R}_\Psi$ is state-independent, this notion of a classical-quantum state is state-dependent. When we identify $\mathcal{A}_{r,0} \rtimes \mathbb{R}_\Omega$ with $\mathcal{A}_{r,0} \rtimes \mathbb{R}_\Psi$ by conjugating the algebra of observables by $u'_{\Psi|\Omega}(P)$, we must also act on the states with the same operator $u'_{\Psi|\Omega}(P)$. But $u'_{\Psi|\Omega}(P)$ does not map a classical-quantum state of the class just described to a state of the same class.

It turns out to be useful to understand modular theory for classical-quantum states of the special form $\widehat{\Psi} = \Psi \otimes g^{1/2}(X)$, where the quantum part of $\widehat{\Psi}$ is the same state Ψ that was used to define the crossed-product. Thus, given the modular operator Δ_Ψ for a state Ψ of an algebra \mathcal{A} , there is a simple and informative formula for the modular operator $\widehat{\Delta}_{\widehat{\Psi}}$ of $\widehat{\Psi}$. We assume here that $g^{1/2}(X)$ is everywhere positive, which ensures that $\widehat{\Psi}$ is cyclic separating.

We will use the following characterization of the modular operator Δ_Ψ of an algebra \mathcal{A} with cyclic separating state Ψ . Δ_Ψ is characterized by the fact that, for all $\mathbf{a}, \mathbf{b} \in \mathcal{A}$,

$$\langle \Psi | \mathbf{a} \mathbf{b} | \Psi \rangle = \langle \Psi | \mathbf{b} \Delta_\Psi \mathbf{a} | \Psi \rangle. \quad (3.23)$$

To verify eq. (3.23), one computes

$$\begin{aligned} \langle \Psi | \mathbf{b} \Delta_\Psi \mathbf{a} | \Psi \rangle &= \langle \Psi | \mathbf{b} S_\Psi^\dagger S_\Psi \mathbf{a} | \Psi \rangle = \langle \mathbf{b}^\dagger \Psi | S_\Psi^\dagger | S_\Psi \mathbf{a} \Psi \rangle \\ &= \langle S_\Psi \mathbf{a} \Psi | S_\Psi | \mathbf{b}^\dagger \Psi \rangle = \langle \mathbf{a}^\dagger \Psi | \mathbf{b} \Psi \rangle = \langle \Psi | \mathbf{a} \mathbf{b} | \Psi \rangle. \end{aligned} \quad (3.24)$$

We used the definition of the adjoint of an antilinear operator, $\langle \Psi | S_\Psi^\dagger | \Psi' \rangle = \langle \Psi' | S_\Psi | \Psi \rangle$. We will also need the KMS condition. Set $\widehat{h}_\Psi = -\log \Delta_\Psi$, and for $\mathbf{a} \in \mathcal{A}$, let $\mathbf{a}_u = e^{\widehat{h}_\Psi u} \mathbf{a} e^{-\widehat{h}_\Psi u}$. The KMS condition is a property of functions such as $\langle \mathbf{a}_u \mathbf{b} \rangle = \langle \Psi | \mathbf{a}_u \mathbf{b} | \Psi \rangle$; if Ψ is a thermofield double state, then \mathbf{a}_u is a time translate of \mathbf{a} and these are real time thermal correlation functions. Replacing \mathbf{a} with \mathbf{a}_u in eq. (3.23) we get the KMS condition

$$\langle \Psi | \mathbf{a}_u \mathbf{b} | \Psi \rangle = \langle \Psi | \mathbf{b} \mathbf{a}_{u+i} | \Psi \rangle. \quad (3.25)$$

We used $\Delta_\Psi \Psi = \Psi$ to get $\langle \Delta_\Psi \mathbf{a}_u | \Psi \rangle = \langle \Delta_\Psi^{is} \mathbf{a}_u \Delta_\Psi^{-is} | \Psi \rangle$ with $s = -i$, and we interpreted this as $\mathbf{a}_{u+i} \Psi$. This explanation has been rather cavalier; see for example section 4.2 of [8] for more detail. A precise statement of the KMS condition is that the function $\langle \Psi | \mathbf{a}_u \mathbf{b} | \Psi \rangle$, originally defined for real u , analytically continues to a function holomorphic in the strip $-1 \leq \text{Im } u \leq 0$, and the function $\langle \Psi | \mathbf{b} \mathbf{a}_u | \Psi \rangle$, likewise initially defined for real u , analytically

continues to a function holomorphic in the strip $0 \leq \text{Im } u \leq 1$, in such a way that eq. (3.25) holds. Because $\widehat{h}_\Psi \Psi = 0$, these functions satisfy

$$\langle \Psi | \mathbf{a}_u \mathbf{b}_v | \Psi \rangle = \langle \Psi | \mathbf{a}_{u+s} \mathbf{b}_{v+s} | \Psi \rangle. \quad (3.26)$$

In the context of the thermofield double, this is time translation symmetry.

We want to find the operator $\widehat{\Delta}_{\widehat{\Psi}}$ that obeys

$$\langle \widehat{\Psi} | \widehat{\mathbf{a}} \widehat{\mathbf{b}} | \widehat{\Psi} \rangle = \langle \widehat{\Psi} | \widehat{\mathbf{b}} \widehat{\Delta}_{\widehat{\Psi}} \widehat{\mathbf{a}} | \widehat{\Psi} \rangle, \quad \forall \widehat{\mathbf{a}}, \widehat{\mathbf{b}} \in \mathcal{A} \rtimes \mathbb{R}_\Psi. \quad (3.27)$$

It turns out that

$$\widehat{\Delta}_{\widehat{\Psi}} = \Delta_\Psi \cdot g(\widehat{h}_\Psi + X)(g(X))^{-1}. \quad (3.28)$$

In proving this, it is convenient to introduce the Fourier transform of $g(X)$:

$$\begin{aligned} \widetilde{g}(w) &= \int_{-\infty}^{\infty} dX e^{iwX} g(X) \\ g(X) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dw e^{-iwX} \widetilde{g}(w). \end{aligned} \quad (3.29)$$

So another way to write the formula for $\widehat{\Delta}_{\widehat{\Psi}}$ is

$$\widehat{\Delta}_{\widehat{\Psi}} = \frac{\Delta_\Psi}{2\pi g(X)} \int_{-\infty}^{\infty} dw e^{-iw(\widehat{h}_\Psi + X)} \widetilde{g}(w). \quad (3.30)$$

The operators Δ_Ψ , \widehat{h}_Ψ , and X all commute, so there are no subtleties of operator ordering.

It is enough to verify (3.27) for operators that form an additive basis of $\mathcal{A} \rtimes \mathbb{R}_\Psi$, so we can take $\widehat{\mathbf{a}} = \mathbf{a} e^{is(\widehat{h}_\Psi + X)}$, $\widehat{\mathbf{b}} = \mathbf{b} e^{it(\widehat{h}_\Psi + X)}$, with $\mathbf{a}, \mathbf{b} \in \mathcal{A}$ and $s, t \in \mathbb{R}$. We have

$$\begin{aligned} \langle \widehat{\Psi} | \widehat{\mathbf{a}} \widehat{\mathbf{b}} | \widehat{\Psi} \rangle &= \int_{-\infty}^{\infty} dX g(X) \langle \Psi | \mathbf{a} e^{is(\widehat{h}_\Psi + X)} \mathbf{b} e^{it(\widehat{h}_\Psi + X)} | \Psi \rangle \\ &= \int_{-\infty}^{\infty} dX g(X) e^{i(s+t)X} \langle \Psi | \mathbf{a} \mathbf{b}_s | \Psi \rangle = \widetilde{g}(s+t) \langle \Psi | \mathbf{a} \mathbf{b}_s | \Psi \rangle. \end{aligned} \quad (3.31)$$

On the other hand, eq. (3.30) leads to

$$\begin{aligned} \langle \widehat{\Psi} | \widehat{\mathbf{b}} \widehat{\Delta}_{\widehat{\Psi}} \widehat{\mathbf{a}} | \widehat{\Psi} \rangle &= \int_{-\infty}^{\infty} dX \frac{1}{2\pi} \int_{-\infty}^{\infty} dw \widetilde{g}(w) \langle \Psi | \mathbf{b} e^{it(\widehat{h}_\Psi + X)} \Delta e^{-iw(\widehat{h}_\Psi + X)} \mathbf{a} e^{is(\widehat{h}_\Psi + X)} | \Psi \rangle \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} dw \widetilde{g}(w) \int_{-\infty}^{\infty} dX e^{i(s+t-w)X} \langle \Psi | \mathbf{b} \mathbf{a}_{t-w+i} | \Psi \rangle \\ &= \widetilde{g}(s+t) \langle \Psi | \mathbf{b} \mathbf{a}_{i-s} | \Psi \rangle = \widetilde{g}(s+t) \langle \Psi | \mathbf{a}_{-s} \mathbf{b} | \Psi \rangle \\ &= \widetilde{g}(s+t) \langle \Psi | \mathbf{a} \mathbf{b}_s | \Psi \rangle. \end{aligned} \quad (3.32)$$

We integrated over X and w and used the KMS condition and time translation symmetry.

This completes the proof of the formula for $\widehat{\Delta}_{\widehat{\Psi}}$.

Eq. (3.28) implies that we can factor $\widehat{\Delta}_{\widehat{\Psi}}$ as

$$\widehat{\Delta}_{\widehat{\Psi}} = \widetilde{K} K, \quad (3.33)$$

with

$$\begin{aligned}
 K &= \Delta e^{-X} g(\widehat{h}_\Psi + X) = e^{-(\widehat{h}_\Psi + X)} g(\widehat{h}_\Psi + X) \\
 \widetilde{K} &= \frac{e^X}{g(X)}.
 \end{aligned}
 \tag{3.34}$$

The point of this factorization is as follows. Since it is only a function of $\widehat{h}_\Psi + X$, K is contained in the crossed product algebra $\mathcal{A} \rtimes \mathbb{R}_\Psi$. (This statement assumes that the function $g(X)$ has the property that $g(X)e^{-X}$ is bounded; otherwise we should say that K is affiliated to $\mathcal{A} \rtimes \mathbb{R}_\Psi$, meaning that bounded functions of K are in $\mathcal{A} \rtimes \mathbb{R}_\Psi$.) Likewise \widetilde{K} is a function only of X , so it commutes with $\mathcal{A} \rtimes \mathbb{R}_\Psi$ and belongs to the commutant $(\mathcal{A} \rtimes \mathbb{R}_\Psi)'$.

Existence of such a factorization means that the modular automorphism group of $\widehat{\Psi}$ is inner. Indeed for $\mathfrak{a} \in \mathcal{A} \rtimes \mathbb{R}_\Psi$, $\widehat{\Delta}_\Psi^{\text{is}} \mathfrak{a} \widehat{\Delta}_\Psi^{-\text{is}} = K^{\text{is}} \mathfrak{a} K^{-\text{is}}$, because \mathfrak{a} commutes with \widetilde{K} . Since $K \in \mathcal{A} \rtimes \mathbb{R}_\Psi$, the automorphism $\mathfrak{a} \rightarrow K^{\text{is}} \mathfrak{a} K^{-\text{is}}$ is inner. For an algebra of Type III, the modular automorphism group is never inner. So for any \mathcal{A} , the crossed product algebra $\mathcal{A} \rtimes \mathbb{R}_\Psi$ is always of Type I or Type II.

A noteworthy fact is that the factorization in eq. (3.33) is not quite unique. If we shift $X \rightarrow X + c$, we get

$$K \rightarrow e^{-c} K, \quad \widetilde{K} \rightarrow e^c \widetilde{K}.
 \tag{3.35}$$

The group of constant shifts of X , $X \rightarrow X + c$, is an outer automorphism group of the crossed product algebra; it was already introduced in section 3.1. There is no further indeterminacy in the factorization of $\widehat{\Delta}$ provided the crossed product algebra is a factor. As already remarked, this happens precisely in the case of interest to us that \mathcal{A} is of Type III₁.

3.4 Traces

A trace on an algebra is a linear function on the algebra such that $\text{Tr } \mathfrak{a} \mathfrak{b} = \text{Tr } \mathfrak{b} \mathfrak{a}$ for any two elements $\mathfrak{a}, \mathfrak{b}$ in the algebra. For example, for any positive integer N , the algebra of $N \times N$ matrices has such a trace. A more subtle case is the Type I_∞ von Neumann algebra of all bounded operators on a separable (infinite-dimensional) Hilbert space \mathcal{H} . This algebra has a trace, but it is not defined for all elements of the algebra, only for a subalgebra consisting of operators that are “trace class.” For example, the trace of the identity element in an algebra of Type I_∞ is not defined; it would be $+\infty$.

The factorization of the modular operator for a classical-quantum state $\widehat{\Psi}$ leads immediately to the existence of a trace, albeit in general one that is defined only for a subalgebra. For $\widehat{\mathfrak{a}} \in \mathcal{A} \rtimes \mathbb{R}_\Psi$, one defines

$$\text{Tr } \widehat{\mathfrak{a}} = \langle \widehat{\Psi} | \widehat{\mathfrak{a}} K^{-1} | \widehat{\Psi} \rangle.
 \tag{3.36}$$

Making use of eq. (3.23), we get

$$\text{Tr } \widehat{\mathfrak{a}} \widehat{\mathfrak{b}} = \langle \widehat{\Psi} | \widehat{\mathfrak{a}} \widehat{\mathfrak{b}} K^{-1} | \widehat{\Psi} \rangle = \langle \widehat{\Psi} | \widehat{\mathfrak{b}} K^{-1} \widehat{\Delta}_{\widehat{\Psi}} \widehat{\mathfrak{a}} | \widehat{\Psi} \rangle.
 \tag{3.37}$$

Since $\widehat{\Psi} = \widehat{\Delta}_{\widehat{\Psi}}^{-1} \Psi$, the right hand side of eq. (3.37) is $\langle \widehat{\Psi} | \widehat{\mathfrak{b}} K^{-1} \widehat{\Delta}_{\widehat{\Psi}} \widehat{\mathfrak{a}} \widehat{\Delta}_{\widehat{\Psi}}^{-1} | \widehat{\Psi} \rangle$. Writing $\widehat{\Delta}_{\widehat{\Psi}} = K \widetilde{K}$, where \widetilde{K} commutes with K and with $\widehat{\mathfrak{a}}$, we get

$$\text{Tr } \widehat{\mathfrak{a}} \widehat{\mathfrak{b}} = \langle \widehat{\Psi} | \widehat{\mathfrak{b}} K^{-1} K \widehat{\mathfrak{a}} K^{-1} | \widehat{\Psi} \rangle = \langle \widehat{\Psi} | \widehat{\mathfrak{b}} \widehat{\mathfrak{a}} K^{-1} | \widehat{\Psi} \rangle = \text{Tr } \widehat{\mathfrak{b}} \widehat{\mathfrak{a}},
 \tag{3.38}$$

completing the proof that this function is a trace.

Using eq. (3.34) for K , along with $\Delta_\Psi \Psi = \Psi$, $\widehat{h}_\Psi |\Psi\rangle = 0$, the definition of the trace can be simplified to

$$\mathrm{Tr} \widehat{a} = \left\langle \widehat{\Psi} \left| \widehat{a} \frac{e^X}{g(X)} \right| \widehat{\Psi} \right\rangle = \int_{-\infty}^{\infty} dX e^X \langle \Psi | \widehat{a} | \Psi \rangle. \quad (3.39)$$

In writing the last formula, we view \widehat{a} as a function of X with values in operators on \mathcal{H} , so that the matrix element $\langle \Psi | \widehat{a} | \Psi \rangle$ is a function of X , which goes into the integral over X . This formula shows that the trace on $\mathcal{A} \rtimes \mathbb{R}_\Psi$ is not defined for all elements of this algebra, since the integral over X may not converge. For example, it does not converge if \widehat{a} is an element of the original algebra \mathcal{A} .

A noteworthy fact is that in the definition of the trace, the dependence on the function $g(X)$ has disappeared. In other words, we used the function $g(X)$ to define a classical-quantum state that in turn motivated the formula for the trace, but the trace that we have defined does not depend on the choice of $g(X)$. That does not mean that the definition of the trace is completely canonical. The outer automorphism of the crossed product that shifts X to $X + c$ has a simple action on the trace:

$$\mathrm{Tr} \widehat{a} \rightarrow e^c \mathrm{Tr} \widehat{a}. \quad (3.40)$$

It is no accident, of course, that the only indeterminacy of the trace is a simple rescaling. When a von Neumann algebra is a factor (its center consists only of complex scalars), a trace on this algebra, if it exists, is unique up to a scalar multiple.

Among those von Neumann algebras that are factors, Type III algebras do not admit any trace, and Type I and Type II₁ algebras admit a trace that is not rescaled by any outer automorphism. The only case of a factor that admits a trace that is rescaled by an outer automorphism is a factor of¹³ Type II_∞. This is the case of primary interest in the present article.

We can use eq. (3.39) to get a useful formula for the trace for elements of a certain subalgebra of $\mathcal{A} \rtimes \mathbb{R}_\Psi$. We recall that $\mathcal{A} \rtimes \mathbb{R}_\Psi$ contains elements $a e^{iu(\widehat{h}_\Psi + X)}$ for any $u \in \mathbb{R}$, and likewise contains linear combinations $\sum_i a_i e^{iu_i(\widehat{h}_\Psi + X)}$. Taking limits, $\mathcal{A} \rtimes \mathbb{R}_\Psi$ contains operators of the form $\widehat{a} = \int_{-\infty}^{\infty} du a(u) e^{iu(\widehat{h}_\Psi + X)}$ with a smooth function $a(u)$ that vanishes for large u . Let us impose a further condition that the function $a(u)$ is holomorphic for $0 \leq \mathrm{Im} u \leq 1$. In this case, we can get a convenient formula for $\mathrm{Tr} \widehat{a}$:

$$\mathrm{Tr} \widehat{a} = \int_{-\infty}^{\infty} dX \langle \widehat{\Psi} | \widehat{a} | \widehat{\Psi} \rangle = \int_{-\infty}^{\infty} dX du e^{X+iuX} \langle \Psi | \widehat{a}(u) | \Psi \rangle. \quad (3.41)$$

After shifting the contour in the u integral from \mathbb{R} to $i + \mathbb{R}$ and integrating first over X to get a delta function in u , we find

$$\mathrm{Tr} \widehat{a} = 2\pi \langle \Psi | a(i) | \Psi \rangle. \quad (3.42)$$

The condition for $a(u)$ to be holomorphic in the strip is rather special, but elements of $\mathcal{A} \rtimes \mathbb{R}_\Psi$ associated with functions that have that property form a subalgebra and it is

¹³See [7], section 3.6, for another explanation of why a Type II_∞ factor has a trace whose normalization cannot be canonically determined.

interesting to check the cyclic property of the trace for elements of this subalgebra. Suppose that $\widehat{\mathbf{a}} = \int_{-\infty}^{\infty} du \mathbf{a}(u) e^{iu(\widehat{h}_{\Psi} + X)}$, $\widehat{\mathbf{b}} = \int_{-\infty}^{\infty} du \mathbf{b}(u) e^{iu(\widehat{h}_{\Psi} + X)}$, where $\mathbf{a}(u)$ and $\mathbf{b}(u)$ are both holomorphic in the strip. Then defining $\mathbf{b}_u(v) = e^{i\widehat{h}_{\Psi}u} \mathbf{b}(v) e^{-i\widehat{h}_{\Psi}u}$, we get

$$\widehat{\mathbf{a}}\widehat{\mathbf{b}} = \int_{-\infty}^{\infty} du dv \mathbf{a}(u) \mathbf{b}_u(v) e^{i(u+v)(\widehat{h}_{\Psi} + X)} = \int_{-\infty}^{\infty} dw du \mathbf{a}(u) \mathbf{b}_u(w - u) e^{iw(\widehat{h}_{\Psi} + X)}. \quad (3.43)$$

So eq. (3.42) gives in this case

$$\text{Tr } \widehat{\mathbf{a}}\widehat{\mathbf{b}} = 2\pi \int_{-\infty}^{\infty} du \langle \Psi | \mathbf{a}(u) \mathbf{b}_u(i - u) | \Psi \rangle. \quad (3.44)$$

On the other hand,

$$\begin{aligned} \text{Tr } \widehat{\mathbf{b}}\widehat{\mathbf{a}} &= 2\pi \int_{-\infty}^{\infty} du \langle \Psi | \mathbf{b}(u) \mathbf{a}_u(i - u) | \Psi \rangle \\ &= 2\pi \int_{-\infty}^{\infty} du \langle \Psi | \mathbf{a}_{u-i}(i - u) \mathbf{b}(u) | \Psi \rangle \\ &= 2\pi \int_{-\infty}^{\infty} du \langle \Psi | \mathbf{a}(i - u) \mathbf{b}_{i-u}(u) | \Psi \rangle \\ &= 2\pi \int_{-\infty}^{\infty} du \langle \Psi | \mathbf{a}(u) \mathbf{b}_u(i - u) | \Psi \rangle. \end{aligned} \quad (3.45)$$

We successively used the KMS condition (3.25), translation invariance (3.26), and a change of variables $u \rightarrow i - u$.

3.5 Density matrices and entropies

Let \mathcal{A} be an algebra that acts on a Hilbert space \mathcal{H} . Suppose also that \mathcal{A} is equipped with a nondegenerate trace Tr that is positive in the sense that $\text{Tr } \mathbf{a}^\dagger \mathbf{a} > 0$ for all nonzero \mathbf{a} . Under these circumstances, one can define a density matrix. The density matrix ρ of a state $\Psi \in \mathcal{H}$ is an element $\rho \in \mathcal{A}$ such that for $\mathbf{a} \in \mathcal{A}$, $\langle \Psi | \mathbf{a} | \Psi \rangle = \text{Tr } \rho \mathbf{a}$. Such a ρ will exist because of nondegeneracy of the trace. ρ will be positive in the sense that $\text{Tr } \rho \mathbf{a}^\dagger \mathbf{a} \geq 0$ for all $\mathbf{a} \in \mathcal{A}$. It will satisfy $\text{Tr } \rho = 1$.

Such a ρ is quite analogous to a density matrix in ordinary quantum mechanics. It describes the outcomes when an arbitrary $\mathbf{a} \in \mathcal{A}$ is measured in the state Ψ .

Once one has a trace and a notion of a density matrix, one can define a version of the von Neumann entropy:

$$S(\rho) = -\text{Tr } \rho \log \rho. \quad (3.46)$$

In the case of the crossed product algebra $\mathcal{A} \rtimes \mathbb{R}_{\Psi}$, the definition of the trace is not completely canonical, since $\mathcal{A} \rtimes \mathbb{R}_{\Psi}$ has a group of outer automorphisms that conjugates X to $X + c$, $c \in \mathbb{R}$. As observed in section 3.4, this rescales the trace by $\text{Tr} \rightarrow e^c \text{Tr}$. To preserve the condition $\text{Tr } \rho = 1$, we have to compensate by rescaling the density matrix, $\rho \rightarrow e^{-c} \rho$, and this has the effect of shifting the entropy by the same constant c :

$$S(\rho) \rightarrow S(\rho) + c. \quad (3.47)$$

Thus the indeterminacy in the entropy is an overall additive constant, the same for all states. This is similar to the situation in classical physics, where entropy differences are

defined by relations such as $dE = TdS$, but there is no natural way to fix an additive constant in the entropy.

One can eliminate the arbitrary additive constant by taking the differential of eq. (3.46). This gives

$$dS = -\text{Tr} d\rho \log \rho, \quad (3.48)$$

by analogy with the classical $dE = TdS$ and also by analogy with what is sometimes called the first law of entanglement entropy [22]. Entropy differences defined this way should be physically sensible. Another analogy is with the many problems, such as those studied in [23, 24], in which entanglement entropy differences in quantum field theory are better defined than entanglement entropies.

Part of the analogy with classical physics is actually that the entropy defined in eq. (3.46) is not positive-definite, and in fact it is not bounded below.¹⁴ The entropy of a classical harmonic oscillator goes to $-\infty$ as the temperature goes to 0, because the system gets compressed in a smaller and smaller phase space volume. As explained in section 3.6 of [7], entropy in a Type II algebra, such as the crossed product algebra that we have encountered in black hole physics, can be made arbitrarily negative by disentangling qubits.

Let us compute the entropy of the classical-quantum state $\widehat{\Psi} = \Psi \otimes g(X)^{1/2}$ that we used to motivate the definition of the trace. In this case, upon replacing \widehat{a} by $\widehat{a}K$ in the definition $\text{Tr} \widehat{a} = \langle \widehat{\Psi} | \widehat{a} K^{-1} | \widehat{\Psi} \rangle$, we get $\text{Tr} \widehat{a} K = \langle \widehat{\Psi} | \widehat{a} | \widehat{\Psi} \rangle$. So the density matrix of $\widehat{\Psi}$ is just K . From eq. (3.34), we find $K \log K = e^{-(\widehat{h}_\Psi + X)} (-\widehat{h}_\Psi + X) g(\widehat{h}_\Psi + X) + \log g(\widehat{h}_\Psi + X)$. Using eq. (3.39) and recalling that $\widehat{h}_\Psi | \widehat{\Psi} \rangle = 0$, we find the entropy of the density matrix K to be

$$S(K) = \int_{-\infty}^{\infty} dX (Xg(X) - g(X) \log g(X)). \quad (3.49)$$

The normalization condition for the state $\widehat{\Psi}$ is

$$\int_{-\infty}^{\infty} dX g(X) = 1. \quad (3.50)$$

Clearly, subject to this normalization condition, $S(K)$ is unbounded above and below. If $g(X)$ is strongly peaked around a classical value X_0 , then $S(K)$ will be very close to X_0 (assuming $|X_0| \gg |\log g(X_0)|$).

In the large N limit of the thermofield double state, $g(X)$ is a Gaussian, presented in eq. (2.12) (recall $X = \beta NU$). This Gaussian is peaked near $X = 0$, not near $X = A/4G$, as we would need to get the Bekenstein-Hawking entropy. One must recall that entropy for a Type II $_\infty$ factor involves an arbitrary additive constant, so only entropy differences can really be defined. In effect, with the normalizations we have used, entropy is measured relative to the classical entropy $A/4G$ of a black hole at inverse temperature β . (Somewhat

¹⁴This statement involves taking literally that we are interested in the entropy of a state of a Type II $_\infty$ algebra, and not thinking in terms of an asymptotic expansion in $1/N$. From the standpoint of the $1/N$ expansion, one would call the entropy positive if the leading term for large N is positive, regardless of the nature of the lower order terms. This is likely to be true in physically sensible situations. The remarks in the text are applicable if we think of N as a fixed number at which we have a Type II $_\infty$ algebra; then, the entropy is unbounded below. However, the analysis in this article leading to a Type II $_\infty$ algebra is really only valid in an asymptotic expansion near $N = \infty$.

similarly, entropy in a Type II₁ algebra is defined relative to the entropy of a maximally entangled state. See for example section 3.6 of [7].) If one evaluates $S(K)$ using eq. (2.12) for g , one finds that the $\log g$ term in eq. (3.49) leads, apart from a constant, to a contribution $\frac{1}{2}\partial_\beta \log |\langle H_R \rangle|$ to $S(K)$. This term is actually a universal logarithmic correction to the Bekenstein-Hawking entropy that is associated to energy fluctuations in the canonical ensemble [25].

We have obtained in eq. (3.49) what looks like a classical formula for the entropy because of considering a special sort of classical-quantum state. For a more generic state one would certainly not get a classical formula for the entropy.

4 Other conserved charges

In addition to time translation symmetry, the two-sided eternal black hole solution of $\mathcal{N} = 4$ super Yang-Mills theory has additional rotational and gauge symmetries. The symmetry group is maximized for the case that the angular momentum and conserved charges of the black hole vanish. In this case, the rotational symmetry group is Spin(4) and the gauge group is $SU(4)_R$, which from the boundary point of view is a group of R-symmetries. The global form of the symmetry group is $G = (\text{Spin}(4) \times SU(4)_R)/\mathbb{Z}_2$. Observers on the left and right side boundaries of the black hole spacetime see separate symmetry groups G_L and G_R . In the boundary theory, these symmetry groups are generated by left and right conserved charges Q_L^a and Q_R^a , where a runs over a basis of the Lie algebra \mathfrak{g} of G .

We can proceed precisely as in the discussion of time translations in section 2. The operators Q_L^a and Q_R^a do not separately have large N limits in the thermofield double state Ψ_{TFD} . The difference¹⁵ $\hat{Q}^a = Q_R^a - Q_L^{a*}$ annihilates Ψ_{TFD} and does have a large N limit. The Q^a act on the Hilbert space \mathcal{H}_{TFD} obtained by quantizing the generalized free field theory of nonzero modes that emerges in the large N limit [1, 2] and generate an action of the group G . Let us denote the representation matrices as $W(g)$, $g \in G$.

The operators $U^a = Q_R^a/N$ also have a large N limit. They are central in the large N limit and their correlation functions are Gaussian. To get a Hilbert space on which the U^a can act in addition to the operators of the generalized free field theory, we can extend \mathcal{H}_{TFD} to $\hat{\mathcal{H}}_{\text{TFD}} = \mathcal{H}_{\text{TFD}} \otimes L^2(\mathfrak{g})$. Here $L^2(\mathfrak{g})$ is the space of square-integrable functions of the U^a , with the U^a acting by multiplication. (We also need to extend \mathcal{H}_{TFD} to accommodate time translations, as described in section 2, but we omit this here.)

This gives an adequate framework to describe the large N limit, but to go to higher orders in $1/N$ in the case of a nonabelian symmetry group, a slightly different point of view is preferable. The global symmetry group of the boundary theory is the product $G_L \times G_R$, but the eternal black hole solution is invariant only under a diagonal subgroup G_D of $G_L \times G_R$. This means that actually there is a family of classical solutions, parametrized by the quotient $(G_L \times G_R)/G_D$. This quotient is a copy of G ; we will call that copy $G_{\mathcal{M}}$ (where \mathcal{M} is for “moduli space”). Concretely, $G_{\mathcal{M}}$ parametrizes the choice of “Wilson line” between

¹⁵The left and right systems in the thermofield double are CPT conjugates. The $*$ in Q_L^{a*} is CPT, which acts by complex conjugation if the group generators are defined to be hermitian operators (or as minus complex conjugation if they are defined to be antihermitian).

the left and right boundary; thus, starting with any one given solution, representing a point that we identify with $1 \in G_{\mathcal{M}}$, we make a solution corresponding to an arbitrary $g \in G_{\mathcal{M}}$ by making a gauge transformation by an element of the gauge group (here the group of rotations and R-symmetries) that equals 1 on the left boundary and equals g on the right boundary. The group $G_L \times G_R$ acts on $G_{\mathcal{M}}$ by $g \rightarrow g_L g g_R^{-1}$, $g_L \in G_L$, $g_R \in G_R$. A given point $g \in G_{\mathcal{M}}$ is invariant under a diagonal subgroup of $G_L \times G_R$ defined by $g_L g g_R^{-1} = g$ or $g_L = g g_R g^{-1}$. We will call this group G_g .

For any choice of $g \in G_{\mathcal{M}}$, quantization of small fluctuations in the background of the eternal black hole gives a thermofield double Hilbert space that we will now call $\mathcal{H}_{\text{TFD},g}$, since it depends on g . Each $\mathcal{H}_{\text{TFD},g}$ is a representation of the subgroup G_g of $G_L \times G_R$ that is unbroken at the point $g \in G_{\mathcal{M}}$. As g varies, $\mathcal{H}_{\text{TFD},g}$ varies as the fiber of a bundle \mathcal{V} of Hilbert spaces over $G_{\mathcal{M}}$:

$$\begin{array}{ccc} \mathcal{H}_{\text{TFD},g} & \longrightarrow & \mathcal{V} \\ & & \downarrow \\ & & G_{\mathcal{M}}. \end{array}$$

An “improved” thermofield double Hilbert space $\widehat{\mathcal{H}}_{\text{TFD}}$ that incorporates the moduli in $G_{\mathcal{M}}$ can be defined as the space of L^2 sections of the Hilbert space bundle \mathcal{V} .

However, a simpler description is available. The Hilbert space bundle $\mathcal{V} \rightarrow G_{\mathcal{M}}$ is homogeneous under the action of $G_L \times G_R$ on the base space $G_{\mathcal{M}}$, and we can pick either a G_L -invariant or a G_R -invariant trivialization of this bundle. Picking a G_L -invariant trivialization can be accomplished by picking any trivialization at all over a chosen point $1 \in G_{\mathcal{M}}$ and then extending this, by acting with G_L , to a G_L -invariant trivialization over all of $G_{\mathcal{M}}$. The extension exists and is unique because for any $g \in G_{\mathcal{M}}$, there is a unique element of G_L , namely g , that maps $1 \in G_{\mathcal{M}}$ to $g \in G_{\mathcal{M}}$. This G_L -invariant trivialization is not G_R -invariant. However, following the same procedure with G_R instead of G_L , we can define a G_R -invariant trivialization that is not G_L -invariant.

Either way, once we trivialize the bundle in a G_L - or G_R -invariant fashion, there is a simple description of the Hilbert space $\widehat{\mathcal{H}}_{\text{TFD}}$ and of the action of $G_L \times G_R$ on it. When the bundle is trivialized, $\widehat{\mathcal{H}}_{\text{TFD}}$ becomes a simple tensor product $\widehat{\mathcal{H}}_{\text{TFD}} = \mathcal{H}_{\text{TFD}} \otimes L^2(G_{\mathcal{M}})$. Thus a wavefunction becomes an \mathcal{H}_{TFD} -valued function on $G_{\mathcal{M}}$. However, there are two such formalisms, depending on whether we use the left-invariant or right-invariant trivialization of \mathcal{V} . Let us write $\Psi(g)$ for the wavefunction with the left-invariant trivialization and $\chi(g)$ for the wavefunction with the right-invariant trivialization. To describe concretely the action of $G_L \times G_R$ on $\Psi(g)$ or $\chi(g)$, we will need to remember the action of the diagonal group G on \mathcal{H}_{TFD} , generated by the operators Q^a that were introduced earlier and with the representation matrices $W(g)$.

Let S_{g_L, g_R} be the operator that represents a group element $g_L \times g_R \in G_L \times G_R$ if we use the G_L -invariant trivialization with wavefunction $\Psi(g)$. Concretely, $S_{g_L, g_R}(\Psi)$ is the function on $G_{\mathcal{M}}$ defined by

$$(S_{g_L, g_R}(\Psi))(g) = W(g_R) \Psi \left(g_L^{-1} g g_R \right). \tag{4.1}$$

The idea of this formula is that since Ψ is defined with a G_L -invariant trivialization of the fibration, G_L acts only on the base space $G_{\mathcal{M}}$ of the fibration and not on the fiber

\mathcal{H}_{TFD} , but as the trivialization is not G_R -invariant, G_R acts on the fiber (via the operator $W(g_R)$) as well as on the base. The reader can verify that eq. (4.1) does define an action of $G_L \times G_R$ on $\widehat{\mathcal{H}}_{\text{TFD}}$. If instead we use a wavefunction $\chi(g)$ defined with the G_R -invariant trivialization, then the roles of G_L and G_R are reversed. The action of $G_L \times G_R$ is then described by operators T_{g_L, g_R} such that

$$(T_{g_L, g_R}(\chi))(g) = W(g_L)\chi(g_L^{-1}gg_R). \tag{4.2}$$

The relation between the two formalisms is simply $\chi(g) = W(g)\Psi(g)$, or

$$\chi = \Lambda\Psi, \tag{4.3}$$

where $\Lambda : \widehat{\mathcal{H}}_{\text{TFD}} \rightarrow \widehat{\mathcal{H}}_{\text{TFD}}$ is defined by $(\Lambda\Psi)(g) = W(g)\Psi(g)$. The equivalence of the two formalisms is defined by

$$T_{g_L, g_R} = \Lambda S_{g_L, g_R} \Lambda^{-1}. \tag{4.4}$$

Let t_L^a and t_R^a be the generators of the G_L and G_R actions on $L^2(G_{\mathcal{M}})$. Thus in view of the formulas of the last paragraph, in the formalism based on a G_L -invariant trivialization, the action of G_L on $\widehat{\mathcal{H}}_{\text{TFD}} = \mathcal{H}_{\text{TFD}} \otimes L^2(\mathbb{R})$ is generated by t_L^a , while the action of G_R is generated by $t_R^a + Q^a$. In the formalism based on a G_R -invariant trivialization, the generators are instead $t_L^a + Q^a$ and t_R^a , respectively.

Now we can explain how to incorporate the collective coordinates associated with the G symmetry in the algebra of observables outside the black hole horizon. On the right side of the horizon, for a fixed choice of the collective coordinates, we have an algebra $\mathcal{A}_{r,0}$ acting on \mathcal{H}_{TFD} . To incorporate the collective coordinates in the left-invariant formalism, we replace \mathcal{H}_{TFD} by $\widehat{\mathcal{H}}_{\text{TFD}} = \mathcal{H}_{\text{TFD}} \otimes L^2(G_{\mathcal{M}})$, and we adjoin the G_R generators $t_R^a + Q^a$ to $\mathcal{A}_{r,0}$. In other words, \mathcal{A}_R is the algebra generated by $\mathcal{A}_{r,0}$ together with $t_R^a + Q^a$. What we have just arrived at is the mathematical definition of the crossed product of the algebra $\mathcal{A}_{r,0}$ by a group of automorphisms G . We denote it as $\mathcal{A}_R = \mathcal{A}_{r,0} \rtimes G$.

As for \mathcal{A}_L , it is the commutant of \mathcal{A}_R . It is generated by t_L^a and $\Lambda^{-1}\mathcal{A}_{\ell,0}\Lambda$.

This description treats the left and right sides of the black hole asymmetrically because of using a trivialization that is G_L -invariant but not G_R -invariant. We can reverse the roles of G_L and G_R by conjugating by Λ . Then we get a description in which \mathcal{A}_L is generated by $\mathcal{A}_{\ell,0}$ and $t_L^a + Q^a$, while \mathcal{A}_R is generated by t_R^a and $\Lambda\mathcal{A}_{r,0}\Lambda^{-1}$. In all cases, regardless of the formalism, \mathcal{A}_L and \mathcal{A}_R are isomorphic to the crossed product of $\mathcal{A}_{\ell,0}$ or $\mathcal{A}_{r,0}$ by G .

When we got to essentially this point in the discussion of time translations in section 3.1, we observed that there exists a left-right symmetric formalism obtained by conjugating by $\Lambda^{1/2}$ (eq. (3.5)). In the case of a nonabelian group G , there is in no equally convenient analog of $\Lambda^{1/2}$.

Thus collective coordinates associated to an automorphism group G can always be included in the algebra of observables to the right (or left) of the eternal black hole horizon by replacing the algebra of observables $\mathcal{A}_{r,0}$ defined in a particular background with its crossed product $\mathcal{A}_R = \mathcal{A}_{r,0} \rtimes G$. However, in contrast to the case of time translation symmetry, it seems that the crossed product with a compact automorphism group does not lead to a qualitative change in the algebraic structure. In quantum field theory, for

compact G , $\mathcal{A}_{r,0} \rtimes G$ is always an algebra of Type III₁, like $\mathcal{A}_{r,0}$. A proof of this has been sketched by R. Longo. The main input is the fact that \mathcal{H}_{TFD} contains all representations of G , each with infinite multiplicity.

Acknowledgments

I am greatly indebted to R. Longo for many patient explanations and in particular for explaining the properties of the crossed product. I also thank M. R. Douglas, H. Liu, and R. Mahajan for helpful discussions, N. Lashkari for assistance with the Connes cocycle, and V. Chandrasekharan for a careful reading of the manuscript and pointing out some inaccuracies. Research supported in part by NSF Grant PHY-1911298.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

- [1] S. Leutheusser and H. Liu, *Causal connectability between quantum systems and the black hole interior in holographic duality*, [arXiv:2110.05497](https://arxiv.org/abs/2110.05497) [[INSPIRE](#)].
- [2] S. Leutheusser and H. Liu, *Emergent times in holographic duality*, [arXiv:2112.12156](https://arxiv.org/abs/2112.12156) [[INSPIRE](#)].
- [3] J.M. Maldacena, *Eternal black holes in anti-de Sitter*, *JHEP* **04** (2003) 021 [[hep-th/0106112](#)] [[INSPIRE](#)].
- [4] M. Takesaki, *Duality for Crossed Products and the Structure of von Neumann Algebras of Type III*, *Acta Math.* **131** (1973) 249.
- [5] A. Connes, *Une Classification des Facteurs de Type III*, *Ann. Sci. de L'E.N.S.* **6** (1973) 133.
- [6] A. Connes, *Noncommutative Geometry*, Academic Press, (1994) [ISBN: 9780121858605].
- [7] E. Witten, *Why Does Quantum Field Theory In Curved Spacetime Make Sense? And What Happens To The Algebra of Observables In The Thermodynamic Limit?*, [arXiv:2112.11614](https://arxiv.org/abs/2112.11614) [[INSPIRE](#)].
- [8] E. Witten, *APS Medal for Exceptional Achievement in Research: Invited article on entanglement properties of quantum field theory*, *Rev. Mod. Phys.* **90** (2018) 045003 [[arXiv:1803.04993](https://arxiv.org/abs/1803.04993)] [[INSPIRE](#)].
- [9] C.B. Thorn, *Infinite N_c QCD at finite temperature: Is there an ultimate temperature?*, *Phys. Lett. B* **99** (1981) 458 [[INSPIRE](#)].
- [10] R. Haag, N.M. Hugenholtz and M. Winnink, *On the Equilibrium states in quantum statistical mechanics*, *Commun. Math. Phys.* **5** (1967) 215 [[INSPIRE](#)].
- [11] J.J. Bisognano and E.H. Wichmann, *On the Duality Condition for Quantum Fields*, *J. Math. Phys.* **17** (1976) 303 [[INSPIRE](#)].
- [12] G.L. Sewell, *Quantum fields on manifolds: PCT and gravitationally induced thermal states*, *Annals Phys.* **141** (1982) 201 [[INSPIRE](#)].

- [13] W.G. Unruh, *Notes on black hole evaporation*, *Phys. Rev. D* **14** (1976) 870 [[INSPIRE](#)].
- [14] H.J. Borchers, *Half-Sided Translations and the Type of von Neumann Algebras*, *Lett. Math. Phys.* **44** (1998) 283.
- [15] D.L. Jafferis, A. Lewkowycz, J. Maldacena and S.J. Suh, *Relative entropy equals bulk relative entropy*, *JHEP* **06** (2016) 004 [[arXiv:1512.06431](#)] [[INSPIRE](#)].
- [16] L. Susskind and J. Uglum, *Black hole entropy in canonical quantum gravity and superstring theory*, *Phys. Rev. D* **50** (1994) 2700 [[hep-th/9401070](#)] [[INSPIRE](#)].
- [17] F. Ceyhan and T. Faulkner, *Recovering the QNEC from the ANEC*, *Commun. Math. Phys.* **377** (2020) 999 [[arXiv:1812.04683](#)] [[INSPIRE](#)].
- [18] N. Lashkari, *Modular zero modes and sewing the states of QFT*, *JHEP* **04** (2021) 189 [[arXiv:1911.11153](#)] [[INSPIRE](#)].
- [19] R. Bousso, V. Chandrasekaran and A. Shahbazi-Moghaddam, *From black hole entropy to energy-minimizing states in QFT*, *Phys. Rev. D* **101** (2020) 046001 [[arXiv:1906.05299](#)] [[INSPIRE](#)].
- [20] R. Bousso, V. Chandrasekaran, P. Rath and A. Shahbazi-Moghaddam, *Gravity dual of Connes cocycle flow*, *Phys. Rev. D* **102** (2020) 066008 [[arXiv:2007.00230](#)] [[INSPIRE](#)].
- [21] R. Longo, *A Simple Proof Of The Existence Of Modular Automorphisms In Approximately Finite Dimensional Von Neumann Algebras*, *Pacific J. Math.* **75** (1978) 199.
- [22] D.D. Blanco, H. Casini, L.-Y. Hung and R.C. Myers, *Relative Entropy and Holography*, *JHEP* **08** (2013) 060 [[arXiv:1305.3182](#)] [[INSPIRE](#)].
- [23] H. Casini and M. Huerta, *A Finite entanglement entropy and the c-theorem*, *Phys. Lett. B* **600** (2004) 142 [[hep-th/0405111](#)] [[INSPIRE](#)].
- [24] H. Casini, *Relative entropy and the Bekenstein bound*, *Class. Quant. Grav.* **25** (2008) 205021 [[arXiv:0804.2182](#)] [[INSPIRE](#)].
- [25] S. Das, P. Majumdar and R.K. Bhaduri, *General logarithmic corrections to black hole entropy*, *Class. Quant. Grav.* **19** (2002) 2355 [[hep-th/0111001](#)] [[INSPIRE](#)].