



COMPOSITIO MATHEMATICA

Green's conjecture for curves on arbitrary *K3* surfaces

Marian Aprodu and Gavril Farkas

Compositio Math. **147** (2011), 839–851.

[doi:10.1112/S0010437X10005099](https://doi.org/10.1112/S0010437X10005099)



FOUNDATION
COMPOSITIO
MATHEMATICA

*The London
Mathematical
Society*





Green’s conjecture for curves on arbitrary $K3$ surfaces

Marian Aprodu and Gavril Farkas

ABSTRACT

Green’s conjecture predicts that one can read off special linear series on an algebraic curve, by looking at the syzygies of its canonical embedding. We extend Voisin’s results on syzygies of $K3$ sections, to the case of $K3$ surfaces with arbitrary Picard lattice. This, coupled with results of Voisin and Hirschowitz–Ramanan, provides a complete solution to Green’s conjecture for smooth curves on arbitrary $K3$ surfaces.

1. Introduction

Green’s conjecture on syzygies of canonical curves asserts that one can recognize existence of special linear series on an algebraic curve, by looking at the syzygies of its canonical embedding. Precisely, if C is a smooth algebraic curve of genus g , $K_{i,j}(C, K_C)$ denotes the (i, j) th Koszul cohomology group of the canonical bundle K_C and $\text{Cliff}(C)$ is the Clifford index of C , then Green [Gre84] predicted the vanishing statement

$$K_{p,2}(C, K_C) = 0 \quad \text{for all } p < \text{Cliff}(C). \quad (1)$$

In recent years, Voisin [Voi02, Voi05] achieved a major breakthrough by showing that Green’s conjecture holds for smooth curves C lying on $K3$ surfaces S with $\text{Pic}(S) = \mathbb{Z} \cdot C$. In particular, this establishes Green’s conjecture for general curves of every genus. Using Voisin’s work, as well as a degenerate form of [HR98], it has been proved in [Apr05] that Green’s conjecture holds for any curve C of genus g of gonality $\text{gon}(C) = k \leq (g + 2)/2$, which satisfies the *linear growth condition*

$$\dim W_{k+n}^1(C) \leq n \quad \text{for } 0 \leq n \leq g - 2k + 2. \quad (2)$$

Thus Green’s conjecture becomes a question in Brill–Noether theory. In particular, one can check that condition (2) holds for a general curve $[C] \in \mathcal{M}_{g,k}^1$ in any gonality stratum of \mathcal{M}_g , for all $2 \leq k \leq (g + 2)/2$. Our main result is the following.

THEOREM 1.1. *Let S be a $K3$ surface and $C \subset S$ be a smooth curve with $g(C) = g$ and $\text{gon}(C) = k$. If $k \leq (g + 2)/2$, then C satisfies Green’s conjecture.*

Note that Theorem 1.1 has been established in [Voi02] when $\text{Pic}(S) = \mathbb{Z} \cdot C$. The proof relies (via [Apr05]) on the case of curves of odd genus of maximal gonality. Precisely, when $g(C) = 2k - 3$ and $\text{gon}(C) = k$, Green’s conjecture is due to Voisin [Voi05] combined with results of Hirschowitz–Ramanan [HR98]. Putting together these results and Theorem 1.1, we conclude the following theorem.

Received 2 December 2009, accepted in final form 15 June 2010, published online 15 February 2011.

2000 Mathematics Subject Classification 13D02, 14C20.

Keywords: syzygy, canonical curve, Brill–Noether theory.

This journal is © Foundation Compositio Mathematica 2011.

THEOREM 1.2. *Green’s conjecture holds for every smooth curve C lying on an arbitrary $K3$ surface S .*

In the proof of Theorem 1.1, we distinguish two cases. When $\text{Cliff}(C)$ is computed by a pencil (that is, $\text{Cliff}(C) = \text{gon}(C) - 2$), we use a parameter count for spaces of Lazarsfeld–Mukai bundles [CP95, Laz86], in order to find a smooth curve $C' \in |C|$, such that C' verifies condition (2). Since Koszul cohomology satisfies the Lefschetz hyperplane principle, one has that $K_{p,2}(C, K_C) \cong K_{p,2}(C', K_{C'})$. This proves Green’s conjecture for C .

When $\text{Cliff}(C)$ is no longer computed by a pencil, it follows from [CP95, Knu09] that either C is a smooth plane curve or else a *generalized ELMS example*, in the sense that there exist smooth curves $D, \Gamma \subset S$, with $\Gamma^2 = -2, \Gamma \cdot D = 1$ and $D^2 \geq 2$, such that $C \equiv 2D + \Gamma$ and $\text{Cliff}(C) = \text{Cliff}(\mathcal{O}_C(D)) = \text{gon}(C) - 3$. This case requires a separate analysis, similarly to [AP08], since condition (2) is no longer satisfied, and we refer to §5 for details.

Theorem 1.1 follows by combining results obtained by using the powerful techniques developed in [Voi02, Voi05], with facts about the effective cone of divisors of $\overline{\mathcal{M}}_g$. As pointed out in [Apr05], starting from a k -gonal smooth curve $[C] \in \mathcal{M}_g$ satisfying the Brill–Noether growth condition (2), by identifying pairs of general points $x_i, y_i \in C$ for $i = 1, \dots, g + 3 - 2k$ one creates a *stable* curve

$$[X := C/x_1 \sim y_1, \dots, x_{g+3-2k} \sim y_{g+3-2k}] \in \overline{\mathcal{M}}_{2g+3-2k}$$

having maximal gonality $g + 3 - k$, that is, lying outside the closure of the Hurwitz divisor $\mathcal{M}_{2g+3-2k, g+3-k}^1$ consisting of curves with a pencil \mathfrak{g}_{g+3-k}^1 . Since the class of the virtual failure locus of Green’s conjecture is a multiple of the Hurwitz divisor $\overline{\mathcal{M}}_{2g+3-2k, g+3-k}^1$ on $\overline{\mathcal{M}}_{2g+3-2k}$, see [HR98], Voisin’s theorem can be extended to all irreducible stable curves of genus $2g + 3 - 2k$ and having maximal gonality, in particular to X as well, and *a posteriori* to smooth curves of genus g sitting on $K3$ surfaces with arbitrary Picard lattice. On the other hand, showing that condition (2) is satisfied for a curve $[C] \in \mathcal{M}_g$, is a question of pure Brill–Noether nature.

Theorem 1.1 has strong consequences on Koszul cohomology of $K3$ surfaces. It is known that for any globally generated line bundle L on a $K3$ surface S , the Clifford index of any smooth irreducible curve is constant, equal to, say, c (see [GL87]). Applying Theorem 1.1, Green’s hyperplane section theorem, the duality theorem and finally the Green–Lazarsfeld non-vanishing theorem [Gre84], we obtain a complete description of the distribution of zeros among the Koszul cohomology groups of S with values in L .

THEOREM 1.3. *Suppose $L^2 = 2g - 2 \geq 2$. The Koszul cohomology group $K_{p,q}(S, L)$ is non-zero if and only if one of the following cases occur:*

- (i) $q = 0$ and $p = 0$; or
- (ii) $q = 1, 1 \leq p \leq g - c - 2$; or
- (iii) $q = 2$ and $c \leq p \leq g - 1$; or
- (iv) $q = 3$ and $p = g - 2$.

The analysis of the Brill–Noether loci implies also that the Green–Lazarsfeld gonality conjecture is satisfied for curves of Clifford dimension one on arbitrary $K3$ surfaces, general in their linear systems, see §4 for details.

2. Brill–Noether loci and their dimensions

Throughout this section we fix a $K3$ surface S and a globally generated line bundle $L \in \text{Pic}(S)$. We recall [Sai74], that the assumption that L be globally generated is equivalent to $|L|$ having no base components. We denote by $|L|_s$ the locus of smooth connected curves in $|L|$. For integers $r, d \geq 1$, we consider the morphism $\pi_S : \mathcal{W}_d^r(|L|) \rightarrow |L|_s$ with fibre over a point $C \in |L|_s$ isomorphic to the Brill–Noether locus $W_d^r(C)$. The analysis of the Brill–Noether loci $W_d^r(C)$, for a general curve $C \in |L|$ in its linear system, is equivalent to the analysis of the restricted maps $\pi_S : \mathcal{W} \rightarrow |L|$ over irreducible components \mathcal{W} of $\mathcal{W}_d^r(|L|)$ dominating the linear system. The main ingredient used to study $\mathcal{W}_d^r(|L|)$ is the Lazarsfeld–Mukai bundle [Laz86] associated with a complete linear series. With any pair (C, A) consisting of a curve $C \in |L|_s$ and a base-point-free linear series $A \in W_d^r(C) \setminus W_d^{r+1}(C)$, one associates the Lazarsfeld–Mukai bundle $E_{C,A} := F_{C,A}^\vee$ on S , via an elementary transformation along $C \subset S$:

$$0 \rightarrow F_{C,A} \rightarrow H^0(C, A) \otimes \mathcal{O}_S \xrightarrow{\text{ev}} A \rightarrow 0. \tag{3}$$

Dualizing the sequence (3), we obtain the short exact sequence

$$0 \rightarrow H^0(C, A)^\vee \otimes \mathcal{O}_S \rightarrow E_{C,A} \rightarrow K_C \otimes A^\vee \rightarrow 0. \tag{4}$$

The bundle $E_{C,A}$ comes equipped with a distinguished subspace of sections $H^0(C, A)^\vee \in G(r + 1, H^0(S, E_{C,A}))$. We summarize some characteristics of $E_{C,A}$.

PROPOSITION 2.1. *One has the following:*

- (i) $\det(E_{C,A}) = L$;
- (ii) $c_2(E_{C,A}) = d$;
- (iii) $h^0(S, E_{C,A}) = h^0(C, A) + h^1(C, A)$, $h^1(S, E_{C,A}) = h^2(S, E_{C,A}) = 0$;
- (iv) $\chi(S, E_{C,A} \otimes F_{C,A}) = 2(1 - \rho(g, r, d))$;
- (v) $E_{C,A}$ is globally generated off the base locus of $K_C \otimes A^\vee$.

In particular, $E_{C,A}$ is globally generated if $K_C \otimes A^\vee$ is globally generated. Conversely, if E is a globally generated bundle on S with $\text{rk}(E) = r + 1$ and $\det(E) = L$, there is a rational map $h_E : G(r + 1, H^0(S, E)) \dashrightarrow |L|$ defined in the following way. A general subspace $\Lambda \in G(r + 1, H^0(S, E))$ is mapped to the degeneracy locus of the evaluation map: $\text{ev}_\Lambda : \Lambda \otimes \mathcal{O}_S \rightarrow E$; note that, generically, this degeneracy locus cannot be the whole surface. The image $h_E(\Lambda)$ is a smooth curve $C_\Lambda \in |L|$, and we set $\text{Coker}(\text{ev}_\Lambda) := K_{C_\Lambda} \otimes A_\Lambda^\vee$, where $A_\Lambda \in \text{Pic}(C_\Lambda)$ and $\text{deg}(A_\Lambda) = c_2(E)$.

Remark 2.2. A rank- $(r + 1)$ vector bundle E on S is a Lazarsfeld–Mukai bundle if and only if $H^1(S, E) = H^2(S, E) = 0$ and there exists an $(r + 1)$ -dimensional subspace of sections $\Lambda \subset H^0(S, E)$, such that the degeneracy locus of the morphism ev_Λ is a smooth curve. In particular, being a Lazarsfeld–Mukai vector bundle is an open condition.

Coming back to the original situation when $C \in |L|_s$ and $A \in W_d^r(C) \setminus W_d^{r+1}(C)$ is globally generated, we consider the Petri map

$$\mu_{0,A} : H^0(C, A) \otimes H^0(C, K_C \otimes A^\vee) \rightarrow H^0(C, K_C),$$

whose kernel can be described in terms of Lazarsfeld–Mukai bundles. Let M_A the vector bundle of rank r on C defined as the kernel of the evaluation map

$$0 \rightarrow M_A \rightarrow H^0(C, A) \otimes \mathcal{O}_C \xrightarrow{\text{ev}} A \rightarrow 0. \tag{5}$$

Twisting (5) with $K_C \otimes A^\vee$, we obtain that $\text{Ker}(\mu_{0,A}) = H^0(C, M_A \otimes K_C \otimes A^\vee)$. Note also that there is an exact sequence on C

$$0 \rightarrow \mathcal{O}_C \rightarrow F_{C,A} \otimes K_C \otimes A^\vee \rightarrow M_A \otimes K_C \otimes A^\vee \rightarrow 0,$$

while from the defining sequence of $E_{C,A}$ one obtains the exact sequence on S

$$0 \rightarrow H^0(C, A)^\vee \otimes F_{C,A} \rightarrow E_{C,A} \otimes F_{C,A} \rightarrow F_{C,A} \otimes K_C \otimes A^\vee \rightarrow 0.$$

Since $h^0(S, F_{C,A}) = h^1(S, F_{C,A}) = 0$, one writes that

$$H^0(S, E_{C,A} \otimes F_{C,A}) = H^0(C, F_{C,A} \otimes K_C \otimes A^\vee). \tag{6}$$

We shall use the following deformation-theoretic result [Par95], which is a consequence of Sard’s theorem applied to the projection $\pi_S : \mathcal{W}_d^r(|L|) \rightarrow |L|$.

LEMMA 2.3. *Suppose $\mathcal{W} \subset \mathcal{W}_d^r(|L|)$ is a dominating component, and $(C, A) \in \mathcal{W}$ is a general element such that A is globally generated and $h^0(C, A) = r + 1$. Then the coboundary map $H^0(C, M_A \otimes K_C \otimes A^\vee) \rightarrow H^1(C, \mathcal{O}_C)$ is zero.*

The above analysis can be summarized as follows (compare with [AP08, Corollary 3.3]).

PROPOSITION 2.4. *If $\mathcal{W} \subset \mathcal{W}_d^r(|L|)$ is a dominating component, and $(C, A) \in \mathcal{W}$ is a general element such that A is globally generated and $h^0(C, A) = r + 1$, then $\dim_A W_d^r(C) \leq \rho(g, r, d) + h^0(S, E_{C,A} \otimes F_{C,A}) - 1$. Moreover, equality holds if \mathcal{W} is reduced at (C, A) .*

In particular, if $E_{C,A}$ is a simple bundle, then $\mu_{0,A}$ is injective and \mathcal{W} is reduced at (C, A) of dimension $\rho(g, r, d) + g$. Thus, the problem of estimating $\dim_A W_d^r(C)$, when $(C, A) \in \mathcal{W}$ is suitably general, can be reduced to the case when $E_{C,A}$ is not a simple bundle.

3. Varieties of pencils on K3 sections

Throughout the remaining sections we mix the additive and the multiplicative notation for divisors and line bundles. If L is a line bundle on a smooth projective variety X and $L \in \text{Pic}(X)$ is a line bundle, we write $L \geq 0$ when $H^0(X, L) \neq 0$. If E is a vector bundle on X and $L \in \text{Pic}(X)$, we set $E(-L) := E \otimes L^\vee$.

As in the previous section, we fix a K3 surface S together with a globally generated line bundle L on S . We denote by k the gonality of a general smooth curve in the linear system $|L|$, and set $g := 1 + L^2/2$. Suppose that $\rho(g, 1, k) \leq 0$ (this leaves out one single case, namely $g = 2k - 3$, when $\rho(g, 1, k) = 1$). Our aim is to prove the Koszul vanishing statement

$$K_{g-\text{Cliff}(C)-1,1}(C, K_C) = 0,$$

for any curve $C \in |L|_s$. By duality, this is equivalent to Green’s conjecture for C .

It was proved in [Apr05] that any smooth curve C that satisfies the linear growth condition (2), verifies both the Green and the Green–Lazarsfeld gonality conjectures. By comments made in the previous section, a general curve $C \in |L|_s$ satisfies (2) if and only if for any $n = 0, \dots, g - 2k + 2$, and any irreducible component $W \subset W_{k+n}^1(C)$ such that a general element $A \in W$ is globally generated, has $h^0(C, A) = 2$, and the corresponding Lazarsfeld–Mukai bundle $E_{C,A}$ is not simple, the estimate $\dim W \leq n$ holds.

Condition (2) for curves which are general in their linear system, can be verified either by applying Proposition 2.4, or by estimating directly the dimension of the corresponding irreducible

components of the scheme $\mathcal{W}_{k+n}^1(|L|)$. In our analysis, we need the following description [DM89] of non-simple Lazarsfeld–Mukai bundles, see also [CP95, Lemma 2.1].

LEMMA 3.1. *Let $E_{C,A}$ be a non-simple Lazarsfeld–Mukai bundle. Then there exist line bundles $M, N \in \text{Pic}(S)$ such that $h^0(S, M), h^0(S, N) \geq 2$, N is globally generated, and there exists a zero-dimensional, locally complete intersection subscheme ξ of S such that $E_{C,A}$ is expressed as an extension*

$$0 \rightarrow M \rightarrow E_{C,A} \rightarrow N \otimes I_\xi \rightarrow 0. \tag{7}$$

Moreover, if $h^0(S, M \otimes N^\vee) = 0$, then $\xi = \emptyset$ and the extension splits.

We say that (7) is the *Donagi–Morrison* (DM) extension associated with $E_{C,A}$.

LEMMA 3.2 (Compare with [AP08, Lemma 3.6]). *For any indecomposable non-simple Lazarsfeld–Mukai bundle E on S , the DM extension (7) is uniquely determined by E .*

Proof. We assume that two DM extensions

$$0 \rightarrow M_j \rightarrow E \rightarrow N_j \otimes I_{\xi_j} \rightarrow 0, \quad j = 1, 2,$$

are given. Observe first that $H^0(S, N_1 \otimes M_2^\vee) = H^0(S, N_2 \otimes M_1^\vee) = 0$. Indeed, if $N_1 - M_2 \geq 0$, we use $M_1 - N_1 \geq 0, M_2 - N_2 \geq 0$ (we are in the non-split case), and $M_1 + N_1 = M_2 + N_2 = L$ to get a contradiction. Then $H^0(S, (N_1 \otimes M_2^\vee) \otimes I_{\xi_1}) = H^0(S, (N_2 \otimes M_1^\vee) \otimes I_{\xi_2}) = 0$, so we obtain non-zero maps $M_1 \rightarrow M_2$ and $M_2 \rightarrow M_1$. This implies that $M_1 = M_2$. \square

Remark 3.3. Similarly, one can prove that a *decomposable* Lazarsfeld–Mukai bundle E cannot be expressed as an extension (7) with $\xi \neq \emptyset$. Thus a DM extension is always unique, up to a permutation of factors in the decomposable case. Moreover, E is decomposable if and only if the corresponding DM extension is trivial.

The size of the space of endomorphisms of a non-simple Lazarsfeld–Mukai bundle can be explicitly computed from the corresponding DM extension.

LEMMA 3.4. *Let E be a non-simple Lazarsfeld–Mukai bundle on S with $\det(E) = L$, and M and N the corresponding line bundles from the DM extension. If E is indecomposable, then*

$$h^0(S, E \otimes E^\vee) = 1 + h^0(S, M \otimes N^\vee).$$

If $E = M \oplus N$, then $h^0(S, E \otimes E^\vee) = 2 + h^0(S, M \otimes N^\vee) + h^0(S, N \otimes M^\vee)$.

Proof. The decomposable case being clear, we treat the indecomposable case. Twisting the DM extension by E^\vee and taking cohomology, we obtain the exact sequence

$$0 \rightarrow H^0(S, E^\vee(M)) \rightarrow H^0(S, E \otimes E^\vee) \rightarrow H^0(S, E^\vee(N) \otimes I_\xi).$$

Since $\det(E) = L$, it follows that $E^\vee(M) \cong E(-N)$, and $E^\vee(N) \cong E(-M)$. Therefore, one has that $h^0(S, E^\vee(N) \otimes I_\xi) = h^0(S, E(-M) \otimes I_\xi)$. Using extension (7), we claim that $h^0(S, M \otimes N^\vee) = h^0(S, E(-N))$. Indeed, if $\xi \neq \emptyset$, then $h^0(S, I_\xi) = 0$. If $\xi = \emptyset$, the image of $1 \in H^0(S, \mathcal{O}_S)$ under the map $H^0(S, \mathcal{O}_S) \rightarrow H^1(S, M \otimes N^\vee)$ is precisely the extension class, hence it is non-zero.

Observe that $H^0(S, \mathcal{O}_S) \cong H^0(S, E(-M))$, in particular, $h^0(S, E^\vee(N) \otimes I_\xi) \leq 1$. On the other hand, the morphism $H^0(S, E \otimes E^\vee) \rightarrow H^0(S, E^\vee(N) \otimes I_\xi)$ maps id_E to the arrow $E \rightarrow N \otimes I_\xi$, hence it is non-zero. It follows that $h^0(S, E^\vee(N) \otimes I_\xi) = 1$, and moreover, the map $H^0(S, E \otimes E^\vee) \rightarrow H^0(S, E^\vee(N) \otimes I_\xi)$ is surjective. \square

In order to parameterize all pairs (C, A) with non-simple Lazarsfeld–Mukai bundles, we need a global construction. We fix a non-trivial globally generated line bundle N on S with $H^0(L(-2N)) \neq 0$, and an integer $\ell \geq 0$. We set $M := L(-N)$ and $g := 1 + L^2/2$. Define $\tilde{\mathcal{P}}_{N,\ell}$ to be the family of *vector bundles* of rank two on S given by non-trivial extensions

$$0 \rightarrow M \rightarrow E \rightarrow N \otimes I_\xi \rightarrow 0, \tag{8}$$

where ξ is a zero-dimensional locally complete intersection subscheme of S of length ℓ , and set

$$\mathcal{P}_{N,\ell} := \{[E] \in \tilde{\mathcal{P}}_{N,\ell} : h^1(S, E) = h^2(S, E) = 0\}.$$

Equivalently (by Riemann–Roch), $[E] \in \mathcal{P}_{N,\ell}$ if and only if $h^0(S, E) = g - c_2(E) + 3$ and $h^1(S, E) = 0$. Note that any non-simple Lazarsfeld–Mukai bundle on S with determinant L belongs to some family $\mathcal{P}_{N,\ell}$.

Remark 3.5. Using the Cayley–Bacharach property, we observe that $\tilde{\mathcal{P}}_{N,\ell} \neq \emptyset$ whenever $\text{Ext}_S^1(N \otimes I_\xi, M) \neq 0$.

Remark 3.6. If $\mathcal{P}_{N,\ell} \neq \emptyset$, then $h^1(S, N) = 0$ and $h^0(S, N \otimes I_\xi) = h^0(S, N) - \ell$. Indeed, we choose $[E] \in \mathcal{P}_{N,\ell}$. Then

$$\begin{aligned} h^0(S, E) &= h^0(S, M) + h^0(S, N \otimes I_\xi) - h^1(S, M) \\ &\geq h^0(S, M) + h^0(S, N) - \text{length}(\xi) - h^1(S, M) \\ &\geq \chi(S, M) + \chi(S, N) - \ell = 2 + \frac{1}{2}M^2 + 2 + \frac{1}{2}N^2 - \ell \\ &= 2 + \frac{1}{2}L^2 - M \cdot N + 2 - \ell = g + 3 - c_2(E). \end{aligned}$$

Since $h^0(S, E) = g + 3 - c_2(E)$, all the inequalities are actually equalities, hence $h^1(S, N) = 0$ and $h^0(S, N \otimes I_\xi) = h^0(S, N) - \ell$.

The family $\mathcal{P}_{N,\ell}$, which, *a priori*, might be the empty set, is an open Zariski subset of a projective bundle of the Hilbert scheme $S^{[\ell]}$, as shown below.

LEMMA 3.7. *If $\xi \in S^{[\ell]}$ and $\text{Ext}_S^1(N \otimes I_\xi, M) \neq 0$, then*

$$\dim \text{Ext}_S^1(N \otimes I_\xi, M) = \ell + h^1(S, M \otimes N^\vee) - h^2(S, M \otimes N^\vee).$$

Proof. Let E be a vector bundle given by a non-trivial extension

$$0 \rightarrow M \rightarrow E \rightarrow N \otimes I_\xi \rightarrow 0.$$

Applying $\text{Hom}_S(-, M)$ to this extension, we obtain the exact sequence

$$H^0(S, \mathcal{O}_S) \rightarrow \text{Ext}_S^1(N \otimes I_\xi, M) \rightarrow H^1(S, E^\vee(M)) \rightarrow H^1(S, \mathcal{O}_S) = 0.$$

Since $1 \in H^0(S, \mathcal{O}_S)$ is mapped to the extension class of E which is non-zero, it follows that $\dim \text{Ext}_S^1(N \otimes I_\xi, M) = h^1(S, E^\vee(M)) + 1 = h^1(S, E(-N)) + 1$. We apply the identification $E^\vee(M) \cong E(-N)$ as well as the Riemann–Roch theorem for $E(-N)$ and $M - N$; note that $c_1(E(-N)) = M - N$ and $c_2(E(-N)) = \ell$ (compute the Chern classes from the defining extension twisted with N^\vee):

$$\chi(S, E(-N)) = 4 + \frac{1}{2}(M - N)^2 - \ell = 2 + \chi(S, M - N) - \ell.$$

We note that $h^0(S, E(-N)) = h^0(S, M - N)$. Indeed, if $\ell \geq 1$ then $h^0(S, I_\xi) = 0$, and if $\ell = 0$ use the fact that $1 \in H^0(S, \mathcal{O}_S)$ is mapped to the extension class through $H^0(S, \mathcal{O}_S) \rightarrow H^1(S, M - N)$. Moreover, $h^2(S, E(-N)) = h^0(S, E(-M)) = 1$, and we write

that $\chi(S, E(-M)) = 2 + h^0(S, M \otimes N^\vee) - h^1(S, M \otimes N^\vee) - \ell$, that is,

$$h^1(S, E(-N)) = \ell - 1 + h^1(S, M \otimes N^\vee) - h^2(S, M \otimes N^\vee). \quad \square$$

Assuming that $\mathcal{P}_{N,\ell} \neq \emptyset$, we consider the Grassmann bundle $\mathcal{G}_{N,\ell}$ over $\mathcal{P}_{N,\ell}$ classifying pairs (E, Λ) with $[E] \in \mathcal{P}_{N,\ell}$ and $\Lambda \in G(2, H^0(S, E))$. If $d := c_2(E)$ we define the rational map $h_{N,\ell} : \mathcal{G}_{N,\ell} \dashrightarrow \mathcal{W}_d^1(|L|)$, by setting $h_{N,\ell}(E, \Lambda) := (C_\Lambda, A_\Lambda)$, where $A_\Lambda \in \text{Pic}^d(C_\Lambda)$ is such that the following exact sequence on S holds:

$$0 \rightarrow \Lambda \otimes \mathcal{O}_S \xrightarrow{\text{ev}_\Lambda} E \rightarrow K_{C_\Lambda} \otimes A_\Lambda^\vee \rightarrow 0.$$

LEMMA 3.8. *If $\mathcal{P}_{N,\ell} \neq \emptyset$, then $\dim \mathcal{G}_{N,\ell} = g + \ell + h^0(S, M \otimes N^\vee)$.*

Proof. Let $[E] \in \mathcal{P}_{N,\ell}$. From Proposition 2.1(ii), it is clear that

$$\dim \mathcal{G}_{N,\ell} = 2\ell + \dim \mathbf{P}(\text{Ext}_S^1(N \otimes I_\xi, M)) + 2(g + 1 - c_2(E)).$$

Applying Lemma 3.7, as well as the fact that $\ell = c_2(E) - M \cdot N$, we find that

$$\begin{aligned} \dim \mathcal{G}_{N,\ell} &= 2g - 3M \cdot N + c_2(E) + 1 + h^1(S, M - N) - h^2(S, M - N) \\ &= (g + c_2(E) - M \cdot N) + (g - 2M \cdot N + 1 + h^1(S, M - N) - h^2(S, M - N)). \end{aligned}$$

From Riemann–Roch, we can write

$$\chi(S, M - N) = 2 + \frac{1}{2}(M - N)^2 = 2 + \frac{1}{2}L^2 - 2M \cdot N = g + 1 - 2M \cdot N.$$

The conclusion follows. □

LEMMA 3.9. *Assume that $\mathcal{P}_{N,\ell}$ contains a Lazarsfeld–Mukai vector bundle E on S with $c_2(E) = d$, and let $\mathcal{W} \subset \mathcal{W}_d^1(|L|)$ be the closure of the image of the rational map $h_{N,\ell} : \mathcal{G}_{N,\ell} \dashrightarrow \mathcal{W}_d^1(|L|)$. Then $\dim \mathcal{W} = g + d - M \cdot N = g + \ell$.*

Proof. Clearly \mathcal{W} is irreducible, as $\mathcal{G}_{N,\ell}$ is irreducible. If $(C, A) \in \text{Im}(h_{N,\ell})$, then the fibre $h_{N,\ell}^{-1}(C, A)$ is isomorphic to the projectivization of the space of morphisms from $E_{C,A}$ to $K_C \otimes A^\vee$. From (6), $\text{Hom}(E_{C,A}, K_C \otimes A^\vee)$ is isomorphic to $H^0(S, E_{C,A} \otimes F_{C,A})$, and has dimension $h^0(S, M \otimes N^\vee) + 1$, because of Lemma 3.4. Therefore, the general fibre of $h_{N,\ell}$ has dimension $h^0(S, M \otimes N^\vee)$. We apply now Lemma 3.8. □

LEMMA 3.10. *Suppose that a smooth curve $C \in |L|$ has Clifford dimension one and A is a globally generated line bundle on C with $h^0(C, A) = 2$ and $[E_{C,A}] \in \mathcal{P}_{N,\ell}$. Then $M \cdot N \geq \text{gon}(C)$.*

Proof. By Lemma 3.1 it follows that $M|_C$ contributes to $\text{Cliff}(C)$. From the exact sequence $0 \rightarrow N^\vee \rightarrow M \rightarrow M|_C \rightarrow 0$ and from the observation that $h^1(S, N) = 0$ (see Remark 3.6), we obtain by direct computation that

$$\text{Cliff}(M|_C) = M \cdot N + M^2 - 2h^0(S, M) + 2 = M \cdot N - 2 - 2h^1(S, M) \geq k - 2,$$

that is, $M \cdot N \geq k + 2h^1(S, M) \geq k$. □

Remark 3.11. If we drop the condition on the Clifford dimension in the hypothesis of Lemma 3.10, we obtain the inequality $M \cdot N \geq \text{Cliff}(C) + 2$.

So far, we have taken care of indecomposable non-simple Lazarsfeld–Mukai bundles, and computed the dimensions of the corresponding parameter spaces. The decomposable case is much simpler. Let $E = E_{C,A} = M \oplus N$ be a decomposable Lazarsfeld–Mukai bundle. It was

proved in [Laz89] that the differential of the natural map $h_E : G(2, H^0(S, E)) \dashrightarrow |L|_s$ at a point $[\Lambda]$, with $\Lambda = H^0(C, A)^\vee$, coincides with the multiplication map $\mu_{0,A}$. Hence, if the Grassmannian $G(2, H^0(S, E))$ dominates the linear system, the multiplication map is surjective at a general point and the corresponding irreducible components of the Brill–Noether loci are zero-dimensional. This case can occur only if the Brill–Noether number is non-negative.

All these intermediate results amount to the following.

THEOREM 3.12. *Let S be a K3 surface and L a globally generated line bundle on S , such that general curves in $|L|$ are of Clifford dimension one. Suppose that $\rho(g, 1, k) \leq 0$, where $L^2 = 2g - 2$ and k is the (constant) gonality of all curves in $|L|_s$. Then a general curve $C \in |L|$ satisfies the linear growth condition (2), thus Green’s conjecture is verified for any smooth curve in $|L|$.*

In the case $\rho(g, 1, k) = 1$, Green’s conjecture is also verified for smooth curves in $|L|$, cf. [HR98, Voi05]. To sum up, Green’s conjecture is verified for any curve of Clifford dimension one on a K3 surfaces.

Proof. It suffices to estimate the dimension of dominating irreducible components \mathcal{W} of $\mathcal{W}_{k+n}^1(|L|)$, with $n = 0, \dots, g - k + 2$, with general point corresponding to a non-simple indecomposable Lazarsfeld–Mukai bundle. Lemmas 3.9 and 3.10 yield $\dim \mathcal{W} \leq g + n$, which finishes the proof. \square

Remark 3.13. The proof of Theorem 3.12 shows that for $d > g - k + 2$, every dominating component of $\mathcal{W}_d^1(|L|)$ corresponds to simple Lazarsfeld–Mukai bundles. In particular, for a general curve $C \in |L|$, one has $\dim W_d^1(C) = \rho(g, 1, d)$.

Remark 3.14. The problem of deciding whether Lazarsfeld–Mukai bundles appear in a given space $\mathcal{P}_{N,\ell}$ is a non-trivial one, cf. Remark 2.2.

4. A criterion for the Green–Lazarsfeld gonality conjecture

Along with Green’s conjecture, another statement of similar flavor was proposed by Green and Lazarsfeld, [GL86].

CONJECTURE 4.1 (The gonality conjecture). For any smooth curve C of gonality d , every non-special globally generated line bundle L on C of sufficiently high degree satisfies $K_{h^0(L)-d,1}(C, L) = 0$.

Conjecture 4.1 is equivalent to the seemingly weaker statement that *there exists* a globally generated line bundle $L \in \text{Pic}(C)$ with $h^1(C, L) = 0$ for which the Koszul vanishing holds [Apr02]. On a curve C with the linear growth condition property (2), line bundles of type $K_C(x + y)$, where $x, y \in C$ are general points, verify the gonality conjecture [Apr05]. In particular, Theorem 3.12 implies the following.

COROLLARY 4.2. *Let S be a K3 surface and L a globally generated line bundle on S , such that general curves in $|L|$ are of Clifford dimension one. Then a general curve $C \in |L|$ verifies Conjecture 4.1.*

The main result of this short section is a refinement of the main result of [Apr05].

THEOREM 4.3. *Let C be a smooth curve of Clifford dimension one and $x, y \in C$ be distinct points, and denote*

$$\mathcal{Z}_n := \{A \in W_{k+n}^1(C) : h^0(C, A(-x - y)) \geq 1\}.$$

Suppose that $\dim \mathcal{Z}_n \leq n - 1$, for all $0 \leq n \leq g - 2k + 2$. Then the bundle $K_C(x + y)$ verifies the gonality conjecture.

The condition in the statement of Theorem 4.3 means that passing through the points x and y is a non-trivial condition on any irreducible component of maximal allowed dimension n of the Brill–Noether locus $W_{k+n}^1(C)$, for all $0 \leq n \leq g - 2k + 2$.

Proof. The proof is an almost verbatim copy of the proof of [Apr05, Theorem 2]. Define $\nu := g - 2k + 2$. The idea is to show that for any $0 \leq n \leq \nu$, and for $(n + 1)$ pairs of distinct general points $x_0 + y_0, x_1 + y_1, \dots, x_n + y_n \in C_2$, there is no line bundle $A \in W_{k+n}^1(C)$ with $h^0(C, A(-x_i - y_i)) \neq 0$ for all $1 \leq i \leq n$, such that either $h^0(C, A(-x - y)) \neq 0$ or $h^0(C, A(-x_0 - y_0)) \neq 0$. To this end, consider the incidence varieties

$$\left(\prod_{i=1}^n C_2\right) \times \mathcal{Z}_n \supset \{(x_1 + y_1, \dots, x_n + y_n, A) : h^0(A(-x_i - y_i)) \neq 0, \forall i\},$$

respectively,

$$\left(\prod_{i=1}^{n+1} C_2\right) \times W_{k+n}^1(C) \supset \{(x_0 + y_0, x_1 + y_1, \dots, x_n + y_n, A) : h^0(A(-x_i - y_i)) \neq 0, \forall i\}.$$

The fibres of the projection to \mathcal{Z}_n are n -dimensional, hence the incidence variety is at most $(2n - 1)$ -dimensional and it cannot dominate $\prod_{i=1}^n C_2$. Similarly, the second variety is at most $(2n + 1)$ -dimensional. Note that the condition to pass through a pair of general points is a non-trivial condition on every variety of complete pencils. To conclude, apply [Apr05, Proposition 8]. \square

5. Curves of higher Clifford dimension

We analyze the Koszul cohomology of curves of higher Clifford dimension on a $K3$ surface S . This case has similarities to [AP08], where one focused on $K3$ surfaces with Picard number 2. Since plane curves are known to verify Green's conjecture, the significant cases occur when the Clifford dimension is at least three. Note that, unlike the Clifford index, the Clifford dimension is *not* semi-continuous. An example was given by Donagi–Morrison [DM89]: if $\epsilon : S \rightarrow \mathbf{P}^2$ is a double sextic and $L = \epsilon^*(\mathcal{O}_{\mathbf{P}^2}(3))$, then the general element in $|L|$ is isomorphic to a smooth plane sextic, hence it has Clifford dimension two, while special points correspond to bielliptic curves and are of Clifford dimension one.

It was proved in [CP95, Knu09] that, except for the Donagi–Morrison example, if a globally generated linear system $|L|$ on S contains smooth curves of Clifford dimension at least two, then $L = \mathcal{O}_S(2D + \Gamma)$, where $D, \Gamma \subset S$ are smooth curves, $D^2 \geq 2$ (hence $h^0(S, \mathcal{O}_S(D)) \geq 2$), $\Gamma^2 = -2$ and $D \cdot \Gamma = 1$; the case when L is ample is treated in [CP95], whereas the general case when L is globally generated is settled in [Knu09]. If the genus of D is $r \geq 3$, then the genus of a smooth curve $C \in |L|$ equals $4r - 2 \geq 10$, and $\text{gon}(C) = 2r$, while $\text{Cliff}(C) = 2r - 3$; the Clifford dimension of C is r . From now on, we assume that we are in this situation.

Green's hyperplane section theorem implies that the Koszul cohomology is constant in a linear system. As in [AP08], we degenerate a smooth curve $C \in |2D + \Gamma|$ to a reducible curve $X + \Gamma$ with $X \in |2D|$. In order to be able to carry out this plan, we first analyze the geometry of the curves in $|2D|$. Notably, we shall prove the following theorem.

THEOREM 5.1. *The hypotheses of Theorem 4.3 are verified for a general curve $X \in |2D|$ and the two points of intersection $X \cdot \Gamma$.*

The proof of Theorem 5.1 proceeds in several steps. The first result describes the fundamental invariants of a quadratic complete intersection section of S .

LEMMA 5.2. *Any smooth curve $X \in |2D|$ has genus $4r - 3$, gonality $2r - 2$, and $\text{Cliff}(X) = 2r - 4$.*

Proof. Since $\text{Cliff}(D|_X) = 2r - 4$, we obtain $\text{Cliff}(X) \leq 2r - 4 < (g(X) - 1)/2$, that is, $\text{Cliff}(X)$ is computed by a line bundle $B \in \text{Pic}(S)$, cf. [GL87]. Both bundles B and $B' := \mathcal{O}_S(X) \otimes B^\vee$ are globally generated, hence $B \cdot \Gamma \geq 0$ and $B' \cdot \Gamma \geq 0$. Since $X \cdot \Gamma = 2$, we can assume that $B \cdot \Gamma \leq 1$. We may also assume, cf. [Mar89, Corollary 2.3] that $h^0(S, B) = h^0(X, B|_X)$ and $h^0(S, B') = h^0(X, B'|_X)$. Then if $C \in |L|$ is smooth as above, we obtain the estimate

$$\text{Cliff}(X) = B \cdot X - 2h^0(S, B) + 2 \geq B \cdot C - 2h^0(C, B|_C) + 1 \geq 2r - 4.$$

Since X has Clifford dimension one, it follows that $\text{gon}(X) = 2r - 2$. □

It suffices therefore to analyze the structure of the loci $W^1_{2r-2+n}(X)$ where $n \leq 3 = g(X) - 2 \text{gon}(X) + 2$, and more precisely those components of dimension n .

LEMMA 5.3. *We fix a general $X \in |2D|$, viewed as a half-canonical curve $X \xrightarrow{|D|} \mathbf{P}^r$.*

- *Then $W^1_{2r-2}(X)$ is finite and all minimal pencils \mathfrak{g}^1_{2r-2} on X are given by the rulings of quadrics of rank four in $H^0(\mathbf{P}^r, \mathcal{I}_{X/\mathbf{P}^r}(2))$.*
- *Then X has no base-point-free pencils \mathfrak{g}^1_{2r-1} , that is, $W^1_{2r-1}(X) = X + W^1_{2r-2}(X)$.*
- *For $n = 2, 3$, if $A \in W^1_{2r-2+n}(X)$ is a base-point-free pencil, then the vector bundle $E_{X,A}$ is not simple.*

In all cases for $n \leq 3$, if A belongs to an n -dimensional component of $W^1_{2r-2+n}(X)$, then the corresponding DM extension

$$0 \rightarrow M \rightarrow E_{X,A} \rightarrow N \otimes I_\xi \rightarrow 0$$

verifies $\text{length}(\xi) = n$, $M \cdot N = 2r - 2$ and $M \cdot \Gamma = N \cdot \Gamma = 1$. When $n = 2, 3$, we can take $M = N = \mathcal{O}_X(D)$.

Proof. We use Accola’s lemma, cf. [ELMS89, Lemma 3.1]. If $A \in W^1_{2r-2+n}(X)$ is base point free with $n \leq 3$, then $h^0(X, \mathcal{O}_X(D) \otimes A^\vee) \geq 2 - n/2$. In particular, when $n = 0, 1$, we find that $A' := \mathcal{O}_X(D) \otimes A^\vee$ is a pencil as well. When $n = 1$, we find that $A' \in W^1_{2r-3}(X)$, which is impossible, that is, X carries no base-point-free pencils \mathfrak{g}^1_{2r-1} . If $n = 0$, then $\text{deg}(A) = \text{deg}(A') = 2r - 2$ and this corresponds to a quadric $Q \in H^0(\mathbf{P}^r, \mathcal{I}_{X/\mathbf{P}^r}(2))$ with $\text{rk}(Q) = 4$ and $X \cap \text{Sing}(Q) = \emptyset$, such that the rulings of Q cut out on X , precisely the pencils A and A' respectively. If $n = 2, 3$, we find that $h^0(X, K_X(-2A)) \neq 0$, thus the kernel of the Petri map $\text{Ker } \mu_{0,A} = H^0(X, K_X(-2A)) \neq 0$, and then the Lazarsfeld–Mukai bundle $E_{X,A}$ cannot be simple.

The vector bundle $E = E_{X,A}$ is thus expressible as a DM extension

$$0 \rightarrow M \rightarrow E_{X,A} \rightarrow N \otimes I_\xi \rightarrow 0,$$

and we recall that N is globally generated with $h^1(S, N) = 0$. Suppose first that $n \neq 0$, thus $n \in \{2, 3\}$. Then $h^0(S, M \otimes N^\vee) \neq 0$, for otherwise $\xi = \emptyset$, the extension is split, and the split case only produces zero-dimensional components of the Brill–Noether loci, whilst we are in the

higher-dimensional case. From the exact sequence defining $E_{X,A}$ coupled with Accola's lemma, we obtain the isomorphisms

$$H^0(S, E(-D)) \cong H^0(X, K_X(-A) \otimes \mathcal{O}_X(-D)) = H^0(X, \mathcal{O}_X(D - A)) \neq 0,$$

therefore $H^0(S, M(-D)) \cong H^0(S, E(-D)) \neq 0$. Choose an effective divisor $F \in |M(-D)|$ and then $F \in |\mathcal{O}_S(D)(-N)|$ as well. From Lemmas 3.9 and 3.10 and the generality assumption on X , we find that $\text{length}(\xi) = n$, $M \cdot N = \text{gon}(X) = 2r - 2$ and $h^1(S, M) = 0$. Furthermore, one computes that $F^2 = 0$. Since, by degree reasons, $h^0(S, \mathcal{O}_S(F)) = h^0(X, \mathcal{O}_X(D - A)) = 1$ one obtains that $F \equiv 0$, that is, $M = N = \mathcal{O}_X(D)$.

If $n = 0$, then in the associated DM extension, $\xi = 0$, and $M, N \in \text{Pic}(S)$ are globally generated, $M \cdot N = 2r - 2$ and $h^0(M) = h^0(M|_X)$ and $h^0(N) = h^0(N|_X)$. The intersection of Γ with one of the bundles, M or N , is ≤ 1 ; suppose $M \cdot \Gamma \leq 1$. We choose a smooth curve $C \in |2D + \Gamma|$, and compute

$$\text{Cliff}(M|_C) = M \cdot C - 2h^0(M|_C) + 2 \leq M \cdot X - 2h^0(M) + 2 + 1 = \text{Cliff}(X) + 1,$$

hence M computes $\text{Cliff}(C)$ and $M \cdot \Gamma = N \cdot \Gamma = 1$. □

LEMMA 5.4. *Let $X \in |2D|$ be any smooth curve, and $x, y \in X \cdot \Gamma$. For any integer $n \geq 0$, and any base-point-free pencil $A \in W_{2r-2+n}^1(X)$, the following are equivalent:*

- (i) $h^0(X, A(-x - y)) \neq 0$;
- (ii) $E_{X,A}|_\Gamma \cong \mathcal{O}_\Gamma \oplus \mathcal{O}_\Gamma(2)$.

Proof. The non-vanishing of $H^0(X, A(-x - y))$ is equivalent to

$$h^0(X, A(-x - y)) = 1. \tag{9}$$

Twisting the defining exact sequence of $E_{X,A}$ by $\mathcal{O}_S(\Gamma)$, we obtain

$$0 \rightarrow H^0(X, A)^\vee \otimes \mathcal{O}_S(\Gamma) \rightarrow E_{X,A} \otimes \mathcal{O}_S(\Gamma) \rightarrow K_X \otimes A^\vee(x + y) \rightarrow 0. \tag{10}$$

By Riemann–Roch, $H^1(S, \mathcal{O}_S(\Gamma)) = 0$. By taking cohomology, $h^0(C, A(-x - y)) = 1$ if and only if $h^0(S, E_{X,A} \otimes \mathcal{O}_S(\Gamma)) = 2r + 3 - n$. On the other hand, $h^0(S, E_{X,A}) = 2 + h^1(X, A) = 2r + 2 - n$. Consider the (twisted) exact sequence defining Γ :

$$0 \rightarrow E_{X,A} \rightarrow E_{X,A} \otimes \mathcal{O}_S(\Gamma) \rightarrow E_{X,A}|_\Gamma(-2) \rightarrow 0.$$

We find by taking cohomology that $x, y \in C$ lie in the same fibre of $|A|$ if and only if $h^0(\Gamma, E_{X,A}|_\Gamma(-2)) = 1$. Expressing $E_{X,A}|_\Gamma = \mathcal{O}_\Gamma(a) \oplus \mathcal{O}_\Gamma(b)$, with $a + b = 2$, condition (9) becomes equivalent to $E_{X,A}|_\Gamma \cong \mathcal{O}_\Gamma \oplus \mathcal{O}_\Gamma(2)$. □

Proof of Theorem 5.1. For a base-point-free $A \in W_{2r-2+n}^1(X)$ with $n = 0, 2, 3$, the vector bundle $E := E_{X,A}$ appears as an extension

$$0 \rightarrow M \rightarrow E \rightarrow N \otimes I_\xi \rightarrow 0,$$

with $\text{length}(\xi) = n$, and $M \cdot \Gamma = N \cdot \Gamma = 1$. Recall that being a Lazarsfeld–Mukai bundle is an open condition in any flat family of bundles, Remark 2.2. Hence, LM bundles in a parameter space $\mathcal{P}_{N,n}$ correspond to general cycles $\xi \in S^{[n]}$. For $n = 0$, we see immediately that $E|_\Gamma = \mathcal{O}_\Gamma(1)^{\oplus 2}$. For $n = 2, 3$ the same is true for ξ such that $\xi \cap \Gamma = \emptyset$. To conclude, we apply Lemma 5.4. □

Remark 5.5. The variety $\mathcal{W}_{2r-2}^1(|2D|)$ is birationally equivalent to the parameter space of pairs (Q, Π) , where $Q \in |\mathcal{O}_{\mathbf{P}^r}(2)|$ is a quadric of rank four and $\Pi \subset Q$ is a ruling. In particular, $\mathcal{W}_{2r-2}^1(|2D|)$ is irreducible (and of dimension g).

Theorems 5.1 and 4.3 imply the following.

COROLLARY 5.6. *For a general curve $X \in |2D|$, we have $K_{2r,1}(X, K_X \otimes \mathcal{O}_S(\Gamma)) = 0$.*

The main result of this section is (compare with [AP08]) the following theorem.

THEOREM 5.7. *Smooth curves of Clifford dimension at least three on K3 surfaces satisfy Green's conjecture.*

Proof. As in [AP08, §4.1], for all $p \geq 1$, we have isomorphisms

$$K_{p,1}(X + \Gamma, \omega_{X+\Gamma}) \cong K_{p,1}(X, K_X(\Gamma)).$$

Corollary 5.6 shows that $K_{2r,1}(X + \Gamma, \omega_{X+\Gamma}) = 0$, implying the vanishing of $K_{2r,1}(S, L)$, via Green's hyperplane section theorem. Using the hyperplane section theorem again, we obtain $K_{2r,1}(C, K_C) = 0$, for any smooth curve $C \in |L|$, that is, the vanishing predicted by Green's conjecture for C . \square

Theorems 3.12 and 5.7 altogether complete the proof of Theorem 1.1.

ACKNOWLEDGEMENTS

Research of the first author was partly supported by a PN-II-ID-PCE-2008-2 grant (cod 1189, contract no 530) and by a resumption of a Humboldt Research Fellowship. Research of both authors was partly supported by the Sonderforschungsbereich 'Raum-Zeit-Materie'. MA thanks HU Berlin for the kind hospitality and the excellent atmosphere during the preparation of this work.

REFERENCES

- Apr02 M. Aprodu, *On the vanishing of higher syzygies of curves*, Math. Z. **241** (2002), 1–15.
- Apr05 M. Aprodu, *Remarks on syzygies of d -gonal curves*, Math. Res. Lett. **12** (2005), 387–400.
- AN08 M. Aprodu and J. Nagel, *Koszul cohomology and algebraic geometry*, University Lecture Series, vol. 52 (American Mathematical Society, Providence, RI, 2010).
- AP08 M. Aprodu and G. Pacienza, *The Green conjecture for exceptional curves on a K3 surface*, Int. Math. Res. Not. IMRN (2008), 25.
- AV03 M. Aprodu and C. Voisin, *Green–Lazarsfeld's conjecture for generic curves of large gonality*, C. R. Math. Acad. Sci. Paris **36** (2003), 335–339.
- CP95 C. Ciliberto and G. Pareschi, *Pencils of minimal degree on curves on a K3 surface*, J. Reine Angew. Math. **460** (1995), 15–36.
- DM89 R. Donagi and D. R. Morrison, *Linear systems on K3-sections*, J. Differential Geom. **29** (1989), 49–64.
- ELMS89 D. Eisenbud, H. Lange, G. Martens and F.-O. Schreyer, *The Clifford dimension of a projective curve*, Compositio Math. **72** (1989), 173–204.
- Gre84 M. Green, *Koszul cohomology and the geometry of projective varieties*, J. Differential Geom. **19** (1984), 125–171.
- GL86 M. Green and R. Lazarsfeld, *On the projective normality of complete linear series on an algebraic curve*, Invent. Math. **83** (1986), 73–90.
- GL87 M. Green and R. Lazarsfeld, *Special divisors on curves on a K3 surface*, Invent. Math. **89** (1987), 357–370.
- HR98 A. Hirschowitz and S. Ramanan, *New evidence for Green's conjecture on syzygies of canonical curves*, Ann. Sci. École Norm. Sup. (4) **31** (1998), 145–152.

- Knu09 A. Knutsen, *On two conjectures for curves on $K3$ surfaces*, *Internat. J. Math.* **20** (2009), 1547–1560.
- Laz86 R. Lazarsfeld, *Brill–Noether–Petri without degenerations*, *J. Differential Geom.* **23** (1986), 299–307.
- Laz89 R. Lazarsfeld, *A sampling of vector bundle techniques in the study of linear series*, in *Proceedings of the first college on Riemann surfaces*, Trieste, November 9–December 18, 1987, eds M. Cornalba *et al.* (World Scientific, Teaneck, NJ, 1989), 500–559.
- Mar89 G. Martens, *On curves on $K3$ surfaces*, in *Algebraic curves and projective geometry, Trento, 1988*, *Lecture Notes in Mathematics*, vol. 1389 (Springer, Berlin–New York, 1989), 174–182.
- Par95 G. Pareschi, *A proof of Lazarsfeld's theorem on curves on $K3$ surfaces*, *J. Algebraic Geom.* **4** (1995), 195–200.
- Sai74 D. Saint-Donat, *Projective models of $K3$ surfaces*, *Amer. J. Math.* **96** (1974), 602–639.
- Voi02 C. Voisin, *Green's generic syzygy conjecture for curves of even genus lying on a $K3$ surface*, *J. Eur. Math. Soc. (JEMS)* **4** (2002), 363–404.
- Voi05 C. Voisin, *Green's canonical syzygy conjecture for generic curves of odd genus*, *Compositio Math.* **141** (2005), 1163–1190.

Marian Aprodu aprodu@imar.ro

Institute of Mathematics 'Simion Stoilow' of the Romanian Academy, RO-014700 Bucharest, Romania

and

Școala Normală Superioară București, Calea Griviței 21, Sector 1, RO-010702 Bucharest, Romania

Gavril Farkas farkas@math.hu-berlin.de

Humboldt-Universität zu Berlin, Institut Für Mathematik, 10099 Berlin, Germany