

 Open access • Journal Article • DOI:10.1103/PHYSREVD.72.014016

## Gribov parameter and the dimension two gluon condensate in Euclidean Yang-Mills theories in the Landau gauge — [Source link](#)

David Dudal, R. F. Sobreiro, S.P. Sorella, Henri Verschelde

**Institutions:** Ghent University, Rio de Janeiro State University

**Published on:** 25 Jul 2005 - Physical Review D (American Physical Society)

Related papers:

- [Quantization of non-Abelian gauge theories](#)
- [Renormalizability of the critical limit of lattice gauge theory by BRS invariance](#)
- [Landau gauge within the Gribov horizon.](#)
- [Local and renormalizable action from the gribov horizon](#)
- [New features of the gluon and ghost propagator in the infrared region from the Gribov-Zwanziger approach](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/gribov-parameter-and-the-dimension-two-gluon-condensate-in-4wanc0jd9n>

# The Gribov parameter and the dimension two gluon condensate in Euclidean Yang-Mills theories in the Landau gauge

D. Dudal<sup>a\*</sup>, R.F. Sobreiro<sup>b†</sup>, S.P. Sorella<sup>b‡§</sup>, H. Verschelde<sup>a¶</sup>

<sup>a</sup> *Ghent University  
Department of Mathematical Physics and Astronomy  
Krijgslaan 281-S9, B-9000 Gent, Belgium*

<sup>b</sup> *UERJ, Universidade do Estado do Rio de Janeiro  
Rua São Francisco Xavier 524, 20550-013 Maracanã  
Rio de Janeiro, Brasil*

## Abstract

The local composite operator  $A_\mu^2$  is added to the Zwanziger action, which implements the restriction to the Gribov region  $\bar{\Omega}$  in Euclidean Yang-Mills theories in the Landau gauge. We prove that Zwanziger's action with the inclusion of the operator  $A_\mu^2$  is renormalizable to all orders of perturbation theory, obeying the renormalization group equations. This allows to study the dimension two gluon condensate  $\langle A_\mu^2 \rangle$  by the local composite operator formalism when the restriction to the Gribov region  $\bar{\Omega}$  is taken into account. The resulting effective action is evaluated at one-loop order in the  $\overline{\text{MS}}$  scheme. We obtain explicit values for the Gribov parameter and for the mass parameter due to  $\langle A_\mu^2 \rangle$ , but the expansion parameter turns out to be rather large. Furthermore, an optimization of the perturbative expansion in order to reduce the dependence on the renormalization scheme is performed. The properties of the vacuum energy, with or without the inclusion of the condensate  $\langle A_\mu^2 \rangle$ , are investigated. In particular, it is shown that in the original Gribov-Zwanziger formulation, i.e. without the inclusion of the operator  $A_\mu^2$ , the resulting vacuum energy is always positive at one-loop order, independently from the choice of the renormalization scheme and scale. In the presence of  $\langle A_\mu^2 \rangle$ , we are unable to come to a definite conclusion at the order considered. In the  $\overline{\text{MS}}$  scheme, we still find a positive vacuum energy, again with a relatively large expansion parameter, but there are renormalization schemes in which the vacuum energy is negative, albeit the dependence on the scheme itself appears to be strong. Concerning the behaviour of the gluon and ghost propagators, we recover the well known consequences of the restriction to the Gribov region, and this in the presence of  $\langle A_\mu^2 \rangle$ , i.e. an infrared suppression of the gluon propagator and an enhancement of the ghost propagator. Such a behaviour is in qualitative agreement with the results obtained from the studies of the Schwinger-Dyson equations and from lattice simulations.

---

\*Research Assistant of the Research Foundation - Flanders (FWO-Vlaanderen).

†sobreiro@uerj.br

‡sorella@uerj.br

§Work supported by FAPERJ, Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro, under the program *Cientista do Nosso Estado*, E-26/151.947/2004.

¶david.dudal@ugent.be, henri.verschelde@ugent.be

# 1 Introduction.

The dimension two condensate  $\langle A_\mu^2 \rangle$  has received a great deal of attention in the last few years, see for example [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17]. This condensate was already introduced in [18] in order to analyze the gluon propagator within the Operator Product Expansion (OPE), while in [19] the condensate  $\langle A_i^2 \rangle$  was considered in the Coulomb gauge. A renormalizable effective potential for  $\langle A_\mu^2 \rangle$  has been constructed and evaluated in analytic form up to two-loop order in the Landau gauge within the local composite operator (LCO) formalism in [3, 12]. The output of these investigations is that a non-vanishing condensate is favoured as it lowers the vacuum energy. The renormalizability of the local composite operator formalism, see [20] for an introduction to the method, was proven to all orders of perturbation theory, in the case of  $\langle A_\mu^2 \rangle$ , in [11] using the algebraic renormalization technique [21]. Besides the Landau gauge, the method was extended to other gauges as, for instance, the Curci-Ferrari gauge [22, 23], the linear covariant gauges [24, 25] and, more recently, the maximal Abelian gauge [26].

As a consequence of the existence of a non-vanishing condensate  $\langle A_\mu^2 \rangle$ , a dynamical mass parameter for the gluons can be generated in the gauge fixed Lagrangian, see [3, 12, 25]. We mention that a gluon mass has been proven to be rather useful in the phenomenological context, see e.g. [27, 28, 29]. Moreover, mass parameters are commonly used in the fitting formulas for the data obtained in lattice simulations, where the gluon propagator has been studied to a great extent in the Landau gauge [30, 31, 32, 33, 34, 35, 36].

The lattice results so far obtained have provided firm evidence of the suppression of the gluon propagator in the infrared region, in the Landau gauge. Next to the gluon propagator, also the ghost propagator has been investigated numerically on the lattice [34, 35, 36, 37, 38], exhibiting an infrared enhancement. It is worth remarking that, in agreement with lattice results, this infrared behaviour of the gluon as well as of the ghost propagator has been obtained in the analysis of the Schwinger-Dyson equations, see [39, 40, 41, 42, 43, 44, 45, 46], as well as in a study making use of the exact renormalization group technique [47].

The aim of the present work is to investigate further the condensation of the operator  $A_\mu^2$  in the Landau gauge using the local composite operator formalism. This will be done by taking into account the nonperturbative effects related to the existence of the Gribov ambiguities [48], which are known to affect the Landau gauge fixing condition,  $\partial_\mu A_\mu^a = 0$ . As a consequence of the existence of the Gribov copies, the domain of integration in the path integral has to be restricted in a suitable way. Gribov's original proposal was to restrict the domain of integration to the region  $\Omega$  whose boundary  $\partial\Omega$  is the first Gribov horizon, where the first vanishing eigenvalue of the Faddeev-Popov operator,  $-\partial_\mu (\partial_\mu \delta^{ab} + gf^{acb} A_\mu^c)$ , appears [48]. Within the region  $\Omega$  the Faddeev-Popov operator is positive definite, i.e.  $-\partial_\mu (\partial_\mu \delta^{ab} + gf^{acb} A_\mu^c) > 0$ . One of the main results of Gribov's work [48] was that the gluon, respectively ghost propagator, got suppressed, respectively enhanced, in the infrared due to the restriction to the region  $\Omega$ .

In two previous papers [49, 50], we have already worked out the consequences of the restriction to the Gribov region  $\Omega$  when the dynamical generation of a gluon mass parameter due to  $\langle A_\mu^2 \rangle$  takes place, also finding an infrared suppression of the gluon and an enhancement of the ghost propagator. In [49], we closely followed the setup of Gribov's paper [48]. In this work, we shall rely on the Zwanziger local formulation of the Gribov horizon. In a series of papers

[51, 52], Zwanziger has been able to implement the restriction to the Gribov region  $\Omega$  through the introduction of a nonlocal horizon function appearing in the Boltzmann weight defining the Euclidean Yang-Mills measure. More precisely, according to [51, 52], the starting Yang-Mills measure in the Landau gauge is given by

$$d\mu_\gamma = DA\delta(\partial_\mu A_\mu^a) \det(\mathcal{M})e^{-(S_{\text{YM}}+\gamma^4 H)} , \quad (1.1)$$

where

$$\mathcal{M}^{ab} = -\partial_\mu \left( \partial_\mu \delta^{ab} + g f^{acb} A_\mu^c \right) , \quad (1.2)$$

$$S_{\text{YM}} = \frac{1}{4} \int d^4x F_{\mu\nu}^a F_{\mu\nu}^a , \quad (1.3)$$

and

$$H = \int d^4x h(x) = g^2 \int d^4x f^{abc} A_\mu^b (\mathcal{M}^{-1})^{ad} f^{dec} A_\mu^e , \quad (1.4)$$

is the so-called horizon function, which implements the restriction to the Gribov region. Notice that  $H$  is nonlocal. The parameter  $\gamma$ , known as the Gribov parameter, has the dimension of a mass and is not free, being determined by the horizon condition

$$\langle h(x) \rangle = 4(N^2 - 1) , \quad (1.5)$$

where the expectation value  $\langle h(x) \rangle$  has to be evaluated with the measure  $d\mu_\gamma$ . To the first order, the horizon condition (1.5) reads, in  $d$  dimensions,

$$1 = \frac{N(d-1)}{4} g^2 \int \frac{d^d k}{(2\pi)^d} \frac{1}{k^4 + 2Ng^2\gamma^4} . \quad (1.6)$$

This equation coincides with the original gap equation derived by Gribov for the parameter  $\gamma$  [48].

Albeit nonlocal, the horizon function  $H$  can be localized through the introduction of a suitable set of additional fields. As shown in [51, 52, 53], the resulting local action turns out to be renormalizable to all orders of perturbation theory. Remarkably, we shall be able to prove that this feature is preserved when the local operator  $A_\mu^2$  is introduced in the Zwanziger action. Moreover, the resulting theory turns out to obey a homogeneous renormalization group equation. These important properties will allow us to study the condensation of the operator  $A_\mu^2$  within a local renormalizable framework when the restriction to the Gribov region  $\Omega$  is implemented.

It is worth remarking that the Gribov region is not free from gauge copies [54, 55, 56, 57], i.e. Gribov copies still exist inside  $\Omega$ . To avoid the presence of these additional copies, a further restriction to a smaller region  $\Lambda$ , known as the fundamental modular region, should be implemented. At present, a clear understanding of the role played by these additional copies appears to be a very difficult task. Nevertheless, we should mention that, recently, it has been argued in [58] that the additional copies existing inside  $\Omega$  could have no influence on the expectation values, so that averages calculated over  $\Lambda$  or  $\Omega$  might give the same value.

The paper is organized as follows. In section 2, we give a short account of how the nonlocal horizon functional  $H$  can be localized by means of the introduction of additional fields. In section 3, we prove the renormalizability, to all orders of perturbation theory, of Zwanziger's action in the presence of the operator  $A_\mu^2$ , introduced through the local composite operator formalism. As the model has a rich symmetry structure, translated into several Ward identities,

it turns out that only three independent renormalization factors are necessary. The resulting quantum effective action obeys a homogeneous renormalization group equation, as explicitly verified at one-loop order. From this effective action, two coupled gap equations, associated to the condensate  $\langle A_\mu^2 \rangle$  and to the Gribov parameter  $\gamma$ , are derived. Section 4 is devoted to the study of these gap equations at one-loop order in the  $\overline{\text{MS}}$  renormalization scheme. It is worth mentioning that, under certain conditions, we find that it is possible that the condensate  $\langle A_\mu^2 \rangle$  is positive when the horizon condition is imposed. We prove that in the  $\overline{\text{MS}}$  scheme, and at one-loop order, the solution of the gap equations is necessarily one with  $\langle A_\mu^2 \rangle > 0$ . We recall that without the restriction to the Gribov region  $\Omega$ , the value found for  $\langle A_\mu^2 \rangle$  using the local composite operator formalism is negative, see [3, 12, 25]. Let us also mention here that in [6, 7, 8, 9], a positive estimate for  $\langle A_\mu^2 \rangle$  was obtained when using the OPE in combination with  $\langle A_\mu^2 \rangle$ . These works were based on the observation of a certain discrepancy at relatively large momentum between the expected perturbative behaviour and the obtained lattice behaviour of e.g. the effective strong coupling constant and gluon propagator. This discrepancy could be accounted for by power corrections in  $\frac{1}{q^2}$ , due to a positive  $\langle A_\mu^2 \rangle_{\text{OPE}}$  gluon condensate. The presence of such power corrections has also been discussed in [59]. We do not know if there is a direct connection between the condensate  $\langle A_\mu^2 \rangle$  that we determine, and  $\langle A_\mu^2 \rangle_{\text{OPE}}$ , as the latter is expected to contain only infrared contributions, according to an OPE treatment, while the gap equations fixing the gluon condensate and the Gribov parameter are evaluated using perturbation theory, implying that reliable results are only to be expected at a sufficiently large scale.

Although the expansion parameter proves to be rather large, an attempt to obtain explicit values for the Gribov and gluon mass parameter is still presented. Also, we shall prove that in the original Gribov-Zwanziger model, the vacuum energy is always positive at one-loop order, irrespective of the choice of renormalization scheme and scale. We outline the importance of the sign of the vacuum energy, as it is related to the gauge invariant gluon condensate  $\langle F_{\mu\nu}^2 \rangle$ , via the trace anomaly. From

$$\theta_{\mu\mu} = \frac{\beta(g^2)}{2g^2} F_{\mu\nu}^2, \quad (1.7)$$

the vacuum energy can be traced back to the value of the gluon condensate  $\langle F_{\mu\nu}^2 \rangle$ . In particular, for  $N = 3$ , from this anomaly one deduces

$$\left\langle \frac{g^2}{4\pi^2} F_{\mu\nu}^2 \right\rangle = -\frac{32}{11} E_{\text{vac}}, \quad (1.8)$$

where the one-loop  $\beta$ -function has been used. Hence, a positive vacuum energy implies a negative value for the condensate  $\left\langle \frac{g^2}{4\pi^2} F_{\mu\nu}^2 \right\rangle$ . This is in contradiction with what is found. In QCD, with quarks present, one can extract phenomenological values for  $\left\langle \frac{g^2}{4\pi^2} F_{\mu\nu}^2 \right\rangle$  via the sum rules [60], obtaining positive values for this condensate. It was discussed in [61] how to obtain an estimate for it by means of lattice calculations. In the case of  $N = 3$  Yang-Mills theory without quarks, it was found that

$$\left\langle \frac{g^2}{4\pi^2} F_{\mu\nu}^2 \right\rangle = 0.14 \pm 0.02 \text{GeV}^4. \quad (1.9)$$

Let us mention here that the Yang-Mills  $\beta$ -function is negative up to the (known) four-loop order [62, 63, 64]. Hence,  $E_{\text{vac}}$  and  $\left\langle \frac{g^2}{4\pi^2} F_{\mu\nu}^2 \right\rangle$  will continue to have opposite sign at higher order. From this viewpoint, it seems to us that it would be an asset that the vacuum energy obtained from any kind of calculation is at least negative.

In section 5 we present an optimized expansion in order to reduce the dependence on the choice of renormalization scheme to a single parameter  $b_0$ , related to the coupling constant renormalization. This is achieved by exchanging the mass parameters by their renormalization scale and scheme invariant counterparts and by re-expanding the series in the one-loop coupling constant. For  $b_0 = 0$ , which corresponds to the  $\overline{\text{MS}}$  scheme, we find a positive  $\langle A_\mu^2 \rangle$ , positive  $E_{\text{vac}}$  and hence negative  $\langle F_{\mu\nu}^2 \rangle$ . However, we find that a region of  $b_0$  is existing in which the vacuum energy is negative, but unfortunately the dependence on  $b_0$  in this region happens to be very large. A higher order analysis seems to be required to reach more definite conclusions about the sign of  $\langle A_\mu^2 \rangle$ ,  $E_{\text{vac}}$  or  $\langle F_{\mu\nu}^2 \rangle$ .

For the benefit of the reader, we provide in section 6 an overview of some important consequences stemming from the presence of the Gribov and gluon mass parameters on the gluon and ghost propagators. We point out a particular renormalization property of the Zwanziger action in order to ensure the enhancement of the ghost propagator. Conclusions are written down in section 7, while the technical details of our work have been collected in the Appendices A and B.

## 2 Local action from the restriction to the Gribov region.

As explained in [51, 52], the nonlocal functional  $H$  can be localized by means of the introduction of a suitable set of additional ghost fields. More precisely, for the localized version of the measure  $d\mu_\gamma$  we get,

$$d\mu_\gamma = DADbDcD\bar{c}D\varphi D\bar{\varphi}D\omega D\bar{\omega}e^{-S}, \quad (2.1)$$

where  $S$  is given by<sup>1</sup>

$$S = S_0 - \gamma^2 g \int d^4x \left( f^{abc} A_\mu^a \varphi_\mu^{bc} + f^{abc} A_\mu^a \bar{\varphi}_\mu^{bc} \right), \quad (2.2)$$

while

$$\begin{aligned} S_0 = S_{\text{YM}} &+ \int d^4x \left( b^a \partial_\mu A_\mu^a + \bar{c}^a \partial_\mu (D_\mu c)^a \right) \\ &+ \int d^4x \left( \bar{\varphi}_\mu^{ac} \partial_\nu \left( \partial_\nu \varphi_\mu^{ac} + g f^{abm} A_\nu^b \varphi_\mu^{mc} \right) - \bar{\omega}_\mu^{ac} \partial_\nu \left( \partial_\nu \omega_\mu^{ac} + g f^{abm} A_\nu^b \omega_\mu^{mc} \right) \right. \\ &\left. - g \left( \partial_\nu \bar{\omega}_\mu^{ac} \right) f^{abm} \left( D_\nu c \right)^b \varphi_\mu^{mc} \right). \end{aligned} \quad (2.3)$$

The fields  $(\bar{\varphi}_\mu^{ac}, \varphi_\mu^{ac})$  are a pair of complex conjugate bosonic fields. Each field has  $4(N^2 - 1)^2$  components. Similarly, the fields  $(\bar{\omega}_\mu^{ac}, \omega_\mu^{ac})$  are anticommuting. The local action (2.2) is renormalizable by power counting. More precisely, it has been shown in [51, 52, 53] that the Green functions obtained with the action  $S_0$  with the insertion of the local composite operators  $f^{abc} A_\mu^a \varphi_\mu^{bc}$  and  $f^{abc} A_\mu^a \bar{\varphi}_\mu^{bc}$  are renormalizable, the action  $S_0$  being indeed renormalizable by a multiplicative renormalization of the coupling constant  $g$  and of the fields [51, 52, 53]. We remark that the action  $S_0$  displays a global  $U(f)$  symmetry,  $f = 4(N^2 - 1)$ , with respect to the composite index  $i = (\mu, c) = 1, \dots, f$ , of the additional fields  $(\bar{\varphi}_\mu^{ac}, \varphi_\mu^{ac}, \bar{\omega}_\mu^{ac}, \omega_\mu^{ac})$ . Setting

$$(\bar{\varphi}_\mu^{ac}, \varphi_\mu^{ac}, \bar{\omega}_\mu^{ac}, \omega_\mu^{ac}) = (\bar{\varphi}_i^a, \varphi_i^a, \bar{\omega}_i^a, \omega_i^a), \quad (2.4)$$

---

<sup>1</sup>Our conventions are different from those originally used by Zwanziger. These can be obtained from ours by setting  $\varphi \rightarrow -\varphi$  and  $\omega \rightarrow -\omega$ .

we get

$$\begin{aligned}
S_0 &= S_{\text{YM}} + \int d^4x \left( b^a \partial_\mu A_\mu^a + \bar{c}^a \partial_\mu (D_\mu c)^a \right) \\
&+ \int d^4x \left( \bar{\varphi}_i^a \partial_\nu (D_\nu \varphi_i)^a - \bar{\omega}_i^a \partial_\nu (D_\nu \omega_i)^a - g (\partial_\nu \bar{\omega}_i^a) f^{abm} (D_\nu c)^b \varphi_i^m \right). \quad (2.5)
\end{aligned}$$

For the  $U(f)$  invariance we have

$$\begin{aligned}
U_{ij} S_0 &= 0, \\
U_{ij} &= \int d^4x \left( \varphi_i^a \frac{\delta}{\delta \varphi_j^a} - \bar{\varphi}_j^a \frac{\delta}{\delta \bar{\varphi}_i^a} + \omega_i^a \frac{\delta}{\delta \omega_j^a} - \bar{\omega}_j^a \frac{\delta}{\delta \bar{\omega}_i^a} \right). \quad (2.6)
\end{aligned}$$

The presence of the global  $U(f)$  invariance means that one can make use of the composite index  $i = (\mu, c)$ . By means of the diagonal operator  $Q_f = U_{ii}$ , the  $i$ -valued fields turn out to possess an additional quantum number. As shown in [51, 52, 53], the action  $S_0$  is left invariant by the following nilpotent BRST transformations,

$$\begin{aligned}
sA_\mu^a &= -(D_\mu c)^a, \\
sc^a &= \frac{1}{2} g f^{abc} c^b c^c, \\
s\bar{c}^a &= b^a, \quad sb^a = 0, \\
s\varphi_i^a &= \omega_i^a, \quad s\omega_i^a = 0, \\
s\bar{\omega}_i^a &= \bar{\varphi}_i^a, \quad s\bar{\varphi}_i^a = 0, \quad (2.7)
\end{aligned}$$

with

$$sS_0 = 0. \quad (2.8)$$

For further use, the quantum numbers of all fields entering the action  $S_0$  are displayed in Table 1. It is worth noticing that, when  $f^{abc} A_\mu^a \varphi_\mu^{bc}$  and  $f^{abc} A_\mu^a \bar{\varphi}_\mu^{bc}$  are treated as composite operators, they are introduced in the starting action  $S_0$  coupled to local external sources  $M_\mu^{ai}, V_\mu^{ai}$ , namely

$$- \int d^4x \left( M_\mu^{ai} (D_\mu \varphi_i)^a + V_\mu^{ai} (D_\mu \bar{\varphi}_i)^a \right). \quad (2.9)$$

The horizon condition (1.5) is thus obtained from the quantum action by requiring that, at the end of the computation, the sources  $M_\mu^{ai}, V_\mu^{ai}$  attain the physical values, obtained by setting

$$M_{\mu\nu}^{ab} = V_{\mu\nu}^{ab} = \gamma^2 \delta^{ab} \delta_{\mu\nu}. \quad (2.10)$$

Indeed, expression (2.9) reduces precisely to that of eq.(2.2) when the sources  $M_\mu^{ai}, V_\mu^{ai}$  attain their physical value.

	$A_\mu^a$	$c^a$	$\bar{c}^a$	$b^a$	$\varphi_i^a$	$\bar{\varphi}_i^a$	$\omega_i^a$	$\bar{\omega}_i^a$
dimension	1	0	2	2	1	1	1	1
ghostnumber	0	1	-1	0	0	0	1	-1
$Q_f$ -charge	0	0	0	0	1	-1	1	-1

Table 1: Quantum numbers of the fields.

	$U_\mu^{ai}$	$M_\mu^{ai}$	$N_\mu^{ai}$	$V_\mu^{ai}$	$\eta$	$\tau$	$K_\mu^a$	$L^a$
dimension	2	2	2	2	2	2	3	4
ghostnumber	-1	0	1	0	-1	0	-1	-2
$Q_f$ -charge	-1	-1	1	1	0	0	0	0

Table 2: Quantum numbers of the sources.

### 3 Renormalization of the Zwanziger action in the presence of the composite operator $A_\mu^a A_\mu^a$ .

The purpose of this section is to show that the renormalizability of the local action  $S_0$  is preserved when, besides the operators  $f^{abc} A_\mu^a \varphi_\mu^{bc}$  and  $f^{abc} A_\mu^a \bar{\varphi}_\mu^{bc}$ , also the local composite operator  $A_\mu^a A_\mu^a$  is introduced. This is a remarkable feature of the Zwanziger action, allowing us to discuss the condensation of the operator  $A_\mu^a A_\mu^a$  when the restriction to the Gribov region  $\Omega$  is implemented. To discuss the renormalizability of the model in the presence of  $A_\mu^2$ , we start from the following complete action

$$\Sigma = S_0 + S_s + S_{\text{ext}} , \quad (3.1)$$

where  $S_s$  is the term containing all needed local composite operators with their respective local sources, and is given by

$$S_s = s \int d^4x \left( -U_\mu^{ai} (D_\mu \varphi_i)^a - V_\mu^{ai} (D_\mu \bar{\omega}_i)^a - U_\mu^{ai} V_\mu^{ai} + \frac{1}{2} \eta A_\mu^a A_\mu^a - \frac{1}{2} \zeta \tau \eta \right) , \quad (3.2)$$

where the BRST operator acts as

$$\begin{aligned} sU_\mu^{ai} &= M_\mu^{ai} , & sM_\mu^{ai} &= 0 , \\ sV_\mu^{ai} &= N_\mu^{ai} , & sN_\mu^{ai} &= 0 , \end{aligned} \quad (3.3)$$

and

$$s\eta = \tau , \quad s\tau = 0 . \quad (3.4)$$

Therefore, for  $S_s$  one gets

$$\begin{aligned} S_s &= \int d^4x \left( -M_\mu^{ai} (D_\mu \varphi_i)^a - gU_\mu^{ai} f^{abc} (D_\mu c)^b \varphi_i^c + U_\mu^{ai} (D_\mu \omega_i)^b \right. \\ &\quad - N_\mu^{ai} (D_\mu \bar{\omega}_i)^a - V_\mu^{ai} (D_\mu \bar{\varphi}_i)^a + gV_\mu^{ai} f^{abc} (D_\mu c)^b \bar{\omega}_i^c \\ &\quad \left. - M_\mu^{ai} V_\mu^{ai} + U_\mu^{ai} N_\mu^{ai} + \frac{1}{2} \tau A_\mu^a A_\mu^a + \eta A_\mu^a \partial_\mu c^a - \frac{1}{2} \zeta \tau^2 \right) . \end{aligned} \quad (3.5)$$

As already noticed, the sources  $M_\mu^{ai}$ ,  $V_\mu^{ai}$  are needed to introduce the composite operators  $(D_\mu \varphi_i)^a$  and  $(D_\mu \bar{\varphi}_i)^a$ . The sources  $U_\mu^{ai}$ ,  $N_\mu^{ai}$  define the BRST variations of these operators, given by  $(D_\mu \omega_i)^b$  and  $(D_\mu \bar{\omega}_i)^a$ . The physical value of these sources is given by

$$\begin{aligned} M_{\mu\nu}^{ab} &= V_{\mu\nu}^{ab} = \gamma^2 \delta^{ab} \delta_{\mu\nu} , \\ U_{\mu\nu}^{ab} &= N_{\mu\nu}^{ab} = 0 . \end{aligned} \quad (3.6)$$

The local composite operator  $A_\mu^a A_\mu^a$  and its BRST variation,  $A_\mu^a \partial_\mu c^a$ , are then introduced by means of the local sources  $\tau$ ,  $\eta$ . We also notice that the complete action  $\Sigma$  contains terms quadratic in the external sources, namely  $(M_\mu^{ai} V_\mu^{ai} - U_\mu^{ai} N_\mu^{ai})$  and  $\zeta \tau^2$ . These terms, allowed by power counting, are in fact needed for the multiplicative renormalizability of the model. As



shown in [3], the dimensionless LCO parameter  $\zeta$  of the quadratic term in the source  $\tau$  is needed to account for the divergences present in the correlation function  $\langle A_\mu^2(x)A_\nu^2(y) \rangle$  for  $x \rightarrow y$ . It should be remarked that, unlike for the term quadratic in the external source  $\tau$ , we have not introduced a new free parameter for the quadratic term  $(M_\mu^{ai}V_\mu^{ai} - U_\mu^{ai}N_\mu^{ai})$  in expression (3.5). As we shall see, this term goes through the renormalization without the need of introducing a new parameter for its renormalizability. This is a remarkable feature of the Zwanziger action which plays an important role when the ghost propagator in the presence of the Gribov horizon will be discussed, see section 6.

Finally, the term  $S_{\text{ext}}$  is the source term needed to define the nonlinear BRST transformations of the gauge and ghost fields, i.e.

$$S_{\text{ext}} = \int d^4x \left( -K_\mu^a (D_\mu c)^a + \frac{1}{2} g L^a f^{abc} c^b c^c \right). \quad (3.7)$$

The technical details concerning the algebraic renormalization procedure have been worked out in Appendix A. In summary, the Zwanziger action in the presence of the local operator  $A_\mu^a A_\mu^a$  is multiplicative renormalizable. In turn, this ensures that the quantum effective action obeys the homogeneous renormalization group equations (RGE). This is an important feature of the model, which will be useful when we shall try to obtain estimates for both the Gribov and mass parameter.

The effective action is defined upon setting the sources  $U_{\mu\nu}^{ab}$ ,  $N_{\mu\nu}^{ab}$ ,  $K_\mu^a$ ,  $L^a$  and  $\eta$  equal to zero and implementing the condition (2.10). Doing so, we get

$$\begin{aligned} S &= S_0 + S_\gamma + \int d^4x \left( \frac{\tau}{2} A_\mu^a A_\mu^a - \frac{\zeta}{2} \tau^2 \right), \\ S_\gamma &= \int d^4x \left[ -\gamma^2 g f^{abc} A_\mu^a \varphi_\mu^{bc} - \gamma^2 g f^{abc} A_\mu^a \bar{\varphi}_\mu^{bc} - 4(N^2 - 1) \gamma^4 \right]. \end{aligned} \quad (3.8)$$

The term  $-4(N^2 - 1) \gamma^4$  originates from the quadratic term in the external sources, namely  $(-M_\mu^{ai}V_\mu^{ai} + U_\mu^{ai}N_\mu^{ai})$ , in expression (3.5), evaluated at the physical values given by eq.(2.10).

Following [3, 12, 20, 25], we introduce a Hubbard-Stratonovich field  $\sigma$  by means of the following unity

$$1 = \int [d\sigma] e^{-\frac{1}{2\zeta} \int d^4x \left[ \frac{\sigma}{g} + \frac{1}{2} A_\mu^a A_\mu^a - \zeta \tau \right]^2}, \quad (3.9)$$

to remove the term proportional to  $\tau^2$ . The source  $\tau$  is henceforth linearly coupled to the field  $\sigma$ , as can be directly seen from the action, which now reads

$$\begin{aligned} S &= S_0 + S_\gamma + S_\sigma + \int d^4x \left( -\tau \frac{\sigma}{g} \right), \\ S_\sigma &= \frac{\sigma^2}{2g^2\zeta} + \frac{1}{2} \frac{g\sigma}{g^2\zeta} A_\mu^a A_\mu^a + \frac{1}{8\zeta} (A_\mu^a A_\mu^a)^2. \end{aligned} \quad (3.10)$$

The following identification is easily derived [3, 12, 20, 25]

$$\langle A_\mu^a A_\mu^a \rangle = -\frac{1}{g} \langle \sigma \rangle, \quad (3.11)$$

from which it follows that a nonvanishing vacuum expectation value of the field  $\sigma$  will result in a nonvanishing condensate  $\langle A_\mu^a A_\mu^a \rangle$ .

The quantum action  $\Gamma$  is obtained through the definition

$$e^{-\Gamma} = \int [d\Phi] e^{-S_0 - S_\gamma - S_\sigma} , \quad (3.12)$$

where  $\Phi$  is a shorthand notation for all the relevant fields.

The value for  $\langle \sigma \rangle$  is found through the minimization condition

$$\frac{\partial \Gamma}{\partial \sigma} = 0 . \quad (3.13)$$

The horizon is implemented by the condition [51, 52].

$$\frac{\partial \Gamma}{\partial \gamma^2} = 0 . \quad (3.14)$$

Let us show this here. The following equivalence is readily found

$$\frac{\partial \Gamma}{\partial \gamma^2} = 0 \Leftrightarrow \left\langle g f^{abc} A_\mu^a \varphi_\mu^{bc} \right\rangle + \left\langle g f^{abc} A_\mu^a \overline{\varphi}_\mu^{bc} \right\rangle = -8 (N^2 - 1) \gamma^2 , \quad (3.15)$$

From expressions (1.1) and (2.2), it follows that

$$-2\gamma^2 \langle h \rangle = \left\langle g f^{abc} A_\mu^a \varphi_\mu^{bc} \right\rangle + \left\langle g f^{abc} A_\mu^a \overline{\varphi}_\mu^{bc} \right\rangle . \quad (3.16)$$

The combination of eq.(3.15) with eq.(3.16) gives rise to the horizon condition eq.(1.5). In order to conclude this, it is tacitly assumed that  $\gamma \neq 0$ . We notice that the condition (3.14) does possess the solution  $\gamma = 0$ . This is an artefact of the reformulation of the horizon condition in terms of the equation (3.14), and must be excluded as it does not lead to the horizon condition (1.5). We shall, however, continue to keep this solution of the gap equation (3.14), as  $\gamma \equiv 0$  corresponds to the case where the restriction to the Gribov region  $\Omega$  would not be implemented. In this case, we must only solve the gap equation stemming from eq.(3.13) with  $\gamma \equiv 0$ .

The original Gribov-Zwanziger model, i.e. without the inclusion of the operator  $A_\mu^2$ , is obtained by only retaining the condition (3.14) with  $\sigma \equiv 0$ .

Up to now, the LCO parameter  $\zeta$  is still a free parameter of the theory. We do not intend here to give a complete overview of the LCO formalism, we suffice by saying that  $\zeta$  is fixed by the demand that the action  $\Gamma$  should obey the homogeneous renormalization group equation

$$\left( \overline{\mu} \frac{\partial}{\partial \overline{\mu}} + \beta(g^2) \frac{\partial}{\partial g^2} + \gamma_{\gamma^2}(g^2) \gamma^2 \frac{\partial}{\partial \gamma^2} + \gamma_\sigma(g^2) \sigma \frac{\partial}{\partial \sigma} \right) \Gamma = 0 , \quad (3.17)$$

with

$$\begin{aligned} \overline{\mu} \frac{\partial g^2}{\partial \overline{\mu}} &= \beta(g^2) , \\ \overline{\mu} \frac{\partial \gamma^2}{\partial \overline{\mu}} &= \gamma_{\gamma^2}(g^2) \gamma^2 , \\ \overline{\mu} \frac{\partial \sigma}{\partial \overline{\mu}} &= \gamma_\sigma(g^2) \sigma . \end{aligned} \quad (3.18)$$

This can be accommodated for by making  $\zeta$  a function of the running coupling constant  $g^2$ , in which case it is found that

$$\zeta(g^2) = \frac{\zeta_0}{g^2} + \zeta_1 + \zeta_2 g^2 + \dots . \quad (3.19)$$

We refer to the available literature [3, 12, 20, 22, 25, 26] for a detailed account of the LCO formalism.

### 3.1 Renormalization group invariance of the one-loop effective action in the $\overline{\text{MS}}$ scheme without the inclusion of $A_\mu^2$ .

Before proceeding with the detailed analysis of the horizon condition in the presence of the local operator  $A_\mu^a A_\mu^a$ , let us first derive the horizon condition and check the explicit renormalization group invariance of the quantum action  $\Gamma$  by switching off the source  $\tau$  coupled to the operator  $A_\mu^a A_\mu^a$ . This amounts to consider the original Gribov-Zwanziger model. We consider thus the action

$$S = S_0 + S_\gamma . \quad (3.20)$$

The one-loop effective action  $\Gamma^{(1)}$  is easily obtained from the quadratic part of eq.(3.20)

$$e^{-\Gamma_\gamma^{(1)}} = \int [D\Phi] e^{-S_{\text{quad}}} , \quad (3.21)$$

with  $S_{\text{quad}}$  given by

$$\begin{aligned} S_{\text{quad}} = & \int d^4x \left[ \frac{1}{4} (\partial_\mu A_\nu^a - \partial_\nu A_\mu^a)^2 + \frac{1}{2\alpha} (\partial_\mu A_\mu^a)^2 + \overline{\varphi}_\mu^{ab} \partial^2 \varphi_\mu^{ab} \right. \\ & \left. - \gamma^2 g \left( f^{abc} A_\mu^a \varphi_\mu^{bc} + f^{abc} A_\mu^a \overline{\varphi}_\mu^{bc} \right) - 4(N^2 - 1)\gamma^4 \right] , \end{aligned} \quad (3.22)$$

where the limit  $\alpha \rightarrow 0$  is understood in order to recover the Landau gauge. After a straightforward computation, one gets

$$\Gamma^{(1)} = -4(N^2 - 1)\gamma^4 + \frac{(N^2 - 1)}{2} (d - 1) \int \frac{d^d p}{(2\pi)^d} \ln(p^4 + 2Ng^2\gamma^4) . \quad (3.23)$$

Dimensional regularization, with  $d = 4 - \varepsilon$ , will be employed throughout this work. Taking the derivative of  $\Gamma^{(1)}$ , one reobtains the original gap equation for the Gribov parameter  $\gamma$ , namely

$$\frac{\partial \Gamma^{(1)}}{\partial \gamma} = 0 \Rightarrow 1 = \frac{N(d-1)}{4} g^2 \int \frac{d^d p}{(2\pi)^d} \frac{1}{(p^4 + 2Ng^2\gamma^4)} . \quad (3.24)$$

More precisely, recalling that

$$\int \frac{d^d p}{(2\pi)^d} \ln(p^4 + \rho^2) = -\frac{\rho^2}{32\pi^2} \left( \ln \frac{\rho^2}{\mu^4} - 3 \right) + \frac{1}{\varepsilon} \frac{4\rho^2}{32\pi^2} , \quad (3.25)$$

the one-loop effective action  $\Gamma^{(1)}$  reads

$$\Gamma^{(1)} = -4(N^2 - 1)\gamma^4 - \frac{3(N^2 - 1)}{64\pi^2} (2Ng^2\gamma^4) \left( \ln \frac{2Ng^2\gamma^4}{\mu^4} - \frac{5}{3} \right) , \quad (3.26)$$

where the  $\overline{\text{MS}}$  renormalization scheme has been used.

In order to check the renormalization group invariance of  $\Gamma^{(1)}$ , we need to know the anomalous dimension of the Gribov parameter  $\gamma$ . This is easily obtained from eq.(A.34), yielding

$$\gamma_{\gamma^2}(g^2) = -\frac{1}{2} \left( \frac{\beta(g^2)}{2g^2} - \gamma_A(g^2) \right) , \quad (3.27)$$

where  $\gamma_A(g^2)$  stands for the anomalous dimension of the gauge field  $A_\mu^a$ . Thus, at one-loop order,

$$\bar{\mu} \frac{d\Gamma^{(1)}}{d\bar{\mu}} = \left( 4(N^2 - 1) \left( \frac{\beta^{(1)}(g^2)}{2g^2} - \gamma_A^{(1)}(g^2) \right) + \frac{3(N^2 - 1)}{16\pi^2} 2Ng^2 \right) \gamma^4. \quad (3.28)$$

Furthermore, from (see e.g. [65])

$$\begin{aligned} \beta^{(1)}(g^2) &= -\frac{22}{3} \frac{g^4 N}{16\pi^2}, \\ \gamma_A^{(1)}(g^2) &= -\frac{13}{6} \frac{g^2 N}{16\pi^2}, \end{aligned} \quad (3.29)$$

it follows

$$\bar{\mu} \frac{d\Gamma^{(1)}}{d\bar{\mu}} = 0, \quad (3.30)$$

which establishes the RGE invariance of the effective action at the order considered.

We are now ready to face the more complex case in which the local composite operator  $A_\mu^a A_\mu^a$  is present. This will be the topic of the next section.

## 4 One-loop effective action in the $\overline{\text{MS}}$ scheme with the inclusion of $A_\mu^2$ .

### 4.1 Calculation of the one-loop effective potential.

Let us turn to the explicit one-loop evaluation of the effective action  $\Gamma$  in the presence of  $A_\mu^2$ . At one-loop, it turns out that<sup>2</sup>

$$\Gamma = -4(N^2 - 1)\gamma^4 + \frac{\sigma^2}{2g^2\zeta} + \frac{N^2 - 1}{2} \ln \det \left[ p^2 \delta_{\mu\nu} + \frac{2Ng^2\gamma^4}{p^2} \delta_{\mu\nu} - p_\mu p_\nu \left( 1 - \frac{1}{\alpha} \right) + \frac{g\sigma}{g^2\zeta} \delta_{\mu\nu} \right], \quad (4.1)$$

or

$$\Gamma = -4(N^2 - 1)\gamma^4 + \frac{\sigma^2}{2g^2\zeta} + \frac{N^2 - 1}{2}(d - 1) \int \frac{d^d p}{(2\pi)^d} \ln \left[ p^4 + 2Ng^2\gamma^4 + \frac{g\sigma}{g^2\zeta} p^2 \right]. \quad (4.2)$$

Before calculating the integral, we quote the two gap equations

$$\begin{aligned} \frac{\partial \Gamma}{\partial \sigma} = 0 &\Leftrightarrow \frac{\sigma}{\zeta_0} \left( 1 - \frac{\zeta_1}{\zeta_0} g^2 \right) + \frac{(N^2 - 1)g(d - 1)}{2\zeta_0} \int \frac{d^d p}{(2\pi)^d} \frac{p^2}{p^4 + \frac{g\sigma}{\zeta_0} p^2 + 2Ng^2\gamma^4} = 0, \\ \frac{\partial \Gamma}{\partial \gamma} = 0 &\Leftrightarrow \gamma^3 = \gamma^3 \frac{d - 1}{4} g^2 N \int \frac{d^d p}{(2\pi)^d} \frac{1}{p^4 + \frac{g\sigma}{\zeta_0} p^2 + 2Ng^2\gamma^4}. \end{aligned} \quad (4.3)$$

The second gap equation of (4.3), being the horizon condition, gives rise to the one obtained in the previous paper [49], while the first one describes the condensation of  $A_\mu^2$  when the restriction to the Gribov region  $\Omega$  is implemented. We notice that that the explicit value of the Gribov parameter  $\gamma$  is influenced by the presence of  $\langle A_\mu^2 \rangle$ .

It remains to calculate

$$\mathcal{I} = \int \frac{d^d p}{(2\pi)^d} \ln [p^4 + bp^2 + c], \quad (4.4)$$

---

<sup>2</sup>We shall drop from now on the superscript <sup>(1)</sup> indicating that we are working at one-loop order.

with

$$b = \frac{g\sigma}{\zeta_0}, \quad c = 2Ng^2\gamma^4, \quad (4.5)$$

Since

$$p^4 + bp^2 + c = (p^2 + \omega_1)(p^2 + \omega_2), \quad (4.6)$$

with

$$\omega_1 = \frac{b + \sqrt{b^2 - 4c}}{2}, \quad \omega_2 = \frac{b - \sqrt{b^2 - 4c}}{2}, \quad (4.7)$$

one has

$$\mathcal{I} = \int \frac{d^d p}{(2\pi)^d} \ln(p^2 + \omega_1) + \int \frac{d^d p}{(2\pi)^d} \ln(p^2 + \omega_2). \quad (4.8)$$

To make sense, the expression (4.4) should be real to ensure that the one-loop effective action is real-valued. Therefore, we must demand that  $c \geq 0$ . If  $b \geq 0$ ,  $\mathcal{I}$  is certainly real. However, when  $b^2 - 4c \leq 0$ , then also  $b < 0$  is allowed. We should thus have a positive Gribov parameter  $\gamma^4$ , while the condensate  $\langle A_\mu^2 \rangle$  can be negative or positive, depending on the case.

Using

$$\int \frac{d^d p}{(2\pi)^d} \ln(p^2 + m^2) = \frac{-m^4}{32\pi^2} \left( \frac{2}{\varepsilon} - \ln \frac{m^2}{\mu^2} + \frac{3}{2} \right), \quad (4.9)$$

it holds

$$\mathcal{I} = -\frac{\omega_1^2}{32\pi^2} \left( \frac{2}{\varepsilon} - \ln \frac{\omega_1}{\mu^2} + \frac{3}{2} \right) - \frac{\omega_2^2}{32\pi^2} \left( \frac{2}{\varepsilon} - \ln \frac{\omega_2}{\mu^2} + \frac{3}{2} \right). \quad (4.10)$$

Finally, in the  $\overline{\text{MS}}$  scheme, we obtain

$$\begin{aligned} \Gamma &= -4(N^2 - 1)\gamma^4 + \frac{\sigma^2}{2\zeta_0} \left( 1 - \frac{\zeta_1}{\zeta_0} g^2 \right) + \frac{3(N^2 - 1)}{2} \times \\ &\left[ \frac{\left( \frac{g\sigma}{\zeta_0} + \sqrt{\frac{g^2\sigma^2}{\zeta_0^2} - 8g^2N\gamma^4} \right)^2}{128\pi^2} \left( \ln \frac{\frac{g\sigma}{\zeta_0} + \sqrt{\frac{g^2\sigma^2}{\zeta_0^2} - 8g^2N\gamma^4}}{2\mu^2} - \frac{5}{6} \right) \right. \\ &\left. + \frac{\left( \frac{g\sigma}{\zeta_0} - \sqrt{\frac{g^2\sigma^2}{\zeta_0^2} - 8g^2N\gamma^4} \right)^2}{128\pi^2} \left( \ln \frac{\frac{g\sigma}{\zeta_0} - \sqrt{\frac{g^2\sigma^2}{\zeta_0^2} - 8g^2N\gamma^4}}{2\mu^2} - \frac{5}{6} \right) \right]. \quad (4.11) \end{aligned}$$

To lighten the notation a bit, let us introduce the new variables<sup>3</sup>

$$\lambda^4 = 8g^2N\gamma^4, \quad (4.12)$$

$$m^2 = \frac{g\sigma}{\zeta_0}. \quad (4.13)$$

in which case the action (4.11) can be rewritten as

$$\begin{aligned} \Gamma &= -\frac{(N^2 - 1)\lambda^4}{2g^2N} + \frac{\zeta_0 m^4}{2g^2} \left( 1 - \frac{\zeta_1}{\zeta_0} g^2 \right) \\ &+ \frac{3(N^2 - 1)}{256\pi^2} \left[ \left( m^2 + \sqrt{m^4 - \lambda^4} \right)^2 \left( \ln \frac{m^2 + \sqrt{m^4 - \lambda^4}}{2\mu^2} - \frac{5}{6} \right) \right. \\ &\left. + \left( m^2 - \sqrt{m^4 - \lambda^4} \right)^2 \left( \ln \frac{m^2 - \sqrt{m^4 - \lambda^4}}{2\mu^2} - \frac{5}{6} \right) \right]. \quad (4.14) \end{aligned}$$

---

<sup>3</sup>In comparison with the previous article [49], we have the correspondence  $\lambda^4 = 4\gamma^4$  with the Gribov parameter  $\gamma^4$  as defined there. It is actually this  $\gamma^4$  which will enter the modified propagators, see [49] and further in this paper.

We notice that the foregoing expression is also valid, i.e. real-valued, in the case in which  $m^4 < \lambda^4$ , as  $\ell_+(m, \lambda)$  and  $\ell_-(m, \lambda)$ , defined by,

$$\begin{aligned}\ell_+(m, \lambda) &= \left(m^2 + \sqrt{m^4 - \lambda^4}\right)^2 \left(\ln \frac{m^2 + \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2} - \frac{5}{6}\right) \\ \ell_-(m, \lambda) &= \left(m^2 - \sqrt{m^4 - \lambda^4}\right)^2 \left(\ln \frac{m^2 - \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2} - \frac{5}{6}\right)\end{aligned}\quad (4.15)$$

are complex conjugate<sup>4</sup>.

The horizon condition, eq.(3.14), can be translated to

$$\frac{\partial \Gamma}{\partial \lambda} = 0, \quad (4.16)$$

and the gap equation (3.13) to

$$\frac{\partial \Gamma}{\partial m^2} = 0. \quad (4.17)$$

As a check of this one-loop calculation, the expression (4.14) with  $m^2 \equiv 0$  reduces to the result obtained earlier in eq.(3.26), i.e. the original Gribov-Zwanziger model without the inclusion of  $A_\mu^2$ . If  $\lambda \equiv 0$ , i.e. the case where the condensation of  $A_\mu^2$  is investigated without implementing the restriction to the Gribov region  $\Omega$ , eq.(4.14) coincides with the result of [3, 12, 25]. From [11], one knows that

$$\bar{\mu} \frac{\partial \langle A_\mu^2 \rangle}{\partial \bar{\mu}} = \gamma_{A_\mu^2}(g^2) \langle A_\mu^2 \rangle = - \left( \frac{\beta(g^2)}{2g^2} + \gamma_A(g^2) \right) \langle A_\mu^2 \rangle, \quad (4.18)$$

or, using the relation (3.11) and the definition (4.13),

$$\bar{\mu} \frac{\partial m^2}{\partial \bar{\mu}} = \gamma_{m^2}(g^2) m^2 = \left( \frac{\beta(g^2)}{2g^2} - \gamma_A(g^2) \right) m^2, \quad (4.19)$$

while from eq.(3.27), it can be inferred that

$$\bar{\mu} \frac{\partial \lambda}{\partial \bar{\mu}} = \gamma_\lambda(g^2) \lambda = \frac{1}{4} \left( \frac{\beta(g^2)}{2g^2} + \gamma_A(g^2) \right) \lambda. \quad (4.20)$$

We notice the remarkable fact that the anomalous dimensions of the Gribov parameter and of the operator  $A_\mu^2$  are proportional to each other, to all orders of perturbation theory.

It can now be checked that  $\Gamma$  is renormalization group invariant, namely

$$\bar{\mu} \frac{d}{d\bar{\mu}} \Gamma = 0. \quad (4.21)$$

Finally, taking the derivatives of the action given in eq.(4.14) gives rise to

$$\begin{aligned}\frac{1}{\lambda^3} \frac{\partial \Gamma}{\partial \lambda} &= -\frac{2(N^2 - 1)}{g^2 N} + \frac{3(N^2 - 1)}{256\pi^2} \left[ -4 \frac{(m^2 + \sqrt{m^4 - \lambda^4})}{\sqrt{m^4 - \lambda^4}} \ln \frac{m^2 + \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2} \right. \\ &\quad \left. + 4 \frac{(m^2 - \sqrt{m^4 - \lambda^4})}{\sqrt{m^4 - \lambda^4}} \ln \frac{m^2 - \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2} + \frac{8}{3} \right],\end{aligned}\quad (4.22)$$

---

<sup>4</sup>Using  $\ln(z) = \ln|z| + i \arg(z)$  with  $-\pi < \arg(z) \leq \pi$ .

and

$$\begin{aligned} \frac{\partial \Gamma}{\partial m^2} = & \frac{\zeta_0 m^2}{g^2} \left(1 - \frac{\zeta_1}{\zeta_0} g^2\right) + \frac{3(N^2 - 1)}{256\pi^2} \left[ 2 \left(m^2 + \sqrt{m^4 - \lambda^4}\right) \left(1 + \frac{m^2}{\sqrt{m^4 - \lambda^4}}\right) \ln \frac{m^2 + \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2} \right. \\ & \left. + 2 \left(m^2 - \sqrt{m^4 - \lambda^4}\right) \left(1 - \frac{m^2}{\sqrt{m^4 - \lambda^4}}\right) \ln \frac{m^2 - \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2} - \frac{8}{3} m^2 \right]. \end{aligned} \quad (4.23)$$

## 4.2 Solving the gap equations.

We have now all the ingredients at hand to search for estimates of the mass parameter  $m^2$  and Gribov parameter  $\lambda$  as solutions of the gap equations (4.22) and (4.23). To avoid misinterpretations due to the suggestive use of the notation  $m^2$ , we remark that, due to the presence of  $\lambda$ , the mass parameter does not even appear as a pole in the tree level gluon propagator, see eq.(6.2).

Let us first consider the pure Gribov-Zwanziger case, i.e. we put  $m^2 \equiv 0$  in the expression (4.14). The relevant gap equation (horizon condition) reads

$$\frac{\partial \Gamma}{\partial \lambda} = \lambda^3 \left( -\frac{2(N^2 - 1)}{g^2 N} - \frac{3(N^2 - 1)}{64\pi^2} \left( \ln \frac{\lambda^4}{4\bar{\mu}^4} - \frac{5}{3} \right) - \frac{3(N^2 - 1)}{64\pi^2} \right) = 0. \quad (4.24)$$

We remind here that the solution  $\lambda = 0$  must be rejected. The natural choice for the renormalization scale is to set  $\bar{\mu}^2 = \frac{\lambda^4}{4}$  to kill the logarithms, and we find

$$\frac{g^2 N}{16\pi^2} \Big|_{\bar{\mu}^2 = \frac{\lambda^2}{2}} = 4. \quad (4.25)$$

In principle, from

$$g^2(\bar{\mu}^2) = \frac{1}{\beta_0 \ln \frac{\bar{\mu}^2}{\Lambda_{\overline{\text{MS}}}^2}}, \quad \text{with} \quad \beta_0 = \frac{11}{3} \frac{N}{16\pi^2}, \quad (4.26)$$

eq.(4.25) could be used to determine an estimate for the Gribov parameter, however it might be clear that this is meaningless since the corresponding expansion parameter (4.25) is far too big.

It is interesting to notice that, in a general massless renormalization scheme, the one-loop action with  $m^2 \equiv 0$  would read

$$\Gamma = -\frac{(N^2 - 1)}{2g^2 N} \lambda^4 - \frac{3\lambda^4 (N^2 - 1)}{264\pi^2} \left( \ln \frac{\lambda^4}{4\bar{\mu}^4} + a \right), \quad (4.27)$$

with  $a$  an arbitrary constant. The corresponding gap equation equals

$$\frac{\partial \Gamma}{\partial \lambda} = \lambda^3 \left( -\frac{2(N^2 - 1)}{g^2 N} - \frac{3(N^2 - 1)}{64\pi^2} \left( \ln \frac{\lambda^4}{4\bar{\mu}^4} + a \right) - \frac{3(N^2 - 1)}{64\pi^2} \right) = 0. \quad (4.28)$$

Denoting by  $\lambda_*$  a solution of eq.(4.28), for the vacuum energy corresponding to (4.27) one finds

$$E_{\text{vac}} = \Gamma(\lambda_*) = \frac{3(N^2 - 1)}{64\pi^2} \frac{\lambda_*^4}{4}. \quad (4.29)$$

This expression is valid for all  $\bar{\mu}$  and for all  $a$ . The vacuum energy is thus always nonnegative at one-loop order in the original Gribov-Zwanziger model.

The gap equation (4.23) with  $\lambda \equiv 0$  obviously has the solution already obtained in [3, 12, 25] where the restriction to the Gribov region  $\Omega$  was not taken into account. We recall the values

$$\frac{g^2 N}{16\pi^2} = \frac{36}{187} \approx 0.193, \quad (4.30)$$

$$m^2 = e^{\frac{17}{12}} \Lambda_{\overline{\text{MS}}}^2 \approx (2.031 \Lambda_{\overline{\text{MS}}})^2, \quad (4.31)$$

$$E_{\text{vac}} = -\frac{3}{16\pi^2} e^{\frac{17}{6}} \Lambda_{\overline{\text{MS}}}^4 \approx -0.323 \Lambda_{\overline{\text{MS}}}^4, \quad (4.32)$$

which were obtained upon setting  $\bar{\mu}^2 = m^2$  to kill the logarithms.

We shall now show that, in the  $\overline{\text{MS}}$  scheme, the gap equations (4.22)-(4.23) have no solution with  $m^2 > 0$  when the restriction to the horizon is implemented (i.e. when  $\lambda \neq 0$ ). To this purpose, we introduce for  $m^2 > 0$  the variable

$$t = \frac{\lambda^4}{m^4}. \quad (4.33)$$

Evidently, we should only consider  $t > 0$ .

Dividing the gap equations (4.22)-(4.23) by  $m^2$ , they can be rewritten as<sup>5</sup>

$$\frac{16\pi^2}{g^2 N} = \frac{3}{8} \left( -2 \ln \frac{m^2}{2\bar{\mu}^2} + \frac{2}{3} + \frac{1}{\sqrt{1-t}} \ln \frac{t}{(1+\sqrt{1-t})^2} - \ln t \right), \quad (4.34)$$

and

$$-\frac{24}{13} \left( \frac{16\pi^2}{g^2 N} \right) + \frac{322}{39} = 4 \ln \frac{m^2}{2\bar{\mu}^2} - \frac{4}{3} - \frac{2-t}{\sqrt{1-t}} \ln \frac{t}{(1+\sqrt{1-t})^2} + 2 \ln t, \quad (4.35)$$

where use has been made of the explicit values of  $\zeta_0$  and  $\zeta_1$ , which can be found in [3, 12, 25]

$$\zeta_0 = \frac{9}{13} \frac{N^2 - 1}{N}, \quad \zeta_1 = \frac{161}{52} \frac{N^2 - 1}{16\pi^2}, \quad (4.36)$$

The eqns.(4.35)-(4.36) can be combined to eliminate  $\ln \frac{m^2}{2\bar{\mu}^2}$ , yielding the following condition

$$\frac{68}{39} \left( \frac{16\pi^2}{g^2 N} \right) + \frac{161}{39} = \frac{t}{\sqrt{1-t}} \ln \frac{t}{(1+\sqrt{1-t})^2} \equiv F(t). \quad (4.37)$$

It can be checked that  $F(t)$  is real-valued and negative for  $t > 0$ , thus the r.h.s. of eq.(4.37) is always negative. Since the l.h.s. of eq.(4.37) is necessarily positive for a meaningful result (i.e.  $g^2 \geq 0$ ), there is no solution with  $m^2 > 0$ . As already mentioned, there are a priori also possible solutions with  $m^2 < 0$ .

To investigate the existence of a solution with  $m^2 < 0$ , it might be instructive to look again at the gap equations (4.22) and (4.23) from another perspective. We recall that, if the horizon is not implemented, i.e.  $\lambda \equiv 0$ , the gap equation (4.23) has two solutions, a perturbative one

<sup>5</sup>We have already factored out  $m^2$  or  $\lambda^3$  since these are non-zero in the present case.



corresponding to  $m^2 = 0$  (no condensation) and a non-perturbative one corresponding to the  $m^2$  given in eq.(4.31).

If we momentarily consider  $\lambda$  as a free, adjustable parameter of the theory, eq.(4.23) dictates how  $m^2$  becomes a function of the parameter  $\lambda$ . From the result at  $\lambda = 0$ , we could expect that two branches of solutions would evolve, one starting from the perturbative and one from the non-perturbative value of  $m^2$  at  $\lambda = 0$ . When  $\lambda \equiv 0$ , the choice for the scale  $\bar{\mu}$  is quite obvious from the requirement that all the logarithms  $\ln \frac{m^2}{\bar{\mu}^2}$  are vanishing. However, when  $\lambda \neq 0$ , we notice that there are two kinds of logarithms present, being  $\ln \frac{m^2 + \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2}$  and  $\ln \frac{m^2 - \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2}$ . We opt to set

$$\bar{\mu}^2 = \frac{|m^2 + \sqrt{m^4 - \lambda^4}|}{2}. \quad (4.38)$$

This reduces to  $\bar{\mu}^2 = m^2$  if  $\lambda = 0$ , while it allows for<sup>6</sup>  $m^2 < 0$ . This is possible if  $m^4 \leq \lambda^4$ , as it was mentioned below eq.(4.8). In this case, the size of both logarithms,  $\ln \frac{m^2 + \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2}$  and  $\ln \frac{m^2 - \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2}$ , is determined by their arguments, which are complex conjugate.

Let us specify to the case  $N = 3$ . In Figure 1, we have plotted the behaviour of  $m^2(\lambda^4)$ . We see that next to the “non-perturbative” branch of solutions, starting from  $m^2 \neq 0$ , also a “perturbative” branch of solutions with  $m^2 < 0$  is emerging from  $m^2 = 0$ , in correspondence with our expectation.




Figure 1:  $m^2$  as a function of  $\lambda^4$ , in units  $\Lambda_{\overline{\text{MS}}} = 1$ .

However,  $\lambda^4$  is not a free parameter of the theory. We should require that  $\lambda^4$  is such that the doublet  $(\lambda^4, m^2(\lambda^4))$  is a solution of the gap equation (4.22), i.e. the horizon condition. In Figure 2, we have plotted the value of the horizon condition equation, as a function of  $\lambda^4$ . It is clear that no solution with  $m^2 > 0$  exists as the horizon condition never becomes zero. Of course, this is in correspondence with the foregoing general proof that there is never such a solution, independently of the choice of  $\bar{\mu}$ . However, we see that there is a single solution with  $m^2 < 0$ .

<sup>6</sup>Evidently,  $\bar{\mu}^2$  should be real and positive, hence the modulus in eq.(4.38).




Figure 2: The horizon condition (4.22) as a function of  $\lambda^4$ , in units  $\Lambda_{\overline{\text{MS}}} = 1$ . The top curve corresponds to the solutions of (4.23) with  $m^2 < 0$  and the lower curve to the solutions with  $m^2 > 0$ .

The corresponding values for the expansion parameter, for the Gribov and mass parameter, as well as for the vacuum energy are found to be

$$\frac{g^2 N}{16\pi^2} \approx 1.18, \quad (4.39)$$

$$\lambda^4 \approx 6.351 \Lambda_{\overline{\text{MS}}}^4, \quad (4.40)$$

$$m^2 \approx -0.950 \Lambda_{\overline{\text{MS}}}^2, \quad (4.41)$$

$$E_{\text{vac}} \approx 0.043 \Lambda_{\overline{\text{MS}}}^4, \quad (4.42)$$

### 4.3 Intermediate comments.

Although the  $\overline{\text{MS}}$  expansion parameter (4.39) is too large to speak about reliable results, we nevertheless would like to raise some questions. Apparently, the solution of the coupled gap equations is laying on the “perturbative” branch, being the one with  $m^2 \leq 0$ . This gives rise to a positive value for the mass dimension two gluon condensate  $\langle A_\mu^2 \rangle$ . When the restriction on the domain of integration in the path integral is not implemented, as in the previous papers [3, 12, 25],  $\langle A_\mu^2 \rangle$  was necessarily negative, the reason being that the action should be real-valued, as it was explained below eq.(4.8). As already explained in the Introduction, a finding a bit unfortunate is that the vacuum energy is positive, eq.(4.42), which leads to a negative estimate for the gluon condensate  $\langle \frac{g^2}{4\pi^2} F_{\mu\nu}^2 \rangle$  via the trace anomaly, eq.(1.8). Essentially, we are thus left with the following questions:

- (i.) What is the sign and value of  $m^2$  and thus of  $\langle A_\mu^2 \rangle$ ?
- (ii.) What is the sign and value of  $E_{\text{vac}}$  and the corresponding value for  $\langle \frac{g^2}{4\pi^2} F_{\mu\nu}^2 \rangle$ ?
- (iii.) Are these values better or not when the operator  $A_\mu^2$  is added to the original Gribov-Zwanziger model?

## 5 Changing and reducing the dependence on the renormalization scheme.

We have already shown that the vacuum energy obtained in a one-loop approximation is always positive when the condensation of the operator  $A_\mu^2$  is left out of the discussion, using whatever renormalization scheme.

To answer the foregoing questions (i.)-(iii.), one could investigate what happens at two-loop order. However, due to the already quite complicated structure of the one-loop effective action and to the fact that the calculations at higher loop order will not get any easier, this task is beyond the scope of the present article. Here, we shall mainly focus on the effects of a change of the renormalization scheme at the one-loop order. It could happen that, in a scheme different from the  $\overline{\text{MS}}$  one, the vacuum energy is negative and/or that the coupling constant is small enough to speak about trustworthy results, at least qualitatively.

Since to obtain an optimization of the renormalization scheme and scale dependence is a rather lengthy task, we shall not dwell upon technicalities in this section. The interested reader can find all details in Appendix B. We shall thus focus on the main results obtained after the optimization.

Essentially, what we have done is replacing in the effective action (4.14) the quantities  $m^2$  and  $\lambda^4$  by their order by order renormalization scale and scheme invariant counterparts  $\widehat{m}^2$  and  $\widehat{\lambda}^4$ . The residual freedom in the choice of renormalization scheme can then be reduced to a single parameter  $b_0$ , related to coupling constant renormalization. As the vacuum energy is a physical quantity, it should in principle not depend on  $b_0$ . At the same time, the quantities  $\widehat{m}^2$  and  $\widehat{\lambda}^4$  should be  $b_0$  independent by construction. This provides one with the interesting opportunity to fix the redundant parameter  $b_0$  by demanding a minimal dependence on it.

The final one-loop action turns out to be given by

$$\begin{aligned} \Gamma = & -\frac{(N^2-1)}{2N}x^{-2b}\widehat{\lambda}^4\left(x+B+(1-2b)\left(\frac{\beta_1}{\beta_0}\ln\frac{x}{\beta_0}-b_0\right)\right) \\ & +\frac{\zeta_0}{2}\widehat{m}^4x^{-2a}\left(x+A-\frac{\zeta_1}{\zeta_0}+(1-2a)\left(\frac{\beta_1}{\beta_0}\ln\frac{x}{\beta_0}-b_0\right)\right)+\frac{3(N^2-1)}{256\pi^2}\times \\ & \left[\left(\widehat{m}^2x^{-a}+\sqrt{\widehat{m}^4x^{-2a}-\widehat{\lambda}^4x^{-2b}}\right)^2\left(\ln\frac{\widehat{m}^2x^{-a}+\sqrt{\widehat{m}^4x^{-2a}-\widehat{\lambda}^4x^{-2b}}}{2\overline{\mu}^2}-\frac{5}{6}\right)\right. \\ & \left.+\left(\widehat{m}^2x^{-a}-\sqrt{\widehat{m}^4x^{-2a}-\widehat{\lambda}^4x^{-2b}}\right)^2\left(\ln\frac{\widehat{m}^2x^{-a}-\sqrt{\widehat{m}^4x^{-2a}-\widehat{\lambda}^4x^{-2b}}}{2\overline{\mu}^2}-\frac{5}{6}\right)\right], \end{aligned} \quad (5.1)$$

while the corresponding gap equations read

$$\frac{1}{\widehat{\lambda}^3}\frac{\partial\Gamma}{\partial\widehat{\lambda}}=0, \quad (5.2)$$

$$\frac{1}{\widehat{m}^2}\frac{\partial\Gamma}{\partial\widehat{m}^2}=0. \quad (5.3)$$

The definitions of all quantities appearing in the above expressions can be found in the Appendix B.

In Figure 3, we collected the solutions of the scale invariant quantities  $\widehat{m}^2$  and  $\widehat{\lambda}^4$  as a function

of  $b_0$ , while Figure 4 displays the vacuum energy  $E_{\text{vac}}$  and the relevant expansion parameter, given by  $y \equiv \frac{N}{16\pi^2 x}$ . For completeness, we have also shown the solutions which correspond to higher values of  $\Gamma$  and are as such describing unstable solutions. These are indicated with the thinner lines.




Figure 3: The quantities  $\hat{m}^2$  and  $\hat{\lambda}^4$  as a function of  $b_0$ , in units  $\Lambda_{\overline{\text{MS}}} = 1$ .




Figure 4: The vacuum energy  $E_{\text{vac}}$  and the expansion parameter  $y$  as a function of  $b_0$ , in units  $\Lambda_{\overline{\text{MS}}} = 1$ .

## 5.1 Interpretation of the results.

Let us first have a look at the plot of vacuum energy, on the l.h.s. of Figure 4. We notice that for  $b_0 < -0.33564\dots$ , the vacuum energy becomes negative. However, we cannot attach any definitive meaning to this result. In fact, as it can be seen from the Figures 3 and 4, the values of the vacuum energy and the supposedly minimally  $b_0$ -dependent quantities  $\hat{m}^2$  and  $\hat{\lambda}^4$  are extremely  $b_0$ -dependent. Very small variations in  $b_0$  induce large fluctuations on e.g. the energy. This is indicative of the fact that the equations we have solved are not yet stable against  $b_0$ -variations in the range of the values obtained for  $b_0$ . The behaviour is better for, let us say  $b_0 > -0.2$ . However, in this case, we find again that the vacuum energy is positive. The vacuum energy  $E_{\text{vac}}$ , as well as  $\hat{m}^2$  and  $\hat{\lambda}^4$  fall of to zero for growing  $b_0$ .

As an example, we set  $b_0 = 0$ , which corresponds to use the  $\overline{\text{MS}}$  coupling constant. Then

we find, with the optimized expansion,

$$y \equiv \frac{N}{16\pi^2 x} \approx 0.796, \quad (5.4)$$

$$\widehat{\lambda}^4 x^{-2b} \approx 7.939 \Lambda_{\overline{\text{MS}}}^4, \quad (5.5)$$

$$\widehat{m}^2 x^{-a} \approx -0.814 \Lambda_{\overline{\text{MS}}}^2, \quad (5.6)$$

$$E_{\text{vac}} \approx 0.063 \Lambda_{\overline{\text{MS}}}^4, \quad (5.7)$$

results which are in fair agreement with the naive  $\overline{\text{MS}}$  results (4.39)-(4.42). We notice that the expansion parameter  $y$  is already smaller than 1, but still relatively large, while the vacuum energy is indeed positive.

The conclusion than can be drawn from this section is that we cannot find a reliable result with negative vacuum energy and hence positive gluon condensate  $\langle F_{\mu\nu}^2 \rangle$  using a one-loop approximation. We see therefore that, in order to be able to give a reasonable answer to the questions concerning the sign of  $m^2$  and  $E_{\text{vac}}$  and to get more trustworthy numerical values, the two-loop evaluation of the effective action  $\Gamma$ , at least in the  $\overline{\text{MS}}$  scheme, would be very useful.

## 6 Consequences of a non-vanishing Gribov parameter.

Before turning to the final conclusions, we shall give in this section a brief account of some well known consequences stemming from the presence of the Gribov parameter, to emphasize the important role of this parameter.

### 6.1 The gluon propagator.

If there is no generation of a mass parameter due to  $\langle A_\mu^2 \rangle$ , we can consider just the action (2.2). Then the tree level gluon propagator turns out to be

$$\langle A_\mu^a A_\nu^b \rangle_p = \delta^{ab} \frac{p^2}{p^4 + \frac{\lambda^4}{4}} \left( \delta_{\mu\nu} - \frac{p_\mu p_\nu}{p^2} \right). \quad (6.1)$$

This result, first pointed out in [48], was obtained by retaining only the first term of the nonlocal horizon function (1.4), corresponding to the approximation  $-\partial D \approx -\partial^2$ . The gluon propagator, eq.(6.1), is suppressed in the infrared region due to the presence of the Gribov parameter  $\lambda$ . In particular, the presence of this parameters implies that  $\langle A_\mu^a A_\nu^b \rangle_p$  vanishes at zero momentum,  $p = 0$ . When the possibility of the existence of a dynamical mass parameter in the gluon propagator is included, by investigating the condensation of  $A_\mu^2$ , the tree level gluon propagator reads

$$\langle A_\mu^a A_\nu^b \rangle_p \equiv \delta^{ab} \frac{\mathcal{D}(p^2)}{p^2} \left( \delta_{\mu\nu} - \frac{p_\mu p_\nu}{p^2} \right) = \delta^{ab} \frac{p^2}{p^4 + m^2 p^2 + \frac{\lambda^4}{4}} \left( \delta_{\mu\nu} - \frac{p_\mu p_\nu}{p^2} \right). \quad (6.2)$$

This type of propagator is sometimes called the Stingl propagator, from the author who used it as an ansatz for solving the Schwinger-Dyson equations, see [66] for more details .

However, it should be realized that eq.(6.2) describes only the tree level gluon propagator. In particular, to produce a plot of the form factor  $\mathcal{D}(p^2)$  as a function of the momentum  $p$ , which would allow to make a comparison with the results obtained in lattice simulations, see

e.g. [31] for  $N = 3$  and [32, 34] for  $N = 2$ , one should go beyond the zeroth order approximation, for example by including higher order polarization effects and/or trying to perform a renormalization group improvement. In general, these corrections will also be dependent on the external momentum  $p$ .

## 6.2 The ghost propagator.

Even more prominent is the influence of the Gribov parameter on the infrared behaviour of the ghost propagator, which can be calculated at one-loop order using the modified gluon propagator (6.1) or (6.2) with their respective gap equations (1.6) and (4.3). In both cases, the infrared behaviour of the ghost propagator [48, 49, 50, 51, 52] is shown to be

$$\frac{\delta^{ab}}{N^2 - 1} \langle c^a \bar{c}^b \rangle_{p \approx 0} \equiv \frac{1}{p^2} \mathcal{G}(p^2) \Big|_{p \approx 0} \approx \frac{4}{3Ng^2 \mathcal{J} p^4}, \quad (6.3)$$

where  $\mathcal{J}$  stands for the real, finite integral given by

$$\mathcal{J} = \int \frac{d^4 k}{(2\pi)^4} \frac{1}{k^2 \left( k^4 + m^2 k^2 + \frac{\lambda^4}{4} \right)}. \quad (6.4)$$

The original Gribov-Zwanziger model corresponds to  $m^2 \equiv 0$ . Thus, the ghost propagator is strongly enhanced in the infrared region compared to the perturbative behaviour, if the restriction to the first Gribov region is taken into account. It is important to notice that this behaviour of the ghost propagator is preserved in the present treatment, due to the peculiar form of the gap equation (4.3) implementing the horizon condition. In particular, from the expression for the effective action in eq.(4.2), one sees that, while the term quadratic in the field  $\sigma$ , i.e.  $\frac{\sigma^2}{2g^2\zeta}$ , contains the LCO parameter  $\zeta$ , the first term which depends on the Gribov parameter, i.e.  $-4(N^2 - 1)\gamma^4$ , does not contain any such new LCO parameter. This important feature follows from the fact that no new parameter has to be introduced in order to renormalize the term  $(M_\mu^{ai} V_\mu^{ai} - U_\mu^{ai} N_\mu^{ai})$ , as remarked in eq.(A.35). While the parameter  $\zeta$  is required to take into account the ultraviolet divergences of the vacuum correlator  $\langle A_\mu^2(x) A_\nu^2(y) \rangle$ , which are proportional to  $\tau^2$ , no such a parameter is needed for  $(M_\mu^{ai} V_\mu^{ai} - U_\mu^{ai} N_\mu^{ai})$  which, upon setting the external sources to their physical values, gives rise to term  $-4(N^2 - 1)\gamma^4$  in the expression (4.2). Said otherwise, this term is not affected by the presence of a new parameter which would be required if eq.(A.35) would not hold. As a consequence, the factor “1” appearing in the left hand side of the gap equation (4.3) is, so to speak, left unchanged by the quantum corrections. It is precisely that property which ensures, through a delicate cancelation mechanism, see [48, 49, 51, 52], the infrared enhancement of the ghost propagator.

Analogously to the case of the gluon propagator, a more detailed study of higher order corrections would be needed in order to obtain a plot of the ghost form factor  $\mathcal{G}(p^2)$ .

## 6.3 The strong coupling constant.

Usually, a nonperturbative definition of the renormalized strong coupling constant  $\alpha_R$  can be written down from the knowledge of the gluon and ghost propagators as, see e.g. [39, 34]

$$\alpha_R(p^2) = \alpha_R(\mu) \mathcal{D}(p^2, \mu) \mathcal{G}^2(p^2, \mu), \quad (6.5)$$

where  $\mathcal{D}$  and  $\mathcal{G}$  stand for the gluon and ghost form factors as defined before. This definition represents a kind of nonperturbative extension of the perturbative results (A.33). According to

Schwinger-Dyson studies [41, 42, 43, 44, 45, 46], those form factors satisfy a power law behaviour in the infrared

$$\begin{aligned}\lim_{p \rightarrow 0} \mathcal{D}(p^2) &\propto (p^2)^\theta, \\ \lim_{p \rightarrow 0} \mathcal{G}(p^2) &\propto (p^2)^\omega,\end{aligned}\tag{6.6}$$

where the infrared exponents  $\theta$  and  $\omega$  obey the sum rule

$$\theta + 2\omega = 0.\tag{6.7}$$

Such a sum rule suggests the development of an infrared fixed point for the renormalized coupling constant, (6.5), as also pointed out by lattice simulations for the  $SU(2)$  as well as for the  $SU(3)$  case [34, 35, 36],

$$\lim_{p \rightarrow 0} \alpha(p^2) = \alpha_c.\tag{6.8}$$

The existence of a fixed point in this reasoning is dependent on the sum rule rather than on the precise value of the exponents. We refer to the already quoted literature for more details on the value of these exponents. We end by noticing that the form factors of the gluon and ghost propagator in our zeroth order approximation give rise to the sum rule (6.7), since we have  $\theta = 2$  and  $\omega = -1$ . Moreover, without Gribov parameter, the sum rule (6.7) is lost, and thus there is no indication for an infrared fixed point.

#### 6.4 Positivity violation.

The behaviour of the gluon propagator is sometimes used as an *indication* of confinement of gluons by means of the so called positivity violation, see e.g. [67, 68] and references therein.

Briefly, when the Euclidean gluon propagator  $D(p) \equiv \frac{\mathcal{D}(p^2)}{p^2}$  is written through a spectral representation as

$$D(p) = \int_0^{+\infty} dM^2 \frac{\rho(M^2)}{p^2 + M^2},\tag{6.9}$$

the spectral density  $\rho(M^2)$  should be positive in order to have a Källén-Lehmann representation, making possible the interpretation of the fields in term of stable particles. We refer to [67, 68] for more details. One can define the temporal correlator [68]

$$\mathcal{C}(t) = \int_0^{+\infty} dM \rho(M^2) e^{-Mt},\tag{6.10}$$

which is certainly positive for positive  $\rho(M^2)$ . The inverse is not necessarily true.  $\mathcal{C}(t)$  can be also positive for a  $\rho(M^2)$  attaining negative values. However, if  $\mathcal{C}(t)$  becomes negative for certain  $t$ , then a fortiori  $\rho(M^2)$  cannot be always positive. Using a contour integration argument, it is not difficult to show that  $\mathcal{C}(t)$  can be rewritten as

$$\mathcal{C}(t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-ipt} D(p) dp.\tag{6.11}$$

Let us consider the function  $\mathcal{C}(t)$  using the tree level propagator (6.2), thus using

$$D(p) = \frac{p^2}{p^4 + p^2 m^2 + \frac{\lambda^4}{4}}.\tag{6.12}$$

We can consider several cases<sup>7</sup>:

---

<sup>7</sup>Each of the following expressions for  $\mathcal{C}(t)$  is obtainable via contour integration.

- if  $\lambda = 0$  (thus  $m^2 > 0$ ), one shall find that

$$\mathcal{C}(t) = \frac{e^{-mt}}{2m} . \quad (6.13)$$

This function is always positive.

- if  $m^2 = 0$ ,

$$\mathcal{C}(t) = \frac{e^{-\frac{Lt}{2}}}{2L} \left( \cos \frac{Lt}{2} - \sin \frac{Lt}{2} \right) , \quad (6.14)$$

and clearly, this function will attain negative values for certain  $t$ .

- in any other case, the correlator  $\mathcal{C}(t)$  is found to be

$$\mathcal{C}(t) = \frac{1}{2} \left[ \frac{\sqrt{\omega_1}}{\omega_1 - \omega_2} e^{-\sqrt{\omega_1}t} + \frac{\sqrt{\omega_2}}{\omega_2 - \omega_1} e^{-\sqrt{\omega_2}t} \right] \quad (6.15)$$

where the decomposition

$$\frac{p^2}{p^4 + p^2 m^2 + \frac{\lambda^4}{4}} = \frac{\omega_1}{\omega_1 - \omega_2} \frac{1}{p^2 + \omega_1} - \frac{\omega_2}{\omega_1 - \omega_2} \frac{1}{p^2 + \omega_2} , \quad (6.16)$$

has been employed. It is understood that  $\sqrt{\omega_1}$  ( $\sqrt{\omega_2}$ ) is the root having a positive real part.

If we assume that  $\hat{m}^4 > \lambda^4$ , then  $\omega_1 > \omega_2$  and  $\mathcal{C}(t)$  becomes negative for  $t > \frac{1}{2} \frac{\ln \frac{\omega_1}{\omega_2}}{\omega_1 - \omega_2}$ . In the case that  $\hat{m}^4 = \lambda^4$ , or  $\omega_1 = \omega_2$ , one finds that  $\mathcal{C}(t) = \frac{e^{-\sqrt{\omega_1}t}}{4\sqrt{\omega_1}} (1 - \sqrt{\omega_1}t)$ , which can also become negative. If  $\hat{m}^4 < \lambda^4$ , we can reintroduce the complex polar coordinates  $R$  and  $\phi$  for the complex conjugate quantities  $\omega_1$  and  $\omega_2$ . If  $\cos \frac{\phi}{2} \geq 0$ , eq.(6.15) can be rewritten as

$$\mathcal{C}(t) = \frac{1}{2\sqrt{R} \sin \phi} e^{-\sqrt{R} \cos(\frac{\phi}{2})t} \sin \left( \frac{\phi}{2} - \sqrt{R} \sin \left( \frac{\phi}{2} \right) t \right) \quad (6.17)$$

By choosing an appropriate value of  $t > 0$ , also this expression can be made negative. An analogous expression and conclusion can be derived in case that  $\cos \frac{\phi}{2} < 0$

We conclude that, when the restriction to the Gribov region  $\Omega$  is implemented, the function  $\mathcal{C}(t)$  exhibits a violation of positivity when the tree level propagator is used, with or without the inclusion of  $\langle A_\mu^2 \rangle$ .

The goal of this section was merely to provide some interesting consequences when the restriction to the first Gribov region  $\Omega$  is implemented. Higher loop effects, which shall be momentum dependent, would also influence the behaviour of the gluon and ghost propagator. Hence, to give a sensible interpretation of the behaviour of e.g. the form factors and of the strong coupling constant  $\alpha_R$ , a more detailed analysis than a tree level one is necessary. This is however far beyond the aim of this work.

## 7 Conclusion.

In this work we have considered  $SU(N)$  Euclidean Yang-Mills theories in the Landau gauge,  $\partial_\mu A_\mu = 0$ . We have studied the condensation of the dimension two composite operator  $A_\mu^2$  when the restriction to the Gribov region  $\Omega$  is taken into account. Such a restriction is needed due to



the presence of the Gribov copies [48], which are known to affect the Landau gauge.

In a previous work [49], the consequences of the restriction to the region  $\Omega$  in the presence of a dynamical mass parameter due to the gluon condensate  $\langle A_\mu^a A_\mu^a \rangle$  were studied by following Gribov's seminal work [48]. Here, we have relied on Zwanziger's action [51, 52], which allows to implement the restriction to the Gribov region  $\Omega$  within a local and renormalizable framework. We have been able to show that Zwanziger's action remains renormalizable to all orders of perturbation theory in the presence of the operator  $A_\mu^2$ , introduced through the local composite operator technique [3, 12, 20, 25]. The effective action, constructed via the local composite operator formalism [3] obeys a homogeneous renormalization group. The explicit form of the one-loop effective action has been worked out. We have seen that, considering the original Gribov-Zwanziger model, i.e. without including the operator  $A_\mu^2$ , the vacuum energy is always positive at one-loop order, independently from the choice of the renormalization scheme. A positive vacuum energy would give rise to a negative value for the gauge invariant gluon condensate  $\langle F_{\mu\nu}^2 \rangle$ , through the trace anomaly. Furthermore, by adding the operator  $A_\mu^2$ , we have proven that there is no solution of the two coupled gap equations at the one-loop order in the  $\overline{\text{MS}}$  scheme with  $\langle A_\mu^2 \rangle < 0$ . Nevertheless, when  $\langle A_\mu^2 \rangle > 0$ , a solution of the gap equations was found, although the corresponding expansion parameter was too large and the vacuum energy still positive.

In order to find out what happens in other schemes, we performed a detailed study, at lowest order, of the influence of the renormalization scheme. We have been able to reduce the freedom of the choice of the renormalization scheme to two parameters, namely the renormalization scale  $\bar{\mu}$  and a parameter  $b_0$ , associated to the coupling constant renormalization. We reexpressed the effective action in terms of the mass parameter  $\hat{m}$  and Gribov parameter  $\hat{\lambda}$ , which are renormalization scheme and scale independent order by order. The resulting gap equations for these parameters have been solved numerically. Although a solution with negative vacuum energy was found, we have been unable to attach any definitive meaning to it. This is due to the fact that the results obtained turn out to be strongly dependent from the parameter  $b_0$ . This brought us to the conclusion that we should extend our calculations to a higher order to obtain more sensible numerical estimates.

The mass parameters  $\hat{m}$  and  $\hat{\lambda}$  are of a nonperturbative nature and appear in the gluon and ghost propagator. Even if we lack reliable estimates for these parameters, some already known interesting features can be recovered. For a nonzero mass and Gribov parameter, there is a *qualitative* agreement with the behaviour found in lattice simulations and Schwinger-Dyson studies: a suppressed gluon and enhanced ghost propagator in the infrared, while further consequences of the Gribov parameter are e.g. the possible existence of an infrared fixed point for the strong coupling constant and the violation of positivity related to the gluon propagator.

## Acknowledgments.

The Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq-Brazil), the Faperj, Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro, the SR2-UERJ and the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) are gratefully acknowledged for financial support. D. Dudal would like to acknowledge the warm hospitality at the UERJ, where part of this work was done, while R. F. Sobreiro would like to acknowledge the kind hospitality at the Department of Mathematical Physics and Astronomy of the Ghent University, where this work was completed.

## A Appendix A.

In this Appendix, we have collected all details of the multiplicative renormalization of the Zwanziger action in the presence of the operator  $A_\mu^2$ .

### A.1 Ward identities.

In order to begin with the algebraic characterization of the most general counterterm needed for the renormalizability of the complete action  $\Sigma$  of eq.(3.1), let us first give the set of Ward identities which are fulfilled by  $\Sigma$ . These are

- the Slavnov-Taylor identity

$$\mathcal{S}(\Sigma) = 0, \quad (\text{A.1})$$

with

$$\begin{aligned} \mathcal{S}(\Sigma) = \int d^4x \left( \frac{\delta\Sigma}{\delta K_\mu^a} \frac{\delta\Sigma}{\delta A_\mu^a} + \frac{\delta\Sigma}{\delta L^a} \frac{\delta\Sigma}{\delta c^a} + b^a \frac{\delta\Sigma}{\delta \bar{c}^a} + \bar{\varphi}_i^a \frac{\delta\Sigma}{\delta \bar{\omega}_i^a} + \omega_i^a \frac{\delta\Sigma}{\delta \varphi_i^a} \right. \\ \left. + M_\mu^{ai} \frac{\delta\Sigma}{\delta U_\mu^{ai}} + N_\mu^{ai} \frac{\delta\Sigma}{\delta V_\mu^{ai}} + \tau \frac{\delta\Sigma}{\delta \eta} \right), \end{aligned} \quad (\text{A.2})$$

- the Landau gauge condition and the antighost equation

$$\frac{\delta\Sigma}{\delta b^a} = \partial_\mu A_\mu^a, \quad (\text{A.3})$$

$$\frac{\delta\Sigma}{\delta \bar{c}^a} + \partial_\mu \frac{\delta\Sigma}{\delta K_\mu^a} = 0, \quad (\text{A.4})$$

- the ghost Ward identity

$$\mathcal{G}^a \Sigma = \Delta_{\text{cl}}^a, \quad (\text{A.5})$$

with

$$\mathcal{G}^a = \int d^4x \left( \frac{\delta}{\delta c^a} + g f^{abc} \left( \bar{c}^b \frac{\delta}{\delta b^c} + \varphi_i^b \frac{\delta}{\delta \omega_i^c} + \bar{\omega}_i^b \frac{\delta}{\delta \bar{\varphi}_i^c} + V_\mu^{bi} \frac{\delta}{\delta N_\mu^{ci}} + U_\mu^{bi} \frac{\delta}{\delta M_\mu^{ci}} \right) \right), \quad (\text{A.6})$$

and

$$\Delta_{\text{cl}}^a = g \int d^4x f^{abc} \left( K_\mu^b A_\mu^c - L^b c^c \right). \quad (\text{A.7})$$

Notice that the term  $\Delta_{\text{cl}}^a$ , being linear in the quantum fields  $A_\mu^a$ ,  $c^a$ , is a classical breaking.

- the linearly broken local constraints

$$\frac{\delta\Sigma}{\delta \bar{\varphi}^{ai}} + \partial_\mu \frac{\delta\Sigma}{\delta M_\mu^{ai}} = g f^{abc} A_\mu^b V_\mu^{ci}, \quad (\text{A.8})$$

$$\frac{\delta\Sigma}{\delta \omega^{ai}} + \partial_\mu \frac{\delta\Sigma}{\delta N_\mu^{ai}} - g f^{abc} \bar{\omega}^{bi} \frac{\delta\Sigma}{\delta b^c} = g f^{abc} A_\mu^b U_\mu^{ci}, \quad (\text{A.9})$$

$$\frac{\delta\Sigma}{\delta \bar{\omega}^{ai}} + \partial_\mu \frac{\delta\Sigma}{\delta U_\mu^{ai}} - g f^{abc} V_\mu^{bi} \frac{\delta\Sigma}{\delta K_\mu^c} = -g f^{abc} A_\mu^b N_\mu^{ci}, \quad (\text{A.10})$$

$$\frac{\delta\Sigma}{\delta \varphi^{ai}} + \partial_\mu \frac{\delta\Sigma}{\delta V_\mu^{ai}} - g f^{abc} \bar{\varphi}^{bi} \frac{\delta\Sigma}{\delta b^c} - g f^{abc} \bar{\omega}^{bi} \frac{\delta\Sigma}{\delta \bar{c}^c} - g f^{abc} U_\mu^{bi} \frac{\delta\Sigma}{\delta K_\mu^c} = g f^{abc} A_\mu^b M_\mu^{ci}, \quad (\text{A.11})$$

- the integrated Ward identity

$$\int d^4x \left( c^a \frac{\delta \Sigma}{\delta \omega^{ai}} + \bar{\omega}^{ai} \frac{\delta \Sigma}{\delta \bar{c}^a} + U_\mu^{ai} \frac{\delta \Sigma}{\delta K_\mu^a} \right) = 0, \quad (\text{A.12})$$

- the exact  $\mathcal{R}_{ij}$  symmetry

$$\mathcal{R}_{ij} \Sigma = 0, \quad (\text{A.13})$$

with

$$\mathcal{R}_{ij} = \int d^4x \left( \varphi_i^a \frac{\delta}{\delta \omega_j^a} - \bar{\omega}_j^a \frac{\delta}{\delta \bar{\varphi}_i^a} + V_\mu^{ai} \frac{\delta}{\delta N_\mu^{ai}} - U_\mu^{ai} \frac{\delta}{\delta M_\mu^{ai}} \right). \quad (\text{A.14})$$

## A.2 Algebraic characterization of the counterterm.

Having established all the Ward identities fulfilled by the complete action  $\Sigma$ , we can now turn to the characterization of the most general allowed counterterm  $\Sigma^c$ . Following the algebraic renormalization procedure [21],  $\Sigma^c$  is an integrated local polynomial in the fields and sources with dimension bounded by four, with vanishing ghost number and  $Q_f$ -charge, obeying the following constraints

$$\begin{aligned} \frac{\delta \Sigma^c}{\delta \varphi^{ai}} + \partial_\mu \frac{\delta \Sigma^c}{\delta V_\mu^{ai}} - g f^{abc} \bar{\omega}^{bi} \frac{\delta \Sigma^c}{\delta \bar{c}^c} - g f^{abc} U_\mu^{bi} \frac{\delta \Sigma^c}{\delta K_\mu^c} &= 0, \\ \frac{\delta \Sigma^c}{\delta \bar{\omega}^{ai}} + \partial_\mu \frac{\delta \Sigma^c}{\delta U_\mu^{ai}} - g f^{abc} V_\mu^{bi} \frac{\delta \Sigma^c}{\delta K_\mu^c} &= 0, \\ \frac{\delta \Sigma^c}{\delta \omega^{ai}} + \partial_\mu \frac{\delta \Sigma^c}{\delta N_\mu^{ai}} &= 0, \\ \frac{\delta \Sigma^c}{\delta \bar{\varphi}^{ai}} + \partial_\mu \frac{\delta \Sigma^c}{\delta M_\mu^{ai}} &= 0, \\ \\ \frac{\delta \Sigma}{\delta \bar{c}^a} + \partial_\mu \frac{\delta \Sigma}{\delta K_\mu^a} &= 0, \\ \frac{\delta \Sigma^c}{\delta b^a} &= 0, \end{aligned} \quad (\text{A.15})$$

$$\mathcal{G}^a \Sigma^c = 0, \quad (\text{A.16})$$

$$\int d^4x \left( c^a \frac{\delta \Sigma^c}{\delta \omega^{ai}} + \bar{\omega}^{ai} \frac{\delta \Sigma^c}{\delta \bar{c}^a} + U_\mu^{ai} \frac{\delta \Sigma^c}{\delta K_\mu^a} \right) = 0, \quad (\text{A.17})$$

$$\mathcal{R}_{ij} \Sigma^c = 0, \quad (\text{A.18})$$

and

$$\mathcal{B}_\Sigma \Sigma^c = 0, \quad (\text{A.19})$$

where  $\mathcal{B}_\Sigma$  is the nilpotent linearized Slavnov-Taylor operator

$$\begin{aligned} \mathcal{B}_\Sigma = \int d^4x \left( \frac{\delta \Sigma}{\delta K_\mu^a} \frac{\delta}{\delta A_\mu^a} + \frac{\delta \Sigma}{\delta A_\mu^a} \frac{\delta}{\delta K_\mu^a} + \frac{\delta \Sigma}{\delta L^a} \frac{\delta}{\delta c^a} + \frac{\delta \Sigma}{\delta c^a} \frac{\delta}{\delta L^a} + b^a \frac{\delta}{\delta \bar{c}^a} \right. \\ \left. + \bar{\varphi}_i^a \frac{\delta}{\delta \bar{\omega}_i^a} + \omega_i^a \frac{\delta}{\delta \varphi_i^a} + M_\mu^{ai} \frac{\delta}{\delta U_\mu^{ai}} + N_\mu^{ai} \frac{\delta}{\delta V_\mu^{ai}} + \tau \frac{\delta}{\delta \eta} \right), \end{aligned} \quad (\text{A.20})$$

$$\mathcal{B}_\Sigma \mathcal{B}_\Sigma = 0 . \quad (\text{A.21})$$

As it was shown in [51, 52, 53], the constraints (A.15) imply that  $\Sigma^c$  does not depend on the Lagrange multiplier  $b^a$ , and that the antighost  $\bar{c}^a$  and the  $i$ -valued fields  $\varphi_i^a, \omega_i^a, \bar{\varphi}_i^a, \bar{\omega}_i^a$  can enter only through the combinations

$$\begin{aligned} \tilde{K}_\mu^a &= K_\mu^a + \partial_\mu \bar{c}^a - g f^{abc} \tilde{U}_\mu^{bi} \varphi^{ci} - g f^{abc} V_\mu^{bi} \bar{\omega}^{ci} , \\ \tilde{U}_\mu^{ai} &= U_\mu^{ai} + \partial_\mu \bar{\omega}^{ai} , \\ \tilde{V}_\mu^{ai} &= V_\mu^{ai} + \partial_\mu \varphi^{ai} , \\ \tilde{N}_\mu^{ai} &= N_\mu^{ai} + \partial_\mu \omega^{ai} , \\ \tilde{M}_\mu^{ai} &= V_\mu^{ai} + \partial_\mu \bar{\varphi}^{ai} . \end{aligned} \quad (\text{A.22})$$

Therefore,  $\Sigma^c$  can be parametrized as follows

$$\begin{aligned} \Sigma^c &= S^c(A) + \int d^4x \left( a_1 g f^{abc} L^a c^b c^c + a_2 \tilde{K}_\mu^a \partial_\mu c^a + a_3 g f^{abc} \tilde{K}_\mu^a A_\mu^b c^c + a_4 f^{abc} \tilde{V}_\mu^{ai} \tilde{U}_\mu^{bi} c^c \right. \\ &\quad \left. + a_5 \tilde{V}_\mu^{ai} \tilde{M}_\mu^{ai} + a_6 \tilde{U}_\mu^{ai} \tilde{N}_\mu^{ai} + \frac{a_7}{2} \tau A_\mu^a A_\mu^a + \frac{a_8}{2} \zeta \tau^2 + a_9 \eta A_\mu^a \partial_\mu c^a + a_{10} \eta c^a \partial A^a \right) , \end{aligned} \quad (\text{A.23})$$

where  $S^c(A)$  depends only on the gauge field  $A_\mu^a$ , and with  $a_1, \dots, a_{10}$  arbitrary parameters. Notice, however, that there is no mixing in expression (A.23) between  $\tilde{M}_\mu^{ai}, \tilde{N}_\mu^{ai}, \tilde{V}_\mu^{ai}, \tilde{U}_\mu^{ai}$  and the sources  $\tau, \eta$ . This is due to the dimensionality and to the  $Q_f$ -charge. It is precisely the absence of this mixing that will ensure the renormalizability of the Zwanziger action in the presence of the composite operator  $A_\mu^a A_\mu^a$ . From the ghost equation (A.16) it follows

$$\begin{aligned} a_1 &= a_3 = a_{10} = 0 , \\ a_4 &= -g(a_6 + a_5) . \end{aligned} \quad (\text{A.24})$$

From the equations (A.17) and (A.18) we obtain

$$a_6 = -a_2 . \quad (\text{A.25})$$

Finally, from eq.(A.19) it turns out that

$$\begin{aligned} a_5 &= a_2 , \\ a_9 &= a_7 - a_2 , \end{aligned} \quad (\text{A.26})$$

and

$$S^c(A) = a_0 S_{YM} + a_2 \int d^4x A_\mu^a \frac{\delta S_{YM}}{\delta A_\mu^a} . \quad (\text{A.27})$$

In summary, the most general local invariant counterterm compatible with all Ward identities contains four arbitrary parameters,  $a_0, a_2, a_7, a_8$ , and reads

$$\begin{aligned} \Sigma^c &= a_0 S_{YM} + a_2 \int d^4x \left( A_\mu^a \frac{\delta S_{YM}}{\delta A_\mu^a} + \tilde{K}_\mu^a \partial_\mu c^a + \tilde{V}_\mu^{ai} \tilde{M}_\mu^{ai} - \tilde{U}_\mu^{ai} \tilde{N}_\mu^{ai} \right) \\ &\quad + \int d^4x \left( \frac{a_7}{2} \tau A_\mu^a A_\mu^a + \frac{a_8}{2} \zeta \tau^2 + (a_7 - a_2) \eta A_\mu^a \partial_\mu c^a \right) . \end{aligned} \quad (\text{A.28})$$

### A.3 Stability and renormalization constants.

Having determined the most general local invariant counterterm  $\Sigma^c$  compatible with all Ward identities, it remains to check that the starting action  $\Sigma$  is stable, i.e. that  $\Sigma^c$  can be reabsorbed through the renormalization of the parameters, fields and sources of  $\Sigma$ . According to expression (A.28),  $\Sigma^c$  contains four arbitrary parameters  $a_0, a_2, a_7, a_8$ , which correspond in fact to a multiplicative renormalization of the gauge coupling constant  $g$ , the parameters  $\zeta$ , and of the fields  $\phi = (A_\mu^a, c^a, \bar{c}^a, b^a, \varphi_i^a, \omega_i^a, \bar{\varphi}_i^a, \bar{\omega}_i^a)$  and sources  $\Phi = (K^{a\mu}, L^a, M_\mu^{ai}, N_\mu^{ai}, V_\mu^{ai}, U_\mu^{ai}, \tau, \eta)$ , according to

$$\Sigma(g, \zeta, \phi, \Phi) + \eta \Sigma^c = \Sigma(g_o, \zeta_o, \phi_o, \Phi_o) + O(\eta^2), \quad (\text{A.29})$$

with

$$g_o = Z_g g, \quad \zeta_o = Z_\zeta \zeta, \quad (\text{A.30})$$

and

$$\begin{aligned} \phi_o &= Z_\phi^{1/2} \phi, \\ \Phi_o &= Z_\Phi \Phi. \end{aligned} \quad (\text{A.31})$$

The coefficients  $a_0, a_2$  are easily seen to be related to the renormalization of the gauge coupling constant  $g$  and of the gauge field  $A_\mu^a$ ,

$$\begin{aligned} Z_g &= \left(1 + \eta \frac{a_0}{2}\right), \\ Z_A^{1/2} &= \left(1 + \eta \left(a_2 - \frac{a_0}{2}\right)\right). \end{aligned} \quad (\text{A.32})$$

From expression (A.28) it follows that the Faddeev-Popov ghosts ( $c^a, \bar{c}^a$ ) and the  $i$ -valued fields ( $\varphi_i^a, \omega_i^a, \bar{\varphi}_i^a, \bar{\omega}_i^a$ ) have a common renormalization constant, given by

$$Z_c = Z_{\bar{c}} = Z_\varphi = Z_{\bar{\varphi}} = Z_\omega = Z_{\bar{\omega}} = (1 - \eta a_2) = Z_g^{-1} Z_A^{-1/2}. \quad (\text{A.33})$$

Eq.(A.33) expresses a well-known renormalization property of the Faddeev-Popov ghosts ( $c^a, \bar{c}^a$ ) in the Landau gauge, stemming from the transversality of the gauge propagator and from the factorization of the ghost momentum in the ghost-antighost-gluon vertex. We see therefore that, in the present case, this property holds for the  $i$ -valued fields ( $\varphi_i^a, \omega_i^a, \bar{\varphi}_i^a, \bar{\omega}_i^a$ ) as well. Similarly to the ghost and the  $i$ -valued fields, the renormalization of the sources ( $M_\mu^{ai}, N_\mu^{ai}, V_\mu^{ai}, U_\mu^{ai}$ ) is also determined by the renormalization constants  $Z_g$  and  $Z_A^{1/2}$ , being given by

$$Z_M = Z_N = Z_V = Z_U = Z_g^{-1/2} Z_A^{-1/4}. \quad (\text{A.34})$$

It is worth noticing here that equation (A.34) ensures that the counterterm  $a_2 (V_\mu^{ai} M_\mu^{ai} - U_\mu^{ai} N_\mu^{ai})$  can be automatically reabsorbed by the term  $(-M_\mu^{ai} V_\mu^{ai} + U_\mu^{ai} N_\mu^{ai})$  in the expression (3.5) without the need of introducing new free parameters. Indeed,

$$-M_o V_o = -M V Z_M^2 = -M V Z_g^{-1} Z_A^{-1/2} = -M V + \varepsilon a_2 M V. \quad (\text{A.35})$$

Concerning now the parameters  $a_7, a_8$ , they are easily seen to correspond to a multiplicative renormalization of the local source  $\tau$  and of the parameter  $\zeta$ , according to

$$\begin{aligned} \tau_o &= Z_\tau \tau, \quad Z_\tau = 1 + \eta(a_7 - 2a_2 + a_0), \\ \zeta_o &= Z_\zeta \zeta, \quad Z_\zeta = 1 + \eta(-a_8 - 2a_7 + 4a_2 - 2a_0). \end{aligned} \quad (\text{A.36})$$

Moreover, we would like to underline that there exists even an extra relation, namely

$$Z_\tau = Z_g Z_A^{-1/2} . \quad (\text{A.37})$$

It can be proven by introducing the operator  $A_\mu^2$  through a more sophisticated set of local sources, like it was done in [11]. We will not repeat that analysis here, we only mention that a key ingredient in the proof of relation (A.37) was the presence of the ghost Ward identity, and since the Zwanziger action possesses that identity, eq.(A.5), one can proceed along the lines of [11]. Thus, there are in fact only three independent renormalization factors present.

## B Appendix B.

In this Appendix, we give the detailed analysis of the procedure used to optimize the renormalization scheme and scale dependence, which was summarized in section 5.

### B.1 Preliminaries.

Before coming to the actual computations, let us first discuss some results which will turn out to be useful.

Consider again the action  $S$  of eq.(3.8). Due to the rich symmetry structure of the model, encoded in the Ward identities (A.1)-(A.14), and due to the extra relation (A.37), only three renormalization factors remain to be fixed, namely  $Z_g$ ,  $Z_A$  and  $Z_\zeta$ . Apparently, this means that we would need three renormalization conditions in order to fix a particular renormalization scheme. However, taking a look at the bare action associated with expression eq.(3.8), we would find the following relations

$$\begin{aligned} \zeta_o &= Z_\zeta \zeta , \\ \zeta_o \tau_o^2 &= \bar{\mu}^{-\varepsilon} Z_\zeta \zeta \tau^2 , \\ \tau_o &= Z_\tau \tau , \end{aligned} \quad (\text{B.38})$$

from which it follows that

$$Z_\zeta \zeta = \bar{\mu}^\varepsilon \zeta_o Z_\tau^2 . \quad (\text{B.39})$$

Since the bare quantity  $\zeta_o$  is renormalization scheme and scale independent and since  $\zeta$  always appears in the combination  $Z_\zeta \zeta$  in the action, it follows that only  $Z_g$  and  $Z_A$  are relevant for the effective action, because  $Z_\tau$  can be expressed in terms of these two factors. Consequently, we would only need two renormalization conditions to fix the scheme. Obviously, we can equally well choose to make use of, for example,  $Z_g$  and  $Z_\tau$  as the two independent renormalization factors, corresponding to coupling constant and mass renormalization.

We will change from the  $\overline{\text{MS}}$  to another massless renormalization scheme by means of the following transformations<sup>8</sup>

$$\begin{aligned} \bar{g}^2 &= g^2 (1 + b_0 g^2 + b_1 g^4 + \dots) , \\ \bar{\lambda} &= \lambda (1 + c_0 g^2 + c_1 g^4 + \dots) , \\ \bar{m}^2 &= m^2 (1 + d_0 g^2 + d_1 g^4 + \dots) , \end{aligned} \quad (\text{B.40})$$

---

<sup>8</sup>Barred quantities refer to the  $\overline{\text{MS}}$  scheme.

where the parameters  $b_i$ ,  $c_i$  and  $d_i$  label the new scheme. However, we should keep in mind that the renormalization of the Gribov parameter  $\lambda$  is not independent of that of  $g^2$  and  $m^2$ . Eliminating  $\gamma_A(g^2)$  between eqns.(4.19) and (4.20), yields

$$\gamma_\lambda(g^2) = \frac{1}{4} \left( \frac{\beta(g^2)}{g^2} - \gamma_{m^2}(g^2) \right). \quad (\text{B.41})$$

This relation, valid to all orders of perturbation theory, implies the existence of relationships between the coefficients  $b_i$ ,  $c_i$  and  $d_i$ . For further use, we shall explicitly construct the relation between  $b_0$ ,  $c_0$  and  $d_0$ . Let us adopt as parametrization of  $\beta(g^2)$ ,  $\gamma_{m^2}(g^2)$  and  $\gamma_\lambda(g^2)$

$$\begin{aligned} \beta(g^2) &= -2(\beta_0 g^4 + \beta_1 g^6 + \dots), \\ \gamma_{m^2}(g^2) &= \gamma_0 g^2 + \gamma_1 g^4 + \dots, \\ \gamma_\lambda(g^2) &= \lambda_0 g^2 + \lambda_1 g^4 + \dots, \end{aligned} \quad (\text{B.42})$$

and an analogous one in the case of the  $\overline{\text{MS}}$  scheme. Then, one computes

$$\begin{aligned} \bar{\mu} \frac{\partial \bar{\lambda}}{\partial \bar{\mu}} &= \bar{\mu} \frac{\partial}{\partial \bar{\mu}} [\lambda (1 + c_0 g^2 + \dots)] \\ &= \dots \\ &= \lambda (\lambda_0 g^2 + (\lambda_1 + c_0 \lambda_0 - 2\beta_0 c_0) g^4 + \dots), \end{aligned} \quad (\text{B.43})$$

which can be expressed in terms of  $\gamma_i$  and  $\beta_i$  by exploiting the relation (B.41). We find

$$\bar{\mu} \frac{\partial \bar{\lambda}}{\partial \bar{\mu}} = \lambda \left[ \frac{-2\beta_0 - \gamma_0}{4} g^2 + \left( \frac{-2\beta_1 - \gamma_1}{4} + c_0 \frac{-2\beta_0 - \gamma_0}{4} - 2\beta_0 c_0 \right) g^4 + \dots \right]. \quad (\text{B.44})$$

We can also calculate  $\bar{\mu} \frac{d\bar{\lambda}}{d\bar{\mu}}$  by first exploiting the relation (B.41), obtaining

$$\begin{aligned} \bar{\mu} \frac{\partial \bar{\lambda}}{\partial \bar{\mu}} &= \frac{1}{4} [(-2\beta_0 - \gamma_0) \bar{g}^2 + (-2\beta_1 - \bar{\gamma}_1) \bar{g}^4 + \dots] [\lambda (1 + c_0 g^2 + \dots)] \\ &= \dots \\ &= \frac{1}{4} [(-2\beta_0 - \gamma_0) g^2 + (c_0(-2\beta_0 - \gamma_0) - 2\beta_1 - \gamma_1 - 2\beta_0(-d_0 + b_0)) g^4 + \dots] \end{aligned} \quad (\text{B.45})$$

In the previous expression, we had to express  $\bar{\gamma}_1$  in terms of  $\gamma_1$ ; a task accomplished by using the relation

$$\bar{\gamma}_1 = \gamma_1 - 2\beta_0 d_0 - \gamma_0 b_0, \quad (\text{B.46})$$

which can be obtained along the same lines of the previous calculations. It should also be noted that  $\gamma_0$ ,  $\beta_0$  and  $\beta_1$  are renormalization scheme independent quantities. Thus, the identification of eqns.(B.44) and (B.45) gives the desired relation, given by

$$c_0 = \frac{1}{4} (b_0 - d_0). \quad (\text{B.47})$$

We now perform the transformations (B.40) on the action (4.14), which was calculated in the  $\overline{\text{MS}}$  scheme, to obtain it in a general scheme.

$$\begin{aligned} \Gamma &= -\frac{(N^2 - 1) \lambda^4}{2g^2 N} (1 + 4c_0 g^2 - b_0 g^2) + \frac{\zeta_0 m^4}{2g^2} \left( 1 - \frac{\zeta_1}{\zeta_0} g^2 + 2d_0 g^2 - b_0 g^2 \right) \\ &+ \frac{3(N^2 - 1)}{256\pi^2} \left[ \left( m^2 + \sqrt{m^4 - \lambda^4} \right)^2 \left( \ln \frac{m^2 + \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2} - \frac{5}{6} \right) \right. \\ &\left. + \left( m^2 - \sqrt{m^4 - \lambda^4} \right)^2 \left( \ln \frac{m^2 - \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2} - \frac{5}{6} \right) \right], \end{aligned} \quad (\text{B.48})$$

while the gap equations now read

$$\begin{aligned}
\frac{\partial \Gamma}{\partial \lambda} &= -\frac{2(N^2-1)}{g^2 N} \lambda^3 (1 + 4c_0 g^2 - b_0 g^2) + \frac{3(N^2-1)\lambda^3}{256\pi^2} \left[ \frac{8}{3} \right. \\
&\quad \left. - 4 \frac{(m^2 + \sqrt{m^4 - \lambda^4})}{\sqrt{m^4 - \lambda^4}} \ln \frac{m^2 + \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2} + 4 \frac{(m^2 - \sqrt{m^4 - \lambda^4})}{\sqrt{m^4 - \lambda^4}} \ln \frac{m^2 - \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2} \right], \\
\frac{\partial \Gamma}{\partial m^2} &= \frac{\zeta_0 m^2}{g^2} \left( 1 - \frac{\zeta_1}{\zeta_0} g^2 + 2d_0 g^2 - b_0 g^2 \right) \\
&\quad + \frac{3(N^2-1)}{256\pi^2} \left[ 2 \left( m^2 + \sqrt{m^4 - \lambda^4} \right) \left( 1 + \frac{m^2}{\sqrt{m^4 - \lambda^4}} \right) \ln \frac{m^2 + \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2} \right. \\
&\quad \left. + 2 \left( m^2 - \sqrt{m^4 - \lambda^4} \right) \left( 1 - \frac{m^2}{\sqrt{m^4 - \lambda^4}} \right) \ln \frac{m^2 - \sqrt{m^4 - \lambda^4}}{2\bar{\mu}^2} - \frac{8}{3} m^2 \right].
\end{aligned} \tag{B.49}$$

We mention that, in the case in which  $m^2 \geq 0$ , similar algebraic manipulations as those leading to the condition (4.37), give a more general equation

$$\frac{68}{39} \left( \frac{16\pi^2}{g^2 N} \right) + \frac{161}{39} + \frac{16\pi^2}{N} \left( \frac{32}{3} c_0 - \frac{68}{39} b_0 - \frac{24}{13} d_0 \right) = \frac{t}{\sqrt{1-t}} \ln \frac{t}{(1 + \sqrt{1-t})^2}, \tag{B.50}$$

or, using the relation (B.47),

$$\frac{68}{39} \left( \frac{16\pi^2}{g^2 N} \right) + \frac{161}{39} + \frac{16\pi^2}{N} \left( \frac{12}{13} b_0 - \frac{176}{39} d_0 \right) = \frac{t}{\sqrt{1-t}} \ln \frac{t}{(1 + \sqrt{1-t})^2}. \tag{B.51}$$

From this expression, it is apparent that a sensible solution with  $m^2 > 0$  might exist, depending on the values of the renormalization parameters  $d_0$  ( $\sim$  mass renormalization) and  $b_0$  ( $\sim$  coupling constant renormalization).

Frequently used are the so-called physical renormalization schemes whereby, loosely speaking, one demands that the quantum corrected quantities reduce to the tree level values at a certain scale  $\bar{\mu}$ . However, it turns out that such an approach is not particularly useful to implement in the current case due to the presence of the several scales. Therefore, the question arises how one can make a somewhat motivated choice for the arbitrary parameters, labeling a certain renormalization scheme. In the next subsection we shall discuss a way to reduce the freedom in the choice of the renormalization parameters. The method relies on the possibility of performing an optimization of the renormalization scheme dependence, as illustrated in [70, 71].

## B.2 Optimization of the renormalization scheme.

Consider a quantity  $\varrho$  that runs according to

$$\bar{\mu} \frac{d\varrho}{d\bar{\mu}} = \gamma_\varrho(g^2) \varrho, \tag{B.52}$$

where

$$\gamma_\varrho(g^2) = \gamma_{\varrho,0} g^2 + \gamma_{\varrho,1} g^4 + \dots \tag{B.53}$$



To  $\varrho$ , we can associate a quantity  $\widehat{\varrho}$  that does not depend on the choice of the renormalization scheme and which is scale independent. It is defined as

$$\widehat{\varrho} = \mathcal{F}_\varrho(g^2)\varrho, \quad (\text{B.54})$$

whereby

$$\bar{\mu} \frac{d}{d\bar{\mu}} \mathcal{F}_\varrho(g^2) = -\gamma_\varrho(g^2) \mathcal{F}_\varrho(g^2). \quad (\text{B.55})$$

It is apparent that  $\widehat{\varrho}$  will not depend on the scale  $\bar{\mu}$ . It can also be checked [70, 71] that  $\widehat{\varrho}$  is left unmodified by a change of the renormalization scheme, implemented through transformations analogous to those of eqs.(B.40). The equation (B.55) can be solved in a series expansion in  $g^2$  by noticing that

$$\bar{\mu} \frac{d}{d\bar{\mu}} \mathcal{F}_\varrho(g^2) \equiv \beta(g^2) \frac{d}{dg^2} \mathcal{F}_\varrho(g^2). \quad (\text{B.56})$$

Then, the above differential equation can be solved in a series expansion in  $g^2$ , more precisely by

$$\mathcal{F}_\varrho(g^2) = (g^2)^{\frac{\gamma_{\varrho,0}}{2\beta_0}} \left( 1 + \frac{1}{2} \left( \frac{\gamma_{\varrho,1}}{\beta_0} - \frac{\beta_1 \gamma_{\varrho,0}}{\beta_0^2} \right) g^2 + \dots \right). \quad (\text{B.57})$$

Consider once more the  $\overline{\text{MS}}$  action  $\Gamma$  given in eq.(4.14). We shall now replace the  $\overline{\text{MS}}$  variables  $\overline{m}^2$  and  $\overline{\lambda}$  by their renormalization scheme and scale independent counterparts  $\widehat{m}^2$  and  $\widehat{\lambda}$ , which are obtained as before. By inverting eq.(B.57), one has

$$\overline{m}^2 = (\overline{g}^2)^{-\frac{\gamma_0}{2\beta_0}} \left( 1 - \frac{1}{2} \left( \frac{\overline{\gamma}_1}{\beta_0} - \frac{\beta_1 \gamma_0}{\beta_0^2} \right) \overline{g}^2 + \dots \right) \widehat{m}^2, \quad (\text{B.58})$$

$$\overline{\lambda} = (\overline{g}^2)^{-\frac{\lambda_0}{2\beta_0}} \left( 1 - \frac{1}{2} \left( \frac{\overline{\lambda}_1}{\beta_0} - \frac{\beta_1 \lambda_0}{\beta_0^2} \right) \overline{g}^2 + \dots \right) \widehat{\lambda}. \quad (\text{B.59})$$

Moreover, introducing the notations

$$a = -\frac{\gamma_0}{2\beta_0}, \quad b = -\frac{\lambda_0}{\beta_0}, \quad (\text{B.60})$$

$$A = -\left( \frac{\overline{\gamma}_1}{\beta_0} - \frac{\beta_1 \gamma_0}{\beta_0^2} \right), \quad B = -2 \left( \frac{\overline{\lambda}_1}{\beta_0} - \frac{\beta_1 \lambda_0}{\beta_0^2} \right), \quad (\text{B.61})$$

the one-loop action is rewritten as

$$\begin{aligned} \Gamma = & -\frac{(N^2-1)}{2N} (\overline{g}^2)^{2b} \widehat{\lambda}^4 \left( \frac{1}{\overline{g}^2} + B \right) + \frac{\zeta_0}{2} \widehat{m}^4 (\overline{g}^2)^{2a} \left( \frac{1}{\overline{g}^2} + A - \frac{\zeta_1}{\zeta_0} \right) + \frac{3(N^2-1)}{256\pi^2} \times \\ & \left[ \left( \widehat{m}^2 (\overline{g}^2)^a + \sqrt{\widehat{m}^4 (\overline{g}^2)^{2a} - \widehat{\lambda}^4 (\overline{g}^2)^{2b}} \right)^2 \left( \ln \frac{\widehat{m}^2 (\overline{g}^2)^a + \sqrt{\widehat{m}^4 (\overline{g}^2)^{2a} - \widehat{\lambda}^4 (\overline{g}^2)^{2b}}}{2\bar{\mu}^2} - \frac{5}{6} \right) \right. \\ & \left. + \left( \widehat{m}^2 (\overline{g}^2)^a - \sqrt{\widehat{m}^4 (\overline{g}^2)^{2a} - \widehat{\lambda}^4 (\overline{g}^2)^{2b}} \right)^2 \left( \ln \frac{\widehat{m}^2 (\overline{g}^2)^a - \sqrt{\widehat{m}^4 (\overline{g}^2)^{2a} - \widehat{\lambda}^4 (\overline{g}^2)^{2b}}}{2\bar{\mu}^2} - \frac{5}{6} \right) \right]. \end{aligned} \quad (\text{B.62})$$

The action (B.62) is still written in terms of the  $\overline{\text{MS}}$  coupling  $\overline{g}^2$ . Performing the first transformation of (B.40),  $\Gamma$  can be reexpressed as

$$\begin{aligned}
\Gamma &= -\frac{(N^2 - 1)}{2N} (g^2)^{2b} \widehat{\lambda}^4 \left( \frac{1}{g^2} + B - b_0 + 2bb_0 \right) \\
&+ \frac{\zeta_0}{2} \widehat{m}^4 (g^2)^{2a} \left( \frac{1}{g^2} + A - b_0 + 2ab_0 - \frac{\zeta_1}{\zeta_0} \right) + \frac{3(N^2 - 1)}{256\pi^2} \times \\
&\left[ \left( \widehat{m}^2 (g^2)^a + \sqrt{\widehat{m}^4 (g^2)^{2a} - \widehat{\lambda}^4 (g^2)^{2b}} \right)^2 \left( \ln \frac{\widehat{m}^2 (g^2)^a + \sqrt{\widehat{m}^4 (g^2)^{2a} - \widehat{\lambda}^4 (g^2)^{2b}}}{2\overline{\mu}^2} - \frac{5}{6} \right) \right. \\
&\left. + \left( \widehat{m}^2 (g^2)^a - \sqrt{\widehat{m}^4 (g^2)^{2a} - \widehat{\lambda}^4 (g^2)^{2b}} \right)^2 \left( \ln \frac{\widehat{m}^2 (g^2)^a - \sqrt{\widehat{m}^4 (g^2)^{2a} - \widehat{\lambda}^4 (g^2)^{2b}}}{2\overline{\mu}^2} - \frac{5}{6} \right) \right].
\end{aligned} \tag{B.63}$$

So far, we have constructed an action which is written in terms of renormalization scale and scheme independent variables  $\widehat{\lambda}$  and  $\widehat{m}^2$  and the coupling constant  $g^2(\overline{\mu})$ . This is a certain improvement, since we are not faced anymore with a choice of the parameters  $d_i$ , related to the renormalization of the Gribov and mass parameter. The remaining freedom in the choice of the renormalization scheme resides in the coupling constant, labeled by the parameters  $b_0, b_1, \dots$ , and in the scale  $\overline{\mu}$ . Of course, the higher order coefficients  $b_i$ ,  $i = 1, \dots$  do not show up here, since we have restricted ourselves to the one-loop level. Nevertheless, we will perform one more step, since the dependence on the coupling constant renormalization can be reduced to solely  $b_0$ , by expanding the perturbative series in inverse powers of

$$x \equiv \beta_0 \ln \frac{\overline{\mu}^2}{\Lambda^2}, \tag{B.64}$$

rather than in terms of  $g^2$ . For another illustration of this, see e.g. [70, 71]. The coupling constant  $g^2$  can be replaced by  $x$  since  $g^2$  is explicitly determined by

$$g^2 = \frac{1}{x} \left( 1 - \frac{\beta_1}{\beta_0} \frac{\ln \frac{x}{\beta_0}}{x} + \dots \right). \tag{B.65}$$

In [69], the relation between the scale parameter  $\Lambda$ , corresponding to a certain coupling constant renormalization, and that of the  $\overline{\text{MS}}$  scheme,  $\Lambda_{\overline{\text{MS}}}$ , was found to be

$$\Lambda = e^{-\frac{b_0}{2\beta_0}} \Lambda_{\overline{\text{MS}}}. \tag{B.66}$$

One finally gets the expression (5.1). We notice that this alternative expansion is correct up to order  $\left(\frac{1}{x}\right)^0$ .

In principle, we can solve the two equations (5.2)-(5.3) for the two quantities  $\widehat{m}_*$  and  $\widehat{\lambda}_*$ , which will be functions of the two remaining parameters  $\overline{\mu}$  and  $b_0$ . However, by construction, we know that  $\widehat{m}$  as well as  $\widehat{\lambda}$  should be independent of the renormalization scale and scheme order by order. This gives us an interesting way to fix these parameters by demanding that the solutions  $\widehat{m}_*(\overline{\mu}, b_0)$  and  $\widehat{\lambda}_*(\overline{\mu}, b_0)$  depend minimally on  $b_0$  and  $\overline{\mu}$ . Since this would give a quite complicated set of equations to solve, we can make life somewhat easier by reasonably choosing the scale<sup>9</sup>  $\overline{\mu}$

<sup>9</sup>This can be motivated thanks to the scale independence of the  $\widehat{\phantom{m}}$ -quantities.

in the gap equations (5.2)-(5.3). In analogy to the choice for  $\bar{\mu}^2$  done in the previous equation (4.38), we shall now set

$$\bar{\mu}^2 = \left| \frac{\widehat{m}^2 x^{-a} + \sqrt{\widehat{m}^4 x^{-2a} - \widehat{\lambda}^4 x^{-2b}}}{2} \right|, \quad (\text{B.67})$$

In order to proceed, we still have two quantities at our disposal to fix the remaining parameter  $b_0$ . In fact, we can also take the vacuum energy  $E_{\text{vac}}$  in consideration since, being a physical quantity, it should depend minimally on the renormalization scheme and scale. Therefore, we could determine the value for  $b_0$  by demanding that

$$\Upsilon(b_0) \equiv \left| \frac{\partial \widehat{\lambda}_*^4}{\partial b_0} \right| + \left| \frac{\partial \widehat{m}_*^4}{\partial b_0} \right| + \left| \frac{\partial E_{\text{vac}}}{\partial b_0} \right|, \quad (\text{B.68})$$

is minimal w.r.t. the parameter  $b_0$ . This seems to be a reasonable candidate. When its dependence on  $b_0$  is small, then the dependence of  $\widehat{m}$ ,  $\widehat{\lambda}$  and  $E_{\text{vac}}$  on  $b_0$  is necessarily small too. The ideal situation would be that  $\Upsilon$  is zero for a certain  $b_0$ . If no such an ideal  $b_0$  would exist, we weaken the condition by requiring that  $\Upsilon$  is as small as possible. The condition (B.68) to fix  $b_0$  can be considered as some kind of *principle of minimal sensitivity* à la Stevenson [72]. An alternative that is sometimes used is a *fastest apparent convergence criterion*, where it is demanded that the quantum corrections are as small as possible compared to the tree level value. For example, if we denote by  $\Gamma^{[0]}$  the action to order  $(\frac{1}{x})^{-1}$  and by  $\Gamma^{[1]}$  to order  $(\frac{1}{x})^0$ , we could demand that

$$\left| \frac{\Gamma^{[1]} - \Gamma^{[0]}}{\Gamma^{[0]}} \right| \quad (\text{B.69})$$

is as small as possible when the parameters fulfill the gap equation describing the vacuum of the theory.

Before continuing with explicit calculations, let us just remark here that the other logarithm, namely  $\ln \frac{\widehat{m}^2 x^{-a} - \sqrt{\widehat{m}^4 x^{-2a} - \widehat{\lambda}^4 x^{-2b}}}{2\bar{\mu}^2}$ , could become large for a small argument, thus when  $\widehat{\lambda}^4 x^{-2b}$  would be small compared to  $\widehat{m}^4 x^{-2a}$ . However, it is harmless since it appears in the form of  $u \ln u$ , while we know that  $u \ln u|_{u \approx 0} \approx 0$ .

### B.3 Numerical results.

Let us first give some numerical factors we need. From e.g. [65], we infer that

$$\beta_1 = \frac{34}{3} \left( \frac{N}{16\pi^2} \right)^2, \quad \gamma_0 = -\frac{3}{2} \frac{N}{16\pi^2}, \quad \gamma_1 = -\frac{95}{24} \left( \frac{N}{16\pi^2} \right)^2, \quad (\text{B.70})$$

and hence, from the relation (B.41),

$$\lambda_0 = -\frac{35}{24} \frac{N}{16\pi^2}, \quad \lambda_1 = -\frac{449}{96} \left( \frac{N}{16\pi^2} \right)^2. \quad (\text{B.71})$$

This means that, for any  $N$ , the quantities  $a$  and  $b$  in eq.(B.60) are found to be

$$a = \frac{9}{44}, \quad b = \frac{35}{88}. \quad (\text{B.72})$$

It is instructive to consider once more the original Gribov-Zwanziger model by setting  $\widehat{m} \equiv 0$  and by solving the gap equation (5.2). If  $\widehat{\lambda}_*$  is a solution of this equation, then it is not difficult to show that the corresponding vacuum energy is given by

$$E_{\text{vac}} = \frac{3(N^2 - 1) \widehat{\lambda}_*^4}{64\pi^2 \cdot 4}, \quad (\text{B.73})$$

for any choice of  $\overline{\mu}^2$ . Thus, also with the improved perturbative expansion, the vacuum energy of the original Gribov-Zwanziger is always nonnegative at the lowest order.

Let us return to the model we were investigating. We solved the gap equations stemming from (5.2)-(5.3) numerically.

Let us first search for a possible solution of the gap equation in the region of space determined by  $\widehat{m}^4 x^{-2a} \geq \widehat{\lambda}^4 x^{-2b}$ . Taking a look at the action (5.1), it might be clear that the gap equations derived from it will be coupled and hence quite complicated to solve numerically. From the calculational point of view, it is useful to introduce new variables, defined by

$$\omega_1 = \frac{\widehat{m}^2 x^{-a} + \sqrt{\widehat{m}^4 x^{-2a} - \widehat{\lambda}^4 x^{-2b}}}{2}, \quad (\text{B.74})$$

$$\omega_2 = \frac{\widehat{m}^2 x^{-a} - \sqrt{\widehat{m}^4 x^{-2a} - \widehat{\lambda}^4 x^{-2b}}}{2}, \quad (\text{B.75})$$

with the inverse transformation

$$\begin{aligned} \widehat{m}^2 x^{-a} &= \omega_1 + \omega_2, \\ \widehat{\lambda}^4 x^{-2b} &= 4\omega_1 \omega_2. \end{aligned} \quad (\text{B.76})$$

This defines a mapping from the space  $\widehat{m}^4 x^{-2a} \geq \widehat{\lambda}^4 x^{-2b} > 0$  to  $\omega_1 \geq \omega_2 > 0$ . One checks that the gap equations (5.2)-(5.3) are equivalent to

$$\left( \frac{\omega_1}{\omega_1 - \omega_2} \frac{\partial}{\partial \omega_1} - \frac{\omega_2}{\omega_1 - \omega_2} \frac{\partial}{\partial \omega_2} \right) \Gamma(\omega_1, \omega_2) = 0, \quad (\text{B.77})$$

$$\left( \frac{1}{\omega_1 - \omega_2} \frac{\partial}{\partial \omega_1} - \frac{1}{\omega_1 - \omega_2} \frac{\partial}{\partial \omega_2} \right) \Gamma(\omega_1, \omega_2) = 0. \quad (\text{B.78})$$

We notice that the case in which  $\omega_1$  and  $\omega_2$  would become equal, i.e.  $\widehat{m}^4 x^{-2a} = \widehat{\lambda}^4 x^{-2b}$ , should be treated with some extra care. Let us therefore first assume that  $\omega_1 > \omega_2$ . Then the two equations (B.77)-(B.78) can be recombined to

$$\frac{\partial}{\partial \omega_1} \Gamma = 0, \quad (\text{B.79})$$

$$\frac{\partial}{\partial \omega_2} \Gamma = 0. \quad (\text{B.80})$$

The action  $\Gamma(\omega_1, \omega_2)$  is explicitly given by

$$\begin{aligned} \Gamma &= -2 \frac{(N^2 - 1)}{N} \mathcal{U}_1 \omega_1 \omega_2 + \frac{\zeta_0}{2} \mathcal{U}_2 (\omega_1 + \omega_2)^2 \\ &+ \frac{3(N^2 - 1)}{64\pi^2} \left[ \omega_1^2 \left( \ln \frac{\omega_1}{\overline{\mu}^2} - \frac{5}{6} \right) + \omega_2^2 \left( \ln \frac{\omega_2}{\overline{\mu}^2} - \frac{5}{6} \right) \right]. \end{aligned} \quad (\text{B.81})$$

where

$$\mathcal{U}_1 = x + B + (1 - 2b) \left( \frac{\beta_1}{\beta_0} \ln \frac{x}{\beta_0} - b_0 \right), \quad (\text{B.82})$$

$$\mathcal{U}_2 = x + A - \frac{\zeta_1}{\zeta_0} + (1 - 2a) \left( \frac{\beta_1}{\beta_0} \ln \frac{x}{\beta_0} - b_0 \right). \quad (\text{B.83})$$

It is not difficult to work out the gap equations (B.79)-(B.80), being given by

$$-2 \frac{N^2 - 1}{N} \mathcal{U}_1 \omega_2 + \zeta_0 \mathcal{U}_2 (\omega_1 + \omega_2) + \frac{3(N^2 - 1)\omega_1}{32\pi^2} \left( -\frac{1}{3} + \ln \frac{\omega_1}{\bar{\mu}^2} \right) = 0, \quad (\text{B.84})$$

$$-2 \frac{N^2 - 1}{N} \mathcal{U}_1 \omega_1 + \zeta_0 \mathcal{U}_2 (\omega_1 + \omega_2) + \frac{3(N^2 - 1)\omega_2}{32\pi^2} \left( -\frac{1}{3} + \ln \frac{\omega_2}{\bar{\mu}^2} \right) = 0. \quad (\text{B.85})$$

From the explicit expression of the gap equations and of the action itself in terms of  $\omega_1$  and  $\omega_2$ , the advantages of using these variables should be obvious, since we can decouple the two gap equations. Explicitly, since  $\bar{\mu}^2 = \omega_1$ , one finds from eq.(B.84),

$$\omega_2 = \frac{\frac{N^2 - 1}{32\pi^2} - \zeta_0 \mathcal{U}_2}{-2 \frac{N^2 - 1}{N} \mathcal{U}_1 + \zeta_0 \mathcal{U}_2} \omega_1, \quad (\text{B.86})$$

which can be substituted in the second gap equation (B.85), yielding an equation for  $\omega_1$  which does not contain  $\omega_2$  anymore. The nominator of eq.(B.86) is different from zero, since filling in the numbers gives

$$-2 \frac{N^2 - 1}{N} \mathcal{U}_1 + \zeta_0 \mathcal{U}_2 = \frac{N^2 - 1}{4576} \left( -\frac{975}{\pi^2} - \frac{5984}{N} x \right) \neq 0. \quad (\text{B.87})$$

where we kept in mind that for a meaningful result,  $x \sim \frac{1}{g^2}$ , should be positive.

A numerical investigation of the gap equation (B.85) using eq.(B.86) revealed that there are no zeros. We conclude that there are no solutions with  $\hat{m}^4 x^{-2a} > \hat{\lambda}^4 x^{-2b}$ .

Next, let us find out if a possible solution with  $\hat{m}^4 x^{-2a} = \hat{\lambda}^4 x^{-2b}$  or  $\omega_1 = \omega_2$  might exist. We explicitly evaluate the gap equations (B.77)-(B.78), where now  $\bar{\mu}^2 = \omega_1$ ,

$$\zeta_0 \mathcal{U}_2 (\omega_1 + \omega_2) - \frac{N^2 - 1}{32\pi^2} (\omega_1 + \omega_2) - \frac{3(N^2 - 1)}{32\pi^2} \frac{\omega_2^2}{\omega_1 - \omega_2} \ln \frac{\omega_2}{\omega_1} = 0, \quad (\text{B.88})$$

$$2 \frac{N^2 - 1}{N} \mathcal{U}_1 - \frac{N^2 - 1}{32\pi^2} - \frac{3(N^2 - 1)}{32\pi^2} \frac{\omega_2}{\omega_1 - \omega_2} \ln \frac{\omega_2}{\omega_1} = 0. \quad (\text{B.89})$$

From the foregoing expressions, we infer that the limit  $\omega_1 \rightarrow \omega_2$  exists, giving rise to

$$2\zeta_0 \mathcal{U}_2 - \frac{2(N^2 - 1)}{32\pi^2} + \frac{3(N^2 - 1)}{32\pi^2} = 0, \quad (\text{B.90})$$

$$2 \frac{N^2 - 1}{N} \mathcal{U}_1 - \frac{N^2 - 1}{32\pi^2} + \frac{3(N^2 - 1)}{32\pi^2} = 0. \quad (\text{B.91})$$

This means that we have two equations to solve for the single quantity  $\omega_1$ , which is present in  $\mathcal{U}_1$  and  $\mathcal{U}_2$  through the quantity  $x$ . It would be an extreme coincidence if these two different equations, which can be rewritten as

$$\frac{18}{13} \mathcal{U}_2 = -\frac{N}{32\pi^2}, \quad (\text{B.92})$$

$$\mathcal{U}_1 = -\frac{N}{32\pi^2}. \quad (\text{B.93})$$




Figure 5: The solution  $\omega_1 = \omega_2$  as a function of  $b_0$  of eq.(B.92), top curve, and eq.(B.93), bottom curve, in units  $\Lambda_{\overline{\text{MS}}} = 1$ . Clearly, these two curves do not coincide.

possess a common solution. That this is not the case can be inferred from the numerical solutions of both equations (B.92) and (B.93), shown in Figure 5.

As a final step, we should investigate if there is a solution in the region  $\widehat{m}^4 x^{-2a} < \widehat{\lambda}^4 x^{-2b}$ . We can still define the coordinates  $\omega_1$  and  $\omega_2$  by

$$\omega_1 = \frac{\widehat{m}^2 x^{-a} + i\sqrt{-\widehat{m}^4 x^{-2a} + \widehat{\lambda}^4 x^{-2b}}}{2}, \quad (\text{B.94})$$

$$\omega_2 = \frac{\widehat{m}^2 x^{-a} - i\sqrt{-\widehat{m}^4 x^{-2a} + \widehat{\lambda}^4 x^{-2b}}}{2}. \quad (\text{B.95})$$

In this case,  $\omega_1$  and  $\omega_2$  are complex conjugate. Henceforth, it would be more appropriate to use the modulus  $R$  and the argument  $\phi$ ,  $\phi \in ]-\pi, \pi]$ , defined by

$$Re^{i\phi} = \omega_1, \quad (\text{B.96})$$

$$Re^{-i\phi} = \omega_2, \quad (\text{B.97})$$

If the argument  $\phi$  is so that  $|\phi| > \frac{\pi}{2}$ , then  $\widehat{m}^2 x^{-a} < 0$ . As a consequence, the estimate for  $\langle A_\mu^2 \rangle$  will be positive.

Most of the foregoing analysis can be repeated. The action (B.81) is rewritten in terms of  $R$  and  $\phi$  by

$$\begin{aligned} \Gamma &= -2 \frac{(N^2 - 1)}{N} \mathcal{U}_1 R^2 + 2\zeta_0 \mathcal{U}_2 R^2 \cos^2 \phi \\ &+ \frac{3R^2 (N^2 - 1)}{32\pi^2} \left[ \cos(2\phi) \left( \ln \frac{R}{\overline{\mu}^2} - \frac{5}{6} \right) - \phi \sin(2\phi) \right]. \end{aligned} \quad (\text{B.98})$$

The gap equations (B.84)-(B.85) reduce to

$$-2 \frac{N^2 - 1}{N} \mathcal{U}_1 R e^{-i\phi} + \zeta_0 \mathcal{U}_2 R (e^{i\phi} + e^{-i\phi}) + \frac{3(N^2 - 1) R e^{i\phi}}{32\pi^2} \left( -\frac{1}{3} + i\phi \right) = 0, \quad (\text{B.99})$$

and its complex conjugate. With the parametrization (B.96), we have  $\overline{\mu}^2 = R$ .

We must solve the following two real equations<sup>10</sup> for  $\phi$  and  $R$ .

$$-2\frac{N^2-1}{N}\bar{U}_1 \cos \phi + 2\zeta_0\bar{U}_2 \cos \phi + \frac{3(N^2-1)}{32\pi^2} \left( -\frac{\cos \phi}{3} - \phi \sin \phi \right) = 0, \quad (\text{B.100})$$

$$2\frac{N^2-1}{N}\bar{U}_1 \sin \phi + \frac{3(N^2-1)}{32\pi^2} \left( -\frac{\sin \phi}{3} + \phi \cos \phi \right) = 0. \quad (\text{B.101})$$

We can divide these equations<sup>11</sup> by  $\cos \phi$  to obtain

$$-2\frac{N^2-1}{N}\bar{U}_1 + 2\zeta_0\bar{U}_2 + \frac{3(N^2-1)}{32\pi^2} \left( -\frac{1}{3} - \phi \tan \phi \right) = 0, \quad (\text{B.102})$$

$$2\frac{N^2-1}{N}\bar{U}_1 \tan \phi + \frac{3(N^2-1)}{32\pi^2} \left( -\frac{\tan \phi}{3} + \phi \right) = 0. \quad (\text{B.103})$$

These equations can also be decoupled. The most efficient way to proceed is to eliminate  $R$  between these two equations to obtain an equation for  $\phi$ , as the range in we must search for a solution is limited for this angle. The equation for  $\phi$  finally becomes

$$\frac{-90985N - 107712\pi^2 b_0 + 12N \left( 484\phi \cot \phi + 1734 \ln \left( \frac{-117(50+11\phi \csc \phi \sec \phi)}{8228} \right) - 1573\phi \tan \phi \right)}{107712\pi^2} = 0 \quad (\text{B.104})$$

while the value of  $R$  is obtained from

$$x \equiv \beta_0 \ln \frac{R}{\Lambda_{\text{MS}}^2} + b_0 = -\frac{1950 + 429\phi \csc \phi \sec \phi}{11968\pi^2} N. \quad (\text{B.105})$$

We shall concentrate on the case  $N = 3$ . Depending on the value of the parameter  $b_0$ , there is more than one solution possible. In Figure 6, we have plotted the expression (B.104) for several values of the parameter  $b_0$ , namely  $b_0 = 0.25, 0, -0.25, -0.3, -0.33564\dots, -0.41594\dots, -0.5$ . It is possible to obtain those values of  $b_0$  where the number of solutions change. If we consider the plots of Figure 6, it is apparent that for each  $b_0$ , the corresponding curve possesses two extremal values. The number of solution exactly changes at those values of  $b_0$  where the curve becomes tangent to the  $\phi$ -axis. An explicit evaluation learns that his occurs at  $b_0 = -0.41595\dots$ , where  $\phi = 2.26407\dots$  and at  $b_0 = -0.33564\dots$  where  $\phi = 2.62545$ . It is important to know these numbers to a high enough accuracy, to instruct the computer in which  $\phi$ -interval it can search for a solution. If the initial values are not chosen in an appropriate way, the iterations will jump between the different branches of solutions and there will be no convergence to any of them. There is a single solution  $\phi$  if  $b_0 > -0.33564\dots$  or  $b_0 < -0.41595\dots$ . If  $-0.41595\dots < b_0 < -0.33564\dots$ , there are three solutions, while for  $b_0 = -0.41595\dots$  and  $b_0 = -0.33564\dots$  there are two solutions. In Figure 7, we have displayed the solution for  $\phi$  and  $R$ . To determine the solution  $\phi$  which characterizes the vacuum, we should take that one which gives us the absolute minimum of the energy functional  $\Gamma$ , which was shown in Figure 3.

As a final remark, we would like to notice that the same decomposition as in eq.(4.7) could also be useful for higher loop computations. The effective action  $\Gamma$  will remain symmetric under the exchange of  $\omega_1$  and  $\omega_2$  and equations like (B.77)-(B.80) shall remain valid. This should facilitate at least a bit the two-loop evaluation of the effective action and gap equations. Also, one does not need to evaluate any new anomalous dimension, since these are already known, either from previous calculations [3, 12, 25], or from exploiting relations like eq.(B.41).

<sup>10</sup>The  $R$ -dependence is hidden in  $\bar{U}_1$  and  $\bar{U}_2$

<sup>11</sup>We may assume  $\cos \phi \neq 0$ , otherwise eqns.(B.100)-(B.101) would give  $\phi = 0$ , which is inconsistent with  $\cos \phi = 0$ .




Figure 6: The gap equation (B.104) with  $N = 3$  plotted in function of  $\phi$  for the values  $b_0 = 0.25, 0, -0.25, -0.3, -0.33564\dots, -0.41594\dots, -0.5$  (from bottom to top).




Figure 7: The angle  $\phi$  and scale  $R$  as a function of  $b_0$ , in units  $\Lambda_{\overline{\text{MS}}} = 1$ .

## References.

- [1] F. V. Gubarev and V. I. Zakharov, Phys. Lett. B **501** (2001) 28.
- [2] F. V. Gubarev, L. Stodolsky and V. I. Zakharov, Phys. Rev. Lett. **86** (2001) 2220.
- [3] H. Verschelde, K. Knecht, K. Van Acoleyen and M. Vanderkelen, Phys. Lett. B **516** (2001) 307.
- [4] K. I. Kondo, T. Murakami, T. Shinohara and T. Imai, Phys. Rev. D **65** (2002) 085034.
- [5] K. I. Kondo, Phys. Lett. B **514** (2001) 335.
- [6] P. Boucaud, A. Le Yaouanc, J. P. Leroy, J. Micheli, O. Pene and J. Rodriguez-Quintero, Phys. Rev. D **63** (2001) 114003.
- [7] P. Boucaud, A. Le Yaouanc, J. P. Leroy, J. Micheli, O. Pene and J. Rodriguez-Quintero, Phys. Lett. B **493** (2000) 315.
- [8] E. Ruiz Arriola, P. O. Bowman and W. Broniowski, Phys. Rev. D **70** (2004) 097505.



- [9] P. Boucaud, F. de Soto, J. P. Leroy, A. Le Yaouanc, J. Micheli, H. Moutarde, O. Pene and J. Rodriguez-Quintero, hep-lat/0504017.
- [10] D. Dudal, H. Verschelde, R. E. Browne and J. A. Gracey, Phys. Lett. B **562** (2003) 87.
- [11] D. Dudal, H. Verschelde and S. P. Sorella, Phys. Lett. B **555** (2003) 126.
- [12] R. E. Browne and J. A. Gracey, JHEP **0311** (2003) 029.
- [13] R. E. Browne and J. A. Gracey, Phys. Lett. B **597** (2004) 368.
- [14] J. A. Gracey, Eur. Phys. J. C **39** (2005) 61.
- [15] X. Li and C. M. Shakin, Phys. Rev. D **70** (2004) 114011.
- [16] X. Li and C. M. Shakin, Phys. Rev. D **71** (2005) 074007.
- [17] F. V. Gubarev and S. M. Morozov, Phys. Rev. D, to appear, hep-lat/0503023.
- [18] M. J. Lavelle and M. Schaden, Phys. Lett. B **208** (1988) 297.
- [19] J. Greensite and M. B. Halpern, Nucl. Phys. B **271** (1986) 379.
- [20] K. Knecht and H. Verschelde, Phys. Rev. D **64** (2001) 085006.
- [21] O. Piguet and S. P. Sorella, Lect. Notes Phys. **M28** (1995) 1.
- [22] D. Dudal, H. Verschelde, V. E. R. Lemes, M. S. Sarandy, S. P. Sorella and M. Picariello, Annals Phys. **308** (2003) 62.
- [23] D. Dudal, H. Verschelde, V. E. R. Lemes, M. S. Sarandy, R. F. Sobreiro, S. P. Sorella, M. Picariello and J. A. Gracey, Phys. Lett. B **569** (2003) 57.
- [24] D. Dudal, H. Verschelde, V. E. R. Lemes, M. S. Sarandy, R. F. Sobreiro, S. P. Sorella and J. A. Gracey, Phys. Lett. B **574** (2003) 325.
- [25] D. Dudal, H. Verschelde, J. A. Gracey, V. E. R. Lemes, M. S. Sarandy, R. F. Sobreiro and S. P. Sorella, JHEP **0401** (2004) 044.
- [26] D. Dudal, J. A. Gracey, V. E. R. Lemes, M. S. Sarandy, R. F. Sobreiro, S. P. Sorella and H. Verschelde, Phys. Rev. D **70** (2004) 114038.
- [27] G. Parisi and R. Petronzio, Phys. Lett. B **94** (1980) 51.
- [28] J. H. Field, Phys. Rev. D **66** (2002) 013013.
- [29] F. Giacosa, T. Gutsche and A. Faessler, Phys. Rev. C **71** (2005) 025202.
- [30] A. Cucchieri, Phys. Rev. D **60** (1999) 034508.
- [31] D. R. Bonnet, P. O. Bowman, D. B. Leinweber, A. G. Williams and J. M. Zanotti, Phys. Rev. D **64** (2001) 034501.
- [32] K. Langfeld, H. Reinhardt and J. Gattnar, Nucl. Phys. B **621** (2002) 131.
- [33] A. Cucchieri, T. Mendes and A. R. Taurines, Phys. Rev. D **67** (2003) 091502.
- [34] J. C. R. Bloch, A. Cucchieri, K. Langfeld and T. Mendes, Nucl. Phys. B **687** (2004) 76.

- [35] S. Furui and H. Nakajima, Phys. Rev. D **69** (2004) 074505.
- [36] S. Furui and H. Nakajima, Phys. Rev. D **70** (2004) 094504.
- [37] H. Suman and K. Schilling, Phys. Lett. B **373** (1996) 314.
- [38] T. D. Bakeev, E. M. Ilgenfritz, V. K. Mitrjushkin and M. Mueller-Preussker, Phys. Rev. D **69** (2004) 074507.
- [39] L. von Smekal, R. Alkofer and A. Hauck, Phys. Rev. Lett. **79** (1997) 3591.
- [40] L. von Smekal, A. Hauck and R. Alkofer, Annals Phys. **267** (1998) 1 [Erratum-ibid. **269** (1998) 182].
- [41] D. Atkinson and J. C. R. Bloch, Phys. Rev. D **58** (1998) 094036.
- [42] D. Atkinson and J. C. R. Bloch, Mod. Phys. Lett. A **13** (1998) 1055.
- [43] R. Alkofer and L. von Smekal, Phys. Rept. **353** (2001) 281.
- [44] P. Watson and R. Alkofer, Phys. Rev. Lett. **86** (2001) 5239.
- [45] D. Zwanziger, Phys. Rev. D **65** (2002) 094039.
- [46] C. Lerche and L. von Smekal, Phys. Rev. D **65** (2002) 125006.
- [47] J. M. Pawłowski, D. F. Litim, S. Nedelko and L. von Smekal, Phys. Rev. Lett. **93** (2004) 152002.
- [48] V. N. Gribov, Nucl. Phys. B **139** (1978) 1.
- [49] R. F. Sobreiro, S. P. Sorella, D. Dudal and H. Verschelde, Phys. Lett. B **590** (2004) 265.
- [50] R. F. Sobreiro, S. P. Sorella, D. Dudal and H. Verschelde, AIP Conf. Proc. **739** (2005) 455.
- [51] D. Zwanziger, Nucl. Phys. B **323** (1989) 513.
- [52] D. Zwanziger, Nucl. Phys. B **399** (1993) 477.
- [53] N. Maggiore and M. Schaden, Phys. Rev. D **50** (1994) 6616.
- [54] Semenov-Tyan-Shanskii and V.A. Franke, Zapiski Nauchnykh Seminarov Leningradskogo Otdeleniya Matematicheskogo Instituta im. V.A. Steklov AN SSSR}, Vol. **120** (1982) 159. English translation: New York: Plenum Press 1986.
- [55] G. Dell'Antonio and D. Zwanziger, Commun. Math. Phys. **138** (1991) 291.
- [56] G. Dell'Antonio and D. Zwanziger, *Proceedings of the NATO Advanced Research Workshop on Probabilistic Methods in Quantum Field Theory and Quantum Gravity, Cargèse, August 21-27, 1989, Damgaard and Hueffel (eds.), p.107, New York: Plenum Press.*
- [57] P. van Baal, Nucl. Phys. B **369** (1992) 259.
- [58] D. Zwanziger, Phys. Rev. D **69** (2004) 016002.
- [59] S. Furui and H. Nakajima, hep-lat/0503029.
- [60] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **147** (1979) 385.

- [61] A. Di Giacomo, M. D'Elia, H. Panagopoulos and E. Meggiolaro, hep-lat/9808056.
- [62] T. van Ritbergen, J. A. M. Vermaseren and S. A. Larin, Phys. Lett. B **400** (1997) 379.
- [63] M. Czakon, Nucl. Phys. B **710** (2005) 485.
- [64] K. G. Chetyrkin, Nucl. Phys. B **710** (2005) 499.
- [65] J. A. Gracey, Phys. Lett. B **552** (2003) 101.
- [66] M. Stingl, Phys. Rev. D **34** (1986) 3863 [Erratum-ibid. D **36** (1987) 651].
- [67] R. Alkofer, W. Detmold, C. S. Fischer and P. Maris, Phys. Rev. D **70** (2004) 014014.
- [68] A. Cucchieri, T. Mendes and A. R. Taurines, Phys. Rev. D **71** (2005) 051902.
- [69] W. Celmaster and R. J. Gonsalves, Phys. Rev. D **20** (1979) 1420.
- [70] K. Van Acoleyen and H. Verschelde, Phys. Rev. D **65** (2002) 085006.
- [71] D. Dudal and H. Verschelde, Phys. Rev. D **67** (2003) 025011.
- [72] P. M. Stevenson, Phys. Rev. D **23** (1981) 2916.