



RESEARCH ARTICLE

10.1002/2014PA002752

Key Points:

- The Maritime Continent area has increased 60% since 5 Ma
- More islands of the Maritime Continent strengthened the Walker Circulation
- More exposed basalt drew atmospheric $p\text{CO}_2$ down

Correspondence to:

P. Molnar,
molnar@colorado.edu

Citation:

Molnar, P., and T. W. Cronin (2015), Growth of the Maritime Continent and its possible contribution to recurring Ice Ages, *Paleoceanography*, 30, 196–225, doi:10.1002/2014PA002752.

Received 13 NOV 2014

Accepted 3 FEB 2015

Accepted article online 6 FEB 2015

Published online 12 MAR 2015

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Growth of the Maritime Continent and its possible contribution to recurring Ice Ages

Peter Molnar¹ and Timothy W. Cronin²

¹Department of Geological Sciences, Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, Colorado, USA, ²Department of Earth and Planetary Sciences, Harvard University, Cambridge, Massachusetts, USA

Abstract The areal extent of the Maritime Continent (the islands of Indonesia and surrounding region) has grown larger by ~60% since 5 Ma. We argue that this growth might have altered global climate in two ways that would have contributed to making recurring Ice Ages possible. First, because rainfall over the islands of the Maritime Continent not only is heavier than that over the adjacent ocean but also correlates with the strength of the Walker Circulation, the growth of the Maritime Continent since 5 Ma may have contributed to the cooling of the eastern tropical Pacific since that time. Scaling relationships between the strength of the Walker Circulation and rainfall over the islands of the Maritime Continent and between sea surface temperature (SST) of the eastern tropical Pacific and the strength of easterly wind stress suggest that the increase in areal extent of islands would lead to a drop in that SST of 0.75°C. Although only a fraction of the 3–4°C decrease in SSTs between the eastern and western tropical Pacific, the growth of the Maritime Continent may have strengthened the Walker Circulation, increased the east-west temperature gradient across the Pacific and thereby enabled ice sheets to wax and wane over Canada since 3 Ma. Second, because the weathering of basaltic rock under warm, moist conditions extracts CO_2 from the atmosphere more rapidly than weathering of other rock or of basalt under cooler or drier conditions, the increase in weathering due to increasing area of basalt in the Maritime Continent may have drawn down enough CO_2 from the atmosphere to affect global temperatures. Simple calculations suggest that increased weathering of basalt might have lowered global temperatures by 0.25°C, possibly important for the overall cooling.

1. Introduction

During the past 5 Ma, the Earth has cooled gradually, with that cooling punctuated by recurring Ice Ages manifested as huge ice sheets on Canada and Fennoscandia. Stable isotopes in benthic foraminifera imply global cooling, with measurements of $\delta^{18}\text{O}$ indicating cooling of the abyssal ocean and increases in continental ice volume (Figure 1) [e.g., Lisiecki and Raymo, 2005; Mudelsee and Raymo, 2005; Zachos et al., 2001], and more sparsely sampled Mg/Ca ratios also indicating cooling of the abyssal ocean [Lear et al., 2000]. Alkenones, TEX_{86} , and Mg/Ca ratios have been used to infer changes in ocean temperatures above the thermocline; widespread and monotonic cooling has been found in the eastern tropical Pacific [Dekens et al., 2008; Groeneveld et al., 2006; Lawrence et al., 2006; Wara et al., 2005; Y. G. Zhang et al., 2014], off the coast of Peru [Dekens et al., 2007], off Southern California [Dekens et al., 2007], in the Caribbean [O'Brien et al., 2014], in the North Atlantic [Lawrence et al., 2009], off the west coasts of northern Africa [Herbert and Schuffert, 1998] and southern Africa [Etourneau et al., 2009; Marlow et al., 2000; Rosell-Melé et al., 2014] (though for this last case, Leduc et al. [2014] suggest that cooling applies only to the warm season), in the southeastern Indian Ocean [Karas et al., 2011], and in the South China Sea [O'Brien et al., 2014]. In the western equatorial Pacific and eastern Indian Ocean, long-term cooling is more ambiguous. Using Mg/Ca ratios, Wara et al. [2005] and Karas et al. [2009, 2011] inferred little change in temperatures in the mixed layer in the past 5 Ma, whereas Karas et al. [2009, 2011] reported cooling in subsurface water at 10°S, which they inferred to be Indonesian Throughflow from the Pacific. Collectively, these observations indicate gradual cooling of much of the Earth.

Although the initiation of recurring Ice Ages may require some kind of instability and the crossing of some threshold in the climate system, the gradual cooling implies that a protracted geologic process must underlie the transition from hundreds of million years without continental ice sheets to recurring Ice Ages. Beginning from G. Philander's (personal communication, 2000) aphorism, "Ice is incidental to the Ice Ages," we ignore the recurrence of Ice Ages and explore a geologic process that might underlie that gradual cooling




Figure 1. Values of $\delta^{18}\text{O}$ from benthic foraminifera, from Zachos *et al.* [2001], with red line showing data smoothed using a Gaussian filter with an averaging length of 200 kyr.

(Figure 1): the growth of the Maritime Continent, the islands of Indonesia and surrounding regions (Figure 2). We discuss two parallel mechanisms by which growth of the Maritime Continent could have led to gradual global cooling and an increasingly favorable climate for inception of an ice sheet in Northern Canada: strengthening of the Walker Circulation related to the increasing fraction of islands and drawdown of global CO_2 concentrations by weathering of emergent terrain rich in basalt. We recognize that many consider the closing of the Central American Seaway to have played a crucial role in the development of recurring Ice Ages [e.g., Bartoli *et al.*, 2005; Berggren and Hollister, 1974; Haug and Tiedemann, 1998; Kaneps, 1979; Keigwin, 1982;

Sarnthein *et al.*, 2009; Weyl, 1968]. We discuss this possibility no further here, however, and merely suggest that the mechanisms that we discuss could have aided whatever role the closing of the Central American Seaway might have played.

1.1. Maritime Continent, Walker Circulation, and Laurentide Ice Sheets

Among possible links between gradual changes with distinct regional structure and the possibility of Ice Ages, that between the eastern tropical Pacific and Canada is particularly clear. When the eastern Pacific warms during El Niño events, teleconnections to the southern half of Canada lead to warmer surface temperatures [e.g., Halpert and Ropelewski, 1992; Horel and Wallace, 1981; Kiladis and Diaz, 1986, 1989; Rasmusson and Wallace, 1983; Ropelewski and Halpert, 1986; Trenberth *et al.*, 2002]. Although such teleconnections are strongest in winter, warmer winters make for longer summers throughout Canada, as measured with positive degree days, which are conducive to melting the previous winter's snowfall [e.g., Huybers and Molnar, 2007]. Moreover, general circulation model (GCM) simulations with a warm eastern tropical Pacific consistently show a warmer Canada with more positive degree days than for normal, average, present-day conditions [e.g., Barreiro *et al.*, 2006; Brierley and Fedorov, 2010; Shukla *et al.*, 2009, 2011; Vizcaíno *et al.*, 2010]. Thus, the cooling of the eastern tropical Pacific over the past 4 Ma may have enabled the accumulation of perennial snow and the inception of Canadian Ice Sheets [e.g., Barreiro *et al.*, 2006; Fedorov *et al.*, 2006; Huybers and Molnar, 2007; Molnar and Cane, 2002, 2007; Philander and Fedorov, 2003; Ravelo *et al.*, 2004, 2006].

The inferred climate of 3–5 Ma, wherein the eastern equatorial Pacific was warmer than today by ~3–4°C, has been described as a “permanent El Niño” state [e.g., Barreiro *et al.*, 2006; Brierley and Fedorov, 2010; Brierley *et al.*, 2009; Fedorov *et al.*, 2006, 2010, 2013; Goldner *et al.*, 2011; Lawrence *et al.*, 2006; Molnar and Cane, 2002, 2007; Ravelo and Wara, 2004; Shukla *et al.*, 2009, 2011; Vizcaíno *et al.*, 2010; Wara *et al.*, 2005]. Recently, Y. G. Zhang *et al.* [2014] reported SSTs, inferred using the TEX_{86} temperature proxy, that suggest a 3° in the difference between SSTs the eastern and western equatorial Pacific until ~5–7 Ma, after which the difference gradually increased to 5–6°C, which might be taken to mean that the present-day La Niña like SST distribution has always existed, but become stronger since 4–5 Ma. The analogy with El Niño derives also from the suggestion that the thermocline in the western Pacific was shallower at 5 Ma than today [Chaisson, 1995; Chaisson and Leckie, 1993], as well as the suggestion, from $\delta^{18}\text{O}$, that the temperature gradient across the thermocline in the eastern equatorial Pacific increased at approximately the same time [Cannariato and Ravelo, 1997; Chaisson and Ravelo, 2000; Ford *et al.*, 2012] and qualitative similarities of differences between El Niño air temperature and precipitation teleconnections with differences between regional climate at 3–6 Ma from those today [Molnar and Cane, 2002, 2007]. Although the term “permanent El Niño” is imprecise,




Figure 2. Maps of the Maritime Continent showing (top) present-day land and that for 5 Ma, where (bottom) submerged terrain is shown in red and islands have been moved to their positions at 5 Ma using maps of Hall [2002].

failing to distinguish between changes in the basic state of the tropical Pacific or changes in the variability (e.g., frequency or intensity) of El Niño events themselves, we continue with the view that regardless of the character of the SST anomalies, the teleconnections associated with any El Niño-like SST anomaly pattern provide a framework with which to consider pre-Ice Age climates.

If cooling of the eastern tropical Pacific underlies the transition to recurring Ice Ages, the question then becomes: What caused sea surface temperatures (SSTs) specifically in that region to decrease over the past ~5 Ma? One possibility is that an overall thermocline shoaling, although global in extent, would lead to regionally concentrated cooling in ocean upwelling regions; *Philander and Fedorov* [2003] suggested that as the thermocline shoaled, easterly winds across the tropical Pacific drew that cold water to the surface and created a background La Niña state. Then, the Bjerknes feedback enhanced zonal winds along the equator and, with an increased SST zonal gradient across the Pacific, strengthened that gradient. In regions of marked upwelling of deep water, apparently warm, nutrient-rich water upwelled in early Pliocene time, consistent with a more recent shoaling of the thermocline [*Dekens et al.*, 2007]. We cannot reject the idea that shoaling of the thermocline contributed to the present-day zonal asymmetry of the Pacific SST, but we pursue another possible contributor to an El Niño-like state.

Other suggested changes that might affect eastern Pacific SSTs include possible roles of the Maritime Continent. *Cane and Molnar* [2001] suggested that the steady, gradual northward movement of New Guinea and Australia so as to restrict flow through the Indonesian Seaway might have helped create the warm pool and, aided by Bjerknes feedback, the cold tongue in the eastern Pacific. Ocean GCM simulations with different amounts of closing of that seaway, but with present-day winds, suggest that the closing of the seaway could have led to a difference in SSTs between the eastern and western Pacific of 0.5°C [*Rodgers et al.*, 2000]. Allowing for Bjerknes feedback and an adjustment to the winds, that contribution might be greater, but more

recent coupled GCM runs have challenged this interpretation [Jochum *et al.*, 2009]. Thus, although a restricting of the Indonesian seaway seems likely to have enhanced the zonal SST differences between the eastern and western Pacific, the likely enhancement seems to be too small by itself to account for the apparent drop of $\sim 3\text{--}4^\circ\text{C}$.

Also with the Maritime Continent in mind, Dayem *et al.* [2007] noted that rainfall in that region tends to occur preferentially over islands, a result corroborated by closer investigation [e.g., As-syakur *et al.*, 2013; Sobel *et al.*, 2011]. They showed that variability in the strength of the Walker Circulation correlated with variability of rainfall over the Maritime Continent, but not with that over the warm pool. They suggested that the growth of land area in the Maritime Continent would have enhanced rainfall over the region, which in turn strengthened the Walker Circulation, and again enhanced by Bjerknes feedback, played a role in the cooling of eastern equatorial SSTs in the Pacific over the past 5 Ma. We develop further this last idea.

We argue that the fraction of land area comprising the Maritime Continent has increased by $\sim 60\%$ in the past 5 Myr. Then by exploiting the relationship of rainfall over the Maritime Continent and the strength of the Walker Circulation determined by Dayem *et al.* [2007], we estimate a strengthening of easterly wind stress across the Pacific associated with the expected increase in island area and greater rainfall there. Finally, using numerical model calculations that relate easterly wind strength to eastern equatorial Pacific SSTs, we estimate how much the eastern Pacific might have cooled as the land area of the Maritime Continent grew.

1.2. Weathering of Basalt in the Maritime Continent, CO_2 Drawdown, and Global Cooling

The growth of the Maritime Continent may have played a second role in facilitating Ice Ages. One candidate for explaining global warmth at 5 Ma is higher concentrations of greenhouse gases. Paleoatmospheric $p\text{CO}_2$ proxies extending to ~ 5 Ma include carbon stable isotope ratios in alkenones [e.g., Pagani *et al.*, 2010; Seki *et al.*, 2010] and planktic foraminifera [Raymo *et al.*, 1996], boron stable isotope ratios in carbonate sediment [Pearson and Palmer, 2000] and planktic foraminifera [Bartoli *et al.*, 2011; Seki *et al.*, 2010], B/Ca ratios in foraminifera [Tripathi *et al.*, 2009], and stomatal density indices of well-preserved remains of leaves from continental sediments [Kürschner *et al.*, 1996]. Uncertainty margins within and across this set of studies are large [e.g., Seki *et al.*, 2010] but generally support the notion that any real decline in $p\text{CO}_2$ over the past 5 Ma has been relatively small, with atmospheric $p\text{CO}_2$ not much greater than ~ 400 ppm at 5 Ma, declining slowly to ~ 280 ppm during interglacials.

Is this decline by ~ 120 ppm enough to explain the global cooling of the past 5 Ma? Studies using global climate models answer with a qualified yes. Haywood *et al.* [2013] describe results from the Pliocene Model Intercomparison Project, where a set of coupled atmosphere-ocean models were run with boundary conditions designed to mimic the mid-Pliocene. These boundary conditions include higher $p\text{CO}_2$ of 405 ppm, elimination of the West Antarctic Ice Sheet and much of the Greenland Ice Sheet, accompanied by a ~ 25 m rise in global sea level and a reduction of total land area, and changes in vegetation cover, including poleward shift of tree line at high latitudes, and a reduction of desert extent in the subtropics. The global mean temperature difference between Pliocene and Modern, averaged across eight climate models, was found to be 2.66°C , with nearly 10°C of warming at 80°N . Lunt *et al.* [2010] argued that the inclusion of slowly responding surface changes—such as vegetation and land ice—are in part responsible for a climate sensitivity of the Earth System to $p\text{CO}_2$ that is $\sim 50\%$ larger than has been found for simulations of future climate change. These slow changes, however, must be represented as a prescribed boundary forcing, for they are not feedbacks in the models. This is a relevant distinction because it means that changes in land cover, which are important for the large global climate sensitivity in the results summarized by Haywood *et al.* [2013], cannot be unambiguously viewed as feedbacks in a climate forced primarily by changes in $p\text{CO}_2$. Thus, although decreasing atmospheric $p\text{CO}_2$ has likely been important in the transition to a cooler global climate, work to date cannot definitively attribute the fraction of cooling that has been caused by the drop in $p\text{CO}_2$. Furthermore, the fundamental question of how much higher atmospheric $p\text{CO}_2$ was at 5 Ma is still plagued by large uncertainties in the proxy reconstructions.

Nonetheless, the potential importance of declining $p\text{CO}_2$ for the gradual cooling of global climate raises the question of why $p\text{CO}_2$ might have declined over the past several million years. Part of the decrease in atmospheric $p\text{CO}_2$ may have occurred abruptly, for example, as a consequence of changes in atmosphere/ocean partitioning of carbon during glaciation, but the timing of abrupt changes inferred from various studies varies

from 3.2 Ma [Tripathi *et al.*, 2009] to 2.8 Ma [Seki *et al.*, 2010] to 2 Ma [Bartoli *et al.*, 2011]. Thus, although a step change in $p\text{CO}_2$ associated with glaciation at 2.7 Ma is plausibly a part of the long-term decline, in the absence of agreement on the timing of abrupt changes, it seems likely that at least part of the decline is steady and gradual. We suggest that increases in silicate weathering due to an increasing area of islands in the Maritime Continent contributed to part of this $p\text{CO}_2$ drop. Rates of chemical weathering and consequent CO_2 consumption per unit area in the Maritime Continent are among the highest in the world, owing to the combination of warmth, high rainfall, lithology, and relief. In consideration of the world's 60 largest rivers in terms of discharge, Gaillardet *et al.* [1999] found that four of the top five of them in terms of CO_2 consumption rate per unit area drain basins on the island of New Guinea (#3 is the Irrawaddy of Southeast Asia). Weathering rates per unit area in excess of 50 times the global mean have been found in basins in the Philippines [Schopka *et al.*, 2011]. Dessert *et al.* [2003] noted the importance of small basaltic regions in the total global CO_2 consumption by weathering and estimated that basalt in Southeast Asia and the Maritime Continent contributes ~9% of the total global silicate weathering consumption of CO_2 . Kent and Muttoni [2013] discussed the paleoatmospheric CO_2 concentrations over the past 120 million years, noting that the migration of South Indochina and the Maritime Continent into the "humid tropical belt" between 5°N and 5°S could have contributed to the decline of CO_2 over the past 20 million years. In section 3 we develop further the idea that changes in island area in the Maritime Continent have played a large role in global silicate weathering and the gradual decline in $p\text{CO}_2$ over the past ~5 Ma; we develop a scaling argument that relates changes in the area of the Maritime Continent to changes in $p\text{CO}_2$ and global mean temperature.

2. Growth of the Maritime Continent

Two factors have contributed to the growth of the Maritime Continent: (1) the northward movement of Australia and New Guinea, from high southern latitudes, into the equatorial region [e.g., Hall, 2002], and (2) the emergence of islands or portions of islands from below sea level. We summarize briefly here, but discuss in much more detail in Appendix A, the geologic evidence for emergence of terrain since 5 Ma (Figure 2). Toward this end, we have relied on syntheses by Hall [2001, 2009, 2012] and Wilson [2008], and where possible, we have relied on quantitative paleodepth indicators [e.g., De Smet *et al.*, 1989, 1990; Fortuin and De Smet, 1991; Fortuin *et al.*, 1988, 1990, 1994, 1997].

Most of the emergence of land above sea level has occurred because of crustal shortening and isostatically compensated crustal thickening, in some places enhanced by sedimentation due to erosion of growing high terrain, though a global sea level fall would also contribute where elevations are within tens of meters of sea level. This combination of processes accounts for the emergence of northeastern Sumatra [Barber and Crow, 2005; Barber *et al.*, 2005; De Smet and Barber, 2005] and Java [e.g., Clements *et al.*, 2009; Hall, 2002, p. 371; Hall, 2009, p. 152; Hamilton, 1979, pp. 39–44]. Hall [2009] wrote: "Most of Sumatra and Java were elevated above sea level and emerged to their present size only since 5 Ma." Similar processes have operated in northern and eastern Borneo [e.g., Chambers *et al.*, 2004; Hamilton, 1979, p. 99; Hall and Nichols, 2002; Morley *et al.*, 2003], but this island has grown only slightly since 5 Ma.

The islands along the Banda arc, from east of Java to the small Kai and Aru Islands, seem to have emerged largely in response to the subduction of the Australian continent beneath the arc. By contrast, Timor consists of rock once part of the ancient margin of Australia and scraped off the top of that margin to build an island [e.g., Audley-Charles, 1986; Bowin *et al.*, 1980; Carter *et al.*, 1976; Veevers *et al.*, 1978]. New Guinea also has grown as the Australian continent plunged beneath a subduction zone along its northern edge, but in this case subduction has ceased and a major mountain range has been built [e.g., Baldwin *et al.*, 2012; Crowhurst *et al.*, 1996; Cloos *et al.*, 2005; Dow, 1977; Dow and Sukamto, 1984; Hall, 2009, p. 156; Hall, 2012, p. 57; Hill and Hall, 2003; Smith and Davies, 1976]. Sedimentation on the flanks of the high axial range also contributed to emergence, particularly on the southern flank of the range [e.g., Pigram and Symonds, 1991; Quarles van Ufford and Cloos, 2005].

Between Borneo and New Guinea, Sulawesi has been built by rapid westward movement and Pliocene accretion of small continental fragments [e.g., Charlton, 1986; Davies, 1990; Hall, 2002, p. 416; Hall, 2009, p. 153; Hall, 2012, p. 54; Hamilton, 1979, pp. 159, 173–174, 181–183; Silver *et al.*, 1983; Smith and Silver, 1991]. Concurrently, subduction beneath Seram and Buru has elevated these islands [Charlton, 2000; Pairault *et al.*,

Table 1. Names of Islands in the Maritime Continent, Centroid Locations, Areas at Present and at 5 Ma, and Pertinent References

Name	Latitude	Longitude	Area	Area	References ^a
			Present	at 5 Ma	
			(km ²)	(km ²)	
Alor	8.30°S	124.75°E	2,120	0	H01
Ambon	3.64°S	128.19°E	806	0	P
Atauro	8.25°S	125.60°E	147	0	CV, H01
Bacan	0.57°S	127.58°E	1,900	0	MH96
Bali	8.40°S	115.20°E	5,416	1,000	H01
Banggai	1.59°S	123.53°E	261	0	D90
Bangka	1.34°S	106.00°E	11,413	11,413	BCD05
Basu	0.34°S	103.59°E	409	0	H01
Batam	1.05°N	104.05°E	399	0	H01
Batanta	0.86°S	130.67°E	456	0	Ch91
Belitung	2.92°S	107.96°E	4,478	4,478	BCD05
Bengkalis	1.45°N	102.30°E	929	0	H01
Biak	1.00°S	136.00°E	1,904	0	H01
Bintan	1.05°N	104.50°E	1,173	0	H01
Borneo	1.00°N	113.00°E	748,168	673,351	90%: Bo
Buru	3.45°S	126.56°E	8,473	4,500	Ch00, FD91, F88
Buton	5.00°S	122.96°E	4,408	2,200	FD91, F90, SS91
Dolak	7.92°S	138.50°E	11,742	0	H01
Enggano	5.38°S	102.25°E	397	0	BCD05, S97
Flores	8.70°S	121.00°E	14,154	8,500	60%: H01
Halmahera	0.50°N	128.00°E	18,040	2,700	15% BM96, H88
Java	7.50°S	110.00°E	138,794	104,000	JAVA
Kabaena	5.25°S	121.94°E	873	0	H01
Kai Besar	5.60°S	133.00°E	550	0	Like Kai Kecil
Kai Kecil	5.75°S	132.75°E	399	0	Ch91, vMD
Kangean	6.90°S	115.35°E	430	0	H01
Karakelong (Talaud Islands)	4.28°N	126.84°E	846	0	M81
Kasiruta	0.39°S	127.19°E	473	0	H01
Kobroor (Aru Islands)	6.13°S	134.55°E	1,723	0	H01
Komodo	8.55°S	119.45°E	330	0	H01
Komoren	8.26°S	138.80°E	695	0	H01
Larat (Tanimbar Islands)	7.17°S	131.81°E	216	0	Ch91, FD91
Laut	3.70°S	116.20°E	2,057	0	vLM05
Lingga	0.15°S	104.69°E	889	0	H01
Lomblen	8.40°S	123.50°E	1,270	0	H01
Lombok	8.60°S	116.36°E	4,625	0	H01
Madura	7.00°S	113.35°E	4,429	0	H01
Maikoor (Aru Islands)	6.20°S	134.27°E	398	0	H01
Mangole (Sula Islands)	1.84°S	125.85°E	1,229	0	H01
Maya	1.15°S	109.55°E	992	992	H01
Misool	1.87°S	130.17°E	2,034	0	S09
Morotai	2.34°N	128.50°E	2,266	0	H01
Moyo	8.25°S	117.55°E	330	0	H01
Muna	5.00°N	122.59°E	2,889	1,400	H01.
New Guinea	6.00°S	140.50°E	785,753	196,000	25%: NG
Nias	1.10°N	97.55°E	4,048	0	BCD05, M80, S95, S97
Niur	0.46°S	103.50°E	342	0	H01
Numfor	1.03°S	134.88°E	335	0	H01
Obi	1.57°S	127.78°E	2,542	2,000	BM96
Padang	1.15°N	102.35°E	1,109	0	H01
Pagai Utara	2.67°S	100.10°E	622	0	BCD05, S97
Pantar	8.40°S	124.10°E	720	0	H01
Peleng	1.38°S	123.25°E	2,346	1,200	50%(?): S83
Rangsang	1.00°N	102.95°E	908	0	H01
Rantau	0.90°N	102.60°E	1,598	0	H01
Rote	10.70°S	123.15°E	1,227	0	H09
Rupat	1.85°N	101.60°E	1,490	0	H01

Table 1. (continued)

Name	Latitude	Longitude	Area	Area	References ^a
			Present	at 5 Ma	
			(km ²)	(km ²)	
Salawati	1.15°S	130.92°E	1,623	0	F78, S09
Sanane (Sula Islands)	2.20°S	125.92°E	558	0	H01
Sangihe	3.55°N	125.56°E	552	0	M03
Savu	10.55°S	121.85°E	380	0	H09
Seram	3.26°S	129.50°E	17,454	0	FD91, F88, H02, P
Siberut	1.40°S	98.97°E	3,829	0	BCD05, S97
Simeulue	2.65°N	96.10°E	1,754	0	BCD05, S97
Sipura	2.20°S	99.67°E	601	0	BCD05, S97
Sulabesi (Sula Islands)	2.21°S	125.97°E	558	0	H01
Sulawesi	2.00°S	121.00°E	180,681	145,000	80%: BR, H02, SS91, Su, vLM
Sumatra	0.50°S	102.00°E	443,066	332,000	75%: BCD05, Sum
Sumba	6.65°S	120.00°E	10,711	0	F94, F97
Sumbawa	8.50°S	118.00°E	14,386	8,600	60%
Supiori	0.85°S	135.58°E	659	0	H01
Taliabu (Sula Islands)	1.83°S	124.88°E	2,913	1,700	60%: H01
Tanahbala	0.42°S	98.40°E	468	0	BCD05, S97
Tanahmasa	0.17°S	98.46°E	344	0	BCD05, S97
Timor	9.30°S	125.50°E	28,418	0	H02, Tim
Trangan (Aru Islands)	6.53°S	134.28°E	2,149	0	H01
Utara Selatan	3.03°S	100.17°E	900	0	BCD05, S97
Waigeo	0.22°S	130.84°E	3,154	0	Ch91
Wetar	7.82°S	126.13°E	3,600	0	H01
Wokam (Aru Islands)	5.79°S	134.53°E	1,604	0	H01
Workai (Aru Islands)	6.84°S	134.72°E	152	0	H01
Wuliaru (Tanimbar Islands)	7.45°S	131.06°E	151	0	Ch91, FD91
Yamdena (Tanimbar Islands)	7.57°S	131.44°E	4,350	0	Ch91, FD91
Yapen	1.85°S	136.34°E	2,278	0	H01
<i>Papua New Guinea</i>					
Bougainville	6.20°S	155.50°E	9,318	8,800	BM67
Buka	5.30°S	154.70°E	682	0	BM67
Goodenough	9.50°S	150.15°E	687	0	BDFL, H02
Karkar	4.67°S	146.00°E	400	0	H79
Kiwai	8.58°S	143.45°E	359	0	BDFL, H02
Lavongai (New Hanover)	2.30°S	150.15°E	1,227	0	EM, Ho78
Long	5.33°S	147.08°E	500	0	H79
Manus	2.05°S	146.90°E	1,940	0	EM, Fr
Moratau	9.60°S	150.60°E	1,437	0	BDFL, H02
Mussau	1.50°S	149.66°E	400	0	EM
Muyua	9.10°S	152.80°E	874	0	BDFL, H02
New Britain	5.70°S	150.90°E	35,145	23,430	BR, L06, RC
New Ireland	3.70°S	152.50°E	7,405	0	EM, Ho78, SS88
Normanby	10.00°S	151.00°E	1040	0	BDFL, H02
Sudest	11.70°S	153.60°E	866	0	BDFL, H02
Umboi	5.70°S	148.00°E	930	0	H79
<i>Philippines</i>					
Balabac	7.95°N	117.50°E	319	319	M91, RS91
Basilan	6.50°N	122.00°E	1,266	1,266	H01, R89
Bohol	9.80°N	124.20°E	3,821	0	F03
Cebu	10.30°N	123.75°E	4,468	0	F03
Jolo	5.97°N	121.17°E	869	869	H01, R89
Leyte	10.80°N	125.00°E	7,368	3,700	D09, Saj
Mindanao	7.50°N	125.00°E	97,530	49,000	50%: Min, Saj
Negros	10.00°N	123.00°E	13,075	13,075	F03
Palawan	10.00°N	118.70°E	12,189	12,189	M91, RS91
Panay	11.10°N	122.60°E	12,011	12,011	Y09
Samar	11.90°N	125.30°E	12,849	6,400	T78
Siargao	9.90°N	126.05°E	416	416	H02, Min

Table 1. (continued)

Name	Latitude	Longitude	Area	Area	References ^a
			Present (km ²)	at 5 Ma (km ²)	
Siquijor	9.21°N	123.60°E	334	334	F03
Tawi-Tawi (Sulu Archipelago)	5.20°N	120.00°E	581	0	H01, R89
Melville	11.60°S	130.80°E	5,765	0	H02
North Central Australia	11.80°S	132.90°E	7,180	0	H02
NE Australia	11.60°S	143.00°E	12,850	0	H02
South Vietnam	9.00°N	105.50°E	133,000	133,000	(Ignoring Mekong Delta growth)
Malay Peninsula	6.00°N	101.00°E	187,000	187,000	
Sum			3,102,400	1,944,800	
Region as a whole 95°E to 154°E and 12°S to 12°N: Area = 17.38 × 10 ⁶ km ²					
Percent of region as a whole			17.9%	11.2%	

^aBCD05: Barber et al. [2005]; BDFL: Baldwin et al. [2012], Davies and Warren [1988], Fitz and Mann [2013], and Little et al. [2011]; BM67: Blake and Miezitis [1967]; BM96: Baker and Malaihollo [1996]; Bo: Chambers et al. [2004], Hall et al. [2008], Hamilton [1979, p. 99], McClay et al. [2000], Moss and Chambers [1999], Moss et al. [1997], Satyana et al. [1999], and van de Weerd and Armin [1992]; BR: Bromfield and Renema [2011]; Ch91: Charlton et al. [1991a]; Ch00: Charlton [2000]; CV: Chappell and Veeh [1978]; D90: Davies [1990]; D09: Dimalanta et al. [2009]; EM: Exon and Marlow [1988]; F78: Froidevaux [1978]; F88: Fortuin et al. [1988]; F90: Fortuin et al. [1990]; F94: Fortuin et al. [1994]; F97: Fortuin et al. [1997]; F03: Faustino et al. [2003]; FD91: Fortuin and De Smet [1991]; Fr: Francis [1988]; H79: Hamilton [1979, p. 289]; H88: Hall et al. [1988a, 1988b] and Nichols and Hall [1991]; H01: Hall [2001] and Wilson [2008]; H02: Hall [2002] (In many areas, his map, Figure 24, shows this to be deep marine.); H09: Harris et al. [2009] and Roosmawati and Harris [2009]; Ho78: Hohnen [1978]; JAVA: Burckle [1982], Clements and Hall [2007], Clements et al. [2009], P. Lunt et al. [2009], Saint-Marc and Suminta [1979], Umbgrove [1946], and Van Gorsel and Troelstra [1981]; L06: Lindley [2006]; M80: Moore et al. [1980]; M81: Moore et al. [1981]; M91: Müller [1991]; M03: Macpherson et al. [2003]; MH96: Malaihollo and Hall [1996]; Min: Pubellier et al. [1991], Queaño [2005], and Quebral et al. [1996]; NG: Abbott [1995], Abbott et al. [1994], Bailly et al. [2009], Davies et al. [1996], Dow [1977], Dow and Sukamto [1984], Pieters et al. [1983], Pigram and Symonds [1991], and Pubellier and Ego [2002]; P: Pairault et al. [2003a, 2003b]; R89: Rangin [1989]; RC: Riker-Coleman et al. [2006]; RS91: Rangin and Silver [1991]; Saj: Sajona et al. [1994, 2000]; S83: Silver et al. [1983]; S95: Samuel et al. [1995]; S97: Samuel et al. [1997]; S09: Sapin et al. [2009]; SS88: Stewart and Sandy [1988]; SS91: Smith and Silver [1991]; Su: BouDagher-Fadel [2002], Calvert [2000], Grainge and Davies [1985], Mayall and Cox [1988], Sudarmono [2000], van Leeuwen and Muhardjo [2005], and van Leeuwen et al. [2010]; Sum: Barber and Crow [2005] and De Smet and Barber [2005]; Tim: Audley-Charles [1986], Bowin et al. [1980], Carter et al. [1976], De Smet et al. [1990], Haig, 2012, Keep and Haig [2010], Johnston and Bowin [1981], Nguyen et al. [2013], Quigley et al. [2012], van Marle [1991], and Veevers et al. [1978]; T78: Travaglio et al. [1978]; vLM: van Leeuwen and Muhardjo [2005]; vMD: van Marle and De Smet [1990]; and Y09: Yumul et al. [2009].

2003a, 2003b]. East-west crustal shortening and thickening have elevated eastern Halmahera, while volcanism has built the western part [e.g., Baker and Malaihollo, 1996; Hall et al., 1988b; Nichols and Hall, 1991].

Crustal shortening and thickening within Mindanao [Pubellier et al., 1991, 1996; Sajona et al., 1994, 2000] and across smaller islands of the Philippines have elevated these regions [e.g., Faustino et al., 2003; Travaglio et al., 1978]. Palawan, however, seems to have emerged by middle Miocene time (between ~15 and 10 Ma) [e.g., Holloway, 1982; Rangin and Silver, 1991].

As summarized in Table 1, the region between 95°E and 154°E (from just west of Sumatra to just west of Bougainville) and between 12°N and 12°S, comprises 17.38 × 10⁶ km². Today, 17.9% of this area or 3.10 × 10⁶ km² stands above sea level, but at 5 Ma, only 11.2% or 1.94 × 10⁶ km² had emerged (Figure 2). Thus, the emergent fraction has increased by 60% or 1.16 × 10⁶ km²; this new land is roughly equal to the combined area of France and Spain, or the combined area of the four-corners states in the USA (Utah, Colorado, Arizona, and New Mexico).

In most cases the emergence of land above sea level has also added orography to the region. Although we have made no attempt to quantify changes in mean elevations, as discussed in Appendix A, mountain ranges seem to have grown in many regions, like New Guinea, Halmahera, Seram, Sulawesi, Borneo, Java, and Sumatra, in the past 5 Ma.

3. Silicate Weathering and Maritime Continent Area

Atmospheric pCO₂ evolution on long time scales is determined by a competition between volcanic outgassing and the combined effects of silicate weathering and burial of organic sediment:

$$\frac{dC}{dt} = \mathcal{V} - \mathcal{W} - \mathcal{B}, \quad (1)$$

where C is the reservoir of carbon in the atmosphere, oceans, and biosphere (units: Pg), \mathcal{V} is the volcanic outgassing rate, \mathcal{W} is the sink from silicate weathering, and \mathcal{B} the burial rate of organic sediment (all Pg/yr). If

the volcanic input is fixed, then equilibration of atmospheric $p\text{CO}_2$ requires that the sinks adjust to make the right-hand side of (1) zero. This adjustment is dominated by the increase of silicate weathering under warmer and wetter conditions—a byproduct of increased $p\text{CO}_2$ —and the interaction between the long-term carbon and climate systems acts to stabilize both [e.g., *Kump et al.*, 2000; *Walker et al.*, 1981]. In models of atmospheric $p\text{CO}_2$ evolution over tens or hundreds of millions of years, global silicate weathering \mathcal{W} is typically represented as a product of several functions that relate rates of weathering to factors including atmospheric $p\text{CO}_2$, global mean temperature and runoff, and total land area [e.g., *Berner and Kothavala*, 2001].

The simplest form that will suffice for us relates the relative change in weathering flux $\delta\mathcal{W}/\mathcal{W}_0$ to relative changes in weathering-rate-weighted land area ($\delta A^*/A_0^*$) and to global temperature (δT):

$$\frac{\delta\mathcal{W}}{\mathcal{W}_0} = \frac{\delta A^*}{A_0^*} + \alpha \delta T. \quad (2)$$

In (2), we have assumed that the influence of both runoff and temperature can be subsumed into a single coefficient, α , that scales relative changes in weathering to changes in temperature; values given in *Berner and Kothavala* [2001] imply $\alpha = 0.12 \text{ K}^{-1}$. We consider fractional changes in weathering-rate-weighted land area, because addition or removal of land with very high or low weathering rates can affect the global value of \mathcal{W} more or less than what one would expect from simple fractional changes in land area.

We now assume that the volcanic source of carbon has been relatively constant over the past 5 Ma and that the quasi-steady limit of (1) can be taken: $\mathcal{V} - \mathcal{W} - \mathcal{B} = 0$. This then implies $\delta\mathcal{W}/\mathcal{W} \approx 0$. It follows that we can estimate quantitatively, if only roughly, how increases in the area in the Maritime Continent—where weathering is especially rapid—have been compensated by declining global temperature since 5 Ma. We justify constant volcanism because it is thought that seafloor-spreading rates largely determine \mathcal{V} . It is difficult to reject the null hypothesis that seafloor-spreading rates have not varied significantly in time even over time scales as long as the past 180 Ma, based on seafloor area-age relationships [e.g., *Parsons*, 1982; *Rowley*, 2002]. Moreover, changes in rates since ~6 Ma have been negligible [*Krijgsman et al.*, 1999]. We justify the quasi-steady limit of (1) because the time scale of changes we consider (~5 Ma) is much larger than the relaxation time scale of C in (1), which is on the order of ~0.5 Ma for the current climate. Applying $\delta\mathcal{W}/\mathcal{W} \approx 0$ in (2) gives the scaling relationship

$$\delta T = -\frac{1}{\alpha} \frac{\delta A^*}{A_0^*}. \quad (3)$$

Dessert et al. [2003] estimated that weathering of basalt in SE Asia and Indonesia consumes $1.03 \times 10^{12} \text{ mol C/yr}$; this is ~9% of the total silicate weathering flux of $11.7 \times 10^{12} \text{ mol C/yr}$ [*Gaillardet et al.*, 1999]. Our use of weathering-rate-weighted area would thus give $\delta A^*/A_0^* = -9\%$, if all of this basaltic region were instantaneously submerged. The question is how much of this basalt has emerged since 5 Ma?

We make a crude estimate, starting with Figure 1 of *Amiotte Suchet et al.* [2003], which clearly shows that the vast majority (about ~5/6) of the basaltic area in the “SE Asia and Indonesia” province is on islands—including a large contribution from the Philippines and particularly Mindanao—and not on the mainland of Asia. In the absence of more geographically explicit information, we estimate that changes in the area of island basalts have scaled similarly to changes in the total island area, i.e., that it increased by 60% from 5 Ma to the present or that it was 40% smaller than the present at 5 Ma. Along with previous assumptions, this gives $\delta A^*/A_0^* = 0.09 \times 5/6 \times 0.4 = 0.03$ in (3). Then, using $\alpha = 0.12 \text{ K}^{-1}$ gives a decrease in global temperature of 0.25°C since 5 Ma, due to the drawdown of $p\text{CO}_2$ by weathering from newly exposed basalt in the Maritime Continent. Although small compared to the apparent temperature change between the Pliocene and the present, this cooling would likely be amplified by a factor of ~2 at the poles, or perhaps more if feedbacks in high-latitude land cover are important.

We may also use this 0.25°C estimate of temperature change due to decreased $p\text{CO}_2$ in the atmosphere to infer how much the concentration of CO_2 has changed. Using the top-of-atmosphere energy balance between the radiative forcing of additional CO_2 and the increased longwave emission from a warmer planet, we may write

$$\gamma \ln \left(1 + \frac{\delta C}{C_0} \right) = \lambda \delta T, \quad (4)$$




Figure 3. Daytime rainfall over the Maritime Continent, based on data from the Tropical Rainfall Measurement Mission (TRMM) [from Sobel *et al.*, 2011]. Note the concentration of rainfall over islands.

where $\gamma \approx 5.35 \text{ W m}^{-2}$ is the radiative forcing in associated with an e -fold change in CO_2 concentration [Ramaswamy *et al.*, 2001], δC is the change in concentration of CO_2 , C_0 is the reference-state concentration of CO_2 (both in ppm), λ is the total climate feedback parameter ($\text{W m}^{-2} \text{K}^{-1}$), and $\delta T \approx -0.25^\circ\text{C}$ is the change in temperature. Our estimate of the decline in $p\text{CO}_2$ thus depends on the choice of climate feedback parameter; values of $\lambda \sim 0.75 \text{ W m}^{-2} \text{K}^{-1}$ are implied by “Earth System Sensitivity” estimates of 5°C in Haywood *et al.* [2013], but larger values of $\lambda \sim 1.2 \text{ W m}^{-2} \text{K}^{-1}$ are given by Soden and Held [2006] for a set of simulations of future climate change due to elevated CO_2 . Taking $\lambda \sim 1 \text{ W m}^{-2} \text{K}^{-1}$ and $C_0 \approx 400$ ppm, the estimated decrease in the concentration of CO_2 would be 19 ppm, albeit a small, but perhaps not negligible, fraction of the plausible 120 ppm decrease from 400 to 280 ppm.

4. Rainfall Over the Maritime Continent

As noted above, rain falls more over the islands than over the open ocean. Holland and Keenan [1980, p. 225] wrote about rainfall over the Maritime Continent on one day, 10 December 1978: “This afternoon convection provides a near perfect map of the region; every island and mountain range is delineated by one or more cumulonimbi and there is a rash of towering cumulonimbi interspersed with smaller cumuli over northern Australia.” A particularly striking example is the recurring thunderstorm known as “Hector,” which forms nearly every day during certain seasons over the Tiwi islands, just north of the Australian mainland [Carbone *et al.*, 2000; Keenan *et al.*, 1989, 2000]. For an 8 year period from December 1997 to November 2005, Dayem *et al.* [2007] calculated an average of 6.8 mm/d for rainfall over the islands of the Maritime Continent and 4.3 mm/d for the surrounding ocean, corresponding to 58% more rain over land than sea. Similarly, for 1998–2000, As-syakur *et al.* [2013] reported a mean rainfall rate of 7.62 mm/d over the Indonesian islands, but only 5.47 mm/d over the adjacent oceans or $\sim 40\%$ more rain over islands than ocean. Sobel *et al.* [2011] reported a 25% enhancement of rain over the islands with surface areas of 315 km^2 to 6150 km^2 in the Maritime Continent, relative to the nearby surrounding ocean.

The enhancement of rainfall over islands is intimately tied to the diurnal cycle of surface heating over islands [e.g., Carbone *et al.*, 2000; Cronin *et al.*, 2015; Keenan *et al.*, 2000; Qian, 2008; Qian *et al.*, 2013; Saito *et al.*, 2001; Sato *et al.*, 2009]. The imprint of the diurnal cycle is particularly apparent when rainfall during daytime hours is plotted on a map, as in Figure 3, from Sobel *et al.* [2011]. In addition, although rainfall enhancement can occur over flat islands for reasons related to the diurnal cycle alone, observations and numerical simulations also suggest that orography can further enhance rainfall over tropical islands [e.g., Liberti *et al.*, 2001; Saito *et al.*, 2001; Sobel *et al.*, 2011; Zhou and Wang, 2006].

To understand how the diurnal cycle alone might lead to island rainfall enhancement, Cronin *et al.* [2015] inserted a circular region with low heat capacity, to represent an island, into a cloud-resolving model with a




Figure 4. Annual average upper troposphere (250 mb) temperature across the equatorial Pacific from the National Centers for Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR) reanalysis [Kalnay et al., 1996]. Image provided by the NOAA-Cooperative Institute for Research in Environmental Sciences Climate Diagnostics Center, Boulder, Colorado, from their Web site at <http://www.cdc.noaa.gov/>.

slab ocean and performed several 250 day simulations of radiative-convective equilibrium across a range of island sizes. For a reference island of radius 48 km, they found that rainfall over the low heat capacity island region, 6.17 mm/d was greater than that over the surrounding ocean, 2.94 mm/d. The islands may have a second impact that is less obvious than average rainfall but perhaps more important for large-scale circulation. In their numerical experiments, Cronin et al. [2015] found that as they increased the fraction of area with low heat capacity, the mean temperature of the upper troposphere over the entire region increased. The asymmetry of hot moist air rising during the day, but cooler air simply spreading out near the surface at night led to a rectification of the troposphere’s response to periodic heating, such that with increasing island fraction, up to 20%, the upper troposphere over the entire model atmosphere warmed by up to ~1.5°C.

Because of the weak Coriolis effect in the tropics, lateral differences of 1°C within the tropical free troposphere are difficult to maintain. The mean temperature difference across the equatorial Pacific today is only ~1.5°C (Figure 4). The pressure gradient associated with thermal anomalies drives a large-scale circulation with ascent and divergence aloft in anomalously warm regions and with convergence and descent in anomalously cool regions. Thus, based on the findings of Cronin et al. [2015] for upper tropospheric warming, the strength of the Walker Circulation might be dependent on the fraction of land within the Maritime Continent. Such a suggestion must remain speculative, however, because of many limitations: consideration of only small regions in the cloud-resolving model, idealized characterization of islands without consideration of many differences between land and ocean (e.g., albedo), neglect of ocean circulation, the Coriolis effect, extratropical interactions, etc. Nevertheless, we pursue the idea that the fraction of island area in the Maritime Continent could play a role in the strength of the Walker Circulation.

5. The Strength of the Walker Circulation and SSTs in the Cold Tongue

Dayem et al. [2007] correlated an indicator of Walker Circulation strength and the rainfall over the Maritime Continent and nearby Pacific Warm Pool. To characterize Walker Circulation strength they used the difference in westerly winds between 250 hPa and 850 hPa in the Central Pacific or wind shear: $\Delta u = u_{250} - u_{850}$ (m/s); a larger Δu implies a stronger Walker Circulation. Whereas they found a negligible correlation of such wind shear with rainfall over the warm pool, for rainfall over the Maritime Continent, P_I (mm/d), the sensitivity of wind shear was large:

$$\frac{d\Delta u}{dP_I} \approx 3 \frac{\text{m/s}}{\text{mm/d}}, \tag{5}$$

over the Pacific between approximately 140°E and 140°W (Figure 5a). A climatological view of equatorial winds, between 10°S and 10°N, over the Pacific (Figure 5b), shows approximately equal magnitudes of westerly winds at 250 hPa and easterly winds at 850 hPa. The average wind shear increases from a negligible value (possibly even negative) at 140°E, to a maximum of ~16 m/s near 140°E, with an average from 140°E-140°W of $\Delta u \approx 8$ m/s, and then with a decrease farther east.




Figure 5. (a) Regression of wind shear ($\Delta u = u_{250} - u_{850}$, where u_{250} is westerly wind speed aloft at 250 hPa and u_{850} is low-level easterly wind speed at 850 hPa) against precipitation, P_i over the islands of the Maritime Continent, and (b) east-west cross section of zonal winds along the equatorial Pacific.

Because rain falls over or near the islands, a logical step is to assume that an increase in the fraction of the region occupied by islands would lead to an increase in precipitation, which in turn would strengthen the Walker Circulation. We seek a scaling for the change in the wind shear, $\delta\Delta u$, equal to $\delta\Delta u = (d\Delta u/dP_i)\delta P_i$, for a change of δP_i in rainfall over the islands. Substituting space for time, we then suppose that the fraction of area occupied by land increased by δf . It follows that we would predict an increase in wind shear of

$$\delta\Delta u = \delta f \frac{d\Delta u}{dP_i} \Delta P, \quad (6)$$

where ΔP is the difference in rainfall over islands and surrounding ocean. Using the values of Maritime Continent island fraction of 11.2% at 5 Ma as compared to 17.9% today, then $\delta f = 6.7\%$. Using (5) and $\Delta P \approx 2.5$ mm/d, (6) gives a change in the wind shear of 0.5 m/s, and therefore a reduction of $\sim 6\%$ at 5 Ma as compared to today.

The difference in time mean midtropospheric vertical velocity between island and ocean in the results of Cronin *et al.* [2015] provides another way to estimate how island area has affected changes in Δu . By mass continuity, air that ascends over the Maritime Continent must be balanced by descent elsewhere; idealized dynamics of an equatorial beta-plane suggest that most of the descent from zonally asymmetric ascent near the equator occurs on the equator to the east and resembles the Walker Circulation more than the Hadley Circulation [Gill, 1980]. Given the horizontal half width of the Maritime Continent region, $L \sim 3000$ km, and the pressure scale between the surface and maximum vertical velocity, $\Delta p \sim 500$ hPa, changes in vertical velocity over the Maritime Continent would translate into changes in westerly wind difference as

$$\delta\Delta u = 2L \frac{\delta\omega}{\Delta p}. \quad (7)$$

If we assume that changes in the vertical velocity over the Maritime Continent result from changes in island fraction, multiplied by a land-ocean contrast in vertical pressure velocity of $\Delta\omega \sim 100$ hPa/d (consistent with the findings of *Cronin et al.* [2015]), then we obtain $\delta\Delta u = 2L \delta f \Delta\omega/\Delta p$ or an increase in westerly wind difference of ~ 0.9 m/s since 5 Ma. This is somewhat larger than the estimate of 0.5 m/s above, which used the observationally based regressions of *Dayem et al.* [2007], but is still of the same order of magnitude.

We next exploit results of *Fedorov and Philander* [2001], who used *Zebiak and Cane's* [1987] simple model to examine how sea surface temperature differences depend on wind stress, and of *von der Heydt et al.* [2011], who used a variant of that model. Both showed that a $\sim 40\%$ drop in easterly wind stress over the Pacific would lead to an increase of $\sim 2.5^\circ\text{C}$ in the eastern Pacific SST.

Recall that wind stress, τ , depends quadratically on wind speed, u :

$$\tau = \rho_a C_D u^2, \quad (8)$$

where ρ_a (≈ 1.2 kg/m³) is the density of air and C_D ($\approx 1.5 \times 10^{-3}$) is a dimensionless drag coefficient. Assuming that surface winds scale with winds somewhat above the surface, then relative changes in surface wind speeds are equal to relative changes in Δu (Figure 5b). Taking Δu at 5 Ma to be $\sim 6\%$ smaller than that today, the easterly wind stress would have been smaller by $\sim 12\%$. Assuming that differences in eastern Pacific temperatures vary linearly with wind stress, as calculations by *von der Heydt et al.* [2011] suggest, this simple calculation suggests that the eastern Pacific SST would have been warmer by 0.75°C ($= 2.5^\circ\text{C} \times 12\%/40\%$) at 5 Ma.

This 0.75°C decrease in temperature in the eastern Pacific might translate to a global cooling as well, through interactions between global temperatures and atmospheric circulation. *Pierrehumbert* [1995] noted that overturning circulations in the tropics create very dry regions of the atmosphere, allowing the planet to emit more longwave radiation to space than an atmosphere with horizontally uniform humidity would. This finding would lead us to expect that long-term strengthening of the Walker Circulation could lead to global cooling by drying out the atmosphere of the tropical Pacific. On shorter time scales, the logical inverse of this proposition holds: a weakening of the Walker Circulation warms the Earth. *Chiang and Sobel* [2002] pointed out that the entire tropics warmed by $\sim 1^\circ\text{C}$ in March–May 1998, following the 1997–1998 El Niño, the largest recent El Niño, and the one whose teleconnections best match differences between pre-Ice Age and present-day climates [e.g., *Molnar and Cane*, 2007]. On a global scale, GCM simulations of both future climate [e.g., *Held and Soden*, 2006; *Vecchi and Soden*, 2007] and idealized Walker Circulation [*Merlis and Schneider*, 2011] show that large-scale atmospheric circulation weakens with global warming. Moreover, this weakening follows from a simple theoretical argument that depends on the temperature dependence of evaporation via the Clausius-Clapeyron relationship [*Held and Soden*, 2006; *Merlis and Schneider*, 2011]. From GCM runs, *Vecchi and Soden* [2007] estimated that the strength of the Walker Circulation should increase by 5%–10% per 1°C decrease in global mean temperature. Thus, a strengthening of the Walker Circulation not only should cool of the eastern tropical Pacific but also should contribute to a global cooling, which by positive feedback would contribute to further weakening of that circulation.

Obviously, a drop of 0.75°C , even if enhanced by a positive feedback between global cooling and strengthened Walker Circulation, is only a fraction of the apparent drop of 3–4 $^\circ\text{C}$ in eastern Pacific SSTs discussed above in section 1. We cannot assert that the emergence of islands in the Maritime Continent played the principal role in altering eastern Pacific SSTs. At the same time, with an additional small contribution from oceanographic consequences of the northward movement of New Guinea [e.g., *Cane and Molnar*, 2001; *Rodgers et al.*, 2000], perhaps the growth of the Maritime Continent played an essential role in that change.

6. Discussion

Because the logic exploited here is speculative at all levels, it seems worth reviewing each step. *Hall* [2001], on whose work we have relied heavily, stated, "I believe that the maps of Figures 7 to 10 are generous in assessing areas of possible land and shallow sea." Thus, perhaps we have underestimated the area below sea level at 5 Ma. Conversely, however, his Figure 10 for 5 Ma, Figure 14 of *Hall* [2009], and Figure 3.12 of *Hall* [2012] all show the region between Borneo and the Malaya Peninsula as emergent. With an area of $\sim 300,000$ km², inclusion of this emergent territory would reduce the difference between 5 Ma and the present from 6.7% to $\sim 5\%$, and therefore the estimated effect of the growth of the Maritime Continent on

Eastern Pacific SSTs to 0.56°C. We ignore this possibility, for he did not discuss it, but obviously its inclusion would call for a smaller difference between eastern Pacific SSTs today and at 5 Ma.

In the opposite direction, we have ignored the evidence for growth of high terrain since ~5 Ma, but as discussed in Appendix A, most of the high terrain seems to be young. All high belts and islands seem to be younger than 10 Ma, and most have grown since 5 Ma. *Sobel et al.* [2011] reported an orographic enhancement of rainfall over tropical islands, and calculations of rainfall over islands consistently suggest an orographic enhancement of rainfall [e.g., *Liberti et al.*, 2001; *Saito et al.*, 2001; *Sobel et al.*, 2011; *Zhou and Wang*, 2006]. Thus, we may have underestimated the impact of islands and their topography on rainfall, and therefore on the strength of the Walker Circulation.

We have treated all land as equal, but for that affecting the Walker Circulation, the islands in the eastern part of the Maritime Continent might be the more important ones. The eastern part of the Maritime Continent has emerged more than the western part since 5 Ma (Figure 2).

The correlation of wind shear with precipitation over the islands of the Maritime Continent is stronger in the western part of the equatorial Pacific than the eastern part (Figure 5a). Yet the part of the tropical Pacific that seems likely to have the greatest impact on Ice Ages is the eastern part. The wind shear and low-level easterlies also are strongest in the eastern part (Figure 5b). Thus, for the growth of islands in the Maritime Continent to have affected SSTs in the eastern Pacific, there must be an additional process that links wind shear in the western equatorial Pacific to SSTs in the east.

Last, although the simple calculations of *Fedorov and Philander* [2001] and *von der Heydt et al.* [2011] rest on a strong theoretical foundation, the precise links between wind stress and equatorial SSTs depend on many other model parameters. The quantitative link used here—a 6% decrease in easterly wind strength and therefore a 12% reduction in wind stress leading to a ~0.75°C increase in eastern Pacific SST—is qualitative in the sense that assigning an uncertainty to that temperature difference requires subjectivity.

With regard to the influence of island area the drawdown of atmospheric CO₂ since 5 Ma, our estimate that this has decreased global mean temperatures by ~0.25°C also is subject to several uncertainties. First, our estimate of the change in weathering-rate-weighted area has been crude and has not relied on geographically matching the recently emerged areas with basalt areas of *Amiotte Suchet et al.* [2003]. On the one hand, it is possible that we have overestimated the fractional change in area of basalt. On the other, however, we have not attempted to account for weathering from other rock types in newly emerged segments of the Maritime Continent, which may contribute additionally to an increase in weathering rate since 5 Ma, because weathering rates in the region are high even in granitic terrain. We have also not accounted for the growth of high terrain since 5 Ma, and the role it may have played in increasing CO₂ drawdown, due to the tight coupling between physical and chemical erosion rates [e.g., *Gaillardet et al.*, 1999; *Larsen et al.*, 2014].

The value of δT we have estimated depends on the fractional sensitivity of weathering to global mean temperature change, which may be smaller than the value of $\alpha = 0.12 \text{ K}^{-1}$ that we have assumed. The parameter α represents the sum of fractional changes in weathering rate with temperature (α_T), and fractional changes in global runoff with temperature (α_R). At the microscopic scale, $\alpha_T \sim 0.1 \text{ K}^{-1}$ is justified by measurements of activation energies of silicate dissolution reactions of $E_a \sim 60\text{--}80 \text{ kJ/mol}$ [*Kump et al.*, 2000]; $\alpha_T \approx E_a / (RT_0^2)$ (where R is the universal gas constant and T_0 is a reference temperature in Kelvin). Empirical estimates from a set of basin-scale weathering rates across a range of temperatures, however, tend to be slightly lower, and also making such estimates becomes difficult due to the covariance of runoff and temperature across basins. For the sake of confidence in these microscopic estimates of α_T , it is disconcerting that *Gaillardet et al.* [1999] noted “On a global scale, there is no general correlation of weathering rates (total or cationic) with temperature.” By focusing on the basins with the highest weathering rates at each temperature, however, they obtained an envelope suggesting $\alpha_T \sim 0.08 \text{ K}^{-1}$. *Dessert et al.* [2003] more carefully controlled for runoff and calculated $\alpha_T \sim 0.06 \text{ K}^{-1}$ for a number of basaltic regions. Estimates of $E_a \sim 50 \text{ kJ/mol}$ and thus $\alpha_T \sim 0.07 \text{ K}^{-1}$ have also been obtained from a survey river chemistry in granitic basins [*Oliva et al.*, 2003].

The fractional change in global runoff with temperature, α_R , can be estimated from climate models or historical observations. Observational studies are controversial and challenging to conduct cleanly; the significance of the finding of *Labat et al.* [2004] that global runoff has increased by 0.04 K^{-1} was disputed by

Legates et al. [2005] and *Dai et al.* [2009], both of whom found no statistically significant correlation between historical runoff and temperature changes. Climate modeling studies in which increasing greenhouse gas concentrations force temperature change find $\alpha_R \sim 0.03 \text{ K}^{-1}$; the multimodel mean estimate of α_R does not depend strongly on the specific scenario of greenhouse gas concentrations, but estimates across models range from $>0.02 \text{ K}^{-1}$ to nearly 0.05 K^{-1} [*X. Zhang et al.*, 2014]. Taken together, these results suggest that a central estimate of α should be closer to 0.1 K^{-1} ($\alpha_T \sim 0.07 \text{ K}^{-1}$ and $\alpha_R \sim 0.03 \text{ K}^{-1}$) than the value of $\alpha = 0.12 \text{ K}^{-1}$ used above, which would increase δT slightly to $\sim 0.3^\circ\text{C}$.

There is also a more fundamental structural uncertainty associated with the estimation of α , to which *Kent and Muttoni* [2013] alluded: if temperatures in the low-latitude regions where a large fraction of global weathering takes place are much less sensitive to $p\text{CO}_2$ than is the global mean temperature, then global weathering can be much less sensitive to global temperature than either microscopic or basin-scale estimates would indicate. For instance, suppose the tropical half of global surface area from 30°S – 30°N contributes 75% of the silicate weathering consumption of CO_2 and the extratropical half of global surface area poleward of 30° contributes the remaining 25%. Then, if the tropics warm at 0.5 times the global average, and the extratropics warm at 1.5 times the global average, global weathering rates will increase with global average temperature at only 75% the rate one would expect from a globally uniform warming. Because local weathering rates scale exponentially with temperature, this issue is directly analogous to the response of global mean atmospheric water vapor to global mean temperature, in that the local exponential scaling can be wrong if applied globally, due to the negative correlation between temperature and temperature change [*Back et al.*, 2013].

Along with this latitudinal covariance of weathering rates and temperature changes, covariance between regional weathering rates and runoff changes, particularly over areas with high weathering rates, could conceivably alter the sensitivity of global weathering to global temperature. For example, we have suggested that the Walker Circulation may have increased in strength as the Maritime Continent has expanded; this would imply greater runoff over the Maritime Continent and would thus lead to an increase in per-area weathering rates that could also have a global influence on $p\text{CO}_2$. Following this line of reasoning to quantitative conclusions would require too many speculative assumptions to pursue, but the importance of small basaltic areas for global weathering rates underscores the importance of constraining regional changes in both temperature and hydrology.

7. Conclusions

From a synthesis of published geological observations, we infer that the fraction of land area in the Maritime Continent has increased by $\sim 60\%$, from $1.94 \times 10^6 \text{ km}^2$ at 5 Ma to the present-day $3.10 \times 10^6 \text{ km}^2$. This increased area could have facilitated the shift from a largely ice-free Northern Hemisphere to recurring Ice Ages in two ways.

First, previous work correlated the strength of the Walker Circulation, as quantified by the difference in zonal winds at 250 hPa and 850 hPa, with precipitation over the islands of the Maritime Continent [*Dayem et al.*, 2007]. Because rain falls preferentially over islands [e.g., *As-syakur et al.*, 2013; *Dayem et al.*, 2007; *Holland and Keenan*, 1980; *Sobel et al.*, 2011], the growth of island area could have strengthened the Walker Circulation. Scaling relationships between precipitation and wind shear [*Dayem et al.*, 2007] and between wind stress and eastern Pacific SSTs [e.g., *Fedorov and Philander*, 2001; *von der Heydt et al.*, 2011], with additional assumptions, allow for the increase in islands in the Maritime Continent to have reduced the eastern Pacific SST by $\sim 0.75^\circ\text{C}$. Such a reduction in eastern Pacific SST, if only a fraction of the 3 – 4°C change in the difference between eastern and western Pacific equatorial SSTs since ~ 5 Ma, might have contributed to a cooling of Canada and contraction of summer warmth. Then shorter summers throughout Canada could have prevented winter snow from melting and allowed Ice Ages to become recurring phenomena since that time [e.g., *Barreiro et al.*, 2006; *Fedorov et al.*, 2006; *Huybers and Molnar*, 2007; *Molnar and Cane*, 2002, 2007; *Philander and Fedorov*, 2003; *Ravelo et al.*, 2004, 2006].

Second, increased area in the Maritime Continent since 5 Ma exposed basaltic rock to weathering. Because the weathering of basalt, particularly in warm, moist conditions, extracts CO_2 from the atmosphere rapidly, this increased exposure of basalt could have lowered $p\text{CO}_2$ in the atmosphere. Simple calculations suggest a

lowering of the concentration of CO₂ by ~20 ppm and a drop in global mean temperature of 0.25°C since 5 Ma. Uncertainties in the requisite parameters needed to make this estimate of temperature change render 0.25°C more likely an underestimate than overestimate. Moreover, because the poles respond more strongly than equatorial regions, a small global average should call for a greater high-latitude temperature drop.

We recognize that the many assumptions make these links between islands in the Maritime Continent and Ice Ages speculative. At the same time, we have tried to be conservative in quantifying the links, and we may have underestimated them.

Appendix A: Growth of the Maritime Continent

We discuss the evidence used to infer the area of islands or peninsulas that reach into the Maritime Continent at 5 Ma, starting in the west and, crudely, moving east. Numerical values of both present-day areas and estimates of those at 5 Ma are given in Table 1.

A1. Sumatra and Malay Peninsula

It appears that little change in land area has occurred in the Malay Peninsula or in southern Southeast Asia, except maybe a progradation of the Mekong Delta, which we ignore here. Thermochronologic studies suggest little erosion of the Malay Peninsula for tens of millions of years [e.g., Cottam *et al.*, 2013a]. Hall [2001, 2009, 2012] shows the Malay peninsular emergent at 5 Ma, and we assume no change in area of those regions.

A consensus suggests that more than a third and perhaps as much as half of the island of Sumatra was still submerged as recently as at 5 Ma. The flat terrain northeast of the Barisan Mountains, which lie just inland of the southwest coast, emerged since ~5 Ma [Hall, 2001, Figure 10; Hall, 2009, Figure 14] or 7 Ma [Wilson, 2008]. Relying on Barber and Crow's [2005] comprehensive discussion of individual basins, Barber *et al.* [2005] inferred that roughly half the island lay below sea level in Late Miocene time, but by Early Pliocene time, that fraction was only a quarter of the present-day area of the island.

The northeastern half of the island underwent largely mild tectonic activity in Cenozoic time, with a bit of crustal extension and subsidence. The Barisan Mountains apparently began to form in Late Oligocene time, in response to crustal shortening, but they seem to have been a minor feature, at least a minor source of sediment, until mid-Miocene time [Barber *et al.*, 2005; De Smet and Barber, 2005]. The regression of the sea from the northeastern half of the island occurred in large part because of abundant fluvial sediment from the mountains. In describing the late Miocene to early Pliocene transition from shallow marine to sublittoral and then to deltaic sediment, De Smet and Barber [2005] stated: "The climax of uplift and erosion of the Barisans occurred in the Late Pliocene..." At that time reverse slip became common, and mild NE-SW shortening of the crust under northeastern Sumatra began. We presume that the small islands northeast of Sumatra—Basu, Batam, Bengkalis, Bintan, Lingga, Niur, Padang, Rangsang, Rantau, and Rupert—emerged at the same time.

This emergence of the northeastern side of the island since ~5 Ma does not seem to apply to the islands of Bangka or Belitung, which lie just east of the southeastern end of Sumatra. Barber *et al.* [2005] show them to be either eroding or collecting fluvial sediment in late Miocene and early Pliocene time.

It appears that the islands southwest of Sumatra, in its forearc, emerged in Pliocene time [Barber *et al.*, 2005; Moore *et al.*, 1980; Samuel *et al.*, 1995, 1997]. These include Enggano, Nias, Pagai Utara, Siberut, Simeulue, Sipura, Tanahbala, Tanahmasa, and Utara Selatan. Although Moore and Karig [1980; Moore *et al.*, 1980] and Samuel *et al.* [1995, 1997] disagree on the structural history of Nias, and on details of when Paleogene sediment on the island rose from deep water, they agree that the island emerged in Pliocene time. Although data are sparse from other islands, a concurrent emergence seems likely [Barber *et al.*, 2005; Samuel *et al.*, 1997].

A2. Java

Much of the northeastern side of Java seems to have emerged since Pliocene time [e.g., Hall, 2009, Figure 14]. Built on the southern edge of the Eurasian landmass, the island stands above sea level in part because of late Cenozoic thrust faulting and crustal thickening and because of volcanism [e.g., Hall, 2002, p. 371; Hamilton, 1979, pp. 39–44]. This elevated terrain has then shed abundant sediment that has filled adjacent basins.

In a synthesis of sediment from western Java, *Clements and Hall* [2007] showed that the northern and southern quarters of that part of the island were submerged in late Miocene time. Then in a more comprehensive study, *Clements et al.* [2009] summarized evidence of middle Miocene and younger thrust faulting and north-south crustal shortening across the axis of the island, which flexed the terrain to the north down to form a basin [see also *Hall et al.*, 2007; *Yulianto et al.*, 2007]. This basin is developed best in the east and only modestly developed in the west. *Van Gorsel and Troelstra* [1981] described “an E-W-trending anticlinorium, about 250 km long and 20 km wide [that] during Miocene-Pliocene ... was a deep ‘foreland basin’, situated between the relatively stable Sunda Shelf in the north and the volcanic arc (axial ridge of Java) in the south.” They inferred north-south shortening as recent as Pleistocene and emergence at only 2.5 Ma of what had been deep water, 1000 m, at 5 Ma.

Elsewhere, regarding the date of the emergence of Java, *P. Lunt et al.* [2009] reported that “widespread, shallow marine limestone,” dated at 5.5–7.5 Ma using strontium isotope stratigraphy, overlies a middle-late Miocene unconformity in central and northeastern Java, suggesting that emergence occurred in Pliocene time. *Burckle* [1982] and *Saint-Marc and Suminta* [1979] reported marine diatoms and planktonic foraminifera from exposed late Miocene and Pliocene sediments in northeastern Java. *Umbgrove* [1946] described a rich assemblage of early Pliocene corals from central Java.

From these observations of Pliocene submergence, and following *Hall* [2001, Figure 10, 2009, Figure 14], we assume that approximately the northern quarter of Java and all of Madura and Kangean (the large and the much smaller islands to the northeast of Java) were submerged at 5 Ma.

A3. The Banda Arc, Sumba, and Timor

We found little specific information about the islands along the Banda Arc, but *Hall* [2001, Figure 10, 2009, Figure 14] shows many of the islands east of Java as totally or partly submerged at 5 Ma, as does *Wilson* [2008, Figure 1f] at 7 Ma. From west to east these are Bali, Lombok, Moyo, Sumbawa, Komodo, Flores, Lombok, Pantar, Alor, Wetar, Atauro, the Taninbar Islands of Wuliuru, Yamdena, and Larat, and the Kai Islands of Kai Besar and Kai Kecil.

A bit is known about a few of the smaller islands. Atauro, the small island just north of Timor, is rising rapidly today, at 0.47 mm/yr [*Chappell and Veeh*, 1978], and at that rate it must have emerged in last couple of Ma. Coral reefs at 700 m overlie volcanic rock older than 3.3 Ma [*Ely et al.*, 2011]. The small Tanimbar and Kai Islands, just east of the Banda Arc, also appear to have emerged in Pliocene or Quaternary time. Late Miocene marine sediment on the Taninbar Islands was folded in Pliocene time, presumably in response to the same collision of Australia with the Banda Arc as in Timor, and then overlain unconformably with Pleistocene sediment [*Charlton et al.*, 1991b]. From microorganisms *Fortuin and De Smet* [1991] inferred that sediment currently exposed on those islands lay at depth greater than 2 km at 5 Ma and merged at 2 Ma. Similarly for one of the Kai Islands, *van Marle and De Smet* [1990] reported exposures of Quaternary sediment whose microorganisms commonly live at depths of 1000 m, suggesting emergence since ~1 Ma. Finally, *Hall* [2001, Figure 10, 2009, Figure 14] shows the Aru Islands of Kobroor, Maikoor, Trangan, Wokam, and Workai as having been submerged at 5 Ma.

Abundant evidence demonstrates that beginning near 3 Ma, the northern margin of the Australian continent met the Banda Arc. As subduction of that margin began, slices of it became detached from the underlying lithosphere and were stacked to create the island of Timor [e.g., *Audley-Charles*, 1986; *Bowin et al.*, 1980; *Carter et al.*, 1976; *De Smet et al.*, 1990; *Johnston and Bowin*, 1981; *Veevers et al.*, 1978]. Moreover, in a number of cases, volcanism along the arc seems to have shut down at ~3 Ma [*Abbott and Chamalaun*, 1981]. Elsewhere, the chemistry of the volcanic rock shows contamination by Australian crust underthrust beneath the islands [*Herrington et al.*, 2010]. This same history seems to characterize many of the islands both along the arc and south of it.

Regarding the emergence of Timor, recent work with improved dating techniques and more detailed mapping has refined the history of the island, but not changed the inference that the island emerged since ~3 Ma or maybe a little before that time [e.g., *Haig*, 2012; *Keep and Haig*, 2010; *Nguyen et al.*, 2013; *Quigley et al.*, 2012; *van Marle*, 1991]. Similarly, islands just to the west of Timor, like Savu and Rote, seem to have emerged more recently, closer to 1 Ma [*Harris et al.*, 2009; *Rigg and Hall*, 2011; *Roosmawati and Harris*, 2009], like that of Sumba, yet farther west. *Fortuin et al.* [1994, 1997] inferred that Sumba lay as deep as 4500 m at

~6 Ma, and subsequently rose at an average rate of ~0.7 mm/yr. It did not emerge above sea level before 3 Ma, though deformation beneath it seems to have begun near 7 Ma [Rutherford *et al.*, 2001].

Thus, it appears that most of the islands east of Java along or south of the Banda Arc emerged since ~5 Ma.

A4. Borneo

Although Borneo has not been immune to late Cenozoic tectonics, it appears that the areal extent of the island has increased little since 5 Ma. Despite high erosion rates, inferred from thick sediment surrounding the island [Hall and Nichols, 2002], and the resulting flux of sediment that has rapidly filled accommodation space, folding and thrusting of the sedimentary layers seems to have compensated for an expansion of subaerial terrain. In fact, both Hall [2001, Figure 10, 2009, Figure 14] and Wilson [2008, Figure 1f] show Borneo and the small island of Maya, connected to the Malay Peninsula at 5 Ma and 7 Ma, respectively, suggesting that a larger area may have been emergent at those times than now.

Shortening of the sedimentary cover along the north side of the island has occurred since middle Miocene time [e.g., Hesse *et al.*, 2009; Hinz *et al.*, 1989; Hutchison, 1996; Morley, 2009; Morley *et al.*, 2003; Rangin *et al.*, 1990; Wannier, 2009], though some have questioned whether convergence continues [e.g., Cottam *et al.*, 2013b; Hall, 2013]. Seismic reflection profiles attest to abundant folding, which is exposed clearly on the northwestern margin of the island [Chambers and Daley, 1997; Franke *et al.*, 2008; Hesse *et al.*, 2010; Hinz *et al.*, 1989], and much of the sediment exposed on land is marine and late Cenozoic in age [Back *et al.*, 2005; Balaguru and Nichols, 2004; Hall, 2013; Hutchison, 1996]. Following, if not concurrent with, this folding, the highest mountain on the island, Mount Kinabalu, seems to have grown by intrusion of granite at 7–8 Ma [Cottam *et al.*, 2010, 2013b]. Moreover, GPS measurements show that shortening across this region continues at several millimeters per year [e.g., Simons *et al.*, 2007; Socquet *et al.*, 2006].

On the eastern side of the island, a large sedimentary basin, the Kutai Basin, has grown eastward largely by east-west crustal shortening, the creation of eroding terrain on the inland side, and progradation of deltas [Chambers *et al.*, 2004; Hamilton, 1979, p. 99; McClay *et al.*, 2000; Moss and Chambers, 1999; Moss *et al.*, 1997; Satyana *et al.*, 1999].

South of the Kutai Basin, the Meratus Mountains, which form a range parallel to the southeast coast, have grown since late Miocene time, presumably in response to north-south crustal shortening [Hall *et al.*, 2008; Satyana *et al.*, 1999; van de Weerd and Armin, 1992; Witts *et al.*, 2011]. GPS measurements suggest modest shortening of a few mm/yr across the range [Simons *et al.*, 2007].

Although some eastward and northward expansion of Borneo may have occurred since 5 Ma, the amount seems to be small, no more than 10%. Similarly, although some of the high terrain, Mount Kinabalu in the north and the Meratus Mountains in the southeast, may have grown higher since 5 Ma, we have no way to quantify the amount.

A5. Sulawesi, the Makassar Strait, and the Sula Islands

Sulawesi has undergone a complicated history, with amalgamation of its various arms only in the past few million years. Moreover, the westward movement of Sulawesi toward Borneo has narrowed the Makassar Strait, through which most of the present-day Indonesian Throughflow passes [e.g., Gordon *et al.*, 2003].

Seismic reflection profiles show clear late Miocene and younger folding on the eastern margin of the Makassar Strait [Bergman *et al.*, 1996]. Calvert [2000] reported late Miocene-Pliocene shallow marine sediment now cropping out on the west coast. In a series of schematic cross sections, van Leeuwen and Muhardjo [2005] show Plio-Quaternary emergence of much of the northerly trending belt of mountains that form the western side of the island, and the small island of Laut. They also report Quaternary reef limestone as high as 1000 m both along the western margin and on the northern arm of Sulawesi. In the north arm of Sulawesi, Hennig *et al.* [2014] reported late Miocene to Pliocene granitic intrusions and volcanic rock that has been exhumed rapidly, ~1–4 mm/yr. They infer rapid emergence of this region and cite Plio-Pleistocene sediment adjacent to the region as supporting evidence of rapid recent erosion.

Hall [2002, p. 416] stated “from about 5 Ma that there is clear evidence for contraction and uplift throughout the whole of Sulawesi and the beginning of the rise of the mountains in western Sulawesi,” an inference corroborated by Bellier *et al.* [2006]. GPS measurements of control points on the east and west sides suggest

as much as 10 mm/yr of NW-SE convergence between the margins, and therefore ~7 mm/yr of narrowing of the Makassar Strait [Rangin *et al.*, 1999; Simons *et al.*, 2007]. If this rate operated for the past 5 Myr, then the strait would have been ~15% wider at that time than today.

For the southern arm of the island, *BouDagher-Fadel* [2002] reported late middle Miocene to early Pliocene shallow water deposits and early Pliocene emergence, though the areal extent of this emergence is not clear. Similarly, *Bromfield and Renema* [2011], *Grainge and Davies* [1985], and *Mayall and Cox* [1988] presented evidence of a late Miocene widespread transgression and reef development at the southern end of this arm of Sulawesi, followed by a Pliocene emergence of the region. *Sudarmono* [2000], *van Leeuwen et al.* [2010], and *Camplin and Hall* [2014] reported similar histories—Miocene submergence and Pliocene emergence—for the Bone Basin, which lies between the southern and southeast arms of the island, and the Bone Mountains along the eastern side of the southern arm.

For the southeastern arm, a submerged terrain (Tukang Besi platform) seems to have collided in middle Miocene time with the island of Buton [Smith and Silver, 1991], which lies east of the southeast arm. The island is covered by only mildly deformed younger rock, and in some areas that younger rock consists of “fine-grained pelagic foraminiferal chalk” [Smith and Silver, 1991]. *Fortuin et al.* [1990] reported abundant marine microorganisms dated at 6 Ma from Buton, some coming from as deep as 1000 m [Fortuin and De Smet, 1991]. Thus, they concluded that at least part of the island emerged since ~5 Ma, and we presume the same for the smaller adjacent islands of Kabaena and Muna.

For the eastern arm, rapid westward displacement of small slivers of crust, including the islands of Peleng and Banggai, has led to their collision with the rest of the arm and their accretion on the eastern end of the peninsula [e.g., *Charlton*, 1996; *Ferdian et al.*, 2010; *Hamilton*, 1979, pp. 159, 173–174, 181–183; *Silver et al.*, 1983]. *Van Leeuwen et al.* [2010], *Villeneuve et al.* [1998, 2000, 2001], and *Watkinson et al.* [2011] date this collision as Pliocene. *Davies* [1990] showed that one block (Banggai) collided with the peninsula between 5.3 and 3.8 Ma, when clastic sedimentation on limestone began.

Despite all of the evidence of post-Miocene deformation and emergence, only a relatively small fraction of the island and adjacent islands seems to have been below sea level at 5 Ma. *Hall* [2001, Figure 10, 2009, Figure 14] shows parts of all of the arms as submerged at 5 Ma, but the fraction is only 10–20% of the present-day emergent region. By contrast, *Wilson* [2008, Figure 1f] shows the entire northern arm and much of the eastern arm submerged at 7 Ma. In a cross section across the northern arm of Sulawesi, *van Leeuwen and Muhardjo* [2005] show most of its emergence to be Plio-Quaternary.

Farther east, *Hall* [2001, Figure 10] shows roughly half of the area of the Sula Islands of Mangole, Sanane, Sulabesi, and Taliabu submerged at 5 Ma.

A6. Buru, Seram, and Surrounding Islands

Buru and Seram lie between Sulawesi and New Guinea (Figure 2). Although Buru has undergone only mild folding, and Seram has been caught in a subduction zone, they share similar histories of pre-Pliocene sedimentation [e.g., *Charlton*, 2000].

For Buru, mild deformation apparently has elevated the island [Charlton, 2000]. *Fortuin et al.* [1988] showed emergence at 6 Ma, but in a more recent summary, *Fortuin and De Smet* [1991] reported depths of 1000 m at 3 Ma. In his synthesis, however, *Hall* [2001, Figure 10] shows only the axial part of the island submerged at 5 Ma.

Much, if not all, of Seram and the small island of Ambon to its southeast seem to have emerged since 5 Ma. Although recent work by *Pownall et al.* [2013, 2014] suggests that much of the high-grade rock exposed on Seram has been exhumed by normal faulting, in some places deeper rock has undergone melting in the past few million years [Linthout and Helmers, 1994; Linthout *et al.*, 1996, 1997]. At present, the seafloor north of Seram is underthrusting the island, and from folding of sediment offshore, *Pairault et al.* [2003a, 2003b] inferred a Pliocene onset of that subduction. Consistent with this timing, *Charlton* [2000] reported that the volcanism on Seram and Ambon is Pliocene. Interpretative cross sections given by *Pairault et al.* [2003a, 2003b] show Seram as having emerged entirely in Pliocene time. From sediment cropping out on the southwest side of Seram, *De Smet et al.* [1989] inferred depths of 500 m between ~3.5 and ~1 Ma. *Fortuin and De Smet* [1991] reported this as depths of 2000 m at 2 Ma.

Although data seems sparse, we follow others [e.g., Hall, 2002, p. 417; Pairault *et al.*, 2003a, 2003b] and assume that Seram emerged entirely since 5 Ma, and so did much of Buru.

A7. New Guinea and Adjacent Islands

Much of New Guinea and the islands immediately adjacent to it—Batanta, Biak, Dolak, Kiwai, Komoren, Misool, Numfor, Salawati, Supiori, Waigeo, and Yapen—seem to have emerged since 5 Ma [Hall, 2001, Figure 10, 2009, Figure 14, 2012, Figure 3.12] or since 7 Ma [Wilson [2008, Figure 1f]. Moreover, although the high axial chain of mountains began to grow before that time [e.g., Baldwin *et al.*, 2012; Cloos *et al.*, 2005; Davies, 2012], high topography also seems to postdate 5 Ma [Hall, 2009, p. 156, 2012, p. 57].

In western New Guinea, Dow and Sukanto [1984] reported that deformation began no later than late Miocene, but with the oldest-coarse clastic sediment as early Pliocene, they inferred that mountains developed in Pliocene time. They stated that 8000 m of Pliocene and Quaternary clastic sediment accumulated in the bay east of the Bird's Head, which clearly requires deep erosion of material from nearby. In a synthesis of sedimentary rock from western New Guinea, Pieters *et al.* [1983] showed that limestone covered much of the region until late Miocene time. On the southwest side of the Bird's Head of western New Guinea, late Miocene to early Pliocene folded rock is overlain by Pliocene and younger shallow marine sediment [Bailly *et al.*, 2009; Pubellier and Ego, 2002]. Thus, much of this region seems to have been below sea level until Pliocene time.

Salawati Island, just west of the Bird's Head, was connected to the New Guinea in Miocene time, and Pliocene separation seems to have occurred during modest Pliocene rifting between them [Froidevaux, 1978]. Extensive Miocene to lower Pliocene reefs cover Salawati island, and we presume that emergence of them is young. Similarly, in summary cross sections based on extensive seismic reflection profiling between Misool and Salawati, Sapin *et al.* [2009] showed this area emerging in Pliocene time.

Waigeo Island and the smaller island of Batanta, north of the Bird's Head of western New Guinea, also appear to be young. On Waigeo, ophiolite is capped by 2000 m of Miocene limestone, which shows coral reefs at the base but higher in the section it is rich in benthic foraminifera and then planktonic foraminifera [Charlton *et al.*, 1991a]. Charlton *et al.* [1991a] allow for some of the limestone to be Pliocene in age. Pliocene folding created two broad anticlines and presumably elevated the island above sea level.

The main axial range of New Guinea seems to have started to grow in middle to late Miocene time in the northern part, and the locus of deformation migrated southward [Cloos *et al.*, 2005; Hill and Hall, 2003; Quarles van Ufford and Cloos, 2005; Sapiie and Cloos, 2004]. Thermochronological studies show rapid exhumation of northern New Guinea in its western part at 14–12 Ma [Hill and Hall, 2003] and between 8 and 5 Ma in its eastern part [Crowhurst *et al.*, 1996]. On the south flank, however, thermochronology shows exhumation as young as 4 Ma [Hill and Gleadow, 1989; Hill and Raza, 1999; McDowell *et al.*, 1996] and deep incision since 2 Ma [Weiland and Cloos, 1996]. From the pattern of exhumation and its timing, Weiland and Cloos [1996] inferred that although exhumation had begun by 7 Ma along the crest of the range and south of it, elevations high enough to affect orographic precipitation did not occur until 5 Ma or more recently. Coinciding with this rapid Pliocene exhumation, a “pronounced sedimentological change occurred in the Pliocene-Pleistocene (~5 Ma), when the coarseness of deposits along the flanks of the Central Range increased dramatically for hundreds of kilometers along strike” [Quarles van Ufford and Cloos, 2005]. Relying on Hill's [1991] estimate of 100 km of shortening across the southern part of the range, Crowhurst *et al.* [1996] inferred that that part of the belt formed in Pliocene time. The rise of these mountains then shed sediment southward, and a late Pliocene and Quaternary southward migration of sediment buried a carbonate platform to the south of the mountain belt [Pigram and Symonds, 1991].

On the northeastern coast, abundant coral reefs on the Huon Peninsula attest to rapid emergence of this region [Chappell, 1974]. Moreover, the sudden deposition of coarse clastic material southwest of the Adelbert and Finisterre Ranges in what had been a deep trough, and derived from those mountains, shows that they emerged at 3.0–3.7 Ma [Abbott, 1995; Abbott *et al.*, 1994]. Farther west a large ophiolitic body, the Cyclops ophiolite was emplaced at ~20 Ma, but apparently Pliocene thrust faulting and crustal shortening elevated the region above sea level [Monnier *et al.*, 1999].

Along the Papuan Peninsula of southeastern New Guinea, rapid deformation in Pliocene time seems to have elevated much of this region above sea level [e.g., Davies *et al.*, 1996; Dow, 1977]. Plio-Quaternary coral reefs

surrounding the island can be found at heights of 600 m [Smith and Davies, 1976]. Paleogeographic maps show the nearby D'Entrecasteaux islands (Goodenough, Moratau, Muyua, Normanby, and Sudest) just north of the Papuan Peninsula to have been submerged at 5 Ma [Hall, 2001, Figure 10] and at 7 Ma Wilson [2008, Figure 1f], but the geologic history of these islands is sufficiently complicated that the history of emergence of these islands, or perhaps submergence adjacent to them, is surely not simple [e.g., Baldwin *et al.*, 2012; Davies and Warren, 1988; Fitz and Mann, 2013; Little *et al.*, 2011]. Davies and Warren [1988] and Little *et al.* [2011] inferred that domal uplifts brought deeply exhumed rock above sea level in Pliocene time and since 6 Ma, respectively.

A8. Islands East and Northeast of New Guinea

We have found less information about islands east of New Guinea, perhaps because little change has occurred on these islands. Relatively flat Miocene limestone caps parts of New Britain [e.g., Lindley, 2006], but accurate dates of vertical movements seem to have been studied in few places. Bromfield and Renema [2011] discuss limestone in eastern New Britain, the bottom part of which they dated at 13–14 Ma, but which seems to be as young as 4 Ma in places. Thus, at least part of eastern New Britain seems to have emerged since 5 Ma. Riker-Coleman *et al.* [2006] reported a series of terraces each with corals atop them, and they inferred a present-day rate of emergence of this region between 0.4 and 2.1 mm/yr. Although the rate is not constrained well, the succession of terraces reaching 270 m above sea level attests to sustained emergence. Mostly as a guess, we assume that the island was two thirds its present size at 5 Ma, though neither Hall [2002] nor Wilson [2008] indicates an island much smaller than today at 5 and 7 Ma, respectively. Also, Hamilton [1979, p. 289] stated that Quaternary volcanoes built the islands of Umboi, Long, and Karkar to the west of New Britain.

Early Miocene limestone was deposited on earlier Cenozoic volcanic rock, apparently marking a period of sustained subsidence of the region that includes New Ireland, Lavongai (New Hanover), Mussau, and Manus. Marine deposition, of largely limestone, continued through Mio-Pliocene time, until the sequence that has emerged was tilted apparently in Plio-Quaternary time [Exon and Marlow, 1988; Francis, 1988; Hohnen, 1978; Lindley, 2006; Stewart and Sandy, 1988]. Moreover, raised marine terraces consisting of coral reach hundreds of meters in elevation [Hohnen, 1978]. Thus, we presume that essentially the entire region that includes these islands remained submerged until more recently than 5 Ma.

Although much of Bougainville seems to have submerged in early Miocene time, volcanic rock covers the much of the island and gives little evidence of post-Miocene emergence [Blake and Mieztis, 1967]. Only the northern end of the island and the adjacent island of Buka, which are covered by Quaternary reef limestone seem to have emerged since 5 Ma.

We ignore the region farther east, the Solomon Islands, but we note that abundant evidence suggests emergence of islands in this region since 5 Ma [Brunns *et al.*, 1989; Cowley *et al.*, 2002; Hughes and Turner, 1977; Mann and Asahiko, 2004; Phinney *et al.*, 2004].

A9. Halmahera and Surrounding Islands

Most of Halmahera and the surrounding islands of Obi, Bacan, Kasiruta, and Morotai seem to have emerged since 5 Ma.

Halmahera consists of two different parts: a long narrow belt of volcanoes and volcanic rock in the west and ophiolitic mélange and oceanic crust with carbonate platforms in the east [Hall, 1987; Hall *et al.*, 1988a]. For Bacan, which lies southwest of Halmahera, the ophiolite has been thrust atop rock possibly as old as Precambrian, but otherwise, its history seems to be similar to that of Halmahera [Malaihollo and Hall, 1996]. Volcanism seems to have begun at 11 Ma on Obi and progressed north to Bacan and Halmahera [Baker and Malaihollo, 1996]. Ages from Bacan and Halmahera begin at 7.5 Ma and 7.8 Ma, respectively. Thus, much of these islands did not exist at all until late Miocene time. Hall *et al.* [1988b] reported late Miocene reef limestone at an elevation of 1000 m over the eastern part of Halmahera. The Pliocene sequence includes marl at the base, with sandstone, siltstone, and conglomerate above, and Hall *et al.* [1988b] describe an early Pliocene "change from the stable conditions of carbonate deposition across east and central Halmahera with a transition from limestones to marls, followed rapidly by an increase in the amount of siliciclastic debris which was deposited as submarine fan turbidites." Nichols and Hall [1991] described a major phase of folding that requires 60 km of shortening across the eastern part of the island. In a preliminary summary of the structure of Halmahera, Hall and Nichols [1990] stated that this deformation began at 3 Ma. Later, Hall [2000] showed this to represent 60%

shortening, which almost surely would have raised a surface at sea level, with reefs, to an elevation of 1–2 km. Also, the geology of Morotai, just northeast of Halmahera, resembles that of Halmahera [Hall *et al.*, 1991], and Hall [2001, Figure 10, 2009, Figure 14] showed the island as submerged at 5 Ma.

There seems little doubt that most of Halmahera and surroundings underwent Pliocene crustal shortening [e.g., Hall, 2002, p. 382] and that most of the region emerged since 5 Ma.

A10. Philippines, Palawan, and Adjacent Islands

North of Halmahera, only small islands rise above sea level: the Sangihe Islands along the Sangihe volcanic arc and the Talaud Islands, of which Karakelong is the biggest. Karakelong seems to have emerged only in Pleistocene time [Moore *et al.*, 1981]. Macpherson *et al.* [2003] report that the Sangihe Islands are Quaternary in age.

We have found little information that places quantitative constraints on the emergence above sea level of Mindanao or other islands in the southern part of the Philippines. Nevertheless, several observations suggest that much of the Mindanao may have emerged since the latest Miocene Epoch, if not since 5 Ma. Pubellier *et al.* [1991] described the main late Cenozoic geologic event as a suturing of two blocks that began at circa 5 Ma [Sajona *et al.*, 1994, 2000]. Suturing was completed by early Pliocene time. The western block seems to be underlain continental crust that had undergone extension in early and middle Miocene time, and therefore presumably crustal thinning and submergence. The eastern block includes the northward continuation of the Sangihe Arc but seems to have been emergent during much of the Miocene Epoch. The collision of the two blocks led to folding, some overturned, and thrust faulting that attest to kilometers, if not much more, of crustal shortening [Pubellier *et al.*, 1996]. Moreover, Quebral *et al.* [1996] describe Pliocene unconformities in several places, consistent with erosion of rock that had undergone deformation just before that time. Also, Queaño [2005] described late Miocene-Pliocene turbidites that crop out on the western side of the Pujada Peninsula, southeastern Mindanao, but this is a small area, and it is not clear how much of the region has emerged since ~5 Ma. Based in part on Hall's [2001, Figure 10, 2009, Figure 14] syntheses, we assume that half of the island, plus the small island of Siargao to its northeast, emerged since 5 Ma.

Regarding other islands, Sajona *et al.* [1997] associated a late Miocene to Pliocene unconformity on Leyte with the initiation of volcanic activity associated with subduction at the Philippine trench. Moreover, Dimalanta *et al.* [2009] noted that much of southwestern Leyte is covered by Plio-Quaternary limestone. It is difficult to know what fraction of the island has emerged since 5 Ma, but it would seem that at least half of it has. Similarly, Faustino *et al.* [2003] point out that most of Bohol is covered by Plio-Quaternary limestone and that although the island did not rise uniformly in Quaternary time, it seems to have emerged since ~5 Ma. Faustino *et al.* [2003] also noted that a northerly trending anticline passing through Cebu then began to grow in late Miocene time and exposes the same Plio-Quaternary limestone. Thus, it seems that Cebu, and presumably Siquijor to its south, also emerged since 5 Ma. To the west of Bohol, Los Negros also shares some of the same history as Bohol [Faustino *et al.*, 2003], but it is not apparent that the entire island emerged in the past 5 Ma. Farther north, for Samar Island, Travaglio *et al.* [1978] reported that much of the island was submerged in early Miocene time, when coralline limestone was deposited on sandstone and shale. They reported that emergence began late Miocene time, with different parts of the island rising different amounts at different times. With doubts about the precise date when the island merged, we assume that half of it rose by 5 Ma.

We suspect that as much as half of Mindanao and adjacent islands might have been submerged until as recently as 5 Ma. Hall [2001, Figure 10, 2009, Figure 14] showed much of southern Mindanao submerged at that time, but gave no discussion; after considering presently submerged surrounding terrain that he considers to have been above sea level at 5 Ma, we estimate approximately 30% of the current land above sea level was submerged. By contrast, but again without discussing the region, Wilson [2008, Figure 1f] and Wilson and Moss [1999, Figure 10] showed virtually all of southern Philippines, Mindanao, Leyte, and adjacent islands, as below sea level at 7 and 8 Ma, respectively.

Discussions by Holloway [1982], Müller [1991], and Rangin and Silver [1991] suggest that Palawan had emerged by middle Miocene time. cursory study of Balabac to its southwest [John, 1963] suggests the same. Similarly, Rangin [1989] shows volcanoes on the islands of Basilan and Jolo along the Sulu Ridge as built on islands dating from early Miocene or earlier, like rock exposed on Tawi-Tawi. Farther east in the Philippines, the geologic history of Panay seems to resemble that of the Palawan and shows no evidence of post 5 Ma emergence [Yumul *et al.*, 2009].

Acknowledgments

We thank Kerry Emanuel for several helpful suggestions, and R. Hall, D. V. Kent, and an anonymous reviewer for encouragement and advice. This research was supported in part by the National Science Foundation under grants AG5 1136466 and 1136480. We provide no new data, but everything that we have used is cited (and that makes a long list of references cited).

References

- Abbott, L. D. (1995), Neogene tectonic reconstruction of the Adelbert-Finisterre-New Britain collision, northern Papua New Guinea, *J. Southeast Asian Earth Sci.*, *11*, 33–52.
- Abbott, L. D., E. A. Silver, P. R. Thompson, M. V. Filewicz, C. Schneider, and Abdoerrias (1994), Stratigraphic constraints on the development and timing of arc-continent collision in northern Papua New Guinea, *J. Sediment. Res.*, *B64*, 169–183.
- Abbott, M. J., and F. H. Chamalaun (1981), Geochronology of some Banda Arc volcanics, in *The Geology and Tectonics of Eastern Indonesia*, Spec. Publ. No. 2, edited by A. J. Barber and S. Wiriyosujono, pp. 253–268, Geological Research and Development Centre, Bandung.
- Amiotte Suchet, P., J.-L. Probst, and W. Ludwig (2003), Worldwide distribution of continental rock lithology: Implications for the atmospheric/soil CO₂ uptake by continental weathering and alkalinity river transport to the oceans, *Global Biogeochem. Cycles*, *17*(2), 1038, doi:10.1029/2002GB001891.
- As-syakur, A. R., T. Tanaka, T. Osawa, and M. S. Mahendra (2013), Indonesian rainfall variability observation using TRMM multi-satellite data, *Int. J. Remote Sens.*, *34*, 7723–7738, doi:10.1080/01431161.2013.826837.
- Audley-Charles, M. G. (1986), Rates of Neogene and Quaternary tectonic movements in the southern Banda Arc based on micropalaeontology, *J. Geol. Soc. London*, *143*, 161–175.
- Back, L., K. Russ, Z. Liu, K. Inoue, J. Zhang, and B. Otto-Bliesner (2013), Global hydrological cycle response to rapid and slow global warming, *J. Clim.*, *26*, 8781–8786.
- Back, S., H. J. Tioe, T. X. Thang, and C. K. Morley (2005), Stratigraphic development of synkinematic deposits in a large growth-fault system, onshore Brunei Darussalam, *J. Geol. Soc. London*, *162*, 243–257.
- Bailly, V., M. Pubellier, J. C. Ringenbach, J. de Sigoyer, and F. Sapin (2009), Deformation zone Jumps, in a young convergent setting; the Lengguru fold-and-thrust belt, New Guinea Island, *Lithos*, *113*, 306–317, doi:10.1016/j.lithos.2009.08.013.
- Baker, S., and J. Malaihollo (1996), Dating of Neogene igneous rocks in the Halmahera region: Arc initiation and development, in *Tectonic Evolution of Southeast Asia*, Spec. Publ., vol. 106, edited by R. Hall and D. J. Blundell, pp. 499–509, Geol. Soc., London.
- Balaguru, A., and G. Nichols (2004), Tertiary stratigraphy and basin evolution, southern Sabah (Malaysian Borneo), *J. Asian Earth Sci.*, *23*, 537–554.
- Baldwin, S. L., P. G. Fitzgerald, and L. E. Webb (2012), Tectonics of the New Guinea region, *Annu. Rev. Earth Planet. Sci.*, *40*, 495–520.
- Barber, A. J., and M. J. Crow (2005), Structure and structural history, in *Sumatra: Geology, Resources and Tectonic Evolution*, Mem., vol. 31, edited by A. J. Barber, M. J. Crow, and J. S. Milsom, pp. 175–233, Geol. Soc., London.
- Barber, A. J., M. J. Crow, and M. E. M. De Smet (2005), Tectonic evolution, in *Sumatra: Geology, Resources and Tectonic Evolution*, Mem., vol. 31, edited by A. J. Barber, M. J. Crow, and J. S. Milsom, pp. 234–259, Geol. Soc., London.
- Barreiro, M., G. Philander, R. Pacanowski, and A. Fedorov (2006), Simulations of warm tropical conditions with application to middle Pliocene atmospheres, *Clim. Dyn.*, *26*, 349–365.
- Bartoli, G., M. Sarnthein, M. Weinelt, H. Erlenkeuser, D. Garbe-Schönberg, and D. W. Lea (2005), Final closure of Panama and the onset of Northern Hemisphere glaciation, *Earth Planet. Sci. Lett.*, *237*, 33–44.
- Bartoli, G., B. Hönisch, and R. E. Zeebe (2011), Atmospheric CO₂ decline during the Pliocene intensification of Northern Hemisphere glaciations, *Paleoceanography*, *26*, PA4213, doi:10.1029/2010PA002055.
- Bellier, O., M. Sébrier, D. Seward, T. Beaudouin, M. Villeneuve, and E. Putranto (2006), Fission track and fault kinematics analyses for new insight into the late Cenozoic tectonic regime changes in west-central Sulawesi (Indonesia), *Tectonophysics*, *413*, 201–220.
- Berggren, W. A., and C. D. Hollister (1974), Paleogeography, paleobiogeography and the history of circulation in the Atlantic Ocean, in *Studies in Paleo-Oceanography*, Spec. Publ., vol. 20, edited by W. W. Hay, pp. 126–186, Soc. Econ. Paleontol. Mineral, Tulsa, Okla.
- Bergman, S. C., D. Q. Coffield, J. P. Talbot, and R. A. Garrard (1996), Tertiary tectonic and magmatic evolution of western Sulawesi and Makassar Strait, Indonesia: Evidence for a Miocene continent-continent collision, in *Tectonic Evolution of Southeast Asia*, edited by R. Hall and D. J. Blundell, *Geol. Soc. London Spec. Publ.*, *106*, 391–428.
- Berner, R. A., and Z. Kothavala (2001), GEOCARB III: A revised model of atmospheric CO₂ over Phanerozoic time, *Am. J. Sci.*, *301*, 182–204.
- Blake, D. H., and Y. Miezitis (1967), *Geology of Bougainville and Buka Islands, New Guinea*, Bulletin, vol. 93, 56 pp., Bureau of Mineral Resources, Australia.
- BouDagher-Fadel, M. K. (2002), The stratigraphical relationship between planktonic and larger benthic foraminifera in middle Miocene to lower Pliocene carbonate facies of Sulawesi, Indonesia, *Micropaleontology*, *48*, 153–176.
- Bowin, C., G. M. Purdy, C. Johnston, G. Shor, L. Lawver, H. M. S. Hartono, and P. Jezek (1980), Arc-continent collision in Banda Sea region, *Am. Assoc. Pet. Geol. Bull.*, *64*, 868–915.
- Brierley, C. M., and A. V. Fedorov (2010), Relative importance of meridional and zonal sea surface temperature gradients for the onset of the Ice Ages and Pliocene-Pleistocene climate evolution, *Paleoceanography*, *25*, PA2214, doi:10.1029/2009PA001809.
- Brierley, C. M., A. V. Fedorov, Z.-h. Liu, T. D. Herbert, K. T. Lawrence, and J. P. LaRiviere (2009), Greatly expanded tropical warm pool and weakened Hadley Circulation in the early Pliocene, *Science*, *323*, 1714–1718.
- Bromfield, K., and W. Renema (2011), Comparison of ⁸⁷Sr/⁸⁶Sr isotope and biostratigraphic ages of uplifted fossil reefs in the Indo-Pacific: Indonesia, Papua New Guinea and Fiji, *Aust. J. Earth Sci.*, *58*, 61–73.
- Bruns, T. R., J. G. Vedder, and A. K. Cooper (1989), Geology of the Shortland Basin region, Central Solomons Trough, Solomon Islands—Review and new findings, in *Geology and Offshore Resources of the Pacific Island Arcs—Solomon Islands and Bougainville, Papua New Guinea Regions*, Earth Science Series, vol. 12, edited by J. G. Vedder and T. R. Bruns, pp. 93–123, Circum-Pacific Council for Energy and Mineral Resources, Houston, Tex.
- Burckle, L. H. (1982), Diatom biostratigraphy of late Miocene and Pliocene sediments of eastern Java (Indonesia), *Mar. Micropaleontol.*, *7*, 363–368.
- Calvert, S. J. (2000), The Cenozoic evolution of the Lariang and Karama Basins, in *Proceedings Indonesian Petroleum Association, 27th Annual Convention*, pp. 505–511, Indonesian Petroleum Association, Jakarta.
- Campbell, D. J., and R. Hall (2014), Neogene history of Bone Gulf, Sulawesi, Indonesia, *Mar. Pet. Geol.*, *57*, 88–108.
- Cane, M. A., and P. Molnar (2001), Closing of the Indonesian seaway as a precursor to east African aridification around 3–4 million years ago, *Nature*, *411*, 157–162.
- Cannariato, K. G., and A. C. Ravelo (1997), Plio-Pleistocene evolution of eastern tropical Pacific rate of circulation and thermocline depth, *Paleoceanography*, *12*, 805–820, doi:10.1029/97PA02514.
- Carbone, R. E., J. Wilson, T. D. Keenan, and J. M. Hacker (2000), Tropical island convection in the absence of significant topography. Part I: Life cycle of diurnally forced convection, *Mon. Weather Rev.*, *128*, 3459–3480.
- Carter, D. J., M. G. Audley-Charles, and A. J. Barber (1976), Stratigraphical analysis of island arc-continent collision in eastern Indonesia, *J. Geol. Soc. London*, *132*, 179–198.

- Chaisson, W. (1995), Planktonic foraminiferal assemblages and paleoceanographic change in the trans-tropical Pacific Ocean: A comparison of west (Leg 130) and east (Leg 138), latest Miocene to Pleistocene, in *Proc. ODP, Sci. Results*, vol. 138, edited by N. G. Pisias et al., pp. 555–597, Ocean Drilling Program, College Station, Tex.
- Chaisson, W. P., and R. M. Leckie (1993), High-resolution Neogene planktonic foraminifer biostratigraphy of Site 806, Ontong Java Plateau (western equatorial Pacific), in *Proc. ODP, Sci. Results*, vol. 130, edited by W. H. Berger et al., pp. 137–178, Ocean Drilling Program, College Station, Tex.
- Chaisson, W. P., and A. C. Ravelo (2000), Pliocene development of the east–west hydrographic gradient in the tropical Pacific, *Paleoceanography*, *15*, 497–505, doi:10.1029/1999PA000442.
- Chambers, J. L. C., and T. E. Daley (1997), A tectonic model for the onshore Kutai Basin, East Kalimantan, in *Petroleum Geology of Southeast Asia*, Spec. Publ., vol. 126, edited by A. J. Fraser, S. J. Matthews, and R. W. Murphy, pp. 375–393, Geol. Soc., London.
- Chambers, J. L. C., I. Carter, I. R. Cloke, J. Craig, S. J. Moss, and D. W. Paterson (2004), Thin-skinned and thick-skinned inversion-related thrusting—a structural model for the Kutai Basin, Kalimantan, Indonesia, in *Thrust Tectonics and Hydrocarbon Systems*, Mem., vol. 82, edited by K. R. McClay, pp. 614–634, AAPG, Tulsa, Okla.
- Chappell, J. (1974), Geology of coral terraces, Huon Peninsula, New Guinea: A study of Quaternary tectonic movements and sea-level changes, *Geol. Soc. Am. Bull.*, *85*, 553–570.
- Chappell, J., and H. H. Veeh (1978), Late Quaternary tectonic movements and sea-level changes at Timor and Atauro Island, *Geol. Soc. Am. Bull.*, *89*, 356–368.
- Charlton, T. R. (1986), A plate tectonic model of the eastern Indonesian collision zone, *Nature*, *319*, 394–396.
- Charlton, T. R. (1996), Correlation of the Salawati and Tomori Basins, eastern Indonesia: A constraint on left-lateral displacements of the Sorong fault zone, in *Tectonic Evolution of Southeast Asia*, Spec. Publ., vol. 106, edited by R. Hall and D. J. Blundell, pp. 465–481, Geol. Soc., London.
- Charlton, T. R. (2000), Tertiary evolution of eastern Indonesia collision complex, *J. Asian Earth Sci.*, *18*, 603–631.
- Charlton, T. R., R. Hall, and E. Partoyo (1991a), The geology and tectonic evolution of Waigeo Island, NE Indonesia, *J. Southeast Asian Earth Sci.*, *6*, 289–297.
- Charlton, T. R., M. E. M. de Smet, H. Samodra, and S. J. Kaye (1991b), The stratigraphic and structural evolution of the Tanimbar islands, eastern Indonesia, *J. Southeast Asian Earth Sci.*, *6*, 343–358.
- Chiang, J. C. H., and A. H. Sobel (2002), Tropical tropospheric temperature variations caused by ENSO and their influence on the remote tropical climate, *J. Clim.*, *15*, 2616–2631.
- Clements, B., and R. Hall (2007), Cretaceous to late Miocene stratigraphic and tectonic evolution of west Java, Proceedings, Indonesian Petroleum Association, Thirty-First Annual Convention and Exhibition, Jakarta.
- Clements, B., R. Hall, H. R. Smyth, and M. A. Cottam (2009), Thrusting of a volcanic arc: A new structural model for Java, *Pet. Geosci.*, *15*, 159–174.
- Cloos, M., B. Sapiie, A. Quarles van Ufford, R. J. Weiland, P. Q. Warren, and T. P. McMahon (2005), *Collisional Delamination in New Guinea: The Geotectonics of Subducting Slab Breakoff*, Spec. Pap., vol. 400, 51 pp., Geol. Soc. Am., Boulder, Colo.
- Cottam, M. A., R. Hall, C. Sperber, and R. Armstrong (2010), Pulsed emplacement of the Mount Kinabalu granite, northern Borneo, *J. Geol. Soc. London*, *167*, 49–60.
- Cottam, M. A., R. Hall, and A. A. Ghani (2013a), Late Cretaceous and Cenozoic tectonics of the Malay Peninsula constrained by thermochronology, *J. Asian Earth Sci.*, *76*, 241–257.
- Cottam, M. A., R. Hall, C. Sperber, B. P. Kohn, M. A. Forster, and G. E. Batt (2013b), Neogene rock uplift and erosion in northern Borneo: Evidence from the Kinabalu granite, Mount Kinabalu, *J. Geol. Soc. London*, *170*, 805–816.
- Cowley, S., P. Mann, M. F. Coffin, and T. H. Shipley (2002), Oligocene to recent tectonic history of the Central Solomon intra-arc basin as determined from marine seismic reflection data and compilation of onland geology, *Tectonophysics*, *389*, 267–307.
- Cronin, T. W., K. A. Emanuel, and P. Molnar (2015), Island precipitation enhancement and the diurnal cycle in radiative-convective equilibrium, *Q. J. R. Meteorol. Soc.*, doi:10.1002/qj.2443, in press.
- Crowhurst, P. V., K. C. Hill, D. A. Foster, and A. P. Bennett (1996), Thermochronological and geochemical constraints on the tectonic evolution of northern Papua New Guinea, in *Tectonic Evolution of Southeast Asia*, Spec. Publ., vol. 106, edited by R. Hall and D. J. Blundell, pp. 525–537, Geol. Soc., London.
- Dai, A., T. Qian, K. E. Trenberth, and J. D. Milliman (2009), Changes in continental freshwater discharge from 1948 to 2004, *J. Clim.*, *22*, 2773–2992.
- Davies, H. L. (2012), The geology of New Guinea—The cordilleran margin of the Australian continent, *Episodes*, *35*, 87–102.
- Davies, H. L., and R. G. Warren (1988), Origin of eclogite-bearing, domed, layered metamorphic complexes (“core complexes”) in the D’Entrecasteaux Islands, Papua New Guinea, *Tectonics*, *7*, 1–21, doi:10.1029/TC007i001p00001.
- Davies, H. L., R. D. Winn, and P. KenGemar (1996), Evolution of the Papuan Basin—A view from the orogen, in *Petroleum Exploration, Development and Production in Papua New Guinea, Proceedings of the Third PNG Petroleum Convention*, edited by P. G. Buchanan, pp. 53–62, Papua New Guinea Chamber of Mines and Petroleum, Port Moresby, Papua New Guinea.
- Davies, I. C. (1990), Geological and exploration review of the Tomori PSC, eastern Indonesia, in *Proceeding Indonesian Petroleum Association, Nineteenth Annual Convention*, pp. 41–67, Indonesian Petroleum Association, Jakarta.
- Dayem, K. E., D. C. Noone, and P. Molnar (2007), Tropical western Pacific warm pool and maritime continent precipitation rates and their contrasting relationships with the Walker Circulation, *J. Geophys. Res.*, *112*, D06101, doi:10.1029/2006JD007870.
- Dekens, P. S., A. C. Ravelo, and M. D. McCarthy (2007), Warm upwelling regions in the Pliocene warm period, *Paleoceanography*, *22*, PA3211, doi:10.1029/2006PA001394.
- Dekens, P. S., A. C. Ravelo, M. D. McCarthy, and C. A. Edwards (2008), A 5 million year comparison of Mg/Ca and alkenone paleothermometers, *Geochem. Geophys. Geosyst.*, *9*, Q10001, doi:10.1029/2007GC001931.
- De Smet, M. E. M., and A. J. Barber (2005), Tertiary stratigraphy, in *Sumatra: Geology, Resources and Tectonic Evolution*, Mem., vol. 31, edited by A. J. Barber, M. J. Crow, and J. S. Milsom, pp. 86–97, Geol. Soc., London.
- De Smet, M. E. M., P. A. Sumosastro, I. Siregar, L. J. Van Marle, S. R. Troelstra, and A. R. Fortuin (1989), Late Cenozoic geohistory of Seram, Indonesia, *Geol. Mijnbouw*, *68*, 221–235.
- De Smet, M. E. M., A. R. Fortuin, S. R. Troelstra, L. J. Van Marle, M. Karmini, S. Tjokrosapoetro, and S. Hadiwasastro (1990), Detection of collision-related vertical movements in the outer Banda Arc (Timor, Indonesia), using micropaleontological data, *J. Southeast Asian Earth Sci.*, *4*, 337–356.
- Dessert, C., B. Dupré, J. Gaillardet, L. M. François, and C. J. Allègre (2003), Basalt weathering laws and the impact of basalt weathering on the global carbon cycle, *Chem. Geol.*, *202*, 257–273.

- Dimalanta, C. B., G. P. Yumul Jr., L. O. Suerte, and R. A. Tamayo Jr. (2009), Geology of southern Leyte: Contribution to the understanding of the evolution of Central Philippines, *NRCP Res. J.*, *10*, 39–54.
- Dow, D. B. (1977), *A Geological Synthesis of Papua New Guinea*, Geol. Geophys. Bull., vol. 201, 41 pp., Bureau of Mineral Resources, Australian Govt. Publ. Serv., Canberra.
- Dow, D. B., and R. Sukanto (1984), Western Irian Jaya: The end-product of oblique-plate convergence in the late Tertiary, *Tectonophysics*, *106*, 109–139.
- Ely, K. S., M. Sandiford, M. L. Hawke, D. Phillips, M. Quigley, and J. E. dos Reis (2011), Evolution of Ataúro Island: Temporal constraints on subduction processes beneath the Wetar zone, Banda Arc, *J. Asian Earth Sci.*, *41*, 477–493.
- Etourneau, J., P. Martinez, T. Blanz, and R. Schneider (2009), Pliocene-Pleistocene variability of upwelling activity, productivity, and nutrient cycling in the Benguela region, *Geology*, *37*, 871–874.
- Exon, N. F., and M. S. Marlow (1988), Geology and offshore resource potential of the New Ireland-Manus region—A synthesis, in *Geology and Offshore Resources of Pacific Island Arcs—New Ireland and Manus Region, Papua New Guinea*, Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, vol. 9, edited by M. S. Marlow, S. V. Dadisman, and N. F. Exon, pp. 241–262, Circum-Pacific Council for Energy and Mineral Resources, Houston, Tex.
- Faustino, D. V., G. P. Yumul Jr., J. V. De Jesus, C. B. Dimalanta, J. C. Aitchison, M.-F. Zhou, R. A. Tamayo Jr., and M. M. De Leon (2003), Geology of southeast Bohol, central Philippines: Accretion and sedimentation in a marginal basin, *Aust. J. Earth Sci.*, *50*, 571–583.
- Fedorov, A. V., and S. G. Philander (2001), A stability analysis of tropical ocean–atmosphere interactions: Bridging measurements and theory for El Niño, *J. Clim.*, *14*, 3086–3101.
- Fedorov, A. V., P. S. Dekens, M. McCarthy, A. C. Ravelo, P. B. deMenocal, M. Barreiro, R. C. Pacanowski, and S. G. Philander (2006), The Pliocene paradox (mechanisms for a permanent El Niño), *Science*, *312*, 1485–1489.
- Fedorov, A. V., C. M. Brierley, and K. Emanuel (2010), Tropical cyclones and permanent El Niño in the early Pliocene epoch, *Nature*, *463*, 1066–1070.
- Fedorov, A. V., C. M. Brierley, K. T. Lawrence, Z. Liu, P. S. Dekens, and A. C. Ravelo (2013), Patterns and mechanisms of early Pliocene warmth, *Nature*, *496*, 43–49.
- Ferdian, F., R. Hall, and I. Watkinson (2010), A structural re-evaluation of the North Banggai-Sula area, eastern Indonesia, *Proceeding Indonesian Petroleum Association, Thirty-Fourth Annual Convention and Exhibition*, 20 pp.
- Fitz, G., and P. Mann (2013), Tectonic uplift mechanism of the Goodenough and Fergusson Island gneiss domes, eastern Papua New Guinea: Constraints from seismic reflection and well data, *Geochem. Geophys. Geosyst.*, *14*, 3969–3995, doi:10.1002/ggge.20208.
- Ford, H. L., A. C. Ravelo, and S. Hovan (2012), A deep eastern equatorial Pacific thermocline during the early Pliocene warm period, *Earth Planet. Sci. Lett.*, *355–356*, 152–161.
- Fortuin, A. R., and M. E. M. De Smet (1991), Rates and magnitudes of late Cenozoic vertical movements in the Indonesian Banda Arc and the distinction of eustatic effects, in *Sedimentation, Tectonics and Eustasy - Sea-Level Changes at Active Margins*, Spec. Publ. Int. Ass. Sediment., vol. 12, edited by D. I. M. Macdonald, pp. 79–89, International Association of Sedimentologists, Gent, Belgium.
- Fortuin, A. R., M. E. M. De Smet, P. A. Sumosusastro, L. J. Van Marle, and S. R. Troelstra (1988), Late Cenozoic geohistory of NW Buru, Indonesia and plate tectonic implications, *Geol. Mijnbouw*, *67*, 91–105.
- Fortuin, A. R., M. E. M. De Smet, S. Hadiwasastro, L. J. Van Marle, S. R. Troelstra, and S. Tjokrosapoetro (1990), Late Cenozoic sedimentary and tectonic history of South Buton, Indonesia, *J. Southeast Asian Earth Sci.*, *4*, 107–124.
- Fortuin, A. R., T. B. Roep, and P. A. Sumosusastro (1994), The Neogene sediments of east Sumba, Indonesia products of a lost arc?, *J. Southeast Asian Earth Sci.*, *9*, 67–79.
- Fortuin, A. R., W. Van der Werff, and H. Wensink (1997), Neogene basin history and paleomagnetism of a rifted and inverted forearc region, on- and offshore Sumba, eastern Indonesia, *J. Asian Earth Sci.*, *15*, 61–88.
- Francis, G. (1988), Stratigraphy of Manus Island, western New Ireland Basin, Papua New Guinea, in *Geology and Offshore Resources of Pacific Island Arcs—New Ireland and Manus Region, Papua New Guinea*, Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, vol. 9, edited by M. S. Marlow, S. V. Dadisman, and N. F. Exon, pp. 31–40, Circum-Pacific Council for Energy and Mineral Resources, Houston, Tex.
- Franke, D., U. Barckhausen, I. Heyde, M. Tingay, and N. Ramli (2008), Seismic images of a collision zone offshore NW Sabah/Borneo, *Mar. Pet. Geol.*, *25*, 606–624.
- Froidveaux, C. M. (1978), Tertiary tectonic history of Salawati area, Irian Jaya, Indonesia, *AAPG Bull.*, *62*, 1127–1150.
- Gaillardet, J., B. Dupré, P. Louvat, and C. J. Allègre (1999), Global silicate weathering and CO₂ consumption rates deduced from the chemistry of large rivers, *Chem. Geol.*, *159*, 3–30.
- Gill, A. E. (1980), Some simple solutions for heat-induced tropical circulation, *Q. J. R. Meteorol. Soc.*, *106*, 447–462.
- Goldner, A., M. Huber, N. Diffenbaugh, and R. Caballero (2011), Implications of the permanent El Niño teleconnection “blueprint” for past global and North American hydroclimatology, *Clim. Past*, *7*, 723–743.
- Gordon, A. L., C. F. Giulivi, and A. G. Ilahude (2003), Deep topographic barriers within the Indonesian seas, *Deep Sea Res., Part II*, *50*, 2205–2228.
- Grainge, A. M., and K. G. Davies (1985), Reef exploration in the East Sengkang Basin, Sulawesi, Indonesia, *Mar. Pet. Geol.*, *2*, 142–155.
- Groeneveld, J., S. Steph, R. Tiedemann, D. Garbe-Schönberg, D. Nürnberg, and A. Sturm (2006), Pliocene mixed-layer oceanography for Site 1241, using combined Mg/Ca and $\delta^{18}\text{O}$ analyses of *Globigerinoides sacculifer*, in *Proc. ODP, Sci. Results*, vol. 202, edited by R. Tiedemann, et al., pp. 1–27, Ocean Drilling Program, College Station, Tex., doi:10.2973/odp.proc.sr.202.209.2006
- Haig, D. W. (2012), Palaeobathymetric gradients across Timor during 5.7–3.3 Ma (latest Miocene-Pliocene) and implications for collision uplift, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, *331–332*, 50–59.
- Hall, R. (1987), Plate boundary evolution in the Halmahera region, Indonesia, *Tectonophysics*, *144*, 337–352.
- Hall, R. (2000), Neogene history of collision in the Halmahera region, Indonesia, in *Proceedings of the Indonesian Petroleum Association 27th Annual Convention*, pp. 487–493, Indonesian Petroleum Association, Jakarta.
- Hall, R. (2001), Cenozoic reconstructions of SE Asia and the SW Pacific: Changing patterns of land and sea, in *Faunal and Floral Migrations and Evolution in SE Asia-Australasia*, edited by I. Metcalfe et al., pp. 35–56, A. A. Balkema (Swets & Zeitlinger Publishers), Lisse.
- Hall, R. (2002), Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: Computer-based reconstructions, model and animations, *J. Asian Earth Sci.*, *20*, 353–431.
- Hall, R. (2009), Southeast Asia’s changing palaeogeography, *Blumea*, *54*, 148–161.
- Hall, R. (2012), Sundaland and Wallacea: Geology, plate tectonics and palaeogeography, in *Biotic Evolution and Environmental Change in Southeast Asia*, The Systematics Association, edited by D. J. Gower et al., pp. 32–78, Cambridge Univ. Press, Cambridge.
- Hall, R. (2013), Contraction and extension in northern Borneo driven by subduction rollback, *J. Asian Earth Sci.*, *76*, 399–411.
- Hall, R., and G. J. Nichols (1990), Terrane amalgamation in the Philippine Sea margin, *Tectonophysics*, *181*, 207–222.

- Hall, R., and G. Nichols (2002), Cenozoic sedimentation and tectonics in Borneo: Climatic influences on orogenesis, in *Sediment Flux to Basins: Causes, Controls and Consequences*, Spec. Publ., vol. 111, edited by S. J. Jones and L. Frostick, pp. 5–22, Geol. Soc., London.
- Hall, R., M. G. Audley-Charles, F. T. Banner, S. Hidayat, and S. L. Tobing (1988a), Basement rocks of the Halmahera region, eastern Indonesia: A Late Cretaceous-early Tertiary arc and fore-arc, *J. Geol. Soc. London*, *145*, 65–84.
- Hall, R., M. G. Audley-Charles, F. T. Banner, S. Hidayat, and S. L. Tobing (1988b), Late Paleogene-Quaternary geology of Halmahera, eastern Indonesia: Initiation of a volcanic island arc, *J. Geol. Soc. London*, *145*, 577–590.
- Hall, R., G. Nichols, T. P. Ballantyne, T. Charlton, and J. Ali (1991), The character and significance of basement rocks of the southern Molucca Sea region, *J. Southeast Asian Earth Sci.*, *6*, 249–258.
- Hall, R., B. Clements, H. R. Smyth, and M. A. Cottam (2007), A new interpretation of Java's structure, Proceedings, Indonesian Petroleum Association, Thirty-First Annual Convention and Exhibition, Jakarta.
- Hall, R., M. W. A. van Hattum, and W. Spakman (2008), Impact of India-Asia collision on SE Asia: The record in Borneo, *Tectonophysics*, *451*, 366–389.
- Halpert, M. S., and C. F. Ropelewski (1992), Surface temperature patterns associated with the Southern Oscillation, *J. Clim.*, *5*, 577–593.
- Hamilton, W. (1979), *Tectonics of the Indonesian Region*, U. S. Geol. Surv. Prof. Pap., vol. 1078, 345 pp., U. S. Govt. Print. Off., Washington, D. C.
- Harris, R., M. W. Vorkink, C. Prasetyadi, E. Zobell, N. Roosmawati, and M. Aporthe (2009), Transition from subduction to arc-continent collision: Geologic and neotectonic evolution of Savu Island, Indonesia, *Geosphere*, *5*, 152–171.
- Haug, G. H., and R. Tiedemann (1998), Effect of the formation of the isthmus of Panama on Atlantic thermohaline circulation, *Nature*, *393*, 673–676.
- Haywood, A. M., et al. (2013), Large-scale features of Pliocene climate: Results from the Pliocene Model Intercomparison Project, *Clim. Past*, *9*, 191–209.
- Held, I. M., and B. J. Soden (2006), Robust responses of the hydrological cycle to global warming, *J. Clim.*, *19*, 5675–5699.
- Hennig, J., E. Advokaat, A. Rudyawan, and R. Hall (2014), Large sediment accumulations and major subsidence offshore; rapid uplift on land: Consequences of extension of Gorontalo Bay and northern Sulawesi, in *Proceeding Indonesian Petroleum Association, Thirty-Eighth Annual Convention & Exhibition*, 16 pp., Jakarta.
- Herbert, T. D., and J. D. Schuffert (1998), Alkenone unsaturation estimates of late Miocene through late Pliocene sea-surface temperatures at Site 958, in *Proc. Ocean Drill. Prog. Sci. Results.*, vol. 159T, edited by J. V. Firth, pp. 17–22, Ocean Drilling Program, College Station, Tex.
- Herrington, R. J., P. M. Scotney, S. Roberts, A. J. Boyce, and D. Harrison (2010), Temporal association of arc-continent collision, progressive magma contamination in arc volcanism and formation of gold-rich massive sulphide deposits on Wetar Island (Banda Arc), *Gondwana Res.*, *19*, 583–593.
- Hesse, S., S. Back, and D. Franke (2009), The deep-water fold-and-thrust belt offshore NW Borneo: Gravity-driven versus basement-driven shortening, *Geol. Soc. Am. Bull.*, *121*, 939–953.
- Hesse, S., S. Back, and D. Franke (2010), The structural evolution of folds in a deepwater fold and thrust belt—A case study from the Sabah continental margin offshore NW Borneo, SE Asia, *Mar. Pet. Geol.*, *27*, 442–454.
- Hill, K. C. (1991), Structure of the Papuan fold belt, Papua New Guinea, *Am. Assoc. Pet. Geol. Bull.*, *75*, 857–872.
- Hill, K. C., and A. J. W. Gleadow (1989), Uplift and thermal history of the Papuan Fold Belt, Papua New Guinea: Apatite fission track analysis, *Aust. J. Earth Sci.*, *36*, 515–539.
- Hill, K. C., and R. Hall (2003), Mesozoic-Cenozoic evolution of Australia's New Guinea margin in a West Pacific context, *Geol. Soc. Am. Spec. Pap.*, *372*, 265–289, Geol. Soc. Aust. Spec. Publ., 22.
- Hill, K. C., and A. Raza (1999), Arc-continent collision in Papua New Guinea: Constraints from fission track thermochronology, *Tectonics*, *18*, 950–966, doi:10.1029/1999TC900043.
- Hinz, K., J. Fritsch, E. H. K. Kempter, A. Manaaf Mohammad, H. Meyer, D. Mohamed, H. Vosberg, J. Weber, and J. J. Benavides (1989), Thrust tectonics along the continental margin of Sabah, Northwest Borneo, *Geol. Rundsch.*, *78*, 705–730.
- Hohnen, P. D. (1978), *Geology of New Ireland, Papua New Guinea*, Bulletin 194, PNG 12, 39 pp., Bureau of Mineral Resources, Geology and Geophysics (Australia), Australian Government Publishing Service, Canberra.
- Holland, G. J., and T. D. Keenan (1980), Diurnal variations of convection over the "maritime continent," *Mon. Weather Rev.*, *108*, 223–225.
- Holloway, N. H. (1982), The stratigraphy and tectonic relationship of Reed Bank, North Palawan and Mindoro to the Asian mainland and its significance in the evolution of the South China Sea, *AAPG Bull.*, *66*, 1355–1385.
- Horel, J. D., and J. M. Wallace (1981), Planetary-scale atmospheric phenomena associated with the Southern Oscillation, *Mon. Weather Rev.*, *109*(4), 813–829.
- Hughes, G. W., and C. C. Turner (1977), Upraised Pacific Ocean floor, southern Malaita, Solomon Islands, *Geol. Soc. Am. Bull.*, *88*, 412–424.
- Hutchison, C. S. (1996), The "Rajang accretionary prism" and "Lupar Line" problem of Borneo, in *Tectonic Evolution of Southeast Asia*, Spec. Publ., vol. 106, edited by R. Hall and D. J. Blundell, pp. 247–261, Geol. Soc., London.
- Huybers, P., and P. Molnar (2007), Tropical cooling and the onset of North American glaciation, *Clim. Past*, *3*, 549–557.
- Jochum, M., B. Fox-Kemper, P. H. Molnar, and C. Shields (2009), Differences in the Indonesian seaway in a coupled climate model and their relevance to Pliocene climate and El Niño, *Paleoceanography*, *24*, PA1212, doi:10.1029/2008PA001678.
- John, T. U. (1963), Geology and mineral deposits of east-central Balabac Island, Palawan Province, Philippines, *Econ. Geol.*, *58*, 107–130.
- Johnston, C. R., and C. O. Bowin (1981), Crustal reactions resulting from the mid-Pliocene to recent continent-island arc collision in the Timor region, *BMR J. Aust. Geol. Geophys.*, *6*, 223–243.
- Kalnay, E., et al. (1996), The NCEP/NCAR 40-Year Reanalysis Project, *Bull. Am. Meteorol. Soc.*, *77*, 437–472.
- Kaneps, A. G. (1979), Gulf stream: Velocity fluctuations during the late Cenozoic, *Science*, *204*, 297–301.
- Karas, C., D. Nürnberg, A. K. Gupta, R. Tiedemann, K. Mohan, and T. Bickert (2009), Mid-Pliocene climate change amplified by a switch in Indonesian subsurface throughflow, *Nat. Geosci.*, *2*, 434–438.
- Karas, C., D. Nürnberg, R. Tiedemann, and D. Garbe-Schönberg (2011), Pliocene Indonesian Throughflow and Leeuwin Current dynamics: Implications for Indian Ocean polar heat flux, *Paleoceanography*, *26*, PA2217, doi:10.1029/2010PA001949.
- Keenan, T. D., B. R. Morton, M. J. Manton, and G. J. Holland (1989), The island thunderstorm experiment (ITEX)—A study of tropical thunderstorms in the maritime continent, *Bull. Am. Meteorol. Soc.*, *70*, 152–159.
- Keenan, T. D., et al. (2000), The Maritime Continent Thunderstorm Experiment (MCTEX): Over view and some results, *Bull. Am. Meteorol. Soc.*, *81*, 2433–2455.
- Keep, M., and D. W. Haig (2010), Deformation and exhumation in Timor: Distinct stages of a young orogeny, *Tectonophysics*, *483*, 93–111.
- Keigwin, L. D., Jr. (1982), Isotope paleoceanography of the Caribbean and East Pacific: Role of Panama uplift in late Neogene time, *Science*, *217*, 350–353.

- Kent, D. V., and G. Muttoni (2013), Modulation of Late Cretaceous and Cenozoic climate by variable drawdown of atmospheric pCO₂ from weathering of basaltic provinces on continents drifting through the equatorial humid belt, *Clim. Past*, *9*, 525–546.
- Kiladis, G. N., and H. F. Diaz (1986), An analysis of the 1877–1878 ENSO episode and comparison with 1982–83, *Mon. Weather Rev.*, *114*, 1035–1047.
- Kiladis, G. N., and H. F. Diaz (1989), Global climatic anomalies associated with extremes in the Southern Oscillation, *J. Clim.*, *2*, 1069–1090.
- Krijgsman, W., F. J. Hilgen, I. Raffi, F. J. Sierro, and D. S. Wilson (1999), Chronology, causes and progression of the Messinian salinity crisis, *Nature*, *400*, 652–655.
- Kump, L. R., S. L. Brantley, and M. A. Arthur (2000), Chemical weathering, atmospheric CO₂, and climate, *Annu. Rev. Earth Planet. Sci.*, *28*, 611–667.
- Kürschner, W. M., J. van der Burgh, H. Visscher, and D. L. Dilcher (1996), Oak leaves as biosensors of late Neogene and early Pleistocene paleoatmospheric CO₂ concentrations, *Mar. Micropaleontol.*, *27*, 299–312.
- Labat, D., Y. Goddérès, J.-L. Probst, and J.-L. Guyot (2004), Evidence for global runoff increase related to climate warming, *Adv. Water Resour.*, *27*, 631–642.
- Larsen, I. J., P. C. Almond, A. Eger, J. O. Stone, D. R. Montgomery, and B. Malcolm (2014), Rapid soil production in the Southern Alps, New Zealand, *Science*, *343*, 637–640.
- Lawrence, K. T., Z.-h. Liu, and T. D. Herbert (2006), Evolution of the eastern tropical Pacific through Plio-Pleistocene glaciation, *Science*, *312*, 79–83.
- Lawrence, K. T., T. D. Herbert, C. M. Brown, M. E. Raymo, and A. M. Haywood (2009), High-amplitude variations in North Atlantic sea surface temperature during the early Pliocene warm period, *Paleoceanography*, *24*, PA2218, doi:10.1029/2008PA001669.
- Lear, C. H., H. Elderfield, and P. A. Wilson (2000), Cenozoic deep-sea temperatures and global ice volumes from Mg/Ca in benthic foraminiferal calcite, *Science*, *287*, 269–272.
- Leduc, G., D. Garbe-Schönberg, M. Regenberg, C. Contoux, J. Etourneau, and R. Schneider (2014), The late Pliocene Benguela upwelling status revisited by means of multiple temperature proxies, *Geochem. Geophys. Geosyst.*, *15*, 475–491, doi:10.1002/2013GC004940.
- Legates, D. R., H. F. Lins, and G. J. McCabe (2005), Comments on “Evidence for global runoff increase related to climate warming” by Labat et al., *Adv. Water Resour.*, *28*, 1310–1315.
- Liberti, G. L., F. Cheruy, and M. Desbois (2001), Land effect on the diurnal cycle of clouds over the TOGA COARE area, as observed from GMS IR data, *Mon. Weather Rev.*, *129*, 1500–1517.
- Lindley, I. D. (2006), Extensional and vertical tectonics in the New Guinea islands: Implications for island arc evolution, *Ann. Geophys.*, *49*, 403–426.
- Linthout, K., and H. Helmers (1994), Pliocene obducted, rotated and migrated ultramafic rocks and obduction-induced anatectic granite, SW Seram and Ambon, Eastern Indonesia, *J. Southeast Asian Earth Sci.*, *9*, 95–109.
- Linthout, K., H. Helmers, J. R. Wijbrans, and J. D. A. M. Van Wees (1996), ⁴⁰Ar/³⁹Ar constraints on obduction of the Seram ultramafic complex: Consequences for the evolution of the southern Banda Sea, in *Tectonic Evolution of Southeast Asia*, Spec. Publ., vol. 106, edited by R. Hall and D. J. Blundell, pp. 455–464, Geol. Soc., London.
- Linthout, K., H. Helmers, and J. Sopaheluwakan (1997), Late Miocene obduction and microplate migration around the southern Banda Sea and the closure of the Indonesian seaway, *Tectonophysics*, *281*, 17–30.
- Lisiecki, L. E., and M. E. Raymo (2005), A Pliocene-Pleistocene stack of 57 globally distributed benthic δ¹⁸O records, *Paleoceanography*, *20*, PA1003, doi:10.1029/2004PA001071.
- Little, T. A., B. R. Hacker, S. M. Gordon, S. L. Baldwin, P. G. Fitzgerald, S. Ellis, and M. Korchinski (2011), Diapiric exhumation of the world’s youngest (UHP) eclogites in the gneiss domes of the D’Entrecasteaux Islands Papua New Guinea, *Tectonophysics*, *510*, 39–68.
- Lunt, D. J., A. M. Haywood, G. A. Schmidt, U. Salzmann, P. J. Valdes, and H. J. Dowsett (2010), Earth system sensitivity inferred from Pliocene modelling and data, *Nat. Geosci.*, *3*, 60–64.
- Lunt, P., G. Burgon, and A. Baky (2009), The Pemali Formation of Central Java and equivalents: Indicators of sedimentation on an active plate margin, *J. Asian Earth Sci.*, *34*, 100–113.
- Macpherson, C. G., E. J. Forde, R. Hall, and M. F. Thirlwall (2003), Geochemical evolution of magmatism in an arc-arc collision: The Halmahera and Sangihe arcs, eastern Indonesia, in *Intra-Oceanic Subduction Systems: Tectonic and Magmatic Processes*, Special Publications, vol. 219, edited by R. D. Larter and P. T. Leat, pp. 207–220, Geological Society, London.
- Malaihollo, J. F. A., and R. Hall (1996), The geology and tectonic evolution of the Bacan region, east Indonesia, in *Tectonic Evolution of Southeast Asia*, Spec. Publ., vol. 106, edited by R. Hall and D. J. Blundell, pp. 483–497, Geol. Soc., London.
- Mann, P., and T. Asahiko (2004), Global tectonic significance of the Solomon Islands and Ontong Java Plateau convergent zone, *Tectonophysics*, *389*, 137–190.
- Marlow, J. R., C. B. Lange, G. Wefer, and A. Rosell-Melé (2000), Upwelling intensification as a part of the Pliocene-Pleistocene climate transition, *Science*, *290*, 2288–2291.
- Mayall, M. J., and M. Cox (1988), Deposition and diagenesis of Miocene limestones, Senkang Basin, Sulawesi, Indonesia, *Sediment. Geol.*, *59*, 77–92.
- McClay, K., T. Dooley, A. Ferguson, and J. Poblet (2000), Tectonic evolution of the Sanga Sanga block, Mahakam delta, Kalimantan, Indonesia, *Am. Assoc. Pet. Geol. Bull.*, *84*, 765–786.
- McDowell, F. W., T. P. McMahon, P. Q. Warren, and M. Cloos (1996), Pliocene Cu-Au-bearing igneous intrusions of the Guning Bijih (Ertsberg) District, Irian Jaya, Indonesia: K-Ar geochronology, *J. Geol.*, *104*, 327–340.
- Merlis, T. M., and T. Schneider (2011), Changes in zonal surface temperature gradients and Walker Circulations in a wide range of climates, *J. Clim.*, *24*, 4757–4768.
- Molnar, P., and M. A. Cane (2002), El Niño’s tropical climate and teleconnections as a blueprint for pre-Ice Age climates, *Paleoceanography*, *17*(2), 1021, doi:10.1029/2001PA000663.
- Molnar, P., and M. A. Cane (2007), Early Pliocene (pre-Ice Age) El Niño-like global climate: Which El Niño?, *Geosphere*, *3*(5), 337–365.
- Monnier, C., J. Girardeau, M. Pubellier, M. Polvé, H. Permana, and H. Bellon (1999), Petrology and geochemistry of the Cyclops ophiolites (Irian Jaya, East Indonesia): Consequences for the Cenozoic evolution of the north Australian margin, *Mineral. Petrol.*, *65*, 1–28.
- Moore, G. F., and D. E. Karig (1980), Structural geology of Nias Island, Indonesia: Implications for subduction zone tectonics, *Am. J. Sci.*, *280*, 193–223.
- Moore, G. F., H. G. Billman, P. E. Hehanussa, and D. E. Karig (1980), Sedimentology and paleobathymetry of Neogene trench-slope deposits, Nias Island, Indonesia, *J. Geol.*, *88*, 161–180.
- Moore, G. F., C. A. Evans, J. W. Hawkins, and K. Kadarisman (1981), Geology of the Talaud Islands, Molucca Sea collision zone, Northeast Indonesia, *J. Struct. Geol.*, *3*, 467–475.

- Morley, C. K. (2009), Geometry of an oblique thrust fault zone in a deepwater fold belt from 3D seismic data, *J. Struct. Geol.*, *31*, 1540–1555.
- Morley, C. K., S. Back, P. van Rensbergen, P. Crevello, and J. J. Lambiase (2003), Characteristics of repeated, detached, Miocene-Pliocene tectonic inversion events, in a large delta province on an active margin, Brunei Darussalam, Borneo, *J. Struct. Geol.*, *25*, 1147–1169.
- Moss, S. J., and J. L. C. Chambers (1999), Tertiary facies architecture in the Kutai Basin, Kalimantan, Indonesia, *J. Asian Earth Sci.*, *17*, 157–181.
- Moss, S. J., J. Chambers, I. Cloke, D. Satria, J. R. Ali, S. Baker, J. Milsom, and A. Carter (1997), New observations on the sedimentary and tectonic evolution of the Tertiary Kutai Basin, East Kalimantan, in *Petroleum Geology of Southeast Asia*, edited by A. J. Fraser, S. J. Matthews, and R. W. Murphy, *Geol. Soc. London Spec. Publ.*, *126*, 395–416.
- Mudelsee, M., and M. E. Raymo (2005), Slow dynamics of the Northern Hemisphere glaciation, *Paleoceanography*, *20*, PA4022, doi:10.1029/2005PA001153.
- Müller, C. (1991), Biostratigraphy and geological evolution of the Sulu Sea and surrounding area, in *Proceedings of the Ocean Drilling Program*, Scientific Results, vol. 124, edited by E. A. Silver, C. Rangin, and M. T. von Breyman, pp. 121–131, Ocean Drilling Program, College Station, Tex.
- Nguyen, N., B. Duffy, J. Shulmeister, and M. Quigley (2013), Rapid Pliocene uplift of Timor, *Geology*, *41*, 179–182.
- Nichols, G. J., and R. Hall (1991), Basin formation and Neogene sedimentation in a back arc setting, Halmahera, eastern Indonesia, *Mar. Pet. Geol.*, *8*, 50–61.
- O'Brien, C. L., G. L. Foster, M. A. Martínez-Botí, R. Abell, J. W. B. Rae, and R. D. Pancost (2014), High sea surface temperatures in tropical warm pools during the Pliocene, *Nat. Geosci.*, *7*, 606–610.
- Oliva, P., J. Viers, and B. Dupré (2003), Chemical weathering in granitic environments, *Chem. Geol.*, *202*, 225–256.
- Pagani, M., Z.-h. Liu, J. LaRiviere, and A. C. Ravelo (2010), High Earth-system climate sensitivity determined from Pliocene carbon dioxide concentrations, *Nat. Geosci.*, *3*, 27–30.
- Pairault, A. A., R. Hall, and C. F. Elders (2003a), Tectonic evolution of the Seram Trough, Indonesia, in *Proceedings, Indonesian Petroleum Association, Twenty-Ninth Annual Convention and Exhibition, October 2003*, pp. 355–370, Indonesian Petroleum Association, Jakarta.
- Pairault, A. A., R. Hall, and C. F. Elders (2003b), Structural styles and tectonic evolution of the Seram Trough, Indonesia, *Mar. Pet. Geol.*, *20*(10), 1141–1160.
- Parsons, B. (1982), Causes and consequences of the relation between area and age of the ocean floor, *J. Geophys. Res.*, *87*, 437–448.
- Pearson, P. N., and M. R. Palmer (2000), Atmospheric carbon dioxide concentrations over the past 60 million years, *Nature*, *406*, 695–699.
- Philander, S. G., and A. V. Fedorov (2003), Role of tropics in changing the response to Milankovich forcing some three million years ago, *Paleoceanography*, *18*(2), 1045, doi:10.1029/2002PA000837.
- Phinney, E. J., P. Mann, M. F. Coffin, and T. H. Shipley (2004), Sequence stratigraphy, structural style, and age of deformation of the Malaita accretionary prism (Solomon Arc-Ontong Java Plateau convergent zone), *Tectonophysics*, *389*, 221–246.
- Pierrehumbert, R. T. (1995), Thermostats, radiator fins and local runaway greenhouse, *J. Atmos. Sci.*, *52*, 405–416.
- Pieters, P. E., C. J. Pigram, D. S. Trail, D. B. Dow, N. Ratman, and R. Sukanto (1983), The stratigraphy of western Irian Jaya, *Bull. Geol. Res. Dev. Cent., Bandung*, *8*, 14–48.
- Pigram, C. J., and P. A. Symonds (1991), A review of the major tectonic events in the New Guinea orogen, *J. Southeast Asian Earth Sci.*, *6*, 307–318.
- Pownall, J. M., R. Hall, and I. M. Watkinson (2013), Extreme extension across Seram and Ambon, eastern Indonesia: Evidence for Banda slab rollback, *Solid Earth*, *4*, 277–314.
- Pownall, J. M., R. Hall, R. A. Armstrong, and M. A. Forster (2014), Earth's youngest known ultrahigh-temperature granulites discovered on Seram, eastern Indonesia, *Geology*, *42*, 279–282.
- Pubellier, M., and F. Ego (2002), Anatomy of an escape tectonic zone: Western Irian Jaya (Indonesia), *Tectonics*, *21*(4), 1019, doi:10.1029/2001TC901038.
- Pubellier, M., R. Quebral, C. Rangin, B. Deffontaine, C. Muller, and J. Butterlin (1991), The Mindanao collision zone: A soft collision event within a continuous Neogene strike-slip setting, *J. Southeast Asian Earth Sci.*, *6*, 239–248.
- Pubellier, M., R. Quebral, M. Aurelio, and C. Rangin (1996), Docking and post-docking escape tectonics in the southern Philippines, in *Tectonic Evolution of Southeast Asia*, Spec. Publ., vol. 106, edited by R. Hall and D. J. Blundell, pp. 511–523, Geol. Soc., London.
- Qian, J.-H. (2008), Why precipitation is mostly concentrated over islands in the Maritime Continent, *J. Atmos. Sci.*, *65*(4), 1428–1441.
- Qian, J.-H., A. W. Robertson, and V. Moron (2013), Diurnal cycle in different weather regimes and rainfall variability over Borneo associated with ENSO, *J. Clim.*, *26*, 1772–1790.
- Quarles van Ufford, A., and M. Cloos (2005), Cenozoic tectonics of New Guinea, *Am. Assoc. Pet. Geol. Bull.*, *89*, 119–140.
- Queaño, K. L. (2005), Upper Miocene to lower Pliocene Sigaboy Formation turbidites, on the Pujada Peninsula, Mindanao, Philippines: Internal structures, composition, depositional elements and reservoir characteristics, *J. Asian Earth Sci.*, *25*, 387–402.
- Quebral, R., M. Pubellier, and C. Rangin (1996), The onset of movement on the Philippine Fault in eastern Mindanao: A transition from a collision to strike slip environment, *Tectonics*, *15*, 713–726, doi:10.1029/95TC00480.
- Quigley, M. C., B. Duffy, J. Woodhead, J. Hellstrom, L. Moody, T. Horton, J. Soares, and L. Fernandes (2012), U/Pb dating of a terminal Pliocene coral from the Indonesian seaway, *Mar. Geol.*, *311–314*, 57–62.
- Ramaswamy, V., O. Boucher, J. Haigh, D. Haughlustaine, J. Haywood, G. Myhre, T. Nakajima, G. Y. Shi, and S. Solomon (2001), Radiative forcing of climate change, in *Climate Change 2001: The Scientific Basis, Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, edited by J. T. Houghton et al., 349–416, Cambridge Univ. Press, Cambridge, U. K., and New York.
- Rangin, C. (1989), The Sulu Sea, a marginal basin setting within a collision zone, *Tectonophysics*, *161*, 119–141.
- Rangin, C., and E. A. Silver (1991), Neogene tectonic evolution of the Celebes-Sulu Basins: New insights from Leg 124 drilling, in *Proceedings of the Ocean Drilling Program*, Scientific Results, vol. 124, edited by E. A. Silver, C. Rangin, and M. T. von Breyman, pp. 121–131, Ocean Drilling Program, College Station, Tex.
- Rangin, C., H. Bellon, F. Benard, J. Letouzey, C. Muller, and S. Tahir (1990), Neogene arc-continent collision in Sabah, N. Borneo (Malaysia), *Tectonophysics*, *183*, 305–319.
- Rangin, C., X. Le Pichon, S. Mazzotti, M. Pubellier, N. Chamot-Rooke, M. Aurelio, A. Walpersdorf, and R. Quebral (1999), Plate convergence measured by GPS across the Sundaland/Philippine Sea plate deformed boundary: The Philippines and eastern Indonesia, *Geophys. J. Int.*, *139*, 296–316.
- Rasmusson, E. M., and J. M. Wallace (1983), Meteorological aspects of the El Niño/Southern Oscillation, *Science*, *222*, 1195–1202.
- Ravelo, A. C., and M. W. Wara (2004), Role of the tropical oceans on global climate during a warm period and a major climate transition, *Oceanography*, *17*, 32–41.
- Ravelo, A. C., D. H. Andreasen, M. Lyle, A. Olivarez Lyle, and M. W. Wara (2004), Regional climate shifts caused by gradual global cooling in the Pliocene epoch, *Nature*, *429*, 263–267.

- Ravelo, A. C., P. S. Dekens, and M. McCarthy (2006), Evidence for El Niño-like conditions during the Pliocene, *GSA Today*, *16*(3), 4–11.
- Raymo, M. E., B. Grant, M. Horowitz, and G. H. Rau (1996), Mid-Pliocene warmth: Stronger greenhouse and stronger conveyor, *Mar. Micropaleontol.*, *27*, 313–326.
- Rigg, J. W. D., and R. Hall (2011), Structural and stratigraphic evolution of the Savu Basin, Indonesia, in *The SE Asian Gateway: History and Tectonics of the Australia-Asia Collision*, Special Publications, vol. 355, edited by R. Hall, M. A. Cottam, and M. E. J. Wilson, pp. 225–240, Geological Society, London.
- Riker-Coleman, K. E., C. D. Gallup, L. M. Wallace, J. M. Webster, H. Cheng, and R. L. Edwards (2006), Evidence of Holocene uplift in east New Britain, Papua New Guinea, *Geophys. Res. Lett.*, *33*, L18612, doi:10.1029/2006GL026596.
- Rodgers, K. B., M. Latif, and S. Legutke (2000), Sensitivity of equatorial Pacific and Indian Ocean watermasses to the position of the Indonesian throughflow, *Geophys. Res. Lett.*, *27*, 2941–2944, doi:10.1029/1999GL002372.
- Roosmawati, N., and R. Harris (2009), Surface uplift history of the incipient Banda arc-continent collision: Geology and synorogenic foraminifera of Rote and Savu Islands, Indonesia, *Tectonophysics*, *479*, 95–110.
- Ropelewski, C. F., and M. S. Halpert (1986), North American precipitation and temperature patterns associated with the El Niño/Southern Oscillation, *Mon. Weather Rev.*, *114*, 2352–2362.
- Rosell-Melé, A., A. Martínez-García, and E. L. McClymont (2014), Persistent warmth across the Benguela upwelling system during the Pliocene epoch, *Earth Planet. Sci. Lett.*, *386*, 10–20.
- Rowley, D. B. (2002), Rate of plate creation and destruction: 180 Ma to present, *Geol. Soc. Am. Bull.*, *114*, 927–933.
- Rutherford, E., K. Burke, and J. Lytvyn (2001), Tectonic history of Sumba Island Indonesia, since the Late Cretaceous and its rapid escape into the forearc in the Miocene, *J. Asian Earth Sci.*, *19*, 453–479.
- Saint-Marc, P., and Suminta (1979), Biostratigraphy of late Miocene and Pliocene deep water sediments of eastern Java, Indonesia, *J. Foraminiferal Res.*, *9*, 106–117.
- Saito, K., T. Keenan, G. Holland, and K. Puri (2001), Numerical simulation of the diurnal evolution of tropical island convection over the Maritime Continent, *Mon. Weather Rev.*, *129*, 378–400.
- Sajona, F. G., H. Bellon, R. C. Maury, M. Pubellier, J. Cotton, and C. Rangin (1994), Magmatic response to abrupt changes in geodynamic settings: Pliocene-Quaternary calc-alkaline and Nb-enriched lavas from Mindanao (Philippines), *Tectonophysics*, *237*, 47–72.
- Sajona, F. G., H. Bellon, R. C. Maury, M. Pubellier, R. D. Quebral, J. Cotten, F. E. Bayon, E. Pagadog, and P. Pamatian (1997), Tertiary and Quaternary magmatism in Mindanao and Leyte (Philippines): Geochronology, geochemistry and tectonic setting, *J. Asian Earth Sci.*, *15*, 121–153.
- Sajona, F. G., R. C. Maury, M. Pubellier, J. Leterrier, H. Bellon, and J. Cotten (2000), Magmatic source enrichment by slab-derived melts in a young post-collision setting, central Mindanao (Philippines), *Lithos*, *54*, 173–206.
- Samuel, M. A., N. A. Harbury, M. E. Jones, and S. J. Matthews (1995), Inversion of an outer-arc ridge: The Sumatran Forearc, Indonesia, in *Basin Inversion*, Special Publication, vol. 88, edited by J. H. Buchanan and P. G. Buchanan, pp. 473–492, Geological Society, London.
- Samuel, M. A., N. A. Harbury, A. Bakri, F. T. Banner, and L. Hartono (1997), A new stratigraphy for the islands of the Sumatran Forearc, Indonesia, *J. Asian Earth Sci.*, *15*, 339–380.
- Sapiie, B., and M. Cloos (2004), Strike-slip faulting in the core of the Central Range of west New Guinea: Ertsberg Mining District, Indonesia, *Geol. Soc. Am. Bull.*, *116*, 277–293.
- Sapin, F., M. Pubellier, J.-C. Ringenbach, and V. Bailly (2009), Alternating thin vs. thick-skinned decollements, example in a fast tectonic setting: The Misool-Onin-Kumawa Ridge (west Papua), *J. Struct. Geol.*, *31*, 444–459.
- Sarnthein, M., G. Bartoli, M. Prange, A. Schmittner, B. Schneider, M. Weinelt, N. Andersen, and D. Garbe-Schönberg (2009), Mid-Pliocene shifts in ocean overturning circulation and the onset of Quaternary-style climates, *Clim. Past*, *5*, 269–283.
- Sato, T., H. Miura, M. Satoh, Y. N. Takayabu, and Y.-q. Wang (2009), Diurnal cycle of precipitation in the tropics simulated in a global cloud-resolving model, *J. Clim.*, *22*(18), 4809–4826.
- Satyana, A. H., D. Nugroho, and I. Surantoko (1999), Tectonic controls on the hydrocarbon habitats of the Barito, Kutei and Tarakan Basins, Eastern Kalimantan, Indonesia: Major dissimilarities in adjoining basins, *J. Asian Earth Sci.*, *17*, 99–122.
- Schopka, H. H., L. A. Derry, and C. A. Arcilla (2011), Chemical weathering, river geochemistry and atmospheric carbon fluxes from volcanic and ultramafic regions on Luzon Island, the Philippines, *Geochim. Cosmochim. Acta*, *75*, 978–1002.
- Seki, O., G. L. Foster, D. N. Schmidt, A. Mackensen, K. Kawamura, and R. D. Pancost (2010), Alkenone and boron based Pliocene $p\text{CO}_2$ records, *Earth Planet. Sci. Lett.*, *292*, 201–211.
- Shukla, S. P., M. A. Chandler, J. Jonas, L. E. Sohl, K. Mankoff, and H. Dowsett (2009), Impact of a permanent El Niño (El Padre) and Indian Ocean Dipole in warm Pliocene climates, *Paleoceanography*, *24*, PA2221, doi:10.1029/2008PA001682.
- Shukla, S. P., M. A. Chandler, D. Rind, L. E. Sohl, J. Jonas, and J. Lerner (2011), Teleconnections in a warmer climate: The Pliocene perspective, *Clim. Dyn.*, *37*, 1869–1887.
- Silver, E. A., R. McCaffrey, Y. Joyodiwiro, and S. Stevens (1983), Ophiolite emplacement by collision between the Sula Platform and the Sulawesi island arc, Indonesia, *J. Geophys. Res.*, *88*, 9419–9435, doi:10.1029/JB088iB11p09419.
- Simons, W. J. F., et al. (2007), A decade of GPS in Southeast Asia: Resolving Sundaland motion and boundaries, *J. Geophys. Res.*, *112*, B06420, doi:10.1029/2005JB003868.
- Smith, I. E., and H. L. Davies (1976), *Geology of the Southeast Papuan Mainland*, Bull., vol. 165, 86 pp., Bureau of Mineral Resources, Geology and Geophysics, Australian Govt. Publ. Serv., Canberra.
- Smith, R. B., and E. A. Silver (1991), Geology of a Miocene collision complex, Buton, eastern Indonesia, *Geol. Soc. Am. Bull.*, *103*, 660–678.
- Sobel, A. H., C. D. Burleyson, and S. E. Yuter (2011), Rain on small tropical islands, *J. Geophys. Res.*, *116*, D08102, doi:10.1029/2010JD014695.
- Socquet, A., W. Simons, C. Vigny, R. McCaffrey, C. Subarya, D. Sarsito, B. Ambrosius, and W. Spakman (2006), Microblock rotations and fault coupling in SE Asia triple junction (Sulawesi, Indonesia) from GPS and earthquake slip vector data, *J. Geophys. Res.*, *111*, B08409, doi:10.1029/2005JB003963.
- Soden, B. J., and I. M. Held (2006), An assessment of climate feedbacks in coupled atmosphere-ocean models, *J. Clim.*, *19*, 3354–3360.
- Stewart, W. D., and M. J. Sandy (1988), Geology of New Ireland and Djaul Islands, northeastern Papua New Guinea, in *Geology and Offshore Resources of Pacific Island Arcs—New Ireland and Manus Region, Papua New Guinea*, Circum-Pacific Council for Energy and Mineral Resources Earth Science Series, vol. 9, edited by M. S. Marlow, S. V. Dadisman, and N. F. Exon, pp. 13–30, Circum-Pacific Council for Energy and Mineral Resources, Houston, Tex.
- Sudarmono (2000), Tectonic and Stratigraphic evolution of the Bone Basin, Indonesia: Insights to the Sulawesi collision complex, in *Proceedings Indonesian Petroleum Association, 27th Annual Convention*, pp. 531–543, Indonesian Petroleum Association, Jakarta.
- Travaglio, C., A. F. Baes, and L. M. Tomas (1978), *Geology of Samar Island, Soil and Land Resources Appraisal and Trading Project Philippines, AGO/PHI/74/003 Field Document 1, Annex 6*, vol. 1, 149 pp., Bureau of Soils, Department of Agriculture, United Nations Development Programme, Food and Agriculture Administration of the United Nations, Manila.

- Trenberth, K. E., J. M. Caron, D. P. Stepaniak, and S. Worley (2002), Evolution of El Niño–Southern Oscillation and global atmospheric surface temperatures, *J. Geophys. Res.*, *107*(D8), 4065, doi:10.1029/2000JD000298.
- Tripati, A. K., C. D. Roberts, and R. A. Eagle (2009), Coupling of CO₂ and ice sheet stability over major climate transitions of the last 20 million years, *Science*, *326*, 1394–1397.
- Umbgrove, J. H. F. (1946), Corals from a Lower Pliocene patch reef in Central Java, *J. Paleontol.*, *20*, 521–542.
- van de Weerd, A., and R. A. Armin (1992), Origin and evolution of the Tertiary hydrocarbon bearing basins in Kalimantan (Borneo), Indonesia, *Am. Assoc. Pet. Geol. Bull.*, *76*, 1778–1803.
- Van Gorsel, J. T., and S. R. Troelstra (1981), Late Neogene planktonic foraminiferal biostratigraphy and climatostratigraphy of the Solo River section (Java, Indonesia), *Mar. Micropaleontol.*, *6*, 183–209.
- van Leeuwen, T. M., and Muhandjo (2005), Stratigraphy and tectonic setting of the Cretaceous and Paleogene volcanic-sedimentary successions in northwest Sulawesi, Indonesia: Implications for the Cenozoic evolution of western and northern Sulawesi, *J. Asian Earth Sci.*, *25*, 481–511.
- van Leeuwen, T. M., E. S. Susanto, S. Maryanto, S. Hadiwisastra, Sudijono, Muhandjo, and Prihardjo (2010), Tectonostratigraphic evolution of Cenozoic marginal basin and continental margin successions in the Bone Mountains, southwest Sulawesi, Indonesia, *J. Asian Earth Sci.*, *38*, 233–254.
- van Marle, L. J. (1991), Late Cenozoic palaeobathymetry and geohistory analysis of central west Timor, eastern Indonesia, *Mar. Pet. Geol.*, *8*, 22–34.
- van Marle, L. J., and M. E. M. De Smet (1990), Note on the late Cenozoic history of the Kai Islands, Indonesia, *Geol. Mijnbouw*, *69*, 93–103.
- Vecchi, G. A., and B. J. Soden (2007), Global warming and the weakening of the tropical circulation, *J. Clim.*, *20*, 4316–4340.
- Veevers, J. J., D. A. Falvey, and S. Robins (1978), Timor Trough and Australia: Facies show topographic wave migrated 80 km during the past 3 m.y., *Tectonophysics*, *45*, 217–227.
- Villeneuve, M., J.-P. Réhault, J.-J. Cornée, C. Honthaas, W. Gunawan, et le groupe Géobanda (1998), Évolution géodynamique de l'Indonésie orientale, de l'Éocène au Pliocène, *C. R. Acad. Sci., Ser. IIa Sci. Terre Planètes*, *327*, 291–302.
- Villeneuve, M., J.-J. Cornée, W. Gunawan, M.-C. Janin, J. Butterlin, P. Saint-Marc, and H. Samodra (2000), Continental block collision in the eastern arm of Sulawesi (Indonesia). Structure and geodynamic interpretation, *C. R. Acad. Sci., Ser. IIa Sci. Terre Planètes*, *330*, 371–378.
- Villeneuve, M., J.-J. Cornée, W. Gunawan, R. Martini, G. Tronchetti, M.-C. Jamin, P. Saint-Marc, and L. Zaninetti (2001), La succession lithostratigraphique du bloc de Banda dans la région de Kolonodale (Sulawesi central, Indonésie), *Bull. Soc. Geol. Fr.*, *172*, 59–68.
- Vizcaíno, M., S. Rupper, and J. C. H. Chiang (2010), Permanent El Niño and the onset of Northern Hemisphere glaciations: Mechanism and comparison with other hypotheses, *Paleoceanography*, *25*, PA2205, doi:10.1029/2009PA001733.
- von der Heydt, A. S., A. Nnafie, and H. A. Dijkstra (2011), Cold tongue/warm pool and ENSO dynamics in the Pliocene, *Clim. Past*, *7*, 903–915.
- Walker, J. C. G., P. B. Hays, and J. F. Kasting (1981), The negative feedback mechanism for long-term stabilization of Earth's surface temperature, *J. Geophys. Res.*, *86*, 9776–9782, doi:10.1029/JC086iC10p09776.
- Wannier, M. (2009), Carbonate platforms in wedge-top basins: An example from the Gunung Mulu National Park, Northern Sarawak (Malaysia), *Mar. Pet. Geol.*, *26*, 177–207.
- Wara, M. W., A. C. Ravelo, and M. L. Delaney (2005), Permanent El Niño-like conditions during the Pliocene warm period, *Science*, *309*, 758–761.
- Watkinson, I. M., R. Hall, and F. Ferdian (2011), Tectonic re-interpretation of the Banggai-Sula-Molucca Sea margin, Indonesia, in *The SE Asian Gateway: History and Tectonics of Australia-Asia Collision*, Spec. Publ., vol. 355, edited by R. Hall, M. A. Cottam, and M. E. J. Wilson, pp. 203–224, Geol. Soc., London.
- Weiland, R. J., and M. Cloos (1996), Pliocene–Pleistocene asymmetric unroofing of the Irian fold belt, Irian Jaya, Indonesia: Apatite fission-track thermochronology, *Geol. Soc. Am. Bull.*, *108*, 1438–1449.
- Weyl, P. K. (1968), The role of the oceans in climatic change: A theory of the Ice Ages, *Meteorol. Monogr.*, *8*, 37–62.
- Wilson, M. E. J. (2008), Global and regional influences on equatorial shallow-marine carbonates during the Cenozoic, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, *255*, 262–274.
- Wilson, M. E. J., and S. J. Moss (1999), Cenozoic palaeogeographic evolution of Sulawesi and Borneo, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, *145*, 303–337.
- Witts, D., R. Hall, R. J. Morley, and M. K. BouDagher-Fadel (2011), Stratigraphy and sediment provenance, Barito Basin, southeast Kalimantan, Proceeding Indonesian Petroleum Association, Thirty-Fifth Annual Convention and Exhibition, 18 pp.
- Yulianto, I., R. Hall, B. Clements, and C. R. Elders (2007), Structural and stratigraphic evolution of the offshore Malingping Block, West Java, Indonesia, Proceedings, Indonesian Petroleum Association, Thirty-First Annual Convention and Exhibition.
- Yumul, G. P., Jr., C. B. Dimalanta, E. J. Marquez, and K. L. Queaño (2009), Onland signatures of the Palawan microcontinental block and Philippine mobile belt collision and crustal growth process: A review, *J. Asian Earth Sci.*, *34*, 610–623.
- Zachos, J. C., M. Pagani, L. Sloan, E. Thomas, and K. Billups (2001), Trends, rhythms, and aberrations in global climate 65 Ma to Present, *Science*, *292*, 686–693.
- Zebiak, S. E., and M. A. Cane (1987), A model El Niño–Southern Oscillation, *Mon. Weather Rev.*, *115*, 2262–2278.
- Zhang, X., Q. Tang, X. Zhang, and D. P. Lettenmaier (2014), Runoff sensitivity to global mean temperature change in the CMIP5 Models, *Geophys. Res. Lett.*, *41*, 5492–5498, doi:10.1002/2014GL060382.
- Zhang, Y. G., M. Pagani, and Z.-h. Liu (2014), A 12-million-year temperature history of the tropical Pacific Ocean, *Science*, *344*, 84–87.
- Zhou, L., and Y.-q. Wang (2006), Tropical Rainfall Measuring Mission observation and regional model study of precipitation diurnal cycle in the New Guinean region, *J. Geophys. Res.*, *111*, D17104, doi:10.1029/2006JD007243.