

# **Growth stages of mono-and dicotyledonous plants**

**BBCH Monograph**

2. Edition, 2001

Edited by Uwe Meier

**Federal Biological Research Centre for Agriculture and Forestry**

The code has been jointly by

- German Federal Biological Research Centre for Agriculture and Forestry (BBA)
- German Federal Office of Plant Varieties (BSA)
- German Agrochemical Association (IVA)
- Institute for Vegetables and Ornamentals in Grossbeeren/Erfurt, Germany

### **Members of the BBCH working group**

H. Bleiholder und Frau E.  
Weber  
BASF AG  
Landwirtschaftliche Ver-  
suchsstation  
Carl-Bosch-Strasse 64  
D-67117 Limburgerhof

C. Feller  
Institut für Gemüse & Zier-  
pflanzenbau  
Theodor-Echtermeyer-Weg 1  
D-14979 Grossbeeren

M. Hess und H. Wicke  
Aventis  
D-65926 Frankfurt/Main

U. Meier  
Biologische Bundesanstalt  
für Land- und Forstwirtschaft  
Messeweg 11/12  
D-38104 Braunschweig

T. van den Boom  
Bayer AG  
Landwirtschaftszentrum  
Monheim  
Alfred-Nobel-Strasse 50  
D-51368 Leverkusen-  
Bayerwerk

P. D. Lancashire  
Bayer plc.  
Eastern Way  
Bury St. Edmunds  
Suffolk IP 32 7 AH, UK

Frau L. Buhr  
Biologische Bundesanstalt  
für Land- und Forstwirtschaft  
Stahnsdorfer Damm 81  
D-14532 Kleinmachnow

H. Hack  
Industrieverband Agrar (IVA)  
Theodor-Storm-Weg 2  
D-51519 Odenthal

Frau R. Klose  
Bundessortenamt  
Osterfelddamm 80  
D-30604 Hannover

R. Stauss  
Ministerium für ländliche  
Räume, Landwirtschaft,  
Ernährung und Tourismus  
des Landes Schleswig-  
Holstein  
Düsternbrooker Weg 104  
D-24105 Kiel

**General Scale**

---

**Cereals, Rice, Maize**

---

**Oilseed rape, Faba bean, Sunflower**

---

**Beta beets**

---

**Potato**

---

**Fruits**

---

**Citrus, Olive, Coffee, Banana**

---

**Grapevine**

---

**Soybean, Cotton, Peanuts**

---

**Hop**

---

**Vegetable crops I**

---

**Vegetable crops II**

---

**Weeds**

---

**BBCH-Publications**

## Foreword

As all branches of science, the individual disciplines in agricultural plant research also work more closely together, and, in addition, have become more international. The exchange of new findings and joint work on projects presuppose, however, that all those involved have the same understanding of the terms they use. This calls the need for an extensively standardised description of plant development stages in order of their phenological characteristics and their coding.

The phenological development stages of plants are also used in agricultural practice, agrometeorology and agricultural insurance, each with its own varying individual objectives. Moreover, the applied botanical sciences also make use of phenological development stages.

This book about plant development stages – and their corresponding codes – aims at satisfying all these demands. Of particular significance is the fact that the work appears in four languages and thus contributes to a large extent to reducing linguistic communication problems. It thus fulfils in a special way the intertwinement in research, trade, production and service present today.

The book thanks its existence to the close co-operation between scientists from variously oriented agricultural authorities and companies from the chemical industry with agricultural research departments. It is hoped that this fruitful co-operation, in the course of which knowledge has been gathered by all sides over many years, will contribute to furnishing decision makers with more surety, and will promote international co-operation.

Prof. Dr. F. Klingauf

President of the  
Federal Biological Research Centre  
for Agriculture and Forestry,  
Berlin and Braunschweig

## **Note of Thanks**

This book was made possible by a joint initiative of the Institute for Vegetables and Ornamentals in Großbeeren/Erfurt (IGZ), the German Federal Office of Plant Varieties (BSA), the concerns Aventis, Bayer, BASF and Syngenta, the German Agrochemical Association (IVA) and the German Federal Biological Research Centre for Agriculture and Forestry (BBA). Not only have these institutions and concerns provided substantial funds, but also scientific and technical employees, whose joint work over several years has made this book possible. Indispensable however has been the work of numerous cooperating scientists and co-authors who have enabled the description of the development stages of the crops with their crop specific knowledge. I would like to thank them at this point for their help.

An essential aim of this book is to facilitate scientific communication on an international level. Fortunately I was able to find specialist translators. The Spanish translations were done by Mr. Enrique Gonzales Medina, Bogotá/Kolumbien, Mr. José Antonio Guerra, Ciba, Barcelona, as well as Dr. Herrman Bleiholder, BASF AG, who earns the credit for unifying versions in the Spanish language from South America and Spain. The French text was translated by Mrs. Sybil Rometsch, Université de Lausanne. The English text was corrected by Dr. P. D. Lancashire, Bayer, Bury St. Edmunds, UK.

Mr. Ernst Halwaß from Nossen agreed as commercial artist to produce the main part of the graphical representations, enabling the clear visual portrayal of the important development stages of the particular crops. Thanks go to him for the creation of drawings of all vegetable and fruit plants, and those of the beet, potato, cotton, peanut, hop, faba bean, sunflower, maize, soybean, musacea, coffe, olive and grapevine. Thanks also to the Gesellschaft der Freunde und Förderer der Biologischen Bundesanstalt for financing the pictures and Mr. Tottman and Mrs. Broad for their drawings of cereals.

The authors would also like to thank those colleagues who carried out the necessary technical work with so much patience and perseverance.

The production of a book in this form requires substantial financial means. We would therefore like to thank the following companies and institutions at this point for the financial resources provided: Aventis, BASF, Bayer, Syngenta, Dow Elanco, Du Pont de Nemours, Rhône-Poulenc Agro, and the German Agrochemical Association (IVA).

Uwe Meier

# The extended BBCH-scale

Hack et al., 1992

---

The extended BBCH-scale is a system for a uniform coding of phenologically similar growth stages of all mono- and dicotyledonous plant species.

It results from teamwork between the German Federal Biological Research Centre for Agriculture and Forestry (BBA), the German Federal Office of Plant Varieties (BSA), the German Agrochemical Association (IVA) and the Institute for Vegetables and Ornamentals in Grossbeeren/Erfurt, Germany (IGZ). The decimal code, which is divided into principal and secondary growth stages, is based on the well-known cereal code developed by ZADOKS et al. (1974) in order to avoid major changes from this widely used phenological key. The abbreviation **BBCH** derives from **B**iologische **B**undesanstalt, **B**undessortenamt and **C**hemical industry.

## The basic principles of the scale

- The general scale forms the framework within which the individual scales are developed. It can also be used for those plant species for which no special scale is currently available.
- Similar phenological stages of each plant species are given the same code.
- For each code, a description is given, and for some important stages, drawings are included.
- For the description of the phenological development stages, clear and easily recognised (external) morphological characteristics are used.
- Except where stated otherwise, only the development of the main stem is taken into consideration.
- The growth stages refer to representative individual plants within the crop stand. Crop stand characteristics may also be considered.
- Relative values relating to species- and/or variety-specific ultimate sizes are used for the indication of sizes.
- The secondary growth stages 0 to 8 correspond to the respective ordinal numbers or percentage values. For example stage 3 could represent: 3rd true leaf, 3rd tiller, 3rd node or 30% of the final length or size typical of the species or 30% of the flowers open.
- Post harvest or storage treatment is coded **99**.
- Seed treatment before planting is coded **00**.

## Organisation of the scale

The entire developmental cycle of the plants is subdivided into ten clearly recognizable and distinguishable longer-lasting developmental phases. These **principal growth stages** are described using numbers from 0 to 9 in ascending order (see Figures 1a and b). The principal growth stages are described in Table 1. Owing to the very many different plant species there may be shifts in the course of the development or certain stages may even be omitted.

The principal growth stages need not proceed in the strict sequence defined by the ascending order of the figures, but can occasionally also proceed in parallel.

**Table 1:**

Principal growth stages

Stage	Description
0	Germination / sprouting / bud development
1	Leaf development (main shoot)
2	Formation of side shoots / tillering
3	Stem elongation or rosette growth / shoot development (main shoot)
4	Development of harvestable vegetative plant parts or vegetatively propagated organs / booting (main shoot)
5	Inflorescence emergence (main shoot) / heading
6	Flowering (main shoot)
7	Development of fruit
8	Ripening or maturity of fruit and seed
9	Senescence, beginning of dormancy

If two or more principal growth stages proceed in parallel, both can be indicated by using a diagonal stroke (example 16/22). If only one stage is to be indicated, either the more advanced growth stage must be chosen or the principal growth stage of particular interest, depending upon the plant species.

The principal growth stages alone are not sufficient to define exactly application or evaluation dates, since they always describe time spans in the course of the development of a plant.

**Secondary stages** are used if **points of time** or steps in the plant development must be indicated precisely. In contrast to the principal growth stages they are defined as short developmental steps characteristic of the respective plant species, which are passed successively during the respective principal growth stage. They are also coded by using the figures 0 to 9. The combination of figures for the principal and the secondary stages, results in the two-digit code.

The two-digit code is a scale which offers the possibility of precisely defining all phenological growth stages for the majority of plant species.

Only in the case of some plant species (e.g. cucumber, onion, potato, soybean, tomato) is further subdivision necessary within a principal growth stage beyond that possible using the secondary stages from 0 to 9.

For these cases a three-digit scale is presented alongside the two-digit scale. This involves the inclusion of the so-called **mesostage** between the principal and the secondary stage, which provides a further subdivision with figures **0** and **1** describing the development on the **main stem** and figures **2** to **9**

that of the side shoots **2nd to 9th order** (see Figures 1a and b). In this way up to 19 leaves can be counted on the main stem or the branching can be described.

The BBCH-scales allow the comparison of individual codes only within one principal growth stage: an arithmetically greater code indicates a plant at a later growth stage. Sorting codes into numerical order therefore allows a listing in order of the stage of plant development.

The time span of certain developmental phases of a plant can be exactly defined and coded by indicating two stages. For this purpose two codes are connected with a hyphen. Thus, for instance, the code 51–69 describes the developmental phase from the appearance of the first inflorescence or flower buds until the end of flowering. This allows the computer-supported monitoring of crop stands.

For a uniform coding which covers the maximum number of plant species, it is necessary to use primarily phenological criteria rather than homologous or analogous stages. Thus, for instance, germination of plants from true seed and sprouting from buds are classified in one principal growth stage, the principal growth stage 0, even though they are completely different biological processes.


In case of the BBCH-scales the descriptions are based on the actual characteristic features of the individual plant. If the scales are used for the definition of the development stage of a plant stand, the description should apply to at least 50% of the plants.

Greater differences in the course of the development of different plant groups have to be taken into consideration for the description of the general scale (see 1.2). This problem is dealt with by offering several definitions for one specific stage wherever the formulation of a uniform text is impossible. The following letters show to which plant group the respective definition refers.


### **Figures 1a and b:**

Subdivision of the developmental cycle of plants into principal and secondary stages (a) and into principal, meso- and secondary stages (b). The mesostages are inserted between the principal and the secondary stages. Modified according to a draft by A. Witzemberger.





**a**



**b**

# The extended BBCH-scale, general Hack et al., 1992

**D** = Dicotyledons

**M** = Monocotyledons

**V** = Development from vegetative parts or propagated organs.

**G** = Gramineae

**P** = Perennial plants

No code letter is used if the description applies to all groups of plants.

---

Code	Description
------	-------------

---

## Principal growth stage 0: Germination, sprouting, bud development

00		Dry seed (seed dressing takes place at stage 00)
	P, V	Winter dormancy or resting period
01		Beginning of seed imbibition;
	P, V	Beginning of bud swelling
03		Seed imbibition complete;
	P, V	End of bud swelling
05		Radicle (root) emerged from seed;
	P, V	Perennating organs forming roots
06		Elongation of radicle, formation of root hairs and/or lateral roots
07	G	Coleoptile emerged from caryopsis;
	D, M	Hypocotyl with cotyledons or shoot breaking through seed coat;
	P, V	Beginning of sprouting or bud breaking
08	D	Hypocotyl with cotyledons growing towards soil surface;
	P, V	Shoot growing towards soil surface
09	G	Emergence: Coleoptile breaks through soil surface;
	D, M	Emergence: Cotyledons break through soil surface (except hypogeal germination);
	D, V	Emergence: Shoot/leaf breaks through soil surface;
	P	Bud shows green tips

---

## Principal growth stage 1: Leaf development (main shoot)

10	G	First true leaf emerged from coleoptile;
	D, M	Cotyledons completely unfolded;
	P	First leaves separated
11		First true leaf, leaf pair or whorl unfolded;
	P	First leaves unfolded
12		2 true leaves, leaf pairs or whorls unfolded
13		3 true leaves, leaf pairs or whorls unfolded
1 .		Stages continuous till . . .
19		9 or more true leaves, leaf pairs or whorls unfolded

---

# The extended BBCH-scale, general Hack et al., 1992

Code	Description
<b>Principal growth stage 2: Formation of side shoots/tillering</b>	
21	First side shoot visible;
G	First tiller visible
22	2 side shoots visible;
G	2 tillers visible
23	3 side shoots visible;
G	3 tillers visible
2 .	Stages continuous till . . .
29	9 or more side shoots visible;
G	9 or more tillers visible
<b>Principal growth stage 3: Stem elongation or rosette growth, shoot development (main shoot)</b>	
31	Stem (rosette) 10% of final length (diameter);
G	1 node detectable
32	Stem (rosette) 20% of final length (diameter);
G	2 nodes detectable
33	Stem (rosette) 30% of final length (diameter);
G	3 nodes detectable
3 .	Stages continuous till . . .
39	Maximum stem length or rosette diameter reached;
G	9 or more nodes detectable
<b>Principal growth stage 4: Development of harvestable vegetative plant parts or vegetatively propagated organs/booting (main shoot)</b>	
40	Harvestable vegetative plant parts or vegetatively propagated organs begin to develop
41	G Flag leaf sheath extending
43	Harvestable vegetative plant parts or vegetatively propagated organs have reached 30% of final size;
G	Flag leaf sheath just visibly swollen (mid-boot)
45	Harvestable vegetative plant parts or vegetatively propagated organs have reached 50% of final size;
G	Flag leaf sheath swollen (late-boot)
47	Harvestable vegetative plant parts or vegetatively propagated organs have reached 70% of final size;
G	Flag leaf sheath opening
49	Harvestable vegetative plant parts or vegetatively propagated organs have reached final size;
G	First awns visible

# The extended BBCH-scale, general Hack et al., 1992

---

Code	Description
------	-------------

---

**Principal growth stage 5: Inflorescence emergence (main shoot) / heading**

51	Inflorescence or flower buds visible;
G	Beginning of heading
55	First individual flowers visible (still closed);
G	Half of inflorescence emerged (middle of heading)
59	First flower petals visible (in petalled forms);
G	Inflorescence fully emerged (end of heading)

---

**Principal growth stage 6: Flowering (main shoot)**

60	First flowers open (sporadically)
61	Beginning of flowering: 10% of flowers open
62	20% of flowers open
63	30% of flowers open
64	40% of flowers open
65	Full flowering: 50% of flowers open, first petals may be fallen
67	Flowering finishing: majority of petals fallen or dry
69	End of flowering: fruit set visible

---

**Principal growth stage 7: Development of fruit**

71	10% of fruits have reached final size or fruit has reached 10% of final size <sup>1</sup>
G	Caryopsis watery ripe
72	20% of fruits have reached final size or fruit has reached 20% of final size <sup>1</sup>
73	30% of fruits have reached final size or fruit has reached 30% of final size <sup>1</sup>
G	Early milk
74	40% of fruits have reached final size or fruit has reached 40% of final size <sup>1</sup>
75	50% of fruits have reached final size or fruit has reached 50% of final size <sup>1</sup>
G	Milky ripe, medium milk
76	60% of fruits have reached final size or fruit has reached 60% of final size <sup>1</sup>
77	70% of fruits have reached final size or fruit has reached 70% of final size <sup>1</sup>
G	Late milk
78	80% of fruits have reached final size or fruit has reached 80% of final size <sup>1</sup>
79	Nearly all fruits have reached final size <sup>1</sup>

---

<sup>1</sup> This stage is not used, if the main fruit growth happens in principal growth stage 8

# The extended BBCH-scale, general Hack et al., 1992

---

Code	Description
<b>Principal growth stage 8: Ripening or maturity of fruit and seed</b>	
81	Beginning of ripening or fruit colouration
85	Advanced ripening or fruit colouration;
G	Dough stage
87	Fruit begins to soften (species with fleshy fruit)
89	Fully ripe: fruit shows fully-ripe colour, beginning of fruit abscission
<b>Principal growth stage 9: Senescence, beginning of dormancy</b>	
91	P Shoot development completed, foliage still green
93	Beginning of leaf-fall
95	50% of leaves fallen
97	End of leaf fall, plants or above ground parts dead or dormant;
P	Plant resting or dormant
99	Harvested product (post-harvest or storage treatment is applied at stage 99)

---

# The extended BBCH-scale, for specific crops

---

## Cereals Witzenberger et al., 1989; Lancashire et al., 1991

### Phenological growth stages and BBCH-identification keys of cereals

(wheat = *Triticum* sp. L., barley = *Hordeum vulgare* L., oat = *Avena sativa* L., rye = *Secale cereale* L.)

---

Code	Description
------	-------------

---

#### Principal growth stage 0: Germination

00	Dry seed (caryopsis)
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from caryopsis
06	Radicle elongated, root hairs and/or side roots visible
07	Coleoptile emerged from caryopsis
09	Emergence: coleoptile penetrates soil surface (cracking stage)

---

#### Principal growth stage 1: Leaf development<sup>1,2</sup>

10	First leaf through coleoptile
11	First leaf unfolded
12	2 leaves unfolded
13	3 leaves unfolded
1 .	Stages continuous till . . .
19	9 or more leaves unfolded

---

#### Principal growth stage 2: Tillering<sup>3</sup>

20	No tillers
21	Beginning of tillering: first tiller detectable
22	2 tillers detectable
23	3 tillers detectable
2 .	Stages continuous till . . .
29	End of tillering. Maximum no. of tillers detectable

---

<sup>1</sup> A leaf is unfolded when its ligule is visible or the tip of the next leaf is visible

<sup>2</sup> Tillering or stem elongation may occur earlier than stage 13; in this case continue with stages 21

<sup>3</sup> If stem elongation begins before the end of tillering continue with stage 30

# Cereals

Witzenberger et al., 1989; Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of cereals

---

Code	Description
<b>Principal growth stage 3: Stem elongation</b>	
30	Beginning of stem elongation: pseudostem and tillers erect, first internode begins to elongate, top of inflorescence at least 1 cm above tillering node
31	First node at least 1 cm above tillering node
32	Node 2 at least 2 cm above node 1
33	Node 3 at least 2 cm above node 2
3 .	Stages continuous till . . .
37	Flag leaf just visible, still rolled
39	Flag leaf stage: flag leaf fully unrolled, ligule just visible

---

### Principal growth stage 4: Booting

41	Early boot stage: flag leaf sheath extending
43	Mid boot stage: flag leaf sheath just visibly swollen
45	Late boot stage: flag leaf sheath swollen
47	Flag leaf sheath opening
49	First awns visible (in awned forms only)

---

### Principal growth stage 5: Inflorescence emergence, heading

51	Beginning of heading: tip of inflorescence emerged from sheath, first spikelet just visible
52	20% of inflorescence emerged
53	30% of inflorescence emerged
54	40% of inflorescence emerged
55	Middle of heading: half of inflorescence emerged
56	60% of inflorescence emerged
57	70% of inflorescence emerged
58	80% of inflorescence emerged
59	End of heading: inflorescence fully emerged

---

### Principal growth stage 6: Flowering, anthesis

61	Beginning of flowering: first anthers visible
65	Full flowering: 50% of anthers mature
69	End of flowering: all spikelets have completed flowering but some dehydrated anthers may remain

---

### Principal growth stage 7: Development of fruit

71	Watery ripe: first grains have reached half their final size
73	Early milk
75	Medium milk: grain content milky, grains reached final size, still green
77	Late milk

---

# Cereals

Witzenberger et al., 1989; Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of cereals

---

Code	Description
------	-------------

---

### Principal growth stage 8: Ripening


- 83 Early dough
  - 85 Soft dough: grain content soft but dry. Fingernail impression not held
  - 87 Hard dough: grain content solid. Fingernail impression held
  - 89 Fully ripe: grain hard, difficult to divide with thumbnail
- 

### Principal growth stage 9: Senescence



- 92 Over-ripe: grain very hard, cannot be dented by thumbnail
  - 93 Grains loosening in day-time
  - 97 Plant dead and collapsing
  - 99 Harvested product
-



# Cereals



# Cereals



# Rice

Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of rice (*Oryza sativa* L.)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Germination

00	Dry seed (caryopsis)
01	Beginning of seed imbibition
03	Seed imbibition complete (pigeon breast)
05	Radicle emerged from caryopsis
06	Radicle elongated, root hairs and/or side roots visible
07	Coleoptile emerged from caryopsis (in water-rice this stage occurs before stage 05)
09	Imperfect leaf emerges (still rolled) at the tip of the coleoptile

---

### Principal growth stage 1: Leaf development<sup>1, 2</sup>

10	Imperfect leaf unrolled, tip of first true leaf visible
11	First leaf unfolded
12	2 leaves unfolded
13	3 leaves unfolded
1 .	Stages continuous till . . .
19	9 or more leaves unfolded

---

### Principal growth stage 2: Tillering<sup>3</sup>

21	Beginning of tillering: first tiller detectable
22	2 tillers detectable
23	3 tillers detectable
2 .	Stages continuous till ...
29	Maximum number of tillers detectable

---

### Principal growth stage 3: Stem elongation

30	Panicle initiation or green ring stage: chlorophyll accumulates in the stem tissue, forming a green ring
32	Panicle formation: panicle 1–2 mm in length
34	Internode elongation or jointing stage: internodes begin to elongate, panicle more than 2 mm long (variety-dependent)
37	Flag leaf just visible, still rolled, panicle moving upwards
39	Flag leaf stage: flag leaf unfolded, collar regions (auricle and ligule) of flag leaf and penultimate leaf aligned (pre-boot stage)

---

<sup>1</sup> A leaf is unfolded when its ligule is visible or the tip of the next leaf is visible

<sup>2</sup> Tillering or stem elongation may occur earlier than stage 13; in this case continue with stages 21 or 30

<sup>3</sup> If stem elongation begins before the end of tillering continue with stage 30

# Rice

Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of rice

Code	Description
<b>Principal growth stage 4: Booting</b>	
41	Early boot stage: upper part of stem slightly thickened, sheath of flag leaf about 5 cm out of penultimate leaf sheath
43	Mid boot stage: sheath of flag leaf 5–10 cm out of the penultimate leaf sheath
45	Late boot stage: flag leaf sheath swollen, sheath of flag leaf more than 10 cm out of penultimate leaf sheath
47	Flag leaf sheath opening
49	Flag leaf sheath open
<b>Principal growth stage 5: Inflorescence emergence, heading<sup>4</sup></b>	
51	Beginning of panicle emergence: tip of inflorescence emerged from sheath
52	20% of panicle emerged
53	30% of panicle emerged
54	40% of panicle emerged
55	Middle of panicle emergence: neck node still in sheath
56	60% of panicle emerged
57	70% of panicle emerged
58	80% of panicle emerged
59	End of panicle emergence: neck node level with the flag leaf auricle, anthers not yet visible
<b>Principal growth stage 6: Flowering, anthesis</b>	
61	Beginning of flowering: anthers visible at top of panicle
65	Full flowering: anthers visible on most spikelets
69	End of flowering: all spikelets have completed flowering but some dehydrated anthers may remain
<b>Principal growth stage 7: Development of fruit</b>	
71	Watery ripe: first grains have reached half their final size
73	Early milk
75	Medium milk: grain content milky
77	Late milk
<b>Principal growth stage 8: Ripening</b>	
83	Early dough
85	Soft dough: grain content soft but dry, fingernail impression not held, grains and glumes still green
87	Hard dough: grain content solid, fingernail impression held
89	Fully ripe: grain hard, difficult to divide with thumbnail

<sup>4</sup> Flowering usually starts before stage 55; continue with principal stage 6

# Rice

Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of rice

---

Code	Description
------	-------------





---

### Principal growth stage 9: Senescence




92	Over-ripe: grain very hard, cannot be dented by thumbnail
97	Plant dead and collapsing
99	Harvested product

---

# Rice



# Rice



# Maize

Weber and Bleiholder, 1990; Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of maize

(*Zea mays* L.)

---

Code	Description
------	-------------

---

**Principal growth stage 0: Germination**

00	Dry seed (caryopsis)
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from caryopsis
06	Radicle elongated, root hairs and/or side roots visible
07	Coleoptile emerged from caryopsis
09	Emergence: coleoptile penetrates soil surface (cracking stage)

---

**Principal growth stage 1: Leaf development<sup>1, 2</sup>**

10	First leaf through coleoptile
11	First leaf unfolded
12	2 leaves unfolded
13	3 leaves unfolded
1 .	Stages continuous till . . .
19	9 or more leaves unfolded

---

**Principal growth stage 3: Stem elongation**

30	Beginning of stem elongation
31	First node detectable
32	2 nodes detectable
33	3 nodes detectable
3 .	Stages continuous till . . .
39	9 or more nodes detectable <sup>3</sup>

---

**Principal growth stage 5: Inflorescence emergence, heading**

51	Beginning of tassel emergence: tassel detectable at top of stem
53	Tip of tassel visible
55	Middle of tassel emergence: middle of tassel begins to separate
59	End of tassel emergence: tassel fully emerged and separated

---

<sup>1</sup> A leaf may be described as unfolded when its ligule is visible or the tip of next leaf is visible

<sup>2</sup> Tillering or stem elongation may occur earlier than stage 19; in this case continue with principal growth stage 3

<sup>3</sup> In maize, tassel emergence may occur earlier, in this case continue with principal growth stage 5



# Maize

Weber and Bleiholder, 1990; Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of maize

---

Code	Description
------	-------------

---

### Principal growth stage 6: Flowering, anthesis

61	Male: stamens in middle of tassel visible Female: tip of ear emerging from leaf sheath
63	Male: beginning of pollen shedding Female: tips of stigmata visible
65	Male: upper and lower parts of tassel in flower Female: stigmata fully emerged
67	Male: flowering completed Female: stigmata drying
69	End of flowering: stigmata completely dry

---

### Principal growth stage 7: Development of fruit

71	Beginning of grain development: kernels at blister stage, about 16% dry matter
73	Early milk
75	Kernels in middle of cob yellowish-white (variety-dependent), content milky, about 40% dry matter
79	Nearly all kernels have reached final size

---

### Principal growth stage 8: Ripening

83	Early dough: kernel content soft, about 45% dry matter
85	Dough stage: kernels yellowish to yellow (variety dependent), about 55% dry matter
87	Physiological maturity: black dot/layer visible at base of kernels, about 60% dry matter
89	Fully ripe: kernels hard and shiny, about 65% dry matter



---

### Principal growth stage 9: Senescence

97	Plant dead and collapsing
99	Harvested product

---

# Maize



# Maize




63



69



79



89

# Oilseed rape Weber and Bleiholder, 1990; Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of oilseed rape

(*Brassica napus* L. ssp. *napus*)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Germination

00	Dry seed
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from seed
07	Hypocotyl with cotyledons emerged from seed
08	Hypocotyl with cotyledons growing towards soil surface
09	Emergence: cotyledons emerge through soil surface

---

### Principal growth stage 1: Leaf development<sup>1</sup>

10	Cotyledons completely unfolded
11	First leaf unfolded
12	2 leaves unfolded
13	3 leaves unfolded
1 .	Stages continuous till . . .
19	9 or more leaves unfolded

---

### Principal growth stage 2: Formation of side shoots

20	No side shoots
21	Beginning of side shoot development: first side shoot detectable
22	2 side shoots detectable
23	3 side shoots detectable
2 .	Stages continuous till . . .
29	End of side shoot development: 9 or more side shoots detectable

---

### Principal growth stage 3: Stem elongation<sup>2</sup>

30	Beginning of stem elongation: no internodes ("rosette")
31	1 visibly extended internode
32	2 visibly extended internodes
33	3 visibly extended internodes
3 .	Stages continuous till . . .
39	9 or more visibly extended internodes

---

<sup>1</sup> Stem elongation may occur earlier than stage stage 19; in this case continue with stage 20

<sup>2</sup> Visibly extended internode n develops between leaf n and leaf n+1

# Oilseed rape Weber and Bleiholder, 1990; Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of oilseed rape

---

Code	Description
------	-------------

---

### Principal growth stage 5: Inflorescence emergence

50	Flower buds present, still enclosed by leaves
51	Flower buds visible from above ("green bud")
52	Flower buds free, level with the youngest leaves
53	Flower buds raised above the youngest leaves
55	Individual flower buds (main inflorescence) visible but still closed
57	Individual flower buds (secondary inflorescences) visible but still closed
59	First petals visible, flower buds still closed ("yellow bud")

---

### Principal growth stage 6: Flowering

60	First flowers open
61	10% of flowers on main raceme open, main raceme elongating
62	20% of flowers on main raceme open
63	30% of flowers on main raceme open
64	40% of flowers on main raceme open
65	Full flowering: 50% flowers on main raceme open, older petals falling
67	Flowering declining: majority of petals fallen
69	End of flowering

---

### Principal growth stage 7: Development of fruit

71	10% of pods have reached final size
72	20% of pods have reached final size
73	30% of pods have reached final size
74	40% of pods have reached final size
75	50% of pods have reached final size
76	60% of pods have reached final size
77	70% of pods have reached final size
78	80% of pods have reached final size
79	Nearly all pods have reached final size

---

# Oilseed rape Weber and Bleiholder, 1990; Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of oilseed rape

---

Code	Description
------	-------------

---

### Principal growth stage 8: Ripening

80	Beginning of ripening: seed green, filling pod cavity
81	10% of pods ripe, seeds dark and hard
82	20% of pods ripe, seeds dark and hard
83	30% of pods ripe, seeds dark and hard
84	40% of pods ripe, seeds dark and hard
85	50% of pods ripe, seeds dark and hard
86	60% of pods ripe, seeds dark and hard
87	70% of pods ripe, seeds dark and hard
88	80% of pods ripe, seeds dark and hard
89	Fully ripe: nearly all pods ripe, seeds dark and hard

---

### Principal growth stage 9: Senescence


97	Plant dead and dry
99	Harvested product

---


# Oilseed rape



10



11



12



13



18



32




51 (Detail)



51


# Oilseed rape




53



53 (Detail)



55




57




61



67



69



79



# Faba bean Weber and Bleiholder, 1990; Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of faba bean

(*Vicia faba* L.)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Germination

00	Dry seed
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from seed
07	Shoot emerged from seed (plumule apparent)
08	Shoot growing towards soil surface
09	Emergence: shoot emerges through soil surface

---

### Principal growth stage 1: Leaf development<sup>1</sup>

10	Pair of scale leaves visible (may be eaten or lost)
11	First leaf unfolded
12	2 leaves unfolded
13	3 leaves unfolded
1 .	Stages continuous till . . .
19	9 or more leaves unfolded

---

### Principal growth stage 2: Formation of side shoots

20	No side shoots
21	Beginning of side shoot development: first side shoot detectable
22	2 side shoots detectable
23	3 side shoots detectable
2 .	Stages continuous till . . .
29	End of side shoot development: 9 or more side shoots detectable

---

### Principal growth stage 3: Stem elongation

30	Beginning of stem elongation
31	One visibly extended internode <sup>2</sup>
32	2 visibly extended internodes
33	3 visibly extended internodes
3 .	Stages continuous till . . .
39	9 or more visibly extended internodes

---

<sup>1</sup> Stem elongation may occur earlier than stage 19; in this case continue with the principal stage 3

<sup>2</sup> First internode extends from the scale leaf node to the first true leaf node

# Faba bean Weber and Bleiholder, 1990; Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of faba bean

---

Code	Description
------	-------------

---

### Principal growth stage 5: Inflorescence emergence

50	Flower buds present, still enclosed by leaves
51	First flower buds visible outside leaves
55	First individual flower buds visible outside leaves but still closed
59	First petals visible, many individual flower buds, still closed

---

### Principal growth stage 6: Flowering

60	First flowers open
61	Flowers open on first raceme
63	Flowers open 3 racemes per plant
65	Full flowering: flowers open on 5 racemes per plant
67	Flowering declining
69	End of flowering

---

### Principal growth stage 7: Development of fruit

70	First pods have reached final length ("flat pod")
71	10% of pods have reached final length
72	20% of pods have reached final length
73	30% of pods have reached final length
74	40% of pods have reached final length
75	50% of pods have reached final length
76	60% of pods have reached final length
77	70% of pods have reached final length
78	80% of pods have reached final length
79	Nearly all pods have reached final length

---

### Principal growth stage 8: Ripening

80	Beginning of ripening: seed green, filling pod cavity
81	10% of pods ripe, seeds dry and hard
82	20% of pods ripe, seeds dry and hard
83	30% of pods ripe and dark, seeds dry and hard
84	40% of pods ripe and dark, seeds dry and hard
85	50% of pods ripe and dark, seeds dry and hard
86	60% of pods ripe and dark, seeds dry and hard
87	70% of pods ripe and dark, seeds dry and hard
88	80% of pods ripe and dark, seeds dry and hard
89	Fully ripe: nearly all pods dark, seeds dry and hard

---

# **Faba bean** Weber and Bleiholder, 1990; Lancashire et al., 1991

## **Phenological growth stages and BBCH-identification keys of faba bean**

---

Code	Description
------	-------------


---

### **Principal growth stage 9: Senescence**

93	Stems begin to darken
95	50% of stems brown or black
97	Plant dead and dry
99	Harvested product

---

# Faba bean



# Sunflower

Weber and Bleiholder, 1990; Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of sunflower

(*Helianthus annuus* L.)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Germination

00	Dry seed (achene)
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from seed
06	Radicle elongated, root hairs developing
07	Hypocotyl with cotyledons emerged from seed
08	Hypocotyl with cotyledons growing towards soil surface
09	Emergence: cotyledons emerge through soil surface

---

### Principal growth stage 1: Leaf development<sup>1</sup>

10	Cotyledons completely unfolded
12	2 leaves (first pair) unfolded
14	4 leaves (second pair) unfolded
15	5 leaves unfolded
16	6 leaves unfolded
17	7 leaves unfolded
18	8 leaves unfolded
19	9 or more leaves unfolded

---

### Principal growth stage 3: Stem elongation

30	Beginning of stem elongation
31	1 visibly extended internode
32	2 visibly extended internodes
33	3 visibly extended internodes
3 .	Stages continuous till . . .
39	9 or more visibly extended internodes

---

### Principal growth stage 5: Inflorescence emergence

51	Inflorescence just visible between youngest leaves
53	Inflorescence separating from youngest leaves, bracts distinguishable from foliage leaves
55	Inflorescence separated from youngest foliage leaf
57	Inflorescence clearly separated from foliage leaves
59	Ray florets visible between the bracts; inflorescence still closed

---

<sup>1</sup> Stem elongation may occur earlier than stage 19; in this case continue with the principal stage 3

# Sunflower

Weber and Bleiholder, 1990; Lancashire et al., 1991

## Phenological growth stages and BBCH-identification keys of sunflower

---

Code	Description
<b>Principal growth stage 6: Flowering</b>	
61	Beginning of flowering: ray florets extended, disc florets visible in outer third of inflorescence
63	Disc florets in outer third of inflorescence in bloom (stamens and stigma visible)
65	Full flowering: disc florets in middle third of inflorescence in bloom (stamens and stigma visible)
67	Flowering declining: disc florets in inner third of inflorescence in bloom (stamens and stigma visible)
69	End of flowering: most disc florets have finished flowering, ray florets dry or fallen

---

### Principal growth stage 7: Development of fruit

71	Seeds on outer edge of the inflorescence are grey and have reached final size
73	Seeds on outer third of the inflorescence are grey and have reached final size
75	Seeds on middle third of the inflorescence are grey and have reached final size
79	Seeds on inner third of the inflorescence are grey and have reached final size

---

### Principal growth stage 8: Ripening

80	Beginning of ripening: seeds on outer third of anthocarp black and hard. Back of anthocarp still green
81	Seeds on outer third of anthocarp dark and hard. Back of anthocarp still green
83	Dark of anthocarp yellowish-green, bracts still green. Seeds about 50% dry matter
85	Seeds on middle third of anthocarp dark and hard. Back of anthocarp yellow, bracts brown edged. Seeds about 60% dry matter
87	Physiological ripeness: back of the anthocarp yellow. Bracts marbled brown. Seeds about 75–80% dry matter
89	Fully ripe: seeds on inner third of anthocarp dark and hard. Back of anthocarp brown. Bracts brown. Seeds about 85% dry matter


---

### Principal growth stage 9:


92	Over ripe, seeds over 90% dry matter
97	Plant dead and dry
99	Harvested product

---

# Sunflower



**00**



**10**



**12**



**14**



**59**



**61**



**65**

# Sunflower



18/32




53




57



79



89



92



# Beet

Meier et al., 1993

## Phenological growth stages and BBCH-identification keys of beet

(*Beta vulgaris* L. ssp. *vulgaris*)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Germination

00	Dry seed
01	Beginning of imbibition: seeds begins to take up water
03	Seed imbibition complete (pellet cracked)
05	Radicle emerged from seed (pellet)
07	Shoot emerged from seed (pellet)
09	Emergence: shoot emerges through soil surface

---

### Principal growth stage 1: Leaf development (youth stage)

10	First leaf visible (pinhead-size): cotyledons horizontally unfolded
11	First pair of leaves visible, not yet unfolded (pea-size)
12	2 leaves (first pair of leaves) unfolded
14	4 leaves (2nd pair of leaves) unfolded
15	5 leaves unfolded
1 .	Stages continuous till . . .
19	9 and more leaves unfolded

---

### Principal growth stage 3: Rosette growth (crop cover)

31	Beginning of crop cover: leaves cover 10% of ground
32	Leaves cover 20% of ground
33	Leaves cover 30% of ground
34	Leaves cover 40% of ground
35	Leaves cover 50% of ground
36	Leaves cover 60% of ground
37	Leaves cover 70% of ground
38	Leaves cover 80% of ground
39	Crop cover complete: leaves cover 90% of ground

---

### Principal growth stage 4: Development of harvestable vegetative plant parts Beet root

49	Beet root has reached harvestable size
----	--

---

### Principal growth stage 5: Inflorescence emergence (2nd year of growth)

51	Beginning of elongation of main stem
52	Main stem 20 cm long
53	Side shoot buds visible on main stem
54	Side shoots clearly visible on main stem
55	First individual flower buds on side shoots visible
59	First bracts visible; flower buds still closed

---

# Beet

Meier et al., 1993

## Phenological growth stages and BBCH-identification keys of beet

---

Code	Description
------	-------------

---

**Principal growth stage 6: Flowering**

60	First flowers open
61	Beginning of flowering: 10% of flowers open
62	20% of flowers open
63	30% of flowers open
64	40% of flowers open
65	Full flowering: 50% of flowers open
67	Flowering declining: 70% of flowers open or dry
69	End of flowering: all flowers dry, fruit set visible

---

**Principal growth stage 7: Development of fruit**

71	Beginning of seed development: seeds visible in infructescence
75	Pericarp green; fruit still mouldable; perisperm milky; colour of seed coat: beige

---

**Principal growth stage 8: Ripening**

81	Beginning of ripening: pericarp green-brown, seed coat light brown
85	Pericarp light brown, seed coat reddish brown
87	Pericarp hard, seed coat dark brown
89	Fully ripe: seed coat final colour (specific to variety and species), perisperm hard


---

**Principal growth stage 9: Senescence**


91	Beginning of leaf discolouration
93	Most leaves yellowish
95	50% of leaves brownish
97	Leaves dead
99	Harvested product (seeds)

---


# Beet




10




17




12




52



55



60



87

# Potato

Hack et al., 1993

## Phenological growth stages and BBCH-identification keys of potato

(*Solanum tuberosum* L.)

---

Code	Description of development from tuber	Description of development from seed	
<hr/>			
2- and 3digit			
<b>Principal growth stage 0: Sprouting / Germination</b>			
00	000	Innate or enforced dormancy, tuber not sprouted	Dry seed
01	001	Beginning of sprouting: sprouts visible (< 1 mm)	Beginning of seed imbibition
02	002	Sprouts upright (< 2 mm)	
03	003	End of dormancy: sprouts 2–3 mm	Seed imbibition complete
05	005	Beginning of root formation	Radicle (root) emerged from seed
07	007	Beginning of stem formation	Hypocotyl with cotyledons breaking
08	008	Stems growing towards soil surface, formation of scale leaves in the axils of which stolons will develop later	Hypocotyl with cotyledons growing towards soil surface
09	009	Emergence: stems break through soil surface	Emergence: cotyledons break through soil surface
021–029 <sup>1</sup>			

---

<sup>1</sup> For second generation sprouts

# Potato

Hack et al., 1993

## Phenological growth stages and BBCH-identification keys of potato

---

Code            Description of development from tuber and seed

---

2- and 3digit

### Principal growth stage 1: Leaf development

10	100	From tuber: first leaves begin to extend From seed: cotyledons completely unfolded
11	101	1st leaf of main stem unfolded (> 4 cm)
12	102	2nd leaf of main stem unfolded (> 4 cm)
13	103	3rd leaf of main stem unfolded (> 4 cm)
1 .	10 .	Stages continuous till . . .
19	109	9 or more leaves of main stem unfolded (> 4cm) (2digit); <sup>2</sup> 9 leaves of main stem unfolded (> 4 cm) (3digit)
	110	10th leaf of main stem unfolded (> 4 cm)
	11 .	Stages continuous till . . .
	119	19. leaf of main stem unfolded (> 4 cm)
	121	First leaf of 2nd order branch above first inflorescence unfolded (> 4 cm)
	122	2nd leaf of 2nd order branch above first inflorescence unfolded (> 4 cm)
	12 .	Stages continuous till . . .
	131	First leaf of 3rd order branch above 2nd inflorescence unfolded (> 4 cm)
	132	2nd leaf of 3rd order branch above 2nd inflorescence unfolded (> 4 cm)
	13 .	Stages continuous till . . .
	1NX	Xth leaf of nth order branch above n-1th inflorescence unfolded (> 4 cm)

---

<sup>2</sup> Stem development stops after termination of main stem by an inflorescence. Branches arise from axils of upper leaves of the main stem, exhibiting a sympodial branching pattern

# Potato Hack et al., 1993

## Phenological growth stages and BBCH-identification keys of potato

---

Codes	Description
-------	-------------

---

2- and 3digit

### Principal growth stage 2: Formation of basal side shoots below and above soil surface (main stem)

21	201	First basal side shoot visible (> 5 cm)
22	202	2nd basal side shoot visible (> 5 cm)
23	203	3rd basal side shoot visible (> 5 cm)
2 .	20 .	Stages continuous till . . .
29	209	9 or more basal side shoots visible (> 5 cm)

---

2- and 3digit

### Principal growth stage 3: Main stem elongation (crop cover)

31	301	Beginning of crop cover: 10% of plants meet between rows
32	302	20% of plants meet between rows
33	303	30% of plants meet between rows
34	304	40% of plants meet between rows
35	305	50% of plants meet between rows
36	306	60% of plants meet between rows
37	307	70% of plants meet between rows
38	308	80% of plants meet between rows
39	309	Crop cover complete: about 90% of plants meet between rows

---

2- and 3digit

### Principal growth stage 4: Tuber formation

40	400	Tuber initiation: swelling of first stolon tips to twice the diameter of subtending stolon
41	401	10% of total final tuber mass reached
42	402	20% of total final tuber mass reached
43	403	30% of total final tuber mass reached
44	404	40% of total final tuber mass reached
45	405	50% of total final tuber mass reached
46	406	60% of total final tuber mass reached
47	407	70% of total final tuber mass reached
48	408	Maximum of total tuber mass reached, tubers detach easily from stolons, skin set not yet complete (skin easily removable with thumb)
49	409	Skin set complete: (skin at apical end of tuber not removable with thumb) 95% of tubers in this stage

---

# Potato Hack et al., 1993

## Phenological growth stages and BBCH-identification keys of potato

---

Codes	Description
-------	-------------

---

2- and 3digit

### Principal growth stage 5: Inflorescence (cyme) emergence

51	501	First individual buds (1–2 mm) of first inflorescence visible (main stem)
55	505	Buds of first inflorescence extended to 5 mm
59	509	First flower petals of first inflorescence visible

---

2- and 3digit

### Principal growth stage 5: Inflorescence emergence (continuation)

521	Individual buds of 2nd inflorescence visible (second order branch)
525	Buds of 2nd inflorescence extended to 5 mm open (main stem)
529	First flower petals of 2nd inflorescence visible above sepals
531	Individual buds of 3rd inflorescence visible (3rd order branch)
535	Buds of 3rd inflorescence extended to 5 mm
539	First flower petals of 3rd inflorescence visible above sepals
5N .	Nth inflorescence emerging

---

2- and 3digit

### Principal growth stage 6: Flowering

60	600	First open flowers in population
61	601	Beginning of flowering: 10% of flowers in the first inflorescence open (main stem)
62	602	20% of flowers in the first inflorescence open
63	603	30% of flowers in the first inflorescence open
64	604	40% of flowers in the first inflorescence open
65	605	Full flowering: 50% of flowers in the first inflorescence open
66	606	60% of flowers in the first inflorescence open
67	607	70% of flowers in the first inflorescence open
68	608	80% of flowers in the first inflorescence open
69	609	End of flowering in the first inflorescence

---

# Potato Hack et al., 1993

## Phenological growth stages and BBCH-identification keys of potato

---

Codes	Description
-------	-------------

---

2- and 3digit

### Principal growth stage 6: Flowering (continuation)

621	Beginning of flowering: 10% of flowers in the 2nd inflorescence open (second order branch)
625	Full flowering: 50% of flowers in the 2nd inflorescence open
629	End of flowering in the 2nd inflorescence
631	Beginning of flowering: 10% of flowers in the 3rd inflorescence open (third order branch)
635	Full flowering: 50% of flowers in the 3rd inflorescence open
639	End of flowering in the 3rd inflorescence
6N .	Nth inflorescence flowering
6N9	End of flowering

---

2- and 3digit

### Principal growth stage 7: Development of fruit

70	700	First berries visible
71	701	10% of berries in the first fructification have reached full size (main stem)
72	702	20% of berries in the first fructification have reached full size
73	703	30% of berries in the first fructification have reached full size
7 .	70 .	Stages continuous till . . .
	721	10% of berries in the 2nd fructification have reached full size (second order branch)
	7N .	Development of berries in nth fructification
	7N9	Nearly all berries in the nth fructification have reached full size (or have been shed)

---

2- and 3digit

### Principal growth stage 8: Ripening of fruit and seed

81	801	Berries in the first fructification still green, seed light-coloured (main stem)
85	805	Berries in the first fructification ochre-coloured or brownish
89	809	Berries in the first fructification shrivelled, seed dark
	821	Berries in the 2nd fructification still green, seed light-coloured (second order branch)
	8N .	Ripening of fruit and seed in nth fructification

---



# Potato

Hack et al., 1993

## Phenological growth stages and BBCH-identification keys of potato

---

Codes	Description
-------	-------------

---

2- and 3digit

### Principal growth stage 9: Senescence


91	901	Beginning of leaf yellowing
93	903	Most of the leaves yellowish
95	905	50% of the leaves brownish
97	907	Leaves and stem dead, stems bleached and dry
99	909	Harvested product

---

# Potato

## The 2-digit decimal code

0 Sprouting			1 Leaf development			5 Inflorescence emergence			6 Flowering			7 Development of fruit			8 Ripening of fruit and seed			9 Senescence							
01	05	09	11	15	19	51	55	59	61	65	69	71	75	79	81	85	89	91	93	95	97				
Tuber formation						40						43			45			47			48			49	%



0 Sprouting			1 Leaf development main stem			2nd order			3rd order			4th order					
001	005	009	101	105	109	111	115	119	121	125	129	131	135	139	141	145	149

5 Inflorescence emergence main stem			2nd order			3rd order			4th order		
501	505	509	521	525	529	531	535	539	541	545	549

## The 3-digit decimal code


6 Flowering main stem			2nd order			3rd order		
601	605	609	621	625	629	631	635	639

7 Development of fruit main stem			2nd order		
701	705	709	721	725	729


8 Ripening of fruit and seed main stem			9 Senescence			
801	805	809	901	903	905	907

4 Tuber formation															
400					403			405			407			408	409


# Potato




00




03




05




09




19 bzw.  
112




51




40



59



65



79

# Pome fruit Meier et al., 1994

## Phenological growth stages and identification keys of pome fruit

(apple = *Malus domestica* Borkh., pear = *Pyrus communis* L.)

---

Code	Description
------	-------------

---

**Principal growth stage 0: Sprouting / Bud development**

00	Dormancy: leaf buds and the thicker inflorescence buds closed and covered by dark brown scales
01	Beginning of leaf bud swelling: buds visibly swollen, bud scales elongated, with light coloured patches
03	End of leaf bud swelling: bud scales light coloured with some parts densely covered by hairs
07	Beginning of bud break: first green leaf tips just visible
09	Green leaf tips about 5 mm above bud scales

---

**Principal growth stage 1: Leaf development**

10	Mouse-ear stage: Green leaf tips 10 mm above the bud scales; first leaves separating
11	First leaves unfolded (others still unfolding)
15	More leaves unfolded, not yet at full size
19	First leaves fully expanded

---

**Principal growth stage 3: Shoot development<sup>1</sup>**

31	Beginning of shoot growth: axes of developing shoots visible
32	Shoots about 20% of final length
33	Shoots about 30% of final length
3 .	Stages continuous till . . .
39	Shoots about 90% of final length

---

**Principal growth stage 5: Inflorescence emergence**

51	Inflorescence buds swelling: bud scales elongated, with light coloured patches
52	End of bud swelling: light coloured bud scales visible with parts densely covered by hairs
53	Bud burst: green leaf tips enclosing flowers visible
54	Mouse-ear stage: green leaf tips 10 mm above bud scales; first leaves separating
55	Flower buds visible (still closed)
56	Green bud stage: single flowers separating (still closed)
57	Pink bud stage: flower petals elongating; sepals slightly open; petals just visible
59	Most flowers with petals forming a hollow ball

---

<sup>1</sup> From terminal bud

# Pome fruit Meier et al., 1994

## Phenological growth stages and identification keys of pome fruit

---

Code	Description
------	-------------

---

### Principal growth stage 6: Flowering

60	First flowers open
61	Beginning of flowering: about 10% of flowers open
62	About 20% of flowers open
63	About 30% of flowers open
64	About 40% of flowers open
65	Full flowering: at least 50% of flowers open, first petals falling
67	Flowers fading: majority of petals fallen
69	End of flowering: all petals fallen

---

### Principal growth stage 7: Development of fruit

71	Fruit size up to 10 mm; fruit fall after flowering
72	Fruit size up to 20 mm
73	Second fruit fall
74	Fruit diameter up to 40 mm; fruit erect (T-stage: underside of fruit and stalk forming a T)
75	Fruit about half final size
76	Fruit about 60% final size
77	Fruit about 70% final size
78	Fruit about 80% final size
79	Fruit about 90% final size

---

### Principal growth stage 8: Maturity of fruit and seed

81	Beginning of ripening: first appearance of cultivar-specific colour
85	Advanced ripening: increase in intensity of cultivar-specific colour
87	Fruit ripe for picking
89	Fruit ripe for consumption: fruit have typical taste and firmness


---

### Principal growth stage 9: Senescence, beginning of dormancy

91	Shoot growth completed; terminal bud developed; foliage still fully green
92	Leaves begin to discolour
93	Beginning of leaf fall
95	50% of leaves discoloured
97	All leaves fallen
99	Harvested product

---

# Pome fruit



1 Leave bud smaller and slimer, directly on the long sprout

# Stone fruit Meier et al., 1994

## Phenological growth stages and BBCH-identification keys of stone fruit

(cherry = *Prunus cerasus* L., plum = *Prunus domestica* L. ssp. *domestica*, peach = *Prunus persica* Batsch., apricot = *Prunus ameriaca* L.)

Code	Description
<b>Principal growth stage 0: Sprouting/Bud development</b>	
00	Dormancy: leaf buds and the thicker inflorescence buds closed and covered by dark brown scales
01	Beginning of bud swelling (leaf buds); light brown scales visible, scales with light coloured edges
03	End of leaf bud swelling: scales separated, light green bud sections visible
09	Green leaf tips visible: brown scales fallen, buds enclosed by light green scales
<b>Principal growth stage 1: Leaf development</b>	
10	First leaves separating: green scales slightly open, leaves emerging
11	First leaves unfolded, axis of developing shoot visible
19	First leaves fully expanded
<b>Principal growth stage 3: Shoot development 1</b>	
31	Beginning of shoot growth: axes of developing shoots visible
32	Shoots about 20% of final length
33	Shoots about 30% of final length
3 .	Stages continuous till . . .
39	Shoots about 90% of final length
<b>Principal growth stage 5: Inflorescence emergence</b>	
51	Inflorescence buds swelling: buds closed, light brown scales visible
53	Bud burst: scales separated, light green bud sections visible
54	Inflorescence enclosed by light green scales, if such scales are formed (not all cultivars)
55	Single flower buds visible (still closed) borne on short stalks, green scales slightly open
56	Flower pedicel elongating; sepals closed; single flowers separating
57	Sepals open: petal tips visible; single flowers with white or pink petals (still closed)
59	Most flowers with petals forming a hollow ball

<sup>1</sup> From terminal bud

# Stone fruit Meier et al., 1994

## Phenological growth stages and BBCH-identification keys of stone fruit

---

Code	Description
------	-------------

---

### Principal growth stage 6: Flowering

60	First flowers open
61	Beginning of flowering: about 10% of flowers open
62	About 20% of flowers open
63	About 30% of flowers open
64	About 40% of flowers open
65	Full flowering: at least 50% of flowers open, first petals falling
67	Flowers fading: majority of petals fallen
69	End of flowering: all petals fallen

---

### Principal growth stage 7: Development of fruit

71	Ovary growing; fruit fall after flowering
72	Green ovary surrounded by dying sepal crown, sepals beginning to fall
73	Second fruit fall
75	Fruit about half final size
76	Fruit about 60% of final size
77	Fruit about 70% of final size
78	Fruit about 80% of final size
79	Fruit about 90% of final size

---

### Principal growth stage 8: Maturity of fruit and seed

81	Beginning of fruit colouring
85	Colouring advanced
87	Fruit ripe for picking
89	Fruit ripe for consumption: fruit have typical taste and firmness

---




### Principal growth stage 9: Senescence, beginning of dormancy

91	Shoot growth completed; foliage still fully green
92	Leaves begin to discolour
93	Beginning of leaf fall
95	50% of leaves discoloured or fallen
97	All leaves fallen
99	Harvested product

---



# Stone fruit



# Currants

Meier et al., 1994

## Phenological growth stages and BBCH-identification keys of currants

(black currant = *Ribes nigrum* L., red currant = *Ribes rubrum* L.)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Sprouting/Bud development

00	Dormancy: leaf buds and the thicker inflorescence buds closed and covered by dark brown scales
01	Beginning of bud swelling: bud scales elongated
03	End of bud swelling: edges of bud scales light coloured
07	Beginning of bud burst: first green or red leaf tips just visible
09	Leaf tips extended beyond scales

---

### Principal growth stage 1: Leaf development

10	Leaf tips above the bud scales: first leaves separating
11	First leaves unfolded (others still unfolding)
15	More leaves unfolded, not yet full size
19	First leaves fully expanded

---

# Currants

Meier et al., 1994

## Phenological growth stages and BBCH-identification keys of currants

---

Code	Description
------	-------------

---

### Principal growth stage 3: Shoot development<sup>1</sup>

31	Beginning of shoot growth: axes of developing shoots visible
32	Shoots about 20% of final length
33	Shoots about 30% of final length
3 .	Stages continuous till . . .
39	Shoots about 90% of final length

---

### Principal growth stage 5: Inflorescence emergence

51	Inflorescence buds and leaf buds swelling: buds closed, light brown scales visible
53	Bud burst: scales separated light green but sections visible
54	Green or red leaf tips above bud scales
55	First flower buds (compact raceme) visible beside unfolded leaves
56	Beginning of raceme elongation
57	First flower bud separated on elongating raceme
59	Grape stage: all flower buds separated

---

### Principal growth stage 6: Flowering

60	First flowers open
61	Beginning of flowering: about 10% of flowers open
65	Full flowering: at least 50% of flowers open, first petals falling
67	Flowers fading: majority of petals fallen
69	End of flowering: all petals fallen

---

### Principal growth stage 7: Development of fruit

71	Beginning of fruit growth: first fruits visible at raceme base
72	20% of fruits formed
73	30% of fruits formed
74	40% of fruits formed
75	50% of fruits formed
76	60% of fruits formed
77	70% of fruits formed
78	80% of fruits formed
79	90% of fruits formed

---

<sup>1</sup> From terminal bud

# Currants

Meier et al., 1994

## Phenological growth stages and BBCH-identification keys of currants

---

Code	Description
<b>Principal growth stage 8: Maturity of fruit and seed</b>	
81	Beginning of ripening: change to cultivar-specific fruit color
85	Advanced ripening: first berries at base of racemes have cultivar-specific color
87	Fruit ripe for picking: most berries ripe
89	Berries at base of racemes tending to drop (beginning of fruit abscission)
<b>Principal growth stage 9: Senescence, beginning of dormancy</b>	
91	Shoot growth completed; terminal bud developed; foliage still fully green
92	Leaves begin to discolour
93	Beginning of leaf fall
95	50% of leaves discoloured or fallen
97	All leaves fallen
99	Harvested product

---

# Currants



# Strawberry

Meier et al., 1994

## Phenological growth stages and BBCH-identification keys of strawberry

(*Fragaria ananassa* Duch.)

---

Code	Description
------	-------------

---

**Principal growth stage 0: Sprouting/Bud development**

00	Dormancy: Leaves prostrate and partly dead
03	Main bud swelling

---

**Principal growth stage 1: Leaf development**

10	First leaf emerging
11	First leaf unfolded
12	2nd leaf unfolded
13	3rd leaf unfolded <sup>1</sup>
1 .	Stages continuous till . . .
19	9 or more leaves unfolded

---

**Principal growth stage 4: Development of stolons and young plants**

41	Beginning of stolon (runner) formation: stolons visible (about 2 cm long)
42	First daughter plant visible
43	Beginning of root development in first daughter plant
45	First daughter plant with roots (ready for planting)
49	Several daughter plants with roots (ready for planting)

---

**Principal growth stage 5: Inflorescence emergence**

55	First set flowers at the bottom of the rosette
56	Inflorescence elongating
57	First flower buds emerged (still closed)
58	Early balloon stage: first flowers with petals forming a hollow ball
59	Most flowers with petals forming a hollow ball

---

**Principal growth stage 6: Flowering**

60	First flowers open (primary or A-flower)
61	Beginning of flowering: about 10% of flowers open
65	Full flowering: secondary (B) and tertiary (C) flowers open, first petals falling
67	Flowers fading: majority of petals fallen

---

<sup>1</sup> Normally after the three leaf stage the bud development occurs in principal growth stage 5

# Strawberry

Meier et al., 1994

## Phenological growth stages and BBCH-identification keys of strawberry

---

Code	Description
------	-------------

---

### Principal growth stage 7: Development of fruit

71	Receptacle protruding from sepal whorl
73	Seeds clearly visible on receptacle tissue

---

### Principal growth stage 8: Maturity of fruit

81	Beginning of ripening: most fruits white in colour
85	First fruits have cultivar-specific colour
87	Main harvest: more fruits coloured
89	Second harvest: more fruits coloured




---

### Principal growth stage 9: Senescence, beginning of dormancy

91	Beginning of axillary bud formation
92	New leaves with smaller lamina and shortened stalk visible
93	Old leaves dying, young leaves curling; old leaves of cultivarspecific colour
97	Old leaves dead

---

# Strawberries





# Citrus

Agusti et al., 1995

## Phenological growth stages and BBCH-identification keys of citrus

(*Citrus* spp. L.),

---

Code	Description
------	-------------

---

**Principal growth stage 0: Sprouting/Bud development**

00	Dormancy: leaf and inflorescence buds undifferentiated, closed and covered by green scales
01	Beginning of bud swelling
03	End of bud swelling: green scales slightly separated
07	Beginning of bud burst
09	Green leaf tips visible

---

**Principal growth stage 1: Leaf development**

10	First leaves separating: green scales slightly open, leaves emerging
11	First leaves visible <sup>1</sup>
15	More leaves visible, not yet at full size
19	First leaves fully expanded

---

**Principal growth stage 3: Shoot development**

31	Beginning of shoot growth: axes of developing shoots visible
32	Shoots about 20% of final length
39	Shoots about 90% of final length

---

**Principal growth stage 5: Inflorescence emergence**

51	Inflorescence buds swelling: buds closed, light green scales visible
53	Bud burst: scales separated, floral tips visible
55	Flowers visible, still closed (green bud), borne on single or multiflowered leafy or leafless inflorescences
56	Flower petals elongating; sepals covering half corolla (white bud)
57	Sepals open: petal tips visible; flowers with white or purplish petals, still closed
59	Most flowers with petals forming a hollow ball

---

<sup>1</sup> In Citrus the term "visible" replaces "unfolded" used for other fruit species. Leaf unfolding takes place precociously in citrus

# Citrus Agusti et al., 1995

## Phenological growth stages and BBCH-identification keys of citrus

---

Code	Description
------	-------------

---

**Principal growth stage 6: Flowering**

60	First flowers open
61	Beginning of flowering: about 10% of flowers open
65	Full flowering: 50% of flowers open; first petals falling
67	Flowers fading: majority of petals fallen
69	End of flowering: all petals fallen

---

**Principal growth stage 7: Development of fruit**

71	Fruit set; beginning of ovary growth; beginning of fruitlets abscission
72	Green fruit surrounded by sepal crown
73	Some fruits slightly yellow: beginning of physiological fruit drop
74	Fruits about 40% of final size. Dark green fruit: end of physiological fruit drop
79	Fruits about 90% of final size

---

**Principal growth stage 8: Maturity of fruit**

81	Beginning of fruit colouring (colour-break)
83	Fruit ripe for picking; fruit has not yet developed variety-specific colour
85	Advanced ripening; increase in intensity of variety-specific colour
89	Fruit ripe for consumption; fruit has typical taste and firmness; beginning of senescence and fruit abscission











---

**Principal growth stage 9: Senescence, beginning of dormancy**

91	Shoot growth complete; foliage fully green
93	Beginning of senescence and abscission of old leaves
97	Winter dormancy period

---

# Citrus



# Olive Sanz-Cortés et al., 2002

## Phenological growth stages and BBCH-identification keys of olive tree (*Olea europaea* L.)

Code	Description
<b>Principal growth stage 0: Bud development</b>	
00	Foliar buds at the apex of shoots grown the previous crop-year are completely closed, sharp-pointed, stemless and ochre-coloured (Fig. 1: 00).
01	Foliar buds start to swell and open, showing the new foliar primordia.
03	Foliar buds lengthen and separate from the base.
07	External small leaves open, not completely separated, remaining joined by apices (Fig. 1: 07).
09	External small leaves opening further with their tips inter crossing (Fig. 1: 09)
<b>Principal growth stage 1: Leaf development</b>	
11	First leaves completely separated. Grey-greenish coloured (Fig. 1: 11)
15	The leaves are more separated without reaching their final size.
19	First leaves turn greenish on the upperside. Leaves get the typical variety size and shape.
<b>Principal growth stage 3: Shoot development</b>	
31	Shoots reach 10 % of final size.
33	Shoots reach 30 % of final size (Fig. 1: 33).
37	Shoots reach 70 % of final size.
<b>Principal growth stage 5: Inflorescence emergence.</b>	
50	Inflorescence buds in leaf axiles are completely closed. They are sharp-pointed, stemless and ochre-coloured.
51	Inflorescence buds start to swell on its stem.
52	Inflorescence buds open. Flower cluster development starts (Fig 1: 53).
54	Flower cluster growing
55	Flower cluster totally expanded. Floral buds start to open (Fig 1: 55).
57	The corolla, green-coloured, is longer than calyx (Fig 1: 57).
59	The corolla changes from green to white colour.

# Olive

Sanz-Cortés et al., 2002

## Phenological growth stages and BBCH-identification keys of olive tree (*Olea europaea* L.)

---

Code	Description
------	-------------

---

**Principal growth stage 6: Flowering**

60	First flowers open (Fig 1: 60).
61	Beginning of flowering: 10 % of flowers open.
65	Full flowering: at least 50 % of flowers open (Fig 1: 65).
67	First petals falling.
68	Majority of petals fallen or faded (Fig 1: 68).
69	End of flowering, fruit set, non-fertilized ovaries fallen.

---

**Principal growth stage 7: Fruit development**

71	Fruit size about 10 % of final size (Fig 1: 71).
75	Fruit size about 50 % of final size. Stone starts to lignificate (it shows cutting resistance).
79	Fruit size about 90 % of final size. Fruit suitable for picking green olives (Fig 1: 79).

---

**Principal growth stage 8: Maturity of fruit**

80	Fruit deep green colour becomes light green, yellowish.
81	Beginning of fruit colouring (Fig 1: 81).
85	Increasing of specific fruit colouring.
89	Harvest maturity: fruits get the typical variety colour, remaining turgid, suitable for oil extraction (Fig. 1: 89).




---


**Principal growth stage 9: Senescence**

92	Overripe: fruits lose turgidity and start to fall (Fig 1: 92).
----	--


---

# Olive







53




55




57




60




65




68




71




79



81



89



92



# Coffee

Arcila-Pulgarín et al., 2002

## Phenological growth stages and BBCH-identification keys of the coffee plant (*Coffea sp.*)

---

Code	Description
<b>Principal growth stage 0: Germination, vegetative propagation</b>	
00	Dry seed (11-12% moisture content), beige color if parchment present or bluish-green if parchment and silver skin removed. Cutting (orthotropic, mononodal, 60 mm long, two half trimmed leaves). Stump with bulky nodes and no buds visible
01	Beginning of seed imbibition, bean swollen, whitish, no radicle visible. Cutting planted in rooting media, no shoots visible, no callus visible
02	Seed imbibition complete, bean whitish, small swelling visible at one end of bean where the embryo is located. Callus formation begins on cuttings. Bud burst start on stumps
05	Seed radicle protrusion and hooking. Shoot and root formation on the cuttings. Green, rounded buds visible on the stumps
06	Elongation of radicle, formation of root hairs and lateral roots on seeds and cuttings.
07	Hypocotyl with cotyledons breaking through the seed coat. Cuttings have formed shoots and branched roots.
09	<i>Emergence:</i> Seeds have emerged from soil and show the hypocotile with cotyledons still enclosed in the parchment. The cuttings present roots 6-7 cm. long and shoots with 1-2 nodes. Stumps show sprouts with first leaf initials.

---

## Principal growth stage 1: Leaf development on main shoot of the young plant, and branches of the coffee tree

10	Cotyledons completely unfolded. First pair of true leaves separating on shoot or first pair of true leaves separating on branch of the coffee tree
11	first leaf pair unfolded, not yet at full size. Leaves are light green or bronze
12	2 leaf pairs unfolded, not yet at full size. Leaves are light green or bronze
13	3 leaf pairs unfolded, not yet full size. The third leaf pair from apex is dark green
14	4 leaf pairs unfolded. The fourth leaf pair from apex is dark green and has reached full size
1.	Stages continues till...
19	9 or more leaf pairs unfolded

---

# Coffee

Arcila-Pulgarín et al., 2002

## Phenological growth stages and BBCH-identification keys of the coffee plant (*Coffea sp.*)

---

Code	Description
------	-------------

---

### Principal growth stage 2: Formation of branches (only for plants in the field)

20	First pair of primary branches are visible
21	10 pair of primary branches visible
22	20 pair of primary branches visible
23	30 pair of primary branches visible
2.	Stages continues till...
29	90 or more pairs of primary branches visible

---

### Principal growth stage 3: Branch elongation

31	10 nodes present in the branch(es)
32	20 nodes present in the branch(es)
3.	Stages continues till...
39	90 or more nodes present in the branch(es)

---

### Principal growth stage 5: Inflorescence emergence

51	Inflorescence buds swelling in leaf axils
53	Inflorescence buds burst and covered by brown mucilage; no flowers visible
57	Flowers visible, still closed and tightly join, borne on multiflowered inflorescence (3-4 flowers per inflorescence)
58	Flowers visible, untight, still closed, petals 4-6 mm long and green (dormant stage)
59	Flowers with petals elongated ( 6-10 mm long), still closed and white color.

---

### Principal growth stage 6: Flowering

60	First flowers open
61	10% of flowers open
63	30% of flowers open
65	50% of flowers open
67	70% of flowers open
69	90% of flowers open

---

# Coffee

Arcila-Pulgarín et al., 2002


## Phenological growth stages and BBCH-identification keys of the coffee plant (*Coffea sp.*)








---

Code	Description
<b>Principal growth stage 7: Development of fruit</b>	
70	Fruits visible as small yellowish berries
71	<i>Fruit set</i> : Beginning of berry growth. Fruits have reached 10% of final size (pinheads).
73	Fruits are light green and contents are liquid and crystalline. Fruits have reached 30% of final size (fast growth).
75	Fruits are light green and its contents are liquid and crystalline. Fruits have reached 50% of final size.
77	Fruits are dark green and its contents are solid and white. Fruits have reached 70% of final size.
79	Fruits are pale green and its contents are solid and white. Physiological maturity is complete. Fruits have reached 90% of final size.
<b>Principal growth stage 8: Ripening of fruit and seed</b>	
81	Beginning of change of fruit coloration from pale green to yellow or red
85	Increase in intensity (variety-specific), yellow or red , fruit color; fruit not yet ready for picking.
88	Fruit is fully-ripe color and ready for picking.
89	Overripe; beginning of darkening or drying; fruits stay on the tree or abscission begins.
<b>Principal growth stage 9: Senescence</b>	
90	Shoots have completed their development; the plant appears of an intense dark green color, leaves are of normal size and harvest locates at the bottom part of the plant.
93	Older leaves change its color from deep green to yellow with red spots, and fall specially at harvesting time.
94	The foliage changes to a pale green color. Defoliation is observed on the bottom part of the main stem and lower branches.
97	The production zone has moved towards the upper parts in the main shoot and outer parts of branches, leaves are of smaller size than normal, strong defoliation is observed on the bottom and inner part of the plant, some dead branches are observed at the bottom.
98	The production zone is limited to a very few branches on the top of the shoot and a very few nodes on the tip of these branches, and the plant is heavily defoliated. A high-degree of senescence has been reached. 90% or more of the harvest completed.
99	Post harvest or storage treatments

---

# Coffee





# Musaceae

Gonzales et al., 2002

## Phenological growth stages and BBCH-identification keys of *edible Musaceae*

---

Code	Description
2- 3- 4-digit	
<b>Principal stage 0: Sprouting or emergence</b>	
00 000 0000	Recently planted material (plants from tissue cultures and corns) without visible growth
05 005 0005	Emergence of the 1st new leaf in plants from tissue cultures or of the foliar shoot of the corn
<b>Principal stage 1: Leaf development</b>	
10 100 1000	Formation of the 1st leaf of the planted corn or the candela leaf in tissue culture plants (candela stage 0)
	1002 Leaf 1 at candela stage 2
	1004 Leaf 1 at candela stage 4
	1006 Leaf I at candela stage 6
	1008 Leaf I at candela stage 8
11 101 1010	One leaf completely open and the youngest leaf at candela stage 0
	1012 One leaf completely open and the youngest leaf at candela stage 2
	1014 One leaf completely open and the youngest leaf at candela stage 4
	1016 One leaf completely open and the youngest leaf at candela stage 6
	1018 One leaf completely open and the youngest leaf at candela stage 8
12 102 1020	Two leaves completely open and the youngest leaf at candela stage 0
	1022 Two leaves completely open and the youngest leaf at candela stage 2
	1024 Two leaves completely open and the youngest leaf at candela stage 4
	1026 Two leaves completely open and the youngest leaf at candela stage 6
	1028 Two leaves completely open and the youngest leaf at candela stage 8

---

# Musaceae

Gonzales et al., 2002

## Phenological growth stages and BBCH-identification keys of *of edible Musaceae*

---

Codes			Description
2-	3-	4-digit	
<b>Principal stage 1: Leaf development (cont.)</b>			
13	103	1030	Three leaves completely open and the youngest leaf at candela stage 0
		1032	Three leaves completely open and the youngest leaf at candela stage 2
		1034	Three leaves completely open and the youngest leaf at candela stage 4
		1036	Three leaves completely open and the youngest leaf at candela stage 6
		1038	Three leaves completely open and the youngest leaf at candela stage 8 stages continue till...
19	109	1090	9 or more leaves (only 2 digit code ) or nine leaves completely open and the youngest leaf at candela stage 0
		1092	Nine leaves completely open and the youngest leaf at candela stage 2
		1094	Nine leaves completely open and the youngest leaf at candela stage 4
		1096	Nine leaves completely open and the youngest leaf at candela stage 6
		1098	Nine leaves completely open and the youngest leaf at candela stage 8 stages continue till...
	119	1190	Nineteen or more leaves completely open and the youngest leaf at candela stage 0
		1192	Nineteen or more leaves completely open and the youngest leaf at candela stage 2
		1194	Nineteen or more leaves completely open and the youngest leaf at candela stage 4
		1196	Nineteen or more leaves completely open and the youngest leaf at candela stage 6
		1198	Nineteen or more leaves completely open and the youngest leaf at candela stage 8

---

# Musaceae

Gonzales et al., 2002

## Phenological growth stages and BBCH-identification keys of *of edible Musaceae*

---

Code			Description
2-	3-	4-digit	
<b>Principal stage 2: Sucker formation</b>			
21	201	2010	1st sucker with visible leaf
		2011	1st sucker with visible sword leaf
		2012	1st sucker with visible water leaf
22	202	2020	2nd sucker with visible leaf
		2021	2nd sucker with visible sword leaf
		2022	2nd sucker with visible water leaf
23	203	2030	3rd sucker with visible leaf
		2031	3rd sucker with visible sword leaf
		2032	3rd sucker with visible water leaf
			stages continue till...
29	209	2090	nine or more suckers with visible leaf
		2091	nine or more suckers with visible sword leaf
		2092	nine or more suckers with visible water leaf
<hr/>			
2-	3-	4-digit	
<b>Principal stage 3: Pseudostem elongation</b>			
35	305	3050	The pseudostem reaches 50 % of its typical thickness and length according to the genome or clone
39	309	3090	The maximum length and thickness of the pseudostem are reached according to the genome or clone and the formation of new leaves of normal size has been terminated

---



# Musaceae

Gonzales et al., 2002

## Phenological growth stages and BBCH-identification keys of *edible Musaceae*

---

Code	Description
2- 3- 4-digit	
<b>Principal stage 4: Leaf development of the sucker (sword sucker)</b>	
40 400 4000	Sub-phase of dependent growth: the sucker becomes visible and develops the leaf shoot
41 401 4011	Development of the 1st lanceolate leaf
4012	Development of the 2nd lanceolate leaf
4013	Development of the 3rd lanceolate leaf
4014	Development of the 4th lanceolate leaf
4015	Development of the 5th lanceolate leaf
4016	Development of the 6th lanceolate leaf
4017	Development of the 7th lanceolate leaf
4018	Development of the 8th lanceolate leaf
4019	Development of the 9th or more lanceolate leaves
45 405 4050	Sub-phase of independent growth: leaves of approx. 10 cm width are developed (original leaf/zero leaf/F10)
4051	Development of the 1st leaf of approx. 10 cm width
4052	Development of the 2nd leaf of approx. 10 cm width
4053	Development of the 3rd leaf of approx. 10 cm width
4054	Development of the 4th leaf of approx. 10 cm width
4055	Development of the 5th leaf of approx. 10 cm width
4056	Development of the 6th leaf of approx. 10 cm width
4057	Development of the 7th leaf of approx. 10 cm width
4058	Development of the 8th leaf of approx. 10 cm width
4059	Development of the 9th or more leaves of approx. 10 cm width
49 409 4090	End of this phase is reached with the development of the first leaf with characteristics of the genome or clone (length width ratio, leaf surface index). Beginning of synchronised development of "normal" leaves (FM)
2- 3- 4-digit	
<b>Principal stage 5: Emergence of inflorescence</b>	
50 500 5000	The development of new normal leaves has been terminated and the flower bract emergence
51 501 5010	Flower bract at candela stage 2
52 502 5020	Flower bract at candela stage 4
53 503 5030	Flower bract at candela stage 6
54 504 5040	Flower bract at candela stage 8
55 505 5050	Flower bract completely open
59 509 5090	Emergence of the last bract leaf or first sterile bract protecting the flower

---

# Musaceae

Gonzales et al., 2002

## Phenological growth stages and BBCH-identification keys of *edible Musaceae*

---

Code			Description
2-	3-	4-digit	
<b>Principal stage 6: Flowering</b>			
60	600	6000	The stage begins with the emergence of the flower protected by the last bract leaf (1st sterile bract)
61	601	6010	A bract which does not protect any hand of flowers rises (2nd sterile bract) and the rachis or flower stalk takes a pendulum position
62	602	6020	The bract rises which protects the first hand of female or pistillate flowers
63	603	6030	The bract rises which protects the second hand of female or pistillate flowers
64	604	6040	The bract rises which protects the third hand of female or pistillate flowers
65	605	6050	Full bloom: at least 50 % of the hands of females flowers are developed
69	609	6090	The bracts which protect the hands wither and fall off and the fingers are bent into a direction perpendicular to the rachis
<hr/>			
2-	3-	4-digit	
<b>Principal stage 7: Development of the fruit</b>			
70	700	7000	At least 50 % of the fingers show an upwards curvature and the fruits (fingers) begin to fill
71	701	7010	Total exposure of the fingers or female flowers (protective bracts fallen off or bent and withered above the hands)
72	702	7020	The fingers of the hands show the characteristic curvature of the fruit (upwards and almost parallel to the axis or rachis)
73	703	7030	From the first two hands up to 30 % of the hands have reached the maximum thickness of the fruit
74	704	7040	Up to 40 % of the hands have reached the maximum thickness of the fruit
75	705	7050	Up to 50 % of the hands have reached the maximum thickness of the fruit
76	706	7060	Up to 60 % of the hands have reached the maximum thickness of the fruit
77	707	7070	Up to 70 % of the hands have reached the maximum thickness of the fruit
78	708	7080	Up to 80 % of the hands have reached the maximum thickness of the fruit
79	709	7090	All hands have reached the maximum thickness of the fruit and no hand shows a loss of weight

---

# Musaceae

Gonzales et al., 2002


## Phenological growth stages and BBCH-identification keys of *edible Musaceae*

Code	Description		
2-	3-	4-digit	
<b>Principal stage 8: Ripening of the fruit</b>			
80	800	8000	Ripening starts when the fruit has reached the maximum thickness, begins to lose weight and shows changes of the colour by which the degrees of maturity are defined
81	801	8010	Degree of maturity 1: green. Normal colour of the fresh fruit
82	802	8020	Degree of maturity 2: tinge of yellow. First modification of colour during the ripening cycle
83	803	8030	Degree of maturity 3: more green than yellow
84	804	8040	Degree of maturity 4: more yellow than green
85	805	8050	Degree of maturity 5: tinge of green
86	806	8060	Degree of maturity 6: all yellow
87	807	8070	Degree of maturity 7: yellow with brown specks. Fruit is completely ripe, has the best flavour and a high nutritive value
88	808	8080	Degree of maturity 8: 20-50 % of surface discoloured brown or spoiled
89	809	8090	Degree of maturity 9: More than 50 % of the surface of the fruits is discoloured brown and spoiled
2-	3-	4-digit	
<b>Principal stage 9: Senescence and death</b>			
90	900	9000	More than 50 % of the surface of the fruits are discoloured brown
91	901	9010	The leaves the plant shows have died off acropetally and the male flowers have withered, are necrotic and/or have fallen off
93	903	9030	Total rot and necrosis of the fruits
95	905	9050	Degeneration (necrosis) of the flower
97	907	9070	The sheaths enclosed in the pseudostem become brittle which indicates the beginning of necrosis of the pseudostem. The pseudostem turns to be brown
98	908	9080	Total decomposition of the tissues and fall down of the pseudostem


\* Harvested product

\* post-harvest or storage treatments take place at stage 99, 909 or 9090

# Musacea




**0 Sprouting**




**1 Leaf development**



**2 Sucker formation**



**3 Pseudostem elongation**



**4 Leaf development  
of the sucker**



**5 Emergence of inflorescence**



**6 Flowering**




**7 Development of the fruit**




**8 Ripening of the fruit**




**9 Senescence and death**




**IXX0**




**IXX2**



**IXX4**




**IXX6**



**IXX8**

**XX = 00 to 19 indicate  
the number of leaves**







**20**




**2x1**

**2x2**







60




61




62




65




69



70




72




79







81




83




84



85




86




87



88



89



91



93



95



98

# Grapevine Lorenz et al., 1994

## Phenological growth stages and BBCH-identification keys of grapevine

(*Vitis vinifera* L. ssp. *vinifera*)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Sprouting/Bud development

00	Dormancy: winter buds pointed to rounded, light or dark brown according to cultivar; bud scales more or less closed according to cultivar
01	Beginning of bud swelling: buds begin to expand inside the bud scales
03	End of bud swelling: buds swollen, but not green
05	"Wool stage": brown wool clearly visible
07	Beginning of bud burst: green shoot tips just visible
08	Bud burst: green shoot tips clearly visible

---

### Principal growth stage 1: Leaf development

11	First leaf unfolded and spread away from shoot
12	2nd leaves unfolded
13	3rd leaves unfolded
1 .	Stages continuous till . . .
19	9 or more leaves unfolded

---

### Principal growth stage 5: Inflorescence emerge

53	Inflorescences clearly visible
55	Inflorescences swelling, flowers closely pressed together
57	Inflorescences fully developed; flowers separating

---

### Principal growth stage 6: Flowering

60	First flowerhoods detached from the receptacle
61	Beginning of flowering: 10% of flowerhoods fallen
62	20% of flowerhoods fallen
63	Early flowering: 30% of flowerhoods fallen
64	40% of flowerhoods fallen
65	Full flowering: 50% of flowerhoods fallen
66	60% of flowerhoods fallen
67	70% of flowerhoods fallen
68	80% of flowerhoods fallen
69	End of flowering

---

# Grapevine

Lorenz et al., 1994

## Phenological growth stages and BBCH-identification keys of grapevine

---

Code	Description
------	-------------

---

### Principal growth stage 7: Development of fruits

71	Fruit set: young fruits begin to swell, remains of flowers lost
73	Berries goat-sized, bunches begin to hang
75	Berries pea-sized, bunches hang
77	Berries beginning to touch
79	Majority of berries touching

---

### Principal growth stage 8: Ripening of berries

81	Beginning of ripening: berries begin to develop variety-specific colour
83	Berries developing colour
85	Softening of berries
89	Berries ripe for harvest




---

### Principal growth stage 9: Senescence

91	After harvest; end of wood maturation
92	Beginning of leaf discolouration
93	Beginning of leaf-fall
95	50% of leaves fallen
97	End of leaf-fall
99	Harvested product

---

# Grapevine



# Soybean Munger et al., 1997

## Phenological growth stages and BBCH-identification keys of the soybean

(*Glycine max* L. Merr.)

---

Code	Description
------	-------------

---

2- and 3digit

### Principal growth stage 0: Germination

00	000	Dry seed
01	001	Beginning of seed imbibition
03	003	Seed imbibition complete
05	005	Radicle emerged from seed
06	006	Elongation of radicle; formation of root hairs
07	007	Hypocotyl with cotyledons breaking through seed coat
08	008	Hypocotyl reaches the soil surface; hypocotyl arch visible
09	009	Emergence: hypocotyl with cotyledons emerged above soil surface ("cracking stage")

---

2- and 3digit

### Principal growth stage 1: Leaf development (Main shoot)

10	100	Cotyledons completely unfolded
11	101	First pair of true leaves unfolded (unifoliolate leaves on the first node)
12	102	Trifoliolate leaf on the 2nd node unfolded
13	103	Trifoliolate leaf on the 3rd node unfolded
1 .	10 .	Stages continuous till . . .
19	109	Trifoliolate leaf on the 9th node unfolded. No side shoots visible <sup>1</sup>
	110	Trifoliolate leaf on the 10th node unfolded <sup>1</sup>
	111	Trifoliolate leaf on the 11th node unfolded <sup>1</sup>
	112	Trifoliolate leaf on the 12th node unfolded <sup>1</sup>
	113	Trifoliolate leaf on the 13th node unfolded <sup>1</sup>
	11 .	Stages continuous till . . .
	119	Trifoliolate leaf on the 19th node unfolded <sup>1</sup>

---

<sup>1</sup>The side shoot development may occur earlier, in this case continue with the principal growth stage 2

# Soybean Munger et al., 1997

## Phenological growth stages and BBCH-identification keys of the soybean

---

Code	Description
------	-------------

---

2- and 3digit

### Principal growth stage 2: Formation of side shoots

21	201	First side shoot visible
22	202	2nd side shoot of first order visible
23	203	3rd side shoot of first order visible
2 .	20 .	Stages continuous till . . .
29	209	9 or more side shoots of first order visible (2 digit)
		9th side shoot of first order visible (3 digit)
	210	10th side shoot of first order visible
	221	First side shoot of 2nd order visible
	22 .	Stages continuous till . . .
	229	9th side shoot of 2nd order visible
	2N1	First side shoot of Nth order visible
	2N9	9th side shoot of Nth order visible

---

2- and 3digit

### Principal growth stage 4: Development of harvestable vegetative plant parts **Main shoot**

49	409	Harvestable vegetative plant parts have reached final size (Cutting of soybean plants for feeding purposes)
----	-----	---

---

2- and 3digit

### Principal growth stage 5: Inflorescence emergence **(Main shoot)**

51	501	First flower buds visible
55	505	First flower buds enlarged
59	509	First flower petals visible; flower buds still closed

---

# Soybean Munger et al., 1997

## Phenological growth stages and BBCH-identification keys of the soybean

---

Code	Description
------	-------------

---

2- and 3digit

### Principal growth stage 6: Flowering (Main shoot)

60	600	First flowers opened (sporadically in population)
61	601	Beginning of flowering: about 10% of flowers open <sup>2</sup> Beginning of flowering <sup>3</sup>
62	602	About 20% of flowers open <sup>2</sup>
63	603	About 30% of flowers open <sup>2</sup>
64	604	About 40% of flowers open <sup>2</sup>
65	605	Full flowering: about 50% of flowers open <sup>2</sup> Main period of flowering <sup>3</sup>
66	606	About 60% of flowers open <sup>2</sup>
67	607	Flowering declining <sup>2</sup>
69	609	End of flowering: first pods visible (approx. 5 mm length) <sup>2</sup>

---

2- and 3digit

### Principal growth stage 7: Development of fruits and seeds

70	700	First pod reached final length (15–20 mm)
71	701	About 10% of pods have reached final length (15–20 mm) <sup>2</sup> Beginning of pod development <sup>3</sup>
72	702	About 20% of pods have reached final length (15–20 mm) <sup>2</sup>
73	703	About 30% of pods have reached final length (15–20 mm) <sup>2</sup> Beginning of pod filling <sup>3</sup>
74	704	About 40% of pods have reached final length (15–20 mm) <sup>2</sup>
75	705	About 50% of pods have reached final length (15–20 mm). Continuation of pod filling. <sup>2</sup> Main period of pod development. Continuation of pod filling <sup>3</sup>
77	707	About 70% of pods have reached final length (15–20 mm); Advanced pod filling. <sup>2</sup> Advanced pod filling <sup>3</sup>
79	709	Approx. all pods have reached final length (15–20 mm). Seeds filling the cavity of the majority of pods <sup>2,3</sup>

---

<sup>2</sup> This definition refers to determinate varieties

<sup>3</sup> This definition refers to indeterminate varieties



# Soybean Munger et al., 1997

## Phenological growth stages and BBCH-identification keys of the soybean

---

Code	Description
------	-------------

---

2- and 3digit

### Principal growth stage 8: Ripening of fruits and seeds

80	800	First pod ripe, beans final colour, dry and hard
81	801	Beginning of ripening; about 10% of pods are ripe, beans final colour, dry and hard. <sup>2</sup> Beginning of pod and seed ripening <sup>3</sup>
82	802	About 20% of pods are ripe; beans final colour, dry and hard <sup>2</sup>
83	803	About 30% of pods are ripe; beans final colour, dry and hard <sup>2</sup>
84	804	About 40% of pods are ripe; beans final colour, dry and hard <sup>2</sup>
85	805	Advanced ripening; about 50% of pods are ripe ; beans final colour, dry and hard. <sup>2</sup> Main period of pod and seed ripening <sup>3</sup>
86	806	About 60% of pods are ripe; beans final colour, dry and hard <sup>2</sup>
87	807	About 70% of pods are ripe; beans final colour, dry and hard <sup>2</sup>
88	808	About 80% of pods are ripe; beans final colour, dry and hard <sup>2</sup>
89	809	Full maturity: approx. all pods are ripe; beans final colour, dry and hard (= Harvest maturity). <sup>2</sup> Majority of pods are ripe; beans final colour, dry and hard <sup>3</sup>

---

2- and 3digit

### Principal growth stage 9: Senescence


91	901	About 10% of leaves discoloured or fallen
92	902	About 20% of leaves discoloured or fallen
93	903	About 30% of leaves discoloured or fallen
94	904	About 40% of leaves discoloured or fallen
95	905	About 50% of leaves discoloured or fallen
96	906	About 60% of leaves discoloured or fallen
97	907	Above ground parts of plants dead
99	909	Harvested product (seeds)

---

<sup>2</sup> This definition refers to determinate varieties

<sup>3</sup> This definition refers to indeterminate varieties

# Soybean



# Cotton Munger et al., 1998

## Phenological growth stages and BBCH-identification keys of the cotton

(*Gossypium hirsutum* L.)

---

Code	Description
<b>Principal growth stage 0: Germination</b>	
00	Dry seed
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from seed
06	Elongation of radicle
07	Hypocotyl with cotyledons breaking through seed coat
08	Hypocotyl with cotyledons growing towards soil surface
09	Emergence: hypocotyl with cotyledons breaking through soil surface ("crook stage")

---

### Principal growth stage 1: Leaf development (Main shoot)

10	Cotyledons completely unfolded <sup>1</sup>
11	First true leaf unfolded <sup>1</sup>
12	2nd true leaf unfolded <sup>1</sup>
13	3rd true leaf unfolded <sup>1</sup>
1 .	Stages continuous till . . .
19	9 or more true leaves unfolded; <sup>1</sup> no side shoots visible <sup>2</sup>

---

### Principal growth stage 2: Formation of side shoots<sup>3</sup>

21	First vegetative side shoot (2nd order) visible
22	2 vegetative side shoots (2nd order) visible
23	3 vegetative side shoots (2nd order) visible
2 .	Stages continuous till . . .
29	9 or more vegetative side shoots (2nd order) visible

---

<sup>1</sup> Leaves are counted from the cotyledon node (= node 0)

<sup>2</sup> Side shoot development may occur earlier, if there is a vegetative side shoot continue with principal growth stage 2. If there is a reproductive side shoot (fruiting branch) continue with the principal growth stage 5

<sup>3</sup> Vegetative side shoots are counted from the cotyledon node

# Cotton

Munger et al., 1998

## Phenological growth stages and BBCH-identification keys of the cotton

---

Code	Description
------	-------------

---

### Principal growth stage 3: Main stem elongation (Crop cover)

31	Beginning of crop cover: 10% of plants meet between rows
32	20% of plants meet between rows
33	30% of plants meet between rows
34	40% of plants meet between rows
35	50% of plants meet between rows
36	60% of plants meet between rows
37	70% of plants meet between rows
38	80% of plants meet between rows
39	Canopy closure: 90% of the plants meet between rows

---

### Principal growth stage 5: Inflorescence emergence (Main shoot)

51	First floral buds detectable ("pin-head square") <sup>4</sup>
52	First floral buds visible ("match-head square") <sup>4</sup>
55	Floral buds distinctly enlarged
59	Petals visible: floral buds still closed

---

### Principal growth stage 6: Flowering

60	First flowers opened (sporadically within the population)
61	Beginning of flowering ("Early bloom"): 5–6 blooms / 25 ft of row (= 5–6 blooms / 7,5 meter of row)
65	Full flowering: ("Mid bloom"): 11 and more blooms / 25 ft of row = 11 and more blooms / 7,5 meter of row
67	Flowering finishing: majority of flowers faded ("Late bloom")
69	End of flowering

---

### Principal growth stage 7: Development of fruits and seeds

71	About 10% of bolls have attained their final size
72	About 20% of bolls have attained their final size
73	About 30% of bolls have attained their final size
74	About 40% of bolls have attained their final size
75	About 50% of bolls have attained their final size
76	About 60% of bolls have attained their final size
77	About 70% of bolls have attained their final size
78	About 80% of bolls have attained their final size
79	About 90% of bolls have attained their final size

---

<sup>4</sup> "pin-head square" or "match-head square" is the first square which forms at the first fruiting position of the first fruiting branch

# Cotton

Munger et al., 1998

## Phenological growth stages and BBCH-identification keys of the cotton

---

Code	Description
------	-------------

---

### Principal growth stage 8: Ripening of fruits and seeds

80	Firstst open bolls on the first fruiting branches
81	Beginning of boll opening: about 10% of bolls open. Nodes Above White Flower (NAWF)
82	About 20% of bolls open
83	About 30% of bolls open. Nodes Above Cracked Boll (NACB)
84	About 40% of bolls open
85	About 50% of bolls open
86	About 60% of bolls open
87	About 70% of bolls open
88	About 80% of bolls open
89	About 90% of bolls open




---

### Principal growth stage 9: Senescence

91	About 10% of leaves discoloured or fallen
92	About 20% of leaves discoloured or fallen
93	About 30% of leaves discoloured or fallen
94	About 40% of leaves discoloured or fallen
95	About 50% of leaves discoloured or fallen
96	About 60% of leaves discoloured or fallen
97	Above ground parts of plant dead; plant dormant
99	Harvested product (bolls and seeds)

---

# Cotton



# Peanut Munger et al., 1998a

## Phenological growth stages and BBCH-identification keys of the peanut

(*Arachis hypogaea* L.)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Germination

00	Dry seed
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from seed
07	Hypocotyl with cotyledons breaking through seed coat
08	Hypocotyl reaches the soil surface; hypocotyl arch visible
09	Emergence: hypocotyl with cotyledons arising above soil surface ("cracking stage")

---

### Principal growth stage 1: Leaf development (Main shoot)

10	Cotyledons completely unfolded <sup>1</sup>
11	First true leaf (pinnate) unfolded <sup>1</sup>
12	2nd true leaf (pinnate) unfolded <sup>1</sup>
13	3rd true leaf (pinnate) unfolded <sup>1</sup>
1 .	Stages continuous till . . .
19	9 or more true leaves unfolded. <sup>1</sup> No side shoots visible <sup>2</sup>

---

### Principal growth stage 2: Formation of side shoots<sup>3</sup>

21	1st side shoot visible
22	2nd side shoot visible
23	3rd side shoot visible
2 .	Stages continuous till . . .
29	9 or more side shoots visible

---

### Principal growth stage 3: Main stem elongation (Crop cover)

31	Beginning of crop cover: 10% of plants meets between rows
32	20% of plants meets between rows
33	30% of plants meets between rows
34	40% of plants meets between rows
35	50% of plants meets between rows
36	60% of plants meets between rows
37	70% of plants meets between rows
38	80% of plants meets between rows
39	Crop cover complete: 90% of plants meets between rows

---

<sup>1</sup> Leaves are counted from the cotyledon node (= node 0)

<sup>2</sup> Side shoot development may occur earlier; in this case continue with principal growth stage 2

# Peanut

Munger et al., 1998a

## Phenological growth stages and BBCH-identification keys of the peanut

---

Code	Description
------	-------------

---

### Principal growth stage 5: Inflorescence emergence

51	First inflorescence buds visible
55	First individual flower buds visible
59	First flower petals visible. Flower buds still closed

---

### Principal growth stage 6: Flowering

61	Beginning of flowering
62	First carpophore pegs visible
63	Continuation of flowering
64	First carpophore pegs visibly elongated
65	Full flowering
66	First carpophore pegs penetrating the soil
67	Flowering declining <sup>4</sup>
68	Tip of first carpophore pegs growing horizontally in the soil
69	End of flowering <sup>4</sup>

---

### Principal growth stage 7: Development of fruits and seeds

71	Beginning of pod development: tip of first carpophore pegs swollen (at least twice the original diameter)
73	Continuation of pod development: beginning of pod filling: first pods have attained final size and are ripening
75	Main phase of pod development: continuation of pod filling
77	Advanced pod filling
79	Fresh seeds fill the cavity of the pods which have attained their final size

---

<sup>4</sup> Only for varieties with a determinate flowering period



# Peanut

Munger et al., 1998a

## Phenological growth stages and BBCH-identification keys of the peanut

---

Code	Description
<b>Principal growth stage 8: Ripening of fruits and seeds<sup>5</sup></b>	
81	Beginning of ripening: about 10% of pods developed to final size are ripe
82	About 20% of pods developed to final size are ripe
83	Continuation of ripening: about 30% of pods developed to final size are ripe
84	About 40% of pods developed to final size are ripe
85	Main phase of ripening: about 50% of pods developed to final size are ripe
86	About 60% of pods developed to final size are ripe
87	Advanced ripening: about 70% of pods developed to final size are ripe
88	About 80% of pods developed to final size are ripe
89	Full maturity: nearly all pods developed to final size are ripe

---


## Principal growth stage 9: Senescence

91	About 10% of above ground parts of plant dry
92	About 40% of above ground parts of plant dry
93	About 30% of above ground parts of plant dry
94	About 40% of above ground parts of plant dry
95	About 50% of above ground parts of plant dry
96	About 60% of above ground parts of plant dry
97	Above ground parts of plant dead
99	Harvested product

---

<sup>5</sup> Criteria of maturity: Pericarp hard, with distinct texture, can be split open easily; Testa (seed coat) dry, with cultivar-specific dark colour

# Peanut



# Hop

Rossbauer et al., 1995

## Phenological growth stages and BBCH-identification keys of hop

(*Humulus lupulus* L.)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Sprouting

00	Dormancy: rootstock without shoots (uncut)
01	Dormancy: rootstock without shoots (cut)
07	Rootstock with shoots (uncut)
08	Beginning of shoot-growth (rootstock cut)
09	Emergence: first shoots emerge at the soil surface

---

### Principal growth stage 1: Leaf development

11	First pair of leaves unfolded
12	2nd pair of leaves unfolded (beginning of twining)
13	3rd pair of leaves unfolded
1 .	Stages continuous till . . .
19	9 and more pairs of leaves unfolded

---

### Principal growth stage 2: Formation of side shoots

21	First pair of side shoots visible
22	2nd pair of side shoots visible
23	3rd pair of side shoots visible
2 .	Stages continuous till . . .
29	Nine and more pairs of side shoots visible (secondary side shoots occur)

---

### Principal growth stage 3: Elongation of bines

31	Bines have reached 10% of top wire height
32	Bines have reached 20% of top wire height
33	Bines have reached 30% of top wire height
3 .	Stages continuous till . . .
38	Plants have reached the top wire
39	End of bine growth

---

### Principal growth stage 5: Inflorescence emergence

51	Inflorescence buds visible
55	Inflorescence buds enlarged

---

# Hop

Rossbauer et al., 1995

## Phenological growth stages and BBCH-identification keys of hop

---

Code	Description
------	-------------

---

### Principal growth stage 6: Flowering

61	Beginning of flowering: about 10% of flowers open
62	About 20% of flowers open
63	About 30% of flowers open
64	About 40% of flowers open
65	Full flowering: about 50% of flowers open
66	About 60% of flowers open
67	About 70% of flowers open
68	About 80% of flowers open
69	End of flowering

---

### Principal growth stage 7: Development of cones

71	Beginning of cone development: 10% of inflorescences are cones
75	Cone development half way: all cones visible, cones soft, stigmas still present
79	Cone development complete: nearly all cones have reached full size

---

### Principal growth stage 8: Maturity of cones

81	Beginning of maturity: 10% of cones are compact
82	20% of cones are compact
83	30% of cones are compact
84	40% of cones are compact
85	Advanced maturity: 50% of cones are compact
86	60% of cones are compact
87	70% of cones are compact
88	80% of cones are compact
89	Cones ripe for picking: cones closed; lupulin golden; aroma potential fully developed




---

### Principal growth stage 9: Senescence, entry into dormancy



92	Overripeness: cones yellow-brown discoloured, aroma deterioration
97	Dormancy: leaves and stems dead

---

# Hop



# Hop



# Bulb vegetables Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of bulb vegetables

(Onion = *Allium cepa* L., leek = *Allium porrum* L., garlic = *Allium sativum* L., shallot = *Allium ascalonicum* auct. non L.)

---

Code	Description
------	-------------

---

2- and 3digit

### Principal growth stage 0: Germination

00	000	Dry seed, <sup>1</sup> dormant bulb <sup>2</sup>
01	000	Beginning of seed imbibition <sup>1</sup>
03	003	Seed imbibition complete <sup>1</sup>
05	005	Radicle emerged from seed. <sup>1</sup> Roots appearing <sup>2</sup>
07	007	Cotyledon breaking through seed coat <sup>1</sup>
09	009	Emergence: cotyledon breaks through soil surface. <sup>1</sup> Green shoot visible <sup>2</sup>
	010	Cotyledon visible as hook <sup>1</sup>
	011	Hook stage: hooked cotyledon green <sup>1</sup>
	012	Whip stage: cotyledon has whip-like form <sup>1</sup>

---

2- and 3digit

### Principal growth stage 1: Leaf development (Main shoot)

10	100	Advanced whip stage: whip begins to die off <sup>1</sup>
11	101	First leaf (> 3 cm) clearly visible
12	102	2nd leaf (> 3 cm) clearly visible
13	103	3rd leaf (> 3 cm)
1 .	10 .	Stages continuous till . . .
19	109	9 or more leaves clearly visible

---

### Principal growth stage 4: Development of harvestable vegetative plant parts

41	401	Leaf bases begin to thicken or extend
43	403	30% of the expected bulb or shaft diameter reached
45	405	50% of the expected bulb or shaft diameter reached
47	407	Bolting begins; in 10% of the plants leaves bent over <sup>3</sup> 70% of the expected shaft length and diameter reached <sup>4</sup>
48	408	Leaves bent over in 50% of plants <sup>3</sup>
49	409	Leaves dead, bulb top dry; dormancy <sup>3</sup> Growth complete; length and stem diameter typical for variety reached <sup>4</sup>

---

<sup>1</sup> Seed sown

<sup>2</sup> Onion sets, shallot and garlic

<sup>3</sup> For onions, garlic

<sup>4</sup> For leek

# Bulb vegetables Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of bulb vegetables

---

Code	Description
------	-------------

---

2- and 3digit

### Principal growth stage 5: Inflorescence emergence

51	501	Onion bulb begins to elongate
53	503	30% of the expected length of flower stem reached
55	505	Flower stem at full length; sheath closed
57	507	Sheath burst open
59	509	First flower petals visible; flowers still closed

---

2- and 3digit

### Principal growth stage 6: Flowering

60	600	First flowers open (sporadically)
61	601	Beginning of flowering: 10% of flowers open
62	602	20% of flowers open
63	603	30% of flowers open
64	604	40% of flowers open
65	605	Full flowering: 50% of flowers open
67	607	Flowering finishing: 70% of petals fallen or dry
69	609	End of flowering

---

2- and 3digit

### Principal growth stage 7: Development of fruit

71	701	First capsules formed
72	702	20% of capsules formed
73	703	30% of capsules formed
74	704	40% of capsules formed
75	705	50% of capsules formed
76	706	60% of capsules formed
77	707	70% of capsules formed
78	708	80% of capsules formed
79	709	Capsule development complete; seeds pale

---

2- and 3digit

### Principal growth stage 8: Ripening of fruit and seed

81	801	Beginning of ripening: 10% of capsules ripe
85	805	First capsules bursting
89	809	Fully ripe: seeds black and hard

---



# Bulb vegetables Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of bulb vegetables

---

Code	Description
------	-------------

---


2- and 3digit

### Principal growth stage 9: Senescence

92	902	Leaves and shoots beginning to discolour
95	905	50% of leaves yellow or dead
97	907	Plants or above ground parts dead
99	909	Harvested product (seeds)

---

# Bulb vegetables



# Root and stem vegetables Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of root and stem vegetables

(Carrot = *Daucus carota* L. ssp. *sativus*, celeriac = *Apium graveolens* L. var. *rapaceum* Gaud., kohlrabi = *Brassica oleracea* L. var. *gongylodes*, chicory = *Cichorium intybus* var. *foliosum*, radish = *Raphanus sativus* L. ssp., swede = *Brassica napus* L. ssp. *rapifera* Metzg., scorzonera = *Scorzonera hispanica* L.)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Germination

00	Dry seed
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from seed
07	Hypocotyl with cotyledons breaking through seed coat
09	Emergence: cotyledons break through soil surface

---

### Principal growth stage 1: Leaf development (Main shoot)

10	Cotyledons completely unfolded; growing point or true leaf initial visible
11	First true leaf unfolded
12	2nd true leaf unfolded
13	3rd true leaf unfolded
1 .	Stages continuous till . . .
19	9 or more true leaves unfolded

---

### Principal growth stage 4: Development of harvestable vegetative plant parts

41	Roots beginning to expand (diameter > 0,5 cm)
42	20% of the expected root diameter reached
43	30% of the expected root diameter reached
44	40% of the expected root diameter reached
45	50% of the expected root diameter reached
46	60% of the expected root diameter reached
47	70% of the expected root diameter reached
48	80% of the expected root diameter reached
49	Expansion complete; typical form and size of roots reached

---

### Principal growth stage 5: Inflorescence emergence

51	Main shoot begins to elongate
53	30% of the expected height of the main shoot reached
55	First individual flowers of main inflorescence visible (still closed)
57	First individual flowers of secondary inflorescences visible (still closed)
59	First flower petals visible; flowers still closed

---

# Root and stem vegetables Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of root and stem vegetables

---

Code	Description
------	-------------

---

### Principal growth stage 6: Flowering

60	First flowers open (sporadically)
61	Beginning of flowering: 10% of flowers open
62	20% of flowers open
63	30% of flowers open
64	40% of flowers open
65	Full flowering: 50% of flowers open
67	Flowering finishing: majority of petals fallen or dry
69	End of flowering

---

### Principal growth stage 7: Development of fruit

71	First fruits formed
72	20% of fruits have reached typical size
73	30% of fruits have reached typical size
74	40% of fruits have reached typical size
75	50% of fruits have reached typical size
76	60% of fruits have reached typical size
77	70% of fruits have reached typical size
78	80% of fruits have reached typical size
79	Fruits have reached typical size

---

### Principal growth stage 8: Ripening of fruit and seed

81	Beginning of ripening: 10% of fruits ripe, or 10% of seeds of typical colour, dry and hard
85	50% of the fruits ripe, or 50% of seeds of typical colour, dry and hard
89	Fully ripe: seeds on the whole plant of typical colour and hard


---

### Principal growth stage 9: Senescence

92	Leaves and shoots beginning to discolour
95	50% of leaves yellow or dead
97	Plants or above ground parts dead
99	Harvested product (seeds)

---

# Root and stem vegetables



# Leaf vegetables (forming heads) Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of leaf vegetables (forming heads)

(cabbage = *Brassica oleracea* L. var. *capitata* f. *alba* and *rubra*, chinese cabbage = *Brassica chinensis* L., lettuce = *Lactuca sativa* L. var. *capitata*, endive = *Cichorium endivia* L.)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Germination

00	Dry seed
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from seed
07	Hypocotyl with cotyledons breaking through seed coat
09	Emergence: cotyledons break through soil surface

---

### Principal growth stage 1: Leaf development (Main shoot)

10	Cotyledons completely unfolded; growing point or true leaf initial visible
11	First true leaf unfolded
12	2nd true leaf unfolded
13	3rd true leaf unfolded
1 .	Stages continuous till . . .
19	9 or more true leaves unfolded

---

### Principal growth stage 4: Development of harvestable vegetative plant parts

41	Heads begin to form: the two youngest leaves do not unfold
42	20% of the expected head size reached
43	30% of the expected head size reached
44	40% of the expected head size reached
45	50% of the expected head size reached
46	60% of the expected head size reached
47	70% of the expected head size reached
48	80% of the expected head size reached
49	Typical size, form and firmness of heads reached

---

### Principal growth stage 5: Inflorescence emergence

51	Main shoot inside head begins to elongate
53	30% of the expected height of the main shoot reached
55	First individual flowers of main inflorescence visible (still closed)
57	First individual flowers of secondary inflorescences visible (still closed)
59	First flower petals visible; flowers still closed

---

# Leaf vegetables (forming heads) Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of leaf vegetables (forming heads)

---

Code	Description
------	-------------

---

### Principal growth stage 6: Flowering

60	First flowers open (sporadically)
61	Beginning of flowering: 10% of flowers open
62	20% of flowers open
63	30% of flowers open
64	40% of flowers open
65	Full flowering: 50% of flowers open
67	Flowering finishing: majority of petals fallen or dry
69	End of flowering

---

### Principal growth stage 7: Development of fruit

71	First fruits formed
72	20% of fruits have reached typical size
73	30% of fruits have reached typical size
74	40% of fruits have reached typical size
75	50% of fruits have reached typical size
76	60% of fruits have reached typical size
77	70% of fruits have reached typical size
78	80% of fruits have reached typical size
79	Fruits have reached typical size

---

### Principal growth stage 8: Ripening of fruit and seed

81	Beginning of ripening: 10% of fruits ripe, or 10% of seeds of typical colour, dry and hard
82	20% of fruits ripe, or 20% of seeds of typical colour, dry and hard
83	30% of fruits ripe, or 30% of seeds of typical colour, dry and hard
84	40% of fruits ripe, or 40% of seeds of typical colour, dry and hard
85	50% of the fruits ripe, or 50% of seeds of typical colour, dry and hard
86	60% of fruits ripe, or 60% of seeds of typical colour, dry and hard
87	70% of fruits ripe, or 70% of seeds of typical colour, dry and hard
88	80% of fruits ripe, or 80% of seeds of typical colour, dry and hard
89	Fully ripe: seeds on the whole plant of typical colour and hard

---

# Leaf vegetables (forming heads) Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of leaf vegetables (forming heads)

---

Code	Description
------	-------------

---


### Principal growth stage 9: Senescence

92	Leaves and shoots beginning to discolour
95	50% of leaves yellow or dead
97	Plants dead
99	Harvested product (seeds)

---



# Leaf vegetables (forming heads)



# Leaf vegetables (not forming heads)

Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of leaf vegetables not forming heads

(spinach = *Spinacia oleracea* L., loosehead lettuce = *Lactuca sativa* L. var. *crispa*, kale = *Brassica oleracea* L. var. *sabellica*)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Germination

00	Dry seed
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from seed
07	Hypocotyl with cotyledons breaking through seed coat
09	Emergence: cotyledons break through soil surface

---

### Principal growth stage 1: Leaf development (Main shoot)

10	Cotyledons completely unfolded; growing point or true leaf initial visible
11	First true leaf unfolded
12	2nd true leaf unfolded
13	3rd true leaf unfolded
1 .	Stages continuous till . . .
19	9 or more true leaves unfolded

---

### Principal growth stage 3: Stem elongation of rosette growth

33	Leaf rosette has reached 30% of the expected diameter typical for the variety. <sup>1</sup> Main shoot has reached 30% of the expected height typical for the variety <sup>2</sup>
35	Leaf rosette has reached 50% of the expected diameter typical for the variety. <sup>1</sup> Main shoot has reached 50% of the expected height typical for the variety <sup>2</sup>
37	Leaf rosette has reached 70% of the expected diameter typical for the variety. <sup>1</sup> Main shoot has reached 70% of the expected height for the variety <sup>2</sup>
39	Rosette development completed <sup>1</sup> Main shoot has reached the height typical for the variety <sup>2</sup>

---

<sup>1</sup> For lettuce varieties without head, spinach and species with rosette-type growth

<sup>2</sup> For kale and species without rosette growth

# Leaf vegetables (not forming heads)

Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of leaf vegetables (not forming heads)

---

Code	Description
------	-------------

---

### Principal growth stage 4: Development of harvestable vegetative plant parts

41	10% of the leaf mass typical for the variety reached
42	20% of the leaf mass typical for the variety reached
43	30% of the leaf mass typical for the variety reached
44	40% of the leaf mass typical for the variety reached
45	50% of the leaf mass typical for the variety reached
46	60% of the leaf mass typical for the variety reached
47	70% of the leaf mass typical for the variety reached
48	80% of the leaf mass typical for the variety reached
49	Typical leaf mass reached

---

### Principal growth stage 5: Inflorescence emergence

51	Main shoot begins to elongate <sup>1</sup> Main inflorescence visible between uppermost leaves <sup>2</sup>
53	30% of the expected height of the main shoot reached
55	First individual flowers of main inflorescence visible (still closed)
59	First flower petals visible; flowers still closed

---

### Principal growth stage 6: Flowering

60	First flowers open (sporadically)
61	Beginning of flowering: 10% of flowers open
62	20% of flowers open
63	30% of flowers open
64	40% of flowers open
65	Full flowering: 50% of flowers open
67	Flowering finishing: majority of petals fallen or dry
69	End of flowering

---

<sup>1</sup> For letucce varieties without head, spinach and species with rosette-type growth

<sup>2</sup> For kale and species without rosette growth

# Leaf vegetables (not forming heads)

Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of leaf vegetables (not forming heads)

---

Code	Description
------	-------------

---

### Principal growth stage 7: Development of fruit

71	First fruits formed
72	20% of fruits have reached typical size
73	30% of fruits have reached typical size
74	40% of fruits have reached typical size
75	50% of fruits have reached typical size
76	60% of fruits have reached typical size
77	70% of fruits have reached typical size
78	80% of fruits have reached typical size
79	Fruits have reached typical size

---

### Principal growth stage 8: Ripening of fruit and seed

81	Beginning of ripening: 10% of fruits ripe, or 10% of seeds of typical colour, dry and hard
82	20% of fruits ripe, or 20% of seeds of typical colour, dry and hard
83	30% of fruits ripe, or 20% of seeds of typical colour, dry and hard
84	40% of fruits ripe, or 20% of seeds of typical colour, dry and hard
85	50% of fruits ripe, or 50% of seeds of typical colour, dry and hard
86	60% of fruits ripe, or 20% of seeds of typical colour, dry and hard
87	70% of fruits ripe, or 20% of seeds of typical colour, dry and hard
88	80% of fruits ripe, or 20% of seeds of typical colour, dry and hard
89	Fully ripe: seeds on the whole plant of typical colour and hard


---

### Principal growth stage 9: Senescence

92	Leaves and shoots beginning to discolor
95	50% of leaves yellow or dead
97	Plants dead
99	Harvested product (seeds)

---

# Leaf vegetables (not forming heads)



# Other brassica vegetables Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of other brassica vegetables

(Brussels sprout = *Brassica oleracea* L. var. *gemmifera* DC./Zenk., cauliflower = *Brassica oleracea* L. var. *botrytis*, broccoli = *Brassica oleracea* L. var. *italica* Plenck)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Germination

00	Dry seed
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from seed
07	Hypocotyl with cotyledons breaking through seed coat
09	Emergence: cotyledons break through soil surface

---

### Principal growth stage 1: Leaf development (Main shoot)

10	Cotyledons completely unfolded; growing point or true leaf initial visible
11	First true leaf unfolded
12	2nd true leaf unfolded
13	3rd true leaf unfolded
1 .	Stages continuous till . . .
19	9 or more true leaves unfolded

---

### Principal growth stage 2: Formation of side shoots

21	First side shoot visible <sup>1</sup>
22	2nd side shoot visible <sup>1</sup>
23	3rd side shoot visible <sup>1</sup>
2 .	Stages continuous till . . .
29	9 or more side shoots visible <sup>1</sup>

---

<sup>1</sup> For broccoli

# Other brassica vegetables Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of other brassica vegetables

---

Code	Description
<b>Principal growth stage 3: Stem elongation of rosette growth</b>	
31	Main shoot has reached 10% of the expected height typical for the variety <sup>2</sup>
32	Main shoot has reached 20% of the expected height typical for the variety <sup>2</sup>
33	Main shoot has reached 30% of the expected height typical for the variety <sup>2</sup>
34	Main shoot has reached 40% of the expected height typical for the variety <sup>2</sup>
35	Main shoot has reached 50% of the expected height typical for the variety <sup>2</sup>
36	Main shoot has reached 60% of the expected height typical for the variety <sup>2</sup>
37	Main shoot has reached 70% of the expected height typical for the variety <sup>2</sup>
38	Main shoot has reached 80% of the expected height typical for the variety <sup>2</sup>
39	Main shoot has reached the height typical for the variety <sup>2</sup>

---

## Principal growth stage 4: Development of harvestable vegetative plant parts

41	Lateral buds begin to develop <sup>2</sup>
	Cauliflower heads begin to form; width of growing tip > 1 cm <sup>3</sup>
43	First sprouts tightly closed <sup>2</sup>
	30% of the expected head diameter reached <sup>3</sup>
45	50% of the sprouts tightly closed <sup>2</sup>
	50% of the expected head diameter reached <sup>3</sup>
46	60% of the sprouts tightly closed <sup>2</sup>
	60% of the expected head diameter reached <sup>3</sup>
47	70% of the sprouts tightly closed <sup>2</sup>
	70% of the expected head diameter reached <sup>3</sup>
48	80% of the sprouts tightly closed <sup>2</sup>
	80% of the expected head diameter reached <sup>3</sup>
49	Sprouts below terminal bud tightly closed <sup>2</sup>
	Typical size and form reached; head tightly closed <sup>3</sup>

---

<sup>2</sup> For brussels sprout

<sup>3</sup> For cauliflower and broccoli

# Other brassica vegetables Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of other brassica vegetables

---

Code	Description
------	-------------

---

### Principal growth stage 5: Inflorescence emergence

51	Main inflorescence visible between uppermost leaves <sup>2</sup> Branches of inflorescence begin to elongate <sup>3</sup>
55	First individual flowers visible (still closed)
59	First flower petals visible; flowers still closed

---

### Principal growth stage 6: Flowering

60	First flowers open (sporadically)
61	Beginning of flowering: 10% of flowers open
62	20% of flowers open
63	30% of flowers open
64	40% of flowers open
65	Full flowering: 50% of flowers open
67	Flowering finishing: majority of petals fallen or dry
69	End of flowering

---

### Principal growth stage 7: Development of fruit

71	First fruits formed
72	20% of fruits have reached typical size
73	30% of fruits have reached typical size
74	40% of fruits have reached typical size
75	50% of fruits have reached typical size
76	60% of fruits have reached typical size
77	70% of fruits have reached typical size
78	80% of fruits have reached typical size
79	Fruits have reached typical size

---

### Principal growth stage 8: Ripening of fruit and seed

81	Beginning of ripening: 10% of fruits ripe
82	20% of fruits ripe
83	30% of fruits ripe
84	40% of fruits ripe
85	50% of fruits ripe
86	60% of fruits ripe
87	70% of fruits ripe
88	80% of fruits ripe
89	Fully ripe: seeds on the whole plant of typical color and hard

---

<sup>2</sup> For brussels sprout

<sup>3</sup> For cauliflower and broccoli



# Other brassica vegetables Feller et al., 1995 a

## Phenological growth stages and BBCH-identification keys of other brassica vegetables

---

Code	Description
------	-------------

---

### Principal growth stage 9: Senescence

92	Leaves and shoots beginning to discolour
95	50% of leaves yellow or dead
97	Plants dead
99	Harvested product (seeds)

---

# Cucurbits Feller et al., 1995 b

## Phenological growth stages and BBCH-identification keys of cucurbits

(Cucumber = *Cucumis sativus* L., melon = *Cucumis melo* L., pumpkin, marrow, squash = *Cucurbita pepo* L., calabash = *Cucurbita pepo* L. var. *giromontiina* Alef./Greb, water-melon = *Citrullus* var. *vulgaris* Schad.)

---

Code	Description
------	-------------

---

2 -and 3digit

### Principal growth stage 0: Germination

00	000	Dry seed
01	001	Beginning of seed imbibition
03	003	Seed imbibition complete
05	005	Radicle emerged from seed
07	007	Hypocotyl with cotyledons breaking through seed coat
09	009	Emergence: cotyledons break through soil surface

---

2- and 3digit

### Principal growth stage 1: Leaf development

10	100	Cotyledons completely unfolded
11	101	First true leaf on main stem fully unfolded
12	102	2nd true leaf on main stem unfolded
13	103	3rd true leaf on main stem unfolded
1 .	10 .	Stages continuous till . . .
19	109	9 or more leaves on main stem unfolded (2digit) 9th leaf unfolded on main stem (3digit)
-	110	10th leaf on main stem unfolded
-	11 .	Stages continuous till . . .
-	119	19th leaf on main stem unfolded

2 -and 3digit

### Principal growth stage 2: Formation of side shoots

21	201	First primary side shoot visible
22	202	2nd primary side shoot visible
2 .	20 .	Stages continuous till . . .
29	209	9 or more primary side shoots visible
-	221	First secondary side shoot visible
-	22 .	Stages continuous till . . .
-	229	9th secondary side shoot visible
-	231	First tertiary side shoot visible

# Cucurbits Feller et al., 1995 b

## Phenological growth stages and BBCH-identification keys of cucurbits

---

Code	Description
------	-------------

---

2- and 3digit

### Principal growth stage 5: Inflorescence emergence

51	501	First flower initial with elongated ovary visible on main stem
52	502	2nd flower initial with elongated ovary visible on main stem
53	503	3rd flower initial with elongated ovary visible on main stem
5 .	50 .	Stages continuous till . . .
59	509	9 or more flower initials with elongated ovary already visible on main stem
–	510	10 or more flower initials with elongated ovary already visible on main stem
–	51 .	Stages continuous till . . .
–	519	19 ore more flower initials with elongated ovary already visible on main stem
–	521	First flower initial visible on a secondary side shoot
–	531	First flower initial visible on a tertiary side shoot

2 -and 3digit

### Principal growth stage 6: Flowering

61	601	First flower open on main stem
62	602	2nd flower open on main stem
63	603	3rd flower open on main stem
6 .	60 .	Stages continuous till . . .
69	609	9th flower open on main stem or 9 flowers on main stem already open
–	610	10th flower open on main stem or 10 flowers on main stem already open
–	61 .	Stages continuous till . . .
–	619	19th flower open on man stem ore more than 19 flowers on main stem already open
–	621	First flower on secondary side shoot open
–	631	First flower on tertiary side shoot open

# Cucurbits

Feller et al., 1995 b

## Phenological growth stages and BBCH-identification keys of cucurbits

---

Code	Description
------	-------------

---

2- and 3digit

### Principal growth stage 7: Development of fruit

71	701	First fruit on main stem has reached typical size and form
72	702	2nd fruit on main stem has reached typical size and form
73	703	3rd fruit on main stem has reached typical size and form
7.	70.	Stages continuous till . . .
79	709	9 or more fruits on main stem has reached typical size and form
–	721	First fruit on a secondary side shoot has reached typical size and form
–	731	First fruit on a tertiary side shoot has reached typical size and form

2 -and 3digit

### Principal growth stage 8: Ripening of fruit and seed

81	801	10% of fruits show typical fully ripe colour
82	802	20% of fruits show typical fully ripe colour
83	803	30% of fruits show typical fully ripe colour
84	804	40% of fruits show typical fully ripe colour
85	805	50% of fruits show typical fully ripe colour
86	806	60% of fruits show typical fully ripe colour
87	807	70% of fruits show typical fully ripe colour
88	808	80% of fruits show typical fully ripe colour
89	809	Fully ripe: fruits have typical fully ripe colour

---


2- and 3digit

### Principal growth stage 9: Senescence

97	907	Plants dead
99	909	Harvested product (seeds)

---

# Cucurbits



# Solanaceous fruits Feller et al., 1995 b

## Phenological growth stages and BBCH-identification keys of solanaceous fruits

(Tomato = *Lycopersicon esculentum* Mill., aubergine = *Solanum melongena* L., paprika = *Capsicum annuum* L)

---

Code	Description
------	-------------

---

2 -and 3digit

### Principal growth stage 0: Germination

00	000	Dry seeds
01	001	Beginning of seed imbibition
03	003	Seed imbibition complete
05	005	Radicle emerged from seed
07	007	Hypocotyl with cotyledons breaking through seed coat
09	009	Emergence: coryledons break through soil surface

---

2- and 3digit

### Principal growth stage 1: Leaf development

10	100	Cotyledons completely unfolded
11	101	First true leaf on main shoot fully unfolded
12	102	2nd leaf on main shoot unfolded
13	103	3rd leaf on main shoot unfolded
1 .	10 .	Stages continuous till . . .
19	109	9 or more leaves on main shoot unfolded

---

2 -and 3digit

### Principal growth stage 2: Formation of side shoots<sup>1</sup>

21	201	First primary apical side shoot visible
22	202	2nd primary apical side shoot visible
2 .	20 .	Stages continuous till . . .
29	209	9 or more apical primary side shoots visible
-	221	First secondary apical side shoot visible
-	22 .	Stages continuous till . . .
-	229	9th secondary apical side shoot visible
-	231	First tertiary apical side shoot visible
-	23 .	Stages continuous till . . .
-	2NX	Xth apical side shoot of the Nth order visible

---

<sup>1</sup> For tomatoes with determinate stem growth, paprika and aubergines. In tomatoes with indeterminate stem growth and only one sympodial branch at the corresponding axis, the apical side shoot formation occurs concurrently with the emergence of the inflorescence (Principal growth stage 5), so that the coding within principal growth stage 2 is not necessary

# Solanaceous fruits Feller et al., 1995 b

## Phenological growth stages and BBCH-identification keys of solanaceous fruits

---

Code            Description

---

2- and 3digit

### Principal growth stage 5: Inflorescence emergence

51	501	First inflorescence visible (first bud erect) <sup>2</sup> First flower bud visible <sup>3</sup>
52	502	2nd inflorescence visible (first bud erect) <sup>2</sup> 2nd flower bud visible <sup>3</sup>
53	503	3th inflorescence visible (first bud erect) <sup>2</sup> 3th flower bud visible <sup>3</sup>
5 .	50 .	Stages continuous till . . .
59	509	9 or more inflorescences visible (2digit), 9th inflorescence visible(first bud erect) (3digit) <sup>2</sup> 9 or more flower buds already visible (2digit), 9th flower bud visible (3digit) <sup>3</sup>
–	510	10th inflorescence visible (first bud erect) <sup>2</sup> 10th flower bud visible <sup>3</sup>
–	51 .	Stages continuous till . . .
–	519	19th inflorescence visible (first bud erect) <sup>2</sup> 19th flower bud visible <sup>3</sup>

---

2 -and 3digit

### Principal growth stage 6: Flowering

61	601	First inflorescence: first flower open <sup>2</sup> First flower open <sup>3</sup>
62	602	2nd inflorescence: first flower open <sup>2</sup> 2nd flower open <sup>3</sup>
63	603	3rd inflorescence: first flower open <sup>2</sup> 3rd flower open <sup>3</sup>
6 .	60 .	Stages continuous till . . .
69	609	9 or more inflorescences with open flowers (2digit) 9th inflorescence: first flower open (3digit) <sup>2</sup> 9 or more flowers already open (2digit) 9th flower open (3digit) <sup>3</sup>
–	610	10th inflorescence: first flower open <sup>2</sup> 10th flower open <sup>3</sup>
–	61 .	Stages continuous till . . .
–	619	19th inflorescence: first flower open <sup>2</sup> 19th flower open <sup>3</sup>

---

<sup>2</sup> For tomato

<sup>3</sup> For paprika and aubergine

# Solanaceous fruits Feller et al., 1995 b

## Phenological growth stages and BBCH-identification keys of solanaceous fruits

---

Code	Description
------	-------------

---

2- and 3digit

### Principal growth stage 7: Development of fruit

71	701	First fruit cluster: first fruit has reached typical size <sup>2</sup> First fruit has reached typical size and form <sup>3</sup>
72	702	2nd fruit cluster: first fruit has reached typical size <sup>2</sup> 2nd fruit has reached typical size and form <sup>3</sup>
73	703	3rd fruit cluster: first fruit has reached typical size <sup>2</sup> 3rd fruit has reached typical size and form <sup>3</sup>
7.	70.	Stages continuous till . . .
79	709	9 or more fruit clusters with fruits of typical size (2digit) 9th fruit cluster: first fruit has reached typical size (3digit) <sup>2</sup> 9 or more fruits have reached typical size and form (2digit) 9th fruit has reached typical size and form (3digit) <sup>3</sup>
-	710	10th fruit cluster: first fruit has reached typical form and size <sup>2</sup> 10th fruit has reached typical form and size <sup>3</sup>
-	71.	Stages continuous till . . . 19th fruit has reached typical form and size <sup>3</sup>
-	719	19th fruit cluster: first fruit has reached typical form and size <sup>2</sup>

---

2 -and 3digit

### Principal growth stage 8: Ripening of fruit and seed

81	801	10% of fruits show typical fully ripe colour
82	802	20% of fruits show typical fully ripe colour
83	803	30% of fruits show typical fully ripe colour
84	804	40% of fruits show typical fully ripe colour
85	805	50% of fruits show typical fully ripe colour
86	806	60% of fruits show typical fully ripe colour
87	807	70% of fruits show typical fully ripe colour
88	808	80% of fruits show typical fully ripe colour
89	809	Fully ripe: fruits have typical fully ripe colour <sup>3</sup>

---

2- and 3digit

### Principal growth stage 9: Senescence

97	907	Plants dead
99	909	Harvested product


---

<sup>2</sup> For tomato

<sup>3</sup> For paprika and aubergine



# Solanaceae fruits



# Pea

Weber and Bleiholder, 1990; Feller et al., 1995 b

## Phenological growth stages and BBCH-identification keys of pea

(*Pisum sativum* L.)

---

Code	Description
------	-------------

---

### Principal growth stage 0: Germination

00	Dry seed
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from seed
07	Shoot breaking through seed coat
08	Shoot growing towards soil surface; hypocotyl arch visible
09	Emergence: shoot breaks through soil surface ("cracking stage")

---

### Principal growth stage 1: Leaf development

10	Pair of scale leaves visible
11	First true leaf (with stipules) unfolded or first tendrils developed
12	2 leaves (with stipules) unfolded or 2 tendrils developed
13	3 leaves (with stipules) unfolded or 3 tendrils developed
1 .	Stages continuous till . . .
19	9 or more leaves (with stipules) unfolded or 9 or more tendrils developed

---

### Principal growth stage 3: Stem elongation (Main shoot)

30	Beginning of stem elongation
31	1 visibly extended internode <sup>1</sup>
32	2 visibly extended internodes <sup>1</sup>
33	3 visibly extended internodes <sup>1</sup>
3 .	Stages continuous till . . .
39	9 or more visibly extended internodes <sup>1</sup>

---

### Principal growth stage 5: Inflorescence emergence

51	First flower buds visible outside leaves
55	First separated flower buds visible outside leaves but still closed
59	First petals visible, flowers still closed

---

<sup>1</sup> The first internode extends from the scale leaf node to the first true leaf node

# Pea

Weber and Bleiholder, 1990; Feller et al., 1995 b

## Phenological growth stages and BBCH-identification keys of pea

---

Code	Description
------	-------------

---

**Principal growth stage 6: Flowering**

60	First flowers open (sporadically within the population)
61	Beginning of flowering: 10% of flowers open
62	20% of flowers open
63	30% of flowers open
64	40% of flowers open
65	Full flowering: 50% of flowers open
67	Flowering declining
69	End of flowering

---

**Principal growth stage 7: Development of fruit**

71	10% of pods have reached typical length; juice exudes if pressed
72	20% of pods have reached typical length; juice exudes if pressed
73	30% of pods have reached typical length; juice exudes if pressed. Tenderometer value: 80 TE
74	40% of pods have reached typical length; juice exudes if pressed. Tenderometer value: 95 TE
75	50% of pods have reached typical length; juice exudes if pressed. Tenderometer value: 105 TE
76	60% of pods have reached typical length; juice exudes if pressed. Tenderometer value: 115 TE
77	70% of pods have reached typical length. Tenderometer value: 130 TE
79	Pods have reached typical size (green ripe); peas fully formed

---

**Principal growth stage 8: Ripening of fruit and seed**

81	10% of pods ripe, seeds final colour, dry and hard
82	20% of pods ripe, seeds final colour, dry and hard
83	30% of pods ripe, seeds final colour, dry and hard
84	40% of pods ripe, seeds final colour, dry and hard
85	50% of pods ripe, seeds final colour, dry and hard
86	60% of pods ripe, seeds final colour, dry and hard
87	70% of pods ripe, seeds final colour, dry and hard
88	80% of pods ripe, seeds final colour, dry and hard
89	Fully ripe: all pods dry and brown. Seeds dry and hard (dry ripe)


---

**Principal growth stage 9: Senescence**

97	Plants dead and dry
99	Harvested product

---

# Pea



# Bean

Feller et al., 1995 b

## Phenological growth stages and BBCH-identification keys of Bean

(*Phaseolus vulgaris* var. *nanus* L.),

---

Code	Description
------	-------------

---

### Principal growth stage 0: Germination

00	Dry seed
01	Beginning of seed imbibition
03	Seed imbibition complete
05	Radicle emerged from seed
07	Hypocotyl with cotyledons breaking through seed coat
08	Hypocotyl reaches the soil surface; hypocotyl arch visible
09	Emergence: hypocotyl with cotyledons break through soil surface ("cracking stage")

---

### Principal growth stage 1: Leaf development

10	Cotyledons completely unfolded
12	2 full leaves (first leaf pair unfolded)
13	3rd true leaf (first trifoliolate leaf) unfolded
1 .	Stages continuous till . . .
19	9 or more leaves (2 full leaves, 7 or more trifoliolate) unfolded

---

### Principal growth stage 2: Formation of side shoots

21	First side shoot visible
22	2nd side shoot visible
23	3rd side shoot visible
2 .	Stages continuous till . . .
29	9 or more side shoots visible

---

### Principal growth stage 5: Inflorescence emergence

51	First flower buds visible
55	First flower buds enlarged
59	First petals visible, flowers still closed

---

# Bean

Feller et al., 1995 b

## Phenological growth stages and BBCH-identification keys of Bean

Code	Description
<b>Principal growth stage 6: Flowering</b>	
60	First flowers open (sporadically within the population)
61	Beginning of flowering: 10% of flowers open <sup>1</sup> Beginning of flowering <sup>2</sup>
62	20% of flowers open <sup>1</sup>
63	30% of flowers open <sup>1</sup>
64	40% of flowers open <sup>1</sup>
65	Full flowering: 50% of flowers open <sup>1</sup> Main flowering period <sup>2</sup>
67	Flowering finishing: majority of petals fallen or dry <sup>1</sup>
69	End of flowering: first pods visible <sup>1</sup>
<b>Principal growth stage 7: Development of fruit</b>	
71	10% of pods have reached typical length <sup>1</sup> Beginning of pod development <sup>2</sup>
72	20% of pods have reached typical length <sup>1</sup>
73	30% of pods have reached typical length <sup>1</sup>
74	40% of pods have reached typical length <sup>1</sup>
75	50% of pods have reached typical length, beans beginning to fill out <sup>1</sup> Main pod development period <sup>2</sup>
76	60% of pods have reached typical length <sup>1</sup>
77	70% of pods have reached typical length, pods still break cleanly <sup>1</sup>
78	80% of pods have reached typical length <sup>1</sup>
79	Pods: individual beans easily visible <sup>1</sup>
<b>Principal growth stage 8: Ripening of fruit and seed</b>	
81	10% of pods ripe (beans hard) <sup>1</sup> Seeds beginning to mature <sup>2</sup>
82	20% of pods ripe (beans hard) <sup>1</sup>
83	30% of pods ripe (beans hard) <sup>1</sup>
84	40% of pods ripe (beans hard) <sup>1</sup>
85	50% of pods ripe (beans hard) <sup>1</sup> Main period of ripening <sup>2</sup>
86	60% of pods ripe (beans hard) <sup>1</sup>
87	70% of pods ripe (beans hard) <sup>1</sup>
88	80% of pods ripe (beans hard) <sup>1</sup>
89	Fully ripe: pods ripe (beans hard) <sup>1</sup>

1 For varieties with limited flowering period

2 For varieties in which the flowering period is not limited

# Bean

Feller et al., 1995 b

## Phenological growth stages and BBCH-identification keys of Bean

---

Code	Description
------	-------------


---

### Principal growth stage 9: Senescence

97	Plants dead
99	Harvested product

---

# Bean





# Weed species Hess et al., 1997

## Phenological growth stages and BBCH-identification keys of weed species

---

D = Dicotyledons,  
G = Gramineae,  
M = Monocotyledons,  
P = Perennial plants,  
V = Development from vegetative parts or propagated organs.

No code letter is used if the description applies to all groups of plants.

---

Code	Description
------	-------------

---

### Principal growth stage 0: Germination, sprouting, bud development

00	Dry seed
V	Perennating or reproductive organs during the resting period (tuber, rhizome, bulb, stolon)
P	Winter dormancy or resting period
01	Beginning of seed imbibition
P, V	Beginning of bud swelling
03	Seed imbibition complete
P, V	End of bud swelling
05	Radicle (root) emerged from seed
V	Perennating or reproductive organs forming roots
06	Elongation of radicle, formation of root hairs and/or lateral roots
07	G Coleoptile emerged from caryopsis
D, M	Hypocotyl with cotyledons or shoot breaking through seed coat
P, V	Beginning of sprouting or bud breaking
08	D Hypocotyl with cotyledons or shoot growing towards soil surface
V	Shoot growing towards soil surface
09	G Emergence: Coleoptile breaks through soil surface
D, M	Emergence: Cotyledons break through soil surface (except hypogeal germination);
V	Emergence: Shoot/Leaf breaks through soil surface
P	Buds show green tips

---

### Principal growth stage 1: Leaf development (main shoot)

10	G, M First true leaf emerged from coleoptile
D	Cotyledons completely unfolded
P	First leaves separated
11	First true leaf, leaf pair or whorl unfolded
P	First leaves unfolded
12	2 true leaves, leaf pairs or whorls unfolded
13	3 true leaves, leaf pairs or whorls unfolded
1 .	Stages continuous till ...
19	9 or more true leaves, leaf pairs or whorls unfolded

---

# Weed species Hess et al., 1997

## Phenological growth stages and BBCH-identification keys of weed species

---

Code	Description
------	-------------

---

### Principal growth stage 2 : Formation of side shoots / tillering

21	First side shoot visible
G	First tiller visible
22	2 side shoots visible
G	2 tillers visible
23	3 side shoots visible
G	3 tillers visible
2 .	Stages continuous till ...
29	9 or more side shoots visible
G	9 or more tillers visible

---

### Principal growth stage 3: Stem elongation /shoot development (main shoot)

30	Beginning of stem elongation
G	Beginning of shooting
31	1 visibly extended internode
G	1 node stage
32	2 visibly extended internode;
G	2 node stage
33	3 visibly extended internode
G	3 node stage
3 .	Stages continuous till ...
39	9 or more visibly extended internodes
G	9 or more nodes

---

### Principal growth stage 4: vegetative propagation / booting (main shoot)

40	V	Vegetative reproductive organs begin to develop (rhizomes, stolons, tubers, runners, bulbs)
41	G	Flag leaf sheath extending
42	V	First young plant visible
43	G	Flag leaf sheath just visibly swollen (mid-boot)
45	G	Flag leaf sheath swollen (late-boot)
47	G	Flag leaf sheath opening
49	V	Constant new development of young plants; vegetative reproductive organs reach final size
	G	First awns visible

---

# Weed species Hess et al., 1997

## Phenological growth stages and BBCH-identification keys of weed species

---

Code	Description
------	-------------

---

### Principal growth stage 5: Inflorescence emergence (main shoot) / heading

51	Inflorescence or flower buds visible
G	Beginning of heading
55	First individual flowers visible (still closed)
G	Half of inflorescence emerged (middle of heading)
59	First flower petals visible (in petalled forms)
G	Inflorescence fully emerged (end of heading)

---

### Principal growth stage 6: Flowering (main shoot)

60	First flowers open (sporadically)
61	Beginning of flowering: 10% of flowers open
63	30% of flowers open
65	Full flowering: 50% of flowers open, first petals may be fallen
67	Flowering finishing: majority of petals fallen or dry
69	End of flowering: fruit set visible

---

### Principal growth stage 7: Development of fruit

71	Fruits begin to develop
G	Caryopsis watery ripe
79	Nearly all fruits have reached final size normal for the species and location

---

### Principal growth stage 8: Ripening or maturity of fruit and seed

81	Beginning of ripening or fruit coloration
89	Fully ripe

---


### Principal growth stage 9: Senescence, beginning of dormancy

97	End of leaf fall, plants or above ground parts dead or dormant;
P, V	Plant resting or dormant


---

# Weed species


*Agropyron repens* (L.) P. Beauv.




*Cynodon dactylon* (L.) Pers.



*Cyperus rotundus* L.




*Galium aparine* L.




# Weed species


*Polygonum convolvulus* L.



*Ranunculus repens* L.




*Stellaria media* (L.) Vill.




# Weed species


*Solanum nigrum* L.



*Taraxacum officinale* Wiggers



*Veronica hederifolia* L.



## **Cited References • Zitierte Literatur • Bibliografía citada • Bibliographie citée**

- AGUSTI, M., S. ZARAGOZA, H. BLEIHOLDER, L. BUHR, H. HACK, R. KLOSE y R. STAUSS, 1995:** Escala BBCH para la descripción de los estadios fenológicos del desarrollo de los agrios (Gén. Citrus). Levante Agrícola 3, 189-199.
- ARCILA-PULGARÍN J., L. BUHR, H. BLEIHOLDER, H. HACK, U. MEIER and H. WICKE, 2002:** Application of the "Extended BBCH - Scale" for the description of the growth stages of coffee (*Coffea sp.*): Ann app.Biol., 141, 19-27.
- FELLER, C., H. BLEIHOLDER, L. BUHR, H. HACK, M. HESS, R. KLOSE, U. MEIER, R. STAUSS, T. VAN DEN BOOM und E. WEBER, 1995a:** Phänologische Entwicklungsstadien von Gemüsepflanzen: I. Zwiebel-, Wurzel-, Knollen- und Blattgemüse. Nachrichtenbl. Deut. Pflanzenschutzd. 47, 193-206.
- FELLER, C., H. BLEIHOLDER, L. BUHR, H. HACK, M. HESS, R. KLOSE, U. MEIER, R. STAUSS, T. VAN DEN BOOM und E. WEBER, 1995b:** Phänologische Entwicklungsstadien von Gemüsepflanzen: II. Fruchtgemüse und Hülsenfrüchte. Nachrichtenbl. Deut. Pflanzenschutzd. 47, 217-232.
- GONZALES, R., C. RUIZ-SILVERA, L. BUHR, H. BLEIHOLDER, H. HACK, U. MEIER and H. WICKE, 2002:** Proposal for codification of the phenological cycle of edible Musaceae. Proceedings XV Reunión International ACORBAT Meeting 2002 Cartagena/Kolumbien. 412-417
- HACK, H., H. BLEIHOLDER, L. BUHR, U. MEIER, U. SCHNOCK-FRICKE, E. WEBER und A. WITZENBERGER, 1992:** Einheitliche Codierung der phänologischen Entwicklungsstadien mono- und dikotyler Pflanzen – Erweiterte BBCH-Skala, Allgemein -. Nachrichtenbl. Deut. Pflanzenschutzd. 44, 265-270.
- HACK, H., H. GALL, TH. KLEMKE, R. KLOSE, U. MEIER, R. STAUSS und A. WITZENBERGER, 1993:** Phänologische Entwicklungsstadien der Kartoffel (*Solanum tuberosum* L.). Codierung und Beschreibung nach der erweiterten BBCH-Skala mit Abbildungen. Nachrichtenbl. Deut. Pflanzenschutzd. 45, 11-19.
- HESS, M., G. BARRALIS, H. BLEIHOLDER, L. BUHR, TH. EGGERS, H. HACK und R. STAUSS, 1997:** Use of the extended BBCH-scale - general for the description of the growth stages of mono- and dicotyledonous weed species. Weed Research, 37, 433-441
- LANCASHIRE, P. D., H. BLEIHOLDER, P. LANGELÜDDECKE, R. STAUSS, T. VAN DEN BOOM, E. WEBER und A. WITZENBERGER, 1991:** An uniform decimal code for growth stages of crops and weeds. Ann. appl. Biol. 119, 561-601.
- LORENZ, D. H., K. W. EICHHORN, H. BLEIHOLDER, R. KLOSE, U. MEIER und E. WEBER, 1994:** Phänologische Entwicklungsstadien der Weinrebe (*Vitis vinifera* L. ssp. *vinifera*). Vitic. Enol. Sci. 49, 66-70.
- MEIER, U., L. BACHMANN, H. BUHTZ, H. HACK, R. KLOSE, B. MÄRLÄNDER und E. WEBER, 1993:** Phänologische Entwicklungsstadien der Beta-Rüben (*Beta vulgaris* L. ssp.). Codierung und Beschreibung nach der erweiterten BBCH-Skala (mit Abbildungen). Nachrichtenbl. Deut. Pflanzenschutzd. 45, 37-41.

- MEIER, U., H. GRAF, H. HACK, M. HESS, W. KENNEL, R. KLOSE, D. MAPPES, D. SEIPP, R. STAUSS, J. STREIF und T. VAN DEN BOOM, 1994:** Phänologische Entwicklungsstadien des Kernobstes (*Malus domestica* Borkh. und *Pyrus communis* L.), des Steinobstes (Prunus-Arten), der Johannisbeere (Ribes-Arten) und der Erdbeere (*Fragaria x ananassa* Duch.). Nachrichtenbl. Deut. Pflanzenschutzd. 46, 141-153.
- MUNGER, P., H. BLEIHOLDER, H. HACK, M. HESS, R. STAUSS, T. VAN DEN BOOM and E. WEBER, 1997:** Phenological Growth Stages of the Soybean Plant (*Glycine max* (L.) MERR.) – Codification and Description according to the General BBCH Scale – with Figures. Journal of Agronomy and Crop Science 179, 209 - 217.
- MUNGER, P., H. BLEIHOLDER, H. HACK, M. HESS, R. STAUSS, T. VAN DEN BOOM and E. WEBER, 1998:** Phenological Growth Stages of the Cotton plant (*Gossypium hirsutum* L.) Codification and Description according to the BBCH Scale – with figures. Journal of Agronomy and Crop Science 180, 143 - 149.
- MUNGER, P., H. BLEIHOLDER, H. HACK, M. HESS, R. STAUSS, T. VAN DEN BOOM and E. WEBER, 1998:** Phenological Growth Stages of the Peanut plant (*Arachis hypogaea* L.) Codification and Description according to the BBCH Scale – with figures. Journal of Agronomy and Crop Science 180, 101 - 107.
- SANZ-CORTÉS F., J. MARTÍNEZ-CALVO, M.L. BADENES, H. BLEIHOLDER, H. HACK, G. LIÁCER, U. MEIER, 2002:** Phenological growth stages of olive trees (*Olea europaea* L.). Ann. Appl. Biol. 140, 151-157.
- ROSSBAUER, G., L. BUHR, H. HACK, S. HAUPTMANN, R. KLOSE, U. MEIER, R. STAUSS und E. WEBER, 1995:** Phänologische Entwicklungsstadien von Kultur-Hopfen (*Humulus lupulus* L.). 249-253.
- WEBER, E. und H. BLEIHOLDER, 1990:** Erläuterungen zu den BBCH-Dezimal-Codes für die Entwicklungsstadien von Mais, Raps, Faba-Bohne, Sonnenblume und Erbse - mit Abbildungen. Gesunde Pflanzen 42, 308-321.
- WITZENBERGER, A., H. HACK und T. VAN DEN BOOM, 1989:** Erläuterungen zum BBCH-Dezimal-Code für die Entwicklungsstadien des Getreides – mit Abbildungen. Gesunde Pflanzen 41, 384-388.
- ZADOKS, J. C., T. T. CHANG, and C. F. KONZAK, 1974:** A decimal code for the growth stages of cereals. Weed Research 14, 415-421 and Eucarpia Bulletin No. 7, 49-52.



# Additional References

## Weiterführende Literatur

### Bibliografía adicional

### Bibliographie additionel

**ADAS; J., 1976:** Black Currant Early Growth Stage Key No. 71. Ministry of Agriculture, Fisheries and Food (GB).

**ANONYMOUS, 1976:** Manual of plant growth stage and disease assessment keys. Field bean growth stages key No. 4.1. Ministry of Agriculture, Fisheries and Food, Harpenden, UK.

**ANONYMOUS, 1984:** EPPO Crop Growth Stage Keys, Echelles OEPP des stades des développement des plantes cultivées - Grapevine/Vigne. OEPP/EPPO Bulletin **14**, 295-298.

**ANONYMOUS, 1990:** EPPO Crop Growth Stage Keys - Soybean -. EPPO Bulletin **20**, 645-650.

**ARCILA, P. J., and B. CHAVES-CORDOBA, 1995:** Desarrollo foliar del cafeto en tres densidades de siembra. Cenicafé 46(1): 5-20.

**ARCILA, P. J., A. JARAMILLO-ROBLEDO, J. V. BALDION, and A. BUSTILLO-PARDEY, 1993:** La floración del cafeto y su relación con el control de la broca. Avances Técnicos Cenicafé. 193:1-6

**ARCILA P. J., 1988:** Aspectos fisiológicos de la producción de café. In: Comité Departamental de Cafeteros de Caldas. Centro Nacional de Investigaciones de Café, Cenicafé. Tecnología del cultivo del café. Chinchiná, Cenicafé. Pp: 59-111.

**AUTORENKOLLEKTIV, 1978:** Methodische Anleitung zur Durchführung von Versuchen mit Pflanzenschutzmitteln und Mitteln zur Steuerung biologischer Prozesse. Institut für Pflanzenschutzforschung, AdL DDR, Kleinmachnow.

**BACHMANN, L., 1984:** Markante Wachstumsstadien der Zuckerrübe zur Datenerfassung. Feldwirtschaft **25**, 407-409.

**BACHMANN, L., 1986:** Zur Einführung eines zweiziffrigen Codes zur Kennzeichnung der Wachstumsstadien bei Zuckerrüben. Feldwirtschaft **27**, 392-394.

**BÄTZ, W., U. MEIER, W. RADTKE, B. SCHÖBER, L. SEIDWITZ und J. STEINBERGER, 1980:** Entwicklungsstadien der Kartoffel. Biologische Bundesanstalt für Land- und Forstwirtschaft. Merkblatt **27/5**.

**BAGGIOLINI, M., 1952:** Les stades repères dans le développement annuel de la vigne et leur utilisation pratique. Rev. romande Agric. Vitic. Arboric **1**, 4-6.

**BAGGIOLINI, M., 1980:** Stades repères du cerisier - Stades repères du prunier. Stades repères de l'abricotier. Stades repères du pêcher. ACTA. Guide Pratique de Défense des Cultures, Paris.

**BAILLOD, M. und M. BAGGIOLINI, 1993:** Les stades repères de la vigne. Revue suisse Vitic. Arboric. Hortic. **25**, 7-9.

**BARTELS, G., A. VON KRIES, B. ÄRLÄNDER, U. MEIER, W. STEUDEL und I. M. WITT-STOCK, 1984:** Entwicklungsstadien der Zucker- und Futterrübe. Biologische Bundesanstalt für Land- und Forstwirtschaft, Merkblatt **27/6**.

**BERNING, A., H. GRAF, J. MARTIN, U. MEIER, W. KENNEL und W. ZELLER, 1987:** Entwicklungsstadien von Kernobst. Biologische Bundesanstalt für Land- und Forstwirtschaft. Merkblatt **27/15**.

- BERNING, A., K. HEIN, L. KUNZE und U. MEIER, 1988a:** Entwicklungsstadien von Steinobst. Biologische Bundesanstalt für Land- und Forstwirtschaft, Merkblatt 27/16.
- BERNING, A., U. MEIER, D. NAUMANN, E. SEEMÜLLER und D. SEIPP, 1988b:** Entwicklungsstadien der Erdbeere. Biologische Bundesanstalt für Land- und Forstwirtschaft, Merkblatt 27/17.
- BLEIHOLDER, H., T. EGGERS, M. HANF U. MEIER, 1986:** Entwicklungsstadien zweikeimblättriger Unkräuter. Biologische Bundesanstalt für Land- und Forstwirtschaft, Merkblatt 27/9.
- BLEIHOLDER, H., T. VAN DEN BOOM, P. LANGELÜDDECKE und R. STAUSS, 1989:** Einheitliche Codierung der phänologischen Stadien bei Kultur- und Schadpflanzen. *Gesunde Pflanzen* **41**, 381-384.
- BLEIHOLDER, H., H. KIRFEL, P. LANGE-LÜDDECKE und R. STAUSS, 1991:** Codificação unificada dos estádios fenológicos de culturas e ervas daninhas. *Pesq. agropec. bras.*, Brasília **26**, 1423-1429.
- BLEIHOLDER, H., T. VAN DEN BOOM, P. LANGELÜDDECKE y R. STAUSS, 1991:** Codificación uniforme para los estadios fenológicos de las plantas cultivadas y de las malas hierbas. *Phytoma España* **28**, 54- 56.
- BÖHM, J., W. FRIEDT, K. LINDEMANN und U. MEIER, 1988:** Entwicklungsstadien der Sonnenblume. Biologische Bundesanstalt für Land- und Forstwirtschaft. Merkblatt 27/11.
- BOOTE, K. J., 1980:** Stages of development for peanut. *Proc. Amer. Peanut Res. and Ed. Soc.* **12**, 63.
- BOOTE, K. J., 1982:** Growth Stages of Peanut (*Arachis hypogaea* L.). *Peanut Sci.* **9**, 35-40.
- BUHTZ, E., L. BOESE, C. GRUNERT und W. HAMANN, 1990:** Koordinierter Dezimalcode (KDC) der phänologischen Entwicklung für landwirtschaftliche Kulturpflanzen., Gemüse, Obst und Sonderkulturen. *Feldversuchswesen*, 7/1, Berlin, 94 S.
- CAMARGO, P. de A, 1985:** Florescimento e frutificacao de café arabica nas diferentes regiones cafeeiras do Brazil. *Pesq. Agropec. Bras.* 20(7): 831-839.
- CAMAYO, V. G. C. 1995:** Estudio anatómico y morfológico de la diferenciación y desarrollo de las flores del cafeto *Coffea arabica* L. var. Colombia. Popayán, Universidad del Cauca. 164 p. (Tesis: Licenciatura en Biología).
- CAMAYO, V. G. C. 1995:** Estudio anatómico y morfológico de la diferenciación y desarrollo de las flores del cafeto *Coffea arabica* L. var. Colombia. Popayán, Universidad del Cauca. 164 p. (Tesis: Licenciatura en Biología).
- DANERT, S., 1957:** Über den Sproßaufbau und die Blattentwicklung bei der Kartoffel. *Der Züchter* **27**, 22-33.
- DECOURTYE, L., B. LANTIN und P. VILCOT, 1979:** Stades de développement du cassissier. In: Stades de Développement des Plantes Cultivées. ACTA, Paris: 45.
- DENNIS, R. E. and R. E. BRIGGS, 1969:** Growth and Development of the cotton plant in Arizona. University of Arizona, Cooperative Extension Service and Agricultural Experiment Station, Phoenix, Arizona. Bulletin A-64, 21 p.
- EGGERS, T., G. HEIDLER, 1985:** Entwicklungsstadien von Unkräutern. *Nachrichtenbl. Deut. Pflanzenschutzd.* **37**, 71-76.
- EHLE, H., F. GMELCH, H. LIEBEL, W. LÜDERS und K. ZÜRN, 1980:** Entwicklungsstadien von Hopfen. Biologische Bundesanstalt für Land- und Forstwirtschaft, Merkblatt 27/8.

- EICHHORN, K. W.** und **D. H. LORENZ, 1977:** Phänologische Entwicklungsstadien der Rebe. Nachrichtenbl. Deut. Pflanzenschutzd. **29**, 119-120.
- ELSNER, J. E., C. W. SMITH** and **D. F. OWEN, 1979:** Uniform Stage Descriptions in Upland Cotton. Crop Sci. **19**, 361-363.
- FAO, 1977:** Growth Stage Key - Cotton -. In: Crop Loss Assessment Methods. FAO Manual on the evaluation and prevention of losses by pests, diseases and weeds. Supplement **2**, 4.4.5/1, Rome, Italy.
- FAO, 1977:** Growth Stage Key - Soybean -. In: Crop Loss Assessment Methods. FAO Manual on the evaluation and prevention of losses by pests, diseases and weeds. Supplement **2**, 4.4.12, Rome, Italy.
- FECKES, W., 1941:** De tarwe en haar milieu. Versl. techn. Tarwe Comm: **12**, 523-888 and **17**, 560-561.
- FEHR, W. R., C. E. CAVINESS, D. T. BUR-MOOD** and **J. S. PENNINGTON, 1971:** Stage of Development Descriptions for Soybeans, *Glycine max* (L.) Merr. Crop Sci. **11**, 929-931.
- FEHR, W. R.** and **C. E. CAVINESS, 1977:** Stages of Soybean Development. Iowa State University of Science and Technology. Agriculture and Home Economics Experiment Station, Ames, Iowa. Special Report **80**, 11 p.
- FELICI, G., 1979:** Stades de développement du fraisier. In: Stades de Développement des Plantes Cultivées. ACTA, Paris: **45**.
- FELIPE, A.** und **A. RAMOS, 1984:** Estados tipo del almendro. Estaciones de avisos agrícolas. Ministerio de Agricultura. Madrid. In: EPPO/OEPP 1984, No. 6, 567-568.
- FLECKINGER, J., 1948:** Les stades végétatifs des arbres fruitiers, en rapport avec le traitements. Pomologie Française, Supplément 81-93.
- FREER, J. B. S., 1991:** A development stage key for linseed (*Linum usitatissimum*). Asp.appl. Biol. **28**, 33-40.
- GALL, H., 1988:** Code zur Kennzeichnung von Entwicklungsphasen und -stadien der Kartoffel - Grundlage der Bestandesführung. Feldwirtschaft, Berlin, **29**, 338.
- GRIESS, H.** und **A. MOLL, 1985:** Vorschlag eines neuen Systems von Entwicklungsstadien der Kartoffel. Arch. Acker- und Pflanzenbau und Bodenkunde, Berlin, **29**, 303-310.
- GRIESS, H., 1987:** Entwicklungsstadien der Kartoffel (Systeme von Entwicklungsstadien und Beschreibung der Ontogenese). AdL DDR, Berlin, 58 S.
- GRIESS, H., H. GALL, A. MOLL** und **D. KLEINHEMPEL, 1987:** Zur Einführung eines zweiziffrigen Codes zur Kennzeichnung von Entwicklungsphasen und -stadien der Kartoffel. Feldwirtschaft **28**, 42-44.
- HANWAY, J. J., 1963:** Growth stages of corn (*Zea mays* L.). Agr. Jour. **55**, 487-492.
- HANWAY, J. J.** and **H. E. THOMPSON, 1967:** How a soybean plant develops. Iowa State University of Science and Technology. Cooperative Extension Service, Ames, Iowa. Special Report **53**, 18 p.
- HANWAY, J. J., 1970:** Growth stages of maize/corn. In: Crop Loss Assessment Methods 4.4.2/1. FAO, Rome.
- HEATHCOTE, G. D., 1973:** Growth stages of the sugar beet - root crop - seed crop. Crop Loss Assessment Methods, FAO Manual of the evaluation and prevention of losses by pests, diseases and weeds, 4.4.7/1-4.4.7/2.
- JEFFRIES, R. A.** und **H. M. LAWSON, 1991:** A key for the stages of development of potato (*Solanum tuberosum*). Ann. appl. Biol. **119**, 387-399.

- KELLER, C. und M. BAGGIOLINI, 1954:** Les stades repères dans la végétation du blé. Revue Romande D' Agriculture, Lausanne**10**, 17-20.
- KITTLITZ, E. VON, A. VON KRIES, U. MEIER, R. STÜLPNAGEL und L.-M. WITTSTOCK, 1984:** Entwicklungsstadien der Faba-Bohne. Biologische Bundesanstalt für Land- und Forstwirtschaft. Merkblatt**27/10**.
- KNOTT, C. M., 1987:** A key for stage of development of the pea (*Pisum sativum* L.). Ann. appl. Biol. **111**, 233-244.
- KNOTT, C. M., 1990:** A key for stage of development of the faba bean (*Vicia faba*). Ann. appl. Biol. **116**, 391-401
- KOHEL, R. J. and C. F. LEWIS, 1984:** Cotton. Amer. Soc. Agron., Madison, Wisconsin, USA.
- KOLBE, W., 1979:** Jahreszeitlicher Verlauf der Entwicklungsstadien bei Obstarten in Beziehung zu Jahreswitterung und Pflanzenschutzmaßnahmen. Pflanzenschutz-Nachrichten Bayer **32**, 97-163.
- KRUG, H., 1986:** Gemüseproduktion. Ein Lehr- und Nachschlagewerk für Studium und Praxis. Paul-Parey Verlag, Berlin und Hamburg, 544 S.
- KURTZ, L., H. LYRE, J. STEINBERGER und W. WEDLER, 1979:** Entwicklungsstadien bei Getreide - außer Mais -. Biologische Bundesanstalt für Land- und Forstwirtschaft. Merkblatt **27/1**.
- LAGIERE, V. R., 1966:** Le Cotonier. Techniques Agricoles et Productions Tropicales, Vol. **9**, Maisonneuve & Larose, Paris.
- LANDES, A. and J. R. PORTER, 1989:** Comparison of scales used for categorising the development of wheat, barley, rye and oats. Ann. appl. Biol. **115**, 343-360.
- LARGE, E. C., 1954:** Growth stages in cereals. Illustrations of the Feekes scale. Plant Pathol. **3**, 128-129.
- LE BARON, J., 1974:** Developmental Stages of the Common Bean Plant. University of Idaho, College of Agriculture Current Information, Series Nr. 228.
- LEON J. and L. FOURNIER 1962:** Crecimiento y desarrollo del fruto de Coffea arabica L. Turrialba**12**: 65-74.
- MARTINEZ, E. and A. URSINA 1978:** Ciclos vegetativos de las variedades de plátano y banano comerciales en la zona bananera del Magdalena. Revista COMALFI (Col) **5** (3/4): 105-114.
- MAUNEY, J. R., 1968:** Morphology of the Cotton Plant. In: Elliot, F. C., Hoover, Porter, W. K. Jr. (Editors): Advances in production and utilization of quality cotton; principles and practices. Iowa State University Press, Ames, Iowa, 532 p.
- MEIER, U., 1985:** Die Merkblattserie 27 „Entwicklungsstadien von Pflanzen“. Biologische Bundesanstalt für Land- und Forstwirtschaft. Nachrichtenbl. Deut. Pflanzenschutzd. **37**, 76-77.
- MEIER, U. 1997:** BBCH-Monograph. Growth stages of plants - Entwicklungsstadien von Pflanzen - Estadios de las plantas - Développement des Plantes. Blackwell Wissenschaftsverlag, Berlin und Wien. 622 p
- MEIER, U., 1988:** Merkblätter über Entwicklungsstadien von Kernobst, Steinobst und Erdbeeren. Erwerbsobstbau **4**, 117.
- MÜLLER, G., 1968:** Cotton - Cultivation and Fertilization. Ruhr-Stickstoff AG, Bochum.
- NIJDAM, F. E., 1955:** L' analyse morphologique des caractéristiques agricoles des variétés. Acta bot. Neerl. **4**, 452-459.

- PATTEE, H. E., E. B. JOHNS, J. A. SINGLE-TON and T. H. SANDERS, 1974:** Composition Changes of Peanut Fruit Parts during Maturation. *Peanut Science* **1**, 57-62.
- PINKAU, H. und I. HOLLNAGEL, 1987:** Dezimal-Code zur Kennzeichnung der Wachstums- und Entwicklungsstadien bei Kopfkohl und Anwendungsbeispiele. *Gartenbau* **34**, 135-136.
- PRENTICE, A. N., 1972:** Cotton - with special reference to Africa. Longman, London.
- REESTMAN, A. J. und A. SCHEPERS, 1971:** Toepassing van morphologisch gewas-analyse bij het topol-onderzoek van aardappelen. In: Jaarsverlag 1971, P. A. Lelystad. pp. 61-64. Cited in: H. P. Beukema & D. E. van der Zaag: Introduction to Potato Production. Pudoc, Wageningen, 1980, 208 S.
- SALAZAR G M R, B. CHAVES-CORDOBA, N. M. RIAÑO-HERRERA, J. ARCILA-PULGARIN, J. A. JARAMILLO-ROBLEDO 1994:** Crecimiento del fruto de café, *Coffea arabica* L. var. Colombia. *Cenicafé* **45(2)**: 41-50.
- SALAZAR A N, F. J. OROZCO, J. CLAVIJO-PORRAS 1989:** Características morfológicas, productivas y componentes del rendimiento de dos variedades de café: Colombia y Caturra. *Cenicafé*. **39(2)**: 41-63.
- SANTOS B R, M. MAESTRI, M. P. COONS 1978:** The physiology of flowering in coffee. A Review. *J. Coffee. Res.* **8**: 29-73.
- SCHENK, R. U., 1961:** Development of peanut fruit. *Georgia A. E. S. Techn. Bull. N. S.* **22**, 53 pp.
- SCHNEITER, A. A. and J. F. MILLER, 1981:** Description of sunflower growth stages. *Crop Sci.* **21**, 901-903.
- SCHOTT, P. E., M. HANF, D. O' NEAL, K. SCHELBERGER, M. SCHROEDER, T. WARE and T. JOHN, 1987:** A decimal code for the development stages of a soybean plant - prerequisite for progressive bioregulator research and use. Proceedings of the 14th annual meeting of plant Growth Regulator Society for the Chemical Regulation of Plants, Honolulu, Hawaii, USA.
- SCHÜTTE, F., J. STEINBERGER und U. MEIER, 1982:** Entwicklungsstadien des Raps - einschl. Rübsen, Senfarten und Ölrettich -. *Biologische Bundesanstalt für Land- und Forstwirtschaft. Merkblatt* **27/7**.
- SIDDIQUI, M. Q., J. F. BROWN and S. J. ALLEN, 1975:** Growth stages of sunflower and intensity indices for white blister and rust. *Plant Dis. Repter*. **59**, 7-11.
- SOENEN, A., 1951:** Les bases de l' avertissement en culture fruitière. Le développement du bourgeon floral. *Comptes rendus de Recherches, IRSIA* **5**.
- SPARKS, W. C., and G. W. WOODBURY, 1967:** Stages of potato plant growth. *Idaho Agric. Exper. Stat.* **309**, 1-22.
- STAUSS, R., 1994:** Compendium of growth stage identification keys for mono- and dicotyledonous plants, extended BBCH scale. Ciba-Geigy AG, Basel, 99 p.
- STAUSS, R., 1995:** Compendium pour l' identification des stades phénologiques des espèces mono et dicotylédones cultivées, échelle BBCH. Ciba-Geigy AG, Basel, 104 p.
- STAUSS, R., H. BLEIHOLDER, T. VAN DEN BOOM, L. BUHR, H. HACK, M. HESS, R. KLOSE, U. MEIER und E. WEBER, 1994:** Einheitliche Codierung der phänologischen Entwicklungsstadien mono- und dikotyle Pflanzen. Erweiterte BBCH-Skala: Allgemein. Ciba-Geigy AG, Basel, 58 S.
- SYLVESTER-BRADLEY, R., R. J. MAKEPEACE and H. BROAD, 1984:** A code for stages of development in oilseed rape (*Brassica napus* L.) *Asp. appl. Biol.* **6**, Agronomy, physiology, plant breeding and crop protection of oilseed rape, 399-419.

- SYLVESTER-BRADLEY, R., 1985:** Revision of a code for stages of development in oilseed rape (*Brassica napus* L.). Asp. appl. Biol. **10**, Field Trials Methods and Data handling, 395-400.
- THARP, W. H., 1960:** The cotton plant - How it grows and why its growth varies. United States Department of Agriculture, Agricultural Research Service. Agriculture Handbook No. 178, U. S. Government Printing Office, Washington, D. C.
- THEUNISSEN, J. und A. SINS, 1984:** Growth stages of Brassica crops for crop protection purposes. Sci. Horticult. **24**, 1-11.
- TOTTMAN, D. R., 1977:** The identification of growth stages in winter wheat with reference to the application of growth-regulator herbicides. Ann. appl. Biol. **87**, 213-224.
- TOTTMAN, D. R. and R. J. MAKEPEACE, 1979:** An explanation of the decimal code for the growth stages of cereals, with illustrations. Ann. appl. Biol. **93**, 221-234.
- TOTTMAN, D. R. and H. BROAD, 1987:** The decimal code for the growth stages of cereals, with illustrations. Ann. appl. Biol. **110**, 441-454.
- TROITZKY, N. N., 1925:** Vorläufige Untersuchungsmittel der experimentell-biologischen Station für angewandte Entomologie. Leningrad. In: Kolbe, W., 1979: Jahreszeitlicher Verlauf der Entwicklungsstadien bei Obstarten in Beziehung zu Jahreswitterung und Pflanzenschutzmaßnahmen. In: Pflanzenschutz-Nachrichten Bayer **32**, 97-163.
- VOGEL, G., H. FRÖHLICH, G. BANHOLZER und H. PINKAU, 1987:** Vorschläge zur Charakteristik ausgewählter Gemüsearten auf der Grundlage eines Dezimal-Code-Systems. Gartenbau **34**, 132-134.
- WINNER, C., 1974:** Die Jugendentwicklung der Zuckerrübe in ihrer Bedeutung für das spätere Wachstum und den Ertrag. Zucker **27**, 517-527.