

 Open access • Journal Article • DOI:10.1136/ARD.2008.100362

Guidelines for initiation of anti-tumour necrosis factor therapy in rheumatoid arthritis: similarities and differences across Europe — [Source link](#)

Paul Emery, R. van Vollenhoven, Mikkel Østergaard, Ernest Choy ...+8 more authors

Institutions: University of Leeds

Published on: 01 Apr 2009 - Annals of the Rheumatic Diseases (BMJ Publishing Group)

Topics: Rheumatoid arthritis

Related papers:

- [Modified disease activity scores that include twenty-eight-joint counts : development and validation in a prospective longitudinal study of patients with rheumatoid arthritis](#)
- [The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis.](#)
- [EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs](#)
- [Validation of rheumatoid arthritis improvement criteria that include simplified joint counts.](#)
- [Treating rheumatoid arthritis to target: recommendations of an international task force](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/guidelines-for-initiation-of-anti-tumour-necrosis-factor-4uua1za8fh>

Guidelines for initiation of anti-tumour necrosis factor therapy in rheumatoid arthritis: similarities and differences across Europe

P Emery,¹ R Van Vollenhoven,² M Ostergaard,³ E Choy,⁴ B Combe,⁵ W Graninger,⁶ K Krueger,⁷ M Matucci-Cerinic,⁸ F Navarro,⁹ P van Riel,¹⁰ L Settas,¹¹ S Steinfeld¹²

Rheumatoid arthritis (RA) is a chronic inflammatory autoimmune disease. In the majority of patients the course is progressive and leads to destruction of joints, functional disability, reduced quality of life and increased morbidity and mortality. Advances in treatment for RA have been associated with an improvement in the health status of patients with RA in recent years.¹ While the management of RA varies widely across Europe,² the underlying pathophysiology does not differ between countries. A cross-sectional comparison of patients with RA in 15 different countries (largely European) revealed great variation in the drug management of patients with RA, including the delay in starting a disease-modifying antirheumatic drug (DMARD) ranging from <6 to >12 months, use of methotrexate (<80% of patients in the UK and Serbia had received methotrexate) and use of biological agents (>40% in France and Ireland vs. <10% in Serbia and Poland).³ It is also noteworthy that disease activity scores varied and were highest, on average, in Serbia and Poland.³

Although tumour necrosis factor (TNF) inhibitors have undoubtedly significantly advanced the treatment of RA, they are a

higher cost treatment option and may not be suitable for all patients.⁴ Hence, guidelines have been developed in order to assist the appropriate use of TNF inhibitors within individual countries.

In order to establish the level of access to TNF inhibitors across Europe, European guidelines were translated and reviewed with regard to their similarities and differences in the recommended criteria for the initiation of TNF inhibition. Particular attention was paid to recommended duration of disease and prior treatment with DMARDs, as well as disease activity levels.

The guidelines selected for inclusion were those from Belgium,⁵ Czech Republic,⁶ Denmark,⁷ France,^{8,9} Germany,¹⁰ Italy,¹¹ Romania,¹² Spain,¹³ Sweden¹⁴ and the UK.¹⁵ These were chosen to provide a fair geographical spread of countries, including countries with differing population densities and different levels of TNF inhibitor usage. Table 1 provides an overview of the recommendations made in these guidelines.

Duration of disease prior to initiation of anti-TNF therapy

It is clear that the duration of disease prior to the recommended initiation of TNF inhibitors is highly variable. The general consensus from national guidelines suggests that TNF inhibition could be started between 3 and 6 months following confirmed diagnosis of RA and failure of treatment with, usually, two DMARDs (including methotrexate). However, in the UK and Czech Republic, at least a year needs to elapse from RA diagnosis to TNF inhibitor initiation unless the patient cannot tolerate DMARDs, in which case it is possible to start treatment after only a few months.^{6,15} This is in dramatic contrast to the Spanish guidelines, which suggest the

theoretical possibility (although highly unlikely) of initiating TNF inhibition prior to a confirmed diagnosis of RA, if there is a high probability of RA in initial stages with especially aggressive progression suspected.¹⁵ While the British Society for Rheumatology (BSR) guidance acknowledges the existence of evidence for the benefit of TNF inhibitors in patients with early RA and patients that are DMARD naïve, it does not recommend this approach.¹⁵ A recent editorial regarding the need to update the BSR guidelines suggests that the reasons are financial and acknowledges that the UK BSR guidelines are out of kilter with other national guidelines.¹⁶

There is a growing body of evidence to suggest that treatment with disease modifying agents should be initiated early in the course of the disease. Importantly, Nell *et al* investigated the effects of delays to the initiation of DMARD therapy on long-term outcomes for patients with RA.¹⁷ By comparing two groups of patients, one with a median symptom duration of 3 months and another with a median symptom duration of 12 months, they established that very early therapy provided significant improvements in disease activity, joint destruction and functional outcome. Furthermore, these differences were most apparent within the first 3 months of treatment, suggesting a “window of opportunity” in very early disease to rapidly halt disease progression.¹⁸ The Spanish guidelines cite such evidence to justify their stance of initiating DMARD therapy as soon as, or possibly before, the diagnosis of RA is confirmed.^{15,17-20}

Evidence also exists for the efficacy of all the currently available TNF inhibitors early in the course of RA.²¹ The Early Rheumatoid Arthritis (ERA) trial showed that etanercept monotherapy was superior to methotrexate in reducing subsequent radiographic erosions over 2 years in patients with early erosive RA; at 24 months the American College of Rheumatology 20% (ACR20) response in the etanercept group was significantly higher than that for the methotrexate group (72% vs 59%, respectively, $p = 0.005$).²² A subanalysis of patients with early RA in the Anti-TNF Therapy in RA with Concomitant Therapy (ATTRACT) study showed that early initiation of infliximab in combination with methotrexate provided significant reductions in radiological progression over 2 years compared with methotrexate alone.²³ Finally, the PREMIER study demonstrated that combination adalimumab plus methotrexate was more effective than either

¹ University of Leeds, Leeds, UK; ² Karolinska University Hospital, Stockholm, Sweden; ³ Hvidovre University Hospital, Copenhagen, Denmark; ⁴ King's College, London, UK; ⁵ Hospital Lapeyronie, Montpellier, France; ⁶ University of Graz, Graz, Austria; ⁷ University of Munich, Munich, Germany; ⁸ University of Florence, Florence, Italy; ⁹ Hospital General Universitatario Virgen Macarena, Seville, Spain; ¹⁰ Radboud University Nijmegen, Nijmegen, The Netherlands; ¹¹ University of Thessaloniki, Thessaloniki, Greece; ¹² Erasme University Hospital, Brussels, Belgium

Correspondence to: Professor P Emery, Leeds Teaching Hospitals NHS Trust, Academic Section of Musculoskeletal Disease, 2nd Floor, Chapel Allerton Hospital, Chapeltown Road, Leeds LS7 4SA, UK; p.emery@leeds.ac.uk

Table 1 Summary of the recommendations from the guidance for the use of tumour necrosis factor (TNF) inhibitors in rheumatoid arthritis (RA) in selected European countries

	Disease activity level required for initiation	Trial of DMARD	Choice of agent	Switch between agents	Treatment response measures	Time to see response	Alter frequency/dose	Radiographic monitoring	MTX combination recommended
Belgium ⁵ Société Royale Belge de Rhumatologie	Not specified	Two (one = MTX) for 6 months total	Any	Yes	SJC, HAQ	3–6 months	Not specified	Not mentioned	For IFX: Yes
Czech Republic ⁶ Czech Society for Rheumatology, 2007	DAS28 > 5.1	Two (one = MTX) for 6 months each	Any	Yes	DAS28	12 weeks	Yes	x Ray every 6–12 months	Not specified for Ada or Et Yes
Denmark ⁷ Dansk Reumatologisk Selskab, 2000	Persistent synovitis ≥ 6 joints	Two (one = MTX) for 4 months each	Not specified (Ada not included)	Not specified	SJC, TJC, pain scale, patient and physician global assessments, HAQ	4 months	Not specified	x Ray every 6 months for at least first 2 years, then every 2 years	For IFX: Yes
France ⁸ Société Française de Rhumatologie, 2007	DAS28 > 5.1 or DAS28 > 3.2 in presence of corticosteroid	One for 3 months	Any	Yes	EULAR (DAS28)	12 weeks	Yes	x Rays annually (or more if RA present for long time)	Not specified for Et Yes
Haute Autorité de Santé, 2007	Radiographic progression	Anti-TNF+MTX in severe early RA (such as early joint damage)							
Germany ¹⁰ Deutschen Gesellschaft für Rheumatologie, 2006	Not specified	Two (one = MTX) for 6 months total (less if marked disease progression)	Not specified	Not specified	Not specified	3 months	Not specified	x Rays at least annually	Not specified
Italy ¹¹ Società Italiana di Reumatologia, 2006	DAS > 3.7 or DAS28 > 5.1	One for 3 months	Not specified	Yes	EULAR (DAS)	12 weeks	Not specified	Not mentioned	Yes
Romania ¹² Ministry for Health, 2007	DAS28 > 5.1	Two (one = MTX) for 12 weeks each	Not specified	Yes	SJC, TJC, morning stiffness, ESR, CRP, DAS28	3 months	Yes for IFX	Not mentioned	Not specified
Spain ¹³ Sociedad Española de Reumatología, 2006	DAS28 > 3.2 (or lower in some circumstances)	Usually one for 3 months but none in aggressive disease	Any	Yes	DAS28, ESR, global evaluation of the disease by the patient	4 months	Yes	Annual x ray (plus quantitative analysis)	Yes
Sweden ¹⁴ Svensk Reumatologisk Förening, 2004	DAS28 > 3.2	Two for 2–3 months each or one (MTX) if rapid progression	Not specified	Not specified	Defined on individual basis?	2–3 months	Not specified	x Ray frequency not specified	Yes
UK ¹⁵ British Society for Rheumatology, 2004	DAS28 > 5.1	Two (one = MTX) for 6 months each	Any	BSR yes, NICE no (not permitted for failure to response but permitted for toxicity)	DAS28	3 months	Possible, but not generally recommended	Not mentioned	Yes

Ada, adalimumab; BSR, British Society for Rheumatology; CRP, C-reactive protein; DAS(28), (28-joint) Disease Activity Score; DMARD, disease-modifying antirheumatic drug; ESR, erythrocyte sedimentation rate; Et, etanercept; EULAR, European League Against Rheumatism; HAQ, Health assessment questionnaire; IFX, infliximab; MTX, methotrexate; NICE, National Institute for Clinical Excellence; SJC, swollen joint count; TJC, tender joint count.

agent as monotherapy in patients with early, aggressive RA.²⁴

By comparing therapeutic strategies in patients newly diagnosed with RA, the BeSt study showed that initial combination therapy with methotrexate plus a TNF inhibitor provided earlier functional improvement and less progression of radiographic joint damage than either sequential monotherapy or step-up combination therapy (conventional DMARDs).²⁵ There is also evidence, from the Trial of Etanercept and Methotrexate with Radiographic Patient Outcomes (TEMPO) study, that in patients who initially receive and fail DMARDs other than methotrexate, combination etanercept and methotrexate is more effective than either agent alone.²⁶ Furthermore, no increase in toxicity was associated with combination treatment with etanercept plus methotrexate.²⁶

Disease activity

Across Europe the goals of RA treatment with TNF inhibitors vary considerably. While a reduction in disease activity is a common theme, the levels of activity designated for initiation and/or cessation of TNF inhibitors differ.

When specified, a Disease Activity Score using 28 joint counts (DAS28) of over 5.1 is commonly used to define highly active disease that could qualify for TNF inhibition. The BSR guidelines for the prescribing of TNF inhibitors in the UK require two DAS measurements at least 1 month apart to confirm the ongoing activity of disease.¹⁵ In Romania the patient must also have five or more joints with active synovitis (tender and swollen joints) plus two of the following three criteria: morning stiffness for over 60 min; erythrocyte sedimentation rate (ESR) >28 mm/h; C-reactive protein (CRP) >20 mg/litre (quantitative determination).¹² However, in Spain and Sweden moderate disease activity also qualifies for the consideration of TNF inhibitors, defined as DAS28 >3.2.^{13 14} In Spain, patients who in spite of achieving remission or low DAS28 show radiographic progression or persistent inflammation in joints of special functional value, may also be considered for TNF inhibitors.¹³

In order for treatment with TNF inhibitors to continue unaltered, guidelines generally require a European League Against Rheumatism (EULAR) response to be demonstrated. EULAR response is defined as DAS28 <3.2 or DAS28 <5.1 and a decrease in DAS28 by at least 1.2 points. Denmark requires that a satisfactory response is achieved, although it is not clearly specified what this implies.⁷ In

Spain, in keeping with their aggressive treatment strategy, treatment should be altered if a of DAS28 <3.2 is not achieved, however, if even this is not possible, a maximum of 5 swollen and painful joints among 66 and 68 joints, respectively, should be the aim of treatment.¹⁵

CONCLUSIONS

It is clear that there is obvious variability in the framework for initiation of TNF inhibitors across Europe. The clinical evidence base available to the national rheumatological societies concerned is largely the same, although obviously expanding as time progresses. Therefore it is interesting that despite many similarities, there are differences in the criteria for initiation.

It is also apparent that a number of areas are not covered by the guidelines. For example, initial DMARD therapy is a requirement of all the European guidelines, and all state inadequate response as a criterion for changing to anti-TNF therapy. However, guidelines do not provide a clear definition of inadequate response to DMARDs. Such definitions are essential to ensure treatment progresses in a way that provides optimal outcomes for the patient.

Even with the similarities between guidelines, such as they are, TNF inhibitor prescribing patterns vary greatly between countries; it would seem that the prospects for people diagnosed with RA are substantially influenced by geographical location, possibly due to the different methods of funding healthcare provision. Other factors may give rise to such differences in treatment, such as levels of recognition of guidelines, the presence or absence of specialised rheumatological clinics, including early RA clinics, and familiarity with TNF inhibitors.

It should be possible for patients to be certain that they are receiving the best advice regarding their treatment, based on clinical evidence and experience, rather than the financial situation, or other influencing factors, in their respective healthcare organisation. In order to achieve this we propose that European guidance, based upon a thorough evidence review, should be prepared and made universally available in order to promote more consistent access to effective therapy.

Acknowledgements: Synergy provided writing assistance.

Funding: Wyeth provided an unrestricted grant for research and writing of this article.

Competing interests: RVV has received consulting honoraria and speaker's fees from Wyeth. EC has served on advisory boards and speaker bureaus for Abbott,

Schering Plough, Wyeth, UCB Celltech and Roche. WG has served on an advisory board and Speaker Bureaus for Abbott, Schering Plough, Wyeth, UCB Celltech and Roche. SS has served in advisory boards for Abbott, Schering-Plough, Genentech and Wyeth. BC has served as a speaker and/or consultant and/or investigator for Abbott, BMS, GSK, MSD, Roche, Schering, UCB and Wyeth. PE has provided expert advice and undertaken clinical trials for Abbott, BMS, GSK, MSD, Roche, Schering, UCB and Wyeth.

Accepted 11 January 2009

Ann Rheum Dis 2009;**68**:456–459.
doi:10.1136/ard.2008.100362

REFERENCES

- Uhlir T, Heiberg T, Mowinckel P, Kvien TK. Rheumatoid arthritis is milder in the new millennium: health status in patients with rheumatoid arthritis 1994–2004. *Ann Rheum Dis* 2008;**67**:1710–15.
- Combe B, Landewe R, Lukas C, Bolosiu HD, Breedveld F, Dougados M, et al. EULAR recommendations for the management of early arthritis: report of a task force of the European Standing Committee for International Clinical Studies Including Therapeutics (ESCIIT). *Ann Rheum Dis* 2007;**66**:34–45.
- Sokka T, Kautiainen H, Toloza S, Mäkinen H, Verstappen SMM, Hetland ML, et al. QUEST-RA: quantitative clinical assessment of patients with rheumatoid arthritis seen in standard rheumatology care in 15 countries. *Ann Rheum Dis* 2007;**66**:1491–6.
- Sesin CA, Bingham CO. Remission in rheumatoid arthritis: wishful thinking or clinical reality? *Semin Arthritis Rheum* 2005;**35**:185–96.
- Société Royale Belge de Rhumatologie. Belgian guidelines. <http://www.srb.be/> (accessed 30 January 2009).
- Becvar R, Vencovsky J, Nemeč P, Suchý D, Procházková L, Pavelka K, et al. Recommendations of the Czech Society for Rheumatology for treatment of rheumatoid arthritis. Efficacy and strategy for treatment. *Ces Revmatol* 2007;**15**:73–90.
- Dansk Reumatologisk Selskab. *Illustrative guidelines for TNF-alpha inhibiting treatment with rheumatoid arthritis*. Brønshøj, Denmark: Institut for Rational Farmakoterapi, 2000.
- Fautrel B, Pham T, Mouterde G, Le Loet X, Goupille P, Guillemin F et al. Club Rhumatismes et Inflammation; Société Française de Rhumatologie. Recommendations of the French Society for Rheumatology regarding TNFalpha antagonist therapy in patients with rheumatoid arthritis. *Joint Bone Spine* 2007;**74**:627–37.
- French Health Authority. *Rheumatoid polyarthritis (RP): treatment at the established stage. Summary of Professional Recommendations*. Saint-Denis La Plaine, France: Haute Autorité de Santé, 2007.
- Manger B, Michels H, Nüsslein HG, Schneider M, Sieper J, die Kommission Pharmakotherapie der Deutschen Gesellschaft für Rheumatologie. Neufassung der Empfehlungen der Deutschen Gesellschaft für Rheumatologie zur Therapie mit Tumornekrosefaktor-hemmenden Wirkstoffen bei entzündlich-rheumatischen Erkrankungen. Stand März 2006. http://www.dgrh.de/fileadmin/media/Qualitaetsicherung/Therapie-Empfehlungen/TNF-Blocker_06.pdf (accessed 27 January 2009).
- Valesini G, Montecucco C, Cutolo M. Recommendations for the use of biologic (TNF-alpha blocking) agents in the treatment of rheumatoid arthritis in Italy. *Clin Exp Rheumatol* 2006;**24**:413–23.
- Casa Națională de Asigurări de Sănătate. Ordin privind aprobarea Protocolului de practică pentru prescrierea, monitorizarea și decontarea medicamentelor în cazul poliartritei reumatoide pentru persoanele asigurate în sistemul de asigurări sociale de sănătate. *Monitorul Oficial* 2007;**366**:Act 22.
- Rodriguez-Valverde V, Cáliz RC, Alvaro-Gracia Alvaro JM, Marengo de la Fuente JL, Mendoza JM, Molina JT, et al. III actualización del consenso de la Sociedad Española de Reumatología sobre terapia

- biológica en la artritis reumatoide. *Reumatol Clin* 2006;**2**(Suppl):263–70.
14. **Svensk Reumatologisk Förening.** *Riktlinjer för vård och behandling av patienter med reumatoid artrit. Grunddokumentet.* Sweden: Svensk Reumatologisk Förening, 2004.
 15. **Ledingham J,** Deighton C, BSR Standards, Guidelines and Audit Working Group (SGWAG). *Update on the British Society for Rheumatology (BSR) guidelines for prescribing TNF alpha blockers in adults with rheumatoid arthritis (update of previous guidelines April 2001).* London, UK: BSR, 2004.
 16. **Deighton CM,** George E, Kiely PDW, Ledingham J, Luqmani RA, Scott DGI. Updating the British Society for Rheumatology guidelines for anti-tumour necrosis factor therapy in adult rheumatoid arthritis (again). *Rheumatology* 2006;**45**:649–52.
 17. **Nell VPK,** Machold KP, Eberl G, Stamm TA, Uffmann M, Smolen JS. Benefit of very early referral and very early therapy with disease-modifying anti-rheumatic drugs in patients with very early rheumatoid arthritis. *Rheumatology* 2004;**43**:906–14.
 18. **Lard LR,** Visser H, Speyer I, Vander Horst-Bruinsma IE, Zwiderman AH, Breedveld FC, *et al.* Early versus delayed treatment in patients with recent-onset rheumatoid arthritis: comparison of two cohorts who received different treatment strategies. *Am J Med* 2001;**111**:446–51.
 19. **Quinn MA,** Conaghan PG, Emery P. The therapeutic approach of early intervention for rheumatoid arthritis: what is the evidence? *Rheumatology* 2001;**40**:1211–20.
 20. **Boers M.** Understanding the window of opportunity concept in early rheumatoid arthritis. *Arthritis Rheum* 2003;**48**:1771–4.
 21. **Machold KP,** Nell VPK, Stamm TA, Smolen JS. Traditional DMARD therapy: is it sufficient? *Arthritis Res Ther* 2006;**8**:211.
 22. **Bathon JM,** Genovese MC. The early rheumatoid arthritis (ERA) trial comparing the efficacy and safety of etanercept and methotrexate. *Clin Exp Rheumatol* 2003;**21**(Suppl 31):S195–7.
 23. **Breedveld FC,** Emery P, Keystone E, Patel K, Furst DE, Kalden JR, *et al.* Infliximab in active early rheumatoid arthritis. *Ann Rheum Dis* 2004;**63**:149–55.
 24. **Breedveld FC,** Weisman MH, Kavanaugh AF, Cohen SB, Pavelka K, van Vollenhoven R, *et al.* The PREMIER study. A multicenter, randomized, double-blind clinical trial of combination therapy with adalimumab plus methotrexate versus methotrexate alone or adalimumab alone in patients with early, aggressive rheumatoid arthritis who had not had previous methotrexate treatment. *Arthritis Rheum* 2006;**54**:26–37.
 25. **Goekoop-Ruiterman YPM,** de Vries-Bouwstra JK, Allart CF, van Zeben D, Kerstens PJSM, Hazes JMWW, *et al.* Clinical and radiographic outcomes of four different treatment strategies in patients with early rheumatoid arthritis (the BeSt study): a randomized controlled trial. *Arthritis Rheum* 2005;**52**:3381–90.
 26. **Van der Heijde D,** Klareskog L, Rodriguez-Valverde V, Codreanu C, Bolosiu H, Melo-Gomes J, *et al.* Comparison of etanercept and methotrexate, alone and combined, in the treatment of rheumatoid arthritis. Two-year clinical and radiographic results from the TEMPO study, a double-blind, randomized trial. *Arthritis Rheum* 2006;**54**:1063–74.

Corrections

The department of one of the authors who co-authored all of the below papers has found that the affiliations were not correct. The correct affiliations for Professor P Emery, for all of the below articles, are: ¹Section of Musculoskeletal Disease, Leeds Institute of Molecular Medicine, University of Leeds; ²NIHR Leeds Musculoskeletal Biomedical Research Unit, Leeds Teaching Hospitals Trust, Leeds, UK.

- Keystone E**, Emery P, Peterfy CG, *et al.* Rituximab inhibits structural joint damage in patients with rheumatoid arthritis with an inadequate response to tumour necrosis factor inhibitor therapies. *Ann Rheum Dis* 2009;**68**:216–21.
- Doward LC**, McKenna SP, Whalley D, *et al.* The development of the L-QoL: a quality-of-life instrument specific to systemic lupus erythematosus. *Ann Rheum Dis* 2009;**68**:196–200.
- Potter C**, Hyrich KL, Tracey A, *et al.* Association of rheumatoid factor and anti-cyclic citrullinated peptide positivity, but not carriage of shared epitope or PTPN22 susceptibility variants, with anti-TNF response in RA. *Ann Rheum Dis* 2009;**68**:69–74.
- Smolen JS**, Han C, van der Heijde DM, *et al.*; Active-Controlled Study of Patients Receiving Infliximab for the Treatment of Rheumatoid Arthritis of Early Onset (ASPIRE) Study Group. Radiographic changes in rheumatoid arthritis patients attaining different disease activity states with methotrexate monotherapy and infliximab plus methotrexate: the impacts of remission and tumour necrosis factor blockade. *Ann Rheum Dis* 2009;**68**:823–7.
- Buch MH**, Boyle DL, Rosengren S, *et al.* Mode of action of abatacept in rheumatoid arthritis patients having failed tumour necrosis factor blockade: a histological, gene expression and dynamic magnetic resonance imaging pilot study. *Ann Rheum Dis* 2009;**68**:1220–7.
- Emery P**, Van Vollenhoven R, Ostergaard M, *et al.* Guidelines for initiation of anti-tumour necrosis factor therapy in rheumatoid arthritis: similarities and differences across Europe. *Ann Rheum Dis* 2009;**68**:456–9.
- Bejarano V**, Conaghan PG, Proudman SM, *et al.* Long-term efficacy and toxicity of ciclosporin A in combination with methotrexate in poor prognosis rheumatoid arthritis. *Ann Rheum Dis* 2009;**68**:761–3.
- Rudwaleit M**, Landewé R, van der Heijde D, *et al.* The development of Assessment of SpondyloArthritis international Society classification criteria for axial spondyloarthritis (part I): classification of paper patients by expert opinion including uncertainty appraisal. *Ann Rheum Dis* 2009;**68**:770–6.
- Bennett AN**, Marzo-Ortega H, Emery P, *et al.*; Leeds Spondyloarthropathy Group. Diagnosing axial spondyloarthropathy. The new Assessment in SpondyloArthritis international Society criteria: MRI entering centre stage. *Ann Rheum Dis* 2009;**68**:765–7.
- Marzo-Ortega H**, McGonagle D, O'Connor P, *et al.* Baseline and 1-year magnetic resonance imaging of the sacroiliac joint and lumbar spine in very early inflammatory back pain. Relationship between symptoms, HLA-B27 and disease extent and persistence. *Ann Rheum Dis* 2009;**68**:1721–7.
- Gilworth G**, Emery P, Gossec L, *et al.* Adaptation and cross-cultural validation of the rheumatoid arthritis work instability scale (RA-WIS). *Ann Rheum Dis* 2009;**68**:1686–90.
- Gilworth G**, Emery P, Gossec L, *et al.* Adaptation and cross-cultural validation of the RA-WIS (Work Instability Scale). *Ann Rheum Dis* 2009;**68**:1686–90.
- Jarrett SJ**, Sivera F, Cawksell LS, *et al.* MRI and clinical findings in patients with ankylosing spondylitis eligible for anti-tumour necrosis factor therapy after a short course of etoricoxib. *Ann Rheum Dis* 2009;**68**:1466–9.
- Haugeberg G**, Conaghan PG, Quinn M, *et al.* Bone loss in patients with active early rheumatoid arthritis: infliximab and methotrexate compared with methotrexate treatment alone. Exploratory analysis from a 12-month randomised, double-blind, placebo-controlled study. *Ann Rheum Dis* 2009;**68**:1898–901.
- Genovese MC**, Breedveld FC, Emery P, *et al.* Safety of biological therapies following rituximab treatment in rheumatoid arthritis patients. *Ann Rheum Dis* 2009;**68**:1894–7.
- Kekow J**, Moots RJ, Emery P, *et al.* Patient-reported outcomes improve with etanercept plus methotrexate in active early rheumatoid arthritis and the improvement is strongly associated with remission: the COMET trial. *Ann Rheum Dis* 2010;**69**:222–5.
- Furst DE**, Keystone EC, Fleischmann R, *et al.* Updated consensus statement on biological agents for the treatment of rheumatic diseases, 2009. *Ann Rheum Dis* 2010;**69**(Suppl 1):i2–29.
- Freeston JE**, Wakefield RJ, Conaghan PG, *et al.* A diagnostic algorithm for persistence of very early inflammatory arthritis: the utility of power Doppler ultrasound when added to conventional assessment tools. *Ann Rheum Dis* 2010;**69**:417–9.
- Jones E**, Churchman SM, English A, *et al.* Mesenchymal stem cells in rheumatoid synovium: enumeration and functional assessment in relation to synovial inflammation level. *Ann Rheum Dis* 2010;**69**:450–7.
- Alten RE**, Zerbini C, Jeka S, *et al.* Efficacy and safety of pamapimod in patients with active rheumatoid arthritis receiving stable methotrexate therapy. *Ann Rheum Dis* 2010;**69**:364–7.
- Machold KP**, Landewé R, Smolen JS, *et al.* The Stop Arthritis Very Early (SAVE) trial, an international multicentre, randomised, double-blind, placebo-controlled trial on glucocorticoids in very early arthritis. *Ann Rheum Dis* 2010;**69**:495–502.
- Schoels M**, Knevel R, Aletaha D, *et al.* Evidence for treating rheumatoid arthritis to target: results of a systematic literature search. *Ann Rheum Dis* 2010;**69**:638–43.
- Smolen JS**, Aletaha D, Bijlsma JW, *et al.*; T2T Expert Committee. Treating rheumatoid arthritis to target: recommendations of an international task force. *Ann Rheum Dis* 2010;**69**:631–7.
- Burr ML**, Naseem H, Hinks A, *et al.*; BIRAC Consortium; YEAR Consortium. PADI4 genotype is not associated with rheumatoid arthritis in a large UK Caucasian population. *Ann Rheum Dis* 2010;**69**:666–70.
- Emery P**, Durez P, Dougados M, *et al.* Impact of T-cell costimulation modulation in patients with undifferentiated inflammatory arthritis or very early rheumatoid arthritis: a clinical and imaging study of abatacept (the ADJUST trial). *Ann Rheum Dis* 2010;**69**:510–16.
- Bennett AN**, Rehman A, Hensor EM, *et al.* The fatty Romanus lesion: a non-inflammatory spinal MRI lesion specific for axial spondyloarthropathy. *Ann Rheum Dis* 2010;**69**:891–4.
- Nam JL**, Winthrop KL, van Vollenhoven RF, *et al.* Current evidence for the management of rheumatoid arthritis with biological disease-modifying antirheumatic drugs: a systematic literature review informing the EULAR recommendations for the management of RA. *Ann Rheum Dis* 2010;**69**:976–86.
- Smolen JS**, Landewé R, Breedveld FC, *et al.* EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs. *Ann Rheum Dis* 2010;**69**:964–75.
- Tan RJ**, Gibbons LJ, Potter C, *et al.*; BRAGGSS. Investigation of rheumatoid arthritis susceptibility genes identifies association of AFF3 and CD226 variants with response to anti-tumour necrosis factor treatment. *Ann Rheum Dis* 2010;**69**:1029–35.
- Robinson JI**, Barrett JH, Taylor JC, *et al.*; YEAR Consortium; BRAGGSS. Dissection of the FCGR3A association with RA: increased association in men and with autoantibody positive disease. *Ann Rheum Dis* 2010;**69**:1054–7.
- Cohen SB**, Keystone E, Genovese MC, *et al.* Continued inhibition of structural damage over 2 years in patients with rheumatoid arthritis treated with rituximab in combination with methotrexate. *Ann Rheum Dis* 2010;**69**:1158–61.
- Haugeberg G**, Bennett AN, McGonagle D, *et al.* Bone loss in very early inflammatory back pain in undifferentiated spondyloarthropathy: a 1-year observational study. *Ann Rheum Dis* 2010;**69**:1364–6.
- Schoels M**, Aletaha D, Smolen JS, *et al.* Follow-up standards and treatment targets in rheumatoid arthritis: results of a questionnaire at the EULAR 2008. *Ann Rheum Dis* 2010;**69**:575–8.
- Eyre S**, Flynn E, Martin P, *et al.* No evidence for association of the KLF12 gene with rheumatoid arthritis in a large UK cohort. *Ann Rheum Dis* 2010;**69**:1407–8.
- Eyre S**, Hinks A, Flynn E, *et al.* Confirmation of association of the REL locus with rheumatoid arthritis susceptibility in the UK population. *Ann Rheum Dis* 2010;**69**:1572–3.
- Orozco G**, Eyre S, Hinks A, *et al.*; Wellcome Trust Case Control consortium YEAR Consortium. Association of CD40 with rheumatoid arthritis confirmed in a large UK case-control study. *Ann Rheum Dis* 2010;**69**:813–16.
- Emery P**, Durez P, Dougados M, *et al.* Impact of T-cell costimulation modulation in patients with undifferentiated inflammatory arthritis or very early rheumatoid arthritis: a clinical and imaging study of abatacept (the ADJUST trial). *Ann Rheum Dis* 2010;**69**:510–16.
- Saleem B**, Keen H, Goeb V, *et al.* Patients with RA in remission on TNF blockers: when and in whom can TNF blocker therapy be stopped? *Ann Rheum Dis* 2010;**69**:1636–42.
- Barkham N**, Coates LC, Keen H, *et al.* Double-blind placebo-controlled trial of etanercept in the prevention of work disability in ankylosing spondylitis. *Ann Rheum Dis* 2010;**69**:1926–8.
- Emery P**, Deodhar A, Rigby WF, *et al.* Efficacy and safety of different doses and retreatment of rituximab: a randomised, placebo-controlled trial in patients who are biological naive with active rheumatoid arthritis and an inadequate response to methotrexate (Study Evaluating Rituximab's Efficacy in MTX iNadequate rEsponders (SERENE)). *Ann Rheum Dis* 2010;**69**:1629–35.
- Dixon WG**, Hyrich KL, Watson KD, *et al.*; BSRBR Control Centre Consortium; British Society for Rheumatology Biologics Register. Influence of anti-TNF therapy on mortality in patients with rheumatoid arthritis-associated interstitial lung disease: results from the British Society for Rheumatology Biologics Register. *Ann Rheum Dis* 2010;**69**:1086–91.

Ann Rheum Dis 2011;**70**:1519. doi:10.1136/annrheumdis-2011-70-08



Guidelines for initiation of anti-tumour necrosis factor therapy in rheumatoid arthritis: similarities and differences across Europe

P Emery, R Van Vollenhoven, M Ostergaard, et al.

Ann Rheum Dis 2009 68: 456-459
doi: 10.1136/ard.2008.100362

Updated information and services can be found at:

<http://ard.bmj.com/content/68/4/456.full.html>

These include:

References

This article cites 20 articles, 7 of which can be accessed free at:

<http://ard.bmj.com/content/68/4/456.full.html#ref-list-1>

Article cited in:

<http://ard.bmj.com/content/68/4/456.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>