



Gut microbiota interacts with markers of adipose tissue browning, insulin action and plasma acetate in morbid obesity

Journal:	<i>Molecular Nutrition and Food Research</i>
Manuscript ID	mnfr.201700721
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	23-Aug-2017
Complete List of Authors:	Moreno-Navarrete, José María Serino, Matteo Blasco-Baque, Vincent Azalbert, Vincent Barton, Richard Cardellini, Marina Latorre, Jessica Ortega, Francisco; Institut d'Investigació Biomèdica de Girona (IdIBGi), Service of Diabetes, Endocrinology and Nutrition (UDEN); José Ortega, Sabater-Masdeu, Mònica Burcelin, R Dumas, ME Ricart, Wifredo Federici, Massimo Fernández-Real, José Manuel
Keywords:	Adipose tissue, Insulin sensitivity, microbiota, obesity, metabolome

SCHOLARONE™
Manuscripts

1
2
3 **Gut microbiota interacts with markers of adipose tissue browning,**
4
5 **insulin action and plasma acetate in morbid obesity**
6
7

8 José María Moreno-Navarrete^{1*}, Matteo Serino², Vincent Blasco-Baque², Vincent
9 Azalbert², Richard Barton³, Marina Cardellini⁴, Jèssica Latorre¹, Francisco Ortega¹,
10 Mònica Sabater-Masdeu¹, Rémy Burcelin², Marc Dumas³, Wifredo Ricart¹, Massimo
11 Federici⁴, José Manuel Fernández-Real^{1*}
12

13 ¹Department of Diabetes, Endocrinology and Nutrition, Institut d'Investigació Biomèdica de
14 Girona (IdIBGi), CIBEROBN (CB06/03/010) and Instituto de Salud Carlos III (ISCIII), Girona,
15 Spain.
16

17 ²Institut National de la Santé et de la Recherche Médicale (Inserm), U1048, I2MC, Toulouse,
18 France
19

20 ³Faculty of Medicine, Department of Surgery and Cancer, Division of Computational and
21 Systems Medicine, Imperial College London, Sir Alexander Fleming Building, Exhibition
22 Road, South Kensington, London SW7 2AZ, UK.
23

24 ⁴Department of Systems Medicine and Center for Atherosclerosis, University of Rome "Tor
25 Vergata", Rome, Italy.
26

27 **Short title::** Ruminococcaceae and adipose tissue browning
28
29

30
31
32 **Corresponding author and person to whom reprint requests should be addressed:**

33 J.M. Moreno-Navarrete, PhD e-mail: jmoreno@idibgi.org

34 J.M. Fernández-Real, M.D. Ph.D e-mail: jmfreal@idibgi.org

35 Section of Diabetes, Endocrinology and Nutrition

36 Hospital of Girona "Dr Josep Trueta"

37 Carretera de França s/n, 17007, Girona, SPAIN.

38 Phone: 34-972-94 02 00

39 Fax: 34-972-94 02 70
40
41

42
43 **Abbreviations:** HOMA-IR, homeostatic model assessment of insulin resistance; M,
44 insulin action value measured by the euglycemic hyperinsulinemic clamp; RA, relative
45 abundance; RU, relative units of gene expression; SAT, subcutaneous adipose tissue;
46 VAT, visceral adipose tissue.
47
48
49

50
51 **Keywords:** adipose tissue, insulin sensitivity, metabolome, microbiota, obesity.
52
53
54
55
56
57
58
59
60

Abstract

Scope: To examine the potential relationship among gene expression markers of adipose tissue browning, gut microbiota and insulin sensitivity in humans.

Methods and results: Gut microbiota composition and gene markers of browning were analysed in subcutaneous (SAT) and visceral (VAT) adipose tissue from morbidly obese subjects (n=34). Plasma acetate was measured through ^1H NMR and insulin sensitivity using euglycemic hyperinsulinemic clamp. Subjects with insulin resistance showed an increase in the relative abundance (RA) of the phyla Bacteroidetes and Proteobacteria while RA of Firmicutes was decreased. In all subjects, Firmicutes RA was negatively correlated with HbA_{1c} and fasting triglycerides, whereas Proteobacteria RA was negatively correlated with insulin sensitivity. Firmicutes RA was positively associated with markers of brown adipocytes (*PRDM16*, *UCP1* and *DIO2*) in SAT, but not in VAT. Multivariate regression analysis indicated that Firmicutes RA contributed significantly to SAT *PRDM16*, *UCP1* and *DIO2* mRNA variance after controlling for age, BMI, HbA_{1c} or insulin sensitivity. Interestingly, Firmicutes RA, specifically those bacteria belonging to Ruminococcaceae family, was positively associated with plasma acetate levels, which were also linked to SAT *PRDM16* mRNA and insulin sensitivity.

Conclusion: Gut microbiota composition is linked to adipose tissue browning and insulin action in humans, possibly through circulating acetate.

Introduction

The human gut microbiota is composed of 100 trillion microbes that exist in a close symbiotic relationship with human cells, and contain at least 150 times more genes than the whole human genome [1,2]. The metabolic state of the host, diet and medication exert significant effects on the abundance and the composition of the gut microbiota. Experiments in rodents demonstrated that the microbiota can modulate both energy balance (weight gain and loss) and energy stores (fat mass) through the production or the secretion of specific molecules (such as short-chain fatty acids) [3,4].

The dominant phyla in the gut are Firmicutes (~60–65%), Bacteroidetes (~20–25%), Proteobacteria (~5–10%), and Actinobacteria (~3%), which together constitute over 97% of the gut microbe population [4]. In rodents, obesity is associated with an increase in the relative size of the Firmicutes versus Bacteroidetes populations in the gut [5], and a decrease in the diversity of the microbiota that is due to both weight and diet composition. There is much greater variability in human studies. Some studies report a similar increase in the ratio of Firmicutes / Bacteroidetes, as well as a decrease in the gut biodiversity in obese humans, but there are also numerous studies reporting contradictory findings in obese vs. lean humans [6,7]. A recent study demonstrated that cold exposure led to dramatic changes of the microbiota composition, increasing Firmicutes vs. Bacteroidetes relative abundance [8]. Interestingly, this shift in microbiota composition is associated to a phenotype of increased capacity of energy extraction, characterized by increased intestinal absorptive surface area and browning of the white adipose tissue [8]. In mice, browning of white adipose tissue prevents obesity-associated metabolic disturbances and insulin resistance [9,10]. In humans, expression

1
2
3 of markers of browning in subcutaneous adipose tissue has been associated to insulin
4
5 sensitivity, a relatively healthy obese phenotype and adipose tissue function [11-13].
6

7
8 To the best of our knowledge, the interaction between microbiota and adipose tissue
9
10 physiology has not been previously investigated in humans. Given the potential
11
12 importance of adipose tissue inter-relationships with microbiota and insulin action in
13
14 human obesity, here we aimed to examine the associations among adipose tissue
15
16 markers of browning (PRDM16, UCP1 and DIO2), insulin sensitivity (SLC2A4 and
17
18 IRS1) and gut metagenomic microbiota composition in morbidly obese participants. We
19
20 also studied adipose tissue markers of adipogenesis (ADIPOQ, PPARG) and chronic
21
22 proinflammatory activity inflammation (TNF and IL6).
23
24
25
26
27

28 **Materials and Methods**

29 **Recruitment of patients and processing of samples.**

30
31
32
33 The human subject cohort comprised 34 morbidly obese women (n=28) and men (n=6)
34
35 at the Endocrinology Service of the Hospital Universitari de Girona Dr Josep Trueta
36
37 (Girona, Spain) included in the FLORINASH project (www.florinash.org). 11 subjects
38
39 received diabetes, 18 and 9 morbidly obese participants receives hypertension and
40
41 dyslipemia therapy, respectively. 13 morbidly obese patients have type 2 diabetes, of
42
43 who 11 received therapy. Pre-established inclusion criteria: all subjects were of
44
45 Caucasian origin and reported a stable body weight at least three months preceding the
46
47 study, were free of any infections and antibiotic treatment for one month before
48
49 inclusion and had no systemic disease. Pre-established exclusion criteria: subjects with
50
51
52
53
54
55
56
57
58
59
60

1
2
3 liver disease, specifically HCV infection and tumor disease, and subjects with thyroid
4
5 dysfunction were excluded by biochemical workup.

6
7 Subcutaneous (SAT) and visceral adipose tissue (VAT) samples were obtained from all
8
9 subjects during elective gastric bypass surgery and were immediately transported to the
10
11 laboratory (5-10 min). The handling of tissue was carried out under strictly aseptic
12
13 conditions. Adipose tissue samples were washed in PBS, cut off with forceps and
14
15 scalpel into small pieces (100 mg), and immediately flash-frozen in liquid nitrogen
16
17 before being stored at -80 °C.
18
19

20 **Euglycemic hyperinsulinemic clamp**

21
22 Insulin action was determined by the euglycemic hyperinsulinemic clamp one week
23
24 before surgery [14]. After an overnight fast, two catheters were inserted into an
25
26 antecubital vein, one for each arm, used to administer constant infusions of glucose and
27
28 insulin, and to obtain arterialized venous blood samples. A 2-h euglycemic
29
30 hyperinsulinemic clamp was initiated by a two step primed infusion of insulin (80
31
32 mU/m²/min for 5 min, 60 mU/m²/min for 5 min) immediately followed by a
33
34 continuous infusion of insulin at a rate of 40 mU/m²/min (regular insulin; Actrapid,
35
36 Novo Nordisk, NJ). Glucose infusion began at minute 4 at an initial perfusion rate of 2
37
38 mg/kg/min, then was adjusted to maintain plasma glucose concentration at 4.9–5.5
39
40 mmol/L. Blood samples were collected every 5 min for determination of plasma glucose
41
42 and insulin. Insulin sensitivity was assessed as the mean glucose infusion rate during the
43
44 last 40 min. In the stationary equilibrium, the amount of glucose administered (M)
45
46 equals the glucose taken by the body tissues and is a measure of overall insulin
47
48 sensitivity.
49
50

51
52
53
54 Subjects in the upper tertile of glucose infusion rate during the clamp were considered
55
56 as insulin sensitive.
57
58
59
60

1
2
3 All subjects gave written informed consent, validated and approved by the Ethical
4
5 Committee of the Hospital of Girona “Dr Josep Trueta”, after the purpose of the study
6
7 was explained to them.
8

9 10 **Gut microbiota composition**

11 Total DNA was extracted as previously described [15-19]. Briefly, the whole 16S
12
13 bacterial DNA V2 region was targeted by the 28F-519R primers and analysed by
14
15 pyrosequencing by the 454 FLX Roche technologies at Research&Testing Laboratory
16
17 (<http://www.researchandtesting.com/>, Texas, USA). An average of 5000 sequences was
18
19 generated per sample. A full description of bioinformatic filters is available at
20
21 http://www.rtlgenomics.com/docs/Data_Analysis_Methodology.pdf.
22
23
24

25 26 **Metabolomic analyses**

27 All ¹H NMR spectra were acquired using Bruker DRX600 spectrometers (Rheinstetten,
28
29 Germany) running under TopSpin, with either a 5 mm TXI probe operating at 600.13
30
31 MHz or a 5 mm BBI probe operating at 419 600.44 MHz. All runs were carried out
32
33 using Bruker BACS60 sample handling automation; prior to each run the 90° pulse
34
35 length was determined and set for the run. The field frequency was locked on D2O as
36
37 solvent. In all experiments, water suppression was carried out by noise irradiation
38
39 during the 2 s recycle delay (RD). For all experiments, 128 scans were recorded into 32
40
41 K data points with a spectral width of 20 ppm, and an exponential function was applied
42
43 to the FID prior to the Fourier transformation, which resulted in a line broadening of 0.3
44
45 Hz. All plasma NMR spectra were automatically phased, baseline-corrected and
46
47 referenced to the center of the α -glucose anomeric doublet (δ 5.23) for plasma, using an
48
49 in-house MatLab (The MathWorks, Natick, Massachusetts) scripts. Baseline and peak
50
51 alignment quality control was done by individual verification for each spectrum and
52
53
54
55
56 occasionally a spectrum was manually adjusted.
57
58
59
60

RNA expression

RNA purification, gene expression procedures and analyses were carried out, as previously described [13,14]. Briefly, RNA purification was performed using the RNeasy Lipid Tissue Mini Kit (QIAGEN, Izasa SA, Barcelona, Spain) and the integrity was checked by the Agilent Bioanalyzer (Agilent Technologies, Palo Alto, CA). Gene expression was assessed by real-time PCR using a LightCycler[®] 480 Real-Time PCR System (Roche Diagnostics SL, Barcelona, Spain), using TaqMan[®] technology suitable for relative genetic expression quantification. The commercially available and pre-validated TaqMan[®] primer/probe sets used were detailed elsewhere [13,14].

Analytical methods

Serum glucose, insulin, HOMA-IR, HDL-cholesterol, fasting triglycerides and C-reactive protein concentration measurement were performed as previously described [13,14].

Statistical analyses

Statistical analyses were performed using SPSS 12.0 software. Unless otherwise stated, descriptive results of continuous variables are expressed as mean and SD for Gaussian variables or median and interquartile range. Parameters that did not fulfil normal distribution criteria were Log transformed to improve symmetry for subsequent analyses. The relation between variables was analyzed by simple correlation (Pearson's test and Spearman's test) and multiple regression analyses in a stepwise manner. Levels of statistical significance were set at $p < 0.05$.

Results

Anthropometrical and metabolic parameters are described in Table 1. In obese subjects with insulin resistance, gut Firmicutes relative abundance (RA) was decreased, whereas Bacteroidetes and Proteobacteria RA were increased compared with insulin sensitive subjects (Table 1). These observations were in parallel to reduced SAT *SLC2A4*, (GLUT4), a tendency towards reduced VAT IRS-1 ($P=0.06$) and reduced SAT *PRDM16* mRNA among insulin resistant subjects (Table 1).

Firmicutes RA was negatively correlated with HbA_{1c} ($r= -0.37$, $p=0.03$) and fasting triglycerides ($r= -0.35$, $p=0.04$), whereas Proteobacteria RA was positively correlated with HOMA-IR ($r= 0.35$, $p=0.04$) and negatively with insulin sensitivity ($r=-0.38$, $p=0.03$).

Interestingly, Firmicutes RA was positively correlated with markers of brown adipocytes [*PRDM16* ($r=0.42$, $p=0.01$), *UCPI* ($r=0.45$, $p=0.01$) and *DIO2* ($r=0.38$, $p=0.03$)] (Figure 1a-b) in SAT, but not in VAT [*PRDM16* ($r=-0.05$, $p=0.7$), *UCPI* ($r=0.02$, $p=0.9$) and *DIO2* ($r=-0.04$, $p=0.8$)]. Otherwise, Bacteroidetes RA was negatively correlated with *PRDM16* ($r=-0.37$, $p=0.03$), *UCPI* ($r=-0.38$, $p=0.03$) and *DIO2* ($r=-0.35$, $p=0.05$) mRNA levels (Figure 1c-d).

The associations were specific for browning, because no significant correlations were found between Firmicutes RA and markers of adipogenesis [*ADIPOQ* ($r=0.15$, $p=0.4$), *PPARG* ($r=-0.07$, $p=0.6$)], insulin sensitivity [*IRS1* ($r=0.17$, $p=0.3$) and *SLC2A4* ($r=0.13$, $p=0.4$)] or inflammation [TNF ($r=-0.16$, $p=0.4$) and IL6 ($r=0.21$, $p=0.3$)] in SAT or VAT (*ADIPOQ* ($r=-0.16$, $p=0.4$), *PPARG* ($r=-0.02$, $p=0.9$), *SLC2A4* ($r=0.03$, $p=0.8$), *IRS1* ($r=0.24$, $p=0.1$), TNF ($r=-0.03$, $p=0.8$) and IL6 ($r=-0.18$, $p=0.4$)).

1
2
3 *PRDM16* mRNA levels were also negatively correlated with Proteobacteria RA ($r=-$
4 0.37, $p=0.03$), but this association did not remain significant after controlling for age,
5
6 BMI or insulin sensitivity.
7

8
9 Multivariate regression analysis indicated that Firmicutes RA contributed significantly
10 to SAT *PRDM16* and *UCP1* mRNA variance after controlling for age, sex, BMI, insulin
11 sensitivity (M value), HbA_{1c}, and diabetes, hypertension or dyslipemia therapy (Table
12 2). Firmicutes RA also contributed significantly to SAT *DIO2* mRNA variance after
13 controlling for age, sex, insulin sensitivity (M value) and HbA_{1c} (Table 2).
14
15

16
17 To explore the possible mechanism underlying the relationship between microbiota and
18 adipose tissue browning, and considering the effects of circulating acetate on adipocyte
19 browning and fat oxidation [20-24], its major abundance compared to other important
20 gut microbial-derived short-chain fatty acids (propionate and butyrate) [25], and its
21 capacity to transfer from colon into bloodstream [20], circulating acetate levels in
22 plasma were examined. Circulating acetate levels were significantly increased in insulin
23 sensitive obese participants (Figure 2a) in positive correlation with gut Firmicutes RA,
24 SAT *PRDM16* mRNA and insulin sensitivity (Figure 2b-d), and negatively correlated
25 with HOMA-IR and fasting triglycerides (Figure 2e-f).
26
27

28
29 Finally, within families from firmicutes phylum, only those bacteria from
30 Ruminococcaceae family, which was the most abundant (Figure 2g), were positively
31 correlated with insulin sensitivity ($r=0.48$, $p=0.01$), plasma acetate ($r=0.52$, $p=0.005$)
32 and *PRDM16* mRNA levels ($r=0.43$, $p=0.02$), and negatively correlated with fasting
33 triglycerides ($r= -0.37$, $p= 0.04$) (Figure 3). Multivariate regression analysis indicated
34 that plasma acetate contributed significantly to SAT *PRDM16* mRNA variance after
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 controlling for age, sex, BMI, insulin sensitivity (M value) and diabetes, hypertension or
4
5 dyslipemia therapy, but not after controlling for family Ruminococcaceae RA (Table 3).
6
7

8 9 10 **Discussion**

11
12 In the current study, we investigated gut microbial composition and markers of adipose
13 tissue browning in morbidly obese subjects. To the best of our knowledge, no previous
14 studies have explored these associations in humans. The main findings indicate that
15 Firmicutes RA, specifically those bacteria from Ruminococcaceae family, was
16 positively associated with insulin action in parallel to increased markers of browning in
17 SAT. A previous study demonstrated that cold exposure, with its associated increased
18 energy harvest, led to dramatic changes of the microbiota composition in mice,
19 increasing Firmicutes *vs.* Bacteroidetes RA in association with browning of white
20 adipose tissue. These changes resulted in improved insulin sensitivity, increased energy
21 expenditure and fat loss [8]. In humans, this shift in microbial composition has been
22 previously associated with significantly increased energy harvest [26]. A significant
23 reduction in bacterial species from Firmicutes phylum and increased RA of bacterial
24 species from Proteobacteria and Bacteroidetes phyla are known to be associated with
25 insulin resistance and type 2 diabetes [27-33]. Markers of browning (PRDM16 and
26 UCP1) in SAT have been also linked to insulin sensitivity [11-13].
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 The mechanisms to explain the relationship between Ruminococcaceae RA and markers
47 of browning in SAT remain to be elucidated. Gut microbial-derived SCFAs have been
48 demonstrated to improve insulin sensitivity through the enhancing of fat oxidation and
49 energy expenditure [20,25,34]. Acetate seems to exert a crucial role in brown adipocyte
50 differentiation through the induction of mitochondrial biogenesis [21], leading to
51
52
53
54
55
56
57
58
59
60

1
2
3 increased oxygen consumption rate in white and brown adipocytes and skeletal muscle
4
5 from Otsuka Long-Evans Tokushima Fatty rats [22]; and increased browning of SAT in
6
7 parallel to increased insulin sensitivity in mice [23]. Acetate administration also resulted
8
9 in increased expression of browning genes (PRDM16, UCP1 and DIO2) in 3T3-L1 cells
10
11 and in white adipose tissue from obese diabetic KK-Ay mice in parallel to increased
12
13 whole-body oxygen consumption [24], leading to enhanced fat oxidation [25,35,36].

14
15
16 In animal models, reduced Firmicutes RA and serum acetate and increased
17
18 Bacteroidetes RA were concomitant to lower glucose infusion rate during a
19
20 hyperinsulinemic euglycemic clamp in mice [37]. The recovery of serum acetate levels
21
22 (using antibiotics or acetate administration) led to improved insulin sensitivity and
23
24 attenuated adipose tissue macrophage infiltration in adipose tissue [37].
25
26

27
28 In humans, distal colonic acetate infusion resulted in increased whole-body fat
29
30 oxidation [28]. Multivariate regression analysis indicated that the association between
31
32 plasma acetate and SAT *PRDM16* mRNA remained significant after controlling for age,
33
34 sex, BMI, insulin sensitivity (*M* value) and diabetes, hypertension or dyslipemia
35
36 therapy, but not after controlling for family Ruminococcaceae RA, suggesting this
37
38 bacterial family as the most important source of plasma acetate. In agreement with this
39
40 suggestion, one recent study found acetate biosynthesis to be correlated with Firmicutes
41
42 RA [38], being those bacteria from Ruminococcaceae family (which was the most
43
44 abundant) considered as important acetate-producing bacteria in rumen studies [39].
45
46 This is the first study, to our knowledge, to describe associations among glucose
47
48 infusion rate during euglycemic hyperinsulinemic clamp, gut microbiota composition
49
50 and markers of browning in SAT in humans.
51
52
53
54
55
56
57
58
59
60

1
2
3 The absence of association between Firmicutes RA and markers of browning in VAT
4
5 might be explained by the lower capacity of this fat depot to develop browning [9,40].
6
7 The current study extends to humans previous observations in mice regarding gut
8
9 microbiota composition and adipose tissue browning [8], and suggests a possible
10
11 mechanism to explain them through increased plasma acetate levels. Further
12
13 intervention studies exploring the possible role of acetate in adipose tissue browning
14
15 and insulin sensitivity are required to confirm current findings.
16
17

20 21 **Author contributions**

22
23 JMM-N researched data, wrote and edited the manuscript; MS, RB, MD, MF researched data,
24
25 contributed to discussion and reviewed manuscript; VB-B, VA, RBa, MC, JL, FO, MS-M
26
27 researched data; WR contributed to discussion; JMF-R researched data and wrote and reviewed
28
29 the manuscript.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ACKNOWLEDGMENTS

We are indebted to the IDIBGI Biobank, integrated in the Spanish National Biobank Network, for the sample and data procurement. We acknowledge the technical assistance of E. Loshuertos and O. Rovira (both from Endocrinology, IdIBGi, Spain).

This work was partially supported by research grant PI15/01934 from the Instituto de Salud Carlos III from Spain and was also supported by FEDER funds. CIBEROBN Fisiopatología de la Obesidad y Nutrición is an initiative from the Instituto de Salud Carlos III from Spain. The authors declared no conflict of interest.

For Peer Review

References

- [1] Hamady, M., Knight, R., Microbial community profiling for human microbiome projects: Tools, techniques, and challenges. *Genome Res.* 2009, *19*, 1141–1152.
- [2] Ursell, L.K., Haiser, H.J., Van Treuren, W., Garg, N., Reddivari, L., Vanamala, J., Dorrestein, P.C., Turnbaugh, P.J., Knight, R., The intestinal metabolome: an intersection between microbiota and host. *Gastroenterology* 2014, *146*, 1470–1476.
- [3] Tagliabue, A., Eli, M., The role of gut microbiota in human obesity: recent findings and future perspectives. *Nutr. Metab. Cardiovasc. Dis.* 2013, *23*, 160–168.
- [4] Rosenbaum, M., Knight, R., Leibel, R.L., The gut microbiota in human energy homeostasis and obesity. *Trends Endocrinol. Metab.* 2015, *26*, 493–501.
- [5] Ravussin, Y., Koren, O., Spor, A., LeDuc, C., Gutman, R., Stombaugh, J., Knight, R., Ley, R.E., Leibel, R.L., Response of gut microbiota to diet composition and weight loss in lean and obese mice. *Obesity (Silver Spring)* 2012, *20*, 736–747.
- [6] Palleja, A., Kashani, A., Allin, K.H., Nielsen, T., Zhang, C., Li, Y., Brach, T., Liang, S., Feng, Q., Jørgensen, N.B., Bojsen-Møller, K.N., Dirksen, C., Burgdorf, K.S., Holst, J.J., Madsbad, S., Wang, J., Pedersen, O., Hansen, T., Arumugam, M., Roux-en-Y gastric bypass surgery of morbidly obese patients induces swift and persistent changes of the individual gut microbiota. *Genome Med.* 2016, *8*, 67.
- [7] Walters, W.A., Xu, Z., Knight, R., Meta-analyses of human gut microbes associated with obesity and IBD. *FEBS Lett.* 2014, *588*, 4223–4233.
- [8] Chevalier, C., Stojanović, O., Colin, D.J., Suarez-Zamorano, N., Tarallo, V., Veyrat-Durebex, C., Rigo, D., Fabbiano, S., Stevanović, A., Hagemann, S., Montet, X., Seimbille, Y., Zamboni, N., Hapfelmeier, S., Trajkovski, M., Gut Microbiota Orchestrates Energy Homeostasis during Cold. *Cell* 2015, *163*, 1360–1374.

1
2
3 [9] Cohen, P., Levy, J.D., Zhang, Y., Frontini, A., Kolodin, D.P., Svensson, K.J., Lo,
4 J.C., Zeng, X., Ye, L., Khandekar, M.J., Wu, J., Gunawardana, S.C., Banks, A.S.,
5 Camporez, J.P., Jurczak, M.J., Kajimura, S., Piston, D.W., Mathis, D., Cinti, S.,
6 Shulman, G.I., Seale, P., Spiegelman, B.M., Ablation of PRDM16 and beige adipose
7 causes metabolic dysfunction and a subcutaneous to visceral fat switch. *Cell* 2014, *156*,
8 304-316.
9

10
11
12
13
14 [10] Kumari, M., Wang, X., Lantier, L., Lyubetskaya A, Eguchi J, Kang S, Tenen D,
15 Roh HC, Kong X, Kazak L, Ahmad R, Rosen ED. IRF3 promotes adipose inflammation
16 and insulin resistance and represses browning. *J. Clin. Invest.* 2016, *126*, 2839-2854.
17

18
19
20 [11] Stanford, K.I., Middelbeek, R.J., Goodyear, L.J., Exercise Effects on White
21 Adipose Tissue: Beiging and Metabolic Adaptations. *Diabetes* 2015, *64*, 2361-2368.
22

23
24 [12] Sidossis, L., Kajimura, S., Brown and beige fat in humans: thermogenic adipocytes
25 that control energy and glucose homeostasis. *J. Clin. Invest.* 2015, *125*, 478-486.
26

27
28 [13] Moreno-Navarrete, J.M., Ortega, F., Moreno, M., Xifra, G., Ricart, W., Fernández-
29 Real, J.M., PRDM16 sustains white fat gene expression profile in human adipocytes in
30 direct relation with insulin action. *Mol. Cell. Endocrinol.* 2015, *405*, 84-93.
31

32
33 [14] Moreno-Navarrete, J.M., Novelle, M.G., Catalán, V., Ortega, F., Moreno, M.,
34 Gomez-Ambrosi, J., Xifra, G., Serrano, M., Guerra, E., Ricart, W., Frühbeck, G.,
35 Diéguez, C., Fernández-Real, J.M., Insulin resistance modulates iron-related proteins in
36 adipose tissue. *Diabetes Care* 2014, *37*, 1092-1100.
37

38
39 [15] Serino, M., Luche, E., Gres, S., Baylac, A., Bergé, M., Cenac, C., Waget, A.,
40 Klopp, P., Iacovoni, J., Klopp, C., Mariette, J., Bouchez, O., Lluch, J., Ouarné, F.,
41 Monsan, P., Valet, P., Roques, C., Amar, J., Bouloumié, A., Théodorou, V., Burcelin,
42 R., Metabolic adaptation to a high-fat diet is associated with a change in the gut
43 microbiota. *Gut* 2012, *61*, 543-553.
44

45
46 [16] Schmieder, R., Edwards, R., Quality control and preprocessing of metagenomic
47 datasets. *Bioinformatics* 2011, *27*, 863-864.
48
49
50
51
52

1
2
3 [17] Aronesty, E., Command-line tools for processing biological sequencing data. ea-
4 utils: FASTQ processing utilities. URL: <http://code.google.com/p/ea-utils>. 2011
5 (Accessed 2016).
6
7

8
9 [18] Langmead, B., Salzberg, S.L., Fast gapped-read alignment with Bowtie 2. *Nat.*
10 *Methods* 2012, 9, 357–359.
11

12
13 [19] Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D.,
14 Costello, E.K., Fierer, N., Peña, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A.,
15 Kelley, S.T., Knights, D., Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D.,
16 Muegge, B.D., Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A.,
17 Widmann, J., Yatsunenko, T., Zaneveld, J., Knight, R., QIIME allows analysis of high-
18 throughput community sequencing data. *Nat Methods* 2010, 7, 335–336.
19
20
21
22
23

24
25 [20] van der Beek, C.M., Canfora, E.E., Lenaerts, K., Troost, F.J., Olde Damink,
26 S.W.M., Holst, J.J., Masclee, A.A., Dejong, C.H., Blaak, E.E., Distal, not proximal,
27 colonic acetate infusions promote fat oxidation and improve metabolic markers in
28 overweight/obese men. *Clin Sci (Lond)* 2016, 130, 2073-2082.
29
30
31

32
33 [21] Hu, J., Kyrou, I., Tan, B.K., Dimitriadis, G.K., Ramanjaneya, M., Tripathi, G.,
34 Patel, V., James, S., Kawan, M., Chen, J., Randeve, H.S., Short-Chain Fatty Acid
35 Acetate Stimulates Adipogenesis and Mitochondrial Biogenesis via GPR43 in
36 Brown Adipocytes. *Endocrinology* 2016, 157, 1881-1894.
37
38
39

40
41 [22] Yamashita, H., Maruta, H., Jozuka, M., Kimura, R., Iwabuchi, H., Yamato,
42 M., Saito, T., Fujisawa, K., Takahashi, Y., Kimoto, M., Hiemori, M., Tsuji, H.,
43 Effects of acetate on lipid metabolism in muscles and adipose tissues of type 2
44 diabetic Otsuka Long-Evans Tokushima Fatty (OLETF) rats. *Biosci. Biotechnol.*
45 *Biochem.* 2009, 73, 570-576.
46
47
48
49

50
51 [23] Sahuri-Arisoylu, M., Brody, L.P., Parkinson, J.R., Parkes, H., Navaratnam, N.,
52 Miller, A.D., Thomas, E.L., Frost, G., Bell, J.D., Reprogramming of hepatic fat
53 accumulation and 'browning' of adipose tissue by the short-chain fatty acid acetate. *Int*
54 *J Obes (Lond)* 2016, 40, 955-963.
55
56
57
58
59
60

1
2
3 [24] Hanatani, S., Motoshima, H., Takaki, Y., Kawasaki, S., Igata, M., Matsumura, T.,
4 Kondo, T., Senokuchi, T., Ishii, N., Kawashima, J., Kukidome, D., Shimoda, S.,
5 Nishikawa, T., Araki, E., Acetate alters expression of genes involved in beige
6 adipogenesis in 3T3-L1 cells and obese KK-Ay mice. *J. Clin. Biochem. Nutr.* 2016, *59*,
7 207-214.

8
9
10
11
12 [25] Canfora, E.E., Jocken, J.W., Blaak, E.E., Short-chain fatty acids in control of body
13 weight and insulin sensitivity. *Nat. Rev. Endocrinol.* 2015, *11*, 577-591.

14
15
16
17 [26] Jumpertz, R., Le, D.S., Turnbaugh, P.J., Trinidad, C., Bogardus, C., Gordon, J.I.,
18 Krakoff, J., Energy-balance studies reveal associations between gut microbes, caloric
19 load, and nutrient absorption in humans. *Am. J. Clin. Nutr.* 2011, *94*, 58-65.

20
21
22
23 [27] Pedersen, H.K., Gudmundsdottir, V., Nielsen, H.B., Hyotylainen, T., Nielsen, T.,
24 Jensen, B.A., Forslund, K., Hildebrand, F., Prifti, E., Falony, G., Le Chatelier, E.,
25 Levenez, F., Doré, J., Mattila, I., Plichta, D.R., Pöhö, P., Hellgren, L.I., Arumugam, M.,
26 Sunagawa, S., Vieira-Silva, S., Jørgensen, T., Holm, J.B., Trošt, K.; MetaHIT
27 Consortium, Kristiansen, K., Brix, S., Raes, J., Wang, J., Hansen, T., Bork, P., Brunak,
28 S., Oresic, M., Ehrlich, S.D., Pedersen, O., Human gut microbes impact host serum
29 metabolome and insulin sensitivity. *Nature* 2016, *21*, 535:376-381.

30
31
32
33 [28] Haro, C., Montes-Borrego, M., Rangel-Zúñiga, O.A., Alcalá-Díaz, J.F., Gómez-
34 Delgado, F., Pérez-Martínez, P., Delgado-Lista, J., Quintana-Navarro, G.M., Tinahones,
35 F.J., Landa, B.B., López-Miranda, J., Camargo, A., Pérez-Jiménez, F., Two Healthy
36 Diets Modulate Gut Microbial Community Improving Insulin Sensitivity in a Human
37 Obese Population. *J. Clin. Endocrinol. Metab.* 2016, *101*, 233-242.

38
39
40
41 [29] Fugmann, M., Breier, M., Rottenkolber, M., Banning, F., Ferrari, U., Sacco, V.,
42 Grallert, H., Parhofer, K.G., Seissler, J., Clavel, T., Lechner, A., The stool microbiota of
43 insulin resistant women with recent gestational diabetes, a high risk group for type 2
44 diabetes. *Sci. Rep.* 2015, *5*, 13212.

45
46
47
48 [30] Korem, T., Zeevi, D., Suez, J., Weinberger, A., Avnit-Sagi, T., Pompan-Lotan, M.,
49 Matot, E., Jona, G., Harmelin, A., Cohen, N., Sirota-Madi, A., Thaïss, C.A., Pevsner-

1
2
3 Fischer, M., Sorek, R., Xavier, R.J., Elinav, E., Segal, E., Growth dynamics of gut
4 microbiota in health and disease inferred from single metagenomic samples. *Science*
5 2015, *349*, 1101-1106.
6
7

8
9 [31] Delzenne, N.M., Cani, P.D., Everard, A., Neyrinck, A.M., Bindels, L.B., Gut
10 microorganisms as promising targets for the management of type 2 diabetes.
11 *Diabetologia* 2015, *58*, 2206-2217.
12
13

14
15 [32] Karlsson, F.H., Tremaroli, V., Nookaew, I., Bergström, G., Behre, C.J., Fagerberg,
16 B., Nielsen, J., Bäckhed, F., Gut metagenome in European women with normal,
17 impaired and diabetic glucose control. *Nature* 2013, *498*, 99-103.
18
19

20
21 [33] Qin, J., Li, Y., Cai, Z., Li, S., Zhu, J., Zhang, F., Liang, S., Zhang, W., Guan, Y.,
22 Shen, D., Peng, Y., Zhang, D., Jie, Z., Wu, W., Qin, Y., Xue, W., Li, J., Han, L., Lu, D.,
23 Wu, P., Dai, Y., Sun, X., Li, Z., Tang, A., Zhong, S., Li, X., Chen, W., Xu, R., Wang,
24 M., Feng, Q., Gong, M., Yu, J., Zhang, Y., Zhang, M., Hansen, T., Sanchez, G., Raes,
25 J., Falony, G., Okuda, S., Almeida, M., LeChatelier, E., Renault, P., Pons, N., Batto,
26 J.M., Zhang, Z., Chen, H., Yang, R., Zheng, W., Li, S., Yang, H., Wang, J., Ehrlich,
27 S.D., Nielsen, R., Pedersen, O., Kristiansen, K., Wang, J., A metagenome-wide
28 association study of gut microbiota in type 2 diabetes. *Nature* 2012, *490*, 55-60.
29
30
31
32
33
34
35

36 [34] Shirouchi, B., Nagao, K., Umegatani, M., Shiraishi, A., Morita, Y., Kai, S.,
37 Yanagita, T., Ogawa, A., Kadooka, Y., Sato, M., Probiotic *Lactobacillus gasseri*
38 SBT2055 improves glucose tolerance and reduces body weight gain in rats by
39 stimulating energy expenditure. *Br. J. Nutr.* 2016, *116*, 451-458.
40
41
42
43

44 [35] Yamaguchi, Y., Adachi, K., Sugiyama, T., Shimozaoto, A., Ebi, M., Ogasawara, N.,
45 Funaki, Y., Goto, C., Sasaki, M., Kasugai, K., Association of Intestinal Microbiota with
46 Metabolic Markers and Dietary Habits in Patients with Type 2 Diabetes. *Digestion*
47 2016, *94*, 66-72.
48
49
50

51
52 [36] Lu, Y., Fan, C., Li, P., Lu, Y., Chang, X., Qi, K., Short Chain Fatty Acids Prevent
53 High-fat-diet-induced Obesity in Mice by Regulating G Protein-coupled Receptors and
54 Gut Microbiota. *Sci. Rep.* 2016, *6*, 37589.
55
56
57
58
59
60

1
2
3 [37] Carvalho, B.M., Guadagnini, D., Tsukumo, D.M.L., Schenka, A.A., Latuf-Filho,
4 P., Vassallo, J., Dias, J.C., Kubota, L.T., Carvalheira, J.B.C., Saad, M.J.A., Modulation
5 of gut microbiota by antibiotics improves insulin signalling in high-fat fed mice.
6 *Diabetologia* 2012, 55, 2823-2834.
7
8

9
10
11 [38] Goffredo, M., Mass, K., Parks, E.J., Wagner, D.A., McClure, E.A., Graf, J.,
12 Savoye, M., Pierpont, B., Cline, G., Santoro, N., Role of Gut Microbiota and Short
13 Chain Fatty Acids in Modulating Energy Harvest and Fat Partitioning in Youth. *J. Clin.*
14 *Endocrinol. Metab.* 2016, 101, 4367-4376.
15
16

17
18
19 [39] Gagen, E.J., Padmanabha, J., Denman, S.E., McSweeney, C.S., Hydrogenotrophic
20 culture enrichment reveals rumen Lachnospiraceae and Ruminococcaceae acetogens
21 and hydrogen-responsive Bacteroidetes from pasture-fed cattle. *FEMS Microbiol. Lett.*
22 2015, 362, 14.
23
24

25
26
27 [40] Gavaldà-Navarro, A., Moreno-Navarrete, J.M., Quesada-López, T., Cairó, M.,
28 Giralt, M., Fernández-Real, J.M., Villarroya, F., Lipopolysaccharide-binding protein is
29 a negative regulator of adipose tissue browning in mice and humans. *Diabetologia*
30 2016, 59, 2208-2218.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Anthropometric, clinical parameters, microbial abundance and SAT gene expressions according to insulin resistance.

	Insulin sensitive	Insulin resistant	<i>P</i>
N	14	20	
Age (years)	48.7 ± 11.1	47.4 ± 7.6	0.7
BMI (kg/m ²)	42.8 ± 7.1	45.4 ± 5.6	0.2
Waist circumference (cm)	123.9 ± 10.7	129.1 ± 11.1	0.2
Fasting glucose (mg/dl)	95.5 (87-100.2)	102 (92.5-129.5)	0.02
HbA _{1c} (%)	5.5 ± 0.5	6.4 ± 1.6	0.02
HbA _{1c} (mmol/mol)	37 ± 3.2	46 ± 9.5	0.02
HDL-cholesterol (mg/dl)	48.0 ± 10.1	46.6 ± 14.8	0.7
Fasting triglycerides (mg/dl)	88.5 (66-147)	134 (99.5-165)	0.1
HOMA-IR	1.6 (1.3-2.1)	4.8 (4.1-8.3)	0.03
M (mg/kg/min)	5.8 ± 2.7	2.7 ± 1.3	<0.0001
Firmicutes (%)	64.9 (60.4-78.7)	47.3 (42.1-64.5)	0.001
Bacteroidetes (%)	29.3 (18.8-36.8)	36.7 (27.1-49.7)	0.02
Proteobacteria (%)	1.27 (0.85-2.98)	2.32 (1.99-5.89)	0.03
Actinobacteria (%)	0.45 (0.19-0.76)	0.28 (0.14-0.61)	0.2
Fusobacteria (%)	0.47 (0.11-1.43)	0.67 (0.24-3.21)	0.4
Shannon index	3.34 ± 0.17	3.23 ± 0.33	0.2
SAT <i>ADIPOQ</i> (RU)	2.42 ± 0.7	2.37 ± 0.7	0.8
SAT <i>PPARG</i> (RU)	0.036 ± 0.01	0.040 ± 0.02	0.6
SAT <i>SLC2A4</i> (RU)	0.053 ± 0.02	0.036 ± 0.01	0.04
SAT <i>IRS1</i> (RU)	0.0087 ± 0.005	0.0078 ± 0.003	0.4
SAT <i>TNF</i> (RU)	0.0039 ± 0.001	0.0046 ± 0.003	0.4
SAT <i>IL6</i> (RU) 10 ⁻⁴	5 (3-20)	13 (5-44)	0.2
SAT <i>PRDM16</i> (RU)	0.0037 ± 0.001	0.0027 ± 0.001	0.03
SAT <i>UCPI</i> (RU) * 10 ⁻⁵	2.5 (2-3)	2 (1-3)	0.4
SAT <i>DIO2</i> (RU) * 10 ⁻⁴	6.4 (4-7)	6.9 (3-11.1)	0.4
VAT <i>ADIPOQ</i> (RU)	1.53 ± 0.47	1.36 ± 0.5	0.3
VAT <i>PPARG</i> (RU)	0.021 ± 0.008	0.020 ± 0.013	0.8
VAT <i>SLC2A4</i> (RU)	0.032 ± 0.02	0.023 ± 0.01	0.1
VAT <i>IRS1</i> (RU)	0.0087 ± 0.003	0.0069 ± 0.002	0.06
VAT <i>TNF</i> (RU)	0.0052 ± 0.002	0.0078 ± 0.005	0.1
VAT <i>IL6</i> (RU) 10 ⁻⁴	9.9 (7.7-20.8)	10.2 (6.8-39.5)	0.8
VAT <i>PRDM16</i> (RU)	0.0043 ± 0.002	0.0034 ± 0.001	0.09
VAT <i>UCPI</i> (RU) * 10 ⁻⁵	8.5 (4.2-25.2)	5 (3-8.5)	0.08
VAT <i>DIO2</i> (RU) * 10 ⁻⁴	2.6 (1.9-4.6)	2.2 (1.6-3.8)	0.8

Table 2. Multivariate regression analyses to predict SAT *PRDM16*, *UCPI* and *DIO2* mRNA levels.

SAT <i>PRDM16</i> (RU)	β	p	β	p	β	p	β	p
Firmicutes RA (%)	0.45	0.01	0.45	0.01	0.43	0.01	0.43	0.01
Age (years)	-0.01	0.9	-	-	-	-	-	-
BMI (kg/m ²)	-	-	-0.09	0.5	-	-	-	-
M (mg/kg/min)	-	-	-	-	0.26	0.1	-	-
HbA _{1c} (%)	-	-	-	-	-	-	-0.06	0.7
Adjusted R ²	0.149 (14.9%)		0.158 (15.8%)		0.189 (18.9%)		0.153 (15.3%)	
P value	0.03		0.03		0.02		0.03	
SAT <i>PRDM16</i> (RU)	β	p	β	p	β	p	β	p
Firmicutes RA (%)	0.43	0.01	0.41	0.02	0.43	0.01	0.46	0.008
Sex	0.27	0.1	-	-	-	-	-	-
Diabetes therapy	-	-	-0.14	0.3	-	-	-	-
Hypertension therapy	-	-	-	-	-0.12	0.4	-	-
Dyslipemia therapy	-	-	-	-	-	-	0.04	0.8
Adjusted R ²	0.230 (23%)		0.172 (17.2%)		0.167 (16.7%)		0.154 (15.4%)	
P value	0.009		0.02		0.02		0.03	
SAT <i>UCPI</i> (RU)	β	p	β	p	β	p	β	p
Firmicutes RA (%)	0.45	0.01	0.37	0.03	0.41	0.03	0.43	0.02
Age (years)	0.05	0.7	-	-	-	-	-	-
BMI (kg/m ²)	-	-	-0.39	0.03	-	-	-	-
M (mg/kg/min)	-	-	-	-	-0.11	0.6	-	-
HbA _{1c} (%)	-	-	-	-	-	-	-0.06	0.7
Adjusted R ²	0.146 (14.6%)		0.295 (29.5%)		0.119 (11.9%)		0.147 (14.7%)	
P value	0.05		0.005		0.08		0.04	
SAT <i>UCPI</i> (RU)	β	p	β	p	β	p	β	p
Firmicutes RA (%)	0.45	0.01	0.49	0.01	0.46	0.01	0.49	0.01
Sex	0.11	0.5	-	-	-	-	-	-
Diabetes therapy	-	-	0.15	0.4	-	-	-	-
Hypertension therapy	-	-	-	-	0.08	0.6	-	-
Dyslipemia therapy	-	-	-	-	-	-	0.15	0.4
Adjusted R ²	0.145 (14.5%)		0.155 (15.5%)		0.139 (13.9%)		0.156 (15.6%)	
P value	0.05		0.04		0.05		0.04	
SAT <i>DIO2</i> (RU)	β	p	β	p	β	p	β	p
Firmicutes RA (%)	0.39	0.03	0.35	0.06	0.51	0.01	0.40	0.03
Age (years)	-0.36	0.05	-	-	-	-	-	-

BMI (kg/m ²)	-	-	-0.09	0.6	-	-	-	-
M (mg/kg/min)	-	-	-	-	-0.31	0.1	-	-
HbA _{1c} (%)	-	-	-	-	-	-	0.14	0.4
Adjusted R ²	0.201 (20.1%)		0.081 (8.1)		0.222 (22.2%)		0.083 (8.3%)	
P value	0.02		0.16		0.03		0.11	
SAT <i>DIO2</i> (RU)	β	p	β	p	β	p	β	p
Firmicutes RA (%)	0.38	0.04	0.30	0.1	0.27	0.1	0.28	0.1
Sex	-0.26	0.1	-	-	-	-	-	-
Diabetes therapy	-	-	-0.18	0.3	-	-	-	-
Hypertension therapy	-	-	-	-	-0.45	0.01	-	-
Dyslipemia therapy	-	-	-	-	-	-	-0.32	0.09
Adjusted R ²	0.142 (14.2%)		0.097 (9.7%)		0.276 (27.6%)		0.166 (16.6%)	
P value	0.05		0.1		0.008		0.04	

Table 3. Multivariate regression analyses to investigate the contribution of circulating acetate on SAT *PRDM16* mRNA levels.

SAT <i>PRDM16</i> (RU)	β	p	β	p	β	p	β	p
Plasma acetate (ppm)	0.33	0.07	0.42	0.02	0.42	0.02	0.38	0.04
Ruminococcaceae RA	0.30	0.08	-	-	-	-	-	-
Age (years)	-	-	-0.01	0.9	-	-	-	-
BMI (kg/m ²)	-	-	-	-	-0.02	0.8		
M (mg/kg/min)	-	-	-	-	-	-	0.15	0.5
Adjusted R ²	0.230 (23%)		0.124 (12.4%)		0.124 (12.4%)		0.176 (17.6%)	
P value	0.01		0.05		0.05		0.04	
SAT <i>PRDM16</i> (RU)	β	p	β	p	β	p	β	p
Plasma acetate (ppm)	0.39	0.02	0.37	0.04	0.43	0.01	0.42	0.02
Sex	0.26	0.1	-	-	-	-	-	-
Diabetes therapy	-	-	-0.13	0.4	-	-	-	-
Hypertension therapy	-	-	-	-	-0.23	0.1	-	-
Dyslipemia therapy	-	-	-	-	-	-	-0.03	0.8
Adjusted R ²	0.195 (19.5%)		0.152 (15.2%)		0.183 (18.3%)		0.125 (12.5%)	
P value	0.01		0.04		0.02		0.05	





Figure legends

Figure 1. a-d) Bivariate correlations among *PRDM16* and *UCPI* mRNA levels and firmicutes (a-b) and bacteroidetes (c-d) relative abundance.

Figure 2. a) Circulating acetate levels according to insulin resistance. b-f) Bivariate correlations between circulating acetate and firmicutes relative abundance (b), SAT *PRDM16* mRNA levels (c), insulin sensitivity (d), HOMA-IR (e) and fasting triglycerides (f). g) Relative abundance of families belonging to Firmicutes phylum.








Figure 3. HeatMap showing bivariate correlations among the relative abundance of families belonging to Firmicutes phylum and metabolic parameters. * $p < 0.05$ and ** $p < 0.01$.

Molecular Nutrition and Food Research



Molecular Nutrition and Food Research



Figure 2



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

Figure 3

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43



Graphical abstract-text

This figure indicates the association among *Ruminococcaceae* family, plasma acetate levels, SAT expression of browning-related genes and insulin sensitivity, suggesting that increased *Ruminococcaceae*-enhanced acetate biosynthesis might promote SAT browning and systemic insulin sensitivity in morbidly obese subjects.

For Peer Review

Graphical Abstract

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43



In morbidly obese subjects



↑↑ Ruminococcaceae family RA ↑↑



↑↑ Plasma acetate levels ↑↑



↑↑ ↑↑ Expression of browning-related genes in SAT

Insulin sensitivity ↑↑ ↑↑