

# $h^*$ -vectors, Eulerian polynomials and stable polytopes of graphs

Christos A. Athanasiadis

Department of Mathematics  
University of Crete  
71409 Heraklion, Crete, Greece  
caa@math.uoc.gr

Submitted: Jul 5, 2004; Accepted: Sep 22, 2004; Published: Oct 7, 2004

Mathematics Subject Classifications: Primary 52B20; Secondary 05C17, 05E99, 13F20.

## Abstract

Conditions are given on a lattice polytope  $P$  of dimension  $m$  or its associated affine semigroup ring which imply inequalities for the  $h^*$ -vector  $(h_0^*, h_1^*, \dots, h_m^*)$  of  $P$  of the form  $h_i^* \geq h_{d-i}^*$  for  $1 \leq i \leq \lfloor d/2 \rfloor$  and  $h_{\lfloor d/2 \rfloor}^* \geq h_{\lfloor d/2 \rfloor + 1}^* \geq \dots \geq h_d^*$ , where  $h_i^* = 0$  for  $d < i \leq m$ . Two applications to order polytopes of posets and stable polytopes of perfect graphs are included.

## 1 Introduction

Let  $P$  be an  $m$ -dimensional convex polytope in  $\mathbb{R}^N$  having vertices with integer coordinates. It is a fundamental result due to Ehrhart [5, 6] that the function  $i(P, r) = \#(rP \cap \mathbb{Z}^N)$ , counting integer points in the  $r$ -fold dilate of  $P$ , is a polynomial in  $r$  of degree  $m$ , called the *Ehrhart polynomial* of  $P$ . Thus one can write

$$\sum_{r \geq 0} i(P, r) t^r = \frac{h_0^* + h_1^* t + \dots + h_m^* t^m}{(1-t)^{m+1}} \quad (1)$$

for certain integers  $h_i^*$ . Following Stanley [21] we call  $(h_0^*, h_1^*, \dots, h_m^*)$  the  $h^*$ -vector of  $P$  and denote it by  $h^*(P)$ . It is known that  $i(P, r)$  is the Hilbert function of a semistandard graded Cohen-Macaulay normal domain  $R_P$  called the semigroup ring of  $P$ ; see [3, Chapter 6] and [9, Chapter X]. In particular the integers  $h_i^*$  are nonnegative. Recall that a sequence  $(a_0, a_1, \dots, a_n)$  of real numbers is said to be *unimodal* if  $a_0 \leq \dots \leq a_j \geq \dots \geq a_n$  holds for some  $0 \leq j \leq n$ . Although  $h^*$ -vectors are not always unimodal, various results and conjectures concerning the unimodality of  $h^*(P)$  have appeared in the literature [8, 9, 19, 20]. For instance it would follow from [8, Conjecture 1.5] and [19, Conjecture 4a] that  $h^*(P)$  is unimodal if the semigroup ring  $R_P$  is standard and Gorenstein. Moreover,

it was conjectured by Hibi [9, p. 111] that  $h^*(P)$  is unimodal whenever it is symmetric, meaning that  $h_i^* = h_{m-i}^*$  for  $0 \leq i \leq m$ .

General conditions on an integer polytope  $P$ , inspired by the work of V. Reiner and V. Welker on order polytopes of graded posets [15], which guarantee that  $h^*(P)$  is unimodal were given in [1]. More precisely it was shown in [1] that if the pulling triangulation  $\Delta_\tau$  of  $P$  with respect to an ordering  $\tau = (v_1, v_2, \dots, v_p)$  of the vertices of  $P$  is unimodular (see Section 2 for basic definitions on triangulations) and for some  $n$  any facet of  $P$  contains exactly  $n - 1$  elements of  $\{v_1, v_2, \dots, v_n\}$  then  $h^*(P)$  is equal to the  $h$ -vector [24, Section 8.3] of a simplicial  $d$ -dimensional polytope, where  $d = m - n + 1$ , and hence that it is unimodal and satisfies  $h_i^* = h_{d-i}^*$  for  $0 \leq i \leq d$  by McMullen's  $g$ -theorem. The simplex with vertices  $\{v_1, v_2, \dots, v_n\}$  is called a *special simplex* for  $P$  in [1]. If  $v_1, v_2, \dots, v_n$  are any  $n$  vertices of  $P$  which are affinely independent we call their convex hull a *semispecial simplex* for  $P$  if any facet of  $P$  contains at least  $n - 1$  elements of  $\{v_1, v_2, \dots, v_n\}$ . It is the main goal of this paper to prove the following more general statement.

**Theorem 1.1** *Let  $P$  be an  $m$ -dimensional integer polytope with  $h^*$ -vector  $(h_0^*, h_1^*, \dots, h_m^*)$ . If  $\tau = (v_1, v_2, \dots, v_p)$  is a linear ordering of the vertices of  $P$  such that*

- (i) *the pulling triangulation  $\Delta_\tau$  of  $P$  is unimodular,*
- (ii)  *$\{v_1, v_2, \dots, v_n\}$  is the vertex set of a semispecial simplex for  $P$*

*and  $d = m - n + 1$ , then  $h_i^* \geq h_{d-i}^*$  for  $0 \leq i \leq \lfloor d/2 \rfloor$ ,  $h_{\lfloor d/2 \rfloor}^* \geq h_{\lfloor d/2 \rfloor + 1}^* \geq \dots \geq h_d^*$  and  $h_i^* = 0$  for  $d < i \leq m$ .*

The next corollary follows essentially from the case  $n = 1$  of Theorem 1.1.

**Corollary 1.2** *Let  $P$  be an  $m$ -dimensional integer polytope with  $h^*$ -vector  $(h_0^*, h_1^*, \dots, h_m^*)$ . If  $P$  has a unimodular pulling triangulation then  $h_i^* \geq h_{m-i}^*$  for  $1 \leq i \leq \lfloor m/2 \rfloor$  and  $h_{\lfloor m/2 \rfloor}^* \geq h_{\lfloor m/2 \rfloor + 1}^* \geq \dots \geq h_m^*$ .*

Pulling triangulations are specific examples of regular triangulations, so it is natural to ask for inequalities satisfied by  $h^*(P)$  under the existence of a regular unimodular triangulation of  $P$ . I am grateful to Takayuki Hibi [10] for informing me that the following statement was also proved by himself and Richard Stanley in 1999 (unpublished) by essentially the same argument as the one given in Section 2.

**Theorem 1.3** *Let  $P$  be an  $m$ -dimensional integer polytope with  $h^*$ -vector  $(h_0^*, h_1^*, \dots, h_m^*)$ . If  $P$  has a regular unimodular triangulation then  $h_i^* \geq h_{m-i+1}^*$  for  $1 \leq i \leq \lfloor (m+1)/2 \rfloor$ ,*

$$h_{\lfloor (m+1)/2 \rfloor}^* \geq \dots \geq h_{m-1}^* \geq h_m^*$$

and

$$h_i^* \leq \binom{h_1^* + i - 1}{i}$$

for  $0 \leq i \leq m$ . In particular, if  $h^*(P)$  is symmetric and  $P$  has a regular unimodular triangulation then  $h^*(P)$  is unimodal.

It was observed in [9, Example 36.4] that the last inequality in Theorem 1.3 does not hold for some integer polytopes. An example of a 0/1 polytope  $P$  with no regular unimodular triangulations such that  $R_P$  is standard was given in [13].

This paper is structured as follows. Section 2 includes the necessary definitions and background on convex polytopes and their triangulations and  $h^*$ -vectors as well as the proof of Theorem 1.3. The notion of triangulation of a polytope  $P$  we will use does not require that all vertices of the triangulation are necessarily vertices of  $P$  unless the contrary is explicitly stated. In Section 3 we introduce the concept of a semispecial simplex for  $P$  and prove a slight generalization of Theorem 1.1 (see Corollary 3.4). Specifically we drop the assumption that all vertices of a (special or) semispecial simplex and the elements of the sequence  $\tau$  which appears in Theorem 1.1 are vertices of  $P$ . Our proofs are based on a result of Kalai and Stanley (Lemma 2.2) on the  $h$ -vector of a Cohen-Macaulay subcomplex of the boundary complex of a simplicial polytope. Sections 4 and 5 include applications of Theorem 1.1 to order polytopes of (not necessarily graded) posets and to stable polytopes of perfect graphs, respectively. In Section 6 we state an analogue of Theorem 1.1 in the context of the affine semigroup ring of  $P$ .

## 2 Triangulations and $h^*$ -vectors

Before proving Theorem 1.3 we review some basic definitions and background on simplicial complexes and convex polytopes. For undefined terminology and more information on these topics we refer the reader to [7, 9, 22, 23, 24]. A *polytopal complex*  $\mathcal{F}$  [24, Section 8.1] is a finite, nonempty collection of convex polytopes in  $\mathbb{R}^N$  such that (i) any face of a polytope in  $\mathcal{F}$  is also in  $\mathcal{F}$  and (ii) the intersection of any two polytopes in  $\mathcal{F}$  is a (possibly empty) face of both. The elements of  $\mathcal{F}$  are its *faces* and those of dimension 0 are its *vertices*. The *dimension* of  $\mathcal{F}$  is the maximum dimension of a face. The complex  $\mathcal{F}$  is *pure* if all maximal (with respect to inclusion) faces of  $\mathcal{F}$  have the same dimension. The collection  $\mathcal{F}(P)$  of all faces of a polytope  $P$  is a pure polytopal complex, called the *face complex* of  $P$ , as is the collection  $\mathcal{F}(\partial P)$  of proper faces of  $P$ , called the *boundary complex*. The complex  $\mathcal{F}$  is a (geometric) *simplicial complex* if all faces of  $\mathcal{F}$  are simplices. Two simplicial complexes  $\Delta$  and  $\Delta'$  are said to be *combinatorially equivalent* if there exists a bijection  $\rho$  between the sets of faces of  $\Delta$  and  $\Delta'$  such that  $\rho$  and its inverse preserve inclusion. The  $h$ -vector  $h(\Delta) = (h_0, h_1, \dots, h_d)$  of a simplicial complex  $\Delta$  of dimension  $d - 1$  is defined by the formula

$$\sum_{i=0}^d f_{i-1}(x-1)^{d-i} = \sum_{i=0}^d h_i x^{d-i} \quad (2)$$

where  $f_i$  is the number of  $i$ -dimensional faces of  $\Delta$  for  $0 \leq i \leq d - 1$  and  $f_{-1} = 1$ . We say that  $\Delta$  is a *triangulation* of a polytopal complex  $\mathcal{F}$  if the union of the faces of  $\Delta$  is equal to the union of the faces of  $\mathcal{F}$  and every face of  $\Delta$  is contained in a face of  $\mathcal{F}$ . In particular we do not require that all vertices of  $\Delta$  are vertices of  $\mathcal{F}$ . A triangulation of the face complex  $\mathcal{F}(P)$  of a convex polytope  $P \subseteq \mathbb{R}^N$  is called a *triangulation* of  $P$ . Such a

triangulation is *regular* if the faces of  $\Delta$  are the projections, under the map  $\mathbb{R}^{N+1} \rightarrow \mathbb{R}^N$  which forgets the last coordinate, of the *lower* faces of a convex polytope  $Q \subseteq \mathbb{R}^{N+1}$ , meaning those faces of  $Q$  which are visible from any point in  $\mathbb{R}^{N+1}$  with sufficiently large negative last coordinate. A convex polytope  $P \subseteq \mathbb{R}^N$  is called an *integer polytope* if all vertices of  $P$  have integer coordinates. A triangulation  $\Delta$  of such a polytope  $P$  is called *unimodular* if all vertices of  $\Delta$  have integer coordinates and the vertex set of any maximal simplex of  $\Delta$  is a basis of the affine integer lattice  $A \cap \mathbb{Z}^N$ , where  $A$  is the affine span of  $P$  in  $\mathbb{R}^N$ . In the special case of pulling triangulations (which we discuss in the sequel) the following lemma appeared as [16, Corollary 2.5].

**Lemma 2.1** ([2]) *If  $P$  is an integer polytope and  $\Delta$  is any unimodular triangulation of  $P$  then  $h^*(P) = h(\Delta)$ .*

A convex polytope  $P$  is said to be *simplicial* if all its proper faces are simplices, so that  $\mathcal{F}(\partial P)$  is a simplicial complex. The next lemma is a consequence of [21, Lemma 2.2] and the note following that lemma in [21]. The inequalities  $h_i \geq h_{d-i}$  were first proved by Kalai [11].

**Lemma 2.2** ([11, 21]) *If  $\Delta$  is a Cohen-Macaulay subcomplex of the boundary complex of a  $d$ -dimensional simplicial polytope and  $h(\Delta) = (h_0, h_1, \dots, h_d)$  then  $h_i \geq h_{d-i}$  for  $0 \leq i \leq \lfloor d/2 \rfloor$  and  $h_{\lfloor d/2 \rfloor} \geq h_{\lfloor d/2 \rfloor + 1} \geq \dots \geq h_d$ .*

Given a polytope  $Q$  and a sequence  $\tau = (v_1, v_2, \dots, v_p)$  of points containing the vertices of  $Q$  we can construct a simplicial polytope  $Q'$  of the same dimension as  $Q$  obtained from  $Q$  by a *sequence of pullings* with respect to  $\tau$ . More specifically let  $\text{pull}_v(P)$  be the convex hull of the set of vertices of  $P$  and the point obtained by moving  $v$  beyond the hyperplanes supporting exactly those facets of  $P$  which contain  $v$  (see [24, Section 3.1]) if  $v$  lies on the boundary of  $P$  and let  $\text{pull}_v(P) = P$  otherwise. We define  $Q' = Q_p$  where  $Q_i = \text{pull}_{v_i}(Q_{i-1})$  for  $1 \leq i \leq p$  and  $Q_0 = Q$ . If  $\tau$  is a linear ordering of the vertices of  $Q$  then  $Q'$  is the polytope obtained from  $Q$  by *pulling the vertices* of  $Q$  in the order  $\tau$  [7, p. 80] [12, Section 2.5]. In this case the vertices of  $Q'$  can be labeled as  $v'_1, v'_2, \dots, v'_p$  so that if  $v_{i_1}, v_{i_2}, \dots, v_{i_j}$  are the vertices of a  $(j-1)$ -dimensional simplex which is a face of  $Q$  then  $v'_{i_1}, v'_{i_2}, \dots, v'_{i_j}$  are the vertices of a  $(j-1)$ -dimensional simplex which is a face of  $Q'$ .

*Proof of Theorem 1.3.* Let  $\Delta$  be a regular unimodular triangulation of  $P$ . Being regular,  $\Delta$  is combinatorially isomorphic to the complex of lower faces of an  $(m+1)$ -dimensional polytope  $Q$  and we may assume that  $Q$  has as many vertices as  $\Delta$ . Pulling the vertices of  $Q$  in an arbitrary order produces an  $(m+1)$ -dimensional simplicial polytope  $Q'$  such that  $\Delta$  is combinatorially isomorphic to a subcomplex  $\Delta'$  of the boundary complex of  $Q'$ . Then  $h(\Delta) = h(\Delta')$  and  $h^*(P) = h(\Delta)$  by Lemma 2.1. Moreover  $\Delta'$  is topologically a ball, being homeomorphic to  $\Delta$ , and hence Cohen-Macaulay. Lemma 2.2 implies that  $h(\Delta') = h^*(P)$  satisfies  $h_{\lfloor (m+1)/2 \rfloor}^* \geq \dots \geq h_m^* \geq h_{m+1}^* = 0$  and  $h_i^* \geq h_{m-i+1}^*$  for  $1 \leq i \leq \lfloor (m+1)/2 \rfloor$ . Let  $(h_0, h_1, \dots, h_{m+1})$  be the  $h$ -vector of the boundary complex of  $Q'$ . The  $h$ -vector of  $\Delta'$  satisfies  $h_i^* \leq h_i$  for  $0 \leq i \leq m$  by [21, Theorem 2.1]. Moreover  $h_1^* = h_1 = n - m - 1$ ,

where  $n$  is the number of vertices of  $\Delta, \Delta', Q$  or  $Q'$  and the last inequality in the theorem follows from the Upper Bound Theorem  $h_i \leq \binom{h_1+i-1}{i}$  for simplicial polytopes [24, Lemma 8.26]. The last statement in the theorem should be clear.  $\square$

We conclude this section with the background on pulling triangulations of polytopal complexes needed for the proof of Theorem 1.1. For any polytopal complex  $\mathcal{F}$  and set of points  $\sigma$  in  $\mathbb{R}^N$  we denote by  $\mathcal{F} \setminus \sigma$  the subcomplex of faces of  $\mathcal{F}$  which do not contain any of the points in  $\sigma$  and write  $\mathcal{F} \setminus v$  for  $\mathcal{F} \setminus \sigma$  if  $\sigma$  consists of a single point  $v$ . Given a sequence  $\tau = (v_1, v_2, \dots, v_p)$  of points containing the vertices of  $\mathcal{F}$  we define the *reverse lexicographic triangulation* or *pulling triangulation*  $\Delta(\mathcal{F}) = \Delta_\tau(\mathcal{F})$  with respect to  $\tau$  [16] [23, Chapter 8] as follows. We have  $\Delta(\mathcal{F}) = \{v\}$  if  $\mathcal{F}$  consists of a single vertex  $v$  and

$$\Delta(\mathcal{F}) = \Delta(\mathcal{F} \setminus v_1) \cup \bigcup_F \{\text{conv}(\{v_1\} \cup G) : G \in \Delta(\mathcal{F}(F))\}$$

otherwise, where the union runs over the facets  $F$  not containing  $v_1$  of the maximal faces of  $\mathcal{F}$  which contain  $v_1$ . The triangulations  $\Delta(\mathcal{F} \setminus v_1)$  and  $\Delta(\mathcal{F}(F))$  are defined with respect to  $(v_2, \dots, v_p)$  by induction. Equivalently, for  $i_0 > i_1 > \dots > i_t$  the set  $\{v_{i_0}, v_{i_1}, \dots, v_{i_t}\}$  is the vertex set of a maximal simplex of  $\Delta_\tau(\mathcal{F})$  if there exists a maximal flag  $F_0 \subset F_1 \subset \dots \subset F_t$  of faces of  $\mathcal{F}$  such that  $v_{i_j}$  is the first element of  $\tau$  in  $F_j$  for all  $j$  and  $v_{i_j}$  is not in  $F_{j-1}$  for  $j \geq 1$ . If  $\mathcal{F}$  is the boundary complex of a polytope  $Q$  then  $\Delta_\tau(\mathcal{F})$  is combinatorially isomorphic to the boundary complex of the simplicial polytope  $Q'$  obtained from  $Q$  by a sequence of pullings with respect to  $\tau$ .

### 3 Semispecial simplices

Throughout this section  $P$  denotes an  $m$ -dimensional convex polytope in  $\mathbb{R}^N$ . We call an  $(n-1)$ -dimensional simplex  $\Sigma$  a *special simplex* for  $P$  if each facet of  $P$  contains exactly  $n-1$  vertices of  $\Sigma$ . This definition is less restrictive than the one given originally in [1] since we do not require that all vertices of  $\Sigma$  are vertices of  $P$ . We call  $\Sigma$  a *semispecial simplex* if each facet of  $P$  contains at least  $n-1$  vertices of  $\Sigma$ . Thus a semispecial simplex for  $P$  is special if and only if it is not contained in the boundary of  $P$ . Two examples are shown in Figure 1.




Figure 1: A special simplex for a triangle and a semispecial simplex for a square pyramid.

**Remark 3.1** Any set  $\sigma$  of  $n$  points in  $\mathbb{R}^N$  having the property that any facet of  $P$  contains at least  $n - 1$  elements of  $\sigma$  must be affinely independent and hence it is the vertex set of a semispecial  $(n - 1)$ -simplex for  $P$ . Indeed, if  $v \in \sigma$  were in the affine span of  $\sigma \setminus v$  then the affine span of any facet of  $P$  would have to contain  $v$ , which is impossible.

If  $V$  is any linear subspace of  $\mathbb{R}^N$  then the *quotient polytope*  $P/V \subseteq \mathbb{R}^N/V$  is the image of  $P$  under the canonical surjection  $\mathbb{R}^N \rightarrow \mathbb{R}^N/V$ . This is a convex polytope in  $\mathbb{R}^N/V$  linearly isomorphic to the image  $\pi(P)$  of  $P$  under any linear surjection  $\pi : \mathbb{R}^N \rightarrow \mathbb{R}^{N-\dim V}$  with kernel  $V$ . Recall that the *simplicial join*  $\Delta_1 * \Delta_2$  of two geometric simplicial complexes  $\Delta_1$  and  $\Delta_2$  in  $\mathbb{R}^N$  is defined if no two line segments, each joining a point in a face of  $\Delta_1$  to a point in a face of  $\Delta_2$ , intersect in their relative interiors. In this case the maximal faces of  $\Delta_1 * \Delta_2$  are the simplices of the form  $\text{conv}(F_1 \cup F_2)$ , where  $F_1$  and  $F_2$  are maximal faces of  $\Delta_1$  and  $\Delta_2$ , respectively. The simplicial join satisfies  $h(\Delta_1 * \Delta_2, x) = h(\Delta_1, x) h(\Delta_2, x)$ , where  $h(\Delta, x) = \sum_{i=0}^d h_i x^{d-i}$  if  $h(\Delta) = (h_0, h_1, \dots, h_d)$ . In particular  $h(\Delta_1 * \Delta_2) = h(\Delta_2)$  if  $\Delta_1$  is a simplex.

**Lemma 3.2** *If  $P$  has a triangulation of the form  $\Sigma * \Delta$  for some  $(n - 1)$ -simplex  $\Sigma$  and simplicial complex  $\Delta$  then  $\Delta$  is combinatorially isomorphic to a triangulation of a pure  $(m - n)$ -dimensional shellable subcomplex of the boundary complex of a polytope of dimension  $m - n + 1$ .*

*Proof.* Let  $V$  be the linear  $(n - 1)$ -dimensional subspace of  $\mathbb{R}^N$  parallel to the affine span of  $\Sigma$  and let  $Q$  be the corresponding  $(m - n + 1)$ -dimensional quotient polytope  $P/V$ . If  $v$  is the point which is the image of  $\Sigma$  under the canonical surjection  $\mathbb{R}^N \rightarrow \mathbb{R}^N/V$  then  $Q$  inherits a triangulation of the form  $v * \Gamma$  for some simplicial complex  $\Gamma$  which is combinatorially isomorphic to  $\Delta$  and triangulates a subcomplex of the boundary complex  $\mathcal{F}(\partial Q)$ . If  $\Sigma$  is not contained in the boundary of  $P$  then  $v$  is an interior point of  $Q$  and  $\Gamma$  triangulates  $\mathcal{F}(\partial Q)$ , which is pure and shellable [24, Section 8.2]. Otherwise  $v$  lies on the boundary of  $Q$  and  $\Gamma$  triangulates the subcomplex of  $\mathcal{F}(\partial Q)$  consisting of all faces of  $Q$  which do not contain  $v$ . This is the complex of faces of  $Q$  which are not visible from a point beyond the hyperplanes supporting exactly those facets of  $Q$  which contain  $v$  and hence is pure  $(m - n)$ -dimensional and shellable (see Lemma 8.10 and Theorem 8.12 in [24]).  $\square$

**Lemma 3.3** *Suppose that  $P = \text{conv}\{v_1, v_2, \dots, v_p\}$ . Let  $\Sigma = \text{conv}\{v_1, v_2, \dots, v_n\}$  and  $\Delta$  be the pulling triangulation of  $\mathcal{F}(P) \setminus \{v_1, v_2, \dots, v_n\}$  with respect to  $(v_{n+1}, \dots, v_p)$ . If  $\Sigma$  is a semispecial  $(n - 1)$ -simplex for  $P$  and  $\tau = (v_1, v_2, \dots, v_p)$  then*

- (i) *the pulling triangulation  $\Delta_\tau$  of  $P$  is combinatorially isomorphic to the simplicial join  $\Sigma * \Delta$  and*
- (ii)  *$\Delta$  is combinatorially isomorphic to a Cohen-Macaulay subcomplex of the boundary complex of a simplicial polytope of dimension  $m - n + 1$ .*

*Proof.* Part (i) can be proved exactly as part (i) of [1, Lemma 3.4], where  $\Sigma$  is assumed to be a special simplex for  $P$ . Lemma 3.2 implies that  $\Delta$  is combinatorially isomorphic to a triangulation  $\Gamma$  of a pure  $(m - n)$ -dimensional shellable subcomplex of the boundary complex of a convex polytope  $Q$  of dimension  $m - n + 1$ . Clearly  $\Gamma$  is homeomorphic to a ball or a sphere and hence Cohen-Macaulay. Let the isomorphism between  $\Delta$  and  $\Gamma$  be induced by the map  $v_i \rightarrow v'_i$  for  $n + 1 \leq i \leq p$  and let  $Q'$  be the simplicial  $(m - n + 1)$ -dimensional polytope obtained from  $Q$  by any sequence of pullings starting with  $(v'_{n+1}, \dots, v'_p)$ . Since  $\Delta$  is a pulling triangulation with respect to  $(v_{n+1}, \dots, v_p)$ ,  $\Gamma$  is combinatorially isomorphic to a subcomplex of the boundary complex of  $Q'$ . This proves part (ii).  $\square$

Let  $h^*(P) = (h_0^*, h_1^*, \dots, h_m^*)$  be the  $h^*$ -vector of  $P$ . Theorem 1.1 is a special case of the following corollary.

**Corollary 3.4** *Suppose that  $P = \text{conv}\{v_1, v_2, \dots, v_p\}$  and  $v_i \in \mathbb{Z}^N$  for  $1 \leq i \leq p$ . Let  $d = m - n + 1$  and  $\tau = (v_1, v_2, \dots, v_p)$ . If the pulling triangulation  $\Delta_\tau$  of  $P$  is unimodular and  $\Sigma = \text{conv}\{v_1, v_2, \dots, v_n\}$  is a semispecial  $(n - 1)$ -simplex for  $P$  then  $h_i^* \geq h_{d-i}^*$  for  $0 \leq i \leq \lfloor d/2 \rfloor$ ,*

$$h_{\lfloor d/2 \rfloor}^* \geq h_{\lfloor d/2 \rfloor + 1}^* \geq \dots \geq h_d^*$$

and  $h_i^* = 0$  for  $d < i \leq m$ . Moreover  $h_d^* = 0$  if  $\Sigma$  is not special.

*Proof.* Let  $\Delta$  denote the pulling triangulation of  $\mathcal{F}(P) \setminus \{v_1, v_2, \dots, v_n\}$  with respect to  $(v_{n+1}, \dots, v_p)$ . Lemma 2.1 guarantees that  $h^*(P) = h(\Delta_\tau)$ . Then part (i) of Lemma 3.3 implies that  $h(\Delta_\tau) = h(\Sigma * \Delta) = h(\Delta)$  and the result follows from part (ii) of the same lemma and Lemma 2.2. If  $\Sigma$  is not special then  $\Delta$  is homeomorphic to a  $(d - 1)$ -dimensional ball and hence  $h_d^* = 0$ .  $\square$

*Proof of Corollary 1.2.* It follows from the case  $n = 1$  of Corollary 3.4 since a singleton  $\{v_1\}$  is always a semispecial 0-dimensional simplex for  $P$ .  $\square$

Observe that if all vertices of the triangulation in the statement of Corollary 1.2 are vertices of  $P$  then  $\{v_1\}$  is not special and hence  $h_m^* = 0$ . It was proved by F. Santos (unpublished) and Ohsugi and Hibi [14] that if  $P$  is a 0/1 polytope, meaning that its vertices are 0/1 vectors, defined by the system of inequalities

$$\begin{aligned} b_i &\leq \sum_{j=1}^N a_{ij}x_j \leq b_i + \varepsilon_i, & 1 \leq i \leq q \\ 0 &\leq x_j \leq 1, & 1 \leq j \leq N \end{aligned} \tag{3}$$

for some integers  $a_{ij}, b_i$  and  $\varepsilon_i$  with  $\varepsilon_i = 0$  or 1 then all pulling triangulations of  $P$  are unimodular. In view of Corollary 1.2 and the remark after its proof we get the following corollary.

**Corollary 3.5** *If  $P$  is an  $m$ -dimensional 0/1 polytope in  $\mathbb{R}^N$  defined by (3) and  $h^*(P) = (h_0^*, h_1^*, \dots, h_m^*)$  then  $h_i^* \geq h_{m-i}^*$  for  $0 \leq i \leq \lfloor m/2 \rfloor$  and  $h_{\lfloor m/2 \rfloor}^* \geq h_{\lfloor m/2 \rfloor + 1}^* \geq \dots \geq h_m^* = 0$ .*

## 4 Order polytopes and Eulerian polynomials

Let  $\Omega$  be a poset (short for partially ordered set) on the ground set  $[N] := \{1, 2, \dots, N\}$  (see [17, Chapter 3] for an introduction to the theory of partially ordered sets). We will denote the partial order of  $\Omega$  by  $\leq_\Omega$ . Recall that an (order) *ideal* of  $\Omega$  is a subset  $I \subseteq \Omega$  for which  $a <_\Omega b$  and  $b \in I$  imply that  $a \in I$  and that  $b$  *covers*  $a$  in  $\Omega$  if  $a <_\Omega b$  but  $a <_\Omega c <_\Omega b$  holds for no  $c \in \Omega$ . Let  $\mathcal{L}(\Omega)$  be the set of *linear extensions* of  $\Omega$ , meaning the set of permutations  $w = (w_1, w_2, \dots, w_N)$  of  $[N]$  for which  $w_i <_\Omega w_j$  implies  $i < j$ . The  $\Omega$ -*Eulerian polynomial* is defined as

$$W(\Omega, t) = \sum_{w \in \mathcal{L}(\Omega)} t^{\text{des}(w)}$$

where

$$\text{des}(w) = \#\{i \in [N-1] : w_i > w_{i+1}\}$$

is the number of descents of  $w$ . Let  $\Omega^0$  be the poset obtained from  $\Omega$  by adjoining a minimum element  $\hat{0} = 0$ . We define the *ideal height* of  $\Omega$  to be the largest length  $e$  of a chain  $I_0 \subset I_1 \subset \dots \subset I_e$  of nonempty ideals of  $\Omega^0$  such that for  $1 \leq i \leq e$  and for any  $a \in I_{i-1}$  the set of elements covering  $a$  in  $\Omega$  is a nonempty subset of  $I_i$ . Figure 2 shows the Hasse diagram of a poset  $\Omega$  of ideal height 3. Observe that the cardinality of the shortest maximal chain in  $\Omega$  in this example is equal to 4. The poset  $\Omega$  is *naturally labeled* if the identity permutation  $(1, 2, \dots, N)$  is a linear extension. The following theorem is the main result of this section.




Figure 2: A poset of ideal height 3 with 8 elements.

**Theorem 4.1** *Let  $\Omega$  be a naturally labeled poset on  $[N]$ . If  $e$  is the ideal height of  $\Omega$ ,  $W(\Omega, t) = q_0 + q_1 t + \dots + q_N t^N$  is the  $\Omega$ -Eulerian polynomial and  $d = N - e$  then  $q_i \geq q_{d-i}$  for  $1 \leq i \leq \lfloor d/2 \rfloor$ ,  $q_{\lfloor d/2 \rfloor} \geq q_{\lfloor d/2 \rfloor + 1} \geq \dots \geq q_d$  and  $q_i = 0$  for  $d < i \leq N$ .*

To prove this statement we will apply Theorem 1.1 to the order polytope of  $\Omega$ . Let  $\hat{\Omega}$  be the poset obtained from  $\Omega^0$  by adjoining a maximum element  $\hat{1} = N + 1$ . The



order polytope [18] of  $\Omega$ , denoted  $O(\Omega)$ , is the intersection of the hyperplanes  $x_0 = 1$  and  $x_{N+1} = 0$  in  $\mathbb{R}^{N+2}$  with the cone defined by the inequalities  $x_i \geq x_j$  for  $i, j \in \hat{\Omega}$  with  $i <_{\hat{\Omega}} j$ . The vertices of  $O(\Omega)$  are the characteristic vectors of the nonempty ideals of  $\Omega^0$  [18, Corollary 1.3] so, in particular,  $O(\Omega)$  is an  $N$ -dimensional integer polytope. Moreover the facets of  $O(\Omega)$  are defined exactly by the equalities of the form  $x_i = x_j$  when  $j$  covers  $i$  in  $\hat{\Omega}$  (see [18, Theorem 1.2] for a complete description of the facial structure of  $O(\Omega)$ ) and the coefficients of the  $\Omega$ -Eulerian polynomial

$$W(\Omega, t) = q_0 + q_1 t + \cdots + q_N t^N$$

are the entries of the  $h^*$ -vector  $(h_0^*, h_1^*, \dots, h_N^*)$  of  $O(\Omega)$  [17, Section 4.5], that is  $q_i = h_i^*$  for all  $i$ .

*Proof of Theorem 4.1.* Let  $P$  be the order polytope of  $\Omega$ . It was shown by Ohsugi and Hibi [14, Example 1.3 (b)] that all pulling triangulations (with integer vertices) of order polytopes are unimodular. Moreover it follows easily from the description of the facets of  $P$  that if  $I_0 \subset I_1 \subset \cdots \subset I_e$  is a chain of nonempty ideals of  $\Omega^0$  as in the definition of the ideal height for  $\Omega$  then the characteristic vectors of the  $I_i$  are the vertices of a semispecial  $e$ -dimensional simplex for  $P$ . The result follows from Theorem 1.1.  $\square$

We close this section with a different characterization of ideal height. For  $a \in \Omega^0$  let  $\mathcal{C}_a$  denote the set of sequences  $(a_0, a_1, \dots, a_l)$  in  $\hat{\Omega}$  such that  $a_0 = \hat{0}$ ,  $a_l = a$  and for  $1 \leq i \leq l$  either  $a_i$  covers  $a_{i-1}$  in  $\hat{\Omega}$  or  $a_{i-1}$  covers  $a_i$ . For each  $\alpha = (a_0, a_1, \dots, a_l) \in \mathcal{C}_a$  let  $e(\alpha)$  denote the number of indices  $1 \leq i \leq l$  for which  $a_i$  covers  $a_{i-1}$  and  $a_i <_{\hat{\Omega}} \hat{1}$  and let  $e(a)$  denote the minimum value of  $e(\alpha)$  when  $\alpha$  ranges over all sequences in  $\mathcal{C}_a$ .

**Proposition 4.2** *The ideal height of  $\Omega$  is equal to the maximum value of  $e(a)$  for  $a \in \Omega$ .*

*Proof.* Let  $f$  denote the maximum value of  $e(a)$  for  $a \in \Omega$  and  $e$  denote the ideal height of  $\Omega$ . Let  $a$  be any maximal element of  $\Omega$ , let  $\alpha = (a_0, a_1, \dots, a_l) \in \mathcal{C}_a$  and let  $I_0 \subset I_1 \subset \cdots \subset I_e$  be a chain of nonempty ideals of  $\Omega^0$  as in the definition of ideal height. Observe that (i)  $a \notin I_{e-1}$ , (ii) if  $a_j \in \Omega$  and  $a_j \notin I_i$  then  $a_{j-1} \notin I_{i-1}$  and  $a_{j-1} \in I_i$  is possible only when  $a_j$  covers  $a_{j-1}$  and (iii) if  $a_j = \hat{1}$  then  $a_{j-1} \notin I_{e-1}$ . Since  $a_0 \in I_0$  there must be at least  $e$  indices  $j$  for which  $a_j$  covers  $a_{j-1}$  in  $\Omega^0$ . This shows that  $e(a) \geq e$  and hence  $f \geq e$ .

For the other inequality let  $I_i$  denote the set of elements  $a \in \Omega^0$  with  $e(a) \leq i$  for  $0 \leq i \leq f$ . Observe that  $e(\hat{0}) = 0$  and that if  $b$  covers  $a$  in  $\Omega^0$  then  $e(a) \leq e(b) \leq e(a) + 1$ . It follows that  $I_0 \subset I_1 \subset \cdots \subset I_f$  is a chain of nonempty ideals of  $\Omega^0$ . We will show that this chain satisfies the condition in the definition of ideal height, whence  $f \leq e$ . Let  $1 \leq i \leq f$  and  $a \in I_{i-1}$ . Any element  $b$  in  $\Omega$  covering  $a$  satisfies  $e(b) \leq e(a) + 1 \leq i$  and hence  $b \in I_i$ . On the other hand if  $a$  were a maximal element of  $\Omega$  and  $b$  is any element of  $\Omega$ , which we may assume to be maximal, with  $e(b) = f$  then the sequence of coverings  $(b, \hat{1}, a)$  in  $\hat{\Omega}$  shows that  $e(b) = e(a)$ , contradicting the hypothesis that  $e(a) \leq i - 1 < f$ . Hence  $a$  is covered by at least one element of  $\Omega$ , which is necessarily in  $I_i$ . This completes the proof.  $\square$

## 5 Stable polytopes of perfect graphs

We consider simple (with no loops or multiple edges) graphs  $G$  on the finite set of nodes  $[N] := \{1, 2, \dots, N\}$ . For  $W \subseteq [N]$  we denote by  $\rho(W)$  the 0/1 vector  $\sum_{i \in W} e_i \in \mathbb{R}^N$ , where  $e_i$  is the  $i$ th unit coordinate vector in  $\mathbb{R}^N$ , and call  $W$  a *stable set* for  $G$  if no two elements of  $W$  are joined by an edge in  $G$ . The *stable polytope* of  $G$ , introduced in [4] and denoted by  $P(G)$ , is the convex hull of all vectors  $\rho(W)$ , where  $W$  is a stable set for  $G$ . Observe that the empty set and all singleton subsets of  $[N]$  are stable and hence  $P(G)$  has dimension  $N$ . The *chromatic number* of  $G$  is the least number  $r$  of colors that can be assigned to the vertices of  $G$ , one color to each vertex, so that no two adjacent vertices of  $G$  are assigned the same color. The graph  $G$  is called *complete* if any two of its vertices are connected by an edge and *perfect* if for any induced subgraph  $H$  of  $G$  the chromatic number of  $H$  is equal to the number of vertices of the largest complete subgraph of  $H$ . We call  $G$  *semipure* if there exists a positive integer  $j$  such that all maximal complete subgraphs of  $G$  have either  $j - 1$  or  $j$  vertices. Thus if  $G$  is perfect and semipure with chromatic number  $r$  then all maximal complete subgraphs of  $G$  have either  $r - 1$  or  $r$  vertices. The following theorem applies in particular to all bipartite graphs and all perfect graphs of chromatic number three with no isolated vertices.

**Theorem 5.1** *If  $G$  is a semipure perfect graph with  $N$  vertices and chromatic number  $r$  and  $d = N - r + 1$  then the  $h^*$ -vector  $(h_0^*, h_1^*, \dots, h_N^*)$  of the stable polytope  $P(G)$  satisfies  $h_i^* \geq h_{d-i}^*$  for  $1 \leq i \leq \lfloor d/2 \rfloor$ ,  $h_{\lfloor d/2 \rfloor}^* \geq h_{\lfloor d/2 \rfloor + 1}^* \geq \dots \geq h_{d-1}^*$  and  $h_i^* = 0$  for  $d \leq i \leq N$ .*

*Proof.* It was proved in [14] that all pulling triangulations of stable polytopes of perfect graphs are unimodular. In view of Theorem 1.1 and the last statement in Corollary 3.4 it suffices to prove that there exists a semispecial  $(r - 1)$ -simplex for  $P(G)$  with integer vertices which is not special. Consider a coloring of  $G$  with colors  $1, 2, \dots, r$  and for  $1 \leq i \leq r$  let  $W_i$  be the set of vertices of  $G$  colored with  $i$ , so that  $W_i$  is a stable set for  $G$ . We claim that  $\Sigma = \text{conv}\{\rho(W_i) : 1 \leq i \leq r\}$  is such a simplex for  $P(G)$ . Since  $G$  is perfect, by [4, Theorem 3.1] a facet of  $P(G)$  is defined by a linear equality of the form  $x_i = 0$  or  $\sum_{j \in U} x_j = 1$  for some vertex set  $U$  of a maximal complete subgraph of  $G$ . A facet of the first form contains exactly  $r - 1$  of the points  $\rho(W_i)$  since the sets  $W_i$  form a partition of  $[N]$ . Since  $G$  is also semipure, a facet of the second form contains either  $r - 1$  or  $r$  points  $\rho(W_i)$ , where the second case occurs, and the claim follows from Remark 3.1.  $\square$

## 6 Affine semigroup rings and ideals

Let  $P$  be an  $m$ -dimensional integer polytope in  $\mathbb{R}^N$  and  $\mathbb{K}$  be a field. We denote by  $R_P$  the subalgebra of the algebra  $\mathbb{K}[x_1, \dots, x_N, x_1^{-1}, \dots, x_N^{-1}, t]$  of Laurent polynomials over  $\mathbb{K}$  generated by the monomials  $x^\alpha t^r$  for positive integers  $r$  and  $\alpha \in \mathbb{Z}^N$  such that  $\alpha/r \in P$ . The algebra  $R_P$  can be graded by letting  $x^\alpha t^r$  have degree  $r$ . With this grading  $R_P$  is a semistandard graded Cohen-Macaulay normal domain, called the *semigroup ring* of  $P$ , whose Hilbert series is the Ehrhart series (1) of  $P$ . See [3, Chapter 6] for background on

semigroup rings. Let  $\tilde{P} = \{(1, x) : x \in P\}$  be the lift of  $P$  in the hyperplane  $x_0 = 1$  in  $\mathbb{R}^{N+1}$  and denote by  $\mathcal{C}_P$  the cone in  $\mathbb{R}^{N+1}$  generated by  $\tilde{P}$  and by  $E_P$  the semigroup of integer points in  $\mathcal{C}_P$ . Let  $\mathcal{L}$  be the linear span of  $\mathcal{C}_P$  in  $\mathbb{R}^{N+1}$  and  $\mathcal{A}$  be a set of integer linear forms in  $\mathbb{R}^{N+1}$ , one for each facet of  $P$ , defining the cone  $\mathcal{C}_P$  as the set of points  $x \in \mathcal{L}$  satisfying  $g(x) \geq 0$  for all  $g \in \mathcal{A}$ . For  $\mathcal{G} \subseteq \mathcal{A}$  let  $E_{\mathcal{G}}$  be the semigroup of elements  $\alpha$  of  $E_P$  satisfying  $g(\alpha) > 0$  for  $g \in \mathcal{G}$  and  $I_{\mathcal{G}}$  be the ideal of  $R_P$  generated by the monomials  $x^\alpha$  with  $\alpha \in E_{\mathcal{G}}$  (observe that the variable  $t$  has been replaced by  $x_0$ ). We say that  $E_{\mathcal{G}}$  has a *unique minimal element*  $\beta$  if  $\beta + E_P = E_{\mathcal{G}}$  or, equivalently, if  $I_{\mathcal{G}}$  is generated by the monomial  $x^\beta$  as an ideal of  $R_P$ . The proof of the following lemma is similar to that of [1, Corollary 4.1].

**Lemma 6.1** *If  $\mathcal{G} \subseteq \mathcal{A}$  and  $v_1 + v_2 + \cdots + v_n$  is the unique minimal element of  $E_{\mathcal{G}}$  for some integer points  $v_1, v_2, \dots, v_n$  in  $\tilde{P}$  then  $\{v_1, v_2, \dots, v_n\}$  is the vertex set of a semispecial  $(n - 1)$ -simplex for  $\tilde{P}$ .*

*Proof.* Let  $\beta = v_1 + v_2 + \cdots + v_n$  and observe that  $v_1, v_2, \dots, v_n$  are necessarily distinct. Let  $F$  be a facet of  $\tilde{P}$  with corresponding linear form  $f \in \mathcal{A}$ . If  $f$  is not in  $\mathcal{G}$  then there exists a  $v \in E_{\mathcal{G}}$  with  $f(v) = 0$  and, by the minimality of  $\beta$ ,  $f(\beta) = 0$ . Hence  $f(v_i) = 0$ , meaning that  $v_i \in F$ , for all  $1 \leq i \leq n$ . Suppose that  $f \in \mathcal{G}$ , so that  $f(\beta) > 0$ . We need to show that at most one of  $v_1, v_2, \dots, v_n$  satisfies  $f(v_i) > 0$ . Clearly at least one of them has this property. Assume that at least two of  $v_1, v_2, \dots, v_n$  satisfy  $f(v_i) > 0$ , say  $v_j$  is one of them, and let  $f(\beta) = b$  and  $f(v_j) = c$ , so that  $1 \leq c < b$ . Since  $F$  is a facet of  $\tilde{P}$  there exists a point  $x$  in the affine span of  $\tilde{P}$ , which we may assume to have rational coordinates, satisfying  $f(x) < 0$  and  $g(x) > 0$  for all  $g \in \mathcal{A}$  other than  $f$ . By replacing  $x$  with a suitable positive integer multiple we find an integer point  $\alpha$  in  $\mathcal{L}$  satisfying  $f(\alpha) < 0$  and  $g(\alpha) > 0$  for all  $g \in \mathcal{A}$  other than  $f$ . Letting  $a = f(\alpha)$ , we may choose a nonnegative integer  $t$  so that  $0 < a + b + tc < b$ . Then  $\gamma = \alpha + \beta + tv_j$  is in  $E_{\mathcal{G}}$  and satisfies  $f(\gamma) < f(\beta)$ , which contradicts the minimality of  $\beta$ .  $\square$

Recall that  $R_P$  is *standard* if it is generated by its homogeneous elements of degree one or, equivalently, if  $E_P$  is generated as a semigroup by the integer points in  $\tilde{P}$ . Clearly this holds if  $P$  has a unimodular triangulation. In the case  $\mathcal{G} = \mathcal{A}$  assumption (ii) in the following corollary is equivalent to the statement that the ring  $R_P$  is Gorenstein [3, Corollary 6.3.8].

**Corollary 6.2** *If*

- (i) *all pulling triangulations of  $P$  with integer vertices are unimodular and*
- (ii)  *$I_{\mathcal{G}}$  is generated by one element as an ideal of  $R_P$*

*then the conclusion of Corollary 3.4 holds where, in the statement of the corollary,  $n$  is the  $x_0$ -coordinate of the unique minimal element of  $E_{\mathcal{G}}$  and the equality  $h_d^* = 0$  holds if  $\mathcal{G} \subsetneq \mathcal{A}$ .*

*Proof.* Let  $\beta$  be the unique minimal element of  $E_G$ , whose existence is guaranteed by (ii). Assumption (i) implies that  $R_P$  is standard and hence  $\beta = v_1 + v_2 + \cdots + v_n$  for some integer points  $v_1, v_2, \dots, v_n$  in  $\tilde{P}$ , which are the vertices of a semispecial  $(n-1)$ -simplex for  $\tilde{P}$  by Lemma 6.1. Because of (i) any sequence  $(v_1, v_2, \dots, v_p)$  of integer points containing the vertices of  $\tilde{P}$  satisfies the assumptions of Corollary 3.4. The result follows from this corollary observing that  $\beta$  has  $x_0$ -coordinate equal to  $n$ .  $\square$

## References

- [1] C.A. Athanasiadis, *Ehrhart polynomials, simplicial polytopes, magic squares and a conjecture of Stanley*, J. Reine Angew. Math. (to appear).
- [2] U. Betke and P. McMullen, *Lattice points in lattice polytopes*, Monatsh. Math. **99** (1985), 253–265.
- [3] W. Bruns and J. Herzog, *Cohen-Macaulay rings*, Cambridge Studies in Advanced Mathematics **39**, Cambridge University Press, Cambridge, 1998.
- [4] V. Chvátal, *On certain polytopes associated with graphs*, J. Combin. Theory Series B **18** (1975), 138–154.
- [5] E. Ehrhart, *Sur un problème de géométrie diophantienne linéaire. I. Polyèdres et réseaux*, J. Reine Angew. Math. **226** (1967), 1–29.
- [6] E. Ehrhart, *Sur un problème de géométrie diophantienne linéaire. II. Systèmes diophantiens linéaires*, J. Reine Angew. Math. **227** (1967), 25–49.
- [7] B. Grünbaum, *Convex Polytopes*, Wiley-Interscience, New York, 1967; second edition, Graduate Texts in Mathematics **221**, Springer-Verlag, New York, 2002.
- [8] T. Hibi, *Flawless  $O$ -sequences and Hilbert functions of Cohen-Macaulay integral domains*, J. Pure Applied Algebra **60** (1989), 245–251.
- [9] T. Hibi, *Algebraic Combinatorics on Convex Polytopes*, Carslaw Publications, Glebe, Australia, 1992.
- [10] T. Hibi, personal communication, 2004.
- [11] G. Kalai, *The diameter of graphs of convex polytopes and  $f$ -vector theory*, in *Applied Geometry and Discrete Mathematics*, DIMACS Series in Discrete Mathematics and Theoretical Computer Science **4**, Amer. Math. Soc., Providence, RI, 1991, pp. 387–411.
- [12] P. McMullen and G.C. Shephard, *Convex Polytopes and the Upper Bound Conjecture*, London Math. Soc. Lecture Note Series **3**, Cambridge University Press, London/New York, 1971.

- [13] H. Ohsugi and T. Hibi, *A normal  $(0, 1)$ -polytope none of whose regular triangulations are unimodular*, Discrete Comput. Geom. **21** (1999), 201–204.
- [14] H. Ohsugi and T. Hibi, *Convex polytopes all of whose reverse lexicographic initial ideals are squarefree*, Proc. Amer. Math. Soc. **129** (2001), 2541–2546.
- [15] V. Reiner and V. Welker, *On the Charney-Davis and Neggers-Stanley conjectures*, J. Combin. Theory Series A (to appear).
- [16] R.P. Stanley, *Decompositions of rational convex polytopes*, Annals of Discrete Math. **6** (1980), 333–342.
- [17] R.P. Stanley, Enumerative Combinatorics, vol. 1, Wadsworth & Brooks/Cole, Pacific Grove, CA, 1986; second printing, Cambridge University Press, Cambridge, 1997.
- [18] R.P. Stanley, *Two poset polytopes*, Discrete Comput. Geom. **1** (1986), 9–23.
- [19] R.P. Stanley, *Log-concave and unimodal sequences in algebra, combinatorics, and geometry*, in *Graph Theory and its Applications: East and West*, Annals of the New York Academy of Sciences **576**, New York Acad. Sci., New York, 1989, pp. 500–535.
- [20] R.P. Stanley, *On the Hilbert function of a graded Cohen-Macaulay domain*, J. Pure Applied Algebra **73** (1991), 307–314.
- [21] R.P. Stanley, *A monotonicity property of  $h$ -vectors and  $h^*$ -vectors*, European J. Combin. **14** (1993), 251–258.
- [22] R.P. Stanley, Combinatorics and Commutative Algebra, Progress in Mathematics **41**, Birkhäuser, Boston, second edition, 1996.
- [23] B. Sturmfels, Gröbner Bases and Convex Polytopes, University Lecture Series **8**, American Mathematical Society, Providence, RI, 1996.
- [24] G.M. Ziegler, Lectures on Polytopes, Graduate Texts in Mathematics **152**, Springer-Verlag, New York, 1995.