

Published in final edited form as:

Science. 2008 October 24; 322(5901): 587–590. doi:10.1126/science.1162667.

H₂S as a Physiologic Vasorelaxant: Hypertension in Mice with Deletion of Cystathionine γ -Lyase

Guangdong Yang^{1,5}, Lingyun Wu^{2,*}, Bo Jiang¹, Wei Yang¹, Jiansong Qi¹, Kun Cao¹, Qinghe Meng³, Asif K. Mustafa⁴, Weitong Mu^{4,6}, Shengming Zhang⁵, Solomon H. Snyder^{4,*}, and Rui Wang^{1,5,*}

¹Department of Physiology, University of Saskatchewan, Saskatoon, SK S7N 5E5, Canada.

²Department of Pharmacology, University of Saskatchewan, Saskatoon, SK S7N 5E5, Canada.

³Department of Pathology, University of Saskatchewan, Saskatoon, SK S7N 5E5, Canada.

⁴Departments of Neuroscience, Pharmacology and Molecular Sciences and Psychiatry, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA.

⁵Department of Biology, Lakehead University, Thunder Bay, ON P7B 5E1, Canada.

⁶Department of Medicine, Gastroenterology Division, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA.

Abstract

Studies of nitric oxide over the past two decades have highlighted the fundamental importance of gaseous signaling molecules in biology and medicine. The physiological role of other gases such as carbon monoxide and hydrogen sulfide (H₂S) is now receiving increasing attention. Here we show that H₂S is physiologically generated by cystathionine γ -lyase (CSE) and that genetic deletion of this enzyme in mice markedly reduces H₂S levels in the serum, heart, aorta, and other tissues. Mutant mice lacking CSE display pronounced hypertension and diminished endothelium-dependent vasorelaxation. CSE is physiologically activated by calcium-calmodulin, which is a mechanism for H₂S formation in response to vascular activation. These findings provide direct evidence that H₂S is a physiologic vasodilator and regulator of blood pressure.

Nitric oxide (NO) and carbon monoxide (CO) are established physiologic messenger molecules, and NO has an important role as an endothelial cell-derived relaxing factor (EDRF) and regulator of blood pressure (1,2). Indirect evidence has implicated another endogenous gasotransmitter, hydrogen sulfide (H₂S), in similar functions (3–7). H₂S can be produced by cystathionine γ -lyase (CSE) or cystathionine β -synthase (CBS) (3,4), but definitive evidence for either of these enzymes in the physiologic formation of H₂S is lacking.

To investigate the role of H₂S as a physiologic vasorelaxant and determinant of blood pressure, we generated mice with a targeted deletion of the gene encoding CSE (8) (fig. S1, A to C). The homozygous (CSE^{-/-}) and heterozygous (CSE^{+/-}) mutant mice were viable, fertile, and indistinguishable from their control wild-type littermates (CSE^{+/+}) in terms of growth pattern.

*To whom correspondence should be addressed. E-mail: rwang@lakeheadu.ca (R.W.), lily.wu@usask.ca (L.W.), or ssnyder@jhmi.edu (S.H.S.).

Supporting Online Material

www.sciencemag.org/cgi/content/full/322/5901/587/DC1

Materials and Methods

Figs. S1 to S6

References

CSE mRNA and protein were absent in heart, aorta, mesenteric artery, liver, and kidneys of CSE^{-/-} mice (fig. S1, D and E). Endogenous H₂S levels in aorta and heart of homozygous mutant male mice (CSE^{-/-}) were both decreased by about 80% (Fig. 1A), and H₂S levels in aorta and heart of heterozygous mutant male mice (CSE^{+/-}) were both decreased by about 50%. Serum H₂S levels in CSE^{-/-} mice and CSE^{+/-} mice were reduced by about 50 and 20%, respectively (Fig. 1B). Female CSE^{-/-} mice showed a similar decline in H₂S levels (fig. S2, A and B). The residual H₂S in serum may reflect nonenzymatic reduction of elemental sulfur to H₂S or H₂S generated from other tissues that express CBS, another H₂S-generating enzyme (3,5,9).

CSE mutant mice developed age-dependent hypertension. Beginning at 7 weeks of age, both male (Fig. 1C) and female (fig. S2C) CSE^{-/-} mice displayed a higher blood pressure than age-matched wild-type (WT) mice. Blood pressure in the mutant mice peaked at more than 135 mm Hg when the mice were 12 weeks of age; this was almost 18 mm Hg higher than that in control mice. Heterozygous CSE^{+/-} mice also showed elevated blood pressure beginning at 7 weeks of age. The rise in blood pressure was similar in homozygous and heterozygous mice until the mice were 10 weeks of age; after this point, the blood pressure of CSE^{-/-} mice was about 10 mm Hg higher than that of CSE^{+/-} mice. Blood pressure levels assessed by the tail-cuff method were confirmed by direct monitoring of arterial blood pressure through intra-carotid artery catheterization (fig. S3A). Heart rates were similar in mutant and WT mice. In humans, CSE activity increases rapidly after birth, reaching adult levels when infants are about 3 months of age (10, 11). The age-dependent hypertension of the mutant mice paralleled the ontogeny of CSE in mice, increasing to peak adult levels 3 weeks after birth (12). Endogenous H₂S levels in brains from CSE^{-/-} mice were similar to WT mouse values (fig. S3B), consistent with evidence that CSE is not the source of brain H₂S (3, 5, 11, 12), and this similarity suggests that the hypertension in the mutant mice is not due to alterations in the central nervous system. In addition, endothelial NO synthase (eNOS) protein was not decreased in CSE^{-/-} mice, which indicated that the hypertension was not due to a loss in NO-mediated vasorelaxation. Kidney architecture was also preserved in the CSE^{-/-} mice, which signifies that the elevation in blood pressure was not caused by renal damage (fig. S4).

H₂S relaxes blood vessels and lowers blood pressure by opening ATP-sensitive K⁺ channels in vascular smooth muscle (4,13,14). We explored whether exogenous H₂S could influence the hypertension of CSE^{-/-} animals. Intravenous bolus injections of NaHS, an H₂S donor (4, 13,15), elicited dose-dependent transient decreases in systolic blood pressure of both anesthetized CSE^{-/-} and CSE^{+/-} mice (Fig. 1D). The magnitude of decline was greater in mutant versus WT mice, which suggested that the former have a heightened sensitivity to H₂S. NaHS injections did not alter the heart rate of WT or mutant mice. Intravenous bolus injections of ammonium (39 μmol/kg) or pyruvate (39 μmol/kg), the two other products of CSE activity (3,16), did not influence blood pressure or heart rate. Plasma levels of oxobutanoate, an intermediate in the catabolism of cystathionine by CSE, were similar in CSE^{-/-} and CSE^{+/-} mice (fig. S3C).

CSE deficiency may elicit accumulation of homocysteine and diminished levels of L-cysteine (17). In 10-week-old male CSE^{-/-} mice, plasma homocysteine and L-cysteine levels were, respectively, about 18 and 0.8 times the levels seen in age-matched WT mice, whereas the levels in CSE^{+/-} mice were, respectively, about 2.0 and 0.8 times those in WT mice (Fig. 1, E and F). A similar decrease in L-cysteine levels was observed in female CSE^{-/-} mice (fig. S2E). To ascertain whether the hypertension of CSE^{-/-} mice reflects hyperhomocysteinemia (18), we administered L-methionine to WT mice in their drinking water for 6 weeks. This intervention augmented plasma homocysteine levels (fig. S3D), but did not alter blood pressure (fig. S3E). Also, while plasma homocysteine levels in male CSE^{-/-} mice were nine times higher than those in male CSE^{+/-} mice (Fig. 1E), blood pressure in the two genotypes was similar

(Fig. 1C). Moreover, female and male CSE^{-/-} displayed similar blood pressures (fig. S5A), despite females having six times the plasma homocysteine levels and homocysteine/cysteine ratios seen in males (Fig. 1E and figs. S2D and S5B). Thus, homocysteine is unlikely to be the principal determinant of hypertension in the CSE mutant mice.

We next investigated whether hypertension in the CSE^{-/-} mice reflected alterations in the vascular redox state. Analysis of vascular tissue indicated that the levels of superoxide anion, a reactive oxygen species (ROS) that regulates vascular tone, were not significantly different in CSE^{-/-} versus WT mice (fig. S6A). Glutathione (GSH) levels were moderately decreased in the aorta and mesenteric artery beds of the mutant mice (fig. S6B), possibly as a result of the modest decreases seen in L-cysteine levels. As substantially greater decreases of GSH are not associated with hypertension, it is unlikely that GSH plays a major role in the hypertension of CSE mutant mice (19).

To investigate mechanisms underlying CSE^{-/-} hypertension, we examined blood vessel responses of the mutant mice. Phenylephrine contracts blood vessels by activating α -adrenoceptors in vascular smooth muscle, whereas H₂S and NO directly relax the muscle (4, 20). By contrast, relaxation after cholinergic stimulation reflects influences on endothelium (21). Phenylephrine evoked contraction of mesenteric arteries to a similar extent in WT and CSE^{-/-} mice (Fig. 2A), and the NO donor sodium nitroprusside produced a similar vasorelaxation response in mutant and WT mesenteric arteries (Fig. 2B). H₂S more potently relaxed mesenteric arteries of CSE^{-/-} mice, with a median inhibitory concentration (IC₅₀) of 75 μ M, as compared with WT mice (IC₅₀ = 120 μ M), consistent with supersensitivity associated with diminished formation of endogenous H₂S (Fig. 2C). By contrast, methacholine-induced relaxation of mesenteric arteries that had been constricted by phenylephrine was markedly impaired in mutant mice (Fig. 2D), and endothelial removal abolished methacholine relaxation of both WT and mutant arteries (Fig. 2E). Immunohistochemistry experiments revealed that CSE protein predominantly localized to the endothelium, with faint staining in smooth muscle (Fig. 2F). In an earlier study, we had shown that CSE mRNA is expressed in vascular smooth muscle (4). Our reexamination of these data revealed that CSE mRNA is also expressed in the endothelium. Thus, H₂S displays properties characteristic of an EDRF. It is formed in endothelium, and prevention of its synthesis impairs relaxation elicited by a neurotransmitter that acts via the endothelium, but does not impair effects of agents that act directly on smooth muscle. The extent to which H₂S, NO, or CO contribute to EDRF activity in different vascular beds is unclear.

How does endothelial stimulation generate H₂S? eNOS and heme oxygenase-2 (HO-2), the biosynthetic enzymes for NO and CO, respectively, are activated by calcium-calmodulin (1, 22). Thus, endothelial activation by substances such as acetylcholine or bradykinin elicits formation of inositol 1,4,5-trisphosphate, which releases intracellular calcium to stimulate formation of NO or CO. We found a similar mode of regulation for CSE. Using recombinant CSE, we demonstrated its direct binding to calmodulin, which was abolished by the calcium chelator EGTA and the calmodulin antagonist W7 (Fig. 3A). Catalytic activity of pure CSE was increased more than twofold by calcium and calmodulin, but not by either substance alone (Fig. 3B), a level of stimulation similar to the NO- and CO-generating enzymes eNOS (23) and HO-2 (22). Calcium-dependent regulation of CSE was also evident in endothelial cells, which contain abundant levels of CSE (Fig. 3C). H₂S formation by these cells was markedly augmented by the calcium ionophore A23187, with the increase blocked by the calcium chelator BAPTA [1,2-bis(2-aminophenoxy)ethane- *N,N,N',N'*-tetraacetic acid] and by W7 (Fig. 3D). These agents markedly reduced basal levels of H₂S, which indicated that H₂S generation by CSE is physiologically regulated by calcium-calmodulin.

Endothelial-dependent vasorelaxation reflects muscarinic cholinergic activation of eNOS (24). We demonstrated similar regulation of H₂S formation. Thus methacholine treatment of endothelial cells tripled H₂S levels (Fig. 3E), an effect blocked by the anticholinergic drug atropine. Depletion of CSE by RNA interference markedly diminished the enhancement of H₂S formation by methacholine and A23187 and lowered basal levels of H₂S (Fig. 3F).


In summary, we have established CSE as the physiologic source of H₂S in multiple tissues, especially the vascular system. Mice genetically deficient in this enzyme display marked hypertension, comparable to that of eNOS^{-/-} mice (24–26). Our findings are consistent with the previous observation that administration of the CSE inhibitor D,L-propargylglycine elevates blood pressure (27). H₂S has properties in common with physiologic EDRFs. Thus, blood vessel relaxation in response to muscarinic stimulation is profoundly reduced in CSE-deficient mice. Moreover, CSE is predominantly localized to the endothelial layer of blood vessels. The EDRF activity of H₂S reflects muscarinic activation of intracellular calcium release, with calcium-calmodulin physiologically stimulating CSE. NO-mediated EDRF activity arises through a similar mechanism.

Although NO is well established as an EDRF, in numerous vascular beds EDRF activity is only partially diminished by NO synthase inhibitors and in mice lacking the gene for eNOS (28,29). In our experiments, EDRF activity of murine mesenteric arteries from mutant mice lacking CSE was reduced by about 60%, which suggests that H₂S functions as an EDRF in this vascular bed. The similar elevation of blood pressure in mice with CSE and eNOS knockouts implies that H₂S influences vascular systems underlying peripheral resistance to an extent comparable to the action of NO. A physiologic role for H₂S in regulating blood pressure raises the possibility that pharmacologic enhancement of H₂S formation could be an alternative approach for treatment of hypertension.

References and Notes

- Boehning D, Snyder SH. *Annu. Rev. Neurosci* 2003;26:105. [PubMed: 14527267]
- Ndisang JF, Tabien HEN, Wang R. *J. Hypertens* 2004;22:1057. [PubMed: 15167436]
- Wang R. *FASEB J* 2002;16:1792. [PubMed: 12409322]
- Zhao W, Zhang J, Lu Y, Wang R. *EMBO J* 2001;20:6008. [PubMed: 11689441]
- Kimura H. *Mol. Neurobiol* 2002;26:13. [PubMed: 12392053]
- Mok YY, et al. *Br. J. Pharmacol* 2004;143:881. [PubMed: 15504752]
- Fiorucci S, et al. *Gastroenterology* 2005;129:1210. [PubMed: 16230075]
- Materials and methods are available as supporting material on *Science* online.
- Searcy DG, Lee SH. *J. Exp. Zool* 1998;282:310. [PubMed: 9755482]
- Zlotkin SH, Anderson GH. *Pediatr. Res* 1982;16:65. [PubMed: 7070878]
- Heinonen K. *Biochem. J* 1973;136:1011. [PubMed: 4786524]
- Ishii I, et al. *Biochem. J* 2004;381:113. [PubMed: 15038791]
- Zhao W, Wang R. *Am. J. Physiol. Heart Circ. Physiol* 2002;283:H474. [PubMed: 12124191]
- Cheng Y, Ndisang JF, Tang G, Cao K, Wang R. *Am. J. Physiol. Heart Circ. Physiol* 2004;287:H2316. [PubMed: 15191893]
- Zhong G, Chen F, Cheng Y, Tang C, Du J. *J. Hypertens* 2003;21:1879. [PubMed: 14508194]
- Yang G, Cao K, Wu L, Wang R. *J. Biol. Chem* 2004;279:49199. [PubMed: 15347670]
- Wang J, Hegele RA. *Hum. Genet* 2003;112:404. [PubMed: 12574942]
- McCully KS. *Nat. Med* 1996;2:386. [PubMed: 8597939]
- Iwata C, Wang X, Uchida K, Nakanishi N, Hattori Y. *Life Sci* 2007;80:873. [PubMed: 17137603]
- Ignarro LJ. *Biosci. Rep* 1999;19:51. [PubMed: 10888468]
- Cooke CL, Davidge ST. *Cardiovasc. Res* 2003;60:635. [PubMed: 14659809]

22. Boehning D, Sedaghat L, Sedlak TW, Snyder SH. J. Biol. Chem 2004;279:30927. [PubMed: 15175337]
23. Michel JB, Feron O, Sase K, Prabhakar P, Michel T. J. Biol. Chem 1997;272:25907. [PubMed: 9325323]
24. Huang PL, et al. Nature 1995;377:239. [PubMed: 7545787]
25. Lake-Bruse KD, et al. Am. J. Physiol.Heart Circ. Physiol 1999;2:H770.
26. Shesely EG, et al. Proc. Natl. Acad. Sci. U.S.A 1996;93:13176. [PubMed: 8917564]
27. Zhao W, Ndisang JF, Wang R. Can. J. Physiol. Pharmacol 2003;81:848. [PubMed: 14614520]
28. Feletou M, Vanhoutte PM. Ann. Med 2007;39:495. [PubMed: 17852039]
29. Brandes RP, et al. Proc. Natl. Acad. Sci. U.S.A 2000;97:9747. [PubMed: 10944233]
30. We thank C. Watkins for experimental and conceptual assistance. Supported by U.S. Public Health Service grant MH18501 and Research Scientist Award DA00074 to S.H.S., and operating grants of Canadian Institutes of Health Research to L.W. and R.W.

**Fig. 1.**

Phenotype of CSE male knockout mice. (A) Reduced H₂S production from aorta and heart tissues in CSE^{-/-} mice and CSE^{+/-} mice. Number of mice are given for each group; $n = 16$. (B) Reduced serum H₂S level in CSE^{-/-} mice and CSE^{+/-} mice ($n = 8$ to 10). (C) Age-dependent increase in blood pressure of CSE^{-/-} mice and CSE^{+/-} mice ($n = 12$). (D) H₂S administration lowers systolic arterial blood pressure in 10-week-old CSE^{-/-} mice ($n = 13$ to 15). (E) Increased plasma homocysteine level in CSE^{-/-} mice and CSE^{+/-} mice ($n = 19$). (F) Decreased plasma L-cysteine level in CSE^{-/-} mice and CSE^{+/-} mice ($n = 15$). All results are means \pm SEM. * $P < 0.05$ versus WT; # $P < 0.05$ versus heterozygote.





Fig. 2. Impaired endothelial function in CSE mutant mice. Contraction of mesenteric artery evoked by phenylephrine (**A**) and relaxation of mesenteric artery by sodium nitroprusside (**B**), H₂S (**C**), and methacholine (**D**). $n = 15$ for each group. All results are means \pm SEM. * $P < 0.05$ versus WT; # $P < 0.05$ versus heterozygote. (**E**) Endothelial removal abolishes methacholine-induced relaxation of mesenteric artery. No relaxation occurs in vessels of WT or mutant mice after stripping of the endothelium. For CSE^{-/-} mice, $n = 8$; and for CSE^{+/+} mice, $n = 9$. (**F**) Immunohistochemical localization of CSE to arterial endothelium (black arrows) is abolished in CSE^{-/-} mice. Scale bars, 20 μ m.

**Fig. 3.**

CSE is activated by calcium-calmodulin upon muscarinic cholinergic stimulation of vascular endothelial cells. (A) Calmodulin binds CSE in vitro in the presence of calcium (2 mM). The interaction is diminished by the calcium chelator, EGTA (1 mM), as well as the calmodulin antagonist W7 (100 μM). (B) Calcium-calmodulin activates purified CSE in vitro. Calcium (1 mM) or calmodulin (5 μM) separately has no effect on CSE activity. $n = 3$. * $P < 0.05$ versus the control. (C) CSE is endogenously expressed in bovine aortic endothelial cells (BAECs) and human umbilical vein endothelial cells (HUVECs). (D) CSE is activated in BAECs treated with the calcium ionophore A23187 (1 μM) for 10 min. Incubation beforehand with the acetoxymethyl ester of the intracellular calcium chelator BAPTA (BAPTA-AM, 50 μM) or W7 (50 μM) for 30 min prevents CSE activation. $n = 3$. * $P < 0.05$ versus A23187 treatment; # $P < 0.05$ versus control. (E) CSE is strongly activated in BAECs treated with the muscarinic agonist methacholine (MCh, 1 μM) for 10 min. The activation is twice as much as with similar concentrations of A23187. The stimulatory effect of MCh is abolished by the muscarinic antagonist atropine (50 μM) ($n = 3$ or 4). * $P < 0.05$ versus control; # $P < 0.05$ versus all other groups. (F) CSE is the endogenous H₂S generator in BAECs. Transfecting cells with 100 nM

CSE-specific short interfering RNA (CSE-siRNA) for 48 hours markedly diminishes the enhanced H₂S production observed with A23187 (1 μ M) or MCh (1 μ M). Western blotting showed that CSE protein is decreased about 60 to 70% by CSE-siRNA ($n = 3$). * $P < 0.05$.